

This is a repository copy of *TWIST1 Integrates Endothelial Responses to Flow in Vascular Dysfunction and Atherosclerosis*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/123766/>

Version: Published Version

Article:

Mahmoud, Marwa M, Kim, Hyejeong Rosemary, Xing, Rouyu et al. (15 more authors) (2016) TWIST1 Integrates Endothelial Responses to Flow in Vascular Dysfunction and Atherosclerosis. *Circulation Research*. pp. 450-62. ISSN: 1524-4571

<https://doi.org/10.1161/CIRCRESAHA.116.308870>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

TWIST1 Integrates Endothelial Responses to Flow in Vascular Dysfunction and Atherosclerosis

Marwa M. Mahmoud,* Hyejeong Rosemary Kim,* Rouyu Xing, Sarah Hsiao, Akiko Mammoto, Jing Chen, Jovana Serbanovic-Canic, Shuang Feng, Neil P. Bowden, Richard Maguire, Markus Ariaans, Sheila E. Francis, Peter D. Weinberg, Kim van der Heiden, Elizabeth A. Jones, Timothy J.A. Chico, Victoria Ridger, Paul C. Evans

Rationale: Blood flow–induced shear stress controls endothelial cell (EC) physiology during atherosclerosis via transcriptional mechanisms that are incompletely understood. The mechanosensitive transcription factor TWIST is expressed during embryogenesis, but its role in EC responses to shear stress and focal atherosclerosis is unknown.

Objective: To investigate whether TWIST regulates endothelial responses to shear stress during vascular dysfunction and atherosclerosis and compare TWIST function in vascular development and disease.

Methods and Results: The expression and function of TWIST1 was studied in EC in both developing vasculature and during the initiation of atherosclerosis. In zebrafish, *twist* was expressed in early embryonic vasculature where it promoted angiogenesis by inducing EC proliferation and migration. In adult porcine and murine arteries, TWIST1 was expressed preferentially at low shear stress regions as evidenced by quantitative polymerase chain reaction and en face staining. Moreover, studies of experimental murine carotid arteries and cultured EC revealed that TWIST1 was induced by low shear stress via a GATA4-dependent transcriptional mechanism. Gene silencing in cultured EC and EC-specific genetic deletion in mice demonstrated that TWIST1 promoted atherosclerosis by inducing inflammation and enhancing EC proliferation associated with vascular leakiness.

Conclusions: TWIST expression promotes developmental angiogenesis by inducing EC proliferation and migration. In addition to its role in development, TWIST is expressed preferentially at low shear stress regions of adult arteries where it promotes atherosclerosis by inducing EC proliferation and inflammation. Thus, pleiotropic functions of TWIST control vascular disease and development. (*Circ Res.* 2016;119:450–462. DOI: 10.1161/CIRCRESAHA.116.308870.)

Key Words: atherosclerosis ■ cholesterol ■ gastrulation ■ obesity ■ transcription factors

Endothelial cells (EC) are exquisitely sensitive to shear stress (mechanical drag), which is imposed on the vessel wall by flowing blood. Although atherosclerosis is promoted by systemic risk factors (eg, cholesterol, smoking, obesity, and age), it develops preferentially near branches and bends exposed to complex blood flow that generates shear stress with low-magnitude and significant variation in direction (eg, oscillations and tangential shear). By contrast, atheroprotected sites are exposed to shear stress with high magnitude and uniform direction.^{1–4} Shear stress controls fundamental processes in EC, including inflammation, proliferation, and migration. Low shear stress promotes atherosclerosis by priming EC for enhanced expression of inflammatory molecules

(eg, intercellular adhesion molecule-1 [ICAM-1] and vascular cell adhesion molecule-1 [VCAM-1]) that coordinate the migration of leukocytes from the blood stream to the vascular wall.^{3–8} Low shear stress is also associated with enhanced EC proliferation,^{9–11} a process where EC lose contact with neighboring cells, thereby enhancing vascular permeability to cholesterol-rich lipoproteins to drive lesion formation.¹² Shear stress controls EC physiology, in part, via transcriptional and post-transcriptional mechanisms that are incompletely characterized.^{4–6,13–18}

A recent microarray study from our group demonstrated that the helix-loop-helix transcription factor TWIST1 and the zinc finger transcription factor GATA4 were enriched in EC at

Original received April 25, 2016; revision received May 20, 2016; accepted May 27, 2016. In April 2016, the average time from submission to first decision for all original research papers submitted to *Circulation Research* was 15.28 days.

From the Department of Infection, Immunity and Cardiovascular Disease, INSIGNEO Institute for In Silico Medicine, and the Bateson Centre, University of Sheffield, Sheffield, United Kingdom (M.M.M., H.R.K., S.H., J.S.-C., S.F., N.P.B., R.M., M.A., S.E.F., T.J.A.C., V.R., P.C.E.); ERASMUS MC, Rotterdam, The Netherlands (R.X., K.v.d.H.); Vascular Biology Program, Department of Surgery (A.M.) and Department of Ophthalmology (J.C.), Boston Children's Hospital, Harvard Medical School, MA; Department of Bioengineering, Imperial College London, London, United Kingdom (P.D.W.); and Department of Cardiovascular Science, Katholieke Universiteit Leuven, Leuven, Belgium (E.A.J.).

*These authors contributed equally to this article.

The online-only Data Supplement is available with this article at <http://circres.ahajournals.org/lookup/suppl/doi:10.1161/CIRCRESAHA.116.308870/-/DC1>.

Correspondence to Paul C. Evans, PhD, Department of Infection, Immunity and Cardiovascular Disease, Medical School, University of Sheffield, Beech Hill Rd, Sheffield S10 2RX, United Kingdom. E-mail paul.evans@sheffield.ac.uk

© 2016 The Authors. *Circulation Research* is published on behalf of the American Heart Association, Inc., by Wolters Kluwer. This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution, and reproduction in any medium, provided that the original work is properly cited.

Circulation Research is available at <http://circres.ahajournals.org>

DOI: 10.1161/CIRCRESAHA.116.308870

Nonstandard Abbreviations and Acronyms

EC	endothelial cells
EGFP	enhanced green fluorescent protein
EndoMT	endothelial–mesenchymal transition
Hpf	hours post fertilization
qPCR	quantitative polymerase chain reaction
Tg	transgenic
WSS	wall shear stress

low shear stress atheroprone regions of the aorta (J. Serbanovic-Canic and P.C. Evans, unpublished data, 2016). *twist* was originally identified in *Drosophila* embryos where it controls gastrulation and other fundamental developmental processes.¹⁹ In vertebrates, TWIST regulates multiple diverse activities including development,^{20,21} epithelial–mesenchymal transition,²² and tumor metastasis.²³ GATA4 is also a key regulator of development. It controls cardiac specification²⁴ and genetic deletion of GATA4 in mice led to multiple cardiac phenotypes.²⁵ Of note, TWIST and GATA4 exert overlapping functions during atrioventricular valve development by inducing endothelial–mesenchymal transition (EndoMT), which describes a program of phenotypic changes including enhanced EC proliferation and migration.^{21,26} Although GATA4 and TWIST have well-recognized roles in some aspects of embryogenesis, their potential function in vascular development and disease is unknown.

Here, we demonstrate for the first time that low shear stress promotes GATA4-dependent induction of TWIST1 in EC. Studies using zebrafish revealed that *twist* was expressed in early embryonic vasculature where it promoted angiogenic sprouting by inducing EC proliferation and migration. In adult mammalian arteries, TWIST1 was expressed preferentially at atheroprone sites exposed to low shear stress where it promoted the development of atherosclerosis by inducing inflammation and EC proliferation. We conclude that TWIST is a shear stress–regulated transcription factor that regulates angiogenesis in embryos and drives focal EC dysfunction and atherosclerosis in adult arteries.

Methods

Mice

Male mice between 2 and 3 months of age were used. For cell tracking studies, transgenic Rosa26-tdTomato mice²⁷ were crossed with endothelial-SCL-Cre-ER^T mice containing a tamoxifen-inducible EC-specific Cre.²⁸ To activate Cre, tamoxifen was administered for 5 consecutive days (2 mg/mouse/d). Mice with conditional deletion of TWIST1 (*Twist1*^{CreKO}) were generated by crossing Tie2-Cre-expressing mice (Jackson Laboratory stock 004128) with TWIST1 floxed mice (*Twist1*^{fllox/flox}).²⁹ Mice with conditional EC deletion of GATA4 (called GATA4^{CreKO}) were generated by crossing endothelial-SCL-Cre-ER^T mice with GATA4 floxed mice (*Gata4*^{fllox/flox}). Constrictive cuffs were applied to the right carotid artery of isoflurane-anesthetized mice following published methods.^{6,30} The expression levels of specific proteins were assessed in EC by en face staining as previously described.^{5,6} Hypercholesterolemia was induced by tail vein injection of adeno-associated virus containing a gain-of-function mutated version of proprotein convertase subtilisin/kexin type 9 (rAAV8-D377Y-mPCSK9) gene followed by exposure to a western diet for 6 weeks as previously described.³¹ Lesions were detected by Oil Red O staining.

Cultured EC

Human umbilical vein EC and porcine aortic EC were isolated and cultured.³² Gene silencing was performed using 2 different small interfering RNAs (siRNAs) against TWIST1 (Silencer Select S14523, Ambion, and L-006434-00-0005 ON-TARGETplus SMARTpool; Dharmacon) or GATA4 (Silencer Select s5603, Ambion, and L-008244-00-0005 ON-TARGETplus SMARTpool; Dharmacon). They were exposed to flow using an orbital shaking system or Ibidi parallel-plate system.^{32,33} Quantitative RT-PCR, immunofluorescent staining and chromatin immunoprecipitation,⁶ and assays of permeability³⁴ were performed as described.

Zebrafish Embryos

Studies were performed using wild-type or transgenic zebrafish lines, *Tg(fli1:EGFP)* (endothelial EGFP), *Tg(kdr1:NLS-EGFP)* (endothelial nuclear EGFP), or *Tg(gata1:dsRed)* (red blood cell expression of dsRed). Whole-mount in situ hybridization was performed as described.³⁵ The *twist1b* gene was mutated using the CRISPR/Cas9 system and *twist1a* expression was reduced using a translation-blocking morpholino.³⁶

Statistics

Differences between samples were analyzed using an unpaired or paired Student *t* test or ANOVA (**P*<0.05, ***P*<0.01, and ****P*<0.001).

Results

Low Shear Stress Induced TWIST1 via a GATA4-Dependent Mechanism

A microarray study from our laboratory comparing EC at low shear stress (inner curvature of aortic arch) and high shear stress (outer curvature) regions in the porcine aorta revealed >800 differentially expressed genes (J. Serbanovic-Canic and P.C. Evans, unpublished data, 2016). They included TWIST1 and GATA4 that were enriched at the low shear stress site (subanalysis shown in Figure 1A). This observation was validated by quantitative polymerase chain reaction (qPCR) studies of an independent cohort of pigs (Figure 1B). Similarly, en face staining of the murine aortic endothelium demonstrated that TWIST1 and GATA4 proteins were expressed at higher levels at the inner curvature of the aortic arch (low shear stress) compared with the outer curvature (high shear stress; Figure 1C). Moreover, both of these transcription factors localized to the nucleus, suggesting that they are active at the low shear site.

Atherosusceptible regions of arteries are associated with increased inflammation and altered transport of substances to and from the arterial wall as well as low shear stress. Therefore, we used in vitro and in vivo models to examine whether shear stress per se is responsible for enhanced expression of TWIST1 and GATA4. Cultured EC were exposed to flow using 2 complementary systems, an orbital-plate and a parallel-plate apparatus. Computational fluid dynamic analysis demonstrated that the orbiting 6-well plate system generates low mean shear stress (4.8 dyn/cm²) with rapid variations in direction at the center and high mean shear stress (11.1 dyn/cm²) with uniform direction at the periphery.^{32,33} Using the orbital system, TWIST1 and GATA4 expression was elevated in Human umbilical vein EC or porcine aortic EC exposed to low (center) compared with high shear stress (periphery) or static conditions (Figure 2A, top, and Figure 2B; Online Figure I). Indeed, the majority of cells exposed to low shear stress expressed TWIST1 and GATA4, which localized almost




Figure 1. TWIST1 and GATA4 were preferentially expressed at low shear atherosusceptible sites. **A** and **B**, Expression of TWIST1 and GATA4 was studied at low wall shear stress (WSS; inner curvature) and high WSS (outer curvature) regions of the porcine aorta. **A**, Differentially expressed genes were identified using microarrays ($n=5$ pigs) and the expression patterns of TWIST1 and GATA4 are presented as a heat map with red indicating enrichment in expression. **B**, Results were validated by qRT-PCR of a different cohort of pigs ($n=5$). The expression level at the low WSS site is presented relative to the expression at the high WSS site (normalized to 1; dotted line). Mean levels \pm SEM are shown. **C**, Expression levels of TWIST1 or GATA4 in EC were assessed by en face staining of low (inner curvature) or high (outer curvature) WSS regions of the aorta in C57BL/6 mice (red). Endothelial cells (ECs) were identified by costaining with anti-CD31 antibodies conjugated to FITC (fluorescein isothiocyanate; green). Cell nuclei were identified using TO-PRO-3 (blue). Representative images and quantification of TWIST1 or GATA4 (mean \pm SEM) are shown. * $P<0.05$ and ** $P<0.01$, using a paired t test.

exclusively to the nucleus (Figure 2B). To delineate between the effects of shear stress magnitude and direction, a parallel-plate system was used to compare gene expression under low oscillatory (± 4 dyn/cm²), low unidirectional (4 dyn/cm²), and high unidirectional (13 dyn/cm²) flow. It revealed that TWIST1 and GATA4 expression was higher in EC exposed

to low compared with high or low oscillatory shear stress (Figure 2A, bottom), indicating that low-magnitude shear stress drives GATA4 and TWIST1 expression in cultured EC. We determined whether shear stress regulates GATA4 and TWIST1 expression in vivo by modifying flow in the murine carotid artery. This was achieved using a constrictive cuff that causes tapering of the lumen to generate high shear stress at the stenosis, low shear stress upstream, and low/oscillatory shear stress downstream.^{6,30} Cuff placement for 14 days led to enhanced expression of GATA4 and TWIST1 at the low shear stress site (Figure 2C), which is consistent with our in vitro data. By contrast, GATA4 was induced in the absence of TWIST1 at the low oscillatory shear stress site (Figure 2C). This apparent discrepancy with the in vitro data (where low oscillatory shear had no effect on GATA4 expression) may be because of differences in the frequency of oscillation or features of the flow waveform between in vitro and in vivo systems.³⁰

We hypothesized that GATA4 and TWIST1 may be cross-regulated because both were induced by low wall shear stress (WSS; Figures 1 and 2) and have overlapping functions in other systems.²⁶ Potential cross-regulation was tested by silencing of GATA4 or TWIST1 using 2 different siRNA sequences and validation by qPCR and Western blotting (Online Figure II). GATA4 was required for TWIST1 expression in EC exposed to low shear stress (Figure 3A), whereas TWIST1 silencing did not alter GATA4 expression (data not shown). We hypothesized that GATA4 may positively regulate transcription of TWIST1 because putative GATA4-binding sites were identified in its promoter (Figure 3B). Consistent with this, chromatin immunoprecipitation followed by qPCR demonstrated that TWIST1 promoter sequences coprecipitate with anti-GATA4 antibodies but not with isotype-matched irrelevant IgG (Figure 3B), indicating that GATA4 binds to the promoter region of TWIST1 to induce transcriptional activation. To validate these observations, we assessed the effects of genetic deletion of GATA4 on endothelial expression of TWIST1 in the murine aorta. GATA4 deletion from EC was achieved by crossing GATA4^{fllox/fllox} mice with endothelial-SCL-Cre-ER transgenics followed by 5 days of tamoxifen treatment (generating GATA4^{cKO}). To validate genetic deletion, it was shown by en face staining that GATA4 expression was absent from EC in GATA4^{cKO} mice (Online Figure IIIB). En face staining revealed that the expression of TWIST1 at the low shear stress site was reduced in GATA4^{cKO} compared with GATA4^{fllox/fllox} mice (Figure 3C), indicating that GATA4 positively regulates TWIST1 in atherosusceptible endothelium. Thus, we conclude that GATA4 induces TWIST1 in cells exposed to low shear stress, and this pathway is suppressed in cells exposed to high shear stress. Our observation that TWIST1 was not expressed in GATA4-expressing EC that were exposed to low oscillatory shear stress in vivo (Figure 2C) indicates that factors other than GATA4 are required for TWIST1 expression. Candidate coregulators include Notch1 that positively regulates TWIST1 in other systems^{37,38} and was induced by low (but not low oscillatory shear stress) in modified carotid arteries (Online Figure IV). Collectively, our in vitro and in vivo data reveal that low-magnitude WSS positively regulates TWIST1 expression via




Figure 2. Low shear stress induced TWIST1 and GATA4. **A**, Human umbilical vein endothelial cells (HUVECs) were exposed to orbital flow to generate low (center) or high (periphery) wall shear stress (WSS) or were cultured under static conditions. Alternatively, HUVECs were exposed to high, low, or low/oscillatory WSS using a parallel-plate system. After 72 h, levels of TWIST1 or GATA4 transcripts were quantified by qRT-PCR. **B**, HUVECs were exposed to orbital flow to generate low (center) or high (periphery) WSS for 72 h or were cultured under static conditions. Expression of TWIST1 and GATA4 was determined by immunofluorescent staining (green) and costaining using DAPI (4',6-diamidino-2-phenylindole; blue). Bar=50 μ m. Fluorescence intensities were measured in multiple cells in 3 independent experiments, and mean values \pm SEM are shown. **C**, Flow-altering, constrictive cuffs were placed on the right carotid arteries of C57BL/6 mice. They generated anatomically distinct regions exposed to low, high, and low oscillatory WSS (as indicated). Right (experimental) and left (sham-operated) carotid arteries were harvested after 14 days, and en face staining was performed using anti-TWIST1 or anti-GATA4 antibodies (red), anti-CD31 antibodies conjugated to FITC (fluorescein isothiocyanate; green), and the nuclear counter stain TO-PRO-3 (blue). Representative images and quantification of TWIST1 or GATA4 expression (mean \pm SEM) are shown. Bar=10 μ m. Data were pooled from 5–6 independent experiments. * P <0.05 and ** P <0.01, using a 1-way ANOVA.




Figure 3. GATA4 induced TWIST1 in response to low shear stress. **A**, Human umbilical vein endothelial cells (HUVECs) were transfected with 2 different small interfering RNAs (siRNAs) targeting GATA4 or with scrambled sequences and exposed to orbital flow for 72 h. Cells exposed to low wall shear stress (WSS; center of well) were collected, and TWIST1 transcript levels were quantified by qRT-PCR. **B**, GATA4-binding sites (A/T)GATAA(G) were predicted in the TWIST1 5' untranslated region, and their position (blue text) in relation to the translational start site (red text) is shown. HUVECs were exposed to orbital flow for 72 h before chromatin immunoprecipitation using anti-GATA4 or irrelevant isotype control antibodies. The levels of TWIST1 promoter DNA were assessed by qPCR and fold enrichment in anti-GATA4 compared with control precipitates was calculated. Mean levels \pm SEM are shown. **A** and **B**, Data from at least 3 independent experiments were pooled. * P <0.05 and ** P <0.01 using an unpaired t test. **C**, Endothelial cells (ECs) at low WSS (susceptible) or high WSS (protected) regions of the aorta in *GATA4^{cko}* or *GATA4^{flox/flox}* mice were studied by en face staining. TWIST1 (red) expression was assessed by en face staining. Representative images are shown, and TWIST1 expression was quantified (mean fluorescence \pm SEM is shown). Bar=10 μ m. * P <0.05 using a 2-way ANOVA.

a transcriptional pathway that requires GATA4 among other factors. We conclude that low shear induces TWIST1 via increasing its synthesis but cannot rule out additional mechanisms that reduce the rate of degradation.

Finally, we used a Cre-based cell-tracking system to investigate whether low shear stress induces TWIST1 expression in fully differentiated EC and/or whether it promotes the homing of TWIST1-positive endothelial progenitor cells to the vascular wall. Cell tracking was performed using transgenic SCL-Cre-ER^T/R26R-tdTomato mice where EC were labeled with tdTomato in response to tamoxifen treatment. Notably, transient tamoxifen treatment allowed mature EC that reside in vessels (tdTomato-positive; Online Figure V) to be distinguished from those generated by subsequent homing of

progenitor cells (tdTomato-negative).²⁸ Using this system, we labeled EC with tdTomato before placement of a constrictive cuff on the carotid artery for 2 weeks and analysis of GATA4 and TWIST1 expression. We observed that EC at the low shear stress site that expressed GATA4 or TWIST1 were tdTomato-positive (Online Figure VB), indicating that GATA4–TWIST1 signaling can be induced by low shear stress in fully differentiated EC of adult arteries and that progenitor EC homing was negligible.

Twist Promoted EC Proliferation and Migration in Embryonic Vasculature

Given that TWIST is expressed in embryogenesis^{20,21} and at atheroprone regions of adult arteries (Figure 1), we wished




Figure 4. *twist* promoted intersegmental vessel sprouting in zebrafish embryos. **A**, The expression of *twist1a*, *twist1b*, and *twist2* was studied at 24 to 75 hours post fertilization (hpf) by qPCR of the trunk and tail of embryos. Data were pooled from $n \geq 15$ embryos studied in 3 independent experiments and mean values \pm SEs are shown. **B**, The expression of *twist1a*, *twist1b*, and *twist2* was studied at 24 and 52 hpf by in situ hybridization. Data are representative of the majority of embryos analyzed (proportion indicated lower right of each part) and were closely similar in at least 3 independent experiments. Higher magnification insets are shown (marked in **top**). Bar=100 μ m. **C**, Zebrafish embryos (wild-type, *Tg(fli1:EGFP)*, or *Tg(kdr1:NLS-EGFP)*) were treated with *twist1b* mRNA (to enforce expression) or treated with mCherry mRNA as a control. They were studied at 24 to 27 hpf using confocal microscopy to visualize endothelial cell (EC) nuclei (*Tg(kdr1:NLS-EGFP)*; **top**) and angiogenic sprouts (*Tg(fli1:EGFP)*; **bottom**; arrows). Representative images are shown. Cell numbers and the length of intersegmental vessels (ISVs; third to fifth vessels in the field view) were quantified in multiple embryos, and mean values \pm SEM are shown (**right**). **D**, The *twist1b* coding sequence was mutated by introduction of a 4 bp deletion causing a frameshift and premature (Continued)

to compare its function in EC during vascular development and disease. The potential role of TWIST in vascular development was studied using zebrafish that possess 3 orthologs of mammalian TWIST genes: *twist1b*, *twist1a*, and *twist2*. Quantitative RT-PCR and in situ hybridization revealed that each of them was expressed in the trunk vasculature at 24 hours post fertilization and subsequently declined in expression at later developmental stages (Figures 4A and 4B). The function of *twist* was studied by enhancing its expression in vessels via injection of mRNA (gain of function). Transgenic fish expressing green fluorescent protein in EC (*Tg(fli1:EGFP)*) or endothelial nuclei (*Tg(kdr1:NLS-EGFP)*) were used to facilitate visualization of the vasculature and cell counting, respectively. Dynamic imaging and analysis of fixed embryos demonstrated that *twist1b* overexpression led to enhanced sprouting of intersegmental vessels associated with increased EC migration and proliferation (Online Movies I and II; Figure 4C). Conversely, suppression of *twist1* expression (loss-of-function) reduced intersegmental vessel formation (Figure 4D). The latter study was performed by mutating *twist1b* using CRISPR/Cas9 genome editing to introduce a 4 bp deletion that caused a frame shift and introduced a stop codon before the helix-loop-helix domain (designated *twist1b^{sh423}*). Intersegmental vessel sprouting was not altered in *twist1b^{sh423}* mutants (data not shown), and we hypothesized that this was because of compensation from *twist1a* that is closely homologous. To test this, *twist1b^{sh423/+}* fish were incrossed to generate *twist1b^{sh423/sh423}* mutant (25%), heterozygotes (50%), and *twist1b^{+/+}* wild-type (25%) embryos that were subsequently treated with a morpholino directed against *twist1a*. Embryos displayed variable phenotypes and were classified into those with minimal sprouting (severe phenotype) and those with intermediate levels of sprouting (mild phenotype). Notably, the severe phenotype was significantly enriched in the homozygous mutant group compared with wild types, indicating that mutation of *twist1b* enhanced the *twist1a* knock down effect (Figure 4D). Overall, these data suggest that *twist1b* and *twist1a* overlap at a functional level to promote angiogenic sprouting. Similar observations were made in a different vascular bed, the developing subintestinal vein. At this site, *twist* genes were expressed in early developing vasculature and enforced expression of *twist1b* enhanced angiogenic sprouting (Online Figure VI). Collectively, these data indicate that *twist* promotes developmental angiogenesis by inducing EC migration and proliferation.

Figure 4 Continued. stop (mutant allele designated *twist1b^{sh423}*). *twist1b^{sh423/+} Tg(fli1:EGFP)* fish were incrossed, and embryos were treated with a morpholino directed against *twist1a*. Sprouting of ISV was assessed at 34 hpf. Embryos were classified into those that displayed minimal sprouting (severe phenotype) and those with intermediate levels of sprouting (mild phenotype). Genotyping was subsequently performed, and the proportion of *twist1b* homozygous mutants (*twist1b^{sh423/sh423}*) and *twist1b* homozygous wild-types (*twist1b^{+/+}*) in the severe phenotype group was calculated (% indicated). Representative images are shown. Data were closely similar in 3 independent experiments. *** $P < 0.001$ and ** $P < 0.01$ using a 1-way ANOVA (**A**) or unpaired *t* test (**C** and **D**). CVP indicates caudal vein plexus.




Figure 5. TWIST1 and GATA4 promoted proliferation in endothelial cells (ECs) exposed to low shear stress. Human umbilical vein endothelial cells (HUVECs) were treated with 2 different small interfering RNAs (siRNAs) targeting TWIST1 or GATA4 (designated i and ii) or with scrambled nontargeting siRNA or remained untransfected. Cells were subsequently cultured in 6-well plates before exposure to orbital flow to generate low (center, C) or high (periphery, P) wall shear stress (WSS) for 72 h. Alternatively, cells were maintained under static (S) conditions. **A**, Cell proliferation was quantified by immunofluorescent staining using anti-proliferating cell nuclear antigen (PCNA) antibodies and costaining using DAPI (4',6-diamidino-2-phenylindole). Images are representative of those generated in 3 independent experiments using 1 version of the gene-specific siRNA or scrambled control sequences (bar=50 μ m). The percentage of PCNA-positive cells were calculated for multiple fields of view in at least 3 independent experiments, and mean values \pm SEM are shown. * P <0.05 using a 2-way ANOVA. **B–F**, The expression of cell cycle regulators and endothelial–mesenchymal transition genes was quantified using qRT-PCR. **B**, The expression level in cells at the center (low WSS) is presented relative to the expression at the periphery (high WSS; normalized to 1; dotted line). **C–F**, Transfected cells were exposed to low WSS (center). The expression level in cells transfected with gene-targeting siRNA is presented relative to the expression in cells transfected with scrambled control siRNA (normalized to 1; dotted line). Data were pooled from 3 independent experiments, and mean values \pm SEM are shown. * P <0.05, ** P <0.01, and *** P <0.001 using an unpaired t test.

TWIST1 and GATA4 Promote EC Proliferation and Inflammation Under Low Shear Stress Conditions

We wished to know whether GATA4–TWIST1 signaling influences EC dysfunction and the initiation of atherosclerosis at low shear sites. This was initially studied using cultured EC exposed to flow under well-controlled experimental conditions. We hypothesized that TWIST1 and GATA4 promote proliferation in EC exposed to low shear because they regulate cell cycle progression in other contexts.^{20,21} Consistent with this concept, gene silencing of TWIST1 or GATA4 using 2 different siRNAs significantly reduced proliferation of EC exposed to low shear using the orbital system (Figure 5A, center; compare 2 with 5, 8, 11, and 14) but did not alter EC exposed to high shear (periphery; compare 3 with 6, 9, 12, and 15) or static conditions (compare 1 with 4, 7, 10, and 13). These data were validated using a parallel-plate system to demonstrate that silencing of TWIST1 or GATA4 reduced proliferation of EC that were exposed to low but not high shear stress conditions (Online Figure VII). Thus, we conclude that low

shear stress activation of GATA4–TWIST1 signaling drives EC proliferation.

We investigated whether the mechanism for enhanced proliferation involved induction of regulators of cell division. qPCR revealed that cyclin D1, cyclin G2, and cyclin-dependent kinase 4 were induced in cultured EC by the application of low shear stress (Figure 5B), and their expression was reduced by silencing of GATA4 or TWIST1 using 2 different siRNAs to ensure specificity (Figures 5C and 5D). Thus, GATA4–Twist1 signaling under low shear stress induces cyclins and cyclin-dependent kinase 4. We also investigated whether GATA4 and TWIST1 promote the expression of regulators of EndoMT (a proproliferative cellular transition) under low shear stress conditions. qPCR demonstrated that the expression of Slug, Snail, N-cadherin, and α -smooth muscle actin was elevated in EC exposed to low WSS compared with EC exposed to high WSS (Figure 5B). Silencing of GATA4 or TWIST1 (using 2 different siRNAs) significantly reduced the expression of Snail but did not influence Slug, N-cadherin, or




Figure 6. TWIST1 and GATA4 promoted permeability in endothelial cells (ECs) exposed to low shear stress. The influence of TWIST1 and GATA4 on EC permeability under low wall shear stress (WSS) was studied. **A**, Cells cultured on transwell inserts were exposed to orbital flow (low WSS) or static conditions for 72 h before assessment of endothelial permeability under static conditions using rhodamine (Rd) albumin as a tracer. A schematic is shown (left). The concentration of Rd-albumin in the lower compartment was measured, and mean values±SEM are shown (right). **B**, Human umbilical vein endothelial cells (HUVECs) were treated with 2 different small interfering RNAs (siRNAs) targeting TWIST1 or GATA4 (designated i and ii) or with scrambled nontargeting siRNA. Transfected cells cultured on transwell inserts were exposed to orbital flow (low WSS) or static conditions for 72 h before assessment of endothelial permeability using Rd-albumin. The concentration of Rd-albumin in the lower compartment was measured in 3 independent experiments, and mean values±SEM are shown. * $P<0.05$ using a paired t test.

α -smooth muscle actin (Figures 5E and 5F). Thus, we conclude that GATA4 and TWIST1 are required for the induction of Snail but that other pathways are also necessary for EndoMT under low shear stress. Collectively, our observations reveal that GATA4–TWIST1 signaling promotes EC proliferation under low shear conditions by inducing cyclins and regulators of EndoMT.

Atherosclerosis is driven by vascular inflammation involving EC expression of adhesion molecules and cytokines. The influence of GATA4 and TWIST1 on inflammatory activation was studied using cultured EC exposed to low or high shear stress. qPCR revealed that inflammatory VCAM-1 and ICAM-1 were expressed at higher levels in EC exposed to low shear (center of orbiting plate) compared with high shear, and their expression under low shear was reduced by silencing of GATA4 or TWIST1 (Online Figure VIII). Thus, GATA4–TWIST1 signaling under low shear stress induces inflammatory adhesion molecules. We also assessed whether GATA4–TWIST1 signaling influences the expression of several molecules involved in mechanosensing (Piezo, CD31, and VEGFR2). However, the expression of Piezo, CD31, and VEGFR2 was not significantly altered under low or high shear stress (Online Figure IX) or by silencing of GATA4 or TWIST1 (Online Figure IX), suggesting that GATA4–TWIST1 signaling may not alter the responsiveness of EC to shear stress.

Given the links between EC turnover, inflammatory activation, and vascular leakiness,¹² we next determined whether GATA4 and TWIST1 regulate EC permeability using EC cultured on transwell inserts. Exposure of EC to low shear stress (orbited) for 72 hours enhanced permeability to rhodamine albumin compared with static conditions as previously described³⁴ (Figure 6A). Silencing of TWIST1 or GATA4 (using 2 different siRNAs) reduced permeability in EC exposed to low WSS (orbited) but not in cells exposed to static conditions (Figure 6B), indicating that both genes promote permeability in response to low shear stress. Thus, TWIST1 and GATA4 are positive regulators of proliferation, inflammatory activation, and permeability under low shear conditions.

TWIST1 and GATA4 Promote Atherogenesis by Enhancing EC Proliferation and Inflammation at Low Shear Stress Regions

Given their roles in regulating inflammation and EC proliferation in response to low shear stress, we hypothesized that GATA4 and TWIST1 may influence the initiation of atherosclerosis. Thus, the function of both molecules was studied at low shear stress sites in adult murine arteries using genetic deletion approaches. TWIST1 was deleted from EC by crossing floxed mice (TWIST1^{lox/flox}) with Tie2-Cre transgenics (generating TWIST1^{CKO}; Online Figure IIIA), whereas GATA4 deletion from EC was studied using GATA4^{CKO} mice (as above, Online Figure III). En face staining of Ki67 revealed that EC proliferation was enhanced in EC exposed to low compared with high shear stress as reported,¹¹ and deletion of GATA4 or TWIST1 from EC significantly reduced proliferation at low shear stress sites in the aorta (Figure 7A). Deletion of TWIST1 from EC reduced the number of cells per field of view at the low shear site, whereas deletion of GATA4 did not (Figure 7A), suggesting that these molecules

may have divergent effects on factors that influence cell density, for example, viability, migration, or shape. Parallel en face staining studies demonstrated that inflammatory ICAM-1 expression was enhanced at the low shear site, and its expression was reduced by deletion of TWIST1 from EC (Online Figure VIII). Plaque formation was studied by inducing hypercholesterolemia via adeno-associated virus delivery of PCSK9 followed by high fat feeding for 6 weeks. Genetic deletion of GATA4 or TWIST1 in EC reduced lesion area in the aorta, indicating that GATA4–TWIST1 signaling contributes to atherosclerosis (Figure 7B). It is unlikely that either molecule promotes atherogenesis by altering lipid metabolism because their deletion from EC did not influence cholesterol or triglyceride levels in plasma (Table 1). Collectively, these data indicate that GATA4–TWIST1 signaling promotes EC proliferation, inflammation, and lesion formation at low shear regions of arteries.

Discussion

TWIST1 can be regulated by mechanical force during *Drosophila* development³⁹ and in response to tumor stiffness.⁴⁰ Here, we demonstrate that mechanical forces also regulate TWIST expression in vascular endothelial cells. Specifically, low shear stress induced TWIST1 in EC via the transcription factor GATA4, thereby enriching TWIST expression at atheroprone sites of arteries. TWIST1 expression contributed to atherosclerosis by enhancing vascular inflammation and driving EC proliferation associated with vascular leakiness. Thus, although TWIST is a central coordinator of embryogenesis, this transcription factor also contributes to the initiation of focal atherosclerosis in adult arteries.

The first part of our study focused on the regulation of TWIST1 by shear stress using in vitro systems and modified murine carotid arteries. Notably, it was observed that TWIST was induced in EC exposed to low-magnitude shear stress with uniform direction but was not enhanced in cells under low oscillatory shear. The mechanism underlying the specificity of TWIST1 induction by low shear stress is uncertain, but it is plausible that unidirectional and oscillatory flow activates different sets of mechanoreceptors that have divergent effects on TWIST1 expression. Our findings are consistent with previous demonstrations that low unidirectional and low oscillatory shear have distinct effects on vascular physiology. Of particular relevance, the imposition of low shear stress in carotid arteries of hypercholesterolemic mice leads to the formation of atherosclerotic plaques with features of instability including a thin fibrous cap and inflammation, whereas the imposition of low oscillatory shear induces stable lesions.³⁰ Thus, it will be of interest in future studies to determine whether the induction of TWIST by low shear stress contributes to subsequent plaque inflammation and instability.

A combination of gene silencing, chromatin immunoprecipitation, and genetic deletion studies demonstrated that GATA4 is required for the induction of TWIST by low shear. However, although GATA4 and TWIST were functionally linked in EC exposed to low shear, they were uncoupled in cells exposed to low oscillatory shear where GATA4 was induced in the absence of TWIST expression. This observation




Figure 7. TWIST1 and GATA4 promote endothelial cell (EC) proliferation and lesions at low shear stress sites. A, EC at low wall shear stress (WSS; susceptible) or high WSS (protected) regions of the aorta were studied by en face staining in *Twist1*^{cKO} or *Twist1*^{flox/flox} mice (**top**) or in *Gata4*^{cKO} or *Gata4*^{flox/flox} mice (**bottom**). EC proliferation was quantified by anti-Ki67 staining (red). EC were identified by costaining with anti-CD31 antibodies (green). Cell nuclei were identified using TOPRO3 (blue). The proportion of Ki67-positive cells, and number of cells was calculated and mean values±SEM are shown. Bar=10 μm. Data were pooled from 5 independent experiments. **B**, *Twist1*^{cKO} or *Twist1*^{flox/flox} mice (**right**) or *Gata4*^{cKO} or *Gata4*^{flox/flox} mice (**left**) were treated with AAV-PCSK9. After 1 wk, they were exposed to a Western diet for 6 wk. Lesions were stained using Oil Red O and quantified using image J software. Representative images are shown (Bar=1 mm). The percentage of lesion coverage was calculated. Data were pooled from multiple mice and mean values±SEM are shown. ***P*<0.01 and **P*<0.05 using a 2-way ANOVA (**A**) or unpaired *t* test (**B**).

Table 1. Deletion of TWIST1 or GATA4 From Endothelial Cells Did Not Alter Plasma Lipids

	GATA4 ^{flx/flx}	GATA4 ^{CKO}	P Value
Cholesterol	9.0±2.9	12.9±1.8	0.3
Triglycerides	1.4±0.4	1.7±0.2	0.4
	TWIST1 ^{flx/flx}	TWIST1 ^{CKO}	P Value
Cholesterol	25.9±2.0	20.5±6.7	0.3
Triglycerides	3.74±0.8	4.4±2.4	0.4

TWIST1^{CKO} or TWIST1^{flx/flx} mice (right) or GATA4^{CKO} or GATA4^{flx/flx} mice (left) were treated with AAV-PCSK9. After 1 wk, they were exposed to a western diet for 6 wk. The levels of total cholesterol and triglycerides in plasma were measured. Data were pooled from multiple mice, and mean values±SEM are shown.

indicates that GATA4 is necessary but not sufficient for the induction of TWIST1, suggesting the involvement of other factors that are activated under specific shear stress environments. Notch is a candidate coregulator of TWIST1 because it promotes TWIST expression during development^{35,36} and was induced specifically by low shear stress in murine carotid arteries. This model explains the expression of TWIST1 under low shear (where GATA4 and Notch are both regulated) and its absence under low oscillatory shear (where GATA4 was activated without Notch). Future studies are required to understand the mechanical regulation of GATA4/Notch cross talk and whether these factors cooperate to induce TWIST1 at atheroprone regions. A further outstanding question relates to the mechanoreceptors that regulate TWIST induction in response to low shear stress. Bone morphogenic proteins are candidates because they can be upregulated by low shear stress⁴¹ and cross talk with TWIST1.⁴² Their potential interaction with TWIST1 in vascular EC should now be investigated.

The second part of our study focused on the function of TWIST1 in EC. Using conditional knockout approaches, it was demonstrated that deletion of GATA4 or TWIST1 from EC significantly reduced lesion formation at low shear stress sites in hypercholesterolemic mice. GATA4–TWIST1 signaling in EC drives atherosclerosis through several mechanisms. First, GATA4 and TWIST1 activation was shown to enhance focal endothelial inflammatory activation, which drives lesion formation by promoting leukocyte recruitment to the vascular wall.^{3–7} Second, GATA4–TWIST1 signaling at low shear stress regions induced EC proliferation, which can enhance the permeability of arteries to cholesterol-containing lipoproteins.¹² Consistent with this, we observed that GATA4 and TWIST1 enhanced permeability in EC monolayers exposed to low shear stress. The mechanism linking GATA4–TWIST1 signaling to EC proliferation involves cyclins, which were induced by low WSS via a GATA4- and TWIST1-dependent mechanism. EC proliferation at atheroprone sites is regulated by several other signaling intermediaries including p53^{9,32}, JNK1,⁴³ and miR-126-5p,¹¹ and it will be important in future studies to assess how these pathways interact with the GATA4–TWIST1 signaling pathway.

Our study also revealed that GATA4 and TWIST1 induced molecules involved in EndoMT (including Snail, α -smooth muscle actin, and N-cadherin) in EC exposed to low shear. EndoMT is an example of cellular plasticity characterized by a program of morphological and physiological changes that involve loss of apical/basal polarity, disruption of intercellular junctions, increased proliferation, and migration of cells into

surrounding tissue.²⁶ Thus, activation of EndoMT by GATA4–TWIST1 signaling may contribute to enhanced proliferation at low shear sites. The canonical function of EndoMT is in the development of valves from EC in the atrioventricular canal²¹; however, EndoMT has also been implicated in disease processes including cerebral cavernous malformations.⁴⁴ Of particular relevance, 2 recent studies demonstrated that EndoMT can be induced by disturbed flow and is a driver of atherosclerosis.^{45,46} Thus, our observation that low shear induces GATA4–TWIST1 signaling provides a molecular mechanism to explain the induction of EndoMT by low shear stress during atherosclerosis. Our study emphasizes the complex heterogeneous phenotypes of EC at disease-prone sites. We previously reported that atheroprone sites contain $\approx 1\%$ senescent cells, which cannot participate in vascular repair because they are irreversibly growth arrested and exhibit defective migration.³² Our current study indicates that atheroprone sites also contain TWIST1-positive EC that exhibit enhanced proliferation. Further studies are required to determine whether specific mechanical signatures are required for the induction of EC proliferation or senescence and the interplay between these 2 populations during atherogenesis.

Our study revealed that the function of TWIST is, at least in part, conserved in developing vasculature and atheroprone regions of adult arteries because it enhances EC proliferation in both contexts. However, the consequence of TWIST activation differs between embryogenic and adult arteries because EC proliferation has an essential role in vascular development but promotes lesion initiation in mature arteries by enhancing permeability to lipoproteins. Thus, the induction of TWIST by low shear stress may be an example of antagonistic pleiotropy because it has beneficial effects during development but contributes to arterial disease in adults. Although EC in developing vasculature are phenotypically distinct from those in adult vessels, our study reveals that some aspects of EC physiology, including TWIST induction of proliferation, are conserved between embryonic and adult tissues. This is consistent with the demonstration that several other genes (eg, bone morphogenic proteins⁴¹ and Notch⁴⁷) and processes (eg, EndoMT^{45,46}) with a well-recognized role in development also contribute to focal EC dysfunction and atherosclerosis. We suggest therefore that the behavior of EC at atheroprone sites is a reflection of an early developmental stage and that developmental genes including GATA4 and TWIST1 could be novel therapeutic targets in atherosclerosis.

Sources of Funding

This study was funded by the British Heart Foundation (RG/13/1/30042).

Disclosures

None.

References

- Kwak BR, Bäck M, Bochaton-Piallat ML, et al. Biomechanical factors in atherosclerosis: mechanisms and clinical implications. *Eur Heart J*. 2014;35:3013–3020, 3020a. doi: 10.1093/eurheartj/ehu353.
- Suo J, Ferrara DE, Sorescu D, Gulberg RE, Taylor WR, Giddens DP. Hemodynamic shear stresses in mouse aortas: implications for atherogenesis. *Arterioscler Thromb Vasc Biol*. 2007;27:346–351. doi: 10.1161/01.ATV.0000253492.45717.46.
- Dai G, Kaazempur-Mofrad MR, Natarajan S, Zhang Y, Vaughn S, Blackman BR, Kamm RD, García-Cardeña G, Gimbrone MA Jr. Distinct endothelial phenotypes evoked by arterial waveforms derived from atherosclerosis-susceptible and -resistant regions of human vasculature. *Proc Natl Acad Sci USA*. 2004;101:14871–14876. doi: 10.1073/pnas.0406073101.
- Passerini AG, Polacek DC, Shi C, Francesco NM, Manduchi E, Grant GR, Pritchard WF, Powell S, Chang GY, Stoeckert CJ Jr, Davies PF. Coexisting proinflammatory and antioxidative endothelial transcription profiles in a disturbed flow region of the adult porcine aorta. *Proc Natl Acad Sci USA*. 2004;101:2482–2487.
- Hajra L, Evans AI, Chen M, Hyduk SJ, Collins T, Cybulsky MI. The NF- κ B signal transduction pathway in aortic endothelial cells is primed for activation in regions predisposed to atherosclerotic lesion formation. *Proc Natl Acad Sci USA*. 2000;97:9052–9057.
- Cuhlmann S, Van der Heiden K, Saliba D, Tremoleda JL, Khalil M, Zakkar M, Chaudhury H, Luong le A, Mason JC, Udalova I, Gsell W, Jones H, Haskard DO, Krams R, Evans PC. Disturbed blood flow induces RelA expression via c-Jun N-terminal kinase 1: a novel mode of NF- κ B regulation that promotes arterial inflammation. *Circ Res*. 2011;108:950–959. doi: 10.1161/CIRCRESAHA.110.233841.
- White SJ, Hayes EM, Lehoux S, Jeremy JY, Horrevoets AJ, Newby AC. Characterization of the differential response of endothelial cells exposed to normal and elevated laminar shear stress. *J Cell Physiol*. 2011;226:2841–2848. doi: 10.1002/jcp.22629.
- Yamawaki H, Lehoux S, Berk BC. Chronic physiological shear stress inhibits tumor necrosis factor-induced proinflammatory responses in rabbit aorta perfused ex vivo. *Circulation*. 2003;108:1619–1625. doi: 10.1161/01.CIR.0000089373.49941.C4.
- Lin K, Hsu PP, Chen BP, Yuan S, Usami S, Shyy JY, Li YS, Chien S. Molecular mechanism of endothelial growth arrest by laminar shear stress. *Proc Natl Acad Sci USA*. 2000;97:9385–9389. doi: 10.1073/pnas.170282597.
- Guo D, Chien S, Shyy JY. Regulation of endothelial cell cycle by laminar versus oscillatory flow: distinct modes of interactions of AMP-activated protein kinase and Akt pathways. *Circ Res*. 2007;100:564–571. doi: 10.1161/01.RES.0000259561.23876.c5.
- Schober A, Nazari-Jahantigh M, Wei Y, Bidzhekov K, Gremse F, Grommes J, Megens RT, Heyll K, Noels H, Hristov M, Wang S, Kiessling F, Olson EN, Weber C. MicroRNA-126-5p promotes endothelial proliferation and limits atherosclerosis by suppressing Dlk1. *Nat Med*. 2014;20:368–376. doi: 10.1038/nm.3487.
- Cancel LM, Tarbell JM. The role of mitosis in LDL transport through cultured endothelial cell monolayers. *Am J Physiol Heart Circ Physiol*. 2011;300:H769–H776. doi: 10.1152/ajpheart.00445.2010.
- Dekker RJ, van Soest S, Fontijn RD, Salamanca S, de Groot PG, VanBavel E, Pannekoek H, Horrevoets AJ. Prolonged fluid shear stress induces a distinct set of endothelial cell genes, most specifically lung Krüppel-like factor (KLF2). *Blood*. 2002;100:1689–1698. doi: 10.1182/blood-2002-01-0046.
- Senbanerjee S, Lin ZY, Atkins GB, Greif DM, Rao RM, Feinberg MW, Chen ZP, Simon DI, Lusinskas FW, Michel TM, Garcia-Cardena G, Gimbrone MA, Jain MK. KLF2 as a novel transcriptional regulator of endothelial cell function. *Circulation*. 2003;108:169–169.
- Ni CW, Qiu H, Rezvan A, Kwon K, Nam D, Son DJ, Visvader JE, Jo H. Discovery of novel mechanosensitive genes in vivo using mouse carotid artery endothelium exposed to disturbed flow. *Blood*. 2010;116:e66–e73. doi: 10.1182/blood-2010-04-278192.
- Dunn J, Qiu H, Kim S, Jjingo D, Hoffman R, Kim CW, Jang I, Son DJ, Kim D, Pan C, Fan Y, Jordan IK, Jo H. Flow-dependent epigenetic DNA methylation regulates endothelial gene expression and atherosclerosis. *J Clin Invest*. 2014;124:3187–3199. doi: 10.1172/JCI74792.
- Zeng L, Zampetaki A, Margariti A, Pepe AE, Alam S, Martin D, Xiao Q, Wang W, Jin ZG, Cockerill G, Mori K, Li YS, Hu Y, Chien S, Xu Q. Sustained activation of XBP1 splicing leads to endothelial apoptosis and atherosclerosis development in response to disturbed flow. *Proc Natl Acad Sci USA*. 2009;106:8326–8331. doi: 10.1073/pnas.0903197106.
- Boon RA, Hergenreider E, Dimmeler S. Atheroprotective mechanisms of shear stress-regulated microRNAs. *Thromb Haemost*. 2012;108:616–620. doi: 10.1160/TH12-07-0491.
- Thisse B, Stoetzel C, Gorostiza-Thisse C, Perrin-Schmitt F. Sequence of the twist gene and nuclear localization of its protein in endomesodermal cells of early Drosophila embryos. *EMBO J*. 1988;7:2175–2183.
- Gitelman I. Twist protein in mouse embryogenesis. *Dev Biol*. 1997;189:205–214. doi: 10.1006/dbio.1997.8614.
- Chakraborty S, Wrigg EE, Hinton RB, Merrill WH, Spicer DB, Yutzy KE. Twist1 promotes heart valve cell proliferation and extracellular matrix gene expression during development in vivo and is expressed in human diseased aortic valves. *Dev Biol*. 2010;347:167–179. doi: 10.1016/j.ydbio.2010.08.021.
- Lovisa S, LeBleu VS, Tampe B, Sugimoto H, Vadrnaga K, Carstens JL, Wu CC, Hagos Y, Burckhardt BC, Pentcheva-Hoang T, Nischal H, Allison JP, Zeisberg M, Kalluri R. Epithelial-to-mesenchymal transition induces cell cycle arrest and parenchymal damage in renal fibrosis. *Nat Med*. 2015;21:998–1009. doi: 10.1038/nm.3902.
- Yang J, Mani SA, Donaher JL, Ramaswamy S, Itzykson RA, Come C, Savagner P, Gitelman I, Richardson A, Weinberg RA. Twist, a master regulator of morphogenesis, plays an essential role in tumor metastasis. *Cell*. 2004;117:927–939. doi: 10.1016/j.cell.2004.06.006.
- Kuo CT, Morrissey EE, Anandappa R, Sigrist K, Lu MM, Parmacek MS, Soudais C, Leiden JM. GATA4 transcription factor is required for ventral morphogenesis and heart tube formation. *Genes Dev*. 1997;11:1048–1060.
- Watt AJ, Battle MA, Li J, Duncan SA. GATA4 is essential for formation of the proepicardium and regulates cardiogenesis. *Proc Natl Acad Sci USA*. 2004;101:12573–12578. doi: 10.1073/pnas.0400752101.
- Lim J, Thiery JP. Epithelial-mesenchymal transitions: insights from development. *Development*. 2012;139:3471–3486. doi: 10.1242/dev.071209.
- Madisen L, Zwingman TA, Sunkin SM, Oh SW, Zariwala HA, Gu H, Ng LL, Palmiter RD, Hawrylycz MJ, Jones AR, Lein ES, Zeng H. A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. *Nat Neurosci*. 2010;13:133–140. doi: 10.1038/nn.2467.
- Göthert JR, Gustin SE, van Eekelen JA, Schmidt U, Hall MA, Jane SM, Green AR, Götgens B, Izon DJ, Begley CG. Genetically tagging endothelial cells in vivo: bone marrow-derived cells do not contribute to tumor endothelium. *Blood*. 2004;104:1769–1777. doi: 10.1182/blood-2003-11-3952.
- Li J, Liu CH, Sun Y, Gong Y, Fu Z, Evans LP, Tian KT, Juan AM, Hurst CG, Mammoto A, Chen J. Endothelial TWIST1 promotes pathological ocular angiogenesis. *Invest Ophthalmol Vis Sci*. 2014;55:8267–8277. doi: 10.1167/iovs.14-15623.
- Cheng C, Tempel D, van Haperen R, van der Baan A, Grosveld F, Daemen MJ, Krams R, de Crom R. Atherosclerotic lesion size and vulnerability are determined by patterns of fluid shear stress. *Circulation*. 2006;113:2744–2753. doi: 10.1161/CIRCULATIONAHA.105.590018.
- Björklund MM, Hollensen AK, Hagensen MK, Dagnaes-Hansen F, Christoffersen C, Mikkelsen JG, Bentzen JF. Induction of atherosclerosis in mice and hamsters without germline genetic engineering. *Circ Res*. 2014;114:1684–1689. doi: 10.1161/CIRCRESAHA.114.302937.
- Warboys CM, de Luca A, Amini N, et al. Disturbed flow promotes endothelial senescence via a p53-dependent pathway. *Arterioscler Thromb Vasc Biol*. 2014;34:985–995. doi: 10.1161/ATVBAHA.114.303415.
- Dardik A, Chen L, Frattini J, Asada H, Aziz F, Kudo FA, Sumpio BE. Differential effects of orbital and laminar shear stress on endothelial cells. *J Vasc Surg*. 2005;41:869–880. doi: 10.1016/j.jvs.2005.01.020.
- Warboys CM, Eric Berson R, Mann GE, Pearson JD, Weinberg PD. Acute and chronic exposure to shear stress have opposite effects on endothelial permeability to macromolecules. *Am J Physiol Heart Circ Physiol*. 2010;298:H1850–H1856. doi: 10.1152/ajpheart.00114.2010.
- Oxtoby E, Jowett T. Cloning of the zebrafish krox-20 gene (krox-20) and its expression during hindbrain development. *Nucleic Acids Res*. 1993;21:1087–1095.
- Das A, Crump JG. Brms and id2a act upstream of Twist1 to restrict ectomesenchyme potential of the cranial neural crest. *PLoS Genet*. 2012;8:e1002710. doi: 10.1371/journal.pgen.1002710.
- Tapanes-Castillo A, Baylies MK. Notch signaling patterns Drosophila mesodermal segments by regulating the bHLH transcription factor twist. *Development*. 2004;131:2359–2372.

38. Tian Y, Xu Y, Fu Q, Chang M, Wang Y, Shang X, Wan C, Marymont JV, Dong Y. Notch inhibits chondrogenic differentiation of mesenchymal progenitor cells by targeting Twist1. *Mol Cell Endocrinol*. 2015;403:30–38. doi: 10.1016/j.mce.2015.01.015.
39. Desprat N, Supatto W, Pouille PA, Beaupaire E, Farge E. Tissue deformation modulates twist expression to determine anterior midgut differentiation in *Drosophila* embryos. *Dev Cell*. 2008;15:470–477. doi: 10.1016/j.devcel.2008.07.009.
40. Wei SC, Fattet L, Tsai JH, Guo Y, Pai VH, Majeski HE, Chen AC, Sah RL, Taylor SS, Engler AJ, Yang J. Matrix stiffness drives epithelial-mesenchymal transition and tumour metastasis through a TWIST1-G3BP2 mechanotransduction pathway. *Nat Cell Biol*. 2015;17:678–688. doi: 10.1038/ncb3157.
41. Sorescu GP, Song H, Tressell SL, Hwang J, Dikalov S, Smith DA, Boyd NL, Platt MO, Lassègue B, Griendling KK, Jo H. Bone morphogenic protein 4 produced in endothelial cells by oscillatory shear stress induces monocyte adhesion by stimulating reactive oxygen species production from a nox1-based NADPH oxidase. *Circ Res*. 2004;95:773–779. doi: 10.1161/01.RES.0000145728.22878.45.
42. Hayashi M, Nimura K, Kashiwagi K, Harada T, Takaoka K, Kato H, Tamai K, Kaneda Y. Comparative roles of Twist-1 and Id1 in transcriptional regulation by BMP signaling. *J Cell Sci*. 2007;120:1350–1357. doi: 10.1242/jcs.000067.
43. Chaudhury H, Zakkar M, Boyle J, Cuhlmann S, van der Heiden K, Luong le A, Davis J, Platt A, Mason JC, Krams R, Haskard DO, Clark AR, Evans PC. c-Jun N-terminal kinase primes endothelial cells at atheroprone sites for apoptosis. *Arterioscler Thromb Vasc Biol*. 2010;30:546–553. doi: 10.1161/ATVBAHA.109.201368.
44. Maddaluno L, Rudini N, Cuttano R, et al. EndMT contributes to the onset and progression of cerebral cavernous malformations. *Nature*. 2013;498:492–496. doi: 10.1038/nature12207.
45. Chen PY, Qin L, Baeyens N, Li G, Afolabi T, Budatha M, Tellides G, Schwartz MA, Simons M. Endothelial-to-mesenchymal transition drives atherosclerosis progression. *J Clin Invest*. 2015;125:4514–4528. doi: 10.1172/JCI82719.
46. Moonen JR, Lee ES, Schmidt M, Maleszewska M, Koerts JA, Brouwer LA, van Kooten TG, van Luyn MJ, Zeebregts CJ, Krenning G, Harmsen MC. Endothelial-to-mesenchymal transition contributes to fibro-proliferative vascular disease and is modulated by fluid shear stress. *Cardiovasc Res*. 2015;108:377–386. doi: 10.1093/cvr/cvv175.
47. Lehoux S, Jones EA. Shear stress, arterial identity and atherosclerosis. *Thromb Haemost*. 2016;115:467–473. doi: 10.1160/TH15-10-0791.

Novelty and Significance

What Is Known?

- Atherosclerosis develops at branches and bends of arteries exposed to disturbed patterns of blood flow.
- Endothelial cells at atheroprone sites display enhanced proliferation, permeability, inflammation, and expression of mesenchymal genes.

What New Information Does This Article Contribute?

- Disturbed flow induced endothelial expression of the transcription factor TWIST1 at atheroprone sites of arteries.
- TWIST1 promoted atherosclerosis by enhancing inflammation, endothelial cell proliferation, and vascular permeability and by inducing endothelial-to-mesenchymal transition.
- TWIST1 also promoted developmental angiogenesis by enhancing endothelial proliferation and migration.

Atherosclerosis is a focal disease of arteries that develops at branches and bends exposed to disturbed patterns of blood flow. Endothelial cells at atheroprone sites display several features that have been linked to lesion formation including excessive

rates of proliferation, enhanced permeability, inflammation, and expression of mesenchymal genes. The molecular mechanisms that induce endothelial dysfunction at sites of disturbed flow are incompletely characterized. This article reveals for the first time that disturbed flow regulates endothelial function by activating the developmental transcription factor TWIST1 at atheroprone sites of arteries. TWIST1 expression contributed to atherosclerosis by enhancing vascular inflammation and by driving endothelial cell proliferation and permeability. TWIST1 also promoted endothelial-to-mesenchymal transition in response to disturbed flow. In addition to its role in atherosclerosis, TWIST1 was also expressed in embryos and contributed to angiogenesis by enhancing endothelial proliferation and migration. Thus, TWIST1 is a coordinator of vascular development and also contributes to the initiation of focal atherosclerosis in adult arteries. We suggest therefore that some mechanosensitive pathways operate both at atheroprone sites of adult arteries and in developing vessels and that developmental genes including TWIST1 could be novel therapeutic targets in atherosclerosis.

TWIST1 Integrates Endothelial Responses to Flow in Vascular Dysfunction and Atherosclerosis

Marwa M. Mahmoud, Hyejeong Rosemary Kim, Rouyu Xing, Sarah Hsiao, Akiko Mammoto, Jing Chen, Jovana Serbanovic-Canic, Shuang Feng, Neil P. Bowden, Richard Maguire, Markus Ariaans, Sheila E. Francis, Peter D. Weinberg, Kim van der Heiden, Elizabeth A. Jones, Timothy J.A. Chico, Victoria Ridger and Paul C. Evans

Circ Res. 2016;119:450-462; originally published online May 31, 2016;
doi: 10.1161/CIRCRESAHA.116.308870

Circulation Research is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231
Copyright © 2016 American Heart Association, Inc. All rights reserved.
Print ISSN: 0009-7330. Online ISSN: 1524-4571

The online version of this article, along with updated information and services, is located on the
World Wide Web at:

<http://circres.ahajournals.org/content/119/3/450>

Free via Open Access

Data Supplement (unedited) at:

<http://circres.ahajournals.org/content/suppl/2016/05/31/CIRCRESAHA.116.308870.DC1>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation Research* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Circulation Research* is online at:
<http://circres.ahajournals.org/subscriptions/>

EXPANDED METHODS

Antibodies and reagents. Antibodies targeting human and murine TWIST1 (ABD29, Millipore), GATA4 (PA1-102, ThermoFisher) and, Snail (ab180714, Abcam), PCNA (ab 18197, Abcam), CD31 (102514, Biolegend), Ki67 (ab15580, Abcam), Activated-Notch 1 (ab8925) were obtained commercially. AlexaFluor-conjugated secondary antibodies, TO-PRO-3 and aqueous mounting media (ProLong® Gold Antifade Mountant) were from Invitrogen. All other reagents were from Sigma-Aldrich unless otherwise stated.

Mouse lines. Mice were housed under specific-pathogen free conditions and studied according to UK Home Office Regulations and the UK Animals (Scientific Procedures) Act 1986. Mice were weaned at 4 weeks of age and maintained on a normal chow diet. Mice between 2 and 3 months of age were used for experimentation in groups of at least 4 animals. Littermates were used as controls. TWIST1 and GATA4 mice were not back crossed onto a C57BL/6 background for 6 generations. For cell tracing studies, transgenic Rosa26-tdTomato¹ mice were crossed with endothelial-SCL-Cre-ER^T mice containing a tamoxifen-inducible EC-specific Cre². Mice with Tie2-specific deletion of TWIST1 (*TWIST1^{ckO}*) were generated by crossing Tie2-Cre expressing mice (Jackson Laboratory stock #004128) with TWIST1 floxed mice (*TWIST1^{flox/flox}*)³. Mice with EC deletion of GATA4 (*GATA4^{ckO}*) were generated by crossing endothelial-SCL-Cre-ER^T mice with GATA4 floxed mice (*GATA4^{flox/flox}*). To activate Cre, tamoxifen was administered for 5 consecutive days (160 mg/kg). Constrictive cuffs with an internal diameter of 400 µm tapering to 200 µm over 1.5 mm were manufactured from PTFE and applied to the right carotid artery of isoflurane-anaesthetized mice following published methods^{4,5}.

En face staining of murine endothelium. The expression levels of specific proteins were assessed in EC at regions of the inner curvature (susceptible site) and outer curvature (protected site) of murine aortae and in carotid arteries by *en face* staining. Animals were killed by I.P injection of pentobarbital or by isoflurane overdose. Aortae were perfused *in situ* with PBS (at a pressure of approximately 100 mm Hg) and then perfusion-fixed with 4% Paraformaldehyde prior to harvesting. Fixed aortae were tested by immunostaining using specific primary antibodies and Alexafluor568-conjugated secondary antibodies (red). EC were identified by co-staining using anti-CD31 antibodies conjugated to the fluorophore FITC (green). Nuclei were identified using a DNA-binding probe with far-red emission (To-Pro-3). Stained vessels were mounted prior to visualization of endothelial surfaces *en face* using confocal laser-scanning microscopy (Zeiss LSM510 NLO inverted microscope). Isotype-matched monoclonal antibodies raised against irrelevant antigens or pre-immune rabbit sera were used as experimental controls for specific staining. The expression of particular proteins at each site was assessed by quantification of fluorescence intensity for multiple cells (at least 50 per site) using Image J (1.49p) and calculation of mean fluorescence intensities with standard error of the mean.

Lesion analysis.

Hypercholesterolemia was induced in mice by a single Intraperitoneal injection of adenoassociated virus containing PCSK9 cDNA (rAAV8-D377Y-mPCSK9; 4.9 x10¹¹ pfu) and subsequent exposure to a high fat Western diet (Western RD (P) Diet, 829100; SDS) for 6 weeks⁶. The diet consisted of (w/w) Milk Fat Anhydrous (20%), cholesterol (0.15%), sucrose (33.94%), cornstarch (5%), corn oil (1%), cellulose (5%), casein (19.5%), choline chloride (0.2%), L-Cystine (0.3%), Calcium Carbonate (0.4%), Choline Bitartrate (0.2%), Antioxidant (0.01%), AIN-76A-MX (3.5%), AIN-76A-VX (1%) total fat content (21.4%). Analysis for lipoprotein profiles

and serum total cholesterol and triglycerides, and *en face* lipid staining was carried out using Oil Red O as described⁷. After termination, perfusion fixation was performed using 10% Formalin at a pressure of 110 cm water for 5 min via a cannula inserted into the left ventricle. Each aorta was isolated and adventitial fat was removed before staining using Oil Red O. After cutting along the outer curvature, the aorta was mounted and imaged. Lesion coverage in aortae was analyzed as previously described (Steiner et al., 2014). Lesion coverage was calculated as a percentage of total aortic area.

Isolation of EC from porcine aortae. Pig aortas from 4-6 month old animals (weight approximately 80kg) were obtained immediately after slaughter from a local abattoir. They were cut longitudinally along the outer curvature to expose the lumen. EC exposed to high (outer curvature) or low (inner curvature) WSS were harvested using collagenase (1 mg/ml for 10 minutes at room temperature) prior to gentle scraping. RNA was extracted using an RNeasy MiniKit (Qiagen) and concentrated using an RNeasy MinElute Cleanup kit (Qiagen) and the purity and integrity of total RNA samples was assessed using a Bioanalyser (Agilent).

EC culture and exposure to WSS. HUVEC and PAEC were isolated using collagenase digestion. EC were cultured on 1% gelatin and maintained in M199 growth medium (HUVEC) or DMEM (PAEC) supplemented with foetal bovine serum (20%), L-glutamine (4 mmol/L), endothelial cell growth supplement (30 µg/ml), penicillin (100 U/ml), streptomycin (100 µg/ml) and heparin (10 IU/ml). EC at passage 3-5 were cultured until confluent in 6 well plates and exposed to flow using an orbital shaking platform (PSU-10i; Grant Instruments) housed inside a cell culture incubator. The radius of orbit of the orbital shaker was 10mm and the rotation rate was set to 210 rpm. This motion caused swirling of the culture medium over the cell surface generating low tangential WSS at the centre and high uniform WSS at the periphery. Alternatively, HUVEC were cultured on Ibidi® gelatin-coated µ-Slides (Ibidi GmbH) until they reached confluency. Flowing medium was then applied using the Ibidi® pump system to generate low (4 dyn/cm²), low oscillatory (+/- 4 dyn/cm², 0.5 Hz) or high (13 dyn/cm²) WSS. The slides and pump apparatus were enclosed in a cell culture incubator warmed to 37°C.

RNA interference. Cell cultures were transfected with siRNA sequences that are known to silence TWIST1 (Silencer® Select S14523, Ambion® and L-006434-00-0005 ON-TARGETplus SMARTpool, Dharmacon) or GATA4 (Silencer® Select s5603, Ambion® and L-008244-00-0005 ON-TARGETplus SMARTpool, Dharmacon) using the Lipofectamine® RNAiMAX transfection system (13778-150, Invitrogen) following the manufacturer's instructions. Two different siRNAs were used to test each gene to ensure specificity. Final siRNA concentration used was 25nM⁸. After knockdown, cells were then incubated in complete M199 growth medium for 2 h before exposure to flow. Non-targeting scrambled sequences were used as a control (D-001810-01-50 ON-TARGETplus Non targeting siRNA#1, Dharmacon).

Comparative real time PCR. RNA was extracted using the RNeasy Mini Kit (74104, Qiagen) and reverse transcribed into cDNA using the iScript cDNA synthesis kit (1708891, Bio-Rad). The levels of human, porcine or zebrafish transcripts were assessed using quantitative real time PCR (qRT-PCR) using gene-specific primers (Supplementary Table 1). Reactions were prepared using SsoAdvanced universal SYBR®Green supermix (172-5271, Bio-rad) and following the manufacturer's instructions, and were performed in triplicate. Relative gene expression was calculated by comparing the number of thermal cycles that were necessary to generate threshold amounts of product. Fold changes were calculated using the

$\Delta\Delta C_t$ method. Data were pooled from at least three independent experiments and mean values were calculated with SEM.

Western blotting Total cell lysates were isolated using lysis buffer (containing 2% SDS, 10% Glycerol and 5% β -mercaptoethanol). Western blotting was carried out using specific antibodies against GATA4 (Sc 1237 (C-20), Santa Cruz), TWIST1 (ABD29, Millipore) and Calnexin (610524, BD Transduction Laboratories) horse radish peroxidase-conjugated secondary antibodies obtained commercially from Dako and chemiluminescent detection was carried out using ECL Prime[®] (GE Healthcare). Membranes were imaged using the C-DiGit scanner (LI-COR Biosciences).

Immunofluorescent staining of cultured EC. The expression levels of proteins were assessed by immunostaining using specific antibodies followed by widefield fluorescence microscopy (LeicaDMI4000B). HUVEC were fixed with Paraformaldehyde (4%) and permeabilised with Triton X-100 (0.1%). Following blocking with goat serum for 30 min monolayers were incubated for 16 h with primary antibodies against TWIST1, GATA4, Snail, or PCNA and AlexaFluor488- or Alexafluor568-conjugated secondary antibodies. Nuclei were identified using the DNA-binding probe DAPI (Sigma). Image analysis was performed using Image J software (1.49p) to calculate average fluorescence. Isotype controls or omission of the primary antibody was used to control for non-specific staining.

Promoter analysis and chromatin immunoprecipitation (ChIP). Genomic sequences (www.epd.vital-it.ch) were interrogated for putative GATA4 binding sites (A/T) GATAA(G) using TFSEARCH data base (<http://www.cbrc.jp/research/db/TFSEARCH.html>). For ChIP, HUVECs were fixed using formaldehyde (1% for 10 min). ChIP was carried out using the SimpleChIP[®]Plus Enzymatic Chromatin IP Kit (#9005; Cell signalling) following the manufacturer's instructions. ChIP-grade GATA4 antibodies (C-20 sc-1237, Santa Cruz) or isotype-matched IgG control were used. Precipitation of specific genomic DNA fragments was assessed by qRT-PCR using primers that amplify regions of the TWIST1 promoter.

Assay of permeability. The permeability of EC monolayers exposed to flow was determined using rhodamine-labelled albumin⁹. HUVEC were cultured in transwell inserts overnight and then exposed to orbital shaking for 72 h. The media in the upper compartment was then replaced with 10% serum-supplemented DMEM containing 1% BPA and rhodamine-labeled albumin (1 mg/ml). Media in the lower compartment was sampled at 1 h and fluorescence was measured using a fluorimeter (Varioskan, Thermoscientific) with excitation at 570 nm and emission at 600 nm. Data were pooled from five experiments and mean concentrations of rhodamine albumin +/- SEM were calculated.

Zebrafish lines. The experiments were performed in accordance with the UK Animal (Scientific Procedures) Act, 1986 and the NIH Guide for the Care and Use of Laboratory Animals. Adult zebrafish were maintained on a 14 h light / 10 h dark cycle at 28°C in UK Home Office approved facilities in the Bateson Centre at the University of Sheffield. Studies were carried out on wild-type zebrafish or transgenic zebrafish lines, *Tg(fli1:EGFP)* (endothelial EGFP), *Tg(kdr1:NLS-EGFP)* (endothelial nuclear EGFP) or *Tg(gata1:dsRed)* (red blood cell expression of dsRed). To generate mutants, a Cas9 expression plasmid (Addgene) was linearised with Apal and mRNA for injection was generated with mMACHINE SP6 Kit (Life Technologies). For gRNA synthesis, a *twist1b* CRISPR target site was selected with

the sequence 5'-GG-TTCCGATAGCTCGACCCT-NGG-3' which contains a BsaJI restriction enzyme site used for genotyping. An ultamer of the sequence (AAAGCACCGACTCGGTGCCACTTTTTCAAGTTGATAACGGACTAGCCTTATTTT AACTTGCTATTTCTAGCTCTAAAACAGGGTCGAGCTATCGGAATCCTATAGTGAG TCGTATTACGC) for gRNA was *in vitro* transcribed with the MEGAshortscript T7 kit (Life technologies). This was injected at a concentration of 2.4 µg/µl along with 0.5 µg/µl Cas9 mRNA into newly fertilized embryos. Injected embryos were raised to maturity (F0) and their progenies were genotyped for *twist1b* mutation. The embryos harbouring mutation were raised into adults (F1) and genotyped by PCR analysis of DNA from fin clips.

Gain- and loss-of-function studies in zebrafish. *Twist1a* translation blocking morpholino (5'-ACCTCTGGAAAAGCTCAGATTGCGG-3') was purchased from Gene Tools LLC (Philomath, OR, USA) and injected at a concentration of 0.5mM and in a volume of 1nl into one cell stage embryos. To generate *twist1b* mRNA, the *twist1b* coding sequence was cloned into pCS2+ expression vector. The plasmid was linearized with NotI and transcribed with SP6 polymerase. The SP6 mMessage Machine kit (Ambion, Texas, USA) was used for *in vitro* transcription. Transgenic or mutant embryos were injected at the 1-cell stage with MO or capped messenger RNA using a Narishige IM-300 micromanipulator. Live embryos were analysed embedded in 0.7% low melting agarose using spinning disk confocal microscope (Perkin Elmer) and Leica stereoscope (M165FC)

In situ hybridisation and immunofluorescence staining of zebrafish embryos. The whole mount *in situ* hybridization of zebrafish embryos was performed using digoxigenin (DIG) labelled *twist1a*, *twist1b*, *twist2* probes synthesized using a DIG labelling mix (Roche) and T7 or SP6 polymerase (Roche). Imaging was performed using an Olympus FV1000 laser scanning confocal microscope.

Statistics. Differences between samples were analysed using an unpaired or paired Student's t-test or ANOVA (*p<0.05, **p<0.01, ***p<0.001).


REFERENCES FOR EXPANDED METHODS

1. Madisen L, Zwingman TA, Sunkin SM, Oh SW, Zariwala HA, Gu H, Ng LL, Palmiter RD, Hawrylycz MJ, Jones AR, Lein ES and Zeng H. A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. *Nature Neurosci.* 2010;13:133-140.
2. Gothert JR, Gustin SE, van Eekelen JA, Schmidt U, Hall MA, Jane SM, Green AR, Gottgens B, Izon DJ and Begley CG. Genetically tagging endothelial cells in vivo: bone marrow-derived cells do not contribute to tumor endothelium. *Blood.* 2004;104:1769-1777.
3. Li J, Liu CH, Sun Y, Gong Y, Fu Z, Evans LP, Tian KT, Juan AM, Hurst CG, Mammoto A and Chen J. Endothelial TWIST1 promotes pathological ocular angiogenesis. *Investig Ophthalmol Vis Sci.* 2014;55:8267-8277.
4. Cheng C, Tempel D, van Haperen R, van der Baan A, Grosveld F, Daemen MJ, Krams R and de Crom R. Atherosclerotic lesion size and vulnerability are determined by patterns of fluid shear stress. *Circulation.* 2006;113:2744-2753.
5. Cuhlmann S, Van der Heiden K, Saliba D, Tremoleda JL, Khalil M, Zakkar M, Chaudhury H, Le AL, Mason JC, Udalova I, Gsell W, Jones H, Haskard DO, Krams R and Evans PC. Disturbed Blood Flow Induces RelA Expression via c-Jun N-Terminal Kinase 1 A Novel Mode of NF-kappa B Regulation That Promotes Arterial Inflammation. *Circ Res.* 2011;108:950-959.
6. Bjorklund MM, Hollensen AK, Hagensen MK, Dagnaes-Hansen F, Christoffersen C, Mikkelsen JG and Bentzon JF. Induction of Atherosclerosis in Mice and Hamsters Without Germline Genetic Engineering. *Circ Res.* 2014;114:1684-1689.
7. Steiner T, Francescut L, Byrne S, Hughes T, Jayanthi A, Guschina I, Harwood J, Cianflone K, Stover C and Francis S. Protective Role for Properdin in Progression of Experimental Murine Atherosclerosis. *Plos One.* 2014;9.
8. Zografou S, Basagiannis D, Papafotika A, Shirakawa R, Horiuchi H, Auerbach D, Fukuda M and Christoforidis S. A complete Rab screening reveals novel insights in Weibel-Palade body exocytosis. *J Cell Sci.* 2012;125:4780-4790.
9. Warboys CM, Berson R, Mann GE, Pearson JD and Weinberg PD. Acute and chronic exposure to shear stress have opposite effects on endothelial permeability to macromolecules. *Am J Physiol-Heart Circ Physiol.* 2010;298:H1850-H1856.

Supplementary Table PCR primer sequences


Gene	Forward primer	Reverse primer
GATA4 (human)	TCCCAGACGTTCTCAGTCAG	GGAGCTGGTCTGTGGAGACT
TWIST1 (human)	CGGACAAGCTGAGCAAGAT	CTGGAGGACCTGGTAGAGGA
TWIST1 promoter (human)	AGCAATCCCAAATCGGCCC	TGGCAACAGCTTCTACACAGT
SLUG (human)	CTGGCCAAACACAAGCAG	ACCCAGGCTCACATATTCCT
N-cadherin (human)	GCACAGATGTGGACAGGATT	CAGCACAAGGATAAGCAGGA
α -SMA (human)	TTTCAGCTTCCCTGAACACCA	GGGCAACACGAAGCTCATTG
Snail (human)	TGCAGGACTCTAATCCAGAGTTT	GACAGAGTCCCAGATGAGCA
Cyclin D1 (human)	TGA CCC CGC ACG ATT TCA TT	TGA GGC GGT AGT AGG ACA GG
Cyclin G2 (human)	TTG GAC AGG TTC TTG GCT CT	TCA ACT ATT CTA GCA GCC AGC
Cdk4 (human)	CAA TGT TGT CCG GCT GAT GG	GCT CAA ACA CCA GGG TTA CC
VCAM-1 (human)	CATTGACTTGCAGCACCACA	AGATGTGGTCCCCTCATTCG
ICAM-1 (human)	CACAAGCCACGCCTCCCTGAAC CTA	TGTGGGCCTTTGTGTTTTGAT GCTA
CD31 (human)	GGTGTGGTGGAAGGAGTG	GGGACAGAACAGTTGACCCT
Piezo1 (human)	ATCGCCATCATCTGGTTCCC	TGGTGAACAGCGGCTCATAG
VEGFR2 (human)	CTCGGGTCCATTTCAAATCT	GCTGTCCCAGGAAATTCTG T
HPRT (human)	TTGGTCAGGCAGTATAATCC	GGGCATATCCTACAACAAC
GATA4 (porcine)	GGAAGGCAGAGAGTGTGTCA	GCTGATGCCATTTCATCTTGT
TWIST1 (porcine)	GGGAGTCCGCAGTCCTAC	TGGATCTTGCTCAGCTTGTC
twist1a (zebrafish)	CGCGTTTTCTGTGTGGAGAA	CCGAGAATCATGCTGCATCA
twist1b (zebrafish)	CGCTTGGTCCATGTCTGC	CTCGTGTTTTCCCAGCTCAC
twist2 (zebrafish)	CGTCCTGATAATGCCGAACG	GCCACGTCACTGAGGTTTAG

SUPPLEMENTAL MATERIAL




Online Figure I Low shear induced TWIST1 and GATA4 in cultured PAEC.

PAEC were exposed to orbital flow to generate low (Centre) or high (Periphery) wall shear stress (WSS). After 72 h, transcript levels were quantified by qRT-PCR and mean values \pm SEM are shown.




Online Figure II Validation of TWIST1 and GATA4 gene silencing.

Silencing of TWIST1 or GATA4 was performed using two different siRNAs for each gene (designated i and ii). HUVEC were treated with siRNA targeting GATA4 or TWIST1, or with scrambled non-targeting siRNA as a control. After knockdown, cells were then cultured in 6 well plates prior to the application of orbital flow for 72 h. (A) Transcript levels of TWIST1 (left) or GATA4 (right) in cells at the centre of the well (low WSS) were quantified by qRT-PCR. Data were pooled from 3 independent experiments and mean levels \pm SEM are shown. (B) Expression of TWIST1 and GATA4 protein at the centre of the well (low WSS) was determined by Western blotting using anti-Calnexin antibodies to control for total protein levels.




Online Figure III Validation of genetic targeting of TWIST1 and GATA4 in endothelial cells of transgenic mice.

(A) *TWIST1*^{flox/flox} mice were crossed with Tie2-Cre transgenics (generating *TWIST1*^{ckO}). (B) *GATA4*^{flox/flox} mice were crossed with endothelial-SCL-Cre-ER^T transgenics followed by 5 days of tamoxifen treatment (generating *GATA4*^{ckO}). (A, B) To validate genetic deletion of *TWIST1* and *GATA4* from EC in the descending aorta *en face* staining was performed using anti-*TWIST1* or anti-*GATA4* antibodies (red). Co-staining was carried out using anti-CD31 antibodies (green) and the nuclear counter stain TOPRO3 (blue). Representative images are shown. Scale bar 10 μ m.




Online Figure IV Low shear stress induced Notch1 in experimental carotid arteries.



Flow-altering, constrictive cuffs were placed on the right carotid arteries of C57BL/6 mice. They generated anatomically distinct regions exposed to low, high and low oscillatory wall shear stress (WSS; as indicated). Right (experimental) and left (sham-operated) carotid arteries were harvested after 14 days and *en face* staining was performed using anti-Notch1 antibodies (red), anti-CD31 antibodies conjugated to FITC (green) and the nuclear counter stain TO-PRO-3 (blue). Representative images and quantitation of Notch1 expression (mean \pm SEM) are shown. Scale bar, 10 μ m. Data were pooled from five-six independent experiments. * $p < 0.05$ using a one-way ANOVA.



Online Figure V Low shear stress induced TWIST1 in fully differentiated EC *in vivo*. Transgenic SCL-Cre-ER^T/R26RtdTomato mice were used to track endothelial cells. The administration of tamoxifen induces nuclear localization of Cre, which subsequently excises a floxed STOP signal from the tdTomato promoter thus inducing tdTomato expression. (A) To validate the system, SCL-Cre-ER^T/R26RtdTomato mice were treated with tamoxifen for 5 consecutive days or with vehicle as a control. After a further 7 days, carotid arteries were harvested and *en face* staining was performed using anti-CD31 antibodies (green), and the nuclear counter stain TOPRO3 (blue). TdTomato was expressed in the majority of EC in mice treated with tamoxifen but not in vehicle-treated mice. (B) SCL-Cre-ER^T/R26RtdTomato mice were treated with tamoxifen for 5 consecutive days. One week later, flow-altering constrictive cuffs were placed on the right carotid arteries to generate regions exposed to low, high and low oscillatory wall shear stress (WSS; as indicated). Right (experimental) and left (sham-operated) carotid arteries were harvested after 14 days and *en face* staining was performed using anti-TWIST1 or anti-GATA4 antibodies (green), and the nuclear counter stain TO-PRO-3 (blue). Representative images and quantitation of TWIST1 or GATA4 expression (mean \pm SEM) are shown. Scale bars, 10 μ m. * p <0.05 using a one-way ANOVA.




Online Figure VI *twist1* regulated sub-intestinal vein sprouting in embryos. (A) *In situ* hybridisation analysis of *twist1a*, *twist1b* or *twist2* expression in embryos at 52 or 75 hpf. Scale bar, 100 μ m. SIV region is indicated using a broken line. Data shown are representative of the majority of embryos analysed (proportion indicated lower right in each panel). (B) Embryos (*Tg(fli1:EGFP);gata1:dsRed*) were treated with *twist1b* mRNA (to enforce expression) or with mCherry mRNA as a control. They were studied at 75 hpf (flow was established as evidenced by *gata1*-positive red blood cells). Confocal microscopy was used to visualise angiogenic sprouts (arrows). The number of angiogenic sprouts was quantified for multiple embryos and mean values \pm SEM are shown. Representative images are shown. Scale bar, 500 μ m. *** $p < 0.001$ using an unpaired t-test.




Online Figure VII TWIST1 and GATA4 promoted proliferation in EC exposed to low shear stress using a parallel plate system.

HUVEC were treated with two different siRNAs targeting TWIST1 or GATA4 (designated i and ii), or with scrambled non-targeting siRNA. Transfected cells were exposed to low or high WSS for 72 h using a parallel plate system. Cell proliferation was quantified by immunofluorescent staining using anti-PCNA antibodies and co-staining using DAPI. Images are representative of those generated in three independent experiments using one version of the gene-specific siRNA or scrambled control sequences. The % PCNA-positive cells were calculated for multiple fields of view in at least three independent experiments and mean values \pm SEM are shown. Scale bar, 50 μ m.



Online Figure VIII TWIST1 and GATA4 induced inflammatory genes in EC exposed to low shear stress.

(A) HUVEC were cultured in 6 well plates prior to exposure to orbital flow to generate low (Centre) or high (Periphery) wall shear stress for 72 h. The expression of VCAM-1 and ICAM-1 was quantified using qRT-PCR. (B) HUVEC were treated with two different siRNAs targeting TWIST1 or GATA4 (designated i and ii), or with scrambled non-targeting siRNA. Transfected cells were exposed to low WSS (centre). The expression of VCAM-1 and ICAM-1 was quantified using qRT-PCR. The expression level in cells transfected with gene-targeting siRNA is presented relative to the expression in cells transfected with scrambled control siRNA (normalised to 1; dotted line). Data were pooled from three independent experiments and mean values \pm SEM are shown. * $p < 0.05$, *** $p < 0.001$ using an unpaired t-test. (C) The influence of TWIST1 on ICAM-1 expression in EC at low (susceptible) or high (protected) WSS regions of the aorta was studied by *en face* staining using *TWIST1^{cKO}* or *TWIST1^{flox/flox}* mice. Staining was performed using anti-ICAM-1 antibodies (red), anti-CD31 antibodies (green) and the nuclear counter stain TO-PRO-3 (blue). Mean fluorescence values were calculated from multiple cells in 5 independent experiments and mean values \pm SEM are shown. Scale bar, 10 μ m.



Online Figure IX TWIST1 and GATA4 did not regulate the expression of mechanoreceptors in EC exposed to shear stress.

(A) HUVEC were cultured in 6 well plates prior to exposure to orbital flow to generate low (Centre) or high (Periphery) wall shear stress for 72 h. The expression of Piezo, CD31 and VEGFR2 was quantified using qRT-PCR. (B) HUVEC were treated with two different siRNAs targeting TWIST1 or GATA4 (designated i and ii), or with scrambled non-targeting siRNA. Transfected cells were exposed to low WSS (centre). The expression of Piezo, CD31 and VEGFR2 was quantified using qRT-PCR. The expression level in cells transfected with gene-targeting siRNA is presented relative to the expression in cells transfected with scrambled control siRNA (normalised to 1; dotted line). Data were pooled from three independent experiments and mean values \pm SEM are shown. Data were not significant using an unpaired t-test.

Legends for Online Videos

Online Video I: Intersegmental vessel (ISV) sprouting in control embryos

*Online Video II: Enhanced ISV sprouting in *twist1b* overexpressing embryos*

Zebrafish embryos (wild-type, Tg(*fli1*-EGFP)) were treated with *twist1b* mRNA or mCherry mRNA (control) and studied at 24-27 hpf, a time where flow is established in trunk vasculature. The migration during ISV sprouting was promoted in embryos overexpressing *twist1b* (Online Video II) in comparison to mCherry mRNA injected control embryos (Online Video I). The movies were taken for the equal duration of 4 hours.