

This is a repository copy of *Integral solutions to boundary quantum Knizhnik–Zamolodchikov equations*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/123824/>

Version: Accepted Version

Article:

Reshetikhin, Nicolai, Stokman, Jasper and Vlaar, Bart Hendrik Maarten orcid.org/0000-0002-0792-720X (2018) Integral solutions to boundary quantum Knizhnik–Zamolodchikov equations. *Advances in Mathematics*. ISSN: 0001-8708

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

INTEGRAL SOLUTIONS TO BOUNDARY QUANTUM KNIZHNIK-ZAMOLODCHIKOV EQUATIONS

NICOLAI RESHETIKHIN, JASPER STOKMAN, BART VLAAR

ABSTRACT. We construct integral representations of solutions to the boundary quantum Knizhnik-Zamolodchikov equations. These are difference equations taking values in tensor products of Verma modules of quantum affine \mathfrak{sl}_2 , with the K-operators acting diagonally. The integrands in question are products of scalar-valued elliptic weight functions with vector-valued trigonometric weight functions (boundary Bethe vectors). These integrals give rise to a basis of solutions of the boundary qKZ equations over the field of quasi-constant meromorphic functions in weight subspaces of the tensor product.

1. INTRODUCTION

The boundary quantum Knizhnik-Zamolodchikov (qKZ) equations have roots in representation theory, in the theory of solvable models in statistical mechanics and integrable quantum field theory. They first appeared in [1] where the qKZ equation of type A [4] was generalized to other affine Weyl types. What we call here the boundary qKZ equation corresponds to type C. Shortly after the work [1] the boundary qKZ equation appeared in [7] as an equation for correlation functions in the 6-vertex model with reflecting boundary conditions [15]. The boundary qKZ equations also appear as an equation for form-factors in such models and in quantum integrable field theories on the half-line [5]. In [20] the precise relation between the boundary qKZ equations and Sklyanin's commuting transfer matrices [15] was established.

Recall that the boundary qKZ equations related to quantum affine \mathfrak{sl}_2 require the following data (we refer the reader to Section 2 for more details). Fix highest weights $\ell_1, \dots, \ell_N \in \mathbb{C}$ of Verma modules $V^{\ell_1}, \dots, V^{\ell_N}$ of $U_q(\widehat{\mathfrak{sl}}_2)$ and a step size $\tau \in \mathbb{C}$. Recall that the Yang-Baxter relation for linear operators $R^{\ell_1 \ell_2}(x) : V^{\ell_1} \otimes V^{\ell_2} \rightarrow V^{\ell_1} \otimes V^{\ell_2}$ is the identity

$$(1.1) \quad R_{12}^{\ell_1 \ell_2}(x) R_{13}^{\ell_1 \ell_3}(x+y) R_{23}^{\ell_2 \ell_3}(y) = R_{23}^{\ell_2 \ell_3}(y) R_{13}^{\ell_1 \ell_3}(x+y) R_{12}^{\ell_1 \ell_2}(x),$$

for linear operators in $V^{\ell_1} \otimes V^{\ell_2} \otimes V^{\ell_3}$. Here we use standard tensor leg notations, e.g., $R_{12}^{\ell_1 \ell_2}(x) = R^{\ell_1 \ell_2}(x) \otimes \text{Id}_{V^{\ell_3}}$. Given $R^{\ell_1 \ell_2}(x)$ satisfying (1.1), the left and right reflection equations are identities in $V^{\ell_1} \otimes V^{\ell_2}$ for two linear operators $K^{\pm, \ell}(x) : V^{\ell} \rightarrow V^{\ell}$ ($\ell = \ell_1, \ell_2$), namely

$$(1.2) \quad \begin{aligned} R^{\ell_1 \ell_2}(x-y) K_1^{+, \ell_1}(x) P^{\ell_2 \ell_1} R^{\ell_2 \ell_1}(x+y) P^{\ell_1 \ell_2} K_2^{+, \ell_2}(y) &= \\ &= K_2^{+, \ell_2}(y) R^{\ell_1 \ell_2}(x+y) K_1^{+, \ell_1}(x) P^{\ell_2 \ell_1} R^{\ell_2 \ell_1}(x-y) P^{\ell_1 \ell_2}, \\ P^{\ell_2 \ell_1} R^{\ell_2 \ell_1}(x-y) P^{\ell_1 \ell_2} K_1^{-, \ell_1}(x) R^{\ell_1 \ell_2}(x+y) K_2^{-, \ell_2}(y) &= \\ &= K_2^{-, \ell_2}(y) P^{\ell_2 \ell_1} R^{\ell_2 \ell_1}(x+y) P^{\ell_1 \ell_2} K_1^{-, \ell_1}(x) R^{\ell_1 \ell_2}(x-y), \end{aligned}$$

respectively; here $P^{\ell_1 \ell_2} : V^{\ell_1} \otimes V^{\ell_2} \rightarrow V^{\ell_2} \otimes V^{\ell_1}$ is the flip of tensor factors.

The corresponding boundary qKZ equations are the compatible system of difference equations

$$\begin{aligned}
 (1.3) \quad \Psi(t_1, \dots, t_r + \tau, \dots, t_N) &= R_{r, r+1}^{\ell_r, \ell_{r+1}}(t_r - t_{r+1} + \tau) \cdots R_{r, N}^{\ell_r, \ell_N}(t_r - t_N + \tau) \\
 &\times K_r^{+, \ell_r}(t_r + \frac{\tau}{2}) R_{N, r}^{\ell_N, \ell_r}(t_N + t_r) \cdots R_{r+1, r}^{\ell_{r+1}, \ell_r}(t_{r+1} + t_r) \\
 &\times R_{r-1, r}^{\ell_{r-1}, \ell_r}(t_{r-1} + t_r) \cdots R_{1, r}^{\ell_1, \ell_r}(t_1 + t_r) K_r^{-, \ell_r}(t_r) \\
 &\times (R_{1, r}^{\ell_1, \ell_r}(t_1 - t_r))^{-1} \cdots (R_{r-1, r}^{\ell_{r-1}, \ell_r}(t_{r-1} - t_r))^{-1} \Psi(t_1, \dots, t_r, \dots, t_N),
 \end{aligned}$$

with $r \in \{1, \dots, N\}$ for meromorphic $V^{\ell_1} \otimes \cdots \otimes V^{\ell_N}$ -valued functions $\Psi(\mathbf{t})$ in $\mathbf{t} = (t_1, \dots, t_N) \in \mathbb{C}^N$. The Yang-Baxter equation (1.1) and the reflection equations (1.2) guarantee the compatibility of the system (1.3).

Special solutions to the boundary qKZ equations are known. First results go back to the papers [6, 7] where the Heisenberg algebra realization of q-vertex operators was used to construct a specific solution corresponding to correlation functions in the 6-vertex model. This method was applied to other models, to obtain special solutions to boundary qKZ for other R-matrices, see [8] for an overview of some results in this direction. A family of solutions to the boundary qKZ equations was constructed in our earlier papers [13, 14] as Jackson integrals (bilateral series), cf. the works [12, 19] which dealt with the same topic in type A. In [17] Laurent-polynomial solutions were found in terms of nonsymmetric Koornwinder polynomials associated to principal series modules of the affine Hecke algebra of type C.

1.1. Summary of main results. This paper can be regarded as a "type C counterpart" of [18] where a basis of integral solutions to the qKZ equations of type A was constructed. From the perspective of the Heisenberg XXZ spin chains, the present paper deals with (diagonal) integrable reflecting boundary conditions as opposed to quasi-periodic boundary conditions in [18].

We consider the boundary qKZ equation (1.3) for the tensor product of N finite-dimensional representations of quantum affine \mathfrak{sl}_2 and/or Verma modules over quantum \mathfrak{sl}_2 , for K-matrices which are diagonal in the weight basis. For a fixed total weight space in the tensor product of the representations, which is naturally labelled by a nonnegative integer M , we construct a basis for the associated meromorphic solutions $\{\Psi_{\mathbf{k}}\}_{\mathbf{k}}$ of the boundary qKZ equations over the field of $\tau\mathbb{Z}^N$ -periodic meromorphic functions admitting an explicit integral representation

$$(1.4) \quad \Psi_{\mathbf{k}}(\mathbf{t}) = \int_{C_{\mathbf{k}}(\mathbf{t})} w_{\mathbf{k}}(\mathbf{x}; \mathbf{t}) \tilde{\mathcal{B}}(\mathbf{x}; \mathbf{t}) \Omega d^M \mathbf{x}$$

for \mathbf{t} deep enough in the asymptotic sector

$$\{\mathbf{t} \mid \Re(t_1) > \Re(t_2) > \cdots > \Re(t_N) > 0\},$$

where $\Re(t)$ is the real part of t . The index set consists of M -tuples $\mathbf{k} = (k_1, \dots, k_M)$ of integers satisfying $1 \leq k_1 \leq k_2 \leq \cdots \leq k_M \leq N$, in case of Verma modules. All the elements appearing in these integrals will be precisely defined in Section 3; we only emphasize some key points here. The integration contours $C_{\mathbf{k}}(\mathbf{t})$ are Cartesian products of line segments in the complex plane of length π parallel to the imaginary axis, whose location depends on the variables t_{k_1}, \dots, t_{k_M} . The

weight functions $w_{\mathbf{k}}$ are similar to those used in [13, 14]; more precisely, they are solutions to the same difference equations, but differ by exponential factors and quotients of Jacobi theta functions. The $\tilde{\mathcal{B}}(\mathbf{x}; \mathbf{t})\Omega$ are so-called off-shell Bethe vectors for reflecting boundary conditions as introduced in [15]; they are, up to a scalar factor, those used in [13, 14]. The proof of completeness and of independence of solutions is done by a careful study of asymptotic behaviour of solutions when $\Re(t_1) \gg \Re(t_2) \gg \cdots \gg \Re(t_N) \gg 0$. It is this asymptotic analysis that relates the integral solutions of the present paper to power series solutions of boundary qKZ equations, as studied before in, e.g., [10] and [16].

1.2. Open problems. It would be very interesting to understand the precise relation between the integral solutions and the solutions considered in [6, 7] in terms of matrix elements of vertex operators with respect to so-called boundary states, as well as the Jackson integral solutions from [13, 14]. See [9] for some results in this direction for type A. Another important problem is to verify the compatibility of these solutions $\Psi_{\mathbf{k}}(\mathbf{t})$ with fusion (cf. [14, Sec. 8]) and study their classical and rational limits.

Among other open problems it would be a natural continuation of present work to find integral solutions for the boundary qKZ equations (1.3) for different R- and K-matrices. It would be nice in all these cases to construct a basis of solutions, to understand their representation-theoretical meaning and to compare with special cases known from physics literature.

1.3. Outline. In Section 2 we review the construction of the boundary qKZ equations for tensor products of Verma modules over quantum \mathfrak{sl}_2 . We restrict throughout the paper to the special case that the associated solutions of the reflections equations are diagonal with respect to the standard basis of the tensor product of Verma modules. Section 3 is the heart of the paper: here we introduce the building blocks necessary for our solutions, namely weight functions defined in terms of q-shifted factorials, the boundary Bethe vectors and the integration contours. We also define the integral solutions and state the main result of this paper, namely, that for a fixed total weight the integral solutions form a basis of the meromorphic solutions to (1.3) over the field of $\tau\mathbb{Z}^N$ -periodic meromorphic functions, taking values in the corresponding (finite-dimensional) total weight space of the tensor product of Verma modules.

The proof of the main statement is spread out over Section 4. In Section 5 we consider the boundary qKZ equations for functions taking values in the tensor product of finite-dimensional modules over quantum affine \mathfrak{sl}_2 . The appendices A and B provide technical statements underpinning the proof of the main result.

1.4. Acknowledgements. The research of N.R. was supported by the Chern-Simons chair and by the NSF grant DMS-1201391; he also acknowledges support from QGM at Aarhus, where an important part of the work has been done. B.V. is grateful to the University of California and the University of Amsterdam for hospitality; he was supported in his work by an NWO free competition grant when employed at the University of Amsterdam and EPSRC grant EP/L000865/1 when employed at the University of Nottingham.

2. REPRESENTATION THEORY AND INTEGRABLE DATA

Here we will use conventions from [14, Section 2.1] and refer to this publication for more detail and references.

2.1. The quantum affine algebra $\mathcal{U}_\eta(\widehat{\mathfrak{sl}}_2)$ and the universal R-matrix. Let $\eta \in \mathbb{C}$ such that e^η is not a root of unity and write

$$\begin{pmatrix} a_{00} & a_{01} \\ a_{10} & a_{11} \end{pmatrix} = \begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}.$$

We start with the unital Hopf algebra $\widehat{\mathcal{U}}_\eta := \mathcal{U}_\eta(\widehat{\mathfrak{sl}}_2)$ with deformation parameter e^η and generators e_0, e_1, f_0, f_1 and e^h ($h \in \mathfrak{h} := \mathbb{C}h_0 \oplus \mathbb{C}h_1$) with defining relations

$$\begin{aligned} e^0 &= 1, & e^{h+h'} &= e^h e^{h'}, \\ e^h e_i e^{-h} &= e^{\alpha_i(h)} e_i, & e^h f_i e^{-h} &= e^{-\alpha_i(h)} f_i, & [e_i, f_j] &= \delta_{i,j} \frac{e^{\eta h_i} - e^{-\eta h_i}}{e^\eta - e^{-\eta}} \end{aligned}$$

for $h, h' \in \mathfrak{h}$ and $i, j = 0, 1$, and with the Serre relations

$$\begin{aligned} e_i^3 e_j - (e^{2\eta} + 1 + e^{-2\eta}) e_i^2 e_j e_i + (e^{2\eta} + 1 + e^{-2\eta}) e_i e_j e_i^2 - e_j e_i^3 &= 0 \\ f_i^3 f_j - (e^{2\eta} + 1 + e^{-2\eta}) f_i^2 f_j f_i + (e^{2\eta} + 1 + e^{-2\eta}) f_i f_j f_i^2 - f_j f_i^3 &= 0 \end{aligned}$$

for $i \neq j$. Here the roots $\alpha_i \in \mathfrak{h}^*$ are defined by $\alpha_i(h_j) := a_{ij}$,

The coproduct Δ and the counit ϵ are determined by their action on generators:

$$\begin{aligned} \Delta(e^h) &= e^h \otimes e^h, \\ \Delta(e_i) &= e_i \otimes 1 + e^{-\eta h_i} \otimes e_i, \\ \Delta(f_i) &= f_i \otimes e^{\eta h_i} + 1 \otimes f_i \end{aligned}$$

and

$$\epsilon(e^h) = 1, \quad \epsilon(e_i) = 0, \quad \epsilon(f_i) = 0$$

for $h \in \mathfrak{h}$ and $i = 0, 1$.

Let $\widetilde{\mathcal{U}}_\eta$ be the extension of $\widehat{\mathcal{U}}_\eta$ with generators e^{xd} ($x \in \mathbb{C}$) satisfying $[e^h, e^{xd}] = 0$, $e^{xd} e_i = e^{x\delta_{i,0}} e_i e^{xd}$ and $e^{xd} f_i = e^{-x\delta_{i,0}} f_i e^{xd}$. Then $\widetilde{\mathcal{U}}_\eta$ turns into a quantized Kac-Moody algebra in a natural way (see [3, Chpt. 9]). In particular, there is a universal R-matrix $R \in \widetilde{\mathcal{U}}_\eta \widehat{\otimes} \widetilde{\mathcal{U}}_\eta$ [2, 3, 4], where $\widehat{\otimes}$ is suitably completed tensor product. The universal R-matrix has the form

$$R = \exp(\eta(c \otimes d + d \otimes c)) \mathcal{R}$$

where $c = h_0 + h_1$ and $\mathcal{R} \in \widehat{\mathcal{U}}_\eta \widehat{\otimes} \widehat{\mathcal{U}}_\eta$ is the “truncated universal R-matrix”. We denote the opposite coproduct by Δ^{op} . In the category of modules where c acts by zero (zero-level representations), \mathcal{R} satisfies all properties of the universal R-matrix:

$$\begin{aligned} \mathcal{R} \Delta(a) &= \Delta^{\text{op}}(a) \mathcal{R}, \\ (\Delta \otimes \text{id})(\mathcal{R}) &= \mathcal{R}_{13} \mathcal{R}_{23}, \quad (\text{id} \otimes \Delta)(\mathcal{R}) = \mathcal{R}_{13} \mathcal{R}_{12} \end{aligned}$$

2.2. Evaluation representations of quantum affine \mathfrak{sl}_2 . $\widehat{\mathcal{U}}_\eta$ contains quantum \mathfrak{sl}_2 as the Hopf subalgebra $\mathcal{U}_\eta = \langle e_1, f_1, e^{zh_1} \rangle$. For $\ell \in \mathbb{C}$, define a representation π^ℓ of \mathcal{U}_η on $V^\ell = \bigoplus_{d=0}^\infty \mathbb{C}v_d^\ell$ by means of

$$\begin{aligned} \pi^\ell(e^{zh_1})(v_d^\ell) &= e^{2(\ell-d)z}v_d^\ell, \\ \pi^\ell(e_1)(v_d^\ell) &= \frac{\sinh(d\eta)\sinh((2\ell+1-d)\eta)}{\sinh(\eta)^2}v_{d-1}^\ell, \quad \text{with } v_{-1}^\ell := 0, \\ \pi^\ell(f_1)(v_d^\ell) &= v_{d+1}^\ell, \end{aligned}$$

for $d \in \mathbb{Z}_{\geq 0}$. The \mathcal{U}_η -module (π^ℓ, V^ℓ) is the Verma module with highest weight ℓ and highest weight vector v_0^ℓ .

Precisely if $\ell \in \frac{1}{2}\mathbb{Z}_{\geq 0}$, the maximal \mathcal{U}_η -submodule W^ℓ is given by $\bigoplus_{d=2\ell+1}^\infty \mathbb{C}v_d^\ell$. We write $\overline{V}^\ell := V^\ell/W^\ell$ for the irreducible finite-dimensional quotient. Denote by pr^ℓ the canonical map: $V^\ell \twoheadrightarrow \overline{V}^\ell$. The cosets

$$\overline{v}_d^\ell := \text{pr}^\ell(v_d^\ell) = v_d^\ell + W^\ell \quad \text{for } d \in \{0, 1, \dots, 2\ell\}$$

form a weight basis of \overline{V}^ℓ . If $\ell = \frac{1}{2}$ we write $\overline{V} = \overline{V}^{\frac{1}{2}}$ and $\overline{v}_d = \overline{v}_d^{\frac{1}{2}}$ for $d \in \{0, 1\}$.

For each $x \in \mathbb{C}$ there exists a unique unit-preserving algebra homomorphism $\phi_x : \widehat{\mathcal{U}}_\eta \rightarrow \mathcal{U}_\eta$ satisfying

$$\begin{aligned} \phi_x(e_0) &= e^{-x}f_1, & \phi_x(e_1) &= e^{-x}e_1, \\ \phi_x(f_0) &= e^x e_1, & \phi_x(f_1) &= e^x f_1, \\ \phi_x(e^{zh_0}) &= e^{-zh_1}, & \phi_x(e^{zh_1}) &= e^{zh_1}, \end{aligned}$$

using which we define the evaluation representation $\pi_x^\ell = \pi^\ell \circ \phi_x : \widehat{\mathcal{U}}_\eta \rightarrow \text{End}(V^\ell)$.

Remark 2.1. In [14] we wrote M^ℓ instead of V^ℓ and $\{m_1^\ell, \dots, m_{2\ell+1}^\ell\}$ instead of $\{v_0^\ell, \dots, v_{2\ell}^\ell\}$. Also, V^ℓ in [14] corresponds to the present notation \overline{V}^ℓ . The purpose of this is to have simplified notation for infinite-dimensional objects (by placing a bar over the symbol to denote the corresponding finite-dimensional objects) as we will focus on these in most of this paper.

2.3. R-matrices. We now pass from the universal R-matrix to its image under the evaluation representation. For details see e.g. [4]. Let $\ell_1, \ell_2 \in \mathbb{C}$ and let $R^{\ell_1 \ell_2}(x-y) \in \text{End}(V^{\ell_1} \otimes V^{\ell_2})$ be the scalar multiple of $(\pi_x^{\ell_1} \otimes \pi_y^{\ell_2})(\mathcal{R})$ normalized by the condition

$$R^{\ell_1 \ell_2}(x)(v_0^{\ell_1} \otimes v_0^{\ell_2}) = v_0^{\ell_1} \otimes v_0^{\ell_2}.$$

Then $R^{\ell_1 \ell_2}(x)$ satisfies the Yang-Baxter equation (1.1). Furthermore, it satisfies unitarity,

$$(2.1) \quad R^{\ell_1 \ell_2}(x)^{-1} = P^{\ell_2 \ell_1} R^{\ell_2 \ell_1}(-x) P^{\ell_1 \ell_2}$$

as well as P -symmetry,

$$(2.2) \quad P^{\ell_2 \ell_1} R^{\ell_2 \ell_1}(x) P^{\ell_1 \ell_2} = R^{\ell_1 \ell_2}(x).$$

See, for example, [14, Lemma 2.1] for a proof of P -symmetry. Another important property is the (*higher-spin*) *ice rule* according to which $R^{\ell_1 \ell_2}(x)$ preserves the total weight spaces

$$(2.3) \quad \begin{aligned} (V^{\ell_1} \otimes V^{\ell_2})(M) &:= \{v \in V^{\ell_1} \otimes V^{\ell_2} \mid (\pi^{\ell_1} \otimes \pi^{\ell_2})(\Delta(e^{h_1}))v = e^{2(\ell_1 + \ell_2 - M)}v\} \\ &= \text{span}\{v_{d_1}^{\ell_1} \otimes v_{d_2}^{\ell_2} \mid d_1, d_2 \in \mathbb{Z}_{\geq 0} \text{ \& } d_1 + d_2 = M\} \end{aligned}$$

for all $M \in \mathbb{Z}_{\geq 0}$.

In the case where $\ell_1 \in \frac{1}{2}\mathbb{Z}_{\geq 0}$, there is a unique linear operator $L^{\ell_1, \ell_2}(x)$ on $\bar{V}^{\ell_1} \otimes V^{\ell_2}$ satisfying

$$(2.4) \quad (\text{pr}^{\ell_1} \otimes \text{Id}_{V^{\ell_2}}) R^{\ell_1, \ell_2}(x) = L^{\ell_1, \ell_2}(x) (\text{pr}^{\ell_1} \otimes \text{Id}_{V^{\ell_2}}).$$

Furthermore, if $\ell_1, \ell_2 \in \frac{1}{2}\mathbb{Z}_{\geq 0}$, there is a unique linear operator $\bar{R}^{\ell_1, \ell_2}(x)$ on $\bar{V}^{\ell_1} \otimes \bar{V}^{\ell_2}$ satisfying

$$(2.5) \quad (\text{pr}^{\ell_1} \otimes \text{pr}^{\ell_2}) R^{\ell_1, \ell_2}(x) = \bar{R}^{\ell_1, \ell_2}(x) (\text{pr}^{\ell_1} \otimes \text{pr}^{\ell_2}).$$

In particular, we have the spin-half R-matrix $\bar{R}(x) := \bar{R}^{\frac{1}{2}, \frac{1}{2}}(x)$ satisfying, for $d_1, d_2 \in \{0, 1\}$,

$$(2.6) \quad \bar{R}(x)(\bar{v}_{d_1} \otimes \bar{v}_{d_2}) = \begin{cases} \bar{v}_{d_1} \otimes \bar{v}_{d_2}, & \text{if } d_1 = d_2, \\ \frac{\sinh(x)}{\sinh(x+\eta)} \bar{v}_{d_1} \otimes \bar{v}_{d_2} + \frac{\sinh(\eta)}{\sinh(x+\eta)} \bar{v}_{d_2} \otimes \bar{v}_{d_1}, & \text{if } d_1 \neq d_2. \end{cases}$$

The operators $\bar{R}^{\ell_1, \ell_2}(x)$ can be recursively defined in terms of $\bar{R}(x)$ through what is known as R-matrix fusion, see [14, Sections 3.2 and 3.3] and references therein.

Given $\ell \in \mathbb{C}$ we also have the spin-half L-operators $L^\ell(x) := L^{\frac{1}{2}, \ell}(x)$, meromorphically depending on $x \in \mathbb{C}$; from (1.1), (2.4) and (2.5) it follows that they satisfy

$$(2.7) \quad \bar{R}_{00'}(x-y) L_0^\ell(x; \mathbf{t}) L_{0'}^\ell(y; \mathbf{t}) = L_{0'}^\ell(y; \mathbf{t}) L_0^\ell(x; \mathbf{t}) \bar{R}_{00'}(x-y),$$

an identity for operators in $\bar{V} \otimes \bar{V} \otimes V^\ell$, with the first copy of \bar{V} labelled 0 and the second labelled $0'$. They also satisfy *crossing symmetry*

$$(2.8) \quad L^\ell(x) = \frac{\sinh(x + (\frac{1}{2} - \ell)\eta)}{\sinh(x + (\frac{1}{2} + \ell)\eta)} \sigma_0^y L^\ell(-x - \eta)^{t_0} \sigma_0^y,$$

an identity for operators in $\bar{V} \otimes V^\ell$, with \bar{V} labelled 0, $\sigma^y = \begin{pmatrix} 0 & -\sqrt{-1} \\ \sqrt{-1} & 0 \end{pmatrix}$ and t_0 transposition relative to \bar{V} , cf. [14, Eq. (2.8)].

Remark 2.2. We emphasize that the notation $R^{\ell_1, \ell_2}(x)$ in this paper corresponds to $\mathcal{R}^{\ell_1, \ell_2}(x)$ in [14] whereas the present notation $\bar{R}^{\ell_1, \ell_2}(x)$ is called $R^{\ell_1, \ell_2}(x)$ in [14].

2.4. K-matrices. We highlight here the key points from [14, Section 4], which the reader may consult for more details and references. Let $R^{\ell_1, \ell_2}(x)$ be as in Section 2.3. Because of P-symmetry (2.2), both reflection equations (1.2) simplify to

$$(2.9) \quad \begin{aligned} R^{\ell_1, \ell_2}(x-y) K_1^{\ell_1}(x) R^{\ell_1, \ell_2}(x+y) K_2^{\ell_2}(y) = \\ K_2^{\ell_2}(y) R^{\ell_1, \ell_2}(x+y) K_1^{\ell_1}(x) R^{\ell_2, \ell_1}(x-y). \end{aligned}$$

There is a one-parameter family $K^\ell(x; \xi) \in \text{End}(V^\ell)$ of solutions to (2.9) which are diagonal in the weight basis. They act on arbitrary weight vectors according to

$$(2.10) \quad K^\ell(x; \xi) v_d^\ell = \left(\prod_{j=1}^d \frac{\sinh(\xi - x + (\ell + \frac{1}{2} - j)\eta)}{\sinh(\xi + x + (\ell + \frac{1}{2} - j)\eta)} \right) v_d^\ell,$$

where $d \in \mathbb{Z}_{\geq 0}$. In particular, they satisfy

$$K^\ell(x; \xi)(v_0^\ell) = v_0^\ell$$

and the unitarity condition

$$K^\ell(x; \xi)^{-1} = K^\ell(-x; \xi).$$

For $\ell \in \frac{1}{2}\mathbb{Z}_{\geq 0}$, the natural projection pr^ℓ to the quotient representation \bar{V}^ℓ applied to $K^\ell(x; \xi)$ gives the corresponding solution $\bar{K}^\ell(x; \xi) \in \text{End}(\bar{V}^\ell)$ to (2.9); furthermore we write $\bar{K}(x; \xi) = \bar{K}^{1/2}(x; \xi)$ as for the R-matrix. The operators $\bar{K}^\ell(x; \xi)$ can be recursively defined in terms of $\bar{R}(x)$ and $\bar{K}(x; \xi)$ through what is known as K-matrix (bulk-boundary) fusion, see e.g. [11].

Remark 2.3. Similar to the R-matrices, the notation $K^\ell(x)$ in this paper corresponds to $\mathcal{K}^\ell(x)$ in [14] whereas the present notation $\bar{K}^\ell(x)$ is called $K^\ell(x)$ in [14].

2.5. Tensor products of evaluation representations. Let $N \in \mathbb{Z}_{\geq 0}$ and fix $\ell = (\ell_1, \dots, \ell_N) \in \mathbb{C}^N$. We will be considering linear operators on

$$V^\ell := V^{\ell_1} \otimes \dots \otimes V^{\ell_N}$$

and write an arbitrary element of its natural basis as $v_{\mathbf{d}}^\ell = v_{d_1}^{\ell_1} \otimes \dots \otimes v_{d_N}^{\ell_N}$, where $\mathbf{d} = (d_1, \dots, d_N) \in \mathbb{Z}_{\geq 0}^N$. Taking into account the weight decomposition $V^\ell = \bigoplus_{d \geq 0} \mathbb{C}v_d^\ell$ with respect to the action of e^{h_1} we have the total weight decomposition

$$V^\ell = \bigoplus_{M=0}^{\infty} V^\ell(M)$$

with

$$V^\ell(M) := \bigoplus_{\mathbf{d} \in P_N(M)} \mathbb{C}v_{\mathbf{d}}^\ell,$$

$$P_N(M) := \{\mathbf{d} = (d_1, \dots, d_N) \in \mathbb{Z}_{\geq 0}^N \mid \sum_{s=1}^N d_s = M\}$$

(compare with (2.3), which is the special case $N = 2$).

The tensor product basis $\{v_{\mathbf{d}}^\ell\}_{\mathbf{d} \in P_N(M)}$ of the weight subspaces $V^\ell(M)$ can be labelled in another natural way. For $M, N \in \mathbb{Z}_{\geq 0}$, define

$$(2.11) \quad I_{M,N} = \{(k_1, \dots, k_M) \in \{1, \dots, N\}^M \mid k_1 \leq k_2 \leq \dots \leq k_M\}.$$

The following map is a bijection between $P_N(M)$ and $I_{M,N}$:

$$\zeta_{M,N} : I_{M,N} \xrightarrow{\sim} P_N(M) : \mathbf{k} \mapsto (n_{\mathbf{k}}(1), \dots, n_{\mathbf{k}}(N)),$$

where we have introduced the notation

$$(2.12) \quad n_{\mathbf{k}}(r) := \#\{i \in \{1, \dots, M\} \mid k_i = r\}$$

for $\mathbf{k} \in \{1, \dots, N\}^M$ and $r \in \{1, \dots, N\}$. The fact that $\zeta_{M,N}$ is a bijection is clear if we write an N -tuple from $I_{M,N}$ as follows:

$$(k_1, \dots, k_M) = (\underbrace{1, \dots, 1}_{n_{\mathbf{k}}(1)}, \underbrace{2, \dots, 2}_{n_{\mathbf{k}}(2)}, \underbrace{3, \dots, 3}_{n_{\mathbf{k}}(3)}, \dots, \underbrace{N, \dots, N}_{n_{\mathbf{k}}(N)}).$$

Note also that $\sum_{r=1}^N n_{\mathbf{k}}(r) = M$ for all $\mathbf{k} \in I_{M,N}$. Using $\zeta_{M,N}$ we can parametrize the tensor product basis of the weight subspace $V^\ell(M)$ by elements in $I_{N,M}$:

$$V^\ell(M) = \bigoplus_{\mathbf{k} \in I_{M,N}} \mathbb{C}\Omega_{\mathbf{k}}$$

where

$$\Omega_{\mathbf{k}} := v_{\zeta_{M,N}(\mathbf{k})}^\ell = v_{n_{\mathbf{k}}(1)}^{\ell_1} \otimes \dots \otimes v_{n_{\mathbf{k}}(N)}^{\ell_N}.$$

2.6. Boundary qKZ equations. Fix $\ell \in \mathbb{C}^N$ and $\tau, \xi_+, \xi_- \in \mathbb{C}$. Given the R- and K-matrices defined in Sections 2.3 and 2.4, the spin- ℓ boundary qKZ equations are the following equations for meromorphic V^ℓ -valued functions $\Psi(\mathbf{t})$ in $\mathbf{t} \in \mathbb{C}^N$:

$$(2.13) \quad \Psi(\mathbf{t} + \tau \mathbf{e}_r) = \mathcal{A}_r(\mathbf{t}) \Psi(\mathbf{t}), \quad r \in \{1, \dots, N\},$$

with $\{\mathbf{e}_r\}_{r=1}^N$ the standard orthonormal basis of \mathbb{C}^N and the transport operators given by

$$(2.14) \quad \begin{aligned} \mathcal{A}_r(\mathbf{t}) &= R_{r,r+1}^{\ell_r, \ell_{r+1}}(t_r - t_{r+1} + \tau) \cdots R_{r,N}^{\ell_r, \ell_N}(t_r - t_N + \tau) \\ &\quad \times K_r^{\ell_r}(t_r + \frac{\tau}{2}; \xi_+) R_{N,r}^{\ell_N, \ell_r}(t_N + t_r) \cdots R_{r+1,r}^{\ell_{r+1}, \ell_r}(t_{r+1} + t_r) \\ &\quad \times R_{r-1,r}^{\ell_{r-1}, \ell_r}(t_{r-1} + t_r) \cdots R_{1,r}^{\ell_1, \ell_r}(t_1 + t_r) K_r^{\ell_r}(t_r; \xi_-) \\ &\quad \times (R_{1,r}^{\ell_1, \ell_r}(t_1 - t_r))^{-1} \cdots (R_{r-1,r}^{\ell_{r-1}, \ell_r}(t_{r-1} - t_r))^{-1} \end{aligned}$$

for $r \in \{1, \dots, N\}$. The Yang-Baxter equation (1.1) and the reflection equation (2.9) yield the consistency conditions

$$\mathcal{A}_r(\mathbf{t} + \tau \mathbf{e}_s) \mathcal{A}_s(\mathbf{t}) = \mathcal{A}_s(\mathbf{t} + \tau \mathbf{e}_r) \mathcal{A}_r(\mathbf{t}) \quad \text{for } r, s \in \{1, \dots, N\}.$$

When $\ell_s \in \frac{1}{2}\mathbb{Z}_{\geq 0}$ the equation projects to the corresponding quotient space $V^{\ell_1} \otimes \dots \otimes V^{\ell_{s-1}} \otimes \bar{V}^{\ell_s} \otimes V^{\ell_{s+1}} \otimes \dots \otimes V^{\ell_N}$, see [14].

The finite-dimensional subspaces $V^\ell(M) \subseteq V^\ell$ ($M \geq 0$) are invariant subspaces for the transport operators $\mathcal{A}_r(\mathbf{t})$ ($r = 1, \dots, N$) since the R -matrices satisfy the ice rule and since the K -matrices act diagonally with respect to the weight basis of the Verma module. It follows that the meromorphic solutions of (2.13) are of the form $\Psi = \sum_{M \geq 0} \Psi_M$ with Ψ_M a meromorphic $V^\ell(M)$ -valued solution of (2.13). In the remainder of the paper we therefore focus on the construction of a basis of $V^\ell(M)$ -valued meromorphic solutions of the boundary qKZ equations (2.13) for a fixed value of M .

We now first discuss the existence of power series solutions of the boundary qKZ equations (2.13), for fixed value of $M \in \mathbb{Z}_{\geq 0}$. Suppose $\Re(\tau) < 0$ and consider the sector

$$\mathbb{A} := \{(t_1, \dots, t_N) \in \mathbb{C}^N \mid \Re(t_1) > \Re(t_2) > \dots > \Re(t_N) > 0\}.$$

We will write $\mathbf{t} \xrightarrow{\mathbb{A}} \infty$ when

$$\Re(t_s - t_{s+1}) \rightarrow \infty \text{ for } s \in \{1, \dots, N-1\} \quad \text{and} \quad \Re(t_N) \rightarrow \infty.$$

Note that if $\mathbf{t} \xrightarrow{\mathbb{A}} \infty$ the real parts of the arguments of all R- and K-matrices in (2.14) go to $+\infty$.

In order to study the asymptotics of the bqKZ equations (2.13) and their solutions as $\mathbf{t} \xrightarrow{\mathbb{A}} \infty$ more precisely, we rely on appendix A and [3, §9.6].

Let $Q_+ \subseteq \mathbb{R}^N$ be the cone

$$(2.15) \quad Q_+ := \bigoplus_{i=1}^{N-1} \mathbb{Z}_{\geq 0}(\mathbf{e}_i - \mathbf{e}_{i+1}) \oplus \mathbb{Z}_{\geq 0} \mathbf{e}_N.$$

For $\beta = (\beta_1, \dots, \beta_N) \in \mathbb{Z}^N$ we write $\mathbf{e}^{(\beta, \mathbf{t})} = \prod_{i=1}^N \mathbf{e}^{\beta_i t_i}$. Let $r \in \{1, \dots, N\}$. It follows from [3, §9.6] and the explicit form of the diagonal K -matrices that there

exist $\mathcal{A}_{\alpha;r} \in \text{End}(V^\ell(M))$ for $\alpha \in Q^+$ such that

$$\mathcal{A}_r(\mathbf{t}) = \sum_{\alpha \in Q_+} \mathcal{A}_{\alpha;r} e^{-(\alpha, \mathbf{t})}$$

as linear operator on $V^\ell(M)$, with the power series converging normally on compact sets for \mathbf{t} deep enough in the sector \mathbb{A} . Write $\mathbf{0} = (0, \dots, 0) \in \mathbb{C}^N$. Lemma A.2 implies that $\mathcal{A}_{\mathbf{0};r}$ acts on $V^\ell(M)$ by the formula

$$(2.16) \quad \mathcal{A}_{\mathbf{0};r} \Omega_{\mathbf{m}} = \varphi_{\mathbf{m};r} \Omega_{\mathbf{m}}, \quad \mathbf{m} \in I_{M,N},$$

where

$$(2.17) \quad \varphi_{\mathbf{m};r} := \left(\prod_{\substack{i=1 \\ m_i=r}}^M e^{2(\eta - \xi_+ - \xi_-) + 4(M-i - \sum_{s \geq r} \ell_s) \eta} \right) e^{-4\ell_r \# \{i \in \{1, \dots, M\} \mid m_i > r\} \eta}.$$

Let $\Phi_{\mathbf{k}}$ be a nonzero meromorphic function on \mathbb{C}^N satisfying the scalar difference equations

$$(2.18) \quad \Phi_{\mathbf{k}}(\mathbf{t} + \tau \mathbf{e}_r) = \varphi_{\mathbf{k};r} \Phi_{\mathbf{k}}(\mathbf{t}), \quad r \in \{1, \dots, N\}.$$

Then it follows that

$$\Psi_{\mathbf{k}}^\infty(\mathbf{t}) := \Phi_{\mathbf{k}}(\mathbf{t}) \Omega_{\mathbf{k}}$$

satisfies the asymptotic boundary qKZ equations

$$\Psi_{\mathbf{k}}^\infty(\mathbf{t} + \tau \mathbf{e}_r) = \mathcal{A}_{\mathbf{0};r} \Psi_{\mathbf{k}}^\infty(\mathbf{t}), \quad r = 1, \dots, N.$$

So it then makes sense to look for power series solutions $\Psi_{\mathbf{k}}(\mathbf{t})$ of (2.13) tending to $\Psi_{\mathbf{k}}^\infty(\mathbf{t})$ as $\mathbf{t} \xrightarrow{\mathbb{A}} \infty$.

It is easy to construct an explicit solution $\Phi_{\mathbf{k}}$ of (2.18) as a quotient of products of renormalized theta functions. We first need to introduce some notations. We write $q = e^\tau$ and we will assume $\Re(\tau) < 0$, so that $0 < |q| < 1$. The q^2 -shifted factorial

$$(z; q^2)_\infty := \prod_{m \geq 0} (1 - zq^{2m})$$

is a holomorphic function of z satisfying the property $(q^2 z; q^2)_\infty = (1-z)^{-1} (z; q^2)_\infty$. We will employ the notation

$$(z_1, z_2; q^2)_\infty := (z_1; q^2)_\infty (z_2; q^2)_\infty.$$

The renormalized Jacobi theta function is the holomorphic function defined by

$$\theta(z) := (z, q^2 z^{-1}; q^2)_\infty.$$

It satisfies the quasi-periodicity condition $\theta(q^2 z) = -z^{-1} \theta(z)$. We will use in formulas the notation $a(\pm x)$ to stand for $a(x)a(-x)$. For instance, $(e^{\pm x}; q^2)_\infty$ stands for $(e^x, e^{-x}; q^2)_\infty$.

An explicit nonzero meromorphic solution $\Phi_{\mathbf{k}}$ of (2.18) is now given by

$$\Phi_{\mathbf{k}}(\mathbf{t}) = \prod_{r=1}^N \frac{\theta(e^{2t_r})}{\theta(\varphi_{\mathbf{k};r} e^{2t_r})}.$$

Any other choice differs from $\Phi_{\mathbf{k}}$ by a nonzero $\tau \mathbb{Z}^N$ -periodic meromorphic function.

If Ψ is a $V^\ell(M)$ -valued meromorphic solution of the boundary qKZ equations (2.13) and $\mathbf{k} \in I_{M,N}$ is fixed, then

$$\tilde{\Theta}_{\mathbf{k}} := \Phi_{\mathbf{k}}^{-1} \Psi$$

is a $V^\ell(M)$ -valued meromorphic solution of the rescaled boundary qKZ equations

$$(2.19) \quad \tilde{\Psi}(\mathbf{t} + \tau \mathbf{e}_r) = \tilde{\mathcal{A}}_r^{\mathbf{k}}(\mathbf{t}) \tilde{\Psi}(\mathbf{t}), \quad r \in \{1, \dots, N\},$$

with rescaled transport operators $\tilde{\mathcal{A}}_r^{\mathbf{k}}(\mathbf{t}) := \varphi_{\mathbf{k};r}^{-1} \mathcal{A}_r(\mathbf{t})$. Now the leading coefficient $\tilde{\mathcal{A}}_{0;r}^{\mathbf{k}}$ of the power series expansion

$$\tilde{\mathcal{A}}_r^{\mathbf{k}}(\mathbf{t}) = \sum_{\alpha \in Q_+} \tilde{\mathcal{A}}_{\alpha;r}^{\mathbf{k}} e^{-(\alpha, \mathbf{t})}$$

is acting on $V^\ell(M)$ by

$$\tilde{\mathcal{A}}_{0;r}^{\mathbf{k}} \Omega_{\mathbf{m}} = \frac{\varphi_{\mathbf{m};r}}{\varphi_{\mathbf{k};r}} \Omega_{\mathbf{m}}, \quad \mathbf{m} \in I_{M,N}.$$

Then [10, Appendix] guarantees, for generic parameters, the existence and uniqueness of a $V^\ell(M)$ -valued meromorphic solution $\tilde{\Theta}_{\mathbf{k}}$ of the rescaled boundary qKZ equations (2.19) such that

$$\tilde{\Theta}_{\mathbf{k}}(\mathbf{t}) = \sum_{\alpha \in Q_+} \tilde{L}_{\alpha}^{\mathbf{k}} e^{-(\alpha, \mathbf{t})}, \quad \tilde{L}_{\alpha}^{\mathbf{k}} \in V^\ell(M)$$

for \mathbf{t} deep enough in the sector \mathbb{A} , with the $V^\ell(M)$ -valued power series normally converging on compact sets and with leading coefficient

$$\tilde{L}_0^{\mathbf{k}} = \Omega_{\mathbf{k}}.$$

Our main goal is to find an explicit integral expression of $\tilde{\Theta}_{\mathbf{k}}$ deep enough in the sector \mathbb{A} .

3. INTEGRAL SOLUTIONS OF THE BOUNDARY qKZ EQUATIONS

We will exhibit $V^\ell(M)$ -valued solutions $\Psi_{\mathbf{k}}(\mathbf{t})$ of (2.13) for $\mathbf{k} = (k_1, \dots, k_M) \in I_{M,N}$ admitting an integral representation of the form

$$\Psi_{\mathbf{k}}(\mathbf{t}) = \int_{C_{\mathbf{k}}(\mathbf{t})} w_{\mathbf{k}}(\mathbf{x}; \mathbf{t}) \tilde{\mathcal{B}}(\mathbf{x}; \mathbf{t}) \Omega d^M \mathbf{x},$$

on some subsector $\tilde{\mathbb{A}} \subset \mathbb{A}$. We will show that, up to an explicit multiplicative constant, $\Psi_{\mathbf{k}}$ equals $\Phi_{\mathbf{k}} \tilde{\Theta}_{\mathbf{k}}$, which provides the link with the power series solutions of the boundary qKZ equations from the previous subsection.

The Bethe vectors $\tilde{\mathcal{B}}(\mathbf{x}; \mathbf{t}) \Omega$ are elements of $V^\ell(M)$ with trigonometric dependence on $\mathbf{x} \in \mathbb{C}^M$ and $\mathbf{t} \in \mathbb{C}^N$. They will be discussed in Subsection 3.1. The scalar weight functions $w_{\mathbf{k}}(\mathbf{x}; \mathbf{t})$ will be defined in Subsection 3.2. We will specify the integration contour $C_{\mathbf{k}}(\mathbf{t})$ in Subsection 3.3 before stating the main theorem in Subsection 3.4.

3.1. Bethe vectors. Like the qKZ transport operators $\mathcal{A}_r(\mathbf{t})$, the objects $\tilde{\mathcal{B}}(\mathbf{x}; \mathbf{t})$ are linear operators on V^ℓ constructed in terms of the R- and K-matrices introduced in Section 2, but according to a different procedure, first conceived for quantum integrable systems with reflecting boundaries by Sklyanin [15]. We first introduce linear operators acting on $\bar{V} \otimes V^\ell$, where the solitary tensor factor $\bar{V} \cong \mathbb{C}^2$ is called auxiliary space and is labelled 0.

Recall the L-operators $L^\ell(x) = L^{\frac{1}{2}\ell}(x) \in \text{End}(\bar{V} \otimes V^\ell)$ defined through (2.4). Fix $\ell = (\ell_1, \dots, \ell_N) \in \mathbb{C}^N$. Define the (type A) monodromy matrix

$$T(x; \mathbf{t}) = L_{01}^{\ell_1}(x - t_1) \cdots L_{0N}^{\ell_N}(x - t_N) \in \text{End}(\bar{V} \otimes V^\ell).$$

From (2.7) it follows that the $T(x; \mathbf{t})$ satisfy

$$(3.1) \quad \bar{R}_{00'}(x-y)T_0(x; \mathbf{t})T_{0'}(y; \mathbf{t}) = T_{0'}(y; \mathbf{t})T_0(x; \mathbf{t})\bar{R}_{00'}(x-y),$$

an identity in $\text{End}(\bar{V} \otimes \bar{V} \otimes V^\ell)$, with the first copy of \bar{V} labelled 0 and the second labelled $0'$. Fix $\xi_- \in \mathbb{C}$ and define

$$\mathcal{T}(x; \mathbf{t}) = T(-x; \mathbf{t})^{-1}(\bar{K}(x; \xi_-) \otimes \text{Id}_{V^\ell})T(x; \mathbf{t}) \in \text{End}(\bar{V} \otimes V^\ell).$$

Given that $T(x; \mathbf{t})$ satisfies (3.1) and $\bar{K}(x; \xi_-)$ satisfies the reflection equation in $\text{End}(\bar{V} \otimes \bar{V})$, it can be straightforwardly verified that $\mathcal{T}(x; \mathbf{t})$ satisfies the reflection equation in $\text{End}(\bar{V} \otimes \bar{V} \otimes V^\ell)$:

$$(3.2) \quad \begin{aligned} \bar{R}_{00'}(x-y)\mathcal{T}_0(x; \mathbf{t})\bar{R}_{00'}(x+y)\mathcal{T}_{0'}(y; \mathbf{t}) = \\ \mathcal{T}_{0'}(y; \mathbf{t})\bar{R}_{00'}(x+y)\mathcal{T}_0(x; \mathbf{t})\bar{R}_{00'}(x-y). \end{aligned}$$

We introduce an operator $\mathcal{B}(x; \mathbf{t}) \in \text{End}(V^\ell)$ by means of

$$(3.3) \quad \mathcal{T}(x; \mathbf{t}) = \begin{pmatrix} * & \mathcal{B}(x; \mathbf{t}) \\ * & * \end{pmatrix},$$

i.e. for all $u \in V^\ell$ we have

$$\mathcal{T}(x; \mathbf{t})(\bar{v}_1 \otimes u) = \bar{v}_0 \otimes \mathcal{B}(x; \mathbf{t})(u) + \bar{v}_1 \otimes (\text{some element of } V^\ell).$$

In the rest of the paper we will use the shorthand notations

$$\tilde{\xi}_+ := \xi_+ - \frac{\eta}{2} - \frac{\tau}{2}, \quad \tilde{\xi}_- := \xi_- - \frac{\eta}{2}.$$

It is convenient, as will become apparent in Appendix B.2, to use a slightly modified version of \mathcal{B} , namely

$$\tilde{\mathcal{B}}(x; \mathbf{t}) := \left(\prod_{s=1}^N \frac{\sinh(t_s - x + \ell_s \eta)}{\sinh(t_s - x - \ell_s \eta)} \right) \frac{\sinh(2x)}{\sinh(2x + \eta)} \frac{\sinh(\tilde{\xi}_- - x)}{\sinh(\eta)} \mathcal{B}(-x - \frac{\eta}{2}; \mathbf{t}).$$

Fix $M \in \mathbb{Z}_{\geq 0}$ and define, for $\mathbf{x} = (x_1, \dots, x_M)$ and $\mathbf{t} = (t_1, \dots, t_N)$,

$$\tilde{\mathcal{B}}(\mathbf{x}; \mathbf{t}) = \tilde{\mathcal{B}}(x_1; \mathbf{t}) \cdots \tilde{\mathcal{B}}(x_M; \mathbf{t}) \in \text{End}(V^\ell)$$

(we do not specify the depth M in the notation of $\tilde{\mathcal{B}}$, it will be clear from context).

The *off-shell spin- ℓ boundary (or type C) Bethe vectors* are the elements $\tilde{\mathcal{B}}(\mathbf{x}; \mathbf{t})\Omega \in V^\ell(M)$, where Ω is the tensor product of highest weight vectors:

$$\Omega := \Omega_\emptyset = v_0^{\ell_1} \otimes \cdots \otimes v_0^{\ell_N} \in V^\ell.$$

One of the results of this paper is an explicit decomposition of the boundary Bethe vectors in terms of the tensor product basis of $V^\ell(M)$.

Proposition 3.1. *As meromorphic $V^\ell(M)$ -valued functions in $(\mathbf{x}, \mathbf{t}) \in \mathbb{C}^M \times \mathbb{C}^N$, we have*

$$(3.4) \quad \tilde{\mathcal{B}}(\mathbf{x}; \mathbf{t})\Omega = \sum_{\mathbf{k} \in I_{M,N}} \beta_{\mathbf{k}}(\mathbf{x}; \mathbf{t})\Omega_{\mathbf{k}}$$

where

$$\begin{aligned} \beta_{\mathbf{k}}(\mathbf{x}; \mathbf{t}) = & e^{\sum_i \left(\frac{n_{\mathbf{k}}(k_i)}{2} - \ell_{k_i} \right) \eta} \sum_{\mathbf{m} \in S_M} \sum_{\mathbf{k} \in \{\pm\}^M} \left(\prod_i \frac{\epsilon_i \sinh(\epsilon_i x_i - \tilde{\xi}_-)}{\sinh(t_{m_i} + \epsilon_i x_i - \ell_{m_i} \eta)} \right. \\ & \times \left(\prod_{s > m_i} \frac{\sinh(t_s + \epsilon_i x_i + \ell_s \eta)}{\sinh(t_s + \epsilon_i x_i - \ell_s \eta)} \right) \left(\prod_s \frac{\sinh(t_s - \epsilon_i x_i + \ell_s \eta)}{\sinh(t_s - \epsilon_i x_i - \ell_s \eta)} \right) \\ & \times \left(\prod_{\substack{i,j \\ i < j}} \frac{\sinh(\epsilon_i x_i + \epsilon_j x_j + \eta)}{\sinh(\epsilon_i x_i + \epsilon_j x_j)} \right) \left(\prod_{\substack{i,j \\ m_i < m_j}} \frac{\sinh(\epsilon_i x_i - \epsilon_j x_j - \eta)}{\sinh(\epsilon_i x_i - \epsilon_j x_j)} \right). \end{aligned}$$

Essentially, the result owes to the higher-spin ice rule of the R -operators and the diagonality of $\bar{K}(x; \xi_-)$. The proof of the Theorem is given in appendix B.

Note that Proposition 3.1 implies that the off-shell boundary Bethe vectors $\tilde{\mathcal{B}}(\mathbf{x}; \mathbf{t})\Omega$ are $\pi\sqrt{-1}$ -periodic in each x_i .

3.2. Weight functions. Fix $\tau, \eta, \xi_+, \xi_- \in \mathbb{C}$ and $\ell \in \mathbb{C}^N$. Suppose that $\Re(\tau) < 0$ and set $q = e^\tau$. For $\mathbf{t} \in \mathbb{C}^N$, introduce the single-variable meromorphic $\pi\sqrt{-1}$ -periodic functions $F(\cdot; \mathbf{t})$, g , h as follows:

$$\begin{aligned} (3.5) \quad F(x; \mathbf{t}) &:= \prod_{s=1}^N \frac{(e^{-2(t_s \pm x - \ell_s \eta)}; q^2)_\infty}{(e^{-2(t_s \pm x + \ell_s \eta)}; q^2)_\infty}, \\ g(x) &:= \frac{(q^2 e^{2(\tilde{\xi}_+ - x)}, q^2 e^{2(\tilde{\xi}_- - x)}; q^2)_\infty}{(e^{2(-\tilde{\xi}_+ - x)}, e^{2(-\tilde{\xi}_- - x)}; q^2)_\infty}, \\ h(x) &:= (1 - e^{-2x}) \frac{(q^2 e^{-2(x+\eta)}; q^2)_\infty}{(e^{-2(x-\eta)}; q^2)_\infty}. \end{aligned}$$

For $i \in \{1, \dots, M\}$ and $\mathbf{k} \in I_{M,N}$ define the single-variable meromorphic $\pi\sqrt{-1}$ -periodic function $u_{\mathbf{k};i}(\cdot; \mathbf{t})$ by

$$(3.6) \quad u_{\mathbf{k};i}(x; \mathbf{t}) := e^{-t_{k_i}} \theta(e^{2(x-t_{k_i}+\psi_{\mathbf{k};i})}) \frac{\prod_{s > k_i} \theta(e^{2(x-t_s-\ell_s \eta)})}{\prod_{s \geq k_i} \theta(e^{2(x-t_s+\ell_s \eta)})}$$

where

$$\psi_{\mathbf{k};i} = \tilde{\xi}_+ + \tilde{\xi}_- + \tau + \ell_{k_i} \eta + 2 \left(\sum_{s > k_i} \ell_s - M + i \right) \eta.$$

We use these functions as building blocks to define a meromorphic weight function $w_{\mathbf{k}}$ on $\mathbb{C}^M \times \mathbb{C}^N$ by

$$(3.7) \quad w_{\mathbf{k}}(\mathbf{x}; \mathbf{t}) := \Phi_{\mathbf{k}}(\mathbf{t}) \left(\prod_{i=1}^M F(x_i; \mathbf{t}) g(x_i) u_{\mathbf{k};i}(x_i; \mathbf{t}) \right) \prod_{1 \leq i < j \leq M} h(x_i \pm x_j)$$

By writing out the theta-functions as products of q^2 -shifted factorials, cancellation with q^2 -shifted factorials coming from F takes place. It leads to the expression

$$\begin{aligned}
 (3.8) \quad w_{\mathbf{k}}(\mathbf{x}; \mathbf{t}) = & \Phi_{\mathbf{k}}(\mathbf{t}) \left\{ \prod_{i=1}^M \frac{\prod_{s < k_i} (e^{2(x_i - t_s + \ell_s \eta)}; q^2)_{\infty}}{\prod_{s \leq k_i} (e^{2(x_i - t_s - \ell_s \eta)}; q^2)_{\infty}} e^{-t_{k_i}} \theta(e^{2(x_i - t_{k_i} + \psi_{\mathbf{k}; i})}) \right. \\
 & \times \frac{\prod_{s > k_i} (q^2 e^{2(t_s - x_i + \ell_s \eta)}; q^2)_{\infty}}{\prod_{s \geq k_i} (q^2 e^{2(t_s - x_i - \ell_s \eta)}; q^2)_{\infty}} \left(\prod_s \frac{(e^{-2(t_s + x_i - \ell_s \eta)}; q^2)_{\infty}}{(e^{-2(t_s + x_i + \ell_s \eta)}; q^2)_{\infty}} \right) \\
 & \times \frac{(q^2 e^{2(\tilde{\xi}_+ - x_i)}, q^2 e^{2(\tilde{\xi}_- - x_i)}; q^2)_{\infty}}{(e^{2(-\tilde{\xi}_+ - x_i)}, e^{2(-\tilde{\xi}_- - x_i)}; q^2)_{\infty}} \Big\} \\
 & \times \left\{ \prod_{\substack{i, j=1 \\ i < j}}^M (1 - e^{-2(x_i \pm x_j)}) \frac{(q^2 e^{-2(x_i \pm x_j + \eta)}; q^2)_{\infty}}{(e^{-2(x_i \pm x_j - \eta)}; q^2)_{\infty}} \right\}.
 \end{aligned}$$

The poles of $w_{\mathbf{k}}(\mathbf{x}; \mathbf{t})$ in x_i are unilateral sequences whose real parts tend to either ∞ or $-\infty$ in steps of size $-\Re(\tau)$. Also, $w_{\mathbf{k}}(\mathbf{x}; \mathbf{t})$ is $\pi\sqrt{-1}$ -periodic in each x_i .

3.3. Integration contours. We fix $\tau \in \mathbb{C}$ with $\Re(\tau) < 0$ and set $q = e^{\tau}$. We furthermore fix $N \in \mathbb{Z}_{>0}$ and $M \in \mathbb{Z}_{\geq 0}$. Write $\ell = (\ell_1, \dots, \ell_N)$ for the N -tuple of highest weights.

Definition 3.2. For $\mathbf{k} \in I_{M,N}$ we write $\mathcal{D}_{M,N}^{\mathbf{k}}$ for the set of parameters $(\ell, \eta) \in \mathbb{C}^N \times \mathbb{C}$ satisfying

$$\Re(\ell_r \eta) > \max\left(0, \frac{n_{\mathbf{k}}(r) - 1}{2} \Re(\eta)\right) \quad \forall r \in \{1, \dots, N\}.$$

Furthermore, set

$$(3.9) \quad \mathcal{D}_{M,N} := \bigcap_{\mathbf{k} \in I_{M,N}} \mathcal{D}_{M,N}^{\mathbf{k}}.$$

For $\mathbf{k} \in I_{M,N}$ we write

$$(3.10) \quad i_{\mathbf{k}}(m; r) := \sum_{s < r} n_{\mathbf{k}}(s) + m$$

for $r \in \{1, \dots, N\}$ and $m \in \{1, \dots, n_{\mathbf{k}}(r)\}$, so that $k_i = r$ if and only if $i = i(m; r)$ for some $1 \leq m \leq n_{\mathbf{k}}(r)$. We now define for $\mathbf{k} \in I_{M,N}$ and $(\ell, \eta) \in \mathcal{D}_{M,N}^{\mathbf{k}}$ the set $\Gamma_{M,N}^{\mathbf{k}}$ of base points of the integration cycle as the set of M -tuples $\gamma = (\gamma_1, \dots, \gamma_M) \in \mathbb{C}^M$ satisfying

$$\begin{aligned}
 & -\Re(\ell_{k_i} \eta) < \Re(\gamma_i) < \Re(\ell_{k_i} \eta), \\
 & \Re(\gamma_{i(s+1;r)}) \leq \Re(\gamma_{i(s;r)}), \\
 & \Re(\gamma_{i(s+1;r)}) + \Re(\eta) < \Re(\gamma_{i(s;r)})
 \end{aligned}$$

for $1 \leq i \leq M$, $1 \leq r \leq N$ and $1 \leq s < n_{\mathbf{k}}(r)$. Note that $\Gamma_{M,N}^{\mathbf{k}}$ is nonempty and path-connected if $(\ell, \eta) \in \mathcal{D}_{M,N}^{\mathbf{k}}$.

Let $\mathbf{k} \in I_{M,N}$, $(\ell, \eta) \in \mathcal{D}_{M,N}^{\mathbf{k}}$ and $\gamma \in \Gamma_{M,N}^{\mathbf{k}}$. The integration cycles that will be used in the definition of the integral solution of the boundary qKZ equations (2.13) are of the form

$$C_{\mathbf{k}}^{\gamma}(\mathbf{t}) := (t_{k_1} + \gamma_1 + \sqrt{-1}[0, \pi]) \times \dots \times (t_{k_M} + \gamma_M + \sqrt{-1}[0, \pi])$$

for $\mathbf{t} = (t_1, \dots, t_N) \in \mathbb{C}^N$.

3.4. Main result. The set $\{\mathbf{e}_i - \mathbf{e}_{i+1}\}_{i=1}^{N-1} \cup \{\mathbf{e}_N\}$ is a choice of simple roots of the standard realization of the root system of type B_N in \mathbb{R}^N . Let Q_+ be the cone in \mathbb{Z}^N generated by these simple roots, see (2.15). Denote

$$\Lambda_+ := \max_{s \in \{1, \dots, N\}} \Re(\ell_s \eta).$$

The main result of the paper can now be stated as follows.

Theorem 3.3. Fix $N \in \mathbb{Z}_{>0}$ and $M \in \mathbb{Z}_{\geq 0}$. Set $q = e^\tau$ with $\Re(\tau) < 0$.

a. Let $\mathbf{k} \in I_{M,N}$ and $(\ell, \eta) \in \mathcal{D}_{M,N}^{\mathbf{k}}$. For $\gamma \in \Gamma_{M,N}^{\mathbf{k}}$ the integral

$$(3.11) \quad \Theta_{\mathbf{k}}(\mathbf{t}) := \int_{C_{\mathbf{k}}^{\gamma}(\mathbf{t})} \frac{w_{\mathbf{k}}(\mathbf{x}; \mathbf{t})}{\Phi_{\mathbf{k}}(\mathbf{t})} \tilde{\mathcal{B}}(\mathbf{x}; \mathbf{t}) \Omega d^M \mathbf{x}$$

defines a $V^\ell(M)$ -valued holomorphic function in $\mathbf{t} \in \tilde{\mathbb{A}}$, where

$$\tilde{\mathbb{A}} := \left\{ \mathbf{t} \in \mathbb{A} \mid \begin{aligned} &\Re(t_s - t_{s+1}) > 2\Lambda_+ + \max(\Re(\eta), 0) \text{ for } 1 \leq s < N, \\ &\Re(t_N) > \Lambda_+ + \max(\Re(\tfrac{\eta}{2}), 0, \Re(-\tilde{\xi}_+), \Re(-\tilde{\xi}_-)) \end{aligned} \right\}.$$

The integral $\Theta_{\mathbf{k}}(\mathbf{t})$ does not depend on the choice of $\gamma \in \Gamma_{M,N}^{\mathbf{k}}$.

b. Let $\mathbf{k} \in I_{M,N}$ and $(\ell, \eta) \in \mathcal{D}_{M,N}^{\mathbf{k}}$. There exists a unique $V^\ell(M)$ -valued meromorphic solution $\Psi_{\mathbf{k}}$ of the boundary qKZ equations (2.13) on \mathbb{C}^N such that, on the sector $\tilde{\mathbb{A}}$,

$$(3.12) \quad \Psi_{\mathbf{k}}(\mathbf{t}) = \Phi_{\mathbf{k}}(\mathbf{t}) \Theta_{\mathbf{k}}(\mathbf{t}) = \int_{C_{\mathbf{k}}^{\gamma}(\mathbf{t})} w_{\mathbf{k}}(\mathbf{x}; \mathbf{t}) \tilde{\mathcal{B}}(\mathbf{x}; \mathbf{t}) \Omega d^M \mathbf{x}.$$

c. Let $\mathbf{k} \in I_{M,N}$ and $(\ell, \eta) \in \mathcal{D}_{M,N}^{\mathbf{k}}$. The $V^\ell(M)$ -valued integral $\Theta_{\mathbf{k}}(\mathbf{t})$ has a $V^\ell(M)$ -valued series expansion for $\mathbf{t} \in \tilde{\mathbb{A}}$ of the form

$$(3.13) \quad \Theta_{\mathbf{k}}(\mathbf{t}) = \sum_{\alpha \in Q_+} L_{\alpha}^{\mathbf{k}} e^{-(\alpha, \mathbf{t})}, \quad L_{\alpha}^{\mathbf{k}} \in V^\ell(M),$$

with the series converging normally for \mathbf{t} in compact subsets of $\tilde{\mathbb{A}}$. The leading coefficient is given by $L_{\mathbf{0}}^{\mathbf{k}} = \nu_{\mathbf{k}} \Omega_{\mathbf{k}}$, where

$$(3.14) \quad \begin{aligned} \nu_{\mathbf{k}} &:= (\pi \sqrt{-1} e^{\xi_-})^M \left(\prod_{1 \leq r < s \leq N} e^{2\ell_r n_{\mathbf{k}}(s) \eta} \right) \\ &\times \prod_{r=1}^N \prod_{m=1}^{n_{\mathbf{k}}(r)} \frac{(q^2 e^{-2m\eta}, q e^{2((m-1-\ell_r)\eta \pm \omega_{\mathbf{k};r})}; q^2)_{\infty}}{(q^2, q^2 e^{-2\eta}, e^{2(m-1-2\ell_r)\eta}; q^2)_{\infty}}, \end{aligned}$$

$$(3.15) \quad \omega_{\mathbf{k};r} := \xi_+ + \xi_- + \tau + (\ell_r - n_{\mathbf{k}}(r))\eta + 2 \sum_{s>r} (\ell_s - n_{\mathbf{k}}(s))\eta.$$

d. Let $(\ell, \eta) \in \mathcal{D}_{M,N}$. Then $\{\Psi_{\mathbf{k}} \mid \mathbf{k} \in I_{M,N}\}$ is a linear basis of the space of $V^\ell(M)$ -valued meromorphic solutions of the boundary qKZ equations (2.13) over the field of $\tau \mathbb{Z}^N$ -periodic meromorphic functions.

Remark 3.4. We may write

$$\Psi_{\mathbf{k}}(\mathbf{t}) = \sum_{\mathbf{m} \in I_{M,N}} \left(\int_{C_{\mathbf{k}}^{\gamma}(\mathbf{t})} w_{\mathbf{k}}(\mathbf{x}; \mathbf{t}) \beta_{\mathbf{m}}(\mathbf{x}; \mathbf{t}) d^M \mathbf{x} \right) \Omega_{\mathbf{m}}$$

for $\mathbf{t} \in \tilde{\mathbb{A}}$ with the coefficients $\beta_{\mathbf{m}}$ as given in Theorem 3.1. It allows for a direct comparison with the integral solutions to type A qKZ in [18]. The coefficients $\beta_{\mathbf{m}}(\mathbf{x}; \mathbf{t})$ of the Bethe vectors are the boundary analogons of the “trigonometric weight functions” appearing in [18], whereas $w_{\mathbf{k}}(\mathbf{x}; \mathbf{t})$ corresponds to the product of the “short phase function” and the “elliptic weight function” and $\Phi_{\mathbf{k}}(\mathbf{t})$ is the direct counterpart of the “adjusting factor” of the elliptic weight function. More precisely, our integrand is the direct analogon of the integrand $F(t)$ considered in [18, p.43, Proof of Theorem 6.6]. Here we add that in [18] the elliptic weight function, like the trigonometric weight function, is defined as an orbit sum. However, owing to invariance properties under an action of the symmetric group, its terms all contribute the same to the integral. In [9], this type A elliptic weight function was re-defined as a single term in order to make a connection with solutions to the qKZ equations in terms of formulae derived from free-field realizations of intertwiners of quantum affine \mathfrak{sl}_2 .

The version of Theorem 3.3 for solutions of the boundary qKZ equations (2.13) taking values in the tensor product of finite dimensional modules over quantum \mathfrak{sl}_2 is discussed in Section 5.

4. PROOF OF THE MAIN RESULTS

Here we will prove the various statements made in Theorem 3.3. We start by a helpful lemma listing the poles of the integrand of $\Theta_{\mathbf{k}}(\mathbf{t})$.

Lemma 4.1. *Let $j \in \{1, \dots, M\}$. Fix $x_i \in t_{k_i} + \gamma_i + \sqrt{-1}[0, \pi]$ for $i \neq j$. The poles of the integrand $w_{\mathbf{k}}(\mathbf{x}; \mathbf{t})\tilde{\mathcal{B}}(\mathbf{x}; \mathbf{t})\Omega$ as a function of x_j are contained in*

$$\left((P_{\mathbf{k};j}^+(\mathbf{t}) - \tau\mathbb{Z}_{\geq 0}) \cup (P_{\mathbf{k};j}^-(\mathbf{t}) + \tau\mathbb{Z}_{\geq 0}) \right) + \pi\sqrt{-1}\mathbb{Z},$$

where

$$\begin{aligned} P_{\mathbf{k};j}^+(\mathbf{t}) &= \{t_s + \ell_s\eta\}_{s \leq k_j} \cup \{x_i - \eta\}_{i < j} \\ P_{\mathbf{k};j}^-(\mathbf{t}) &= \{t_s - \ell_s\eta\}_{s \geq k_j} \cup \{-t_s - \ell_s\eta\}_s \cup \{-\tilde{\xi}_+, -\tilde{\xi}_-\} \cup \\ &\quad \cup \{x_i + \eta\}_{i > j} \cup \{-x_i + \eta\}_{i \neq j}. \end{aligned}$$

Proof. Since $\Re(\tau) < 0$, from (3.8) we see that the sequences of poles of $x_j \mapsto w_{\mathbf{k}}(\mathbf{x}; \mathbf{t})$ whose real parts tend to $+\infty$ are given by

$$\left(\{t_s + \ell_s\eta \mid 1 \leq s \leq k_j\} \cup \{x_i - \eta \mid i < j\} \right) - \tau\mathbb{Z}_{\geq 0} + \pi\sqrt{-1}\mathbb{Z}$$

whereas the sequences of poles whose real parts tend to $-\infty$ are given by

$$\begin{aligned} &\left(\{t_s - \ell_s\eta + \tau \mid k_j \leq s \leq N\} \cup \{-t_s - \ell_s\eta \mid 1 \leq s \leq N\} \cup \{-\tilde{\xi}_+, -\tilde{\xi}_-\} \cup \right. \\ &\quad \left. \cup \{x_i + \eta \mid i > j\} \cup \{-x_i + \eta \mid i \neq j\} \right) + \tau\mathbb{Z}_{\geq 0} + \pi\sqrt{-1}\mathbb{Z}. \end{aligned}$$

From Proposition 3.1 we see that the poles of $x_j \mapsto \beta_{\mathbf{k}'}(\mathbf{x}; \mathbf{t})$ are contained in

$$\left(\{\pm(t_s - \ell_s\eta) \mid 1 \leq s \leq N\} \cup \{\pm x_i \mid i \neq j\} \right) + \pi\sqrt{-1}\mathbb{Z}.$$

Using (3.8) again, we see that all of these poles except those of the form $t_s - \ell_s\eta$ for $1 \leq s \leq k_j$ are cancelled by zeros of $w_{\mathbf{k}}(\mathbf{x}; \mathbf{t})$. We obtain the desired statement. \square

In the following four subsections we prove the four parts of Theorem 3.3.

4.1. Holomorphicity of the $\Theta_{\mathbf{k}}$. Here and in Section 4.3 we will use the short-hand notation

$$\mathbf{t}_{\mathbf{k}} := (t_{k_1}, \dots, t_{k_M}) \in \mathbb{C}^M$$

for $\mathbf{t} = (t_1, \dots, t_N) \in \mathbb{C}^N$ and $\mathbf{k} = (k_1, \dots, k_M) \in I_{M,N}$. We may substitute $x_i = y_i + t_{k_i}$ for $i \in \{1, \dots, M\}$ in the defining formula (3.11), so that

$$(4.1) \quad \Theta_{\mathbf{k}}(\mathbf{t}) = \int_{C_{\mathbf{k}}^{\gamma}(\mathbf{0})} \frac{w_{\mathbf{k}}(\mathbf{t}_{\mathbf{k}} + \mathbf{y}; \mathbf{t})}{\Phi_{\mathbf{k}}(\mathbf{t})} \tilde{\mathcal{B}}(\mathbf{t}_{\mathbf{k}} + \mathbf{y}; \mathbf{t}) \Omega d^M \mathbf{y}.$$

Note that the integration over y_i is over the line segment $\gamma_i + \sqrt{-1}[0, \pi]$.

From Lemma 4.1 we see that, for any $j \in \{1, \dots, M\}$, the poles of the integrand as a function of y_j are avoided if the real part of every element of $P_{\mathbf{k};j}^+(\mathbf{t})$ (with $x_i = y_i + t_{k_i}$ for $i \neq j$) exceeds $\Re(t_{k_j} + y_j)$, and the real part of every element of $P_{\mathbf{k};j}^-(\mathbf{t})$ (with $x_i = y_i + t_{k_i}$ for $i \neq j$) is less than $\Re(t_{k_j} + y_j)$. Note that $\Re(y_i) = \Re(\gamma_i)$ for all $i \in \{1, \dots, N\}$. This yields the inequalities

$$\begin{aligned} |\Re(\gamma_j)| &< \Re(\ell_{k_j} \eta), \\ \Re(\gamma_j - \gamma_j) &> \Re(\eta), & \text{for } i < j \text{ and } k_i = k_j, \\ \Re(\gamma_j - \gamma_i) &> \Re(\eta), & \text{for } i > j \text{ and } k_i = k_j, \\ \Re(t_s - t_{k_j}) &> \Re(\gamma_j - \ell_s \eta), & \text{for } 1 \leq s < k_j, \\ \Re(t_{k_j} - t_s) &> \Re(-\gamma_j - \ell_s \eta), & \text{for } k_j < s \leq N, \\ \Re(t_s + t_{k_j}) &> \Re(-\gamma_j - \ell_s \eta), & \text{for } 1 \leq s \leq N, \\ \Re(t_{k_i} - t_{k_j}) &> \Re(\gamma_j - \gamma_i + \eta), & \text{for } i < j \text{ and } k_i < k_j, \\ \Re(t_{k_j} - t_{k_i}) &> \Re(\gamma_i - \gamma_j + \eta), & \text{for } i > j \text{ and } k_i > k_j, \\ \Re(t_{k_i} + t_{k_j}) &> \Re(-\gamma_i - \gamma_j + \eta), & \text{for } i \neq j, \\ \Re(t_{k_j}) &> \Re(-\tilde{\xi}_+ - \gamma_j), \\ \Re(t_{k_j}) &> \Re(-\tilde{\xi}_- - \gamma_j). \end{aligned}$$

The inequalities independent of \mathbf{t} are simple consequences of the condition $\gamma \in \Gamma_{M,N}^{\mathbf{k}}$. For the inequalities involving sums and differences of the t_s we also need $\Re(t_r - t_{r+1}) > 2\Lambda_+ + \max(\Re(\eta), 0)$ for $1 \leq r < N$, $\Re(t_N) > \Lambda_+ + \max(\Re(\frac{\eta}{2}), 0)$ and $\Re(t_N) > 0$. The final two inequalities rely on $\Re(t_N) > \Lambda_+ + \max(\Re(-\tilde{\xi}_+), \Re(-\tilde{\xi}_-))$.

Since the integration in (4.1) is over a compact set which is independent of \mathbf{t} the integrals define $V^{\ell}(M)$ -valued holomorphic functions $\Theta_{\mathbf{k}}$ on $\tilde{\mathbb{A}}$.

By virtue of Cauchy's integral theorem, these functions $\Theta_{\mathbf{k}}$ do not depend on $\gamma \in \Gamma_{M,N}^{\mathbf{k}}$ since $\Gamma_{M,N}^{\mathbf{k}}$ is path-connected and the separation of poles by the contours is unaltered for different choices of γ . We obtain part **a** of Theorem 3.3.

4.2. Integral solutions of the boundary qKZ equations. It is convenient to define $F_{\mathbf{k}}$ to be the $V^{\ell}(M)$ -valued meromorphic function $\Phi_{\mathbf{k}} \Theta_{\mathbf{k}}$ on $\tilde{\mathbb{A}}$. Consider the subsector

$$\begin{aligned} \tilde{\mathbb{A}}_{\tau} := \left\{ \mathbf{t} \in \mathbb{A} \mid \Re(t_s - t_{s+1}) > 2\Lambda_+ + \max(\Re(\eta), 0) - \Re(\tau) \text{ for } 1 \leq s < N, \right. \\ \left. \Re(t_N) > \Lambda_+ + \max(\Re(\frac{\eta}{2}), 0, \Re(-\tilde{\xi}_+), \Re(-\tilde{\xi}_-)) - \Re(\tau) \right\} \end{aligned}$$

of $\tilde{\mathbb{A}}$. Note that if $\mathbf{t} \in \tilde{\mathbb{A}}_{\tau} \subset \tilde{\mathbb{A}}$ then $\mathbf{t} + \tau \mathbf{e}_r \in \tilde{\mathbb{A}}$ for all $r \in \{1, \dots, N\}$ so that both sides of the boundary qKZ equations (2.13) with $\Psi = F_{\mathbf{k}}$ as given by (3.12) are well-defined on $\tilde{\mathbb{A}}_{\tau}$. Here we will show that they are equal on $\tilde{\mathbb{A}}_{\tau}$.

To show this we need to make additional assumption on the step size τ , which we can later remove by meromorphic continuation.

Note that the definition of the parameter set $\mathcal{D}_{M,N}^{\mathbf{k}}$ does not depend on τ . We can and will therefore restrict to parameters $(\ell, \eta) \in \mathcal{D}_{M,N}^{\mathbf{k}}$ and $\tau \in \mathbb{C}$ with $\Re(\tau) < 0$ satisfying the additional conditions

$$(4.2) \quad -\Re(\tau) \leq \min_{1 \leq r \leq N} \Re(\ell_r \eta) \text{ and } -\Re(\tau) < \min_{1 \leq r \leq N} \frac{\Re((2\ell_r + 1 - n_{\mathbf{k}}(r))\eta)}{n_{\mathbf{k}}(r) + 1}$$

(which is possible since $(\ell, \eta) \in \mathcal{D}_{M,N}^{\mathbf{k}}$). We also take the base point γ in the restricted set $\Gamma_{M,N;\tau}^{\mathbf{k}}$ consisting of the base points $\gamma \in \Gamma_{M,N}^{\mathbf{k}}$ satisfying

$$\begin{aligned} -\Re(\ell_{k_i} \eta + \tau) &< \Re(\gamma_i) < \Re(\ell_{k_i} \eta + \tau), \\ \Re(\gamma_{i(s+1;r)}) + \Re(\eta - \tau) &< \Re(\gamma_{i(s;r)}) \end{aligned}$$

for $1 \leq i \leq M$, $1 \leq r \leq N$ and $1 \leq s < n_{\mathbf{k}}(r)$. Note that $\Gamma_{M,N;\tau}^{\mathbf{k}}$ is nonempty as a consequence of $(\ell, \eta) \in \mathcal{D}_{M,N}^{\mathbf{k}}$ and (4.2).

We will use the proof [14, Section 7] of the analogous statement for Jackson integral solutions of the boundary qKZ equations. These were defined as summations over $\mathbf{x} \in \mathbf{x}_0 + \tau \mathbb{Z}^M$ for some base point \mathbf{x}_0 . Hence each variable x_j could be replaced by $x_j - \tau$ without affecting the overall value of the sum; such shifts provided a key step to the proof.

To mimic the proof for Jackson integrals we will shift integration variables by $-\tau$ in the integrals using Cauchy's theorem (recall that $\Re(\tau) < 0$). We use the following standard observation. Let $f(\mathbf{x})$ be a meromorphic function in $\mathbf{x} \in \mathbb{C}^M$ and view it, for fixed x_i ($i \neq j$), as meromorphic function in x_j . Suppose it is $\pi\sqrt{-1}$ -periodic in x_j and holomorphic for x_j in the vertical strip

$$(4.3) \quad S_\tau(z) := \{y \in \mathbb{C} \mid \Re(z) \leq \Re(x) \leq \Re(z - \tau)\}.$$

Then

$$\int_{z+[0,\pi]\sqrt{-1}} f(\mathbf{x}) dx_j = \int_{z+[0,\pi]\sqrt{-1}} f(\mathbf{x} - \mathbf{e}_j \tau) dx_j$$

by a direct application of Cauchy's theorem. So when in [14] a summation variable is shifted by $-\tau$ (this occurs in Lemma 7.6 in *ibid.*), we replace this by the procedure above based on the application of Cauchy's theorem.

The only other difference with the Jackson integral case arises because in the definition of $F_{\mathbf{k}}(\mathbf{t})$, the contours depend on \mathbf{t} . In other words, for $\mathbf{t} \in \tilde{\mathbb{A}}_\tau$ we have

$$F_{\mathbf{k}}(\mathbf{t} + \tau \mathbf{e}_r) = \int_{C_{\mathbf{k}}^\gamma(\mathbf{t} + \tau \mathbf{e}_r)} w_{\mathbf{k}}(\mathbf{x}; \mathbf{t} + \tau \mathbf{e}_r) \tilde{\mathcal{B}}(\mathbf{x}; \mathbf{t} + \tau \mathbf{e}_r) \Omega d^M \mathbf{x},$$

with the shift in τ possibly appearing in one of the integration contours as well as the integrand. If $r \neq k_j$ for all $j \in \{1, \dots, M\}$, then the shift in τ does not affect the contour, hence

$$(4.4) \quad F_{\mathbf{k}}(\mathbf{t} + \tau \mathbf{e}_r) = \int_{C_{\mathbf{k}}^\gamma(\mathbf{t})} w_{\mathbf{k}}(\mathbf{x}; \mathbf{t} + \tau \mathbf{e}_r) \tilde{\mathcal{B}}(\mathbf{x}; \mathbf{t} + \tau \mathbf{e}_r) \Omega d^M \mathbf{x}.$$

We first claim that (4.4) is also true if $r = k_j$ for some $j \in \{1, \dots, M\}$. We can apply the above procedure involving Cauchy's theorem (without the subsequent variable substitution) with $z = t_{k_j} + \gamma_j + \tau$. A subtlety arises if $n_{\mathbf{k}}(r) > 1$, in which case Cauchy's theorem needs to be successively applied for all integration variables x_j with $k_j = r$ starting with the variable x_j with $j = i_{\mathbf{k}}(1; r)$ and working our way up to $i_{\mathbf{k}}(n_{\mathbf{k}}(r); r)$. Hence, at each step we must assume that $x_i \in t_{k_i} + \gamma_i + [0, \pi]\sqrt{-1}$

if $k_i = k_j$ and $i < j$, and $x_i \in t_{k_i} + \gamma_i + \tau + [0, \pi]\sqrt{-1}$ if $k_i = k_j$ and $i > j$. To show that Cauchy's theorem indeed can be applied, we use the following lemma.

Lemma 4.2. *Let $\mathbf{k} \in I_{M,N}$. Let $(\ell, \eta) \in \mathcal{D}_{M,N}^{\mathbf{k}}$ and $\tau \in \mathbb{C}$ satisfying $\Re(\tau) < 0$ and (4.2). Choose $\gamma \in \Gamma_{M,N;\tau}^{\mathbf{k}}$. Let $1 \leq j \leq M$. Fix $\mathbf{t} \in \tilde{\mathbb{A}}_\tau$, $x_i \in t_{k_i} + \gamma_i + [0, \pi]\sqrt{-1}$ if $i < j$ and $x_i \in t_{k_i} + \gamma_i + \delta_{k_i, k_j} \tau + [0, \pi]\sqrt{-1}$ if $i > j$, then*

$$\frac{w_{\mathbf{k}}(\mathbf{x}; \mathbf{t} + \tau \mathbf{e}_{k_j})}{\Phi_{\mathbf{k}}(\mathbf{t} + \tau \mathbf{e}_{k_j})} \tilde{\mathcal{B}}(\mathbf{x}; \mathbf{t} + \tau \mathbf{e}_{k_j}) \Omega$$

as a function of x_j is $\pi\sqrt{-1}$ -periodic and has no poles in $S_\tau(t_{k_j} + \gamma_j + \tau)$.

Proof. The periodicity condition follows immediately from the analogous properties of $w_{\mathbf{k}}(\mathbf{x}; \mathbf{t})$ and $\tilde{\mathcal{B}}(\mathbf{x}; \mathbf{t})\Omega$. Using Lemma 4.1, the desired result on the poles holds if the real parts of all elements of $P_{\mathbf{k};j}^+(\mathbf{t} + \tau \mathbf{e}_{k_j})$ are strictly greater than $\Re(t_{k_j} + \gamma_j)$ and the real parts of all elements of $P_{\mathbf{k};j}^-(\mathbf{t} + \tau \mathbf{e}_{k_j})$ are strictly less than $\Re(t_{k_j} + \gamma_j + \tau)$. This imposes the inequalities

$$\begin{aligned} \Re(\gamma_j) &< \Re(\ell_{k_j} \eta + \tau), \\ -\Re(\gamma_j) &< \Re(\ell_{k_j} \eta), \\ \Re(\gamma_i - \gamma_j) &> \Re(\eta), & \text{for } i < j \text{ and } k_i = k_j, \\ \Re(\gamma_j - \gamma_i) &> \Re(\eta), & \text{for } i > j \text{ and } k_i = k_j, \\ \Re(t_s - t_{k_j}) &> \Re(\gamma_j - \ell_s \eta), & \text{for } 1 \leq s < k_j, \\ \Re(t_{k_j} - t_s) &> \Re(-\gamma_j - \ell_s \eta - \tau), & \text{for } k_j < s \leq N, \\ \Re(t_s + t_{k_j}) &> \Re(-\gamma_j - \ell_s \eta - \tau), & \text{for } 1 \leq s \leq N, \\ \Re(t_{k_i} - t_{k_j}) &> \Re(\gamma_j - \gamma_i + \eta), & \text{for } i < j \text{ and } k_i < k_j, \\ \Re(t_{k_j} - t_{k_i}) &> \Re(\gamma_i - \gamma_j + \eta - \tau), & \text{for } i > j \text{ and } k_i > k_j, \\ \Re(t_{k_i} + t_{k_j}) &> \Re(-\gamma_i - \gamma_j + \eta - \tau), & \text{for } i \neq j \text{ and } k_i \neq k_j, \\ \Re(2t_{k_j}) &> \Re(-\gamma_i - \gamma_j + \eta - \tau), & \text{for } i < j \text{ and } k_i = k_j, \\ \Re(2t_{k_j}) &> \Re(-\gamma_i - \gamma_j + \eta - 2\tau), & \text{for } i > j \text{ and } k_i = k_j, \\ \Re(t_{k_j}) &> \Re(-\tilde{\xi}_+ - \tau - \gamma_j), \\ \Re(t_{k_j}) &> \Re(-\tilde{\xi}_- - \tau - \gamma_j). \end{aligned}$$

The inequalities independent of \mathbf{t} immediately follow from $\gamma \in \Gamma_{M,N;\tau}^{\mathbf{k}}$ (the inequality $\Re(\gamma_j) < \Re(\ell_{k_j} \eta + \tau)$ is the only instance in this proof where we actually require $\gamma \in \Gamma_{M,N;\tau}^{\mathbf{k}}$ as opposed to $\gamma \in \Gamma_{M,N}^{\mathbf{k}}$). For the inequalities involving \mathbf{t} we also need the condition $|\Re(\gamma_j)| < \Re(\ell_{k_j} \eta)$ and in addition conditions implied by $\mathbf{t} \in \tilde{\mathbb{A}}_\tau$. Namely, the first three inequalities involving \mathbf{t} follow from the conditions $\Re(t_r - t_{r+1}) > 2\Lambda_+ - \Re(\tau)$ for $1 \leq r < N$ and $\Re(t_N) > 0$. The inequalities involving $t_{k_i} \pm t_{k_j}$ are a consequence of the conditions $\Re(t_r - t_{r+1}) > 2\Lambda_+ + \Re(\eta - \tau)$, $\Re(t_N) > \Lambda_+ + \Re(\frac{\eta}{2} - \tau)$, $\Re(t_N) > 0$ (and $\Re(\tau) < 0$), and the final two inequalities are a consequence of $\Re(t_N) > \Lambda_+ + \max(\Re(-\tilde{\xi}_+), \Re(-\tilde{\xi}_-)) - \Re(\tau)$. \square

Applying Lemma 4.2 for j running through $\{i \in \{1, \dots, M\} \mid k_i = r\}$ from low to high values, we obtain (4.4). Hence the two sides of the boundary qKZ equation (2.13) for $\Psi(\mathbf{t}) = F_{\mathbf{k}}(\mathbf{t})$ have the same contour $C_{\mathbf{k}}^\gamma(\mathbf{t})$ and differ in the integrands in the same way as the summands did at the start of the proof in [14, Section 7]. Thus we follow that proof taking care to correctly use Cauchy's theorem when dealing with replacements $x_j \rightarrow x_j - \tau$. We recall that the strategy is to shift the integration variable by τ in appropriate terms in (4.4); these terms are essentially due to the expansion of $\tilde{\mathcal{B}}(\mathbf{x}; \mathbf{t})$ by means of (3.4). Next, we use the conditions (4.5)

on the weight function $w_{\mathbf{k}}$ to establish that the terms on the left- and right-hand sides of (4.4) can be matched.

We need the following functional equations

$$\begin{aligned}
 F(x; \mathbf{t} + \tau \mathbf{e}_r) &= e^{4\ell_r \eta} \frac{\sinh(t_r \pm x - \ell_r \eta + \tau)}{\sinh(t_r \pm x + \ell_r \eta + \tau)} F(x; \mathbf{t}), \\
 F(x - \tau; \mathbf{t}) &= \left(\prod_{s=1}^N \frac{\sinh(t_s - x - \ell_s \eta + \tau)}{\sinh(t_s - x + \ell_s \eta + \tau)} \frac{\sinh(t_s + x + \ell_s \eta)}{\sinh(t_s + x - \ell_s \eta)} \right) F(x; \mathbf{t}), \\
 g(x - \tau) &= q^{-2} e^{-2(\tilde{\xi}_+ + \tilde{\xi}_-)} \frac{\sinh(x + \tilde{\xi}_+)}{\sinh(x - \tilde{\xi}_+ - \tau)} \frac{\sinh(x + \tilde{\xi}_-)}{\sinh(x - \tilde{\xi}_- - \tau)} g(x), \\
 h(x - \tau) &= e^{2\eta} \frac{\sinh(x - \tau)}{\sinh(x)} \frac{\sinh(x - \eta)}{\sinh(x + \eta - \tau)} h(x), \\
 u_{\mathbf{k};i}(x; \mathbf{t} + \tau \mathbf{e}_r) &= u_{\mathbf{k};i}(x; \mathbf{t}) \times \begin{cases} 1 & \text{if } r < k_i, \\ e^{2(\xi_+ + \xi_- - \eta) + 4(\sum_{s > k_i} \ell_s - M + i)\eta} & \text{if } r = k_i, \\ e^{-4\ell_r \eta} & \text{if } r > k_i, \end{cases} \\
 u_{\mathbf{k};i}(x - \tau; \mathbf{t}) &= q^2 e^{2(\tilde{\xi}_+ + \tilde{\xi}_-) - 4(M-i)\eta} u_{\mathbf{k};i}(x; \mathbf{t})
 \end{aligned}$$

where $k, r \in \{1, \dots, N\}$ and $i \in \{1, \dots, M\}$. From these and (2.18) one derives

$$\begin{aligned}
 \frac{w_{\mathbf{k}}(\mathbf{x}; \mathbf{t} + \tau \mathbf{e}_r)}{w_{\mathbf{k}}(\mathbf{x}; \mathbf{t})} &= \left(\prod_{i=1}^M \frac{\sinh(t_r \pm x_i - \ell_r \eta + \tau)}{\sinh(t_r \pm x_i + \ell_r \eta + \tau)} \right), \\
 \frac{w_{\mathbf{k}}(\mathbf{x} - \tau \mathbf{e}_j; \mathbf{t})}{w_{\mathbf{k}}(\mathbf{x}; \mathbf{t})} &= \left(\prod_{s=1}^N \frac{\sinh(t_s + x_j + \ell_s \eta)}{\sinh(t_s + x_j - \ell_s \eta)} \frac{\sinh(t_s - x_j - \ell_s \eta + \tau)}{\sinh(t_s - x_j + \ell_s \eta + \tau)} \right) \\
 &\quad \times \frac{\sinh(x_j + \tilde{\xi}_+)}{\sinh(x_j - \tilde{\xi}_+ - \tau)} \frac{\sinh(x_j + \tilde{\xi}_-)}{\sinh(x_j - \tilde{\xi}_- - \tau)} \\
 &\quad \times \left(\prod_{i \neq j} \frac{\sinh(x_j \pm x_i - \tau)}{\sinh(x_j \pm x_i)} \frac{\sinh(x_j \pm x_i - \eta)}{\sinh(x_j \pm x_i + \eta - \tau)} \right).
 \end{aligned} \tag{4.5}$$

We emphasize that these are the same equations as those satisfied by the weight function $w(\mathbf{x}; \mathbf{t})$ appearing in the Jackson integral in [14, Theorem 6.2]; in particular they do not depend on \mathbf{k} .

We denote $e_r \mathbf{t} = (t_1, \dots, t_{r-1}, -t_r, t_{r+1}, \dots, t_N)$ for $\mathbf{t} = (t_1, \dots, t_N) \in \mathbb{C}^N$ and $r \in \{1, \dots, N\}$ and $J^c = \{1, \dots, M\} \setminus J$ for $J \subseteq \{1, \dots, M\}$. Define, for $\epsilon \in \{\pm\}^M$, $J \subseteq \{1, \dots, M\}$, $r \in \{1, \dots, N\}$ and $\mathbf{x} \in C_{\mathbf{k}}^\gamma(\mathbf{t})$:

$$\begin{aligned}
 m_r^{\epsilon, J}(\mathbf{x}; \mathbf{t}) &:= (-1)^{\#J^c} \left(\prod_{i=1}^M \epsilon_i \sinh(\tilde{\xi}_- - \epsilon_i x_i) \prod_{s \neq r} \frac{\sinh(t_s - \epsilon_i x_i + \ell_s \eta)}{\sinh(t_s - \epsilon_i x_i - \ell_s \eta)} \right) \\
 &\quad \times \left(\prod_{i' \in J^c} \frac{1}{\sinh(t_r + \epsilon_{i'} x_{i'} + \ell_r \eta)} \right) \left(\prod_{\substack{i \in J \\ i' \in J^c}} \frac{\sinh(\pm x_i + \epsilon_{i'} x_{i'} + \eta)}{\sinh(\pm x_i + \epsilon_{i'} x_{i'})} \right) \\
 &\quad \times \left(\prod_{\substack{i < i' \\ i, i' \in J \text{ or } i, i' \in J^c}} \frac{\sinh(\epsilon_i x_i + \epsilon_{i'} x_{i'} + \eta)}{\sinh(\epsilon_i x_i + \epsilon_{i'} x_{i'})} \right),
 \end{aligned}$$

which defines a $\pi\sqrt{-1}$ -periodic function of each x_j with $j \in J^c$; this is the same function as in [14, Eqn. (7.8)].

We require some further notation. For $d \in \{0, \dots, M\}$ consider a subset $J \subseteq \{1, \dots, M\}$ of cardinality $M - d$. Write $J = \{i_1, \dots, i_{M-d}\}$ with $1 \leq i_1 < i_2 < \dots < i_{M-d} \leq M$. Then for given $\mathbf{k} = (k_1, \dots, k_M) \in I_{M,N}$, $\mathbf{x} = (x_1, \dots, x_M) \in C_{\mathbf{k}}^\gamma(\mathbf{t})$ and $\boldsymbol{\epsilon} = (\epsilon_1, \dots, \epsilon_M) \in \{\pm\}^M$, we denote $\mathbf{k}_J := (k_{i_1}, \dots, k_{i_{M-d}})$, $\mathbf{x}_J := (x_{i_1}, \dots, x_{i_{M-d}})$ and $\boldsymbol{\epsilon}_J := (\epsilon_{i_1}, \dots, \epsilon_{i_{M-d}})$. Note that $\mathbf{x}_J \in C_{\mathbf{k}_J}(\mathbf{t})$.

For the remainder of this subsection, we fix $r \in \{1, \dots, N\}$, $d \in \{0, \dots, M\}$, $J \subset \{1, \dots, M\}$ such that $\#J = M - d$ and the subtuples \mathbf{k}_J , \mathbf{x}_J and $\boldsymbol{\epsilon}_J$ as above. For $\boldsymbol{\epsilon}_{J^c} \in \{\pm\}^d$, set

$$\Lambda_{r, \boldsymbol{\epsilon}_{J^c}}(\mathbf{t}) := \int_{C_{\mathbf{k}_{J^c}}(\mathbf{t})} w_{\mathbf{k}}(\mathbf{x}; \mathbf{t}) m_r^{\boldsymbol{\epsilon}, J}(\mathbf{x}; e_r \mathbf{t}) d^d \mathbf{x}_{J^c},$$

$$\Upsilon_{r, \boldsymbol{\epsilon}_{J^c}}(\mathbf{t}) := \int_{C_{\mathbf{k}_{J^c}}(\mathbf{t})} w_{\mathbf{k}}(\mathbf{x}; \mathbf{t}) m_r^{\boldsymbol{\epsilon}, J}(\mathbf{x}; \mathbf{t} + \tau e_r) d^d \mathbf{x}_{J^c}.$$

They are the analogons of their namesakes introduced in [14, Sec. 7.3] with the summation over $\mathbf{x}_{J^c} \in \tau\mathbb{Z}^d$ replaced by an integral over $\mathbf{x}_{J^c} \in C_{\mathbf{k}_{J^c}}(\mathbf{t})$.

A careful inspection of the proof in [14, Section 7] shows that the only statement which does not immediately generalize from the Jackson integral case to the integral case is in fact in the proof of [14, Lemma 7.6] where in the Jackson integral a certain variable x_j in the summand is replaced by $x_j - \tau$. In order to adapt that proof to one suitable for our integral solutions, we apply the argument based on Cauchy's theorem again. Hence, the following lemma replaces the start of the proof of [14, Lemma 7.6].

Lemma 4.3. *Let $\mathbf{k} \in I_{M,N}$. Let $(\ell, \eta) \in \mathcal{D}_{M,N}^{\mathbf{k}}$ and $\tau \in \mathbb{C}$ satisfying $\Re(\tau) < 0$ and (4.2). Choose $\boldsymbol{\gamma} \in \Gamma_{M,N;\tau}^{\mathbf{k}}$. Let $j \in J^c$ such that $\epsilon_j = +$. For fixed $\mathbf{t} \in \tilde{\mathbb{A}}_\tau$ and $x_i \in t_{k_i} + \gamma_i + [0, \pi]\sqrt{-1}$ ($i \neq j$), the functions*

$$x_j \mapsto w_{\mathbf{k}}(\mathbf{x}; \mathbf{t}) m_r^{\boldsymbol{\epsilon}, J}(\mathbf{x}; e_r \mathbf{t}),$$

$$x_j \mapsto w_{\mathbf{k}}(\mathbf{x}; \mathbf{t}) m_r^{\boldsymbol{\epsilon}, J}(\mathbf{x}; \mathbf{t} + \tau e_r)$$

are $\pi\sqrt{-1}$ -periodic and have no poles in $S_\tau(t_{k_j} + \gamma_j)$.

Proof. The periodicity statement follows from the fact that both $x_j \mapsto w_{\mathbf{k}}(\mathbf{x}; \mathbf{t})$ and $x_j \mapsto m_r^{\boldsymbol{\epsilon}, J}(\mathbf{x}; \mathbf{t})$ are $\pi\sqrt{-1}$ -periodic for any \mathbf{t} .

Using that $\epsilon_j = +$ we see that the poles of $m_r^{\boldsymbol{\epsilon}, J}(\mathbf{x}; e_r \mathbf{t})$ as a function of x_j are contained in the set

$$\left(\{t_s - \ell_s \eta \mid 1 \leq s \leq N\} \cup \{\pm x_i \mid i \neq j\} \right) + \pi\sqrt{-1}\mathbb{Z}.$$

Similarly, the poles of $m_r^{\boldsymbol{\epsilon}, J}(\mathbf{x}; \mathbf{t} + \tau e_r)$ as a function of x_j are contained in the set

$$\left(\{t_s - \ell_s \eta \mid 1 \leq s \leq N, s \neq r\} \cup \{-t_r - \tau - \ell_r \eta\} \cup \{\pm x_i \mid i \neq j\} \right) + \pi\sqrt{-1}\mathbb{Z}.$$

Considering (3.8), we see that all x_j -dependent poles are cancelled by zeros of $w_{\mathbf{k}}(\mathbf{x}; \mathbf{t})$, as well as the poles of the form $t_s - \ell_s \eta$ for $s < k_j$. It follows that all poles of $w_{\mathbf{k}}(\mathbf{x}; \mathbf{t}) m_r^{\boldsymbol{\epsilon}, J}(\mathbf{x}; e_r \mathbf{t})$ and $w_{\mathbf{k}}(\mathbf{x}; \mathbf{t}) m_r^{\boldsymbol{\epsilon}, J}(\mathbf{x}; \mathbf{t} + \tau e_r)$ are contained in

$$\left((Q_{\mathbf{k};j}^+ - \tau\mathbb{Z}_{\geq 0}) \cup (Q_{\mathbf{k};j}^- + \tau\mathbb{Z}_{\geq 0}) \right) + \pi\sqrt{-1}\mathbb{Z},$$

where

$$\begin{aligned} Q_{\mathbf{k};j}^+(\mathbf{t}) &= \{t_s + \ell_s \eta\}_{s \leq k_j} \cup \{x_i - \eta\}_{i < j} \\ Q_{\mathbf{k};j}^-(\mathbf{t}) &= \{t_s - \ell_s \eta\}_{s \geq k_j} \cup \{-t_s - \ell_s \eta - \tau\}_s \cup \{-\tilde{\xi}_+, -\tilde{\xi}_-\} \cup \\ &\quad \cup \{x_i + \eta\}_{i > j} \cup \{-x_i + \eta\}_{i \neq j}. \end{aligned}$$

The conclusion of the lemma is justified if the real parts of all elements of $Q_{\mathbf{k};j}^+$ are strictly greater than $\Re(t_{k_j} + \gamma_j - \tau)$ and the real parts of all elements of $Q_{\mathbf{k};j}^-$ are strictly less than $\Re(t_{k_j} + \gamma_j)$. This yields the inequalities

$$\begin{aligned} \Re(\gamma_j) &> -\Re(\ell_{k_j} \eta), \\ \Re(\gamma_j) &< \Re(\ell_{k_j} \eta + \tau), \\ \Re(\gamma_i - \gamma_j) &> \Re(\eta - \tau), & \text{for } i < j \text{ and } k_i = k_j, \\ \Re(\gamma_j - \gamma_i) &> \Re(\eta - \tau), & \text{for } i > j \text{ and } k_i = k_j, \\ \Re(t_s - t_{k_j}) &> \Re(\gamma_j - \ell_s \eta - \tau), & \text{for } 1 \leq s < k_j, \\ \Re(t_{k_j} - t_s) &> \Re(-\gamma_j - \ell_s \eta), & \text{for } k_j < s \leq N, \\ \Re(t_s + t_{k_j}) &> \Re(-\gamma_j - \ell_s \eta - \tau), & \text{for } 1 \leq s \leq N, \\ \Re(t_{k_i} - t_{k_j}) &> \Re(\gamma_j - \gamma_i + \eta - \tau), & \text{for } i < j \text{ and } k_i < k_j, \\ \Re(t_{k_j} - t_{k_i}) &> \Re(\gamma_i - \gamma_j + \eta), & \text{for } i > j \text{ and } k_i > k_j, \\ \Re(t_{k_i} + t_{k_j}) &> \Re(-\gamma_i - \gamma_j + \eta), & \text{for } i \neq j \text{ and } k_i \neq k_j, \\ \Re(2t_{k_j}) &> \Re(-\gamma_i - \gamma_j + \eta), & \text{for } i \neq j \text{ and } k_i = k_j, \\ \Re(t_{k_j}) &> \Re(-\tilde{\xi}_+ - \gamma_j), \\ \Re(t_{k_j}) &> \Re(-\tilde{\xi}_- - \gamma_j). \end{aligned}$$

We get similar conditions as in the proof of Lemma 4.2. However, the inequality for the differences on γ_j is stronger as $\Re(\tau) < 0$. All these conditions are again a consequence of $\gamma \in \Gamma_{M,N;\tau}^{\mathbf{k}}$, $\Re(\tau) < 0$ and $\mathbf{t} \in \tilde{\mathbb{A}}_\tau$. \square

In Fig. 1 we illustrate a typical arrangement of poles with respect to the vertical strip $S_\tau(t_{k_i} + \gamma_i)$ as pertains to Lemma 4.3.

Let $\mathbf{k} \in I_{M,N}$ and $(\ell, \eta) \in \mathcal{D}_{M,N}^{\mathbf{k}}$. Following [14] it follows that for τ satisfying $\Re(\tau) < 0$ and (4.2) that

$$(4.6) \quad F_{\mathbf{k}}(\mathbf{t} + \tau \mathbf{e}_r) = \mathcal{A}_r(\mathbf{t}) F_{\mathbf{k}}(\mathbf{t}), \quad 1 \leq r \leq N$$

as meromorphic functions in $\mathbf{t} \in \tilde{\mathbb{A}}_\tau$.

Next we show that the $V^\ell(M)$ -valued meromorphic function $F_{\mathbf{k}} = \Phi_{\mathbf{k}} \Theta_{\mathbf{k}}$ on $\tilde{\mathbb{A}}_\tau$ uniquely extends to a meromorphic $V^\ell(M)$ -valued meromorphic function $\Psi_{\mathbf{k}}$ on \mathbb{C}^N satisfying (2.13).

Let $\{\mathcal{A}_\alpha(\mathbf{t})\}_{\alpha \in \mathbb{Z}^N}$ be the unique family of linear operators on V^ℓ depending meromorphically on $\mathbf{t} \in \mathbb{C}^N$ satisfying

$$\begin{aligned} \mathcal{A}_\alpha(\mathbf{t} + \beta \tau) \mathcal{A}_\beta(\mathbf{t}) &= \mathcal{A}_{\alpha+\beta}(\mathbf{t}), & \alpha, \beta \in \mathbb{Z}^N, \\ \mathcal{A}_0(\mathbf{t}) &= \text{Id}_{V^\ell}, \\ \mathcal{A}_{\mathbf{e}_r}(\mathbf{t}) &= \mathcal{A}_r(\mathbf{t}), & r = 1, \dots, N, \end{aligned}$$

with $\mathcal{A}_r(\mathbf{t})$ the transport operators (2.14) of the boundary qKZ equations (the compatibility of the transport operators guarantees the existence of the cocycle $\{\mathcal{A}_\alpha(\mathbf{t})\}_{\alpha \in \mathbb{Z}^N}$).




FIGURE 1. Integration contour $t_1 + \gamma_1 + [0, \pi]\sqrt{-1}$ and t_1 -dependent poles of $x_1 \mapsto w_{\mathbf{k}}(\mathbf{x}; \mathbf{t})\tilde{\mathcal{B}}(\mathbf{x}; \mathbf{t})\Omega$ for $k_1 = 1$ and γ_1 satisfying $|\Re(\gamma_1)| < \Re(\ell_{k_1}\eta + \tau)$ as in the proof of Lemma 4.3. For each pole sequence, the pole closest to the vertical strip $S_\tau(t_1 + \gamma_1)$ (4.3), which is the shaded area in the figure, is indicated. The poles consist of (unilateral) sequences entirely to the left or entirely to the right of $S_\tau(t_1 + \gamma_1)$. In the vertical strip we have marked the closed contour with respect to which Cauchy's theorem is used in this subsection.

Consider the cone

$$P_+ := \{\lambda \in \mathbb{Z}^N \mid \lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_N \geq 0\}.$$

Note that $\mathbf{t} - \lambda\tau \in \tilde{\mathbb{A}}_\tau$ for $\mathbf{t} \in \tilde{\mathbb{A}}_\tau$ and $\lambda \in P_+$. It follows from (4.6) that for all $\lambda \in P_+$,

$$(4.7) \quad F_{\mathbf{k}}(\mathbf{t} - \lambda\tau) = \mathcal{A}_{-\lambda}(\mathbf{t})F_{\mathbf{k}}(\mathbf{t})$$

as meromorphic $V^\ell(M)$ -valued functions in $\mathbf{t} \in \tilde{\mathbb{A}}_\tau$.

If $\Psi_{\mathbf{k}}$ is a $V^\ell(M)$ -valued meromorphic function on \mathbb{C}^N satisfying the boundary qKZ equations and coinciding with $F_{\mathbf{k}}$ on $\tilde{\mathbb{A}}_\tau$, then

$$(4.8) \quad \Psi_{\mathbf{k}}(\mathbf{t}) = \mathcal{A}_\alpha(\mathbf{t} - \alpha\tau)F_{\mathbf{k}}(\mathbf{t} - \alpha\tau)$$

for $\mathbf{t} \in \mathbb{C}^N$ and $\alpha \in \mathbb{Z}^N$ such that $\mathbf{t} - \alpha\tau \in \tilde{\mathbb{A}}_\tau$. Now (4.8) can be used to prove the existence of the meromorphic extension of $F_{\mathbf{k}}$ to \mathbb{C}^N . For this we need to show that the right hand side of (4.8) does not depend on the choice of α such that $\mathbf{t} - \alpha\tau \in \tilde{\mathbb{A}}_\tau$.

Suppose $\beta \in \mathbb{Z}^N$ also satisfies $\mathbf{t} - \beta\tau \in \tilde{\mathbb{A}}_\tau$. Let $\lambda, \mu \in P_+$ such that

$$\lambda + \alpha = \mu + \beta \in P_+.$$

Then repeated application of (4.7) gives

$$\begin{aligned}\mathcal{A}_\beta(\mathbf{t} - \beta\tau)F_{\mathbf{k}}(\mathbf{t} - \beta\tau) &= \mathcal{A}_\beta(\mathbf{t} - \beta\tau)\mathcal{A}_{-\mu}(\mathbf{t} - \beta\tau)^{-1}F_{\mathbf{k}}(\mathbf{t} - (\mu + \beta)\tau) \\ &= \mathcal{A}_\beta(\mathbf{t} - \beta\tau)\mathcal{A}_{-\mu}(\mathbf{t} - \beta\tau)^{-1}\mathcal{A}_{-\lambda}(\mathbf{t} - \alpha\tau)F_{\mathbf{k}}(\mathbf{t} - \alpha\tau) \\ &= \mathcal{A}_\alpha(\mathbf{t} - \alpha\tau)F_{\mathbf{k}}(\mathbf{t} - \alpha\tau),\end{aligned}$$

where the last equality follows from the cocycle property. By a similar computation one shows that the resulting $V^\ell(M)$ -valued meromorphic function $\Psi_{\mathbf{k}}$ on \mathbb{C}^N satisfies the boundary qKZ equations (2.13).

The identity $\Psi_{\mathbf{k}} = F_{\mathbf{k}}$ on $\tilde{\mathbb{A}}_\tau$ extends to $\tilde{\mathbb{A}}$ by meromorphic continuation. Finally, the extra assumption (4.2) on the step size can be removed by meromorphic continuation. This completes the proof of part **b** of Theorem 3.3.

4.3. Asymptotics. In this subsection we prove part **c** of Theorem 3.3. Recall the notation $\mathbf{t}_{\mathbf{k}} := (t_{k_1}, \dots, t_{k_M}) \in \mathbb{C}^M$ for $\mathbf{t} \in \mathbb{C}^N$, $\mathbf{k} \in I_{M,N}$. We will investigate the asymptotics of $\Psi_{\mathbf{k}}(\mathbf{t})$ as $\mathbf{t} \xrightarrow{\mathbb{A}} \infty$ using the formula (4.1). It is convenient to study first the asymptotics of $e^{\sum_i t_{k_i} \frac{w_{\mathbf{k}}(\mathbf{t}_{\mathbf{k}} + \mathbf{y}; \mathbf{t})}{\Phi_{\mathbf{k}}(\mathbf{t})}}$ and of

$$e^{-\sum_i t_{k_i}} \tilde{\mathcal{B}}(\mathbf{t}_{\mathbf{k}} + \mathbf{y}; \mathbf{t}) \Omega = e^{-\sum_i t_{k_i}} \sum_{\mathbf{m} \in I_{M,N}} \beta_{\mathbf{m}}(\mathbf{t}_{\mathbf{k}} + \mathbf{y}; \mathbf{t}) \Omega_{\mathbf{m}}.$$

Lemma 4.4. *Let $\mathbf{k} \in I_{M,N}$.*

a. *There exists unique continuous functions $\Delta_{\alpha}^{\mathbf{k}}$ on $C_{\mathbf{k}}^{\gamma}(\mathbf{0})$ ($\alpha \in Q_+$) such that*

$$e^{\sum_{i=1}^M t_{k_i} \frac{w_{\mathbf{k}}(\mathbf{t}_{\mathbf{k}} + \mathbf{y}; \mathbf{t})}{\Phi_{\mathbf{k}}(\mathbf{t})}} = \sum_{\alpha \in Q_+} \Delta_{\alpha}^{\mathbf{k}}(\mathbf{y}) e^{-2(\alpha, \mathbf{t})}$$

for $(\mathbf{y}, \mathbf{t}) \in C_{\mathbf{k}}^{\gamma}(\mathbf{0}) \times \tilde{\mathbb{A}}$, with the series converging normally on compact sets. Furthermore,

$$\begin{aligned}\Delta_0^{\mathbf{k}}(\mathbf{y}) &= \prod_{i=1}^M \frac{\theta(e^{2(y_i + \psi_{\mathbf{k}; i})})}{(e^{2(y_i - \ell_{k_i} \eta)}, q^2 e^{-2(y_i + \ell_{k_i} \eta)}; q^2)_{\infty}} \\ &\quad \times \prod_{\substack{1 \leq i < j \leq M \\ k_i = k_j}} (1 - e^{2(y_j - y_i)}) \frac{(q^2 e^{2(y_j - y_i - \eta)}; q^2)_{\infty}}{(e^{2(y_j - y_i + \eta)}; q^2)_{\infty}}.\end{aligned}$$

b. *Let $\mathbf{m} \in I_{M,N}$. There exists unique continuous functions $f_{\alpha}^{\mathbf{k}, \mathbf{m}}$ on $C_{\mathbf{k}}^{\gamma}(\mathbf{0})$ ($\alpha \in Q_+$) such that*

$$e^{-\sum_{i=1}^M t_{k_i}} \beta_{\mathbf{m}}(\mathbf{t}_{\mathbf{k}} + \mathbf{y}; \mathbf{t}) = \sum_{\alpha \in Q_+} f_{\alpha}^{\mathbf{k}, \mathbf{m}}(\mathbf{y}) e^{-(\alpha, \mathbf{t})}$$

for $(\mathbf{y}, \mathbf{t}) \in C_{\mathbf{k}}^{\gamma}(\mathbf{0}) \times \tilde{\mathbb{A}}$, with the series converging normally on compact sets. Furthermore,

$$f_0^{\mathbf{k}, \mathbf{m}}(\mathbf{y}) = \delta_{\mathbf{k}, \mathbf{m}} \prod_{i=1}^M \frac{-e^{\xi_- + 2 \sum_{s < k_i} \ell_s \eta}}{1 - e^{-2(y_i + \ell_{k_i} \eta)}}.$$

Proof. **a.** This follows from the explicit expression

$$\begin{aligned}
& e^{\sum_{i=1}^M t_{k_i}} \frac{w_{\mathbf{k}}(\mathbf{t}_{\mathbf{k}} + \mathbf{y}; \mathbf{t})}{\Phi_{\mathbf{k}}(\mathbf{t})} = \\
& = \prod_{i=1}^M \left\{ \left(\prod_{s < k_i} \frac{(e^{2(y_i + t_{k_i} - t_s + \ell_s \eta)}; q^2)_{\infty}}{(e^{2(y_i + t_{k_i} - t_s - \ell_s \eta)}; q^2)_{\infty}} \right) \frac{\theta(e^{2(y_i + \psi_{\mathbf{k}; i})})}{(e^{2(y_i - \ell_{k_i} \eta)}, q^2 e^{-2(y_i + \ell_{k_i} \eta)}; q^2)_{\infty}} \right. \\
& \quad \times \left(\prod_{s > k_i} \frac{(q^2 e^{2(t_s - t_{k_i} - y_i + \ell_s \eta)}; q^2)_{\infty}}{(q^2 e^{2(t_s - t_{k_i} - y_i - \ell_s \eta)}; q^2)_{\infty}} \right) \left(\prod_{s=1}^N \frac{(e^{-2(t_s + t_{k_i} + y_i - \ell_s \eta)}; q^2)_{\infty}}{(e^{-2(t_s + t_{k_i} + y_i + \ell_s \eta)}; q^2)_{\infty}} \right) \\
& \quad \times \left. \frac{(q^2 e^{2(\tilde{\xi}_+ - t_{k_i} - y_i)}, q^2 e^{2(\tilde{\xi}_- - t_{k_i} - y_i)}; q^2)_{\infty}}{(e^{-2(\tilde{\xi}_+ + t_{k_i} + y_i)}, e^{-2(\tilde{\xi}_- + t_{k_i} + y_i)}; q^2)_{\infty}} \right\} \\
& \quad \times \prod_{1 \leq i < j \leq M} \left\{ (1 - e^{-2((y_i + t_{k_i}) \pm (y_j + t_{k_j}) + \eta)}) \frac{(q^2 e^{-2((y_i + t_{k_i}) \pm (y_j + t_{k_j}) + \eta)}; q^2)_{\infty}}{(e^{-2((y_i + t_{k_i}) \pm (y_j + t_{k_j}) - \eta)}; q^2)_{\infty}} \right\}.
\end{aligned}$$

b. We use the explicit expression from Proposition 3.1. It gives

$$\begin{aligned}
& e^{-\sum_i t_{k_i}} \beta_{\mathbf{m}}(\mathbf{t}_{\mathbf{k}} + \mathbf{y}; \mathbf{t}) = \\
& e^{\sum_i \left(\frac{n_{\mathbf{m}}(m_i)}{2} - \ell_{m_i} \right) \eta} \sum_{\mathbf{p} \in S_M \mathbf{m}} \sum_{\boldsymbol{\epsilon} \in \{\pm\}^M} \left\{ \prod_{i=1}^M \left\{ \frac{\epsilon_i e^{-t_{k_i}} \sinh(\epsilon_i t_{k_i} + \epsilon_i y_i - \tilde{\xi}_-)}{\sinh(t_{p_i} + \epsilon_i t_{k_i} + \epsilon_i y_i - \ell_{p_i} \eta)} \right. \right. \\
(4.9) \quad & \times \left(\prod_{s > p_i} \frac{\sinh(t_s + \epsilon_i t_{k_i} + \epsilon_i y_i + \ell_s \eta)}{\sinh(t_s + \epsilon_i t_{k_i} + \epsilon_i y_i - \ell_s \eta)} \right) \left(\prod_{s=1}^N \frac{\sinh(t_s - \epsilon_i y_i - \epsilon_i t_{k_i} + \ell_s \eta)}{\sinh(t_s - \epsilon_i y_i - \epsilon_i t_{k_i} - \ell_s \eta)} \right) \Big\} \\
& \times \left(\prod_{1 \leq i < j \leq M} \frac{\sinh(\epsilon_i t_{k_i} + \epsilon_j t_{k_j} + \epsilon_i y_i + \epsilon_j y_j + \eta)}{\sinh(\epsilon_i t_{k_i} + \epsilon_j t_{k_j} + \epsilon_i y_i + \epsilon_j y_j)} \right) \\
& \times \left. \left(\prod_{\substack{1 \leq i < j \leq M \\ p_i < p_j}} \frac{\sinh(\epsilon_i t_{k_i} - \epsilon_j t_{k_j} + \epsilon_i y_i - \epsilon_j y_j - \eta)}{\sinh(\epsilon_i t_{k_i} - \epsilon_j t_{k_j} + \epsilon_i y_i - \epsilon_j y_j)} \right) \right\}.
\end{aligned}$$

For fixed $\mathbf{p} \in S_M \mathbf{m}$ and $\boldsymbol{\epsilon} \in \{\pm\}^M$, a direct computation shows that

$$\prod_{i=1}^M \frac{\epsilon_i e^{-t_{k_i}} \sinh(\epsilon_i t_{k_i} + \epsilon_i y_i - \tilde{\xi}_-)}{\sinh(t_{p_i} + \epsilon_i t_{k_i} + \epsilon_i y_i - \ell_{p_i} \eta)} = \sum_{\alpha \in Q_+} d_{\alpha}^{\mathbf{p}, \boldsymbol{\epsilon}}(\mathbf{y}) e^{-(\alpha, \mathbf{t})}$$

with leading coefficient

$$d_0^{\mathbf{p}, \boldsymbol{\epsilon}}(\mathbf{y}) = \delta_{\mathbf{p}, \mathbf{k}} \delta_{\boldsymbol{\epsilon}, (-)^M} \prod_{i=1}^M \frac{-e^{\tilde{\xi}_- - \ell_{k_i} \eta}}{1 - e^{-2(y_i + \ell_{k_i} \eta)}}$$

and $(-)^M := (-, \dots, -)$ the M -tuple of minus signs. Here the convergence is as indicated in the lemma. For all the other factors in the right hand side of (4.9) it is easy to compute the series expansion when $\mathbf{t} \xrightarrow{\mathbb{A}} \infty$. It leads to the result that

$$e^{-\sum_{i=1}^M t_{k_i}} \beta_{\mathbf{m}}(\mathbf{t}_{\mathbf{k}} + \mathbf{y}; \mathbf{t}) = \sum_{\alpha \in Q_+} f_{\alpha}^{\mathbf{k}, \mathbf{m}}(\mathbf{y}) e^{-(\alpha, \mathbf{t})}$$

with

$$f_0^{\mathbf{k}, \mathbf{m}}(\mathbf{y}) = \delta_{\mathbf{k}, \mathbf{m}} \left(\prod_{i=1}^M \frac{-e^{\tilde{\xi}_- + \left(\frac{n_{\mathbf{k}}(k_i)}{2} + 2 \sum_{s < k_i} \ell_s \eta\right) \eta}}{1 - e^{-2(y_i + \ell_{k_i} \eta)}} \right) e^{(\#\{(i, j) \mid k_i < k_j\} - \#\{(i, j) \mid i < j\}) \eta}$$

and the convergence as indicated in the lemma. Finally note that the expression of $f_0^{\mathbf{k}, \mathbf{m}}(\mathbf{y})$ coincides with the expression as given in the lemma because

$$\begin{aligned} \#\{(i, j) \mid k_i < k_j\} - \#\{(i, j) \mid i < j\} &= -\#\{(i, j) \mid i < j, k_i = k_j\} \\ &= \frac{M - \#\{(i, j) \mid k_i = k_j\}}{2} = \sum_{i=1}^M \frac{1 - n_{\mathbf{k}}(k_i)}{2}, \end{aligned}$$

where we have used that $k_i \leq k_j$ if $i < j$. \square

From Lemma 4.4 we immediately deduce that there exists unique continuous $V^\ell(M)$ -valued functions $R_\alpha^{\mathbf{k}}$ on $C_{\mathbf{k}}^\gamma(\mathbf{0})$ ($\alpha \in Q_+$) such that

$$(4.10) \quad \frac{w_{\mathbf{k}}(\mathbf{t}_{\mathbf{k}} + \mathbf{y}; \mathbf{t})}{\Phi_{\mathbf{k}}(\mathbf{t})} \tilde{\mathcal{B}}(\mathbf{t}_{\mathbf{k}} + \mathbf{y}; \mathbf{t}) \Omega = \sum_{\alpha \in Q_+} R_\alpha^{\mathbf{k}}(\mathbf{y}) e^{-(\alpha, \mathbf{t})}$$

for $(\mathbf{y}, \mathbf{t}) \in C_{\mathbf{k}}^\gamma(\mathbf{0}) \times \tilde{\mathbb{A}}$ with the $V^\ell(M)$ -valued series converging normally on compact sets, and with leading coefficient given by

$$\begin{aligned} R_0^{\mathbf{k}}(\mathbf{y}) &= \sum_{\mathbf{m} \in I_{M, N}} \Delta_0^{\mathbf{k}}(\mathbf{y}) f_0^{\mathbf{k}, \mathbf{m}}(\mathbf{y}) \Omega_{\mathbf{m}} \\ &= \Delta_0^{\mathbf{k}}(\mathbf{y}) \left(\prod_{i=1}^M \frac{-e^{\xi_- + 2 \sum_{s < k_i} \ell_s \eta}}{1 - e^{-2(y_i + \ell_{k_i} \eta)}} \right) \Omega_{\mathbf{k}} \\ &= \left(\prod_{i=1}^M \frac{-e^{\xi_- + 2 \sum_{s < k_i} \ell_s \eta} \theta(e^{2(y_i + \psi_{\mathbf{k}; i})})}{(e^{2(\pm y_i - \ell_{k_i} \eta)}; q^2)_\infty} \right) \\ &\quad \times \left(\prod_{\substack{1 \leq i < j \leq M \\ k_i = k_j}} (1 - e^{2(y_j - y_i)}) \frac{(q^2 e^{2(y_j - y_i - \eta)}; q^2)_\infty}{(e^{2(y_j - y_i + \eta)}; q^2)_\infty} \right) \Omega_{\mathbf{k}}. \end{aligned}$$

It follows from (4.1), (4.10) and Fubini's theorem that $\Theta_{\mathbf{k}}(\mathbf{t})$ has a $V^\ell(M)$ -valued series expansion

$$\Theta_{\mathbf{k}}(\mathbf{t}) = \sum_{\alpha \in Q_+} L_\alpha^{\mathbf{k}} e^{-(\alpha, \mathbf{t})}$$

normally converging for \mathbf{t} in compact subsets of $\tilde{\mathbb{A}}$ and with the coefficients given by

$$L_\alpha^{\mathbf{k}} = \int_{C_{\mathbf{k}}^\gamma(\mathbf{0})} R_\alpha^{\mathbf{k}}(\mathbf{y}) d^M \mathbf{y}, \quad \alpha \in Q_+.$$

By the explicit expression of $R_0^{\mathbf{k}}(\mathbf{y})$ we have in particular

$$L_0^{\mathbf{k}} = \mu_{\mathbf{k}} \Omega_{\mathbf{k}}$$

with $\mu_{\mathbf{k}}$ the explicit integral given by

$$(4.11) \quad \mu_{\mathbf{k}} = \int_{C_{\mathbf{k}}^{\gamma}(\mathbf{0})} \left(\prod_{i=1}^M \frac{-e^{\xi_- + 2 \sum_{s < k_i} \ell_s \eta} \theta(e^{2(y_i + \psi_{\mathbf{k}; i})})}{(e^{2(\pm y_i - \ell_{k_i} \eta)}; q^2)_{\infty}} \right) \times \left(\prod_{\substack{1 \leq i < j \leq M \\ k_i = k_j}} (1 - e^{2(y_j - y_i)}) \frac{(q^2 e^{2(y_j - y_i - \eta)}; q^2)_{\infty}}{(e^{2(y_j - y_i + \eta)}; q^2)_{\infty}} \right) d^M \mathbf{y}.$$

So to complete the proof of part c of Theorem 3.3, it suffices to prove the following explicit evaluation formula for the integral $\mu_{\mathbf{k}}$.

Lemma 4.5. *Let $\mathbf{k} \in I_{M,N}$, $(\ell, \eta) \in \mathcal{D}_{M,N}^{\mathbf{k}}$ and $\tau \in \mathbb{C}$ with $\Re(\tau) < 0$. Let $\gamma \in \Gamma_{M,N}^{\mathbf{k}}$. Then $\mu_{\mathbf{k}} = \nu_{\mathbf{k}}$, with $\nu_{\mathbf{k}}$ given by (3.14).*

Proof. Fix $\mathbf{k} \in I_{M,N}$. Recall the notation $i_{\mathbf{k}}(m; r)$ from (3.10). It follows that

$$\psi_{\mathbf{k}; i(m; r)} = \omega_{\mathbf{k}; r} + \frac{\tau}{2} + (2m - n_{\mathbf{k}}(r) - 1)\eta,$$

with $\omega_{\mathbf{k}; r}$ given by (3.15).

Let $r \in \{1, \dots, N\}$. We introduce the meromorphic function $f_r^{\mathbf{k}}$ on $\mathbb{C}^{n_{\mathbf{k}}(r)}$ by

$$f_r^{\mathbf{k}}(z) := \left(\prod_{m=1}^{n(r)} \frac{\theta(q e^{2(z_m + \omega_{\mathbf{k}; r} + (2m - n(r) - 1)\eta)})}{(e^{2(\pm z_m - \ell_r \eta)}; q^2)_{\infty}} \right) \times \left(\prod_{1 \leq m < m' \leq n_{\mathbf{k}}(r)} (1 - e^{2(z_{m'} - z_m)}) \frac{(q^2 e^{2(z_{m'} - z_m - \eta)}; q^2)_{\infty}}{(e^{2(z_{m'} - z_m + \eta)}; q^2)_{\infty}} \right)$$

and the cycles

$$\mathcal{C}_{(r)} := (\gamma_{i(1; r)} + \sqrt{-1}[0, \pi]) \times \dots \times (\gamma_{i(n_{\mathbf{k}}(r); r)} + \sqrt{-1}[0, \pi]).$$

Define

$$(4.12) \quad \mu_r^{\mathbf{k}} := \int_{\mathcal{C}_{(r)}} f_r^{\mathbf{k}}(z) d^{n_{\mathbf{k}}(r)} z$$

$$(4.13) \quad \nu_r^{\mathbf{k}} := \prod_{m=1}^{n_{\mathbf{k}}(r)} (-\pi \sqrt{-1}) \frac{(q^2 e^{-2m\eta}, q e^{2((m-1-\ell_r)\eta \pm \omega_{\mathbf{k}; r})}; q^2)_{\infty}}{(q^2, q^2 e^{-2\eta}, e^{2(m-1-2\ell_r)\eta}; q^2)_{\infty}}.$$

Because

$$C_{\mathbf{k}}^{\gamma}(\mathbf{0}) = \mathcal{C}_{(1)} \times \dots \times \mathcal{C}_{(N)}$$

and the product over $i < j$ in (4.11) entails the restriction $k_i = k_j$, we note that the integral $\mu_{\mathbf{k}}$ factorizes into a product of $n_{\mathbf{k}}(r)$ -fold integrals for $r \in \{1, \dots, N\}$:

$$(4.14) \quad \mu_{\mathbf{k}} = \left(\prod_{i=1}^M -e^{\xi_- + 2 \sum_{s < k_i} \ell_s \eta} \right) \prod_{r=1}^N \mu_r^{\mathbf{k}}.$$

In view of (3.14) it therefore suffices to show that $\mu_r^{\mathbf{k}} = \nu_r^{\mathbf{k}}$.

We prove $\mu_r^{\mathbf{k}} = \nu_r^{\mathbf{k}}$ for the following parameter values. We fix $\mathbf{k} \in I_{M,N}$, $r \in \{1, \dots, N\}$ with $n_{\mathbf{k}}(r) \geq 1$ and $\tau \in \mathbb{C}$ with $\Re(\tau) < 0$. Consider the path-connected non-empty parameter domain

$$\mathcal{D} = \left\{ (\ell_r, \eta) \in \mathbb{C}^2 \mid \Re(\ell_r \eta) > \max\left(0, \frac{n_{\mathbf{k}}(r)-1}{2} \Re(\eta)\right) \right\}.$$

For fixed $(\ell_r, \eta) \in \mathcal{D}$ let

$$\gamma_{(r)} = (\gamma_{i(1;r)}, \dots, \gamma_{i(n_{\mathbf{k}}(r);r)}) \in \mathbb{C}^{n_{\mathbf{k}}(r)}$$

be such that

$$\begin{aligned} -\Re(\ell_r \eta) &< \Re(\gamma_{i(s;r)}) < \Re(\ell_r \eta), \\ \Re(\gamma_{i(s'+1;r)}) &\leq \Re(\gamma_{i(s';r)}), \\ \Re(\gamma_{i(s'+1;r)}) + \Re(\eta) &< \Re(\gamma_{i(s';r)}) \end{aligned}$$

for $1 \leq s \leq n_{\mathbf{k}}(r)$ and $1 \leq s' < n_{\mathbf{k}}(r)$. Note that if $\Re(\eta) < 0$ then $\gamma_{(r)} = (0, \dots, 0)$ satisfies the conditions.

The integral $\mu_r^{\mathbf{k}}$ (see (4.12)) does not depend on the choice of $\gamma_{(r)}$. This follows from the fact that the separation of the poles of $f_r^{\mathbf{k}}(\mathbf{z})$ by the cycle $\mathcal{C}_{(r)}$ does not depend on the choice of $\gamma_{(r)}$, and Cauchy's theorem. It now also follows directly that $\mu_r^{\mathbf{k}}$ is holomorphic in $(\ell_r, \eta) \in \mathcal{D}$. The same is true for $\nu_r^{\mathbf{k}}$ (see (4.13)) by a direct inspection. Hence it suffices to prove that $\mu_r^{\mathbf{k}} = \nu_r^{\mathbf{k}}$ for the restricted parameter domain

$$\{(\ell_r, \eta) \in \mathcal{D} \mid \Re(\eta) < 0\} = \{(\ell_r, \eta) \in \mathbb{C}^2 \mid \Re(\ell_r \eta) > 0 \text{ \& } \Re(\eta) < 0\}.$$

But this is the special case of [18, Appendix D - Proof of formula (5.13)] with the associated parameters specialized to

$$a = b = e^{-2\ell_r \eta}, \quad c = q e^{-2\omega_{\mathbf{k};r}}, \quad p = q^2, \quad x = e^{2\eta}. \quad \square$$

4.4. Completeness. Consider the boundary qKZ equations (2.13) as a compatible system of difference equations for $V^\ell(M)$ -valued meromorphic functions on \mathbb{C}^N . The fact that the leading coefficients $\nu_{\mathbf{k}} \Omega_{\mathbf{k}}$ of $\Theta_{\mathbf{k}}(\mathbf{t})$ as $\mathbf{t} \xrightarrow{\mathbb{A}} \infty$ ($\mathbf{k} \in I_{M,N}$) form a linear basis of $V^\ell(M)$ implies that the $\{\Psi_{\mathbf{k}}\}_{\mathbf{k} \in I_{M,N}}$ form a linear basis of the space of $V^\ell(M)$ -valued meromorphic solutions of the boundary qKZ equations over the field of $\tau\mathbb{Z}^N$ -periodic meromorphic functions on \mathbb{C}^N by the arguments of [10, §5.6]. This proves part **d** of Theorem 3.3.

5. INTEGRAL SOLUTIONS FOR FINITE-DIMENSIONAL REPRESENTATIONS OF QUANTUM \mathfrak{sl}_2

Fix $\ell_r \in \frac{1}{2}\mathbb{Z}_{>0}$ for $1 \leq r \leq N$ and write

$$\text{pr}^\ell := \text{pr}^{\ell_1} \otimes \dots \otimes \text{pr}^{\ell_N} : V^\ell \rightarrow \bar{V}^\ell := \bar{V}^{\ell_1} \otimes \dots \otimes \bar{V}^{\ell_N}$$

for the projection onto the finite-dimensional quotient. Write

$$\bar{\Omega}_{\mathbf{k}} := \text{pr}^\ell \Omega_{\mathbf{k}} = \bar{v}_{n_{\mathbf{k}}(1)}^{\ell_1} \otimes \dots \otimes \bar{v}_{n_{\mathbf{k}}(N)}^{\ell_N}$$

for $\mathbf{k} \in I_{M,N}$. Note that

$$\bar{V}^\ell = \bigoplus_{M=0}^{2\sum_{r=1}^N \ell_r} \bar{V}^\ell(M)$$

with

$$\bar{V}^\ell(M) = \bigoplus_{\mathbf{k} \in I_{M,N}^\ell} \mathbb{C} \bar{\Omega}_{\mathbf{k}}$$

and with index set

$$I_{M,N}^\ell := \{\mathbf{k} \in I_{M,N} \mid n_{\mathbf{k}}(r) \leq 2\ell_r \quad \forall r = 1, \dots, N\}.$$

Let $\overline{\mathcal{A}}_r(\mathbf{t}) : \overline{V}^\ell \rightarrow \overline{V}^\ell$ be the linear operators such that

$$\overline{\mathcal{A}}_r(\mathbf{t}) \circ \text{pr}^\ell = \text{pr}^\ell \circ \mathcal{A}_r(\mathbf{t}),$$

with $\mathcal{A}_r(\mathbf{t}) : V^\ell \rightarrow V^\ell$ the transport operators of the boundary qKZ equations (2.13). The boundary qKZ equations for \overline{V}^ℓ -valued meromorphic functions $\overline{\Psi}$ on \mathbb{C}^N are

$$(5.1) \quad \overline{\Psi}(\mathbf{t} + \tau \mathbf{e}_r) = \overline{\mathcal{A}}_r(\mathbf{t}) \overline{\Psi}(\mathbf{t}), \quad r \in \{1, \dots, N\}.$$

Let $\eta \in \mathbb{C}$ with $\Re(\eta) > 0$. Then for all $\mathbf{k} \in I_{M,N}^\ell$ we have $(\ell, \eta) \in \mathcal{D}_{M,N}^{\mathbf{k}}$. Hence Theorem 3.3 immediately gives the following result.

Theorem 5.1. *Let $\tau \in \mathbb{C}$ with $\Re(\tau) < 0$. Let $\ell_r \in \frac{1}{2}\mathbb{Z}_{>0}$ ($1 \leq r \leq N$) and $\eta \in \mathbb{C}$ with $\Re(\eta) > 0$. Let $0 \leq M \leq 2 \sum_{r=1}^N \ell_r$. For $\mathbf{k} \in I_{M,N}^\ell$ let $\Psi_{\mathbf{k}}$ be the $V^\ell(M)$ -valued meromorphic solution of the boundary qKZ equations (2.13) with respect to the parameters (ℓ, η) , as defined in Theorem 3.3.*

Then $\{\text{pr}^\ell \Psi_{\mathbf{k}}\}_{\mathbf{k} \in I_{M,N}^\ell}$ is a basis of the $\overline{V}^\ell(M)$ -valued meromorphic solutions of the boundary qKZ equations (5.1) over the field of $\tau\mathbb{Z}^N$ -invariant meromorphic functions.

Remark 5.2. If $\ell_1 = \dots = \ell_N = \frac{1}{2}$ then $\overline{V}^\ell \cong (\mathbb{C}^2)^{\otimes N}$ and $\overline{V}^\ell(M)$ is spanned by vectors $\overline{\Omega}_{\mathbf{k}}$ with $\mathbf{k} \in I_{M,N}$ such that $n_{\mathbf{k}}(r) \leq 1$ for all $r \in \{1, \dots, N\}$. For all such \mathbf{k} we may take $\gamma = (0, 0, \dots, 0) \in \Gamma_{M,N}^{\mathbf{k}}$, since $k_i = k_j$ for $i \neq j$ does not occur.

APPENDIX A. ASYMPTOTICS OF THE BOUNDARY qKZ EQUATIONS

Lemma A.1. *Let $\ell_1, \ell_2 \in \mathbb{C}$. The limits*

$$(A.1) \quad R_\infty^{\ell_1 \ell_2} := \lim_{\Re(x) \rightarrow \infty} R^{\ell_1 \ell_2}(x), \quad K_\infty^{\ell_1}(\xi) := \lim_{\Re(x) \rightarrow \infty} K^{\ell_1}(x; \xi).$$

exist and for all $d_1, d_2 \in \mathbb{Z}_{\geq 0}$ we have

$$(A.2) \quad \begin{aligned} R_\infty^{\ell_1 \ell_2}(v_{d_1}^{\ell_1} \otimes v_{d_2}^{\ell_2}) &= e^{2(d_1 d_2 - \ell_1 d_2 - d_1 \ell_2)\eta} v_{d_1}^{\ell_1} \otimes v_{d_2}^{\ell_2} \\ K_\infty^{\ell_1}(\xi)(v_{d_1}^{\ell_1}) &= (-1)^{d_1} e^{-d_1(2\xi + (2\ell_1 - d_1)\eta)} v_{d_1}^{\ell_1}. \end{aligned}$$

Proof. Let \mathcal{R} be the truncated universal R-matrix for $\widehat{\mathcal{U}}_\eta$, see Section 2.1. After twisting it by $z^{\frac{d}{2}} \otimes f_1$ we will have an element $\mathcal{R}(z) \in \widehat{\mathcal{U}}_\eta^{\otimes 2}[[z]]$ of the form $\mathcal{R}(z) = \exp(\frac{\eta}{2} h_1 \otimes h_1)(1 + O(z))$. The evaluation of $\mathcal{R}(z)$ in the tensor product of two irreducible representations $V^{\ell_1} \otimes V^{\ell_2}$ gives the R-matrix $e^{2\ell_1 \ell_2 \eta} R^{\ell_1 \ell_2}(x)$ where $z = e^x$; the extra factor is due to the normalized action on the tensor product of highest weight vectors. The limit $\Re(x) \rightarrow -\infty$ corresponds to $z \rightarrow 0$. Thus, for the R-matrices in question we have:

$$R^{\ell_1 \ell_2}(x) \rightarrow e^{2\ell_1 \ell_2 \eta} (\pi^{\ell_1} \otimes \pi^{\ell_2}) \exp\left(-\frac{\eta}{2} h_1 \otimes h_1\right)$$

as $\Re(x) \rightarrow -\infty$, and owing to unitarity (2.1) we obtain

$$R^{\ell_1 \ell_2}(x) \rightarrow e^{-2\ell_1 \ell_2 \eta} (\pi^{\ell_1} \otimes \pi^{\ell_2}) \exp\left(\frac{\eta}{2} h_1 \otimes h_1\right)$$

as $\Re(x) \rightarrow \infty$. When applied to $v_{d_1}^{\ell_1} \otimes v_{d_2}^{\ell_2}$, this gives the desired formula.

The asymptotic formula for $K^\ell(x; \xi)$ follows immediately from (2.10). \square

As a consequence of Lemma A.1 we have

Lemma A.2. *Let $\ell \in \mathbb{C}^N$ and $r \in \{1, \dots, N\}$. Then the limit $\mathcal{A}_{\infty, r}$ as defined in (2.16) exists and for all $\mathbf{k} \in I_{M, N}$ and all $r \in \{1, \dots, N\}$ we have*

$$(A.3) \quad \mathcal{A}_{\infty, r}(\Omega_{\mathbf{k}}) = \varphi_{\mathbf{k}; r} \Omega_{\mathbf{k}},$$

with $\varphi_{\mathbf{k}; r}$ given by (2.17).

Proof. According to Lemma A.1, the desired limit exists and is equal to

$$\begin{aligned} \mathcal{A}_{\infty, r} &= (R_{\infty}^{\ell_{r+1}})_{r+1} \cdots (R_{\infty}^{\ell_N})_{rN} \left(K_{\infty}^{\ell_r}(\xi_+) \right)_r (R_{\infty}^{\ell_r})_{Nr} \cdots (R_{\infty}^{\ell_{r+1}})_{r+1} \\ &\quad \times (R_{\infty}^{\ell_{r-1}})_{r-1} \cdots (R_{\infty}^{\ell_1})_{1r} \left(K_{\infty}^{\ell_r}(\xi_-) \right)_r (R_{\infty}^{\ell_1})_{1r}^{-1} \cdots (R_{\infty}^{\ell_{r-1}})_{r-1}^{-1} \\ &= (K_{\infty}^{\ell_r}(\xi_+) K_{\infty}^{\ell_r}(\xi_-))_r \prod_{s>r} (R_{\infty}^{\ell_s})_{rs}^2, \end{aligned}$$

where we have used (2.2) and the diagonality of both $R_{\infty}^{\ell_s}$ and $K_{\infty}^{\ell}(\xi)$. Applying (A.2), for $\mathbf{d} \in \mathbb{Z}_{\geq 0}^N$ we have

$$\begin{aligned} \mathcal{A}_{\infty, r}(v_{\mathbf{d}}) &= e^{-2d_r(\xi_+ + \xi_- + (2\ell_r - d_r)\eta)} \left(\prod_{s>r} e^{4(d_r d_s - \ell_r d_s - d_r \ell_s)\eta} \right) v_{\mathbf{d}} \\ &= e^{2d_r(\eta - \xi_+ - \xi_-) + 4d_r \left(\sum_{s>r} d_s + \frac{d_r - 1}{2} - \sum_{s \geq r} \ell_s \right) \eta} \left(\prod_{s>r} e^{-4\ell_r d_s \eta} \right) v_{\mathbf{d}}. \end{aligned}$$

Using the bijection $\zeta_{M, N} : I_{M, N} \rightarrow P_N(M)$ we arrive at the following formula

$$\begin{aligned} \mathcal{A}_{\infty, r}(\Omega_{\mathbf{k}}) &= e^{2n_{\mathbf{k}}(r)(\eta - \xi_+ - \xi_-) + 4n_{\mathbf{k}}(r) \left(\sum_{s>r} n_{\mathbf{k}}(s) + \frac{n_{\mathbf{k}}(r) - 1}{2} - \sum_{s \geq r} \ell_s \right) \eta} \\ &\quad \times \left(\prod_{s>r} e^{-4\ell_r n_{\mathbf{k}}(s)\eta} \right) \Omega_{\mathbf{k}} \end{aligned}$$

for $\mathbf{k} \in I_{M, N}$. We denote $\max_{\mathbf{k}}(r) = \max\{i \in \{1, \dots, M\} \mid k_i = r\}$. Recalling the definition of $n_{\mathbf{k}}(s)$ as given by (2.12) we have $\sum_{s>r} n_{\mathbf{k}}(s) = M - \max_{\mathbf{k}}(r)$. From the formula for the sum of a finite arithmetic progression we infer that

$$n_{\mathbf{k}}(r) \left(\sum_{s>r} n_{\mathbf{k}}(s) + \frac{n_{\mathbf{k}}(r) - 1}{2} \right) = \sum_{\substack{i=1 \\ k_i=r}}^M (M - i).$$

Formula (A.3) now readily follows. \square

APPENDIX B. THE BOUNDARY BETHE VECTORS

In this section we prove Proposition 3.1

B.1. Decomposition of ordinary Bethe vectors. For generic values of $\ell, \mathbf{t} \in \mathbb{C}^N$ and $x \in \mathbb{C}$ we write

$$T^{\ell}(x; \mathbf{t}) = L_{01}^{\ell_1}(x - t_1) \cdots L_{0N}^{\ell_N}(x - t_N) = \begin{pmatrix} * & B^{\ell}(x; \mathbf{t}) \\ * & D^{\ell}(x; \mathbf{t}) \end{pmatrix}$$

with $B^{\ell}(x; \mathbf{t}), D^{\ell}(x; \mathbf{t}) \in \text{End}(V^{\ell})$; for the case $N = 0$ the convention on empty products means $B^{\emptyset}(x; \emptyset) = 0$. For $\ell, \mathbf{t} \in \mathbb{C}^N$ and $\mathbf{x} \in \mathbb{C}^M$, we define

$$B^{\ell}(\mathbf{x}; \mathbf{t}) = B^{\ell}(x_1; \mathbf{t}) \cdots B^{\ell}(x_M; \mathbf{t}) \in \text{End}(V^{\ell})$$

with the convention that for $M = 0$ we have $B^{\ell}(\emptyset; \mathbf{t}) = \text{Id}_{V^{\ell}}$. From (3.1) it follows that $[B^{\ell}(x; \mathbf{t}), B^{\ell}(y; \mathbf{t})] = 0$; as a consequence we have $B^{\ell}(w\mathbf{x}; \mathbf{t})\Omega = B^{\ell}(\mathbf{x}; \mathbf{t})\Omega$ for

any permutation $w \in S_M$. The vectors $B^\ell(\mathbf{x}; \mathbf{t})\Omega$ are called *spin- ℓ ordinary (or type A) Bethe vectors*. A simple induction argument with respect to N , ultimately a consequence of the ice rule for the R -operators, establishes that

$$(B.1) \quad B^\ell(\mathbf{x}; \mathbf{t})\Omega = \sum_{\mathbf{d} \in P_N(M)} b_{\mathbf{d}}^\ell(\mathbf{x}; \mathbf{t}) v_{\mathbf{d}}$$

for some $b_{\mathbf{d}}^\ell(\mathbf{x}; \mathbf{t}) \in \mathbb{C}$, depending meromorphically on $\mathbf{x} \in \mathbb{C}^M$ and $\mathbf{t} \in \mathbb{C}^M$.

It is the aim of this section to provide a self-contained prescription for the closed formula for the coefficients $b_{\mathbf{d}}^\ell(\mathbf{x}; \mathbf{t})$. They are the “trigonometric weight functions” of [18]; we refer to this work, and references therein, for other points of view on $b_{\mathbf{d}}^\ell(\mathbf{x}; \mathbf{t})$ and other derivations for their explicit formulae.

Recall that $J^c := \{1, \dots, M\} \setminus J$ for any $J \subset \{1, \dots, M\}$ and introduce the notations $\widehat{\mathbf{z}} := (z_1, \dots, z_{N-1}) \in \mathbb{C}^{N-1}$ for $\mathbf{z} = (z_1, \dots, z_N) \in \mathbb{C}^N$ and $\mathbf{x}_J := (x_{i_1}, \dots, x_{i_{M-d}})$ for $\mathbf{x} = (x_1, \dots, x_M) \in \mathbb{C}^M$ and $J = \{i_1, \dots, i_{M-d}\}$ with $i_1 < i_2 < \dots < i_{M-d}$. From a statement analogous to [13, Lemma 4.2] it follows that

$$(B.2) \quad B^\ell(\mathbf{x}; \mathbf{t})\Omega = \sum_{J \subset \{1, \dots, M\}} \left(\prod_{i \in J} \frac{\sinh(t_N - x_i - (\frac{1}{2} - \ell_N)\eta)}{\sinh(t_N - x_i - (\frac{1}{2} + \ell_N)\eta)} \right) \\ \times \left(\prod_{\substack{i \in J \\ j \in J^c}} \frac{\sinh(x_i - x_j + \eta)}{\sinh(x_i - x_j)} \right) \left(B^{\widehat{\ell}}(\mathbf{x}_J; \widehat{\mathbf{t}}) \otimes \prod_{i \in J^c} B^{\ell_N}(x_i; t_N) \right) \Omega.$$

Cf. [14, Eq. (5.6)] we have, for $\ell, x, t \in \mathbb{C}$ and $d \in \mathbb{Z}_{\geq 0}$,

$$B^\ell(x; t) v_d^\ell = \frac{-e^{(-\ell + \frac{1}{2} + d)\eta} \sinh(\eta)}{\sinh(t - x - (\frac{1}{2} + \ell)\eta)} v_{d+1}^\ell.$$

Combining this with (B.2) we obtain

$$(B.3) \quad B^\ell(\mathbf{x}; \mathbf{t})\Omega = \sum_{d=0}^M e^{d(\frac{d}{2} - \ell_N)\eta} \sum_{\substack{J \subset \{1, \dots, M\} \\ \# J^c = d}} \left(\prod_{i \in J^c} \frac{-\sinh(\eta)}{\sinh(t_N - x_i - (\frac{1}{2} + \ell_N)\eta)} \right) \\ \times \left(\prod_{i \in J} \frac{\sinh(t_N - x_i - (\frac{1}{2} - \ell_N)\eta)}{\sinh(t_N - x_i - (\frac{1}{2} + \ell_N)\eta)} \right) \\ \times \left(\prod_{\substack{i \in J \\ j \in J^c}} \frac{\sinh(x_i - x_j + \eta)}{\sinh(x_i - x_j)} \right) \left(B^{\widehat{\ell}}(\mathbf{x}_J; \widehat{\mathbf{t}}) \Omega^{\widehat{\ell}} \right) \otimes v_d^{\ell_N}.$$

A collection of functions

$$\{c_{\mathbf{d}}^\ell : \mathbb{C}^M \otimes \mathbb{C}^N \rightarrow \mathbb{C} \text{ meromorphic} \mid \ell \in \mathbb{C}^N, \mathbf{d} \in P_N(M)\}_{M, N \in \mathbb{Z}_{\geq 0}}$$

is said to satisfy *quantum affine \mathfrak{sl}_2 -recursion* if the initial condition

$$(B.4) \quad c_{\emptyset}^\emptyset(\emptyset; \emptyset) = 1$$

and the recurrence relation

$$\begin{aligned}
 c_{\mathbf{d}}^{\ell}(\mathbf{x}; \mathbf{t}) &= e^{d_N(\frac{d_N}{2} - \ell_N)\eta} \sum_{\substack{J \subset \{1, \dots, M\} \\ \#J^c = d_N}} \left(\prod_{i \in J^c} \frac{-\sinh(\eta)}{\sinh(t_N - x_i - (\frac{1}{2} + \ell_N)\eta)} \right) \\
 &\times \left(\prod_{i \in J} \frac{\sinh(t_N - x_i - (\frac{1}{2} - \ell_N)\eta)}{\sinh(t_N - x_i - (\frac{1}{2} + \ell_N)\eta)} \right) \\
 &\times \left(\prod_{\substack{i \in J \\ j \in J^c}} \frac{\sinh(x_i - x_j + \eta)}{\sinh(x_i - x_j)} \right) c_{\mathbf{d}}^{\widehat{\ell}}(\mathbf{x}_J; \widehat{\mathbf{t}})
 \end{aligned}
 \tag{B.5}$$

hold true. Our plan is now as follows. First we will show that the $b_{\mathbf{d}}$ implicitly defined in (B.1) satisfy this recursion. Then we will present another collection of meromorphic functions $a_{\mathbf{d}}^{\ell}$ on $\mathbb{C}^M \times \mathbb{C}^N$, parametrized by $\ell \in \mathbb{C}^N$, $\mathbf{d} \in \mathbb{Z}_{\geq 0}^N$, by means of a closed-form expression, which can be shown to satisfy the same recursion. Since quantum affine \mathfrak{sl}_2 -recursion has a unique solution, we obtain $b_{\mathbf{d}}^{\ell} = a_{\mathbf{d}}^{\ell}$ and an explicit expression for the Bethe vectors in terms of the basis $\{v_{\mathbf{d}}^{\ell}\}_{\mathbf{d} \in \mathbb{Z}_{\geq 0}^N}$ follows.

Lemma B.1. $\{b_{\mathbf{d}}^{\ell} \mid \ell \in \mathbb{C}^N, \mathbf{d} \in P_N(M)\}_{M, N \in \mathbb{Z}_{\geq 0}}$ satisfies quantum affine \mathfrak{sl}_2 -recursion.

Proof. In the case $N = 0$, from $B^0(\mathbf{x}; \emptyset) = \delta_{0, M}$ we obtain (B.4). Combining (B.3) with (B.1) and using that the $v_{\mathbf{d}}^{\ell}$ form a basis for V^{ℓ} we obtain that the $b_{\mathbf{d}}^{\ell}$ also satisfy the recurrence relation (B.5). \square

Let $\mathbf{d} \in P_N(M)$ and define

$$I(\mathbf{d}) := \{\mathbf{m} \in \{1, \dots, N\}^M \mid \forall s \, n_{\mathbf{m}}(s) = d_s\}$$

and

$$\begin{aligned}
 a_{\mathbf{d}}^{\ell}(\mathbf{x}; \mathbf{t}) &= \sum_{\mathbf{m} \in I(\mathbf{d})} \left(\prod_i \frac{-e^{\left(\frac{n_{\mathbf{m}}(m_i)}{2} - \ell_{m_i}\right)\eta} \sinh(\eta)}{\sinh(t_{m_i} - x_i - (\frac{1}{2} + \ell_{m_i})\eta)} \prod_{s > m_i} \frac{\sinh(t_s - x_i - (\frac{1}{2} - \ell_s)\eta)}{\sinh(t_s - x_i - (\frac{1}{2} + \ell_s)\eta)} \right) \\
 &\times \prod_{\substack{i, j \\ m_i < m_j}} \frac{\sinh(x_i - x_j + \eta)}{\sinh(x_i - x_j)}.
 \end{aligned}$$

Lemma B.2. $\{a_{\mathbf{d}}^{\ell} \mid \ell \in \mathbb{C}^N, \mathbf{d} \in P_N(M)\}_{M, N \in \mathbb{Z}_{\geq 0}}$ satisfies quantum affine \mathfrak{sl}_2 -recursion.

Proof. It is immediately seen that the $a_{\mathbf{d}}^{\ell}$ satisfy (B.4) owing to the convention that empty sums are zero and empty products are one. To establish (B.5) for the $a_{\mathbf{d}}^{\ell}$ first note that if $n_{\mathbf{m}}(s) = d_s$ then

$$\sum_{i=1}^M \left(\frac{n_{\mathbf{m}}(m_i)}{2} - \ell_{m_i} \right) = \sum_{s=1}^N \sum_{\substack{i=1 \\ m_i=s}}^M \left(\frac{n_{\mathbf{m}}(s)}{2} - \ell_s \right) = \sum_{s=1}^N d_s \left(\frac{d_s}{2} - \ell_s \right).$$

Note that

$$a_{\widehat{\mathbf{d}}}^{\widehat{\ell}}(\mathbf{x}_J; \widehat{\mathbf{t}}) = e^{\sum_{s=1}^{N-1} d_s \left(\frac{d_s}{2} - \ell_s\right) \eta} \sum_{\mathbf{m} \in I(\widehat{\mathbf{d}})} \left(\prod_{i \in J} \frac{-\sinh(\eta)}{\sinh(t_{m_i} - x_i - (\frac{1}{2} + \ell_{m_i})\eta)} \right) \\ \times \prod_{s=m_i+1}^{N-1} \frac{\sinh(t_s - x_i - (\frac{1}{2} - \ell_s)\eta)}{\sinh(t_s - x_i - (\frac{1}{2} + \ell_s)\eta)} \prod_{\substack{i, j \in J \\ m_i < m_j}} \frac{\sinh(x_i - x_j + \eta)}{\sinh(x_i - x_j)}.$$

Hence, the right-hand side of (B.5) for $c_{\mathbf{d}}^{\ell} = a_{\mathbf{d}}^{\ell}$ is given by

$$e^{\sum_{s=1}^N d_s \left(\frac{d_s}{2} - \ell_s\right) \eta} \sum_{\mathbf{m} \in I(\widehat{\mathbf{d}})} \sum_{\substack{J \subset \{1, \dots, M\} \\ \#J^c = d_N}} \left(\prod_{i \in J^c} \frac{-\sinh(\eta)}{\sinh(t_N - x_i - (\frac{1}{2} + \ell_N)\eta)} \right) \left(\prod_{i \in J} \frac{-\sinh(\eta)}{\sinh(t_{m_i} - x_i - (\frac{1}{2} + \ell_{m_i})\eta)} \right) \\ \times \left(\prod_{i \in J} \frac{\sinh(t_N - x_i - (\frac{1}{2} - \ell_N)\eta)}{\sinh(t_N - x_i - (\frac{1}{2} + \ell_N)\eta)} \prod_{s=m_i+1}^{N-1} \frac{\sinh(t_s - x_i - (\frac{1}{2} - \ell_s)\eta)}{\sinh(t_s - x_i - (\frac{1}{2} + \ell_s)\eta)} \right) \\ \times \left(\prod_{\substack{i \in J \\ j \in J^c}} \frac{\sinh(x_i - x_j + \eta)}{\sinh(x_i - x_j)} \right) \left(\prod_{\substack{i, j \in J \\ m_i < m_j}} \frac{\sinh(x_i - x_j + \eta)}{\sinh(x_i - x_j)} \right).$$

We specify a map

$$I(\widehat{\mathbf{d}}) \times \{J \subset \{1, \dots, M\} \mid \#J = M - d_N\} \rightarrow I(\mathbf{d})$$

by inserting N 's in the $(M - d_N)$ -tuples at the places given by the elements of $J^c = \{1, \dots, M\} \setminus J$, i.e.

$$((m_1, \dots, m_{M-d_N}), J) \mapsto (m_1, \dots, m_{j_1-1}, N, m_{j_1}, \dots, m_{j_2-2}, N, m_{j_2-1}, \dots, \\ \dots, m_{j_{d_N}-d_N}, N, m_{j_{d_N}-d_N+1}, \dots, m_{M-d_N}),$$

where $J^c = \{j_1, \dots, j_{d_N}\}$ with $j_1 < j_2 < \dots < j_{d_N}$; this is evidently injective, and since both sets have finite cardinality $\frac{(M-d_N)!}{\prod_{s=1}^{N-1} d_s!} \binom{M}{d_N} = \frac{M!}{\prod_{s=1}^N d_s!}$ it follows that the map is bijective. Hence the right-hand side of (B.5) equals

$$e^{\sum_{s=1}^N d_s \left(\frac{d_s}{2} - \ell_s\right) \eta} \sum_{\mathbf{m} \in I(\mathbf{d})} \left(\prod_{\substack{i=1 \\ m_i=N}}^M \frac{-\sinh(\eta)}{\sinh(t_N - x_i - (\frac{1}{2} + \ell_N)\eta)} \right) \left(\prod_{\substack{i=1 \\ m_i \neq N}}^M \frac{-\sinh(\eta)}{\sinh(t_{m_i} - x_i - (\frac{1}{2} + \ell_{m_i})\eta)} \right) \\ \times \left(\prod_{\substack{i=1 \\ m_i \neq N}}^M \frac{\sinh(t_N - x_i - (\frac{1}{2} - \ell_N)\eta)}{\sinh(t_N - x_i - (\frac{1}{2} + \ell_N)\eta)} \prod_{s=m_i+1}^{N-1} \frac{\sinh(t_s - x_i - (\frac{1}{2} - \ell_s)\eta)}{\sinh(t_s - x_i - (\frac{1}{2} + \ell_s)\eta)} \right) \\ \times \left(\prod_{\substack{i, j=1 \\ m_i < m_j=N}}^M \frac{\sinh(x_i - x_j + \eta)}{\sinh(x_i - x_j)} \right) \left(\prod_{\substack{i, j=1 \\ m_i < m_j \neq N}}^M \frac{\sinh(x_i - x_j + \eta)}{\sinh(x_i - x_j)} \right),$$

where the remaining products recombine so as to yield $a_{\mathbf{d}}^{\ell}(\mathbf{x}; \mathbf{t})$ as required. \square

Combining Lemmas B.1 and B.2 we deduce

$$\begin{aligned}
 B^\ell(\mathbf{x}; \mathbf{t})\Omega &= \sum_{\mathbf{k} \in I_{M,N}} \sum_{\mathbf{m} \in S_M(\mathbf{k})} \left(\prod_{i=1}^M \frac{-e^{\left(\frac{n_{\mathbf{m}}(m_i)}{2} - \ell_{m_i}\right)\eta} \sinh(\eta)}{\sinh(t_{m_i} - x_i - (\frac{1}{2} + \ell_{m_i})\eta)} \right. \\
 &\quad \times \prod_{s=m_i+1}^N \frac{\sinh(t_s - x_i - (\frac{1}{2} - \ell_s)\eta)}{\sinh(t_s - x_i - (\frac{1}{2} + \ell_s)\eta)} \Bigg) \\
 &\quad \times \left(\prod_{\substack{i,j=1 \\ m_i < m_j}}^M \frac{\sinh(x_i - x_j + \eta)}{\sinh(x_i - x_j)} \right) \Omega_{\mathbf{m}}.
 \end{aligned}
 \tag{B.6}$$

B.2. Decomposition of the boundary Bethe vectors. Having completed the induction, we may now consider ℓ fixed and drop it from the notation. Because of (2.8) we have

$$T(-x; \mathbf{t})^{-1} = \left(\prod_{s=1}^N \frac{\sinh(t_s + x + (\frac{1}{2} - \ell_s)\eta)}{\sinh(t_s + x + (\frac{1}{2} + \ell_s)\eta)} \right) \sigma_0^y T(-x - \eta; \mathbf{t})^{t_0} \sigma_0^y.$$

Hence, from (3.3) it follows that

$$\begin{aligned}
 \mathcal{B}(x; \mathbf{t}) &= \left(\prod_{s=1}^N \frac{\sinh(t_s + x + (\frac{1}{2} - \ell_s)\eta)}{\sinh(t_s + x + (\frac{1}{2} + \ell_s)\eta)} \right) \\
 &\quad \times \left(D(-x - \eta; \mathbf{t}) B(x; \mathbf{t}) - \frac{\sinh(\xi_- - x)}{\sinh(\xi_- + x)} B(-x - \eta; \mathbf{t}) D(x; \mathbf{t}) \right).
 \end{aligned}
 \tag{B.7}$$

From (3.1) we obtain the commutation relation

$$\begin{aligned}
 D(x; \mathbf{t}) B(y; \mathbf{t}) &= \\
 &= \frac{\sinh(x - y + \eta)}{\sinh(x - y)} B(y; \mathbf{t}) D(x; \mathbf{t}) - \frac{\sinh(\eta)}{\sinh(x - y)} B(x; \mathbf{t}) D(y; \mathbf{t}).
 \end{aligned}$$

Using this and a trigonometric identity we infer from (B.7) that

$$\begin{aligned}
 \mathcal{B}(x; \mathbf{t}) &= \frac{\sinh(2x)}{\sinh(2x + \eta)} \left(\prod_{s=1}^N \frac{\sinh(t_s + x + (\frac{1}{2} - \ell_s)\eta)}{\sinh(t_s + x + (\frac{1}{2} + \ell_s)\eta)} \right) \\
 &\quad \times \left(B(x; \mathbf{t}) D(-x - \eta; \mathbf{t}) - \frac{\sinh(\xi_- - x - \eta)}{\sinh(\xi_- + x)} B(-x - \eta; \mathbf{t}) D(x; \mathbf{t}) \right).
 \end{aligned}$$

Hence

$$\tilde{\mathcal{B}}(x; \mathbf{t}) = \sum_{\epsilon \in \{\pm\}} \epsilon \frac{\sinh(\tilde{\xi}_- - \epsilon x)}{\sinh(\eta)} B(-\epsilon x - \frac{\eta}{2}; \mathbf{t}) D(\epsilon x - \frac{\eta}{2}; \mathbf{t}).
 \tag{B.8}$$

(B.8) serves as the base case of an inductive argument analogous to the proof of [13, Prop. 4.1]. It establishes that

$$\begin{aligned}
 \tilde{\mathcal{B}}(\mathbf{x}; \mathbf{t}) &= \sum_{\epsilon \in \{\pm\}^M} \left(\prod_{i=1}^M \epsilon_i \frac{\sinh(\tilde{\xi}_- - \epsilon_i x_i)}{\sinh(\eta)} \right) \left(\prod_{\substack{i,j=1 \\ i < j}}^M \frac{\sinh(\epsilon_i x_i + \epsilon_j x_j + \eta)}{\sinh(\epsilon_i x_i + \epsilon_j x_j)} \right) \\
 &\quad \times \left(\prod_{i=1}^M B(-\epsilon_i x_i - \frac{\eta}{2}; \mathbf{t}) \right) \left(\prod_{i=1}^M D(\epsilon_i x_i - \frac{\eta}{2}; \mathbf{t}) \right).
 \end{aligned}
 \tag{B.9}$$

In [14, Eq. (5.8)] the relation

$$(B.10) \quad D(x; \mathbf{t})\Omega = \left(\prod_{s=1}^N \frac{\sinh(t_s - x - (\frac{1}{2} - \ell_s)\eta)}{\sinh(t_s - x - (\frac{1}{2} + \ell_s)\eta)} \right) \Omega$$

is derived. Combining it with (B.9) we find the following expression for the spin- ℓ boundary Bethe vectors in terms of the spin- ℓ ordinary (type A) Bethe vectors:

$$(B.11) \quad \begin{aligned} \tilde{B}(\mathbf{x}; \mathbf{t})\Omega = & \sum_{\epsilon \in \{\pm\}^M} \left(\prod_{i=1}^M \epsilon_i \frac{\sinh(\tilde{\xi}_i - \epsilon_i x_i)}{\sinh(\eta)} \prod_{s=1}^N \frac{\sinh(t_s - \epsilon_i x_i + \ell_s \eta)}{\sinh(t_s - \epsilon_i x_i - \ell_s \eta)} \right) \\ & \times \left(\prod_{\substack{i,j=1 \\ i < j}}^M \frac{\sinh(\epsilon_i x_i + \epsilon_j x_j + \eta)}{\sinh(\epsilon_i x_i + \epsilon_j x_j)} \right) \left(\prod_{i=1}^M B(-\epsilon_i x_i - \frac{\eta}{2}; \mathbf{t}) \right) \Omega. \end{aligned}$$

In this formula we can then substitute expression (B.6) to arrive at an explicit formula for the coefficients $\beta_{\mathbf{k}}(\mathbf{x}; \mathbf{t})$ in the decomposition (3.4), yielding Theorem 3.1.

REFERENCES

- [1] I. Cherednik, *Quantum Knizhnik-Zamolodchikov equations and affine root systems*, Comm. Math. Phys. **150** (1992), 109–136.
- [2] V. G. Drinfeld, *Quantum groups*, Proceedings ICM, Vol. 1,2 (Berkeley 1986), 798–820, Amer. Math. Soc., Providence, RI, 1987.
- [3] P.I. Etingof, I.B. Frenkel, A.A. Kirillov Jr., *Lectures on Representation Theory and Knizhnik-Zamolodchikov equations*, Math. Surveys and Monographs, Vol. 58, Amer. Math. Soc., 1998.
- [4] I. Frenkel, N. Reshetikhin, *Quantum affine algebras and holonomic difference equations*, Comm. Math. Phys. **146** (1992), 1–60.
- [5] S. Ghoshal, A. Zamolodchikov, *Boundary S-matrix and boundary state in two-dimensional integrable quantum field theory*, Int. J. Mod. Phys. **A9** (1994) 3841–3886.
- [6] M. Jimbo, R. Kedem, T. Kojima, H. Konno, T. Miwa, *XXZ chain with a boundary*, Nucl. Phys. B **441** (1995), no. 3, 437–470.
- [7] M. Jimbo, R. Kedem, H. Konno, T. Miwa, R. Weston, *Difference equations in spin chains with a boundary*, Nucl. Phys. B **448** (1995), no. 3, 429–456.
- [8] T. Kojima, *Free field approach to diagonalization of boundary transfer matrix: recent advances*, arXiv:1103.5526.
- [9] K. Kuroki, A. Nakayashiki, *Free field approach to solutions of the quantum Knizhnik-Zamolodchikov equations*, SIGMA Symmetry Integrability Geom. Methods Appl. **4** (2008), Paper 049, 13 pp.
- [10] M. van Meer, J.V. Stokman, *Double affine Hecke algebras and bispectral quantum Knizhnik-Zamolodchikov equations*, Int. Math. Res. Not. IMRN **2010**, no. 6, 969–1040.
- [11] L. Mezincescu, R.I. Nepomechie, *Fusion procedure for open chains*, J. Phys. A: Math. Gen. **25**, no. 9 (1992), 2533.
- [12] N. Reshetikhin, *Jackson-type integrals, Bethe vectors, and solutions to a difference analog of the Knizhnik-Zamolodchikov system*, Lett. Math. Phys. **26** (1992), 153–165.
- [13] N. Reshetikhin, J.V. Stokman, B. Vlaar, *Boundary quantum Knizhnik-Zamolodchikov equations and Bethe vectors*, Comm. Math. Phys. **336** (2014), 953–986.
- [14] N. Reshetikhin, J.V. Stokman, B. Vlaar, *Boundary quantum Knizhnik-Zamolodchikov equations and fusion*, Ann. Henri Poincaré **17** (2016), no. 1, 137–177.
- [15] E. Sklyanin, *Boundary conditions for integrable quantum systems*, J. Phys. A **21** (1988), no. 10, 2375–2389.
- [16] J.V. Stokman, *Connection problems for quantum affine KZ equations and integrable lattice models*, Comm. Math. Phys. **338** (2015), no. 3, 1363–1409.
- [17] J.V. Stokman, B. Vlaar, *Koornwinder polynomials and the XXZ spin chain*, J. Approx. Theory **197** (2015), 69–100.

- [18] V.O. Tarasov, A.N. Varchenko, *Geometry of q -hypergeometric functions, quantum affine algebras and elliptic quantum groups*, Astérisque **246** (1997), vi+135 pp.
- [19] V.O. Tarasov, A.N. Varchenko, *Jackson integral representations for solutions to the quantized Knizhnik-Zamolodchikov equation*, St. Petersburg Math. J. **6** (1995), no. 2, 275–313.
- [20] B. Vlaar, *Boundary transfer matrices and boundary quantum KZ equations*, J. Math. Phys. **56** (2015), no. 7, 071705, 22 pp.

N.R.: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720, USA & ITMO UNIVERSITY, KRONVERSKII AVE. 49, SAINT PETERSBURG, 197101, RUSSIA & KdV INSTITUTE FOR MATHEMATICS, UNIVERSITY OF AMSTERDAM, SCIENCE PARK 105-107, 1098 XG AMSTERDAM, THE NETHERLANDS

E-mail address: `reshetik@math.berkeley.edu`

J.S.: KdV INSTITUTE FOR MATHEMATICS, UNIVERSITY OF AMSTERDAM, SCIENCE PARK 105-107, 1098 XG AMSTERDAM, THE NETHERLANDS.

E-mail address: `j.v.stokman@uva.nl`

B.V.: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, HESLINGTON, YORK YO10 5DD, UK.

E-mail address: `bart.vlaar@york.ac.uk`