

This is a repository copy of *Discovery of ^{72}Rb : A Nuclear Sandbank beyond the Proton Drip line*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/124091/>

Version: Accepted Version

Article:

Suzuki, H., Sinclair, Laura Frances, Soderstrom, P-A. et al. (33 more authors) (2017) Discovery of ^{72}Rb : A Nuclear Sandbank beyond the Proton Drip line. Physical Review Letters. 192503-1-6. ISSN: 1079-7114

<https://doi.org/10.1103/PhysRevLett.119.192503>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Supplementary material for Discovery of ^{72}Rb : A nuclear sandbank beyond the proton drip-line

H. Suzuki,¹ L. Sinclair,^{1,2} P.-A. Söderström,^{1,3,4,*} G. Lorusso,^{1,5} P. Davies,² L.S. Ferreria,⁶ E. Maglione,⁷ R. Wadsworth,² J. Wu,^{1,8,†} Z.Y. Xu,^{9,‡} S. Nishimura,¹ P. Doornenbal,¹ D.S. Ahn,¹ F. Browne,^{1,10} N. Fukuda,¹ N. Inabe,¹ T. Kubo,^{1,§} D. Lubos,^{1,11} Z. Patel,^{1,12} S. Rice,^{1,12} Y. Shimizu,¹ H. Takeda,¹ H. Baba,¹ A. Estrade,^{13,¶} Y. Fang,^{14,**} J. Henderson,^{2,††} T. Isobe,¹ D. Jenkins,² S. Kubono,¹ Z. Li,⁸ I. Nishizuka,^{15,‡‡} H. Sakurai,^{1,9} P. Schury,^{1,16,§§} T. Sumikama,¹⁵ H. Watanabe,^{17,18} and V. Werner^{19,3}

¹*RIKEN Nishina Center, 2-1 Hirosawa, Wako-shi, Saitama 351-0198, Japan*

²*Department of Physics University of York, Heslington, York YO10 5DD, United Kingdom*

³*Institut für Kernphysik, TU Darmstadt, D-64289 Darmstadt, Germany*

⁴*GSI Helmholtzzentrum für Schwerionenforschung GmbH, 64291 Darmstadt, Germany*

⁵*National Physical Laboratory, Teddington, Middlesex, TW11 0LW, UK*

⁶*Centro de Física e Engenharia de Materiais Avançados CeFEMA,*

Instituto Superior Técnico, Universidade de Lisboa,

Avenida Rovisco Pais, P1049-001 Lisbon, Portugal

⁷*Dipartimento di Fisica e Astronomia “G. Galilei”,*

and Istituto Nazionale di Fisica Nucleare, Via Marzolo 8, I-35131 Padova, Italy

⁸*Department of Physics, Peking University, Beijing 100871, China*

⁹*Department of Physics, University of Tokyo, Hongo, Bunkyo-ku, Tokyo 113-0033, Japan*

¹⁰*School of Computing, Engineering and Mathematics,*

University of Brighton, Brighton BN2 4JG, United Kingdom

¹¹*Physik Department E12, Technische Universität München, D-85748 Garching, Germany*

¹²*Department of Physics, University of Surrey, Guildford, GU2 7XH, UK*

¹³*School of Physics and Astronomy, University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom*

¹⁴*Department of Physics, Osaka University, Machikaneyama-machi 1-1, Osaka 560-0043 Toyonaka, Japan*

¹⁵*Department of Physics, Tohoku University, Aoba, Sendai, Miyagi 980-8578, Japan*

¹⁶*Institute of Physics, University of Tsukuba, Ibaraki 305-8571, Japan*

¹⁷*International Research Center for Nuclei and Particles in the Cosmos, Beihang University, Beijing 100191, China*

¹⁸*School of Physics and Nuclear Energy Engineering, Beihang University, Beijing 100191, China*

¹⁹*Wright Nuclear Structure Laboratory, Yale University, New Haven, CT 06520-8120, USA*

PACS numbers:

In this supplementary material we provide some detailed information about the experimental setup and results that are presented in the paper Discovery of ^{72}Rb : A submerged nuclear island beyond the proton drip-line.

The data were collected in three settings of the BigRIPS spectrometer, which are summarised in Table I.

The full set of cross-sections from this experiment, together with previous experiments in the same region, are shown in Fig. 1 and Fig. 2. As can be seen in these figures, the EPAX cross-sections reproduce the experimental data well in the $A = 70$ region, with the largest discrepancies being close to the drip line. In the $N = Z$ nuclei the cross sections are well known to be overestimated by a factor of 3-5.

*Corresponding author: p-a@ikp.tu-darmstadt.de

†Present address: Argonne National Laboratory, Argonne, Illinois 60439, USA

‡Present address: KU Leuven, Instituut voor Kern- en Stralingsfysica, 3001 Leuven, Belgium

§Present address: National Superconducting Cyclotron Laboratory (NSCL), Michigan State University (MSU), 640 South Shaw Lane, East Lansing, Michigan 48824-1321, USA

¶Present address: Department of Physics, Central Michigan University, Mount Pleasant, Michigan 48859, USA

**Present address: State University of New York at Stony Brook, Department of Physics and Astronomy, Stony Brook, New York, USA

††Present address: TRIUMF, 4004 Westbrook Mall, Vancouver, British Columbia, Canada V6T 2A3

‡‡Present address: GSI Helmholtzzentrum für Schwerionenforschung GmbH, 64291 Darmstadt, Germany

§§Present address: Institute of Particle and Nuclear Studies (IPNS), High Energy Accelerator Research Organization (KEK), Ibaraki 305-0801, Japan

TABLE I: Summary of the three different settings used in this experiment. This table shows the cenral particle, target thickness, magnetic rigidity ($B\rho$), degrader thickness, slit openings relative to the central trajectory, location of the ionization chamber, total beam time, average beam current, and live time for each individual setting. For quantities common to all three settings, these are listed only in the central column.

Setting	1	2	3
Central particle	^{73}Sr		
Target	Be 40.3 mm		
$B\rho_{0,1}$	5.110 Tm		
Degrader F1	Al 3.98 mm		
$B\rho_{1,2}$	4.432 Tm		
Degrader F5	Al 3.50 mm		
$B\rho_{5,7}$	3.606 Tm		
F1 slit	+64.2/0	+64.2/+5	+64.2/-30
	-3.0%/0.0%	-3.0%/+0.23%	-3.0%/+1.40%
F2 slit	+3/-5	+25/+9	+14/+8
F5 slit		± 120	
F7 slit	+2/-10	+8/10	+6/-14
IC for ΔE	F11		
Time	36.4 h	1.74 h	3.72 h
Current	30.2 pA	29.9 pA	30.7 pA
Live time	98.0 %	94.8 %	97.5 %




FIG. 1: Cross-section systematics for even- Z nuclei from the present experiment (red) and the associated cross-section measurement runs (blue), previous xenon-beam campaigns [1] (green), and the ^{100}Sn campaign [2] (magenta). The cross sections are compared to calculations based on EPAX3.1a and EPAX2.15.




FIG. 2: Cross-section systematics for odd- Z nuclei from the present experiment (red) and the associated cross-section measurement runs (blue), previous xenon-beam campaigns [1] (green), and the ^{100}Sn campaign [2] (magenta). The cross sections are compared to calculations based on EPAX3.1a and EPAX2.15.

Acknowledgments

This work was carried out at the RIBF operated by RIKEN Nishina Center, RIKEN and CNS, University of Tokyo. We acknowledge the EUROBALL Owners Committee for the loan of germanium detectors and the Pre-Spec Collaboration for the readout electronics of the clus-

ter detectors. This work was partially funded by UK STFC under grants ST/J000124/1 and ST/L005727/1. PAS was financed by the Japan Society for the Promotion of Science (JSPS) Kakenhi Grant No. 23-01752 and the RIKEN Foreign Postdoctoral Researcher Program. VW was supported by U.S. DOE grant DE-FG02-91ER40609 and NuSTAR.DA BMBF 05P15RDFN1.

-
- [1] H. Suzuki, T. Kubo, N. Fukuda, N. Inabe, D. Kameda, H. Takeda, K. Yoshida, K. Kusaka, Y. Yanagisawa, M. Ohtake, et al., Nucl. Instrum. Meth. **B317**, 756 (2013).
 - [2] I. Čeliković, M. Lewitowicz, R. Gernhäuser, R. Krücken,

S. Nishimura, H. Sakurai, D. Ahn, H. Baba, B. Blank, A. Blazhev, et al., Phys. Rev. Lett. **116**, 162501 (2016).