



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/124356/>

Version: Accepted Version

Article:

Kontopantelis, Evangelos, Mamas, Mamas A, van Marwijk, Harm et al. (2017)
Geographical epidemiology of health and overall deprivation in England, its changes and persistence from 2004 to 2015:a longitudinal spatial population study. Journal of epidemiology and community health. ISSN: 1470-2738

<https://doi.org/10.1136/jech-2017-209999>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

**Geographical epidemiology of health and overall deprivation
 in England, its changes and persistence from 2004 to
 2015:a longitudinal spatial population study**

Journal:	<i>Journal of Epidemiology & Community Health</i>
Manuscript ID	jech-2017-209999.R1
Article Type:	Research report
Date Submitted by the Author:	n/a
Complete List of Authors:	Kontopantelis, Evangelos; University of Manchester, the Farr Institute for Health Informatics Research; NIHR School for Primary Care Research, University of Manchester Mamas, Mamas; manchester heart centre van Marwijk, Harm; EMGO+ Institute of Health Care Research, General Practice and Elderly Care Medicine Ryan, Andrew; University of Michigan, Department of Health Management and Policy, School of Public Health Buchan, Iain; University of Manchester Institute of Population Health, Farr Institute Ashcroft, Darren; University of Manchester Doran, Tim; University of York,
Keywords:	DEPRIVATION, CLUSTERS, SPATIAL ANALYSIS, EPIDEMIOLOGY, Health inequalities

1
2
3 **Geographical epidemiology of health and overall deprivation in England, its changes and**
4 **persistence from 2004 to 2015: a longitudinal spatial population study**
5
6

7
8 Evangelos Kontopantelis^{1,2}, Mamas A. Mamas³, Harm van Marwijk^{1,2}, Andrew M Ryan⁴, Iain E
9 Buchan¹, Darren M. Ashcroft⁵, and Tim Doran⁶
10

11
12 ¹ Faculty of Biology Medicine and Health, University of Manchester, Greater Manchester, UK

13 ² NIHR School for Primary Care Research, University of Manchester, Greater Manchester, UK

14 ³ Centre for Prognosis Research, Institute for Primary care and Health Sciences, Keele

15 University, Stoke-on-Trent, UK

16
17 ⁴ School of Public Health, University of Michigan, MI, USA

18
19 ⁵ Division of Pharmacy and Optometry, Faculty of Biology Medicine and Health, University of
20 Manchester, Manchester Academic Health Science Centre, Manchester, UK

21
22 ⁶ Department of Health Sciences, University of York, Yorkshire, UK
23
24

25 Correspondence to:

26 Evangelos Kontopantelis

27 e-mail: e.kontopantelis@manchester.ac.uk
28
29
30
31
32

33
34 Word count: 3,424
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Background: Socio-economic deprivation is a key determinant for health. In England, the Index of Multiple Deprivation (IMD) is a widely used composite measure of deprivation. However, little is known about its spatial clustering or persistence across time.

Methods: Data for overall IMD and its health domain were analysed for 2004-2015 at a low geographical area (average of 1500 people). Levels and temporal changes were spatially visualised for the whole of England and its 10 administrative regions. Spatial clustering was quantified using Moran's I , correlations over time were quantified using Pearson's ρ .

Results: Between 2004 and 2015 we observed a strong persistence for both overall ($\rho=0.94$) and health-related deprivation ($\rho=0.92$). At the regional level, small changes were observed over time, but with areas slowly regressing towards the mean. However, for the North East, North West and Yorkshire, where health-related deprivation was the highest, the decreasing trend in health-related deprivation reversed and we noticed increases in 2015. Results did not support our hypothesis of increasing spatial clustering over time. However, marked regional variability was observed in both aggregate deprivation outcomes. The lowest autocorrelation was seen in the North East and changed very little over time, while the South East had the highest autocorrelation at all time points.

Conclusions: Overall and health-related deprivation patterns persisted in England, with large and unchanging health inequalities between the North and the South. The spatial aspect of deprivation can inform the targeting of health and social care interventions, particularly in areas with high levels of deprivation clustering.

Keywords: England; deprivation; Index of Multiple Deprivation; IMD; clustering; spatial; persistence; north; south.

What is already known on the subject

- Socio-economic deprivation is a key determinant of health.
- However, very little is known about the spatial clustering of English deprivation or its persistence across time.
- The Index of Multiple Deprivation (IMD) is a comprehensive aggregate measure incorporating seven domains of deprivation. The IMD can provide insight on relative changes in deprivation over time at the regional level.

What this study adds

- Between 2004 and 2015, overall and health-related deprivation in England strongly persisted at a low geographical area level.
- For the North of England, where health-related deprivation was the highest, the decreasing trend in health-related deprivation was reversed by 2015.
- Spatial clustering for both overall and health-related deprivation appears to have decreased over time, with some exceptions in 2015.
- Spatial aspects of deprivation can inform more effective organisation of health services and the targeting of health or social interventions, particularly in areas where there are high levels of deprivation clustering.

Introduction

The impact on health of socio-economic deprivation – for example, lower levels of income, wealth and education – is well established. At the area level, greater deprivation is associated with worse clinical outcomes,^{1 2} higher comorbidity levels,^{3 4} and lower levels of healthcare access and utilisation.^{5 6} Higher individual-level socio-economic deprivation has been linked to poorer health for secondary school students in New Zealand,⁷ higher oral-health-related hospitalisation rates in Australia,⁸ poorer self-reported health in Germany and the US,^{9 10} worse perinatal outcomes in France,¹¹ and higher all-cause and cause-specific mortality in England and Wales.¹²

Socio-economic deprivation is often found to be a more powerful determinant of health outcomes than medical care.¹³ In US studies, only 10-15% of preventable mortality has been estimated to be attributed to medical care.¹⁴ In England, performance of primary care practices on a range of quality of care indicators was not found to be associated with all-cause or cause-specific mortality,¹⁵ whereas local area deprivation was the strongest measured predictor.¹⁶ This highlights the importance of socio-economic deprivation, and the need for accurate and valid measurement to inform policies to address its health impacts.

A major challenge is developing measures that reliably quantify deprivation. Despite international interest, most countries have been slow to create comprehensive deprivation indices; for example, composite indices have only recently been proposed in Canada,^{17 18} and the US has yet to develop its own alternative.^{19 20} In England, the first widely used composite measure of material deprivation, the Townsend Index, was created in the 1980s, incorporating four indicator variables derived from decennial Census data (unemployment, non-car ownership, non-home ownership and household crowding).²¹ A similar measure, the Carstairs Index, was compiled for Scotland,²² and New Zealand developed its own census-based aggregate measure of deprivation.²³

In England, a more comprehensive measure, the Index of Multiple Deprivation (IMD), was developed in the 2000s incorporating routine administrative data covering a wider range of indicators (a total of 37 in 2004).²⁴ Iterations of the index have been reported for 2004,²⁵ 2007,²⁶ 2010,²⁷ and 2015.²⁸ The current measure quantifies relative deprivation across seven domains, known collectively as the English Indices of Deprivation (income, employment, education and skills, health and disability, crime, barriers to housing and services, and living environment), using an area-based model at a low geography (average of 1500 people) although it was originally developed for a higher level.²⁹ The overall IMD is calculated as a weighted mean across the seven domains, with income and employment deprivation given the largest weight (22.5% each), followed by health and education deprivation (13.5% each), and

1
2
3 with the other three domains given equal weights (9.3%). English epidemiological studies
4 routinely incorporate the index or individual component domains.
5
6

7
8 Despite the clinical importance of the English Indices of Deprivation, some aspects remain
9 under-researched. The first is the spatial nature of deprivation, i.e. how clustered it is. The
10 second is the longitudinal trends of clustering and regional deprivation (although the IMD is
11 standardised in each time point and thus does not allow for investigating absolute change,³⁰⁻³²
12 it is possible to assess relative changes over time). And the third is the persistence of
13 deprivation over time. All three are important to help guide the organisation of health care
14 services and to target social and health interventions. In this study we longitudinally and
15 spatially describe health-related (i.e. the health and disability domains) and overall deprivation
16 in England (the IMD). We hypothesised that between 2004 and 2015, and especially after the
17 2008 financial crisis, spatial autocorrelation for deprivation would increase, indicating
18 increasing spatial clustering of both poverty and wealth. We also quantify the longitudinal and
19 spatial properties of deprivation within each of 10 administrative English regions and make
20 comparisons across them. Finally, we quantify the persistence of deprivation over the study
21 period.
22
23
24
25
26
27
28
29
30

31 **Methods**

32 **English Indices of Deprivation**

33
34 Details about the subdomains included in each domain are provided in Table 1. For each
35 domain, the subdomains are normalised on ranking and then standardised, before factor
36 analysis is used to inform on the weighting of each subdomain. Therefore, the subdomain
37 weighting varies over time while the domain weighting does not. The health deprivation
38 domain aggregates information on years of potential life lost, illness and disability, acute
39 morbidity and mood and anxiety disorders. Their respective weights in 2015 were 0.244, 0.287,
40 0.254 and 0.216. The years these indicators cover also vary due to data availability and other
41 reasons. For 2015 for example, years of potential life lost was based on 2008-2012 calendar
42 year data, illness and disability on 2013 calendar year data, acute morbidity on 2011/12-
43 2012/13 financial year data and mood and anxiety disorders on 2012-2013 calendar year data.
44
45
46
47
48
49

50 Given the normalisation and standardisation of the subdomains, the aggregate measures, i.e.
51 overall IMD and each of the seven domains, are relative and cannot account for longitudinal
52 improvement or deterioration at the country level. However, longitudinal analyses can still
53 inform on distributional changes in the aggregate measures over time. Deprivation scores are
54 calculated and assigned to very low UK geographical units. Although small changes to the
55 indices have been implemented over time, comparability has been maintained since the
56
57
58
59
60

1
2
3 methods used are the same and no major changes have been considered necessary.²⁸ Separate
4 deprivation measures exist for other UK countries, with varying degrees of similarity to the
5 English IMD.
6
7

8 9 Lower Super Output Areas

10 The low geographical units to which the indices of deprivation are assigned are called lower
11 super output areas (LSOAs) and they are designed to contain around 1500 inhabitants, on
12 average. To roughly comply with this size-based definition, they can change following a
13 decennial census. Thus, following the 2011 census, English LSOAs were re-organised into
14 32,844 units (from 32,482 after the 2001 census) to better reflect population changes, mainly
15 increases.³³ Nevertheless only 2.5% of English LSOAs merged, split or underwent a more
16 complicated change.³⁴ Census-adjusted population estimates over time and for each English
17 LSOA were obtained from the Office of National Statistics.³⁵ Using the 2015 population
18 estimates we observed that mean LSOA size was 1654 inhabitants (median: 1595; 1st centile
19 1102; 99th centile 2926). Since pre-2011 deprivation indices were only reported using 2001
20 LSOAs we created a weighted means algorithm to assign them to 2011 LSOAs and allow a
21 seamless comparison over time. Spatial coordinates for the 2011 LSOAs were obtained from the
22 ONS open geography portal.³⁶ We used digital vector boundaries generalised to 20 meters and
23 clipped to the coastline to reduce size and improve visualisation. Finally, LSOAs were organised
24 into 10 regions to allow for comparisons within England, based on the 2006 restructuring of
25 Strategic Health Authorities: North East, North West, Yorkshire & the Humber, East Midlands,
26 West Midlands, East of England, London, South East Coast, South Central and South West.³⁷
27
28
29
30
31
32
33
34
35
36

37 Analyses

38 Deprivation outcomes of interest were overall and health-related IMD, in 2004, 2007, 2010 and
39 2015. We also analysed the fourth health-related IMD domains separately: years of potential
40 life lost, illness and disability, acute morbidity and mood and anxiety disorders. Outcomes and
41 their temporal changes were visualised using spatial maps for all of England and each of the 10
42 regions.
43
44
45

46
47 Spatial autocorrelation, or correlation in a signal among nearby locations in space, was
48 assessed and quantified using Moran's I .³⁸ This measure accounts for the multi-dimensional and
49 multi-directional nature of spatial autocorrelation, and can identify the presence or not of
50 deprivation clusters. A higher value than the one expected under a random spatial pattern
51 would indicate that deprived areas are clustered together and hence a deprived LSOA is more
52 likely to have deprived LSOAs as bordering or close neighbours (and similarly for affluent
53 LSOAs), with a value of 1 indicating perfect spatial correlation. We compared the values for
54 Moran's I in the four time points to assess whether existing spatial autocorrelation for
55
56
57
58
59
60

1
2
3 deprivation in 2004 has changed over time. An increasing value for Moran's I would indicate an
4 increase in deprivation-related spatial segregation. The measure was calculated for the whole
5 of England and each of the 10 regions, to allow for within-England comparisons.
6
7

8
9 Pairwise correlations across the four time points were computed (Pearson's ρ) to quantify the
10 persistency of area deprivation. Pairwise correlations between overall and health-related IMD
11 but also across health-related IMD subdomains were computed, to assess whether the two
12 main outcomes and the subdomains are different enough to justify reporting on all. To visualise
13 and compare temporal changes in the average deprivation levels between the 10 English
14 regions, we plotted population weighted boxplots for each of the two main outcomes and the
15 health subdomains. Analyses were executed with Stata v14.1 and R v3.3.1. Due of the size of
16 the dataset, effectively the whole of England, statistical significance is irrelevant; any
17 comparison would be statistically significant and thus we focus on effects sizes wherever
18 possible.
19
20
21
22
23

24 25 26 Results

27
28 Both overall and health-related deprivation was strongly correlated at all time points, indicating
29 small changes over time. Pearson's ρ remained above 0.94 for overall deprivation (2004 vs
30 2007: $\rho = 0.98$; 2004 vs 2010: $\rho = 0.97$; 2004 vs 2015: $\rho = 0.94$) and 0.92 for health-related
31 deprivation (2004 vs 2007: $\rho = 0.97$; 2004 vs 2010: $\rho = 0.93$; 2004 vs 2015: $\rho = 0.92$).
32 Correlation between overall and health-related IMD was strong, as anticipated, but the
33 strength somewhat deteriorated over time from $\rho = 0.88$ in 2004 to $\rho = 0.84$ in 2015. As
34 expected, subdomains of the health-specific deprivation were strongly correlated with overall
35 deprivation, with the lowest values observed for mood and anxiety disorders vs acute morbidity
36 ($\rho = 0.57$ in 2004 and $\rho = 0.63$ in 2015) and the highest for years of potential life lost vs the
37 comparative illness and disability ratio ($\rho = 0.79$ in 2004 and 2015).
38
39
40
41
42
43

44 Spatial maps were plotted for the whole of England and each region, for all years and each
45 deprivation outcome (overall IMD and health-related IMD and its four subdomains). We
46 present overall deprivation for 2015 and the whole of England in the main paper (Figure 1), and
47 for health-related deprivation in Appendix 2 (Supplemental Figure A1). Longitudinal spatial
48 maps for all other regions and the whole of England are provided in Appendices 1a to 1f, while
49 a longitudinal spatial map for London which allows for visual comparisons over time is
50 presented in Appendix 2 (Supplemental Figure A2).
51
52
53
54

55 Temporal changes of deprivation for English regions are presented in Figure 2. Although
56 changes over time are small, areas appear to be very slowly converging towards the English
57
58
59
60

1
2
3 average for both overall and health-related deprivation. However, for the three northern
4 regions where health-related deprivation was the highest (North-East, North West and
5 Yorkshire & the Humber) the decreasing trend between 2004 and 2010 seems to have been
6 reversed and we observed increases in 2015. The findings are similar across all four health
7 subdomains (Appendix 2, Supplemental Figures A3 and A4).
8
9

10
11
12 Spatial autocorrelation results did not support our original hypothesis of increasing spatial
13 clustering over time (Figure 3). Across the whole of England and for overall deprivation,
14 Moran's I was 0.0972 in 2004 (95% CI: 0.0970, 0.0974) but by 2015 it had linearly dropped to
15 0.0686 (95% CI: 0.0683, 0.0688), indicating a reduction in spatial autocorrelation and clustering.
16 However, the picture was different for health deprivation where spatial autocorrelation was
17 higher at a level of 0.1725 in 2004 (95% CI: 0.1723, 0.1728), dropped to 0.1490 in 2007 (95% CI:
18 0.1488, 0.1493) and remained relatively stable thereafter (0.1450 in 2010, 95%CI: 0.1448,
19 0.1453; 0.1505 in 2015, 95%CI: 0.1503, 0.1508). In addition, the levels and trends of spatial
20 autocorrelation were not consistent across the health deprivation subdomains. For years of
21 potential life lost, autocorrelation was relatively stable between 2004 and 2010, but
22 moderately increased in 2015. Spatial autocorrelation was higher for the comparative illness
23 and disability ratio, but it deteriorated over time and in 2015 it was lower than was observed
24 for years of potential life lost. Acute morbidity autocorrelation remained stable over time,
25 although a small increase was seen for 2015. Finally, autocorrelation for mood and anxiety
26 disorders has changed little over time and was the highest observed in 2015. Overall, regarding
27 the health IMD subdomains, spatial clustering appears to be increasing for years of potential
28 life lost and acute morbidity (Figure 3).
29
30
31
32
33
34
35
36
37

38 Marked regional variability was observed in the two aggregate deprivation outcomes, both for
39 2004 spatial autocorrelation levels and for their changes over time (Figure 4). The lowest
40 autocorrelation was seen in the North East and it has changed very little over time, while the
41 South East had the highest autocorrelation at all time points although it reduced over time. For
42 the East of England and the South Central coast, autocorrelation for overall deprivation and
43 especially health related deprivation increased in 2015, in contrast to the trends for the rest of
44 the country. Regional variation in autocorrelation was also observed for the health deprivation
45 subdomains (Appendix 2, Supplemental Figures A5 and A6). The smallest regional variation was
46 seen for years of potential life lost, with the South East reporting the highest levels and the
47 North East reporting the lowest. Changes over time were small for this subdomain, but trends
48 also varied by region. For the comparative illness and disability ratio, regional variability was
49 also modest and the general decreasing trend was seen in all regions, while the South East and
50 South West had consistently the highest level and the North East the lowest. For the other two
51 subdomains, acute morbidity and mood and anxiety disorders, regional changes in
52
53
54
55
56
57
58
59
60

1
2
3 autocorrelation over time are more unstable, with the East of England and the South Central
4 demonstrating high levels of autocorrelation.
5
6
7

8 9 Discussion

10 From 2004 to 2015, overall and health-related deprivation strongly persisted at the LSOA level.
11 Regional changes over time were modest for overall deprivation, with regional levels
12 converging towards the English average. However, the picture was different for health-related
13 deprivation (and within each of its four subdomains: years of potential life lost, illness and
14 disability, acute morbidity, and mood and anxiety disorders), with significant national
15 heterogeneity observed and levels consistently higher in the North East and North West, and
16 regional differences persisting over time. Our findings do not provide evidence of increasing
17 clustering for overall deprivation, with autocorrelation for overall deprivation declining over
18 time for the whole of England and each of its 10 regions, with some exceptions in 2015. Despite
19 this overall decreasing trend, regional variation in overall deprivation autocorrelation was
20 observed, with deprivation in the South coast appearing much more clustered than the North,
21 particularly the North East.
22
23
24
25
26
27

28
29 For health-related deprivation, levels of autocorrelation were consistently higher than what
30 was observed for overall deprivation, especially for the South coast. However, clustering levels
31 of health-related deprivation for the whole of England remained broadly the same since 2007,
32 with some regional variation of, primarily decreasing, trends. Different autocorrelation trends
33 were observed for the subdomains, with clustering increases for years of potential life lost and
34 acute morbidity.
35
36
37
38

39 Strengths and limitations of the study

40 This large longitudinal study that used data for the whole of England (53 million people in
41 2011), to investigate spatial and longitudinal patterns of overall and health-related deprivation.
42 To our knowledge, it is the first study of its kind to provide an insight into the temporal
43 persistence of deprivation and its clustering.
44
45
46
47

48 However, some limitations exist. First, there have been some minor changes in the underlying
49 indicators of deprivation over time, which might have influenced our estimates. However, the
50 trends we observe persist between 2007 and 2010 when there were no changes to the
51 measures, while the health deprivation domain has remained unchanged across the whole time
52 period. Second, the IMD and each of each domains are normalised and standardised at each
53 time point, hence the measure cannot account for longitudinal improvement or deterioration
54 at the country level,³⁹ only for relative regional changes. Third, there are varying levels of lag in
55
56
57
58
59
60

1
2
3 the underlying indicators included in each IMD version (for example data from 2012-13 used for
4 the 2015 IMD), but there is no data overlap across them. Fourth, it is possible that analyses at
5 the LSOA level are masking the relative deprivation trends in smaller and extremely deprived
6 communities.⁴⁰ Finally, there was a boundary change following the 2011 census which may
7 have affected our findings,⁴¹ but only 2.5% of LSOAs were affected while we developed an
8 algorithm to make reasonable population weighted based estimates for these localities,
9 depending on whether they merged, split or another change occurred.
10
11
12
13

14 Findings

15
16 Overcoming material and health deprivation is a global challenge. Neighbourhood connectivity
17 has been shown to be important, both in terms of the housing market and the labour market.⁴²
18 However, a lack of connectivity does not fully explain deprivation, which poses a diverse
19 challenge, a fact that needs to be taken into account by policy interventions. The very high
20 persistence of neighbourhood deprivation over the 11-year study period (2004 to 2015),
21 relative to the rest of England, was expected but is nevertheless of concern and highlights the
22 need for more social interventions to tackle inequality and social mobility in England.
23
24
25
26
27

28 The social inequality gap for the North of England, and how it compares to the South, is well
29 known,⁴³ although our results for overall deprivation indicate it has been slowly closing. In the
30 context of a universal healthcare provider, the National Health Service, persistent deprivation
31 plays an important role in premature mortality in England,^{44 45} but it does not fully explain the
32 persistently higher levels of all-cause mortality in the North,⁴⁶ which has recently increased at
33 an alarming rate for those aged 25-44.⁴⁷ Although health selective migration is likely to be a
34 factor,⁴⁸ and it is likely to influence inequalities by age,⁴⁹ we found that levels of health-related
35 deprivation were consistently higher in the North, not only in premature mortality but also
36 illness and disability, acute morbidity (hospital admissions), and mood and anxiety disorders.
37
38
39
40
41

42 Despite the longitudinal persistence of overall deprivation, the spatial clustering of
43 neighbourhood deprivation appears to have slowly decreased over time, although large
44 regional variation was observed both in levels of clustering and their changes. For most regions,
45 this reducing trend plateaued after 2010, possibly as a consequence of the 2008 financial crisis,
46 with London and the South East the notable exceptions. This pattern may be related to
47 gentrification projects, which have aimed to regenerate deprived inner city neighbourhoods.⁵⁰
48 Gentrification is a controversial and still debated practice, accused of displacement, segregation
49 and social polarisation,⁵¹ while too often the regeneration agenda is not inclusive.⁵²
50
51
52
53
54

55 High levels of spatial clustering for health-related deprivation would have implications for the
56 planning of health services and interventions, since they would imply a non-uniform
57
58
59
60

1
2
3 distribution of need across a region with hotspots of high levels of morbidity. To address such
4 spatial health inequality, services may need to be redesigned to take this factor into account.⁵³
5 For example, general practices may be incentivised to re-locate within such hotspots, and
6 targeted interventions may need to be prioritised in these areas. Similarly, infrastructure and
7 regeneration spending needs to be inclusive and weighted towards the health-deprivation
8 hotspots, rather than uniform. For the South of England we observed modest levels of
9 clustering, which even increased for the South Central region.
10
11

12
13
14
15 The regional clustering of health-related deprivation (and its levels) will be affected by external
16 economic migration, which varies greatly across regions.⁵⁴ On average, external migrants are
17 likely to be more materially deprived than English-born residents but with better health, at
18 least initially.⁵⁵ The age structure of the population will also play an important role, and the
19 population in the North of England is older.⁴⁷
20
21

22 23 24 Conclusions

25 Socio-economic deprivation is a key determinant of health. In order to improve population
26 health, policy makers need to not only invest in health care, but also – and perhaps more
27 importantly – to address material and educational inequalities. The time-lag between effective
28 policy intervention and reductions in inequality highlights the need for urgent action. We found
29 that from 2004 to 2015, overall and health-related deprivation patterns persisted in England at
30 a low geographical level. In terms of relative deprivation comparisons, regional levels of overall
31 deprivation appear to be slowly converging to the mean over time. However, regional variation
32 of health-related deprivation changed little, highlighting large and unchanging health
33 inequalities between the North and the South of England. Although this variation can be
34 partially explained by population differences, more needs to be done to address the issue.
35
36
37
38
39

40
41 Over time, the clustering of overall deprivation, and to a smaller extent of health-related
42 deprivation, decreased, with the regional trends demonstrating high heterogeneity. This
43 decrease and the underlying causes need to be better understood. The spatial aspect of
44 deprivation is often overlooked, but it can provide vital information for the effective
45 organisation of health services and targeting of health or social interventions. This is particularly
46 important for areas such as the South coast of England, where there are high levels of
47 clustering.
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgments

We would like to thank the Office of National Statistics for the wealth of information they have collected and systematically organised, which made this study possible.

Declaration of competing interests

MRC Health eResearch Centre Grant MR/K006665/1 supported the time and facilities of EK and IB. No other relationships or activities that could appear to have influenced the submitted work.

Authorship & contributorship

EK designed the study, extracted the data from all sources, performed the analyses and drafted the manuscript. MM, HvM, AMR, DA, IB and TD critically edited the manuscript.

EK is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Data sharing

The data used in this study are freely available and the authors are willing to share in an organised and cleaned final dataset.

References

1. Pickett KE, Pearl M. Multilevel analyses of neighbourhood socioeconomic context and health outcomes: a critical review. *J Epidemiol Commun H* 2001;55(2):111-22. doi: DOI 10.1136/jech.55.2.111
2. Stafford M, Marmot M. Neighbourhood deprivation and health: does it affect us all equally? *Int J Epidemiol* 2003;32(3):357-66. doi: 10.1093/ije/dyg084
3. Barnett K, Mercer SW, Norbury M, et al. Epidemiology of multimorbidity and implications for health care, research, and medical education: a cross-sectional study. *Lancet* 2012;380(9836):37-43. doi: 10.1016/S0140-6736(12)60240-2
4. Charlton J, Rudisill C, Bhattarai N, et al. Impact of deprivation on occurrence, outcomes and health care costs of people with multiple morbidity. *J Health Serv Res Po* 2013;18(4):215-23. doi: 10.1177/1355819613493772
5. LaVeist TA, Gaskin D, Richard P. Estimating the Economic Burden of Racial Health Inequalities in the United States. *Int J Health Serv* 2011;41(2):231-38. doi: 10.2190/HS.41.2.c
6. Franks P, Muennig P, Lubetkin E, et al. The burden of disease associated with being African-American in the United States and the contribution of socio-economic status. *Soc Sci Med* 2006;62(10):2469-78. doi: 10.1016/j.socscimed.2005.10.035
7. Denny S, Lewycka S, Utter J, et al. The association between socioeconomic deprivation and secondary school students' health: findings from a latent class analysis of a national adolescent health survey. *Int J Equity Health* 2016;15 doi: ARTN 109 10.1186/s12939-016-0398-5
8. Kruger E, Tennant M. Socioeconomic disadvantage and oral-health-related hospital admissions: a 10-year analysis. *BDJ Open* 2016;2:16004. doi: 10.1038/bdjopen.2016.4
9. Voigtländer S, Berger U, Razum O. The impact of regional and neighbourhood deprivation on physical health in Germany: a multilevel study. *BMC Public Health* 2010;10(1) doi: 10.1186/1471-2458-10-403
10. Subramanyam M, Kawachi I, Berkman L, et al. Relative deprivation in income and self-rated health in the United States. *Soc Sci Med* 2009;69(3):327-34. doi: 10.1016/j.socscimed.2009.06.008
11. Henrotin JB, Vaissiere M, Etaix M, et al. Deprivation, occupational hazards and perinatal outcomes in pregnant workers. *Occup Med (Lond)* 2017;67(1):44-51. doi: 10.1093/occmed/kqw148
12. Romeri E, Baker A, Griffiths C. Mortality by deprivation and cause of death in England and Wales, 1999-2003. *Stroke (cerebrovascular diseases)* 2006;60:169.
13. Braveman P, Gottlieb L. The Social Determinants of Health: It's Time to Consider the Causes of the Causes. *Public Health Rep* 2014;129:19-31.
14. McGinnis JM, Williams-Russo P, Knickman JR. The case for more active policy attention to health promotion. *Health Affair* 2002;21(2):78-93. doi: DOI 10.1377/hlthaff.21.2.78
15. Ryan AM, Krinsky S, Kontopantelis E, et al. Long-term evidence for the effect of pay-for-performance in primary care on mortality in the UK: a population study. *Lancet* 2016;388(10041):268-74. doi: 10.1016/S0140-6736(16)00276-2

16. Kontopantelis E, Springate DA, Ashworth M, et al. Investigating the relationship between quality of primary care and premature mortality in England: a spatial whole-population study. *Bmj-Brit Med J* 2015;350 doi: ARTN h904
10.1136/bmj.h904
17. Pampalon R, Hamel D, Gamache P, et al. A deprivation index for health planning in Canada. *Chronic Dis Can* 2009;29(4):178-91.
18. Chan E, Serrano J, Chen L, et al. Development of a Canadian socioeconomic status index for the study of health outcomes related to environmental pollution. *BMC Public Health* 2015;15:714. doi: 10.1186/s12889-015-1992-y
19. Phillips RL, Liaw W, Crampton P, et al. How Other Countries Use Deprivation Indices-And Why The United States Desperately Needs One. *Health Affair* 2016;35(11):1991-98. doi: 10.1377/hlthaff.2016.0709
20. Eibner C, Sturm R. US-based indices of area-level deprivation: Results from HealthCare for Communities. *Soc Sci Med* 2006;62(2):348-59. doi: 10.1016/j.socscimed.2005.06.017
21. Townsend P. Deprivation. *J Soc Policy* 1987;16:125-46.
22. Carstairs VDL, Morris R. Deprivation and health in Scotland 1991.
23. Salmond CE, Crampton P. Development of New Zealand's Deprivation Index (NZDep) and Its Uptake as a National Policy Tool. *Can J Public Health* 2012;103(8):S7-S11.
24. Noble M, Wright G, Smith G, et al. Measuring multiple deprivation at the small-area level. *Environ Plann A* 2006;38(1):169-85. doi: 10.1068/a37168
25. Communities and Local Government. The English Indices of Deprivation 2004 (revised): Department for Communities and Local Government, 2004.
26. Communities and Local Government. The English Indices of Deprivation 2007: Technical Report: Department for Communities and Local Government, 2008.
27. Communities and Local Government. The English Indices of Deprivation 2010: Technical Report: Department for Communities and Local Government, 2011.
28. Communities and Local Government. The English Indices of Deprivation 2015: Technical Report: Department for Communities and Local Government, 2015.
29. Noble M, Smith G, Penhale B, et al. Measuring multiple deprivation at the small area level: the indices of deprivation 2000. *Regeneration research summary* 2000;37
30. Norman P. Identifying Change Over Time in Small Area Socio-Economic Deprivation. *Applied Spatial Analysis and Policy* 2010;3(2):107-38. doi: 10.1007/s12061-009-9036-6
31. Norman P. The Changing Geography of Deprivation in Britain, 1971 to 2011 and Beyond. 2016
32. Norman P. Demographic and Deprivation Change in the UK. In: Stillwell J, Norman P, Thomas C, et al., eds. *Spatial and Social Disparities: Understanding Population Trends and Processes: volume 2*. Dordrecht: Springer Netherlands 2010:17-35.
33. Office for National Statistics. Changes to Output Areas and Super Output Areas in England and Wales, 2001 to 2011, 2012:13.
34. Office for National Statistics. Lower Layer Super Output Area (2001) to Lower Layer Super Output Area (2011) to Local Authority District (2011) Lookup in England and Wales [Available from: http://geoportal.statistics.gov.uk/datasets/6d3b1fc88b284a9bb9d4827530b16da4_02016.

- 1
2
3
4 35. Office for National Statistics. Super Output Area mid-year population estimates for England
5 and Wales, Mid-2011 (Census Based) [Available from:
6 [http://www.ons.gov.uk/ons/publications/re-reference-tables.html?edition=tcm%3A77-](http://www.ons.gov.uk/ons/publications/re-reference-tables.html?edition=tcm%3A77-2856292016)
7 [2856292016](http://www.ons.gov.uk/ons/publications/re-reference-tables.html?edition=tcm%3A77-2856292016).
8
9 36. Office for National Statistics. Open Geography Portal [Available from:
10 <http://geoportal.statistics.gov.uk/2016>.
11
12 37. Department of Health. Strategic Health Authority Configurations 2006 [Available from:
13 [https://web.archive.org/web/20070205232558/http://www.dh.gov.uk/assetRoot/04/1](https://web.archive.org/web/20070205232558/http://www.dh.gov.uk/assetRoot/04/13/37/60/04133760.pdf)
14 [3/37/60/04133760.pdf](https://web.archive.org/web/20070205232558/http://www.dh.gov.uk/assetRoot/04/13/37/60/04133760.pdf).
15
16 38. Moran PAP. Notes on Continuous Stochastic Phenomena. *Biometrika* 1950;37(1-2):17-23.
17 doi: Doi 10.2307/2332142
18
19 39. Norman P, Darlington-Pollock F. The Changing Geography of Deprivation in Great Britain:
20 Exploiting Small Area Census Data, 1971 to 2011. 2017
21
22 40. Exeter D, Flowerdew R, Boyle P. Policy implications of pockets of deprivation in Scotland.
23 *GIS Evid Based Policy Making* 2007;95
24
25 41. Norman P, Rees P, Boyle P. Achieving data compatibility over space and time: creating
26 consistent geographical zones. *Population, Space and Place* 2003;9(5):365-86.
27
28 42. Rae A, Hamilton R, Crisp R, et al. Overcoming deprivation and disconnection in UK cities:
29 Joseph Rowntree Foundation 2016.
30
31 43. Doran T, Drever F, Whitehead M. Is there a north-south divide in social class inequalities in
32 health in Great Britain? Cross sectional study using data from the 2001 census. *Brit Med*
33 *J* 2004;328(7447):1043-45. doi: DOI 10.1136/bmj.328.7447.1043
34
35 44. Norman P, Boyle P, Exeter D, et al. Rising premature mortality in the UK's persistently
36 deprived areas: Only a Scottish phenomenon? *Soc Sci Med* 2011;73(11):1575-84. doi:
37 10.1016/j.socscimed.2011.09.034
38
39 45. Exeter DJ, Boyle PJ, Norman P. Deprivation (im)mobility and cause-specific premature
40 mortality in Scotland. *Soc Sci Med* 2011;72(3):389-97. doi:
41 10.1016/j.socscimed.2010.10.009
42
43 46. Hacking JM, Muller S, Buchan IE. Trends in mortality from 1965 to 2008 across the English
44 north-south divide: comparative observational study. *Brit Med J* 2011;342 doi: ARTN
45 d508
46 10.1136/bmj.d508
47
48 47. Buchan IE, Kontopantelis E, Sperrin M, et al. North-South disparities in English
49 mortality 1965-2015: longitudinal population study. *J Epidemiol Community Health*
50 2017;71(9):928-36. doi: 10.1136/jech-2017-209195 [published Online First: 2017/08/10]
51
52 48. Connolly S, O'Reilly D, Rosato M. Increasing inequalities in health: Is it an artefact caused by
53 the selective movement of people? *Soc Sci Med* 2007;64(10):2008-15. doi:
54 10.1016/j.socscimed.2007.02.021
55
56 49. Norman P, Boyle P. Are health inequalities between differently deprived areas evident at
57 different ages? A longitudinal study of census records in England and Wales, 1991-2001.
58 *Health Place* 2014;26:88-93. doi: 10.1016/j.healthplace.2013.12.010
59
60 50. Lees L. A reappraisal of gentrification: towards a 'geography of gentrification'. *Prog Hum*
Geog 2000;24(3):389-408. doi: Doi 10.1191/030913200701540483

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
51. Lees L. Gentrification and Social Mixing: Towards an Inclusive Urban Renaissance? *Urban Stud* 2008;45(12):2449-70. doi: 10.1177/0042098008097099
 52. Atkinson R. The evidence on the impact of gentrification: new lessons for the urban renaissance? *International Journal of Housing Policy* 2004;4(1):107-31. doi: 10.1080/1461671042000215479
 53. Lorant V, Thomas I, Deliege D, et al. Deprivation and mortality: the implications of spatial autocorrelation for health resources allocation. *Soc Sci Med* 2001;53(12):1711-19. doi: Doi 10.1016/S0277-9536(00)00456-1
 54. Rienzo C, Vargas-Silva C. Migrants in the UK: An Overview. Migration Observatory Briefing, 2016.
 55. Rechel B, Mladovsky P, Ingleby D, et al. Migration and health in an increasingly diverse Europe. *Lancet* 2013;381(9873):1235-45. doi: 10.1016/S0140-6736(12)62086-8

Table 1: Index of multiple deprivation domains, 2015*

Domain (weight)	Subdomains
Income (22.5%)	<ul style="list-style-type: none"> Adults and children in Income Support families Adults and children in income-based Jobseeker's Allowance families Adults and children in income-based Employment and Support Allowance families Adults and children in Pension Credit (Guarantee) families Adults and children in Child Tax Credit and Working Tax Credit families, below 60% median income not already counted Asylum seekers in England in receipt of subsistence support, accommodation support, or both
Employment (22.5%)	<ul style="list-style-type: none"> Claimants of Jobseeker's Allowance, aged 18-59/64 Claimants of Employment and Support Allowance, aged 18-59/64 Claimants of Incapacity Benefit, aged 18-59/64 Claimants of Severe Disablement Allowance, aged 18-59/64 Claimants of Carer's Allowance, aged 18-59/64
Health and disability (13.5%)	<ul style="list-style-type: none"> Years of potential life lost: age/sex standardised measure of premature death Comparative illness and disability ratio: age/sex standardised morbidity/disability ratio Acute morbidity: age/sex standardised rate of emergency admission to hospital Mood and anxiety disorders: composite score based on the rate of adults suffering from mood and anxiety disorders, hospital episodes data, suicide mortality data and health benefits data
Education, Skills & Training (13.5%)	<ul style="list-style-type: none"> Key stage 2 attainment: average points score Key stage 4 attainment: average points score Secondary school absence Staying on in education post 16 Entry to higher education Adults with no or low qualifications, aged 25-59/64 English language proficiency, aged 25-59/64
Crime (9.3%)	<ul style="list-style-type: none"> Recorded crime rates for: Violence; Burglary; Theft; Criminal damage
Barriers to Housing & Services (9.3%)	<ul style="list-style-type: none"> Road distance to: post office; primary school; general store / supermarket; GP surgery Household overcrowding Homelessness Housing affordability
Living Environment (9.3%)	<ul style="list-style-type: none"> Housing in poor condition Houses without central heating Air quality Road traffic accidents

*Details available in the 2015 technical report of the English Indices of Deprivation ²⁸

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1: Overall deprivation for England, 2015



Figure 2: Overall (top) and health-related (bottom) deprivation for English regions, over time

Figure 3: Spatial autocorrelation with Moran's I for deprivation domains and subdomains, for the whole of England over time

Figure 4: Spatial autocorrelation with Moran's I for overall (top) and health-related (bottom) deprivation by region, over time

Confidential: For Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24







Figure 2: Overall (top) and health-related (bottom) deprivation for English regions, over time


203x270mm (300 x 300 DPI)






Deprivation domains and subdomains over time

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37




Regions over time

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37



Online Appendix 2 for “Geographical epidemiology of health and overall deprivation in England, its changes and persistence from 2004 to 2015: a longitudinal spatial study”

Figure A1: Health-related deprivation for England, 2015



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure A2: Overall (top) and health-related (bottom) deprivation for Greater London, over time




Figure A3: Health-related deprivation subdomains for English regions, over time (1 of 2)




Figure A4: Health-related deprivation subdomains for English regions, over time (2 of 2)




Figure A5: Spatial autocorrelation with Moran's I for years of potential life lost (top) and the comparative illness and disability ratio (bottom), by region and over time




Figure A6: Spatial autocorrelation with Moran's I for acute morbidity (top) and mood and anxiety disorders (bottom), by region and over time

