

This is a repository copy of *Badly approximable points on planar curves and winning*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/124399/>

Version: Published Version

Article:

An, Jinpeng, Beresnevich, Victor orcid.org/0000-0002-1811-9697 and Velani, Sanju orcid.org/0000-0002-4442-6316 (2018) Badly approximable points on planar curves and winning. *Advances in Mathematics*. 148–202. ISSN: 0001-8708

<https://doi.org/10.1016/j.aim.2017.11.009>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.




ELSEVIER

Contents lists available at ScienceDirect

Advances in Mathematics

www.elsevier.com/locate/aim



Badly approximable points on planar curves and winning

Jinpeng An^{a,1}, Victor Beresnevich^{b,*,2}, Sanju Velani^{b,3}^a LMAM, School of Mathematical Sciences, Peking University, Beijing, 100871, China^b Department of Mathematics, University of York, Heslington, York, YO10 5DD, England, United Kingdom

ARTICLE INFO

Article history:

Received 11 August 2017

Accepted 6 November 2017

Available online xxxx

Communicated by Kenneth Falconer

MSC:

11J83

11J13

11K60

Keywords:

Inhomogeneous Diophantine approximation

Non-degenerate curves

Schmidt games

ABSTRACT

For any $i, j > 0$ with $i + j = 1$, let $\mathbf{Bad}(i, j)$ denote the set of points $(x, y) \in \mathbb{R}^2$ such that $\max\{\|qx\|^{1/i}, \|qy\|^{1/j}\} > c/q$ for some positive constant $c = c(x, y)$ and all $q \in \mathbb{N}$. We show that $\mathbf{Bad}(i, j) \cap \mathcal{C}$ is winning in the sense of Schmidt games for a large class of planar curves \mathcal{C} , namely, everywhere non-degenerate planar curves and straight lines satisfying a natural Diophantine condition. This strengthens recent results solving a problem of Davenport from the sixties. In short, within the context of Davenport's problem, the winning statement is best possible. Furthermore, we obtain the inhomogeneous generalisations of the winning results for planar curves and lines and also show that the inhomogeneous form of $\mathbf{Bad}(i, j)$ is winning for two dimensional Schmidt games.

© 2017 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

* Corresponding author.

E-mail addresses: anjinpeng@gmail.com (J. An), vb8@york.ac.uk (V. Beresnevich), slv3@york.ac.uk (S. Velani).¹ Research supported by NSFC grant 11322101.² Research supported by EPSRC grant EP/J018260/1.³ Research supported by EPSRC grant EP/J018260/1.

1. Introduction

A real number x is said to be *badly approximable* if there exists a positive constant $c(x)$ such that

$$\|qx\| > c(x) q^{-1} \quad \forall q \in \mathbb{N}.$$

Here and throughout $\|\cdot\|$ denotes the distance of a real number to the nearest integer. It is well-known that the set **Bad** of badly approximable numbers is of Lebesgue measure zero but of maximal Hausdorff dimension; i.e. $\dim \mathbf{Bad} = 1$. In higher dimensions there are various natural generalisations of **Bad**. Restricting our attention to the plane \mathbb{R}^2 , given a pair of real numbers i and j such that

$$0 < i, j < 1 \quad \text{and} \quad i + j = 1, \quad (1.1)$$

a point $(x, y) \in \mathbb{R}^2$ is said to be (i, j) -*badly approximable* if there exists a positive constant $c(x, y)$ such that

$$\max\{\|qx\|^{\frac{1}{i}}, \|qy\|^{\frac{1}{j}}\} > c(x, y) q^{-1} \quad \forall q \in \mathbb{N}.$$

Denote by $\mathbf{Bad}(i, j)$ the set of (i, j) -badly approximable points in \mathbb{R}^2 . In the case $i = j = 1/2$, the set under consideration is the standard set of simultaneously badly approximable points. It easily follows from classical results in the theory of metric Diophantine approximation that $\mathbf{Bad}(i, j)$ is of (two-dimensional) Lebesgue measure zero. Regarding dimension, it was shown by Schmidt [14] in the vintage year of 1966 that $\dim \mathbf{Bad}(\frac{1}{2}, \frac{1}{2}) = 2$. In fact, Schmidt proved the significantly stronger statement that $\mathbf{Bad}(\frac{1}{2}, \frac{1}{2})$ is winning in the sense of his now famous (α, β) -games – see §2.1. Almost forty years later it was proved in [12] that $\dim \mathbf{Bad}(i, j) = 2$ and just recently the first author in [2] has shown that $\mathbf{Bad}(i, j)$ is in fact winning. The latter implies that any countable intersection of $\mathbf{Bad}(i, j)$ sets is of full dimension and thus provides a clean and direct proof of Schmidt's Conjecture – see also [1, 3].

Now let \mathcal{C} be a planar curve. Without loss of generality, we assume that \mathcal{C} is given as a graph

$$\mathcal{C}_f := \{(x, f(x)) : x \in I\}$$

for some function f defined on an interval $I \subset \mathbb{R}$. Throughout we will assume that $f \in C^{(2)}(I)$, a condition that conveniently allows us to define the curvature. Motivated by a problem of Davenport [9, p. 52] from the sixties, the following statement regarding the intersection of $\mathbf{Bad}(i, j)$ sets with any curve \mathcal{C} that is not a straight line segment has recently been established [4, 5].

Theorem A. *Let (i_t, j_t) be a countable number of pairs of real numbers satisfying (1.1) and suppose that*

$$\liminf_{t \rightarrow \infty} \min\{i_t, j_t\} > 0. \quad (1.2)$$

Let $\mathcal{C} := \mathcal{C}_f$ be a $C^{(2)}$ planar curve that is not a straight line segment. Then

$$\dim \left(\bigcap_{t=1}^{\infty} \mathbf{Bad}(i_t, j_t) \cap \mathcal{C} \right) = 1.$$

The theorem implies that there are continuum many points on the parabola $\mathcal{V}_2 := \{(x, x^2) : x \in \mathbb{R}\}$ that are simultaneously badly approximable in the $\mathbf{Bad}(\frac{1}{2}, \frac{1}{2})$ sense and thus provides a solution to the specific problem raised by Davenport in [9]. It is worth mentioning that a consequence of [6, Theorem 1] is that the set $\mathbf{Bad}(i, j) \cap \mathcal{C}$ is of zero (induced) Lebesgue measure on \mathcal{C} . Thus, the fact that it is a set of full dimension is not trivial.

The condition imposed on \mathcal{C} is natural since the statement is not true for all lines. Indeed, let L_a denote the vertical line parallel to the y -axis passing through the point $(a, 0)$ in the (x, y) -plane. Then, it is easily verified, see [3, §1.3] for the details, that $\mathbf{Bad}(i, j) \cap L_a = \emptyset$ for any $a \in \mathbb{R}$ satisfying $\liminf_{q \rightarrow \infty} q^{1/i} \|qa\| = 0$. On the other hand, if the \liminf is strictly positive then $\dim(\mathbf{Bad}(i, j) \cap L_a) = 1$. This is much harder to prove and is at the heart of the original proof of Schmidt's Conjecture established in [3]. Subsequently, it was shown in [1] that $\mathbf{Bad}(i, j) \cap L_a$ is winning. The following non-trivial extension of the full dimensional result to non-vertical lines has recently been established in [4].

Theorem B. *Let (i_t, j_t) be a countable number of pairs of real numbers satisfying (1.1) and (1.2). Given $a, b \in \mathbb{R}$, let $L_{a,b}$ denote the line defined by the equation $y = ax + b$. Suppose there exists $\epsilon > 0$ such that*

$$\liminf_{q \rightarrow \infty} q^{\frac{1}{\sigma} - \epsilon} \|qa\| > 0 \quad \text{where} \quad \sigma := \sup\{\min\{i_t, j_t\} : t \in \mathbb{N}\}. \quad (1.3)$$

Then

$$\dim \left(\bigcap_{t=1}^{\infty} \mathbf{Bad}(i_t, j_t) \cap L_{a,b} \right) = 1.$$

Both Theorem A and Theorem B should be true without the \liminf condition (1.2). Indeed, as pointed out in Remark 2 of [4, § 1.2], it is very tempting and not at all outrageous to assert that $\mathbf{Bad}(i, j) \cap \mathcal{C}$ is winning at least on the part of the curve that is genuinely curved. If true it would imply Theorem A without assuming (1.2). In short, this is precisely what we show in this paper. We also obtain a winning statement for non-vertical lines that not only implies Theorem B without assuming (1.2) but replaces condition (1.3) by a weaker and essentially optimal Diophantine condition. Furthermore, by making use of a simple idea introduced in [7] that provides a natural mechanism for

generalising homogenous badly approximable statements to the inhomogeneous setting, we establish the inhomogeneous generalisation of our winning results. The same idea is also exploited to prove that inhomogeneous $\mathbf{Bad}(i, j)$ is winning.

1.1. Inhomogeneous $\mathbf{Bad}(i, j)$ and our results

For $\theta = (\gamma, \delta) \in \mathbb{R}^2$, let $\mathbf{Bad}_\theta(i, j)$ denote the set of points $(x, y) \in \mathbb{R}^2$ such that

$$\max\{\|qx - \gamma\|^{\frac{1}{i}}, \|qy - \delta\|^{\frac{1}{j}}\} > c(x, y) q^{-1} \quad \forall q \in \mathbb{N}.$$

It is straight forward to deduce that $\mathbf{Bad}_\theta(i, j)$ is of measure zero from the inhomogeneous version of Khintchine's theorem [13, Theorem 1] with varying approximating functions in each co-ordinate. Surprisingly, the fact that $\dim \mathbf{Bad}_\theta(i, j) = 2$ is very much a recent development – see [10] for the symmetric $i = j = 1/2$ case and [7] for the general case.

One of the main goals of this paper is to prove the following full dimension statement which not only implies the inhomogeneous analogue of Theorem A but totally removes the \liminf condition (1.2).

Theorem 1.1. *Let (i, j) be a pair of real numbers satisfying (1.1) and let $\mathcal{C} := \mathcal{C}_f$ be a planar curve such that $f \in C^{(2)}(I)$ and that $f''(x) \neq 0$ for all $x \in I$. Then, for any $\theta = (\gamma, \delta) \in \mathbb{R}^2$ we have that $\mathbf{Bad}_\theta(i, j) \cap \mathcal{C}$ is a winning subset of \mathcal{C} .*

Remark 1. The condition $f''(x) \neq 0$ is often referred to as non-degeneracy at x . Note that if $f \in C^{(2)}(I)$ and $f''(x) \neq 0$ for some point $x \in I$, then, by continuity, there exists an interval $I^* \subset I$ such that $f''(x) \neq 0$ for all $x \in I^*$. In other words, if the curve \mathcal{C}_f is not a straight line segment then it is always possible to find an arc of \mathcal{C}_f that is non-degenerate everywhere. Thus, Theorem 1.1 with $I = I^*$ and $\theta = (0, 0)$ implies Theorem A without assuming (1.2). Of course this makes use of the well know fact that any winning set is of full dimension and that any countable intersection of winning sets is again winning.

Remark 2. Given a subset $X \subset \mathbb{R}^2$, let $\pi(X)$ denote the projection of X onto the x -axis. Regarding Theorem 1.1, we actually prove that $\pi(\mathbf{Bad}_\theta(i, j) \cap \mathcal{C})$ is a $1/2$ -winning subset of I . In other words, for the projected set we prove α -winning with the best possible winning constant; i.e. $\alpha = 1/2$. Now, since the projection map π is bi-Lipschitz on \mathcal{C} and the image of a winning set under a bi-Lipschitz map is again winning [8, Proposition 5.3], it trivially follows that $\mathbf{Bad}_\theta(i, j) \cap \mathcal{C}$ is an α_0 -winning subset of \mathcal{C} for some $\alpha_0 \in (0, 1/2]$. The actual value of α_0 is dependent on the Lipschitz constant κ associated with $(\pi|_{\mathcal{C}})^{-1}$. In fact, if we use the maximum norm on \mathbb{R}^2 it is possible to obtain the winning statement with $\alpha_0 = 1/2$. Essentially, if $\kappa > 1$ we consider the projection of $\mathbf{Bad}_\theta(i, j) \cap \mathcal{C}$ onto the y -axis rather than the x -axis.

For straight lines we prove the following counterpart statement.

Theorem 1.2. *Let (i, j) be a pair of real numbers satisfying (1.1) and given $a, b \in \mathbb{R}$ with $a \neq 0$, let $L_{a,b}$ denote the line defined by the equation $y = ax + b$. Suppose there exists $\epsilon > 0$ such that*

$$\liminf_{q \rightarrow \infty} q^{\frac{1}{\sigma} - \epsilon} \max\{\|qa\|, \|qb\|\} > 0 \quad \text{where } \sigma := \min\{i, j\}. \quad (1.4)$$

Then, for any $\theta = (\gamma, \delta) \in \mathbb{R}^2$ we have that $\mathbf{Bad}_\theta(i, j) \cap L_{a,b}$ is a winning subset of $L_{a,b}$. Moreover, if $a \in \mathbb{Q}$ the statement is true with $\epsilon = 0$ in (1.4).

Remark 3. As with curves, the theorem is actually deduced on showing that the projected set $\pi(\mathbf{Bad}_\theta(i, j) \cap L_{a,b})$ is an $1/2$ -winning subset of \mathbb{R} .

Remark 4. The Diophantine condition (1.4) imposed in the theorem is essentially optimal since

$$\mathbf{Bad}(i, j) \cap L_{a,b} = \emptyset \quad \text{if} \quad \liminf_{q \rightarrow \infty} q^{\frac{1}{\sigma}} \max\{\|qa\|, \|qb\|\} = 0. \quad (1.5)$$

To see that this is the case, assume for the moment that $\mathbf{Bad}(i, j) \cap L_{a,b}$ is nonempty. Then there exists some point $x \in \mathbb{R}$ such that $(x, ax + b) \in \mathbf{Bad}(i, j)$. In terms of the equivalent dual form representation of $\mathbf{Bad}(i, j)$ – see [3, §1.3], this means that there exists a constant $c(x) > 0$ such that

$$\max\{|A|^{\frac{1}{i}}, |B|^{\frac{1}{j}}\} |Ax + B(ax + b) + C| \geq c(x) \quad (1.6)$$

for all $A, B, C \in \mathbb{Z}$ with $(A, B) \neq (0, 0)$. Now, for any given $B \in \mathbb{N}$ we choose $A, C \in \mathbb{Z}$ such that $|Ba + A| = \|Ba\|$ and $|Bb + C| = \|Bb\|$. Then

$$\begin{aligned} |Ax + B(ax + b) + C| &= |(Ba + A)x + (Bb + C)| \leq \|Ba\| |x| + \|Bb\| \\ &\leq (1 + |x|) \max\{\|Ba\|, \|Bb\|\} \end{aligned}$$

and

$$\begin{aligned} \max\{|A|^{\frac{1}{i}}, |B|^{\frac{1}{j}}\} &\leq \max\{(|Ba| + 1)^{\frac{1}{i}}, |B|^{\frac{1}{j}}\} \\ &\leq \max\{(1 + |a|)^{\frac{1}{i}} |B|^{\frac{1}{i}}, |B|^{\frac{1}{j}}\} \leq (1 + |a|)^{\frac{1}{i}} |B|^{\frac{1}{\sigma}}. \end{aligned}$$

Thus, on combining these estimates with (1.6), it follows that

$$|B|^{\frac{1}{\sigma}} \max\{\|Ba\|, \|Bb\|\} \geq \frac{c(x)}{(1 + |a|)^{\frac{1}{i}} (1 + |x|)} \quad \forall B \in \mathbb{N}$$

and so

$$\liminf_{q \rightarrow \infty} q^{\frac{1}{\sigma}} \max\{\|qa\|, \|qb\|\} > 0.$$

This establishes (1.5). Note that in view of (1.5) and the moreover part of the theorem, the Diophantine condition (1.4) with $\epsilon = 0$ is optimal in the case a is rational.

Remark 5. The fact that $a = 0$ is excluded in the statement of the theorem is natural since as in Remark 4, on making use of the equivalent dual form representation of $\mathbf{Bad}(i, j)$ it is easily verified that

$$\mathbf{Bad}(i, j) \cap L_{0,b} = \emptyset \quad \text{if} \quad \liminf_{q \rightarrow \infty} q^{\frac{1}{j}} \|qb\| = 0.$$

On the other hand, if the above \liminf is strictly positive then it was shown in [1] that $\mathbf{Bad}(i, j) \cap L_{0,b}$ is a winning subset of the horizontal line $L_{0,b}$. By making use of the mechanism developed in this paper, it is relatively straightforward to adapt the homogeneous proof given in [1] to show that if $\liminf_{q \rightarrow \infty} q^{1/j} \|qb\| > 0$, then for any $\theta \in \mathbb{R}^2$ the set $\mathbf{Bad}_\theta(i, j) \cap L_{0,b}$ is a winning subset of the horizontal line $L_{0,b}$.

Remark 6. Observe that when it comes to intersecting countably many (i_t, j_t) pairs the Diophantine condition (1.4) imposes the condition that

$$\liminf_{q \rightarrow \infty} q^{\frac{1}{\sigma} - \epsilon} \max\{\|qa\|, \|qb\|\} > 0 \quad \text{where} \quad \sigma := \sup\{\min\{i_t, j_t\} : t \in \mathbb{N}\}. \quad (1.7)$$

This is clearly weaker than condition (1.3) imposed in Theorem B and moreover in view of Remark 4 it is essentially optimal.

The proofs of Theorem 1.1 and Theorem 1.2 rely on first establishing the homogeneous cases and then making use of a natural mechanism that we develop for generalising homogeneous winning statements to the inhomogeneous setup. This mechanism is further exploited to prove that inhomogeneous $\mathbf{Bad}(i, j)$ is winning.

Theorem 1.3. Let (i, j) be a pair of real numbers satisfying (1.1). Then, for any $\theta = (\gamma, \delta) \in \mathbb{R}^2$ we have that $\mathbf{Bad}_\theta(i, j)$ is a $(30\sqrt{2})^{-1}$ -winning subset of \mathbb{R}^2 .

Remark 7. It is worth pointing out that the winning constant $(30\sqrt{2})^{-1}$ is not optimal. Indeed, the ideas used to prove the above theorem and the argument given in [11] can be combined to show that $\mathbf{Bad}_\theta(i, j)$ is hyperplane absolute winning. This is a stronger version of winning and implies that $\mathbf{Bad}_\theta(i, j)$ is α -winning for any $\alpha < 1/2$.

2. The main strategy

In this section we outline the key steps in establishing the homogeneous case ($\theta = (0, 0)$) of [Theorem 1.1](#). The general inhomogeneous statement is obtained by appropriately adapting the homogeneous argument and is carried out in §5. To begin with observe that for any planar curve $\mathcal{C} := \mathcal{C}_f$ and $\theta \in \mathbb{R}^2$

$$\mathbf{Bad}_{\theta}^f(i, j) := \{x \in I : (x, f(x)) \in \mathbf{Bad}_{\theta}(i, j)\} = \pi(\mathbf{Bad}_{\theta}(i, j) \cap \mathcal{C}).$$

Recall, that $\pi : \mathbb{R}^2 \rightarrow \mathbb{R}$ is the projection map onto the x -axis. Also, for convenience and without loss of generality we will assume that $j \leq i$. Thus, the homogeneous case of [Theorem 1.1](#) is easily deduced from the following statement for $\mathbf{Bad}^f(i, j) := \mathbf{Bad}_{(0,0)}^f(i, j)$ – see [Remark 2](#) above for the justification.

Theorem 2.1. *Let (i, j) be a pair of real numbers satisfying $0 < j \leq i < 1$ and $i + j = 1$. Let $I \subset \mathbb{R}$ be a compact interval and $f \in C^{(2)}(I)$ such that $f''(x) \neq 0$ for all $x \in I$. Then $\mathbf{Bad}^f(i, j)$ is a $1/2$ -winning subset of I .*

At this point it is useful to recall the definition of a winning set and the notion of rooted trees – a key ‘structural’ ingredient in establishing the above winning statement.

2.1. Schmidt games and rooted trees

Wolfgang M. Schmidt introduced the games which now bear his name in [\[14\]](#). The simplified account which we are about to present is sufficient for the purposes of this paper. Suppose that $0 < \alpha < 1$ and $0 < \beta < 1$. Consider the following game involving the two arch rivals **Ayesha** and **Bhupen** – often simply referred to as players **A** and **B**. First, **B** chooses a closed ball $\mathbf{B}_0 \subset \mathbb{R}^m$. Next, **A** chooses a closed ball \mathbf{A}_0 contained in \mathbf{B}_0 of diameter $\alpha \rho(\mathbf{B}_0)$ where $\rho(\cdot)$ denotes the diameter of the ball under consideration. Then, **B** chooses at will a closed ball \mathbf{B}_1 contained in \mathbf{A}_0 of diameter $\beta \rho(\mathbf{A}_0)$. Alternating in this manner between the two players, generates a nested sequence of closed balls in \mathbb{R}^m :

$$\mathbf{B}_0 \supset \mathbf{A}_0 \supset \mathbf{B}_1 \supset \mathbf{A}_1 \supset \dots \supset \mathbf{B}_n \supset \mathbf{A}_n \supset \dots$$

with diameters

$$\rho(\mathbf{B}_n) = (\alpha \beta)^n \rho(\mathbf{B}_0) \quad \text{and} \quad \rho(\mathbf{A}_n) = \alpha \rho(\mathbf{B}_n).$$

A subset X of \mathbb{R}^m is said to be (α, β) -winning if **A** can play in such a way that the unique point of intersection

$$\bigcap_{n=0}^{\infty} \mathbf{B}_n = \bigcap_{n=0}^{\infty} \mathbf{A}_n$$

lies in X , regardless of how \mathbf{B} plays. The set X is called α -winning if it is (α, β) -winning for all $\beta \in (0, 1)$. Finally, X is simply called *winning* if it is α -winning for some α . Informally, player \mathbf{B} tries to stay away from the ‘target’ set X whilst player \mathbf{A} tries to land on X . As shown by Schmidt [14], the following are two key consequences of winning.

- If $X \subset \mathbb{R}^m$ is a winning set, then $\dim X = m$.
- The intersection of countably many α -winning sets is α -winning.

In the setting of Theorem 2.1, we have $m = 1$ and $X = \mathbf{Bad}^f(i, j)$. Thus \mathbf{A}_n and \mathbf{B}_n are compact intervals. Note that more generally, we can replace \mathbb{R}^m in the above description of Schmidt games by a m -dimensional Riemannian manifold. It is this slightly more general form that is implicitly referred to within the context of Theorem 1.1.

We now turn our attention to rooted trees. Recall that a *rooted tree* is a connected graph \mathcal{T} without cycles and with a distinguished vertex τ_0 , called the *root* of \mathcal{T} . We identify \mathcal{T} with the set of its vertices. Any vertex $\tau \in \mathcal{T}$ is connected to τ_0 by a unique path. The length of the path is called the *height* of τ . The set of vertices of height n is called the n 'th *level* of \mathcal{T} and is denoted by \mathcal{T}_n . Thus $\mathcal{T}_0 = \{\tau_0\}$. Next, given $\tau, \tau' \in \mathcal{T}$ we write $\tau \prec \tau'$ to indicate that the path between τ_0 and τ passes through τ' and in this case we call τ a *descendant* of τ' and τ' an *ancestor* of τ . By definition, every vertex is a descendant and an ancestor of itself. For $\mathcal{V} \subset \mathcal{T}$, we write $\tau \prec \mathcal{V}$ if \mathcal{V} contains an ancestor of τ . If $\tau \prec \tau'$ and the height of τ is one greater than that of τ' , then τ is called a *successor* of τ' and τ' is called the *predecessor* of τ . Let $\mathcal{T}(\tau)$ denote the rooted tree formed by all descendants of τ . Thus the root of $\mathcal{T}(\tau)$ is τ and we denote by $\mathcal{T}_{\text{suc}}(\tau)$ the set of all successors of τ . More generally, for $\mathcal{V} \subset \mathcal{T}$, we let $\mathcal{T}_{\text{suc}}(\mathcal{V}) := \bigcup_{\tau \in \mathcal{V}} \mathcal{T}_{\text{suc}}(\tau)$. In this paper, we use the convention that a subtree of \mathcal{T} has the same root as \mathcal{T} . As a consequence, $\mathcal{T}(\tau)$ is not regarded as a subtree of \mathcal{T} unless $\tau = \tau_0$.

Let $N \in \mathbb{N}$. We say that a rooted tree is N -regular if every vertex has exactly N successors. Note that an N -regular rooted tree is necessarily infinite. The following statement appears as Proposition 2.1 in [1].

Proposition 2.2. *Let \mathcal{T} be an N -regular rooted tree, $\mathcal{S} \subset \mathcal{T}$ be a subtree, and $1 \leq m \leq N$ be an integer. Suppose that for every m -regular subtree \mathcal{R} of \mathcal{T} , we have that $\mathcal{S} \cap \mathcal{R}$ is infinite. Then \mathcal{S} contains a $(N - m + 1)$ -regular subtree.*

This proposition will be required in establishing Proposition 2.4 below. As shown in §2.2.1, the latter is very much at the heart of the proof of Theorem 2.1.

2.2. The winning strategy for Theorem 2.1

Let $\beta \in (0, 1)$. We want to prove that $\mathbf{Bad}^f(i, j)$ is $(\frac{1}{2}, \beta)$ -winning. In the first round of the game, Bhupen chooses a closed interval $\mathbf{B}_0 \subset I$. Now Ayesha chooses the closed

interval $\mathbf{A}_0 \subset \mathbf{B}_0$ with diameter $\rho(\mathbf{A}_0) = \frac{1}{2}\rho(\mathbf{B}_0)$ such that \mathbf{A}_0 has the same centre as \mathbf{B}_0 . Let $\kappa > 1$ be sufficiently large so that for every $x \in I$ we have that

$$|f'(x)| \leq \kappa - 1 \quad (2.1)$$

and

$$\kappa^{-1} \leq |f''(x)| \leq \kappa. \quad (2.2)$$

Clearly such a $\kappa > 1$ exists by the conditions imposed on f and I . Let

$$R := (2\beta^{-1})^5 \quad \text{and} \quad l := \rho(\mathbf{A}_0).$$

Without loss of generality, we may assume that

$$3\kappa l R^2 < 1. \quad (2.3)$$

The point is that if this is not the case then **Ayesha** will choose her intervals \mathbf{A}_{n-1} ($n \geq 1$) arbitrarily until the interval \mathbf{B}_n chosen by **Bhupen** with diameter $\rho(\mathbf{B}_n) = (\beta/2)^n \rho(\mathbf{B}_0)$ satisfies $6\kappa \rho(\mathbf{B}_n) R^2 < 1$. At this stage **Ayesha** chooses the closed interval \mathbf{A}_n with the same centre as \mathbf{B}_n and half its diameter. We now simply relabel \mathbf{A}_n and \mathbf{B}_n as \mathbf{A}_0 and \mathbf{B}_0 respectively.

Choose $\mu > 0$ such that

$$10\kappa^2 l^{-1} R^{\frac{1}{j} - \frac{j}{6}\mu} \leq 1 \quad (2.4)$$

and define

$$\lambda_0 := 0 \quad \text{and} \quad \lambda_k := \frac{2(1+j)}{j}k + \mu \quad \text{for} \quad k \geq 1. \quad (2.5)$$

In turn, let

$$c := \frac{l^2}{10^3 \kappa^5 R^{4+\lambda_1}} \quad (2.6)$$

and

$$\mathcal{P} := \left\{ P = \left(\frac{p}{q}, \frac{r}{q} \right) : \frac{p}{q} \in I, \left| f\left(\frac{p}{q} \right) - \frac{r}{q} \right| < \frac{\kappa c}{q^{1+j}} \right\}. \quad (2.7)$$

Note that R is large while l and c are small. Indeed, we have the inequalities

$$R > 32, \quad l < 10^{-3}, \quad c < 10^{-10}l$$

which we will use without further reference. Throughout, when a rational point in \mathbb{R}^2 is expressed as $P = (\frac{p}{q}, \frac{r}{q})$, we assume that $q > 0$ and that the integers p, q, r are co-prime. Finally, for each $P = (\frac{p}{q}, \frac{r}{q}) \in \mathbb{Q}^2$ we associate the interval

$$\Delta(P) := \left\{ x \in I : \left| x - \frac{p}{q} \right| < \frac{c}{q^{1+i}} \right\}. \quad (2.8)$$

The following inclusion is a simple consequence of the manner in which the above quantities and objects have been defined.

Lemma 2.3. *Let \mathbf{A}_0 , \mathcal{P} and $\Delta(P)$ be as above. Then*

$$\mathbf{A}_0 \setminus \bigcup_{P \in \mathcal{P}} \Delta(P) \subset \mathbf{Bad}^f(i, j).$$

Proof. Let $x \in \mathbf{A}_0$. Suppose $x \notin \mathbf{Bad}^f(i, j)$. Then there exists $P = (\frac{p}{q}, \frac{r}{q}) \in \mathbb{Q}^2$ such that

$$\left| x - \frac{p}{q} \right| < \frac{c}{q^{1+i}}, \quad \left| f(x) - \frac{r}{q} \right| < \frac{c}{q^{1+j}}.$$

In view of the fact that

$$\left| x - \frac{p}{q} \right| < \frac{c}{q^{1+i}} \leq c \leq \frac{l}{2},$$

it follows that $\frac{p}{q} \in \mathbf{B}_0 \subset I$. Hence, using the Mean Value Theorem together with (2.1) we obtain the following estimate:

$$\begin{aligned} \left| f\left(\frac{p}{q}\right) - \frac{r}{q} \right| &\leq \left| f\left(\frac{p}{q}\right) - f(x) \right| + \left| f(x) - \frac{r}{q} \right| \\ &\leq (\kappa - 1) \left| x - \frac{p}{q} \right| + \frac{c}{q^{1+j}} \\ &< \frac{(\kappa - 1)c}{q^{1+i}} + \frac{c}{q^{1+j}} \leq \frac{\kappa c}{q^{1+j}}. \end{aligned}$$

The upshot is that $x \in \Delta(P)$ with $P \in \mathcal{P}$. This completes the proof of the lemma. \square

Now let \mathcal{T} be an $[R]$ -regular rooted tree with root τ_0 , where $[\cdot]$ denotes the integer part of a real number. We choose and fix an injective map \mathcal{I} from \mathcal{T} to the set of closed subintervals of \mathbf{A}_0 satisfying the following conditions:

- For any $n \geq 0$ and $\tau \in \mathcal{T}_n$, $\rho(\mathcal{I}(\tau)) = lR^{-n}$. In particular, $\mathcal{I}(\tau_0) = \mathbf{A}_0$.
- For $\tau, \tau' \in \mathcal{T}$, if $\tau \prec \tau'$, then $\mathcal{I}(\tau) \subset \mathcal{I}(\tau')$.
- For any $\tau \in \mathcal{T}$, the interiors of the intervals $\{\mathcal{I}(\tau') : \tau' \in \mathcal{T}_{\text{suc}}(\tau)\}$ are mutually disjoint, and $\bigcup_{\tau' \in \mathcal{T}_{\text{suc}}(\tau)} \mathcal{I}(\tau')$ is connected.

Note that for $n \geq 1$ and $\tau \in \mathcal{T}_{n-1}$, any closed subinterval of $\mathcal{I}(\tau)$ of length $2lR^{-n}$ must contain an $\mathcal{I}(\tau')$ for some $\tau' \in \mathcal{T}_{\text{suc}}(\tau)$. Suppose that \mathcal{P} is partitioned into a disjoint union

$$\mathcal{P} = \bigcup_{n=1}^{\infty} \mathcal{P}_n.$$

We inductively define a subtree \mathcal{S} of \mathcal{T} as follows. Let $\mathcal{S}_0 = \{\tau_0\}$. If \mathcal{S}_{n-1} ($n \geq 1$) is defined, we let

$$\mathcal{S}_n := \left\{ \tau \in \mathcal{T}_{\text{suc}}(\mathcal{S}_{n-1}) : \mathcal{I}(\tau) \cap \bigcup_{P \in \mathcal{P}_n} \Delta(P) = \emptyset \right\}.$$

Then

$$\mathcal{S} := \bigcup_{n=0}^{\infty} \mathcal{S}_n$$

is a subtree of \mathcal{T} and by construction

$$\mathcal{I}(\tau) \subset \mathbf{A}_0 \setminus \bigcup_{P \in \mathcal{P}_n} \Delta(P) \quad \forall n \geq 1 \quad \text{and} \quad \tau \in \mathcal{S}_n. \quad (2.9)$$

Thus given a partition \mathcal{P}_n of \mathcal{P} , the intervals $\{\mathcal{I}(\tau) : \tau \in \mathcal{S}_n\}$ serve as possible candidates when it comes to **Ayesha** to turn to make a move. Moreover, we are able to choose the partition \mathcal{P}_n in such a way that \mathcal{S} has the following key feature.

Proposition 2.4. *There exists a partition $\mathcal{P} = \bigcup_{n=1}^{\infty} \mathcal{P}_n$ such that the tree \mathcal{S} has an $([R] - 10)$ -regular subtree.*

Armed with this proposition we are able to describe the winning strategy that **Ayesha** will adopt.

2.2.1. Proof of Theorem 2.1 modulo Proposition 2.4

Let $\mathcal{P} = \bigcup_{n=1}^{\infty} \mathcal{P}_n$ be a partition such that \mathcal{S} has an $([R] - 10)$ -regular subtree, say \mathcal{S}' . We inductively prove that for every $n \geq 0$,

$$\text{Ayesha can choose } \mathbf{A}_{5n} = \mathcal{I}(\tau_n) \text{ for some } \tau_n \in \mathcal{S}'_n. \quad (2.10)$$

Since $\mathbf{A}_0 = \mathcal{I}(\tau_0)$, we trivially have that (2.10) holds when $n = 0$. Assume $n \geq 1$ and that **Ayesha** has chosen $\mathbf{A}_{5(n-1)} = \mathcal{I}(\tau_{n-1})$, where $\tau_{n-1} \in \mathcal{S}'_{n-1}$. We refer to the intervals $\{\mathcal{I}(\tau) : \tau \in \mathcal{T}_{\text{suc}}(\tau_{n-1}) \setminus \mathcal{S}'_{\text{suc}}(\tau_{n-1})\}$ as dangerous intervals – they represent intervals that **Ayesha** needs to avoid. We first prove that

$$\begin{aligned} &\text{For } t \in \{0, 1, 2, 3, 4\}, \text{ Ayesha can play so that } \mathbf{A}_{5(n-1)+t}, \text{ and hence} \\ &\mathbf{B}_{5(n-1)+t+1}, \text{ contains at most } [10 \cdot 2^{-t}] \text{ dangerous intervals.} \end{aligned} \quad (2.11)$$

If $t = 0$, there is nothing to prove. Assume $1 \leq t \leq 4$ and (2.11) holds if t is replaced by $t - 1$. Thus $\mathbf{B}_{5(n-1)+t}$ contains at most $[10 \cdot 2^{-t+1}]$ dangerous intervals. Divide $\mathbf{B}_{5(n-1)+t}$

into two closed subintervals of equal length. Then Ayesha can choose $\mathbf{A}_{5(n-1)+t}$ to be one of the subintervals so that it contains at most $\left\lceil \frac{1}{2}[10 \cdot 2^{-t+1}] \right\rceil \leq [10 \cdot 2^{-t}]$ dangerous intervals. This proves (2.11). By letting $t = 4$ in (2.11), we see that Ayesha can play so that \mathbf{B}_{5n} contains no dangerous intervals. Since \mathbf{B}_{5n} has length $2lR^{-n}$, it contains an $\mathcal{I}(\tau_n)$ for some $\tau_n \in \mathcal{T}_{\text{suc}}(\tau_{n-1})$. It follows that $\tau_n \in \mathcal{S}'_n$. So Ayesha can choose $\mathbf{A}_{5n} = \mathcal{I}(\tau_n)$. This completes the proof of (2.10). In view of (2.10), (2.9) and Lemma 2.3, we have

$$\begin{aligned} \bigcap_{n=0}^{\infty} \mathbf{A}_n &= \bigcap_{n=1}^{\infty} \mathbf{A}_{5n} = \bigcap_{n=1}^{\infty} \mathcal{I}(\tau_n) \subset \bigcap_{n=1}^{\infty} \mathbf{A}_0 \setminus \bigcup_{P \in \mathcal{P}_n} \Delta(P) \\ &= \mathbf{A}_0 \setminus \bigcup_{P \in \mathcal{P}} \Delta(P) \subset \mathbf{Bad}^f(i, j). \end{aligned}$$

This proves the theorem assuming the truth of Proposition 2.4. \square

3. Preliminaries for Proposition 2.4

The following simple but important lemma was established in [3].

Lemma 3.1. *For any point $P = (\frac{p}{q}, \frac{r}{q}) \in \mathbb{Q}^2$ there exist coprime integers A, B, C with $(A, B) \neq (0, 0)$ such that*

$$\begin{aligned} Ap + Br + Cq &= 0, \\ |A| &\leq q^i \quad \text{and} \quad |B| \leq q^j. \end{aligned}$$

Proof. Since the proof is only a few lines we reproduce it here. By Minkowski's theorem for systems of linear forms there is $(A, B, C) \in \mathbb{Z}^3 \setminus \{0\}$ such that

$$|Ap + Br + Cq| < 1, \quad |A| \leq q^i \quad \text{and} \quad |B| \leq q^j.$$

Since $Ap + Br + Cq$ is an integer it must be zero. If $(A, B) = (0, 0)$ then $qC = 0$ and, since $q \neq 0$ we also have that $C = 0$, a contradiction. Hence $(A, B) \neq (0, 0)$ and the proof is complete. \square

In view of Lemma 3.1, to each point $P = (\frac{p}{q}, \frac{r}{q}) \in \mathcal{P}$, we can assign a rational line

$$L_P := \{(x, y) \in \mathbb{R}^2 : A_P x + B_P y + C_P = 0\} \tag{3.1}$$

passing through P where $A_P, B_P, C_P \in \mathbb{Z}$ are co-prime with $(A_P, B_P) \neq (0, 0)$ and such that

$$|A_P| \leq q^i \quad \text{and} \quad |B_P| \leq q^j. \tag{3.2}$$

If there is more than one line satisfying the above conditions, we choose any one. Further, for $P \in \mathcal{P}$ we define the function $F_P : I \rightarrow \mathbb{R}$ by

$$F_P(x) := A_P x + B_P f(x) + C_P$$

and the set

$$\Theta(P) := \left\{ x \in I : |F_P(x)| < \frac{2\kappa c}{q} \right\}. \quad (3.3)$$

In this section we gather basic information regarding the sets $\Theta(P)$ and associated quantities.

Lemma 3.2. *Let $P = (\frac{p}{q}, \frac{r}{q}) \in \mathcal{P}$ and $x \in \Delta(P)$. Then*

$$\left| F_P\left(\frac{p}{q}\right) \right| < \frac{\kappa c}{q} \quad \text{and} \quad \left| F_P(x) - F_P\left(\frac{p}{q}\right) \right| < \frac{\kappa c}{q}.$$

In particular, we have that $\Delta(P) \subset \Theta(P)$. Furthermore, if $x \in \Theta(P)$ then

$$\left| F_P(x) - F_P\left(\frac{p}{q}\right) \right| < \frac{3\kappa c}{q}.$$

Proof. On using the fact that P lies on the line L_P and (2.7) we obtain the following

$$\left| F_P\left(\frac{p}{q}\right) \right| = \left| B_P \left(f\left(\frac{p}{q}\right) - \frac{r}{q} \right) \right| \leq q \left| f\left(\frac{p}{q}\right) - \frac{r}{q} \right| < \frac{\kappa c}{q}.$$

For the second inequality, by the Mean Value Theorem there exists a point $\xi \in I$ such that

$$\left| F_P(x) - F_P\left(\frac{p}{q}\right) \right| = |F'_P(\xi)| \left| x - \frac{p}{q} \right| < (|A_P| + (\kappa - 1)|B_P|) \frac{c}{q^{1+i}} \leq \frac{\kappa c}{q}.$$

In particular, it follows that for $x \in \Delta(P)$, we have that

$$|F_P(x)| \leq \left| F_P\left(\frac{p}{q}\right) \right| + \left| F_P(x) - F_P\left(\frac{p}{q}\right) \right| < \frac{2\kappa c}{q}$$

and so by definition $x \in \Theta(P)$.

Finally, if $x \in \Theta(P)$, then

$$\left| F_P(x) - F_P\left(\frac{p}{q}\right) \right| \leq |F_P(x)| + \left| F_P\left(\frac{p}{q}\right) \right| < \frac{3\kappa c}{q}. \quad \square$$

Our next goal is to describe the structure of $\Theta(P)$, in particular, to estimate its size. To this end, we introduce the following quantities. Let

$$E_P := F'_P\left(\frac{p}{q}\right) = A_P + B_P f'\left(\frac{p}{q}\right) \quad (3.4)$$

and let

$$\mathcal{P}^* := \left\{ P = \left(\frac{p}{q}, \frac{r}{q} \right) \in \mathcal{P} : qE_P^2 < 9\kappa^2 c |B_P| \right\}. \quad (3.5)$$

Then we define the height of P by

$$H(P) := \max\{|q|E_P|, 3\kappa\sqrt{cq|B_P|}\} = \begin{cases} 3\kappa\sqrt{cq|B_P|} & \text{if } P \in \mathcal{P}^* \\ |q|E_P| & \text{if } P \in \mathcal{P} \setminus \mathcal{P}^*. \end{cases} \quad (3.6)$$

Lemma 3.3. Let $P = (\frac{p}{q}, \frac{r}{q}) \in \mathcal{P}$. Then

$$|E_P| \leq \kappa q^i \quad (3.7)$$

$$H(P) \leq \kappa q^{1+i} \quad (3.8)$$

Proof. In view of (2.1) and the fact that $j \leq i$, it follows that

$$|E_P| \leq |A_P| + (\kappa - 1)|B_P| \leq q^i + (\kappa - 1)q^j \leq \kappa q^i,$$

and hence

$$H(P) = \max\{|q|E_P|, 3\kappa\sqrt{cq|B_P|}\} \leq \max\{\kappa q^{1+i}, 3\kappa c^{\frac{1}{2}} q^{\frac{1+i}{2}}\} = \kappa q^{1+i}. \quad \square$$

In the above, to each $P \in \mathcal{P}$ we have attached a rational line L_P with coefficients A_P, B_P, C_P and the function F_P and the quantity E_P . For ease of notation and clarity, we shall drop the subscript P if there is no ambiguity or confusion caused.

Lemma 3.4. If $P \in \mathcal{P}^*$ (resp. $P \in \mathcal{P} \setminus \mathcal{P}^*$), then $\Theta(P)$ is contained in one (resp. at most two) open interval(s) of length at most

$$\frac{42\kappa^3 c}{H(P)}.$$

Proof. Case (1). Suppose $P \in \mathcal{P}^*$. Then $B \neq 0$. For any $x \in \Theta(P)$, it follows from Lemma 3.2 and (3.5) that

$$\begin{aligned} \frac{3\kappa c}{q} &> \left| F(x) - F\left(\frac{p}{q}\right) \right| = \left| E\left(x - \frac{p}{q}\right) + \frac{F''(\xi)}{2} \left(x - \frac{p}{q}\right)^2 \right| \\ &\geq \frac{|B|}{2\kappa} \left| x - \frac{p}{q} \right|^2 - 3\kappa \sqrt{\frac{c|B|}{q}} \left| x - \frac{p}{q} \right|, \end{aligned}$$

where $\xi \in I$. This implies that

$$\left| x - \frac{p}{q} \right| < 7\kappa^2 \sqrt{\frac{c}{q|B|}} = \frac{21\kappa^3 c}{H(P)}.$$

The upshot of this is that $\Theta(P)$ is contained in an open interval of length $42\kappa^3 c/H(P)$.

Case (2). Suppose $P \in \mathcal{P} \setminus \mathcal{P}^*$. If $B = 0$, then $E = A \neq 0$ and $F(x) = Ex + C$. It follows that $\Theta(P)$ is an open interval of length

$$\frac{4\kappa c}{q|E|} = \frac{4\kappa c}{H(P)} < \frac{42\kappa^3 c}{H(P)}.$$

Now suppose $B \neq 0$. Then, by (3.5), we have that $E \neq 0$. Consider the closed interval

$$\Omega := \left\{ x \in I : \left| x - \frac{p}{q} \right| \leq \frac{|E|}{\kappa|B|} \right\}.$$

We first prove that $\Theta(P) \cap \Omega$ is contained in an open interval of length $42\kappa^3 c/H(P)$. If $x \in \Theta(P) \cap \Omega$, then it follows from Lemma 3.2 that

$$\begin{aligned} \frac{3\kappa c}{q} &> \left| F(x) - F\left(\frac{p}{q}\right) \right| = \left| E\left(x - \frac{p}{q}\right) + \frac{1}{2}Bf''(\xi)\left(x - \frac{p}{q}\right)^2 \right| \\ &\geq \left(|E| - \frac{\kappa|B|}{2} \left| x - \frac{p}{q} \right| \right) \left| x - \frac{p}{q} \right| \geq \frac{|E|}{2} \left| x - \frac{p}{q} \right|. \end{aligned}$$

This implies that

$$\left| x - \frac{p}{q} \right| < \frac{6\kappa c}{q|E|} = \frac{6\kappa c}{H(P)}. \quad (3.9)$$

Hence $\Theta(P) \cap \Omega$ is contained in an open interval of length $12\kappa c/H(P) < 42\kappa^3 c/H(P)$. In particular, this implies the desired statement if $\Theta(P) \subset \Omega$.

Suppose that $\Theta(P) \not\subset \Omega$. Then the following three observations imply that $\Theta(P) \setminus \Omega$ is a connected interval.

- $|F(\frac{p}{q})| < \kappa c/q$ by Lemma 3.2.
- If x_0 is an end point of Ω and is contained in the interior of I , then $|F(x_0)| \geq 2\kappa c/q$. To see this, note that the assumption on x_0 and (3.5) imply that

$$\left| x_0 - \frac{p}{q} \right| = \frac{|E|}{\kappa|B|} > \frac{6\kappa c}{q|E|},$$

which together with (3.9) implies that $x_0 \notin \Theta(P)$. Hence the desired inequality follows from (3.3).

- The function $F : I \rightarrow \mathbb{R}$ is either convex or concave.

We next claim that

$$|F'(\xi)| \geq \frac{|E|}{2\kappa} \quad \text{for any } \xi \in \Theta(P) \setminus \Omega. \quad (3.10)$$

Assuming this claim for the moment, it follows that for any $x_1, x_2 \in \Theta(P) \setminus \Omega$

$$\frac{4\kappa c}{q} > |F(x_1) - F(x_2)| = |F'(\xi)| |x_1 - x_2| \geq \frac{|E|}{2\kappa} |x_1 - x_2|,$$

where $\xi \in \Theta(P) \setminus \Omega$. Thus

$$|x_1 - x_2| < \frac{8\kappa^2 c}{q|E|} = \frac{8\kappa^2 c}{H(P)}.$$

Hence $\Theta(P) \setminus \Omega$ is contained in an open interval of length $8\kappa^2 c/H(P) < 42\kappa^3 c/H(P)$ and thereby completes the proof of the lemma modulo (3.10).

We now prove (3.10) in the instance that $f'' > 0$ and $B > 0$. The other cases are similar and left to the reader. Let $\xi \in \Theta(P) \setminus \Omega$, and consider the functions $G : \mathbb{R} \rightarrow \mathbb{R}$ and $H : \mathbb{R} \rightarrow \mathbb{R}$ given by

$$G(x) := \frac{\kappa B}{2} \left(x - \frac{p}{q}\right)^2 + E \left(x - \frac{p}{q}\right) + F\left(\frac{p}{q}\right)$$

and

$$H(x) := \frac{B}{2\kappa} (x - \xi)^2 + F'(\xi)(x - \xi) + F(\xi).$$

It follows that $G(\frac{p}{q}) = F(\frac{p}{q})$, $G'(\frac{p}{q}) = F'(\frac{p}{q})$, $H(\xi) = F(\xi)$, and $H'(\xi) = F'(\xi)$. Moreover, if $x \in I$, then

$$\begin{aligned} G''(x) - F''(x) &= B(\kappa - f''(x)) \geq 0, \\ H''(x) - F''(x) &= B(\kappa^{-1} - f''(x)) \leq 0. \end{aligned}$$

Thus, for $x \in I$ we have that $G(x) \geq F(x) \geq H(x)$. It follows that for any $x \in I$,

$$G(x) \geq H(x) \geq -\frac{\kappa F'(\xi)^2}{2B} + F(\xi) \geq -\frac{\kappa F'(\xi)^2}{2B} - \frac{2\kappa c}{q}. \quad (3.11)$$

Note that $F'' = Bf'' > 0$. So the end point of Ω that is contained in the interval with end points $\frac{p}{q}$ and ξ is equal to $\frac{p}{q} - \frac{E}{\kappa B}$. In particular, we have that $\frac{p}{q} - \frac{E}{\kappa B} \in I$. Thus (3.11) implies that

$$G\left(\frac{p}{q} - \frac{E}{\kappa B}\right) \geq -\frac{\kappa F'(\xi)^2}{2B} - \frac{2\kappa c}{q}.$$

On the other hand, we have that

$$G\left(\frac{p}{q} - \frac{E}{\kappa B}\right) = -\frac{E^2}{2\kappa B} + F\left(\frac{p}{q}\right) \leq -\frac{E^2}{2\kappa B} + \frac{\kappa c}{q}.$$

On combining the previous two inequalities, we find that

$$F'(\xi)^2 \geq \frac{E^2}{\kappa^2} - \frac{6cB}{q} \geq \frac{E^2}{\kappa^2} - \frac{2E^2}{3\kappa^2} = \frac{E^2}{3\kappa^2}$$

and (3.10) follows. This completes the proof of Lemma 3.4. \square

4. Proof of Proposition 2.4

For $n \geq 1$, let

$$H_n := 42\kappa^3 cl^{-1} R^n$$

and

$$\mathcal{P}_n := \left\{ P = \left(\frac{p}{q}, \frac{r}{q} \right) \in \mathcal{P} : H_n \leq H(P) < H_{n+1} \right\}. \quad (4.1)$$

Note that if $P \in \mathcal{P}_n$, then, by (3.8), we have that

$$\kappa q^{1+i} \geq H_n. \quad (4.2)$$

Next let

$$\mathcal{P}_{n,0} := \mathcal{P}_n \cap \mathcal{P}^* \quad (4.3)$$

and

$$\mathcal{P}_{n,k} := \{ P \in \mathcal{P}_n \setminus \mathcal{P}^* : H_n R^{\lambda_{k-1}} \leq \kappa q^{1+i} \leq H_n R^{\lambda_k} \} \quad \text{for } 1 \leq k \leq n, \quad (4.4)$$

where λ_k are defined by (2.5).

Lemma 4.1. *With \mathcal{P}_n and $\mathcal{P}_{n,k}$ as above, we have that*

$$\mathcal{P} = \bigcup_{n=1}^{\infty} \mathcal{P}_n \quad \text{and} \quad \mathcal{P}_n = \bigcup_{k=0}^n \mathcal{P}_{n,k}.$$

Proof. It is easily verified via (2.6) that $H(P) \geq 3\kappa c^{\frac{1}{2}}$ for any $P \in \mathcal{P}$, and that $H_1 = 42\kappa^3 cl^{-1}R \leq 3\kappa c^{\frac{1}{2}}$. Hence, $\mathcal{P} = \bigcup_{n=1}^{\infty} \mathcal{P}_n$.

Since for $P \in \mathcal{P}_n \setminus \mathcal{P}^*$ we have

$$q = \frac{H(P)^2}{qE^2} \leq \frac{H_{n+1}^2}{9\kappa^2 c} = \frac{H_n^2 R^2}{9\kappa^2 c},$$

it follows from (2.4) that

$$\begin{aligned} \frac{\kappa q^{1+i}}{H_n R^{\lambda_n}} &\leq \kappa \frac{q^2}{H_n} R^{-\lambda_n} \leq \kappa \frac{H_n^3 R^4}{81\kappa^4 c^2} R^{-\lambda_n} \\ &= \frac{42^3}{81} \kappa^6 cl^{-3} R^{-(\frac{2(1+i)}{j}-3)n+4-\mu} \\ &\leq (10\kappa^2 l^{-1} R^{2-\frac{\mu}{3}})^3 \leq 1. \end{aligned}$$

This together with (4.2) implies that $\mathcal{P}_n = \bigcup_{k=0}^n \mathcal{P}_{n,k}$. \square

We claim that the partition of \mathcal{P} given by Lemma 4.1 satisfies the requirement of Proposition 2.4. In other words, \mathcal{P} gives rise to a tree \mathcal{S} as described in §2.2 that contains an $([R] - 10)$ -regular subtree. Recall that \mathcal{S} is itself a subtree of an $[R]$ -regular rooted tree \mathcal{T} . The key towards establishing the claim is the following lemma and its corollary. For $k \geq 0$, we let

$$k^+ := \max\{k, 1\}.$$

The following lemma contains a crucial property of the lines L_P defined by (3.1).

Lemma 4.2. *For any $n \geq 1$, $0 \leq k \leq n$ and $\tau \in \mathcal{S}_{n-k^+}$, the map $P \mapsto L_P$ is constant on*

$$\mathcal{P}_{n,k}(\tau) := \{P \in \mathcal{P}_{n,k} : \mathcal{I}(\tau) \cap \Delta(P) \neq \emptyset\}.$$

We postpone the proof for the moment and continue by stating an important consequence of the lemma.

Corollary 4.3. *For any $n \geq 1$, $0 \leq k \leq n$ and $\tau \in \mathcal{S}_{n-k^+}$, we have*

$$\#\left\{\tau' \in \mathcal{T}_n : \mathcal{I}(\tau') \cap \bigcup_{P \in \mathcal{P}_{n,k}(\tau)} \Delta(P) \neq \emptyset\right\} \leq \begin{cases} 2, & \text{if } k = 0, \\ 4, & \text{if } k \geq 1. \end{cases}$$

Proof. We may assume that $\mathcal{P}_{n,k}(\tau) \neq \emptyset$. Let $P_0 = (\frac{p_0}{q_0}, \frac{r_0}{q_0}) \in \mathcal{P}_{n,k}(\tau)$ be such that $q_0 \leq q$ for any $(\frac{p}{q}, \frac{r}{q}) \in \mathcal{P}_{n,k}(\tau)$. By Lemma 4.2, for any $P \in \mathcal{P}_{n,k}(\tau)$ we have $L_P = L_{P_0}$ and so $\Theta(P) \subset \Theta(P_0)$. Thus it follows from Lemma 3.2 that

$$\bigcup_{P \in \mathcal{P}_{n,k}(\tau)} \Delta(P) \subset \bigcup_{P \in \mathcal{P}_{n,k}(\tau)} \Theta(P) \subset \Theta(P_0).$$

Hence

$$\left\{ \tau' \in \mathcal{T}_n : \mathcal{I}(\tau') \cap \bigcup_{P \in \mathcal{P}_{n,k}(\tau)} \Delta(P) \neq \emptyset \right\} \subset \{ \tau' \in \mathcal{T}_n : \mathcal{I}(\tau') \cap \Theta(P_0) \neq \emptyset \}.$$

By [Lemma 3.4](#), if $k = 0$ (resp. $k \geq 1$), then $\Theta(P_0)$ is contained in one (resp. at most two) open interval(s) of length at most

$$\frac{42\kappa^3 c}{H(P_0)} \stackrel{(4.1)}{\leq} \frac{42\kappa^3 c}{H_n} = lR^{-n}.$$

Since the intervals $\{\mathcal{I}(\tau') : \tau' \in \mathcal{T}_n\}$ are of length lR^{-n} and have mutually disjoint interiors, there can be at most 2 (resp. 4) of them that intersect $\Theta(P_0)$. This proves the corollary. \square

We are now in the position to prove [Proposition 2.4](#). In view of [Proposition 2.2](#), it suffices to prove that the intersection of \mathcal{S} with every 11-regular subtree of \mathcal{T} is infinite. Let $\mathcal{R} \subset \mathcal{T}$ be an 11-regular subtree and let

$$\mathcal{R}' := \mathcal{R} \cap \mathcal{S} \quad \text{and} \quad a_n := \#\mathcal{R}'_n \quad (n \geq 0),$$

where \mathcal{R}'_n is the n 'th level of the tree \mathcal{R}' . Then $a_0 = 1$. We prove that \mathcal{R}' is infinite by showing that

$$a_n > 3a_{n-1} \quad (n \geq 1). \tag{4.5}$$

We use induction. For $n \geq 1$, let

$$\mathcal{U}_n := \left\{ \tau \in \mathcal{T}_{\text{suc}}(\mathcal{R}'_{n-1}) : \mathcal{I}(\tau) \cap \bigcup_{P \in \mathcal{P}_n} \Delta(P) \neq \emptyset \right\}.$$

Then

$$\mathcal{R}'_n = \left\{ \tau \in \mathcal{R}_{\text{suc}}(\mathcal{R}'_{n-1}) : \mathcal{I}(\tau) \cap \bigcup_{P \in \mathcal{P}_n} \Delta(P) = \emptyset \right\} = \mathcal{R}_{\text{suc}}(\mathcal{R}'_{n-1}) \setminus \mathcal{U}_n.$$

It follows that

$$a_n \geq 11a_{n-1} - \#\mathcal{U}_n. \tag{4.6}$$

On the other hand,

$$\begin{aligned}
\mathcal{U}_n &= \bigcup_{k=0}^n \left\{ \tau \in \mathcal{T}_{\text{suc}}(\mathcal{R}'_{n-1}) : \mathcal{I}(\tau) \cap \bigcup_{P \in \mathcal{P}_{n,k}} \Delta(P) \neq \emptyset \right\} \\
&\subset \bigcup_{k=0}^n \left\{ \tau \in \mathcal{T}_n : \tau \prec \mathcal{R}'_{n-k+}, \mathcal{I}(\tau) \cap \bigcup_{P \in \mathcal{P}_{n,k}} \Delta(P) \neq \emptyset \right\} \\
&= \bigcup_{k=0}^n \bigcup_{\tau' \in \mathcal{R}'_{n-k+}} \left\{ \tau \in \mathcal{T}_n : \tau \prec \tau', \mathcal{I}(\tau) \cap \bigcup_{P \in \mathcal{P}_{n,k}} \Delta(P) \neq \emptyset \right\} \\
&\subset \bigcup_{k=0}^n \bigcup_{\tau' \in \mathcal{R}'_{n-k+}} \left\{ \tau \in \mathcal{T}_n : \mathcal{I}(\tau) \cap \bigcup_{P \in \mathcal{P}_{n,k}(\tau')} \Delta(P) \neq \emptyset \right\}.
\end{aligned}$$

Thus, [Corollary 4.3](#) implies that

$$\#\mathcal{U}_n \leq 2a_{n-1} + \sum_{k=1}^n 4a_{n-k}. \quad (4.7)$$

On combining [\(4.6\)](#) and [\(4.7\)](#), we obtain that

$$a_n \geq 9a_{n-1} - \sum_{k=1}^n 4a_{n-k}. \quad (4.8)$$

With $n = 1$ in [\(4.8\)](#), we find that $a_1 \geq 5$. Hence, [\(4.5\)](#) holds for $n = 1$. Now assume $n \geq 2$ and that [\(4.5\)](#) holds with n replaced by $1, \dots, n-1$. Then for any $1 \leq k \leq n$, we have that

$$a_{n-k} \leq 3^{-k+1} a_{n-1}.$$

Substituting this into [\(4.8\)](#), gives that

$$a_n \geq 9a_{n-1} - 4a_{n-1} \sum_{k=1}^n 3^{-k+1} > 3a_{n-1}.$$

This completes the induction step and thus establishes [\(4.5\)](#). In turn this completes the proof of [Proposition 2.4](#) modulo the truth of [Lemma 4.2](#). \square

4.1. Proof of [Lemma 4.2](#)

To begin with we prove the following result.

Lemma 4.4. *Let n, k and τ be as in [Lemma 4.2](#), and let $P_1 := (\frac{p_1}{q_1}, \frac{r_1}{q_1}), P_2 := (\frac{p_2}{q_2}, \frac{r_2}{q_2}) \in \mathcal{P}_{n,k}(\tau)$. Denote $A_s = A_{P_s}, B_s = B_{P_s}, C_s = C_{P_s}$ and $F_s = F_{P_s}, s = 1, 2$.*

(1) If $k = 0$ and $q_1 \leq q_2$, then

$$A_2 p_1 + B_2 r_1 + C_2 q_1 = 0 \quad (4.9)$$

$$|A_1 B_2 - A_2 B_1| < q_1. \quad (4.10)$$

(2) If $k = 1$, then

$$A_2 p_1 + B_2 r_1 + C_2 q_1 = 0 = A_1 p_2 + B_1 r_2 + C_1 q_2. \quad (4.11)$$

(3) If $k \geq 2$, then

$$\left| A_1 F_2 \left(\frac{p_1}{q_1} \right) - A_2 F_1 \left(\frac{p_1}{q_1} \right) \right| \leq \frac{c}{2} H_n^{-\frac{j}{1+i}}, \quad (4.12)$$

$$\left| B_1 F_2 \left(\frac{p_1}{q_1} \right) - B_2 F_1 \left(\frac{p_1}{q_1} \right) \right| \leq \frac{c}{2} H_n^{-\frac{i}{1+i}}, \quad (4.13)$$

$$|A_1 B_2 - A_2 B_1| \leq \frac{1}{3\kappa} H_n^{\frac{1}{1+i}} R^{-k}. \quad (4.14)$$

Proof. We first prove that for any $0 \leq k \leq n$,

$$\left| \frac{p_1}{q_1} - \frac{p_2}{q_2} \right| \leq 2lR^{-n+k^+}, \quad (4.15)$$

$$\max \left\{ q_2 \left| F_2 \left(\frac{p_1}{q_1} \right) \right|, q_1 \left| F_1 \left(\frac{p_2}{q_2} \right) \right| \right\} \leq 477\kappa^5 c R^{2k^++2}, \quad (4.16)$$

$$|A_2 p_1 + B_2 r_1 + C_2 q_1| \leq \frac{1}{2} R^{-4-\lambda_1} \left(\frac{q_1}{q_2} R^{2k^++2} + \frac{|B_2|}{q_1^j} \right), \quad (4.17)$$

$$|A_1 p_2 + B_1 r_2 + C_1 q_2| \leq \frac{1}{2} R^{-4-\lambda_1} \left(\frac{q_2}{q_1} R^{2k^++2} + \frac{|B_1|}{q_2^j} \right), \quad (4.18)$$

$$|A_1 B_2 - A_2 B_1| \leq |E_1 B_2| + |E_2 B_1| + 2|B_1 B_2| \kappa l R^{-n+k^+}, \quad (4.19)$$

where $E_1 = E_{P_1}$ and $E_2 = E_{P_2}$.

To establish (4.15), note that for $s = 1, 2$ there exists $x_s \in \mathcal{I}(\tau) \cap \Delta(P_s) \neq \emptyset$ and that $|x_1 - x_2| \leq \rho(\mathcal{I}(\tau)) = lR^{-n+k^+}$. Then, it follows from (2.8) that

$$\begin{aligned} \left| \frac{p_1}{q_1} - \frac{p_2}{q_2} \right| &\leq \left| \frac{p_1}{q_1} - x_1 \right| + |x_1 - x_2| + \left| x_2 - \frac{p_2}{q_2} \right| \\ &\leq \frac{c}{q_1^{1+i}} + lR^{-n+k^+} + \frac{c}{q_2^{1+i}} \stackrel{(4.2)}{\leq} \frac{2\kappa c}{H_n} + lR^{-n+k^+} \\ &= \frac{1}{21\kappa^2} lR^{-n} + lR^{-n+k^+} \leq 2lR^{-n+k^+}. \end{aligned}$$

Regarding (4.16), we expand $F_2(\frac{p_1}{q_1})$ using Taylor's formula at the point $\frac{p_2}{q_2}$ and estimate as follows

$$\begin{aligned}
 q_2 \left| F_2\left(\frac{p_1}{q_1}\right) \right| &\leq q_2 \left| F_2\left(\frac{p_2}{q_2}\right) \right| + q_2 |E_2| \left| \frac{p_1}{q_1} - \frac{p_2}{q_2} \right| + \frac{\kappa q_2 |B_2|}{2} \left| \frac{p_1}{q_1} - \frac{p_2}{q_2} \right|^2 \\
 &\leq \kappa c + H(P_2) \left| \frac{p_1}{q_1} - \frac{p_2}{q_2} \right| + \frac{H(P_2)^2}{18\kappa c} \left| \frac{p_1}{q_1} - \frac{p_2}{q_2} \right|^2 \\
 &\stackrel{(4.15)}{\leq} \kappa c + H_{n+1} \cdot 2lR^{-n+k^+} + \frac{H_{n+1}^2}{18\kappa c} \cdot 4l^2 R^{-2n+2k^+} \\
 &= \kappa c + 84\kappa^3 c R^{k^++1} + 392\kappa^5 c R^{2k^++2} \\
 &\leq 477\kappa^5 c R^{2k^++2}.
 \end{aligned}$$

The above argument can be trivially modified to show that the same upper bound is valid for $q_1 |F_1(\frac{p_2}{q_2})|$. Turning our attention to (4.17), using (2.7) we estimate as follows

$$\begin{aligned}
 |A_2 p_1 + B_2 r_1 + C_2 q_1| &= q_1 \left| F_2\left(\frac{p_1}{q_1}\right) - B_2 \left(f\left(\frac{p_1}{q_1}\right) - \frac{r_1}{q_1} \right) \right| \\
 &\leq q_1 \left| F_2\left(\frac{p_1}{q_1}\right) \right| + q_1 |B_2| \left| f\left(\frac{p_1}{q_1}\right) - \frac{r_1}{q_1} \right| \\
 &\stackrel{(4.16)}{\leq} 477\kappa^5 c \frac{q_1}{q_2} R^{2k^++2} + |B_2| \frac{\kappa c}{q_1^j} \\
 &\leq 477\kappa^5 c \left(\frac{q_1}{q_2} R^{2k^++2} + \frac{|B_2|}{q_1^j} \right) \\
 &\stackrel{(2.6)}{\leq} \frac{1}{2} R^{-4-\lambda_1} \left(\frac{q_1}{q_2} R^{2k^++2} + \frac{|B_2|}{q_1^j} \right).
 \end{aligned}$$

The same argument with obvious modifications yields (4.18). Finally, regarding (4.19), using the definition (3.4) for E_1 and E_2 , we have that

$$\begin{aligned}
 |A_1 B_2 - A_2 B_1| &= \left| \left(E_1 - f'\left(\frac{p_1}{q_1}\right) B_1 \right) B_2 - \left(E_2 - f'\left(\frac{p_2}{q_2}\right) B_2 \right) B_1 \right| \\
 &= \left| E_1 B_2 - E_2 B_1 + B_1 B_2 \left(f'\left(\frac{p_2}{q_2}\right) - f'\left(\frac{p_1}{q_1}\right) \right) \right| \\
 &\leq |E_1 B_2| + |E_2 B_1| + |B_1 B_2| \kappa \left| \frac{p_1}{q_1} - \frac{p_2}{q_2} \right| \\
 &\stackrel{(4.15)}{\leq} |E_1 B_2| + |E_2 B_1| + 2|B_1 B_2| \kappa l R^{-n+k^+}.
 \end{aligned}$$

Having established (4.15)–(4.19), we are now in the position to prove the lemma.

Part (1). Suppose $k = 0$ and $q_1 \leq q_2$. It follows from the definition (4.3) for $\mathcal{P}_{n,0}$ that

$$\begin{aligned}
 |A_2 p_1 + B_2 r_1 + C_2 q_1| &\stackrel{(4.17)}{\leq} \frac{1}{2} R^{-4-\lambda_1} \left(\frac{q_1}{q_2} R^4 + \frac{|B_2|}{q_1^j} \right) \\
 &\leq \frac{1}{2} R^{-4-\lambda_1} \left(R^4 + \frac{q_2 |B_2|}{q_1 |B_1|} \right) \\
 &= \frac{1}{2} R^{-4-\lambda_1} \left(R^4 + \frac{H(P_2)^2}{H(P_1)^2} \right) \\
 &\leq \frac{1}{2} R^{-4-\lambda_1} (R^4 + R^2) \\
 &\leq R^{-\lambda_1} < 1.
 \end{aligned}$$

Since the left hand side of the above inequality is an integer, it follows that $A_2 p_1 + B_2 r_1 + C_2 q_1 = 0$. Regarding (4.10), note that for $s = 1, 2$

$$|E_s| \leq \frac{H(P_s)^2}{q_s H(P_s)} \leq \frac{9\kappa^2 c |B_s|}{H_n} \leq \frac{l R^{-n}}{2\kappa} |B_s|$$

and that

$$\frac{|B_2|}{|B_1|} \leq \frac{q_2 |B_2|}{q_1 |B_1|} = \frac{H(P_2)^2}{H(P_1)^2} \leq \frac{H_{n+1}^2}{H_n^2} = R^2.$$

These inequalities together with (2.3) imply that

$$\begin{aligned}
 |A_1 B_2 - A_2 B_1| &\stackrel{(4.19)}{\leq} |E_1 B_2| + |E_2 B_1| + 2|B_1 B_2| \kappa l R^{-n+1} \\
 &\leq |B_1 B_2| \left(\frac{l R^{-n}}{\kappa} + 2\kappa l R^{-n+1} \right) \\
 &\leq B_1^2 R^2 \cdot 3\kappa l \\
 &\leq 3\kappa l R^2 q_1^{2j} < q_1.
 \end{aligned}$$

Part (2). Suppose $k = 1$. It follows from (4.4) that

$$\max \left\{ \frac{q_1}{q_2}, \frac{q_2}{q_1} \right\} \leq R^{\frac{\lambda_1}{1+i}} < R^{\lambda_1}.$$

This implies that

$$\begin{aligned} |A_2 p_1 + B_2 r_1 + C_2 q_1| &\stackrel{(4.17)}{\leq} \frac{1}{2} R^{-4-\lambda_1} \left(\frac{q_1}{q_2} R^4 + \frac{|B_2|}{q_1^j} \right) \\ &\leq \frac{1}{2} R^{-\lambda_1} \left(\frac{q_1}{q_2} + \frac{q_2^j}{q_1^j} \right) < 1. \end{aligned}$$

The left hand side is an integer and so must be zero. The same argument involving (4.18) rather than (4.17) shows that $|A_1 p_2 + B_1 r_2 + C_1 q_2| = 0$.

Part (3). Suppose $k \geq 2$. We first prove that

$$\max \left\{ \frac{q_1^i}{q_2}, \frac{q_2^j}{q_1} \right\} \leq \frac{1}{10^3 \kappa^5} H_n^{-\frac{j}{1+i}} R^{-2k-2} \quad (4.20)$$

$$\max \left\{ \frac{q_1^j}{q_2}, \frac{q_2^j}{q_1} \right\} \leq \frac{1}{10^3 \kappa^5} H_n^{-\frac{i}{1+i}} R^{-2k-2}. \quad (4.21)$$

It follows from (4.4) that

$$\max\{q_1, q_2\} \leq (\kappa^{-1} H_n)^{\frac{1}{1+i}} R^{\frac{\lambda_k}{1+i}}$$

and

$$\min\{q_1, q_2\} \geq (\kappa^{-1} H_n)^{\frac{1}{1+i}} R^{\frac{\lambda_k-1}{1+i}}.$$

In view of the fact that

$$\frac{j\lambda_k - \lambda_{k-1}}{1+i} \leq \frac{i\lambda_k - \lambda_{k-1}}{1+i} = -\frac{j}{1+i}\mu + \frac{2}{j} - 2k,$$

it follows from (2.4) that

$$\begin{aligned} \max \left\{ \frac{q_1^i}{q_2}, \frac{q_2^j}{q_1} \right\} &\leq (\kappa^{-1} H_n)^{-\frac{j}{1+i}} R^{\frac{i\lambda_k - \lambda_{k-1}}{1+i}} \\ &\leq \kappa H_n^{-\frac{j}{1+i}} R^{-\frac{j}{1+i}\mu + \frac{2}{j} - 2k} \\ &\leq \frac{1}{10^3 \kappa^5} H_n^{-\frac{j}{1+i}} R^{-2k-2} \end{aligned}$$

and that

$$\begin{aligned} \max \left\{ \frac{q_1^j}{q_2}, \frac{q_2^j}{q_1} \right\} &\leq (\kappa^{-1} H_n)^{-\frac{i}{1+i}} R^{\frac{j\lambda_k - \lambda_{k-1}}{1+i}} \\ &\leq \kappa H_n^{-\frac{i}{1+i}} R^{-\frac{j}{1+i}\mu + \frac{2}{j} - 2k} \\ &\leq \frac{1}{10^3 \kappa^5} H_n^{-\frac{i}{1+i}} R^{-2k-2}. \end{aligned}$$

This establishes (4.20) and (4.21). It now follows that

$$\begin{aligned}
 \left| A_1 F_2 \left(\frac{p_1}{q_1} \right) - A_2 F_1 \left(\frac{p_1}{q_1} \right) \right| &\leq |A_1| \left| F_2 \left(\frac{p_1}{q_1} \right) \right| + |A_2| \left| F_1 \left(\frac{p_1}{q_1} \right) \right| \\
 &\stackrel{(4.16)}{\leq} q_1^i \cdot \frac{1}{q_2} 477 \kappa^5 c R^{2k+2} + q_2^i \frac{\kappa c}{q_1} \\
 &\stackrel{(4.20)}{\leq} \frac{1}{10^3 \kappa^5} H_n^{-\frac{j}{1+i}} R^{-2k-2} (477 \kappa^5 c R^{2k+2} + \kappa c) \\
 &\leq \frac{c}{2} H_n^{-\frac{j}{1+i}}
 \end{aligned}$$

and

$$\begin{aligned}
 \left| B_1 F_2 \left(\frac{p_1}{q_1} \right) - B_2 F_1 \left(\frac{p_1}{q_1} \right) \right| &\leq |B_1| \left| F_2 \left(\frac{p_1}{q_1} \right) \right| + |B_2| \left| F_1 \left(\frac{p_1}{q_1} \right) \right| \\
 &\stackrel{(4.16)}{\leq} q_1^j \cdot \frac{1}{q_2} 477 \kappa^5 c R^{2k+2} + q_2^j \frac{\kappa c}{q_1} \\
 &\stackrel{(4.21)}{\leq} \frac{1}{10^3 \kappa^5} H_n^{-\frac{i}{1+i}} R^{-2k-2} (477 \kappa^5 c R^{2k+2} + \kappa c) \\
 &\leq \frac{c}{2} H_n^{-\frac{i}{1+i}}.
 \end{aligned}$$

Finally, since $P_s \notin \mathcal{P}^*$, we have that $q_s E_s^2 \geq 9\kappa^2 c |B_s|$ and $H(P_s) = q_s |E_s|$ for $s = 1, 2$. Then

$$|B_2| \leq \frac{H(P_2)^2}{9\kappa^2 c q_2}$$

and using (3.2) and the fact that $H_{n+1} = R H_n$, we get that

$$\begin{aligned}
 |A_1 B_2 - A_2 B_1| &\stackrel{(4.19)}{\leq} |E_1 B_2| + |E_2 B_1| + 2|B_1 B_2| \kappa l R^{-n+k} \\
 &\leq \frac{H(P_1)}{q_1} q_2^j + \frac{H(P_2)}{q_2} q_1^j + 2q_1^j \frac{H(P_2)^2}{9\kappa^2 c q_2} \kappa l R^{-n+k} \\
 &\stackrel{(4.21)}{\leq} \frac{1}{10^3 \kappa^5} H_n^{-\frac{i}{1+i}} R^{-2k-2} \left(2H_{n+1} + \frac{2H_{n+1}^2}{9\kappa^2 c} \kappa l R^{-n+k} \right) \\
 &\leq \frac{1}{500 \kappa^5} H_n^{\frac{1}{1+i}} R^{-2k-2} (R + 5\kappa^2 R^{k+2}) \\
 &\leq \frac{1}{3\kappa} H_n^{\frac{1}{1+i}} R^{-k}.
 \end{aligned}$$

This thereby completes the proof of Part (3) and thus the lemma. \square

We now proceed with the proof of [Lemma 4.2](#). Let $P_1 = (\frac{p_1}{q_1}, \frac{r_1}{q_1})$ and $P_2 = (\frac{p_2}{q_2}, \frac{r_2}{q_2})$ be distinct points in $\mathcal{P}_{n,k}(\tau)$. We need to prove that $L_{P_1} = L_{P_2}$. We consider three separate cases.

Case (1). Suppose $k = 0$. Without loss of generality, we assume that $q_1 \leq q_2$. Then in view of [\(4.9\)](#) we have that $A_2p_1 + B_2r_1 + C_2q_1 = 0$. Hence L_{P_2} passes through P_1 . We prove that $L_{P_1} = L_{P_2}$ by contradiction. Thus, assume that $L_{P_1} \neq L_{P_2}$. Since the two lines intersect at P_1 , it follows that

$$\frac{p_1}{q_1} = \frac{B_1C_2 - B_2C_1}{A_1B_2 - A_2B_1} \quad \text{and} \quad \frac{r_1}{q_1} = \frac{A_2C_1 - A_1C_2}{A_1B_2 - A_2B_1}.$$

In particular, since p_1, q_1, r_1 are co-prime, the non-zero integer $A_1B_2 - A_2B_1$ is divisible by q_1 . Thus

$$q_1 \leq |A_1B_2 - A_2B_1| \stackrel{(4.10)}{<} q_1,$$

which is of course impossible.

Case (2). Suppose $k = 1$. Then in view of [\(4.11\)](#) we have that $A_2p_1 + B_2r_1 + C_2q_1 = 0 = A_1p_2 + B_1r_2 + C_1q_2$. Hence, L_{P_2} passes through P_1 and L_{P_1} passes through P_2 . By definition, L_{P_2} passes through P_2 and L_{P_1} passes through P_1 . The upshot is that both lines pass through the points P_1 and P_2 , and so we must have that $L_{P_1} = L_{P_2}$.

Case (3). Suppose $k \geq 2$. We prove that $L_{P_1} = L_{P_2}$ by contradiction. Thus, assume that $L_{P_1} \neq L_{P_2}$. We first consider the case where L_{P_1} is parallel to L_{P_2} . Then, it is easily verified that

$$A_1F_2\left(\frac{p_1}{q_1}\right) - A_2F_1\left(\frac{p_1}{q_1}\right) = A_1C_2 - A_2C_1$$

and

$$B_1F_2\left(\frac{p_1}{q_1}\right) - B_2F_1\left(\frac{p_1}{q_1}\right) = B_1C_2 - B_2C_1.$$

Since $H_n \geq c$, it follows via [\(4.12\)](#) and [\(4.13\)](#) that

$$\begin{aligned} 1 &\leq |A_1C_2 - A_2C_1| + |B_1C_2 - B_2C_1| \\ &\leq \left| A_1F_2\left(\frac{p_1}{q_1}\right) - A_2F_1\left(\frac{p_1}{q_1}\right) \right| + \left| B_1F_2\left(\frac{p_1}{q_1}\right) - B_2F_1\left(\frac{p_1}{q_1}\right) \right| \\ &\leq \frac{c}{2}H_n^{-\frac{j}{1+i}} + \frac{c}{2}H_n^{-\frac{i}{1+i}} \\ &\leq \frac{c}{2}(c^{-\frac{j}{1+i}} + c^{-\frac{i}{1+i}}) \\ &= \frac{1}{2}(c^{\frac{2i}{1+i}} + c^{\frac{1}{1+i}}) < 1 \end{aligned}$$

which is of course impossible.

Now suppose L_{P_1} is not parallel to L_{P_2} . Let $P_0 = (\frac{p_0}{q_0}, \frac{r_0}{q_0}) \in \mathbb{Q}^2$ be the point of intersection of L_{P_1} and L_{P_2} . Then it follows that the non-zero integer $A_1B_2 - A_2B_1$ is divisible by q_0 and so

$$q_0 \leq |A_1B_2 - A_2B_1|. \quad (4.22)$$

We first prove that $\Delta(P_1) \subset \Delta(P_0)$ and that $P_0 \in \mathcal{P}$. It is easily verified that

$$\begin{pmatrix} A_1 & B_1 \\ A_2 & B_2 \end{pmatrix} \begin{pmatrix} \frac{p_1}{q_1} - \frac{p_0}{q_0} \\ f(\frac{p_1}{q_1}) - \frac{r_0}{q_0} \end{pmatrix} = \begin{pmatrix} F_1(\frac{p_1}{q_1}) \\ F_2(\frac{p_1}{q_1}) \end{pmatrix}.$$

Hence, on using Cramer's rule, we obtain that

$$|A_1B_2 - A_2B_1| \left| \frac{p_1}{q_1} - \frac{p_0}{q_0} \right| = \left| B_1F_2\left(\frac{p_1}{q_1}\right) - B_2F_1\left(\frac{p_1}{q_1}\right) \right| \stackrel{(4.13)}{\leq} \frac{c}{2} H_n^{-\frac{i}{1+i}}, \quad (4.23)$$

and

$$|A_1B_2 - A_2B_1| \left| f\left(\frac{p_1}{q_1}\right) - \frac{r_0}{q_0} \right| = \left| A_1F_2\left(\frac{p_1}{q_1}\right) - A_2F_1\left(\frac{p_1}{q_1}\right) \right| \stackrel{(4.12)}{\leq} \frac{c}{2} H_n^{-\frac{j}{1+i}}. \quad (4.24)$$

If $x \in \Delta(P_1)$, then (4.14) and (4.23) imply that

$$\begin{aligned} q_0^{1+i} \left| x - \frac{p_0}{q_0} \right| &\stackrel{(4.22)}{\leq} |A_1B_2 - A_2B_1|^{1+i} \left| x - \frac{p_1}{q_1} \right| + |A_1B_2 - A_2B_1|^{1+i} \left| \frac{p_1}{q_1} - \frac{p_0}{q_0} \right| \\ &\leq \frac{H_n}{3\kappa} \cdot \frac{c}{q_1^{1+i}} + H_n^{\frac{i}{1+i}} \cdot \frac{c}{2} H_n^{-\frac{i}{1+i}} \\ &\stackrel{(4.2)}{\leq} \frac{c}{3} + \frac{c}{2} < c. \end{aligned}$$

Thus $x \in \Delta(P_0)$ and the upshot is that $\Delta(P_1) \subset \Delta(P_0)$. In particular,

$$\left| \frac{p_1}{q_1} - \frac{p_0}{q_0} \right| < \frac{c}{q_0^{1+i}}. \quad (4.25)$$

Since $\mathbf{A}_0 \cap \Delta(P_1) \neq \emptyset$, there exists $x \in \mathbf{A}_0$ such that $|x - \frac{p_1}{q_1}| < c/q_1^{1+i}$, and hence it follows that

$$\left| x - \frac{p_0}{q_0} \right| \leq \left| x - \frac{p_1}{q_1} \right| + \left| \frac{p_1}{q_1} - \frac{p_0}{q_0} \right| \leq 2c \leq \frac{l}{2}.$$

This implies that $\frac{p_0}{q_0} \in \mathbf{B}_0 \subset I$. Also note that by (4.14) and (4.24), we have that

$$\begin{aligned}
q_0^{1+j} \left| f\left(\frac{p_0}{q_0}\right) - \frac{r_0}{q_0} \right| &\stackrel{(4.22)}{\leq} q_0^{1+j} \left| f\left(\frac{p_0}{q_0}\right) - f\left(\frac{p_1}{q_1}\right) \right| + |A_1 B_2 - A_2 B_1|^{1+j} \left| f\left(\frac{p_1}{q_1}\right) - \frac{r_0}{q_0} \right| \\
&\leq q_0^{1+i} (\kappa - 1) \left| \frac{p_0}{q_0} - \frac{p_1}{q_1} \right| + H_n^{\frac{j}{1+i}} \cdot \frac{c}{2} H_n^{-\frac{j}{1+i}} \\
&\stackrel{(4.25)}{\leq} (\kappa - 1)c + \frac{c}{2} \\
&< \kappa c.
\end{aligned}$$

Thus $P_0 \in \mathcal{P}$ and so there exists a unique integer $n_0 \geq 1$ such that $P_0 \in \mathcal{P}_{n_0}$. Suppose for the moment that $n_0 \leq n - k$. Then there exists $\tau' \in \mathcal{S}_{n_0}$ such that $\tau \prec \tau'$, and hence

$$\mathcal{I}(\tau) \cap \Delta(P_1) \subset \mathcal{I}(\tau') \cap \Delta(P_0) = \emptyset.$$

This contradicts the fact that $P_1 \in \mathcal{P}_{n,k}(\tau)$. Thus

$$n_0 \geq n - k + 1,$$

and so

$$H(P_0) \geq H_{n_0} \geq H_{n-k+1}.$$

On the other hand, we have that

$$H(P_0) \leq \kappa q_0^{1+i} \stackrel{(4.22)}{\leq} \kappa |A_1 B_2 - A_2 B_1|^{1+i} \stackrel{(4.14)}{\leq} H_n R^{-k} = H_{n-k}.$$

This contradicts the above lower bound for $H(P_0)$ and so completes the proof of Case (3) and indeed the lemma. \square

5. The inhomogeneous case: establishing [Theorem 1.1](#)

[Theorem 1.1](#) is easily deduced from the following statement.

Theorem 5.1. *Let (i, j) be a pair of real numbers satisfying $0 < j \leq i < 1$ and $i + j = 1$. Let $I \subset \mathbb{R}$ be a compact interval and $f \in C^{(2)}(I)$ such that $f''(x) \neq 0$ for all $x \in I$. Then, for any $\theta = (\gamma, \delta) \in \mathbb{R}^2$ we have that $\mathbf{Bad}_\theta^f(i, j)$ is a $1/2$ -winning subset of I .*

We have already established the homogeneous case ($\gamma = \delta = 0$) of [Theorem 5.1](#); namely [Theorem 2.1](#). With reference to §2.2, the crux of the ‘homogeneous’ proof involved constructing a partition \mathcal{P}_n ($n \geq 1$) of \mathcal{P} (given by [Lemma 4.1](#)) such that the subtree \mathcal{S} of an $[R]$ -regular rooted tree \mathcal{T} has an $([R] - 10)$ -regular subtree \mathcal{S}' – the substance of [Proposition 2.4](#). To prove [Theorem 5.1](#), the idea is to merge the inhomogeneous constraints into the homogeneous construction. More precisely, we show that \mathcal{S}'

has an $([R] - 12)$ -regular subtree \mathcal{Q}' that incorporates the inhomogeneous constraints. With this in mind, let

$$c' := \frac{1}{10}cR^{-2}$$

where c is defined in (2.6), and let

$$\mathcal{V} := \left\{ (p, r, q) \in \mathbb{Z}^2 \times \mathbb{N} : \frac{p+\gamma}{q} \in I, \left| f\left(\frac{p+\gamma}{q}\right) - \frac{r+\delta}{q} \right| < \frac{\kappa c'}{q^{1+j}} \right\}.$$

Furthermore, for each $v = (p, r, q) \in \mathbb{Z}^2 \times \mathbb{N}$, we associate the interval

$$\Delta_{\theta}(v) := \left\{ x \in I : \left| x - \frac{p+\gamma}{q} \right| < \frac{c'}{q^{1+i}} \right\}.$$

Then, with $\mathbf{A}_0 \subset \mathbf{B}_0$ as in §2.2, the following is the inhomogeneous analogue of Lemma 2.3.

Lemma 5.2. *Let \mathbf{A}_0 , \mathcal{V} and $\Delta_{\theta}(v)$ be as above. Then*

$$\mathbf{A}_0 \setminus \bigcup_{v \in \mathcal{V}} \Delta_{\theta}(v) \subset \mathbf{Bad}_{\theta}^f(i, j).$$

Proof. The proof is similar to the homogeneous proof but is included for completeness. Let $x \in \mathbf{A}_0$. Suppose $x \notin \mathbf{Bad}_{\theta}^f(i, j)$. Then there exists $v = (p, r, q) \in \mathbb{Z}^2 \times \mathbb{N}$ such that

$$\left| x - \frac{p+\gamma}{q} \right| < \frac{c'}{q^{1+i}}, \quad \left| f(x) - \frac{r+\delta}{q} \right| < \frac{c'}{q^{1+j}}.$$

In view of the fact that

$$\left| x - \frac{p+\gamma}{q} \right| < \frac{c'}{q^{1+i}} \leq c \leq \frac{l}{2},$$

it follows that $\frac{p+\gamma}{q} \in \mathbf{B}_0 \subset I$. Hence

$$\begin{aligned} \left| f\left(\frac{p+\gamma}{q}\right) - \frac{r+\delta}{q} \right| &\leq \left| f\left(\frac{p+\gamma}{q}\right) - f(x) \right| + \left| f(x) - \frac{r+\delta}{q} \right| \\ &\leq (\kappa - 1) \left| x - \frac{p+\gamma}{q} \right| + \frac{c'}{q^{1+j}} \\ &< \frac{(\kappa - 1)c'}{q^{1+i}} + \frac{c'}{q^{1+j}} \leq \frac{\kappa c'}{q^{1+j}}. \end{aligned}$$

Thus $x \in \Delta_{\theta}(v)$ and $v \in \mathcal{V}$. This completes the proof of the lemma. \square

In view of [Lemma 5.2](#), to prove [Theorem 5.1](#), it suffices to show that Ayesha can play the $(\frac{1}{2}, \beta)$ -game such that $\bigcap_{n=0}^{\infty} \mathbf{A}_n \subset \mathbf{A}_0 \setminus \bigcup_{v \in \mathcal{V}} \Delta_{\theta}(v)$. We do this by proving the stronger statement that Ayesha can play such that

$$\bigcap_{n=0}^{\infty} \mathbf{A}_n \subset \mathbf{A}_0 \setminus \left(\bigcup_{P \in \mathcal{P}} \Delta(P) \cup \bigcup_{v \in \mathcal{V}} \Delta_{\theta}(v) \right), \quad (5.1)$$

where \mathcal{P} and $\Delta(P)$ are given in [\(2.7\)](#) and [\(2.8\)](#), respectively. Recall that for the partition $\mathcal{P} = \bigcup_{n=1}^{\infty} \mathcal{P}_n$ given by [Lemma 4.1](#), the $[R]$ -regular rooted tree \mathcal{T} has an $([R] - 10)$ -regular subtree \mathcal{S}' such that

$$\mathcal{I}(\tau) \cap \bigcup_{P \in \mathcal{P}_n} \Delta(P) = \emptyset, \quad \forall n \geq 1 \quad \text{and} \quad \tau \in \mathcal{S}'_n. \quad (5.2)$$

In what follows, we work with the tree \mathcal{S}' and take the inhomogeneous constraints $\Delta_{\theta}(v)$ into account.

For $n \geq 1$, let

$$H'_n := 2c'l^{-1}R^n$$

and

$$\mathcal{V}_n := \{(p, r, q) \in \mathcal{V} : H'_n \leq q^{1+i} < H'_{n+1}\}.$$

Observe that $H'_1 = 2c'l^{-1}R \leq 1$ and so it follows that $\mathcal{V} = \bigcup_{n=1}^{\infty} \mathcal{V}_n$. We inductively define a subtree \mathcal{Q} of \mathcal{S}' as follows. Let $\mathcal{Q}_0 = \{\tau_0\}$. If \mathcal{Q}_{n-1} ($n \geq 1$) is defined, we let

$$\mathcal{Q}_n := \left\{ \tau \in \mathcal{S}'_{\text{suc}}(\mathcal{Q}_{n-1}) : \mathcal{I}(\tau) \cap \bigcup_{v \in \mathcal{V}_n} \Delta_{\theta}(v) = \emptyset \right\}.$$

Then

$$\mathcal{Q} := \bigcup_{n=0}^{\infty} \mathcal{Q}_n$$

is a subtree of \mathcal{S}' and, by construction, we have that

$$\mathcal{I}(\tau) \subset \mathbf{A}_0 \setminus \left(\bigcup_{P \in \mathcal{P}_n} \Delta(P) \cup \bigcup_{v \in \mathcal{V}_n} \Delta_{\theta}(v) \right) \quad \forall n \geq 1 \quad \text{and} \quad \tau \in \mathcal{Q}_n.$$

Armed with the following result, the same arguments as in [§2.2.1](#) with the most obvious modifications enables us to prove [\(5.1\)](#), and hence [Theorem 5.1](#). In view of this the details of the proof of [Theorem 5.1](#) modulo [Proposition 5.3](#) are omitted.

Proposition 5.3. *The tree \mathcal{Q} has an $([R] - 12)$ -regular subtree.*

Since \mathcal{S}' is $([R] - 10)$ -regular, in order to establish the proposition, it suffices to prove the following statement.

Lemma 5.4. *For any $n \geq 1$ and $\tau \in \mathcal{Q}_{n-1}$, there is at most one $v \in \mathcal{V}_n$ such that $\mathcal{I}(\tau) \cap \Delta_{\theta}(v) \neq \emptyset$. Moreover, $\rho(\Delta_{\theta}(v)) \leq lR^{-n}$. Therefore,*

$$\#\left\{\tau' \in \mathcal{S}'_{\text{suc}}(\tau) : \mathcal{I}(\tau') \cap \bigcup_{v \in \mathcal{V}_n} \Delta_{\theta}(v) \neq \emptyset\right\} \leq 2.$$

Proof. Suppose $v_s = (p_s, r_s, q_s) \in \mathcal{V}_n$ and $\mathcal{I}(\tau) \cap \Delta_{\theta}(v_s) \neq \emptyset$, $s = 1, 2$. We need to prove that $v_1 = v_2$. Without loss of generality, assume that $q_1 \geq q_2$. Let $x_s \in \mathcal{I}(\tau) \cap \Delta_{\theta}(v_s)$. Then

$$\left|x_s - \frac{p_s + \gamma}{q_s}\right| < \frac{c'}{q_s^{1+i}}, \quad s = 1, 2$$

and

$$|x_1 - x_2| \leq \rho(\mathcal{I}(\tau)) = lR^{-n+1}.$$

It follows that

$$\begin{aligned} |(q_1 - q_2)x_1 - (p_1 - p_2)| &= \left|q_1\left(x_1 - \frac{p_1 + \gamma}{q_1}\right) - q_2\left(x_2 - \frac{p_2 + \gamma}{q_2}\right) - q_2(x_1 - x_2)\right| \\ &\leq q_1\left|x_1 - \frac{p_1 + \gamma}{q_1}\right| + q_2\left|x_2 - \frac{p_2 + \gamma}{q_2}\right| + q_2|x_1 - x_2| \\ &\leq \frac{c'}{q_1^i} + \frac{c'}{q_2^i} + q_2lR^{-n+1} \\ &\leq \frac{2c'}{q_2^i} + q_2lR^{-n+1}. \end{aligned} \tag{5.3}$$

Moreover,

$$\begin{aligned} &\left|(q_1 - q_2)f\left(\frac{p_1 + \gamma}{q_1}\right) - (r_1 - r_2)\right| \\ &= \left|q_1\left(f\left(\frac{p_1 + \gamma}{q_1}\right) - \frac{r_1 + \delta}{q_1}\right) - q_2\left(f\left(\frac{p_2 + \gamma}{q_2}\right) - \frac{r_2 + \delta}{q_2}\right) \right. \\ &\quad \left. - q_2\left(f\left(\frac{p_1 + \gamma}{q_1}\right) - f\left(\frac{p_2 + \gamma}{q_2}\right)\right)\right| \\ &\leq q_1\left|f\left(\frac{p_1 + \gamma}{q_1}\right) - \frac{r_1 + \delta}{q_1}\right| + q_2\left|f\left(\frac{p_2 + \gamma}{q_2}\right) - \frac{r_2 + \delta}{q_2}\right| \\ &\quad + q_2\left|f\left(\frac{p_1 + \gamma}{q_1}\right) - f\left(\frac{p_2 + \gamma}{q_2}\right)\right| \end{aligned}$$

$$\begin{aligned}
&\leq \frac{\kappa c'}{q_1^j} + \frac{\kappa c'}{q_2^j} + q_2 \kappa \left| \frac{p_1 + \gamma}{q_1} - \frac{p_2 + \gamma}{q_2} \right| \\
&\leq \frac{\kappa c'}{q_1^j} + \frac{\kappa c'}{q_2^j} + q_2 \kappa \left(\frac{c'}{q_1^{1+i}} + \frac{c'}{q_2^{1+i}} + l R^{-n+1} \right) \\
&\leq \frac{4\kappa c'}{q_2^j} + q_2 \kappa l R^{-n+1}.
\end{aligned} \tag{5.4}$$

Suppose for the moment that $q_1 > q_2$ and let

$$P_0 := \left(\frac{p_1 - p_2}{q_1 - q_2}, \frac{r_1 - r_2}{q_1 - q_2} \right).$$

We show that

$$\mathcal{I}(\tau) \cap \Delta(P_0) \neq \emptyset \tag{5.5}$$

and that $P_0 \in \mathcal{P}$, where \mathcal{P} and $\Delta(P_0)$ are defined by (2.7) and (2.8), respectively. In view of (5.3), it follows that

$$\begin{aligned}
(q_1 - q_2)^{1+i} \left| x_1 - \frac{p_1 - p_2}{q_1 - q_2} \right| &\leq (q_1 - q_2)^i \left(\frac{2c'}{q_2^i} + q_2 l R^{-n+1} \right) \\
&\leq 2c' \frac{q_1^i}{q_2^i} + q_1^{1+i} l R^{-n+1} \leq 2c' R + 2c' R^2 < c.
\end{aligned}$$

So $x_1 \in \Delta(P_0)$ and (5.5) follows. Also, the above inequality implies that

$$\left| x_1 - \frac{p_1 - p_2}{q_1 - q_2} \right| < c \leq \frac{l}{2}$$

and since $x_1 \in \mathbf{A}_0$, it follows that $\frac{p_1 - p_2}{q_1 - q_2} \in \mathbf{B}_0 \subset I$. Moreover, by making use of (5.4) we have that

$$\begin{aligned}
(q_1 - q_2)^{1+j} \left| f\left(\frac{p_1 - p_2}{q_1 - q_2}\right) - \frac{r_1 - r_2}{q_1 - q_2} \right| &\leq (q_1 - q_2)^{1+j} \left| f\left(\frac{p_1 - p_2}{q_1 - q_2}\right) - f\left(\frac{p_1 + \gamma}{q_1}\right) \right| \\
&\quad + (q_1 - q_2)^j \left| (q_1 - q_2) f\left(\frac{p_1 + \gamma}{q_1}\right) - (r_1 - r_2) \right| \\
&\leq (q_1 - q_2)^{1+j} \kappa \left| \frac{p_1 - p_2}{q_1 - q_2} - \frac{p_1 + \gamma}{q_1} \right| \\
&\quad + (q_1 - q_2)^j \left(\frac{4\kappa c'}{q_2^j} + q_2 \kappa l R^{-n+1} \right) \\
&= (q_1 - q_2)^j \left(q_2 \kappa \left| \frac{p_1 + \gamma}{q_1} - \frac{p_2 + \gamma}{q_2} \right| + \frac{4\kappa c'}{q_2^j} + q_2 \kappa l R^{-n+1} \right)
\end{aligned}$$

$$\begin{aligned}
&\leq q_1^j \left(q_2 \kappa \left(\frac{c'}{q_1^{1+i}} + lR^{-n+1} + \frac{c'}{q_2^{1+i}} \right) + \frac{4\kappa c'}{q_2^j} + q_2 \kappa l R^{-n+1} \right) \\
&\leq 6\kappa c' \frac{q_1^j}{q_2^j} + 2q_1^{1+j} \kappa l R^{-n+1} \\
&\leq 6\kappa c' R + 4\kappa c' R^2 < \kappa c.
\end{aligned}$$

Thus $P_0 \in \mathcal{P}$ and so there exists a unique integer $n_0 \geq 1$ such that $P_0 \in \mathcal{P}_{n_0}$. Suppose for the moment that $n_0 \leq n-1$. Then there exists $\tau' \in \mathcal{S}'_{n_0}$ such that $\tau \prec \tau'$, and it follows from (5.2) that $\mathcal{I}(\tau) \cap \Delta(P_0) \subset \mathcal{I}(\tau') \cap \Delta(P_0) = \emptyset$ contrary to (5.5). Thus, $n_0 \geq n$ and so

$$H(P_0) \geq H_{n_0} \geq H_n = 42\kappa^3 cl^{-1} R^n.$$

On the other hand, we have that

$$H(P_0) \leq \kappa(q_1 - q_2)^{1+i} \leq \kappa q_1^{1+i} \leq \kappa H'_{n+1} = \frac{1}{5} \kappa cl^{-1} R^{n-1}.$$

This contradicts the above lower bound for $H(P_0)$ and we conclude that $q_1 = q_2$. Since $q_2 \leq H'_{n+1} \leq cl^{-1} R^{n-1}$, it now follows from (5.3) and (5.4) that

$$|p_1 - p_2| \leq \frac{2c'}{q_2^i} + q_2 l R^{-n+1} \leq 2c' + c < 1$$

and

$$|r_1 - r_2| \leq \frac{4\kappa c'}{q_2^j} + q_2 \kappa l R^{-n+1} \leq 4\kappa c' + c\kappa < 1.$$

Thus, $p_1 = p_2$ and $r_1 = r_2$. In other words, $v_1 = v_2$ and this proves the main substance of the lemma. To prove the ‘moreover’ part, it is easily verified that for any $v \in \mathcal{V}_n$ we have that

$$\rho(\Delta_{\theta}(v)) = \frac{2c'}{q^{1+i}} \leq \frac{2c'}{H'_n} = lR^{-n}.$$

The ‘therefore’ part of the lemma is a direct consequence of this and the fact that there is at most one $v \in \mathcal{V}_n$ such that $\mathcal{I}(\tau) \cap \Delta_{\theta}(v)$ is non-empty. \square

As already mentioned, given Proposition 5.3, the proof of Theorem 5.1 follows on adapting the arguments of §2.2.1.

6. The proof of Theorem 1.2

The basic strategy towards establishing the winning result for lines is the same as when considering curves. To begin with observe that for any line $L_{a,b}$ given by

$$y = f(x) := ax + b$$

and $\theta \in \mathbb{R}^2$,

$$\mathbf{Bad}_{\theta}^f(i, j) := \{x \in \mathbb{R} : (x, f(x)) \in \mathbf{Bad}_{\theta}(i, j)\} = \pi(\mathbf{Bad}_{\theta}(i, j) \cap L_{a,b}).$$

As in the case of curves, without loss of generality we will assume that $j \leq i$. Thus, the homogeneous case of Theorem 1.2 is easily deduced from the following statement.

Theorem 6.1. *Let (i, j) be a pair of real numbers satisfying $0 < j \leq i < 1$ and $i + j = 1$. Given $a, b \in \mathbb{R}$, suppose there exists $\epsilon > 0$ such that*

$$\liminf_{q \rightarrow \infty} q^{\frac{1}{j} - \epsilon} \max\{\|qa\|, \|qb\|\} > 0. \quad (6.1)$$

Then $\mathbf{Bad}^f(i, j)$ is a $1/2$ -winning subset of \mathbb{R} . Moreover, if $a \in \mathbb{Q}$ the statement is true with $\epsilon = 0$ in (6.1).

Note that in view of Remark 5 after the statement of Theorem 1.2, we do not require that $a \neq 0$ in Theorem 6.1 since $j \leq i$.

6.1. The winning strategy for Theorem 6.1

Let $\beta \in (0, 1)$. We want to prove that $\mathbf{Bad}^f(i, j)$ is $(\frac{1}{2}, \beta)$ -winning. In the first round of the game, Bhupen chooses a closed interval $\mathbf{B}_0 \subset \mathbb{R}$. Now Ayesha chooses any closed interval $\mathbf{A}_0 \subset \mathbf{B}_0$ with diameter $\rho(\mathbf{A}_0) = \frac{1}{2}\rho(\mathbf{B}_0)$. Let

$$R := (2\beta^{-1})^4, \quad l := \rho(\mathbf{A}_0) \quad \text{and} \quad c_1 := c_0 \max\{|x|_{\max} + l, 1\}^{-1}$$

where

$$|x|_{\max} := \max\{|x| : x \in \mathbf{A}_0\} \quad \text{and} \quad c_0 := \inf_{q \in \mathbb{N}} q^{\frac{1}{j} - \epsilon} \max\{\|qa\|, \|qb\|\} > 0.$$

The fact that $c_0 > 0$ follows from the Diophantine condition (6.1). Recall, that by hypothesis $\epsilon = 0$ if a is rational and $\epsilon > 0$ otherwise. We denote

$$\kappa := |a| + 1,$$

and choose $\mu \geq 1$ such that

$$20\kappa^2 R^{\frac{3}{i}-i\mu} < 1. \quad (6.2)$$

If $a \in \mathbb{Q}$, we also require that

$$R^{\mu-1} \geq \kappa d^2 \quad (6.3)$$

where $d \in \mathbb{N}$ is the smallest positive integer such that $da \in \mathbb{Z}$. Next, if $a \notin \mathbb{Q}$, so that $\epsilon > 0$, we let $\lambda > 0$ be such that

$$R^{\lambda j^{-n}} \geq \kappa c_1^{-1} R^{\frac{1+i}{j\epsilon} n} \quad \text{for } n \geq 1. \quad (6.4)$$

If $a \in \mathbb{Q}$, we simply let $\lambda = 0$. In turn, let

$$\lambda_0 := 0 \quad \text{and} \quad \lambda_k := \lambda j^{-k} + \frac{k}{i} + \mu \quad \text{for } k \geq 1,$$

and let

$$c := \min \left\{ \frac{c_1}{4\kappa} l R^{-1}, l^{-i}, \frac{1}{8\kappa} R^{-2-\frac{\lambda_1}{1+i}} \right\}. \quad (6.5)$$

For each rational point $P = (\frac{p}{q}, \frac{r}{q}) \in \mathbb{R}^2$ we associate the interval

$$\Delta(P) := \left\{ x \in \mathbb{R} : \left| x - \frac{p}{q} \right| < \frac{c}{q^{1+i}} \right\}$$

and we let

$$\mathcal{P} := \left\{ P = \left(\frac{p}{q}, \frac{r}{q} \right) : \mathbf{A}_0 \cap \Delta(P) \neq \emptyset, \left| b + \frac{ap-r}{q} \right| < \frac{\kappa c}{q^{1+j}} \right\}.$$

The following inclusion is a simple consequence of the manner in which the above quantities and objects have been defined.

Lemma 6.2. *Let \mathbf{A}_0 , \mathcal{P} and $\Delta(P)$ be as above. Then*

$$\mathbf{A}_0 \setminus \bigcup_{P \in \mathcal{P}} \Delta(P) \subset \mathbf{Bad}^f(i, j).$$

Proof. Let $x \in \mathbf{A}_0$. Suppose $x \notin \mathbf{Bad}^f(i, j)$. Then there exists $P = (\frac{p}{q}, \frac{r}{q}) \in \mathbb{Q}^2$ such that

$$\left| x - \frac{p}{q} \right| < \frac{c}{q^{1+i}}, \quad \left| f(x) - \frac{r}{q} \right| = \left| ax + b - \frac{r}{q} \right| < \frac{c}{q^{1+j}}.$$

It follows that

$$\begin{aligned}
\left| b + \frac{ap-r}{q} \right| &= \left| \left(ax + b - \frac{r}{q} \right) - a \left(x - \frac{p}{q} \right) \right| \\
&\leq \frac{c}{q^{1+j}} + |a| \frac{c}{q^{1+i}} \\
&\leq \frac{\kappa c}{q^{1+j}}.
\end{aligned}$$

The upshot is that $x \in \Delta(P)$ with $P \in \mathcal{P}$. This completes the proof of the lemma. \square

Next, just as in §2.2, but with \mathbf{A}_0 and \mathcal{P} as above, let

- \mathcal{T} be an $[R]$ -regular rooted tree with root τ_0 ,
- \mathcal{I} be an injective map from \mathcal{T} to the set of closed subintervals of \mathbf{A}_0 ,
- $\mathcal{S} = \bigcup_{n=0}^{\infty} \mathcal{S}_n$ be a subtree of \mathcal{T} associated with a partition \mathcal{P}_n of \mathcal{P} .

The following proposition is the lines analogue of Proposition 2.4. It enables us to deduce Theorem 6.1 (and thus the homogeneous case of Theorem 1.2) by adapting the arguments of §2.2.1 in the most obvious manner. In view of this the details of the proof of Theorem 6.1 modulo Proposition 6.3 are omitted.

Proposition 6.3. *There exists a partition $\mathcal{P} = \bigcup_{n=1}^{\infty} \mathcal{P}_n$ such that the tree \mathcal{S} has an $([R] - 5)$ -regular subtree.*

6.1.1. Proof of Proposition 6.3

As in the ‘curves’ proof, to each point $P = (\frac{p}{q}, \frac{r}{q}) \in \mathcal{P}$, we attach a rational line

$$L_P := \{(x, y) \in \mathbb{R}^2 : Ax + By + C = 0\}$$

passing through P where $A, B, C \in \mathbb{Z}$ are co-prime with $(A, B) \neq (0, 0)$ and such that

$$|A| \leq q^i \quad \text{and} \quad |B| \leq q^j.$$

Associated with each point $P \in \mathcal{P}$, we also consider the quantity

$$E := A + Ba.$$

Then

$$|E| \leq q^i + |a|q^j \leq \kappa q^i. \quad (6.6)$$

Note that if $x \in \Delta(P)$, then

$$|Ex + Bb + C| = \left| E \left(x - \frac{p}{q} \right) + B \left(b + \frac{ap-r}{q} \right) \right| < \frac{2\kappa c}{q}. \quad (6.7)$$

The following statement enables us to construct the desired partition in [Proposition 6.3](#).

Lemma 6.4. *For any $P = (\frac{p}{q}, \frac{r}{q}) \in \mathcal{P}$, we have*

$$q^{1-j\epsilon}|E| \geq c_1. \quad (6.8)$$

Proof. If $B = 0$, then $q^{1-j\epsilon}|E| = q^{1-j\epsilon}|A| \geq 1 \geq c_1$ and we are done. If $B \neq 0$, then it is easily verified that

$$q^{1-j\epsilon} \max\{|E|, |Bb + C|\} \geq |B|^{\frac{1}{j}-\epsilon} \max\{|Ba + A|, |Bb + C|\} \geq c_0.$$

If $|E|$ is the maximum in the above then again we are done. So suppose $q^{1-j\epsilon}|Bb + C| \geq c_0$. Since $P \in \mathcal{P}$, there exists $x \in \mathbf{A}_0 \cap \Delta(P)$ and it follows that

$$\begin{aligned} q^{1-j\epsilon}|Ex| &\geq q^{1-j\epsilon}|Bb + C| - q^{1-j\epsilon}|Ex + Bb + C| \\ &\stackrel{(6.7)}{\geq} c_0 - 2\kappa c \geq c_1(|x|_{\max} + l) - c_1 l = c_1|x|_{\max}. \end{aligned}$$

This proves the lemma. \square

A particular consequence of [\(6.8\)](#) is that $E \neq 0$. Thus every line L_P intersects at the line $L_{a,b}$ given by $y = f(x) = ax + b$ at a single point.

For $n \geq 1$, let

$$H_n := 4\kappa cl^{-1}R^n$$

and

$$\mathcal{P}_n := \left\{ P = \left(\frac{p}{q}, \frac{r}{q} \right) \in \mathcal{P} : H_n \leq q|E| < H_{n+1} \right\}.$$

Note that if $P \in \mathcal{P}_n$, then

$$\kappa q^{1+i} \stackrel{(6.6)}{\geq} q|E| \geq H_n. \quad (6.9)$$

Next let

$$\mathcal{P}_{n,k} := \{ P \in \mathcal{P}_n : H_n R^{\lambda_{k-1}} \leq \kappa q^{1+i} < H_n R^{\lambda_k} \} \quad \text{for } 1 \leq k \leq n.$$

Lemma 6.5. *With \mathcal{P}_n and $\mathcal{P}_{n,k}$ as above, we have that*

$$\mathcal{P} = \bigcup_{n=1}^{\infty} \mathcal{P}_n \quad \text{and} \quad \mathcal{P}_n = \bigcup_{k=1}^n \mathcal{P}_{n,k}.$$

Proof. Note that by (6.8), for any $P \in \mathcal{P}$ we have that $q|E| \geq c_1$ and by definition

$$H_1 = 4\kappa cl^{-1}R \leq c_1. \quad (6.10)$$

Thus, $\mathcal{P} = \bigcup_{n=1}^{\infty} \mathcal{P}_n$.

To prove the second conclusion, we first show that

$$\kappa q^{1+i} < H_n R^{\lambda_n} \quad \text{for } P \in \mathcal{P}_n. \quad (6.11)$$

We consider two separate cases. If $a \notin \mathbb{Q}$, on combining the fact that $q|E| < H_{n+1}$ with (6.8) implies that

$$q^{j\epsilon} < c_1^{-1} H_{n+1}.$$

It then follows that

$$\begin{aligned} \frac{\kappa q^{1+i}}{H_n R^{\lambda_n}} &< \frac{\kappa (c_1^{-1} H_{n+1})^{\frac{1+i}{j\epsilon}}}{H_n R^{\lambda_n}} = \frac{\kappa c_1^{-1} (c_1^{-1} H_1 R^n)^{\frac{1+i}{j\epsilon}-1}}{R^{\lambda_n-1}} \\ &\stackrel{(6.10)}{\leq} \kappa c_1^{-1} R^{(\frac{1+i}{j\epsilon}-1)n-\lambda_j^{-n}} \stackrel{(6.4)}{\leq} 1. \end{aligned}$$

If $a \in \mathbb{Q}$, then $|E| \geq 1/d$, and hence the fact $q|E| < H_{n+1}$ implies that $q < dH_{n+1}$. It follows that

$$\frac{\kappa q^{1+i}}{H_n R^{\lambda_n}} < \frac{\kappa d^2 H_{n+1}^{1+i}}{H_n R^{\frac{n}{i}+\mu}} \stackrel{(6.3)}{\leq} H_1^i R^{n(i-\frac{1}{i})} < 1.$$

This proves (6.11). Now (6.11) together with (6.9) implies that $\mathcal{P}_n = \bigcup_{k=1}^n \mathcal{P}_{n,k}$. \square

We claim that the partition of \mathcal{P} given by Lemma 6.5 satisfies the requirement of Proposition 6.3. The key towards establishing the claim is the following lemma. It is the lines analogue of Lemma 4.2.

Lemma 6.6. *For any $n \geq 1$, $1 \leq k \leq n$ and $\tau \in \mathcal{S}_{n-k}$, the map $P \mapsto L_P$ is constant on*

$$\mathcal{P}_{n,k}(\tau) := \{P \in \mathcal{P}_{n,k} : \mathcal{I}(\tau) \cap \Delta(P) \neq \emptyset\}.$$

Proof. Let $P_1 = (\frac{p_1}{q_1}, \frac{r_1}{q_1})$ and $P_2 = (\frac{p_2}{q_2}, \frac{r_2}{q_2})$ be distinct points in $\mathcal{P}_{n,k}(\tau)$. We need to prove that $L_{P_1} = L_{P_2}$. We let A_s, B_s, C_s and E_s be the respective quantities associated with P_s , $s = 1, 2$, and consider two separate cases.

Case (1). Suppose $k = 1$. Then

$$|A_1 p_2 + B_1 r_2 + C_1 q_2| = q_2 \left| A_1 \left(\frac{p_2}{q_2} - \frac{p_1}{q_1} \right) + B_1 \left(\frac{r_2}{q_2} - \frac{r_1}{q_1} \right) \right|$$

$$\begin{aligned}
&= q_2 \left| E_1 \left(\frac{p_2}{q_2} - \frac{p_1}{q_1} \right) + B_1 \left(\frac{ap_1 - r_1}{q_1} - \frac{ap_2 - r_2}{q_2} \right) \right| \\
&\leq q_2 |E_1| \left(\frac{c}{q_1^{1+i}} + \frac{c}{q_2^{1+i}} + lR^{-n+1} \right) + q_2 |B_1| \left(\frac{\kappa c}{q_1^{1+j}} + \frac{\kappa c}{q_2^{1+j}} \right) \\
&\leq \frac{q_2}{q_1} |E_1| \left(\frac{2\kappa c}{H_n} + lR^{-n+1} \right) + \kappa c \left(\frac{q_2}{q_1} + \frac{q_1^j}{q_2^j} \right) \\
&\leq R^{\frac{\lambda_1}{1+i}} H_{n+1} \cdot \frac{2\kappa c(1+2R)}{H_n} + 2\kappa c R^{\frac{\lambda_1}{1+i}} \\
&< 8\kappa c R^{2+\frac{\lambda_1}{1+i}} \leq 1.
\end{aligned}$$

Since the left hand side of the above inequality is an integer, it follows that $A_1 p_2 + B_1 r_2 + C_1 q_2 = 0$. Similarly, we obtain that $A_2 p_1 + B_2 r_1 + C_2 q_1 = 0$. The upshot is that both the lines L_{P_1} and L_{P_2} pass through both the points P_1 and P_2 , and so we must have that $L_{P_1} = L_{P_2}$.

Case (2). Suppose $k \geq 2$. We prove that $L_{P_1} = L_{P_2}$ by contradiction. Thus, assume that $L_{P_1} \neq L_{P_2}$. We first establish various preliminary estimates. Let

$$m_q := (\kappa^{-1} H_n R^{\lambda_{k-1}})^{\frac{1}{1+i}} \quad \text{and} \quad M_q := (\kappa^{-1} H_n R^{\lambda_k})^{\frac{1}{1+i}}.$$

Then, by definition, for $P = (\frac{p}{q}, \frac{r}{q}) \in \mathcal{P}_{n,k}$ we have that $m_q \leq q < M_q$. Also let

$$M_E := m_q^{-1} H_{n+1} \quad \text{and} \quad M_B := M_q^j.$$

Then $|E_s| \leq M_E$, $|B_s| \leq M_B$, $s = 1, 2$. We claim that

$$M_E^{1+\frac{1}{i}} \leq M_B^j M_E^{2+j} \leq M_B^{1+i} M_E^{1+i} < \frac{1}{5} c l^{-1} R^{n-k}. \quad (6.12)$$

First observe that

$$\frac{M_E}{M_B^{j/j}} = \frac{H_{n+1}}{m_q M_q^i} = \kappa R^{1-\frac{\lambda_{k-1}+i\lambda_k}{1+i}} \leq \kappa R^{1-\mu} \leq 1.$$

It then follows that

$$\frac{M_E^{1+\frac{1}{i}}}{M_B^j M_E^{2+j}} = \left(\frac{M_E}{M_B^{i/j}} \right)^{\frac{j^2}{i}} \leq 1 \quad \text{and} \quad \frac{M_B^j M_E^{2+j}}{M_B^{1+i} M_E^{1+i}} = \left(\frac{M_E}{M_B^{i/j}} \right)^{2j} \leq 1.$$

This establishes the left and middle inequalities within (6.12). Regarding the right inequality, we have that

$$\begin{aligned} M_B^{1+i} M_E^{1+i} &= \left(\frac{H_{n+1}}{M_q^i} \cdot \frac{M_q}{m_q} \right)^{1+i} = \frac{\kappa^i H_{n+1}^{1+i}}{(H_n R^{\lambda_k})^i} \cdot R^{\lambda_k - \lambda_{k-1}} \\ &= \kappa^i H_n R^{j\lambda_k - \lambda_{k-1} + 1 + i} = 4\kappa^{1+i} cl^{-1} R^{n-k + \frac{1}{i} - i\mu + 1 + i} \\ &< \frac{1}{5} cl^{-1} R^{n-k}. \end{aligned}$$

With $s = 1$ or 2 , let $(x_s, ax_s + b)$ denote the intersection point of L_{P_s} with the line $L_{a,b}$. Then

$$E_s \left(x_s - \frac{p_s}{q_s} \right) + B_s \left(b + \frac{ap_s - r_s}{q_s} \right) = 0,$$

and so

$$\left| x_s - \frac{p_s}{q_s} \right| = \frac{|B_s|}{|E_s|} \left| b + \frac{ap_s - r_s}{q_s} \right| \leq \frac{|B_s|}{|E_s|} \frac{\kappa c}{q_s^{1+j}}, \quad s = 1, 2.$$

Hence

$$\begin{aligned} |x_1 - x_2| &\leq \left| x_1 - \frac{p_1}{q_1} \right| + \left| x_2 - \frac{p_2}{q_2} \right| + \left| \frac{p_1}{q_1} - \frac{p_2}{q_2} \right| \\ &\leq \frac{|B_1|}{|E_1|} \frac{\kappa c}{q_1^{1+j}} + \frac{|B_2|}{|E_2|} \frac{\kappa c}{q_2^{1+j}} + \left(\frac{c}{q_1^{1+i}} + \frac{c}{q_2^{1+i}} + lR^{-n+k} \right) \\ &\leq \frac{\kappa c}{q_1 |E_1|} + \frac{\kappa c}{q_2 |E_2|} + \left(\frac{\kappa c}{q_1 |E_1|} + \frac{\kappa c}{q_2 |E_2|} + lR^{-n+k} \right) \\ &\leq \frac{4\kappa c}{H_n} + lR^{-n+k} = lR^{-n} + lR^{-n+k} \\ &\leq 2lR^{-n+k}. \end{aligned} \tag{6.13}$$

This completes the preliminaries. Recall that we are assuming that $L_{P_1} \neq L_{P_2}$ and the name of the game is to obtain a contradiction. We first consider the case that L_{P_1} is parallel to L_{P_2} . Then there exist $(A, B) \in \mathbb{Z}^2 \setminus \{(0, 0)\}$ and nonzero integers t_1, t_2 such that

$$(A_1, B_1) = t_1(A, B) \quad \text{and} \quad (A_2, B_2) = t_2(A, B).$$

Thus

$$x_s = -\frac{B_s b + C_s}{B_s a + A_s} = -\frac{1}{Ba + A} \left(Bb + \frac{C_s}{t_s} \right) \quad s = 1, 2$$

and it follows that

$$|x_1 - x_2| = \frac{|t_1 C_2 - t_2 C_1|}{|t_1 t_2| |Ba + A|} \geq \frac{1}{|t_1 t_2| |Ba + A|} = \frac{1}{|t_1 E_2|}.$$

This together with (6.13) implies that

$$|t_1 E_2| \geq \frac{1}{2} l^{-1} R^{n-k}. \quad (6.14)$$

If $B_1 = 0$, then $|t_1| \leq |A_1| \leq |A_1|^{1/i} = |E_1|^{1/i}$. So

$$|t_1 E_2| \leq M_E^{1+\frac{1}{i}} \stackrel{(6.12)}{<} \frac{1}{5} c l^{-1} R^{n-k}$$

and this contradicts (6.14). If $B_1 \neq 0$, then $|t_1| \leq |B_1|$ and

$$|t_1 E_2|^{1+i} \leq (M_B M_E)^{1+i} \stackrel{(6.12)}{<} \frac{1}{5} c l^{-1} R^{n-k}.$$

This together with (6.14) implies that

$$\left(\frac{1}{2} l^{-1} R^{n-k}\right)^{1+i} < \frac{1}{5} c l^{-1} R^{n-k}.$$

However, this contradicts the fact that $c \leq l^{-i}$. Hence, if L_{P_1} is parallel to L_{P_2} then we must have that $L_{P_1} = L_{P_2}$.

Now suppose L_{P_1} is not parallel to L_{P_2} . Let $P_0 = (\frac{p_0}{q_0}, \frac{r_0}{q_0}) \in \mathbb{Q}^2$ be the intersection of L_{P_1} and L_{P_2} . Then it follows that $A_1 B_2 - A_2 B_1$ is a nonzero integer and is divisible by q_0 and so

$$q_0 \leq |A_1 B_2 - A_2 B_1| = |E_1 B_2 - E_2 B_1|. \quad (6.15)$$

We first prove that $\Delta(P_1) \subset \Delta(P_0)$ and that $P_0 \in \mathcal{P}$. In view of the fact that

$$E_s \left(\frac{p_s}{q_s} - \frac{p_0}{q_0} \right) + B_s \left(\frac{r_s - ap_s}{q_s} - \frac{r_0 - ap_0}{q_0} \right) = 0 \quad s = 1, 2$$

it is easily verified that

$$-(E_1 B_2 - E_2 B_1) \left(\frac{p_1}{q_1} - \frac{p_0}{q_0} \right) = B_1 B_2 \left(\frac{r_1 - ap_1}{q_1} - \frac{r_2 - ap_2}{q_2} \right) + B_1 E_2 \left(\frac{p_1}{q_1} - \frac{p_2}{q_2} \right).$$

It then follows that

$$\begin{aligned} q_0 \left| \frac{p_1}{q_1} - \frac{p_0}{q_0} \right| &\stackrel{(6.15)}{\leq} |B_1 B_2| \left(\frac{\kappa c}{q_1^{1+j}} + \frac{\kappa c}{q_2^{1+j}} \right) + |B_1 E_2| \left(\frac{c}{q_1^{1+i}} + \frac{c}{q_2^{1+i}} + l R^{-n+k} \right) \\ &\leq |B_2 E_1| \frac{\kappa c}{q_1 |E_1|} + |B_1 E_2| \frac{\kappa c}{q_2 |E_2|} + |B_1 E_2| \left(\frac{\kappa c}{q_1 |E_1|} + \frac{\kappa c}{q_2 |E_2|} + l R^{-n+k} \right) \\ &\leq M_B M_E \left(\frac{4\kappa c}{H_n} + l R^{-n+k} \right) \\ &\leq 2M_B M_E l R^{-n+k}. \end{aligned}$$

So if $x \in \Delta(P_1)$, then

$$\begin{aligned}
 q_0^{1+i} \left| x - \frac{p_0}{q_0} \right| &\leq q_0^{1+i} \left| x - \frac{p_1}{q_1} \right| + q_0^{1+i} \left| \frac{p_1}{q_1} - \frac{p_0}{q_0} \right| \\
 &\leq q_0^{1+i} \frac{c}{q_1^{1+i}} + 2q_0^i M_B M_E l R^{-n+k} \\
 &\stackrel{(6.15)}{\leq} 4M_B^{1+i} M_E^{1+i} \frac{\kappa c}{H_n} + 4M_B^{1+i} M_E^{1+i} l R^{-n+k} \\
 &\leq 5M_B^{1+i} M_E^{1+i} l R^{-n+k} \stackrel{(6.12)}{<} c.
 \end{aligned}$$

Thus $x \in \Delta(P_0)$ and the upshot is that $\Delta(P_1) \subset \Delta(P_0)$. In turn, since $\mathbf{A}_0 \cap \Delta(P_1) \neq \emptyset$, it follows that $\mathbf{A}_0 \cap \Delta(P_0) \neq \emptyset$. In view of this, in order to prove that $P_0 \in \mathcal{P}$ we need to show that

$$\left| b + \frac{ap_0 - r_0}{q_0} \right| < \frac{\kappa c}{q_0^{1+j}}. \quad (6.16)$$

Since

$$E_s \left(x_s - \frac{p_0}{q_0} \right) + B_s \left(b + \frac{ap_0 - r_0}{q_0} \right) = 0 \quad s = 1, 2$$

we have that

$$|E_1 B_2 - E_2 B_1| \left| b + \frac{ap_0 - r_0}{q_0} \right| = |E_1 E_2| |x_1 - x_2| \stackrel{(6.13)}{\leq} 2M_E^2 l R^{-n+k}.$$

Hence

$$\begin{aligned}
 q_0^{1+j} \left| b + \frac{ap_0 - r_0}{q_0} \right| &\stackrel{(6.15)}{\leq} |E_1 B_2 - E_2 B_1|^{1+j} \left| b + \frac{ap_0 - r_0}{q_0} \right| \\
 &\leq 4M_B^j M_E^{2+j} l R^{-n+k} \\
 &\stackrel{(6.12)}{<} \kappa c
 \end{aligned}$$

and this established (6.16). Thus $P_0 \in \mathcal{P}$ and so there exists a unique integer $n_0 \geq 1$ such that $P_0 \in \mathcal{P}_{n_0}$. Suppose for the moment that $n_0 \leq n - k$. Then there exists $\tau' \in \mathcal{S}_{n_0}$ such that $\tau \prec \tau'$, and hence

$$\mathcal{I}(\tau) \cap \Delta(P_1) \subset \mathcal{I}(\tau') \cap \Delta(P_0) = \emptyset.$$

This contradicts the fact that $P_1 \in \mathcal{P}_{n,k}(\tau)$. Thus

$$n_0 \geq n - k + 1,$$

and so

$$q_0^{1+i} \stackrel{(6.9)}{\geq} \kappa^{-1} H_{n_0} \geq \kappa^{-1} H_{n-k+1} = 4cl^{-1} R^{n-k+1}.$$

On the other hand, we have that

$$q_0^{1+i} \leq 4(M_B M_E)^{1+i} \stackrel{(6.12)}{<} cl^{-1} R^{n-k}.$$

This contradicts the above lower bound for q_0^{1+i} and so completes the proof of Case (2) and indeed the lemma. \square

An important consequence of Lemma 6.6 is the following analogue of Corollary 4.3.

Corollary 6.7. *For any $n \geq 1$, $1 \leq k \leq n$ and $\tau \in \mathcal{S}_{n-k}$, we have*

$$\#\left\{\tau' \in \mathcal{T}_n : \mathcal{I}(\tau') \cap \bigcup_{P \in \mathcal{P}_{n,k}(\tau)} \Delta(P) \neq \emptyset\right\} \leq 2.$$

Proof. By Lemma 6.6 and (6.7), there exists $(A, B, C) \in \mathbb{Z}^3$ with $E := A + Ba \neq 0$ such that for any $P = (\frac{p}{q}, \frac{r}{q}) \in \mathcal{P}_{n,k}(\tau)$ and $x \in \Delta(P)$,

$$|Ex + Bb + C| < \frac{2\kappa c}{q} \quad \text{and} \quad q|E| \geq H_n.$$

Thus

$$\left|x + \frac{Bb + C}{E}\right| < \frac{2\kappa c}{q|E|} \leq \frac{2\kappa c}{H_n} = \frac{1}{2}lR^{-n}.$$

This implies that $\bigcup_{P \in \mathcal{P}_{n,k}(\tau)} \Delta(P)$ is contained in the open interval

$$\left(-\frac{Bb + C}{E} - \frac{1}{2}lR^{-n}, -\frac{Bb + C}{E} + \frac{1}{2}lR^{-n}\right), \quad (6.17)$$

which has length lR^{-n} . Since the intervals $\{\mathcal{I}(\tau') : \tau' \in \mathcal{T}_n\}$ are of length lR^{-n} and have mutually disjoint interiors, there can be at most 2 of them that intersect the interval (6.17). This proves the corollary. \square

We are now in the position to prove Proposition 6.3. In view of Proposition 2.2, it suffices to prove that the intersection of \mathcal{S} with every 6-regular subtree of \mathcal{T} is infinite. Let $\mathcal{R} \subset \mathcal{T}$ be a 6-regular subtree and let

$$\mathcal{R}' := \mathcal{R} \cap \mathcal{S} \quad \text{and} \quad a_n := \#\mathcal{R}'_n \quad (n \geq 0).$$

Then $a_0 = 1$. We prove that \mathcal{R}' is infinite by showing that

$$a_n > 2a_{n-1} \quad (n \geq 1). \quad (6.18)$$

We use induction. As in §4, for $n \geq 1$, let

$$\mathcal{U}_n := \left\{ \tau \in \mathcal{T}_{\text{suc}}(\mathcal{R}'_{n-1}) : \mathcal{I}(\tau) \cap \bigcup_{P \in \mathcal{P}_n} \Delta(P) \neq \emptyset \right\}.$$

Then

$$\mathcal{R}'_n = \mathcal{R}_{\text{suc}}(\mathcal{R}'_{n-1}) \setminus \mathcal{U}_n,$$

and it follows that

$$a_n \geq 6a_{n-1} - \#\mathcal{U}_n. \quad (6.19)$$

On the other hand, as in §4, we have that

$$\begin{aligned} \mathcal{U}_n &= \bigcup_{k=1}^n \left\{ \tau \in \mathcal{T}_{\text{suc}}(\mathcal{R}'_{n-1}) : \mathcal{I}(\tau) \cap \bigcup_{P \in \mathcal{P}_{n,k}} \Delta(P) \neq \emptyset \right\} \\ &\subset \bigcup_{k=1}^n \bigcup_{\tau' \in \mathcal{R}'_{n-k}} \left\{ \tau \in \mathcal{T}_n : \mathcal{I}(\tau) \cap \bigcup_{P \in \mathcal{P}_{n,k}(\tau')} \Delta(P) \neq \emptyset \right\}. \end{aligned}$$

Thus, Corollary 6.7 implies that

$$\#\mathcal{U}_n \leq \sum_{k=1}^n 2a_{n-k}. \quad (6.20)$$

On combining (6.19) and (6.20), we obtain that

$$a_n \geq 6a_{n-1} - \sum_{k=1}^n 2a_{n-k}. \quad (6.21)$$

With $n = 1$ in (6.21), we find that $a_1 \geq 4$. Hence, (6.18) holds for $n = 1$. Now assume $n \geq 2$ and that (6.18) holds with n replaced by $1, \dots, n-1$. Then for any $1 \leq k \leq n$, we have that

$$a_{n-k} \leq 2^{-k+1} a_{n-1}.$$

Substituting this into (6.21), gives that

$$a_n \geq 6a_{n-1} - 2a_{n-1} \sum_{k=1}^n 2^{-k+1} > 2a_{n-1}.$$

This completes the induction step and thus establishes (6.18). In turn this completes the proof of Proposition 6.3. \square

6.2. The inhomogeneous case: establishing Theorem 1.2

Theorem 1.2 is easily deduced from the following statement.

Theorem 6.8. *Let (i, j) be a pair of real numbers satisfying $0 < j \leq i < 1$ and $i + j = 1$. Given $a, b \in \mathbb{R}$, suppose there exists $\epsilon > 0$ such that (6.1) is satisfied. Then $\mathbf{Bad}_{\theta}^f(i, j)$ is a $1/2$ -winning subset of \mathbb{R} .*

We have already established the homogeneous case ($\gamma = \delta = 0$) of Theorem 6.8; namely Theorem 6.1. With reference to §6.1, the crux of the ‘homogeneous’ proof involved constructing a partition \mathcal{P}_n ($n \geq 1$) of \mathcal{P} (given by Lemma 6.5) such that the subtree \mathcal{S} of an $[R]$ -regular rooted tree \mathcal{T} has an $([R] - 5)$ -regular subtree \mathcal{S}' – the substance of Proposition 6.3. To prove Theorem 6.8, the idea is to merge the inhomogeneous constraints into the homogeneous construction as in the case of curves in §5. More precisely, we show that \mathcal{S}' has an $([R] - 7)$ -regular subtree \mathcal{Q}' that incorporates the inhomogeneous constraints. With this in mind, let

$$c' := \frac{1}{10}cR^{-2}$$

and

$$\mathcal{V} := \left\{ (p, r, q) \in \mathbb{Z}^2 \times \mathbb{N} : \left| f\left(\frac{p+\gamma}{q}\right) - \frac{r+\delta}{q} \right| < \frac{\kappa c'}{q^{1+j}} \right\}.$$

Furthermore, for each $v = (p, r, q) \in \mathbb{Z}^2 \times \mathbb{N}$, we associate the interval

$$\Delta_{\theta}(v) := \left\{ x \in \mathbb{R} : \left| x - \frac{p+\gamma}{q} \right| < \frac{c'}{q^{1+i}} \right\}.$$

Then, with $\mathbf{A}_0 \subset \mathbf{B}_0$ as in §6.1, the following is the inhomogeneous analogue of Lemma 6.2.

Lemma 6.9. *Let \mathbf{A}_0 , \mathcal{V} and $\Delta_{\theta}(v)$ be as above. Then*

$$\mathbf{A}_0 \setminus \bigcup_{v \in \mathcal{V}} \Delta_{\theta}(v) \subset \mathbf{Bad}_{\theta}^f(i, j).$$

Proof. The proof is similar to the homogeneous proof but is included for completeness. Let $x \in \mathbf{A}_0$. Suppose $x \notin \mathbf{Bad}_{\theta}^f(i, j)$. Then there exists $v = (p, r, q) \in \mathbb{Z}^2 \times \mathbb{N}$ such that

$$\left| x - \frac{p+\gamma}{q} \right| < \frac{c'}{q^{1+i}}, \quad \left| f(x) - \frac{r+\delta}{q} \right| < \frac{c'}{q^{1+j}}.$$

It follows that

$$\begin{aligned} \left| f\left(\frac{p+\gamma}{q}\right) - \frac{r+\delta}{q} \right| &\leq \left| f\left(\frac{p+\gamma}{q}\right) - f(x) \right| + \left| f(x) - \frac{r+\delta}{q} \right| \\ &< |a| \frac{c'}{q^{1+i}} + \frac{c'}{q^{1+j}} \leq \frac{\kappa c'}{q^{1+j}}. \end{aligned}$$

Thus $x \in \Delta_{\theta}(v)$ and $v \in \mathcal{V}$. This completes the proof of the lemma. \square

Similar to the proof of [Theorem 5.1](#), we prove [Theorem 6.8](#) by showing that Ayesha can play such that

$$\bigcap_{n=0}^{\infty} \mathbf{A}_n \subset \mathbf{A}_0 \setminus \left(\bigcup_{P \in \mathcal{P}} \Delta(P) \cup \bigcup_{v \in \mathcal{V}} \Delta_{\theta}(v) \right).$$

For $n \geq 1$, let

$$H'_n := 2c'l^{-1}R^n$$

and

$$\mathcal{V}_n := \{(p, r, q) \in \mathcal{V} : H'_n \leq q^{1+i} < H'_{n+1}\}.$$

Observe that $H'_1 = 2c'l^{-1}R \leq 1$ and so it follows that $\mathcal{V} = \bigcup_{n=1}^{\infty} \mathcal{V}_n$. We inductively define a subtree \mathcal{Q} of \mathcal{S}' as follows. Let $\mathcal{Q}_0 = \{\tau_0\}$. If \mathcal{Q}_{n-1} ($n \geq 1$) is defined, we let

$$\mathcal{Q}_n := \left\{ \tau \in \mathcal{S}'_{\text{suc}}(\mathcal{Q}_{n-1}) : \mathcal{I}(\tau) \cap \bigcup_{v \in \mathcal{V}_n} \Delta_{\theta}(v) = \emptyset \right\}.$$

Then

$$\mathcal{Q} := \bigcup_{n=0}^{\infty} \mathcal{Q}_n$$

is a subtree of \mathcal{S}' and by construction

$$\mathcal{I}(\tau) \subset \mathbf{A}_0 \setminus \left(\bigcup_{P \in \mathcal{P}} \Delta(P) \cup \bigcup_{v \in \mathcal{V}_n} \Delta_{\theta}(v) \right) \quad \forall n \geq 1 \quad \text{and} \quad \tau \in \mathcal{Q}_n.$$

Armed with the following result, the same arguments as in [§2.2.1](#) with the most obvious modifications enables us to prove [Theorem 6.8](#). In view of this the details of the proof of [Theorem 6.8](#) modulo [Proposition 6.10](#) are omitted.

Proposition 6.10. *The tree \mathcal{Q} has an $([R] - 7)$ -regular subtree.*

In order to establish the proposition, it suffices to prove the following statement.

Lemma 6.11. *For any $n \geq 1$ and $\tau \in \mathcal{Q}_{n-1}$, there is at most one $v \in \mathcal{V}_n$ such that $\mathcal{I}(\tau) \cap \Delta_{\theta}(v) \neq \emptyset$. Moreover, $\rho(\Delta_{\theta}(v)) \leq lR^{-n}$. Therefore,*

$$\#\left\{\tau' \in \mathcal{S}'_{\text{suc}}(\tau) : \mathcal{I}(\tau') \cap \bigcup_{v \in \mathcal{V}_n} \Delta_{\theta}(v) \neq \emptyset\right\} \leq 2.$$

Proof. Suppose $v_s = (p_s, r_s, q_s) \in \mathcal{V}_n$ and $\mathcal{I}(\tau) \cap \Delta_{\theta}(v_s) \neq \emptyset$, $s = 1, 2$. We need to prove that $v_1 = v_2$. Without loss of generality, assume that $q_1 \geq q_2$. Let $x_1 \in \mathcal{I}(\tau) \cap \Delta_{\theta}(v_1)$. The same arguments as in the proofs of (5.3) and (5.4) show that

$$|(q_1 - q_2)x_1 - (p_1 - p_2)| \leq q_2 l R^{-n+1} + \frac{2c'}{q_2^i} \quad (6.22)$$

and

$$\left| (q_1 - q_2) f\left(\frac{p_1 + \gamma}{q_1}\right) - (r_1 - r_2) \right| \leq q_2 |a| l R^{-n+1} + \frac{4\kappa c'}{q_2^j}. \quad (6.23)$$

Suppose for the moment that $q_1 > q_2$ and let

$$P_0 := \left(\frac{p_1 - p_2}{q_1 - q_2}, \frac{r_1 - r_2}{q_1 - q_2} \right).$$

We show that $\mathcal{I}(\tau) \cap \Delta(P_0) \neq \emptyset$ and that $P_0 \in \mathcal{P}$. Similar to the proof of Lemma 5.4, it follows from (6.22) that

$$(q_1 - q_2)^{1+i} \left| x_1 - \frac{p_1 - p_2}{q_1 - q_2} \right| < c.$$

So $x_1 \in \Delta(P_0)$ and it follows that $\mathcal{I}(\tau) \cap \Delta(P_0) \neq \emptyset$. Moreover, by making use of (6.23) we have that

$$(q_1 - q_2)^{1+j} \left| f\left(\frac{p_1 - p_2}{q_1 - q_2}\right) - \frac{r_1 - r_2}{q_1 - q_2} \right| < \kappa c.$$

Thus $P_0 \in \mathcal{P}$ and so there exists a unique integer $n_0 \geq 1$ such that $P_0 \in \mathcal{P}_{n_0}$. The same argument as in the proof of Lemma 5.4 shows that $n_0 \geq n$, and so

$$(q_1 - q_2) |E_{P_0}| \geq H_{n_0} \geq H_n = 4\kappa c l^{-1} R^n.$$

On the other hand, we have that

$$\begin{aligned} (q_1 - q_2) |E_{P_0}| &\leq \kappa (q_1 - q_2)^{1+i} \leq \kappa q_1^{1+i} \\ &\leq \kappa H'_{n+1} = \frac{1}{5} \kappa c l^{-1} R^{n-1}. \end{aligned}$$

This contradicts the above lower bound for $(q_1 - q_2)|E_{P_0}|$ and we conclude that $q_1 = q_2$. It now follows from (6.22) and (6.23) that

$$|p_1 - p_2| < 1 \quad \text{and} \quad |r_1 - r_2| < 1.$$

Thus, $p_1 = p_2$ and $r_1 = r_2$. In other words, $v_1 = v_2$ and this proves the main substance of the proposition. The proofs of the remaining parts are the same as those for Lemma 5.4. \square

As already mentioned, given Proposition 6.10, the proof of Theorem 6.8 follows on adapting the arguments of §2.2.1.

7. The proof of Theorem 1.3

We need the notion of regular colourings of rooted trees. Let $D \in \mathbb{N}$. A D -colouring of a rooted tree \mathcal{T} is a map $\gamma : \mathcal{T} \rightarrow \{1, \dots, D\}$. For $\mathcal{V} \subset \mathcal{T}$ and $1 \leq i \leq D$, we denote $\mathcal{V}^{(i)} = \mathcal{V} \cap \gamma^{-1}(i)$. Let $N \in \mathbb{N}$ be an integer multiple of D , and suppose that \mathcal{T} is N -regular. We say that a D -colouring of \mathcal{T} is *regular* if for any $\tau \in \mathcal{T}$ and $1 \leq i \leq D$, we have $\#\mathcal{T}_{\text{suc}}(\tau)^{(i)} = N/D$. The following two types of subtrees are of interest to us.

- The subtree \mathcal{S} is of *type (I)* if for any $\tau \in \mathcal{S}$ and $1 \leq i \leq D$, we have $\#\mathcal{S}_{\text{suc}}(\tau)^{(i)} = 1$.
- The subtree \mathcal{S} is of *type (II)* if for any $\tau \in \mathcal{S}$, there exists $1 \leq i(\tau) \leq D$ such that $\mathcal{S}_{\text{suc}}(\tau) = \mathcal{T}_{\text{suc}}(\tau)^{(i(\tau))}$.

Roughly speaking, in the proof of Theorem 1.3, the two types of subtrees correspond to strategies of the two players in Schmidt's game. We will make use of the following criterion for the existence of subtree of type (I). It appears as Proposition 2.2 in [2].

Proposition 7.1. *Let \mathcal{T} be an N -regular rooted tree with a regular D -colouring, and let $\mathcal{S} \subset \mathcal{T}$ be a subtree. Suppose that for every subtree $\mathcal{R} \subset \mathcal{T}$ of type (II), $\mathcal{S} \cap \mathcal{R}$ is infinite. Then \mathcal{S} contains a subtree of type (I).*

7.1. The winning strategy for Theorem 1.3

Let $\alpha_0 := (30\sqrt{2})^{-1}$, $\beta \in (0, 1)$. We want to prove that $\mathbf{Bad}_\theta(i, j)$ is (α_0, β) -winning. In the first round of the game, Bhupen chooses a closed disc $\mathbf{B}_0 \subset \mathbb{R}^2$. Now Ayesha chooses any closed disc $\mathbf{A}_0 \subset \mathbf{B}_0$ with diameter $\rho(\mathbf{A}_0) = \alpha_0 \rho(\mathbf{B}_0)$. Let

$$l := \rho(\mathbf{A}_0), \quad R := (\alpha_0 \beta)^{-1}, \quad m := 15.$$

By a *square* we mean a set of the form

$$\Sigma = \{(x, y) \in \mathbb{R}^2 : x_0 \leq x \leq x_0 + \ell(\Sigma), y_0 \leq y \leq y_0 + \ell(\Sigma)\},$$

where $\ell(\Sigma) > 0$ is the side length of Σ . Let Σ_0 be the circumscribed square of \mathbf{A}_0 . Then $\ell(\Sigma_0) = l$. Let \mathcal{T} be an $m^2[R/m]^2$ -regular rooted tree with a regular $[R/m]^2$ -colouring. We choose and fix an injective map Φ from \mathcal{T} to the set of subsquares of Σ_0 satisfying the following conditions:

- For any $n \geq 0$ and $\tau \in \mathcal{T}_n$, we have $\ell(\Phi(\tau)) = lR^{-n}$. In particular, the root τ_0 of \mathcal{T} is mapped to Σ_0 .
- For $\tau, \tau' \in \mathcal{T}$, if $\tau \prec \tau'$, then $\Phi(\tau) \subset \Phi(\tau')$.
- For any $n \geq 1$ and $\tau \in \mathcal{T}_{n-1}$, the interiors of the squares $\{\Phi(\tau') : \tau' \in \mathcal{T}_{\text{suc}}(\tau)\}$ are mutually disjoint, the union $\bigcup_{\tau' \in \mathcal{T}_{\text{suc}}(\tau)} \Phi(\tau')$ is a square of side length $m[R/m]lR^{-n}$, and for any $1 \leq i \leq [R/m]^2$, the union $\bigcup_{\tau' \in \mathcal{T}_{\text{suc}}(\tau)^{(i)}} \Phi(\tau')$ is a square of side length mlR^{-n} .

Let $c > 0$ be such that

$$c < \min \left\{ \frac{1}{6}lR^{-1}, \frac{1}{16}R^{-12} \right\}, \quad (7.1)$$

and in turn let

$$c' := \frac{1}{6}cR^{-2}. \quad (7.2)$$

For each $P = (\frac{p}{q}, \frac{r}{q}) \in \mathbb{Q}^2$, we associate the rectangle

$$\Delta(P) := \left\{ (x, y) \in \mathbb{R}^2 : \left| x - \frac{p}{q} \right| \leq \frac{c}{q^{1+i}}, \left| y - \frac{r}{q} \right| \leq \frac{c}{q^{1+j}} \right\},$$

and for $v = (p, r, q) \in \mathbb{Z}^2 \times \mathbb{N}$, we associate the rectangle

$$\Delta_{\theta}(v) := \left\{ (x, y) \in \mathbb{R}^2 : \left| x - \frac{p+\gamma}{q} \right| \leq \frac{c'}{q^{1+i}}, \left| y - \frac{r+\delta}{q} \right| \leq \frac{c'}{q^{1+j}} \right\}.$$

Then

$$\mathbb{R}^2 \setminus \left(\bigcup_{P \in \mathbb{Q}^2} \Delta(P) \cup \bigcup_{v \in \mathbb{Z}^2 \times \mathbb{N}} \Delta_{\theta}(v) \right) \subset \mathbf{Bad}(i, j) \cap \mathbf{Bad}_{\theta}(i, j). \quad (7.3)$$

For $n \geq 1$, let

$$H_n := 6cl^{-1}R^n, \quad H'_n := 3c'l^{-1}R^n$$

and define

$$\mathcal{P}_n := \left\{ P = \left(\frac{p}{q}, \frac{r}{q} \right) \in \mathbb{Q}^2 : H_n \leq q \max\{|A_P|, |B_P|\} < H_{n+1} \right\} \quad (7.4)$$

and

$$\mathcal{V}_n := \{v = (p, r, q) \in \mathbb{Z}^2 \times \mathbb{N} : H'_n \leq q^{1+\max\{i,j\}} < H'_{n+1}\}, \quad (7.5)$$

where A_P and B_P are as in §3. In view of (7.1), we have that $H'_1 \leq H_1 = 6cl^{-1}R \leq 1$. Thus

$$\mathbb{Q}^2 = \bigcup_{n=1}^{\infty} \mathcal{P}_n \quad \text{and} \quad \mathbb{Z}^2 \times \mathbb{N} = \bigcup_{n=1}^{\infty} \mathcal{V}_n.$$

We inductively define a subtree \mathcal{S} of \mathcal{T} as follows. Let $\mathcal{S}_0 = \{\tau_0\}$. If $n \geq 1$ and \mathcal{S}_{n-1} is defined, we let

$$\mathcal{S}_n := \left\{ \tau \in \mathcal{T}_{\text{suc}}(\mathcal{S}_{n-1}) : \Phi(\tau) \cap \left(\bigcup_{P \in \mathcal{P}_n} \Delta(P) \cup \bigcup_{v \in \mathcal{V}_n} \Delta_{\theta}(v) \right) = \emptyset \right\}. \quad (7.6)$$

Then $\mathcal{S} = \bigcup_{n=0}^{\infty} \mathcal{S}_n$ is a subtree of \mathcal{T} and by construction

$$\Phi(\tau) \subset \mathbb{R}^2 \setminus \left(\bigcup_{P \in \mathcal{P}_n} \Delta(P) \cup \bigcup_{v \in \mathcal{V}_n} \Delta_{\theta}(v) \right) \quad \forall n \geq 1 \quad \text{and} \quad \tau \in \mathcal{S}_n. \quad (7.7)$$

The following proposition is the key to proving Theorem 1.3.

Proposition 7.2. *The tree \mathcal{S} contains a subtree of type (I).*

7.1.1. Proof of Theorem 1.3 modulo Proposition 7.2

This is essentially the same as the proof of Theorem 1.1 in [2]. However for completeness we have included the short argument. Let \mathcal{S}' be a subtree of \mathcal{S} of type (I). We inductively prove that for every $n \geq 0$,

$$\text{Ayesha can choose } \mathbf{A}_n \text{ to be the inscribed closed disc of } \Phi(\tau_n) \text{ for some } \tau_n \in \mathcal{S}'_n. \quad (7.8)$$

If $n = 0$, there is nothing to prove. Assume $n \geq 1$ and Ayesha has chosen \mathbf{A}_{n-1} as the inscribed closed disc of $\Phi(\tau_{n-1})$, where $\tau_{n-1} \in \mathcal{S}'_{n-1}$. For any choice $\mathbf{B}_n \subset \mathbf{A}_{n-1}$ of Bhupen, the inscribed square of \mathbf{B}_n has side length

$$\frac{\sqrt{2}}{2} \rho(\mathbf{B}_n) = \frac{\sqrt{2}}{2} \beta \rho(\mathbf{A}_{n-1}) = \frac{\sqrt{2}}{2} \beta \ell(\Phi(\tau_{n-1})) = \frac{\sqrt{2}}{2} \beta l R^{-n+1} = 2mlR^{-n}.$$

Thus there exists $1 \leq i \leq [R/m]^2$ such that $\bigcup_{\tau \in \mathcal{T}_{\text{suc}}(\tau_{n-1})^{(i)}} \Phi(\tau) \subset \mathbf{B}_n$. Let τ_n be the unique vertex in $\mathcal{S}'_{\text{suc}}(\tau_{n-1})^{(i)}$. Then $\Phi(\tau_n) \subset \mathbf{B}_n$. Note that the diameter of the inscribed closed disc of $\Phi(\tau_n)$ is equal to

$$\ell(\Phi(\tau_n)) = R^{-1} \ell(\Phi(\tau_{n-1})) = \alpha_0 \beta \rho(\mathbf{A}_{n-1}) = \alpha_0 \rho(\mathbf{B}_n).$$

So Ayesha can choose \mathbf{A}_n to be the inscribed closed disc of $\Phi(\tau_n)$. This proves (7.8).

In view of (7.8), (7.7) and (7.3), we have

$$\begin{aligned} \bigcap_{n=0}^{\infty} \mathbf{A}_n &\subset \bigcap_{n=0}^{\infty} \Phi(\tau_n) \subset \bigcap_{n=1}^{\infty} \mathbb{R}^2 \setminus \left(\bigcup_{P \in \mathcal{P}_n} \Delta(P) \cup \bigcup_{v \in \mathcal{V}_n} \Delta_{\theta}(v) \right) \\ &= \mathbb{R}^2 \setminus \left(\bigcup_{P \in \mathbb{Q}^2} \Delta(P) \cup \bigcup_{v \in \mathbb{Z}^2 \times \mathbb{N}} \Delta_{\theta}(v) \right) \subset \mathbf{Bad}(i, j) \cap \mathbf{Bad}_{\theta}(i, j). \end{aligned}$$

This proves the theorem assuming the truth of Proposition 7.2. \square

7.2. Proof of Proposition 7.2

Let $w > 0$. By a strip of width w , we mean a subset of \mathbb{R}^2 of the form

$$\mathcal{L} := \{\mathbf{x} \in \mathbb{R}^2 : |\mathbf{x} \cdot \mathbf{u} - a| \leq w/2\},$$

where the dot denotes the standard inner product, $\mathbf{u} \in \mathbb{R}^2$ is a unit vector, and $a \in \mathbb{R}$. The following result is proved in [2, Corollary 4.2].

Lemma 7.3. *For any $n \geq 1$, there exists a partition $\mathcal{P}_n = \bigcup_{k=1}^n \mathcal{P}_{n,k}$ such that for any $1 \leq k \leq n$ and $\tau \in \mathcal{S}_{n-k}$, there is a strip of width $\frac{2}{3}lR^{-n}$ which contains all the rectangles*

$$\{\Delta(P) : P \in \mathcal{P}_{n,k}, \Phi(\tau) \cap \Delta(P) \neq \emptyset\}.$$

We now prove a corresponding result which takes into consideration the inhomogeneous approximation aspect.

Lemma 7.4. *For any $n \geq 1$ and $\tau \in \mathcal{S}_{n-1}$, there is at most one $v \in \mathcal{V}_n$ such that $\Phi(\tau) \cap \Delta_{\theta}(v) \neq \emptyset$. Moreover, $\Delta_{\theta}(v)$ is contained in a strip of width $\frac{2}{3}lR^{-n}$.*

Proof. Suppose $v_s = (p_s, r_s, q_s) \in \mathcal{V}_n$ and $\Phi(\tau) \cap \Delta_{\theta}(v_s) \neq \emptyset$, $s = 1, 2$. We need to prove that $v_1 = v_2$. Without loss of generality, assume that $q_1 \geq q_2$. Since $\Phi(\tau) \cap \Delta_{\theta}(v_s) \neq \emptyset$, there exists $(x_s, y_s) \in \Phi(\tau)$ such that

$$|q_s x_s - (p_s + \gamma)| \leq \frac{c'}{q_s^i}, \quad |q_s y_s - (r_s + \delta)| \leq \frac{c'}{q_s^j}.$$

It follows that

$$\begin{aligned} |(q_1 - q_2)x_1 - (p_1 - p_2)| &\leq q_2|x_1 - x_2| + |q_1x_1 - (p_1 + \gamma)| + |q_2x_2 - (p_2 + \gamma)| \\ &\leq q_2lR^{-n+1} + \frac{2c'}{q_2^i}. \end{aligned} \tag{7.9}$$

Similarly, we have that

$$|(q_1 - q_2)y_1 - (r_1 - r_2)| \leq q_2 l R^{-n+1} + \frac{2c'}{q_2^j}. \quad (7.10)$$

We first prove that $q_1 = q_2$. Suppose this is not the case. Then

$$\begin{aligned} \left| x_1 - \frac{p_1 - p_2}{q_1 - q_2} \right| &\leq \frac{1}{q_1 - q_2} \left(q_2 l R^{-n+1} + \frac{2c'}{q_2^j} \right) \\ &\leq \frac{1}{(q_1 - q_2)^{1+i}} \left(q_1^i q_2 l R^{-n+1} + 2c' \frac{q_1^i}{q_2^i} \right) \\ &\leq \frac{1}{(q_1 - q_2)^{1+i}} (H'_{n+1} l R^{-n+1} + 2c' R) \\ &= \frac{1}{(q_1 - q_2)^{1+i}} \left(\frac{c}{2} + \frac{1}{3} c R^{-1} \right) \\ &\leq \frac{c}{(q_1 - q_2)^{1+i}}. \end{aligned}$$

Similarly,

$$\left| y_1 - \frac{r_1 - r_2}{q_1 - q_2} \right| \leq \frac{c}{(q_1 - q_2)^{1+j}}.$$

Thus, if we let

$$P_0 := \left(\frac{p_1 - p_2}{q_1 - q_2}, \frac{r_1 - r_2}{q_1 - q_2} \right),$$

then $(x_1, y_1) \in \Delta(P_0)$. In particular, $\Phi(\tau) \cap \Delta(P_0) \neq \emptyset$. Let $n_0 \geq 1$ be the unique integer such that $P_0 \in \mathcal{P}_{n_0}$. It is easily verified that

$$n_0 \geq n. \quad (7.11)$$

Indeed, if $n_0 \leq n - 1$, then \mathcal{S}_{n_0} contains an ancestor τ' of τ and by (7.6) we have that

$$\Phi(\tau) \cap \Delta(P_0) \subset \Phi(\tau') \cap \Delta(P_0) = \emptyset.$$

This is a contradiction since the left hand side is non-empty. Now, in view of (7.4) and (7.11), we have that

$$(q_1 - q_2)^{1+\max\{i,j\}} \geq (q_1 - q_2) \max\{|A_{P_0}|, |B_{P_0}|\} \geq H_{n_0} \geq H_n = 6cl^{-1}R^n.$$

On the other hand, we have that

$$(q_1 - q_2)^{1+\max\{i,j\}} \leq q_1^{1+\max\{i,j\}} \leq H'_{n+1} = \frac{1}{2} cl^{-1} R^{n-1}.$$

This contradicts the above lower bound and so we must have that $q_1 = q_2$. It then follows via (7.9) and (7.10) that

$$\begin{aligned} |p_1 - p_2| &\leq q_2 l R^{-n+1} + 2c' \leq c + 2c' < 1, \\ |r_1 - r_2| &\leq q_2 l R^{-n+1} + 2c' \leq c + 2c' < 1. \end{aligned}$$

The left hand sides of these inequalities are integers so we must have that $p_1 = p_2$ and $r_1 = r_2$. The upshot of this is that $v_1 = v_2$ and so establishes the main substance of the lemma. Regarding the moreover part, simply observe that for any $v = (p, r, q) \in \mathcal{V}_n$, $\Delta_{\theta}(v)$ is contained in a strip of width

$$\min \left\{ \frac{2c'}{q^{1+i}}, \frac{2c'}{q^{1+j}} \right\} = \frac{2c'}{q^{1+\max\{i,j\}}} \leq \frac{2c'}{H'_n} = \frac{2}{3} l R^{-n}. \quad \square$$

The following result proved in [2, Lemma 4.3] gives an upper bound for the number of certain squares which intersect a thin strip.

Lemma 7.5. *Let $\mathcal{R} \subset \mathcal{T}$ be a subtree of type (II), let $n \geq 1$, and let \mathcal{L} be a strip of width $\frac{2}{3} l R^{-n}$. Then for any $1 \leq k \leq n$ and $\tau \in \mathcal{R}_{n-k}$, we have that*

$$\#\{\tau' \in \mathcal{R}(\tau)_k : \Phi(\tau') \cap \mathcal{L} \neq \emptyset\} \leq (3m - 2)^k.$$

On combining Lemmas 7.3, 7.4 and 7.5, we obtain the following statement.

Corollary 7.6. *Let $\mathcal{R} \subset \mathcal{T}$ be a subtree of type (II) and let $n \geq 1$. Then*

- For any $\tau \in \mathcal{S}_{n-1} \cap \mathcal{R}_{n-1}$, we have

$$\#\left\{\tau' \in \mathcal{R}_{\text{suc}}(\tau) : \Phi(\tau') \cap \left(\bigcup_{P \in \mathcal{P}_{n,1}} \Delta(P) \cup \bigcup_{v \in \mathcal{V}_n} \Delta_{\theta}(v) \right) \neq \emptyset \right\} \leq 2(3m - 2).$$

- For any $2 \leq k \leq n$ and $\tau \in \mathcal{S}_{n-k} \cap \mathcal{R}_{n-k}$, we have

$$\#\left\{\tau' \in \mathcal{R}(\tau)_k : \Phi(\tau') \cap \bigcup_{P \in \mathcal{P}_{n,k}} \Delta(P) \neq \emptyset \right\} \leq (3m - 2)^k.$$

We are now in the position to prove Proposition 7.2. The proof is essentially the same as the proof of Proposition 3.3 in [2]. However for completeness we have included the argument. In view of Proposition 7.1, it suffices to prove that the intersection of \mathcal{S} with every subtree of type (II) is infinite. Let $\mathcal{R} \subset \mathcal{T}$ be a subtree of type (II), and let

$$\mathcal{R}' := \mathcal{R} \cap \mathcal{S}, \quad \text{and} \quad a_n := \#\mathcal{R}'_n \quad (n \geq 0).$$

Then $a_0 = 1$. We prove that \mathcal{R}' is infinite by showing that

$$a_n > 112a_{n-1} \quad (n \geq 1). \quad (7.12)$$

We use induction. For $n \geq 1$, let

$$\mathcal{U}_n := \left\{ \tau \in \mathcal{R}_{\text{suc}}(\mathcal{R}'_{n-1}) : \Phi(\tau) \cap \left(\bigcup_{P \in \mathcal{P}_n} \Delta(P) \cup \bigcup_{v \in \mathcal{V}_n} \Delta_{\theta}(v) \right) \neq \emptyset \right\}.$$

It is easy to see from (7.6) that

$$\mathcal{R}'_n = \mathcal{R}_{\text{suc}}(\mathcal{R}'_{n-1}) \setminus \mathcal{U}_n,$$

and so it follows that

$$a_n = \#\mathcal{R}_{\text{suc}}(\mathcal{R}'_{n-1}) - \#\mathcal{U}_n = m^2 a_{n-1} - \#\mathcal{U}_n. \quad (7.13)$$

On the other hand, we have that

$$\begin{aligned} \mathcal{U}_n &= \left\{ \tau' \in \mathcal{R}_{\text{suc}}(\mathcal{R}'_{n-1}) : \Phi(\tau') \cap \left(\bigcup_{P \in \mathcal{P}_{n,1}} \Delta(P) \cup \bigcup_{v \in \mathcal{V}_n} \Delta_{\theta}(v) \right) \neq \emptyset \right\} \\ &\cup \bigcup_{k=2}^n \left\{ \tau' \in \mathcal{R}_{\text{suc}}(\mathcal{R}'_{n-1}) : \Phi(\tau') \cap \bigcup_{P \in \mathcal{P}_{n,k}} \Delta(P) \neq \emptyset \right\} \\ &\subset \bigcup_{\tau \in \mathcal{R}'_{n-1}} \left\{ \tau' \in \mathcal{R}_{\text{suc}}(\tau) : \Phi(\tau') \cap \left(\bigcup_{P \in \mathcal{P}_{n,1}} \Delta(P) \cup \bigcup_{v \in \mathcal{V}_n} \Delta_{\theta}(v) \right) \neq \emptyset \right\} \\ &\cup \bigcup_{k=2}^n \bigcup_{\tau \in \mathcal{R}'_{n-k}} \left\{ \tau' \in \mathcal{R}(\tau)_k : \Phi(\tau') \cap \bigcup_{P \in \mathcal{P}_{n,k}} \Delta(P) \neq \emptyset \right\}. \end{aligned}$$

Thus, Corollary 7.6 implies that

$$\#\mathcal{U}_n \leq 2(3m-2)a_{n-1} + \sum_{k=2}^n (3m-2)^k a_{n-k}. \quad (7.14)$$

On combining (7.13) and (7.14), we obtain that

$$a_n \geq (m^2 - 3m + 2)a_{n-1} - \sum_{k=1}^n (3m-2)^k a_{n-k} = 182a_{n-1} - \sum_{k=1}^n 43^k a_{n-k}. \quad (7.15)$$

With $n = 1$ in (7.15), we find that $a_1 \geq 139$. Hence, (7.12) holds for $n = 1$. Now assume $n \geq 2$ and that (7.12) holds with n replaced by $1, \dots, n-1$. Then for any $1 \leq k \leq n$, we have that

$$a_{n-k} \leq 112^{-k+1} a_{n-1}.$$

Substituting this into (7.15), gives that

$$a_n \geq \left(182 - 112 \sum_{k=1}^n (43/112)^k\right) a_{n-1} > 112 a_{n-1}.$$

This completes the induction step and thus establishes (7.12). In turn this completes the proof of Proposition 7.2. \square

Acknowledgments

This paper was (finally) completed during the authors' stay at the Isaac Newton Institute for Mathematical Sciences at Cambridge during the programme "Interactions between Dynamics of Group Actions and Number Theory" 9 June–4 July 2014. We thank the organisers for inviting us and for creating a wonderful and relaxed working environment. JA would like to thank the great hospitality provided by the University of York where this work was initiated during his stay in June–July of 2012. SV would like to use this opportunity to acknowledge his school friend and fellow mathematician Sunil Talwar (1964–2014) who died very unexpectedly just short of his fiftieth birthday. We had many memorable times together and shared many lifetimes of laughter – even when discussing analysis. Thank you Sunil!

References

- [1] J. An, Badziahin–Pollington–Velani's theorem and Schmidt's game, *Bull. Lond. Math. Soc.* 45 (4) (2013) 721–733.
- [2] J. An, Two-dimensional badly approximable vectors and Schmidt's game, *Duke Math. J.* 165 (2) (2016) 267–284.
- [3] D. Badziahin, A. Pollington, S. Velani, On a problem in simultaneous Diophantine approximation: Schmidt's conjecture, *Ann. of Math.* (2) 174 (3) (2011) 1837–1883.
- [4] D. Badziahin, S. Velani, Badly approximable points on planar curves and a problem of Davenport, *Math. Ann.* 359 (3–4) (2014) 969–1023.
- [5] V. Beresnevich, Badly approximable points on manifolds, *Invent. Math.* 202 (3) (2015) 1199–1240.
- [6] V. Beresnevich, S. Velani, A note on simultaneous Diophantine approximation on planar curves, *Math. Ann.* 337 (2007) 769–796.
- [7] V. Beresnevich, S. Velani, A note on three problems in metric Diophantine approximation, in: *Recent Trends in Ergodic Theory and Dynamical Systems*, Vadodara, 2012, in: *Contemp. Math.*, vol. 631, Amer. Math. Soc., Providence, RI, 2015, pp. 211–229.
- [8] S. Dani, On badly approximable numbers, Schmidt games and bounded orbits of flows, in: *Number Theory and Dynamical Systems*, York, 1987, in: *London Math. Soc. Lecture Note Ser.*, vol. 134, CUP, 1989, pp. 69–86.
- [9] H. Davenport, A note on Diophantine approximation. II, *Mathematika* 11 (1964) 50–58.
- [10] M. Einsiedler, J. Tseng, Badly approximable systems of affine forms, fractals, and Schmidt games, *J. Reine Angew. Math.* 660 (2011) 83–97.
- [11] E. Nesharim, D. Simmons, **Bad**(s, t) is hyperplane absolute winning, *Acta Arith.* 164 (2) (2014) 145–152.
- [12] A. Pollington, S. Velani, On simultaneously badly approximable numbers, *J. Lond. Math. Soc.* (2) 66 (1) (2002) 29–40.
- [13] W.M. Schmidt, Metrical theorems on fractional parts of sequences, *Trans. Amer. Math. Soc.* 110 (1964) 493–518.
- [14] W.M. Schmidt, On badly approximable numbers and certain games, *Trans. Amer. Math. Soc.* 123 (1966) 178–199.