



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/124640/>

Version: Accepted Version

Article:

Bavula, V.V. (2017) The groups of automorphisms of the Lie algebras of polynomial vector fields with zero or constant divergence. *Communications in Algebra*, 45 (3). pp. 1114-1133. ISSN: 0092-7872

<https://doi.org/10.1080/00927872.2016.1175596>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

The groups of automorphisms of the Lie algebras of polynomial vector fields with zero or constant divergence

V. V. Bavula (Aut-Div.tex)

Abstract

Let $P_n = K[x_1, \dots, x_n]$ be a polynomial algebra over a field K of characteristic zero and \mathfrak{div}_n^0 (respectively, \mathfrak{div}_n^c) be the Lie algebra of derivations of P_n with zero (respectively, constant) divergence. We prove that $\text{Aut}_{\text{Lie}}(\mathfrak{div}_n^0) \simeq \text{Aut}_{K\text{-alg}}(P_n)$ ($n \geq 2$) and $\text{Aut}_{\text{Lie}}(\mathfrak{div}_n^c) \simeq \text{Aut}_{K\text{-alg}}(P_n)$. The Lie algebra \mathfrak{div}_n^c is a maximal Lie subalgebra of $\text{Der}_K(P_n)$. Minimal finite sets of generators are found for the Lie algebras \mathfrak{div}_n^0 and \mathfrak{div}_n^c .

Key Words: Group of automorphisms, derivation, the divergence, Lie algebra, automorphism, locally nilpotent derivation, the Lie algebras of polynomial vector fields with zero or constant divergence.

Mathematics subject classification 2010: 17B40, 17B20, 17B66, 17B65, 17B30.

1 Introduction

In this paper, module means a left module, K is a field of characteristic zero and K^* is its group of units, and the following notation is fixed:

- $P_n := K[x_1, \dots, x_n] = \bigoplus_{\alpha \in \mathbb{N}^n} Kx^\alpha$ is a polynomial algebra over K where $x^\alpha := x_1^{\alpha_1} \dots x_n^{\alpha_n}$ and $Q_n := K(x_1, \dots, x_n)$ is its field of fractions,
- $G_n := \text{Aut}_{K\text{-alg}}(P_n)$ is the group of automorphisms of the polynomial algebra P_n ,
- $\partial_1 := \frac{\partial}{\partial x_1}, \dots, \partial_n := \frac{\partial}{\partial x_n}$ are the partial derivatives (K -linear derivations) of P_n ,
- $D_n := \text{Der}_K(P_n) = \bigoplus_{i=1}^n P_n \partial_i$ is the Lie algebra of K -derivations of P_n where $[\partial, \delta] := \partial\delta - \delta\partial$,
- $\mathbb{G}_n := \text{Aut}_{\text{Lie}}(D_n)$ is the group of automorphisms of the Lie algebra D_n ,
- $\delta_1 := \text{ad}(\partial_1), \dots, \delta_n := \text{ad}(\partial_n)$ are the inner derivations of the Lie algebra D_n determined by $\partial_1, \dots, \partial_n$ (where $\text{ad}(a)(b) := [a, b]$),
- $\mathcal{D}_n := \bigoplus_{i=1}^n K\partial_i$,
- $\mathcal{H}_n := \bigoplus_{i=1}^n KH_i$ where $H_1 := x_1\partial_1, \dots, H_n := x_n\partial_n$,
- $D'_n := \bigoplus_{i=1}^n P_n H_i = \bigoplus_{\alpha \in \mathbb{N}^n} x^\alpha \mathcal{H}_n$,
- $\mathfrak{h} := \bigoplus_{i=1}^n Kh_i$ where $h_1 := \partial_1 x_1, \dots, h_n := \partial_n x_n \in \text{End}_K(P_n)$,
- for a derivation $\partial = \sum_{i=1}^n a_i \partial_i \in D_n$, $\text{div}(\partial) := \sum_{i=1}^n \frac{\partial a_i}{\partial x_i}$ is the *divergence* of ∂ ,
- $\mathfrak{div}_n^0 := \{\partial \in D_n \mid \text{div}(\partial) = 0\}$ is the Lie algebra of polynomial vector fields (derivations) with zero divergence,
- $\mathbf{G}_n := \text{Aut}_{\text{Lie}}(\mathfrak{div}_n^0)$,
- $\mathcal{H}'_n := \bigoplus_{i=1}^{n-1} KH_{i,i+1}$ where $H_{ij} := H_i - H_j$ for $i \neq j$,

- $\mathfrak{div}_n^c := \{\partial \in D_n \mid \text{div}(\partial) \in K\}$ is the Lie algebra of polynomial vector fields (derivations) with constant divergence,
- $\mathbf{G}_n^c := \text{Aut}_{\text{Lie}}(\mathfrak{div}_n^c)$,
- $A_n := K\langle x_1, \dots, x_n, \partial_1, \dots, \partial_n \rangle = \bigoplus_{\alpha, \beta \in \mathbb{N}^n} Kx^\alpha \partial^\beta$ is the n 'th Weyl algebra,

The groups of automorphisms of the Lie algebras \mathfrak{div}_n^0 and \mathfrak{div}_n^c . The aim of the paper is to prove the following two theorems.

16Mar13

Theorem 1.1 $\mathbf{G}_n = \begin{cases} G_1/\text{Sh}_1 \simeq K^* & \text{if } n = 1, \\ G_n & \text{if } n \geq 2. \end{cases}$

Structure of the proof. The case $n = 1$ is trivial (see Section 2 where the group Sh_1 is defined in (1)). So, let $n \geq 2$.

- (i) $G_n \subseteq \mathbf{G}_n$ via the group monomorphism (Lemma 2.9.(3))

$$G_n \rightarrow \mathbf{G}_n, \quad \sigma \mapsto \sigma : \partial \mapsto \sigma(\partial) := \sigma \partial \sigma^{-1}.$$

- (ii) Let $\sigma \in \mathbf{G}_n$. Then $\partial'_1 := \sigma(\partial_1), \dots, \partial'_n := \sigma(\partial_n)$ are commuting, locally nilpotent derivations of the polynomial algebra P_n (Lemma 2.14.(1)).

- (iii) $\bigcap_{i=1}^n \ker_{P_n}(\partial'_i) = K$ (Lemma 2.14.(2)).

- (iv) There exists a polynomial automorphism $\tau \in G_n$ such that $\tau\sigma \in \text{Fix}_{\mathbf{G}_n}(\partial_1, \dots, \partial_n)$ (Corollary 2.16).

- (v) $\text{Fix}_{\mathbf{G}_n}(\partial_1, \dots, \partial_n) = \text{Sh}_n$ (Proposition 2.13.(3)) where

Shndef

$$\text{Sh}_n := \{s_\lambda \in G_n \mid s_\lambda(x_1) = x_1 + \lambda_1, \dots, s_\lambda(x_n) = x_n + \lambda_n\} \quad (1)$$

is the *shift group* of automorphisms of the polynomial algebra P_n and $\lambda = (\lambda_1, \dots, \lambda_n) \in K^n$.

- (vi) By (iv) and (v), $\sigma \in G_n$, i.e. $\mathbf{G}_n = G_n$. \square

A16Mar13

Theorem 1.2 $\mathbf{G}_n^c = G_n$.

Structure of the proof. The case $n = 1$ is trivial (see Section 2). So, let $n \geq 2$.

- (i) $G_n \subseteq \mathbf{G}_n^c$ via the group monomorphism (Lemma 2.9.(4))

$$G_n \rightarrow \mathbf{G}_n^c, \quad \sigma \mapsto \sigma : \partial \mapsto \sigma(\partial) := \sigma \partial \sigma^{-1}.$$

- (ii) $\mathfrak{div}_n^0 = [\mathfrak{div}_n^c, \mathfrak{div}_n^c]$ (Lemma 2.10).

- (iii) The short exact sequence of group homomorphisms

$$1 \rightarrow F := \text{Fix}_{\mathbf{G}_n^c}(\mathfrak{div}_n^0) \rightarrow \mathbf{G}_n^c \xrightarrow{\text{res}} \mathbf{G}_n \rightarrow 1$$

is exact (by (i) and Theorem 1.1) where $\text{res} : \sigma \mapsto \sigma|_{\mathfrak{div}_n^0}$ is the restriction map, see (ii).

- (iv) Since $\mathbf{G}_n = G_n$ (Theorem 1.1) and $G_n \subseteq \mathbf{G}_n^c$ (by (i)), the short exact sequence splits

G=GF

$$\mathbf{G}_n^c \simeq G_n \times F. \quad (2)$$

- (v) $F = \{e\}$ (Lemma 2.17). Therefore, $\mathbf{G}_n^c = G_n$. \square

Theorem 1.1 was announced in [8] where a sketch of the proof is given based on a study of certain Lie subalgebras of \mathfrak{div}_n^0 of finite codimension. Our proof is based on completely different ideas. The groups of automorphisms of infinite dimensional Lie algebras were considered in [2]-[10].

A subalgebra \mathcal{M} of a Lie algebra \mathcal{G} is called a *maximal* Lie subalgebra if $\mathcal{M} \neq \mathcal{G}$ and \mathcal{G} is the only Lie subalgebra of \mathcal{G} properly containing \mathcal{M} .

- (Proposition 2.21) For $n \geq 2$, \mathfrak{div}_n^c is a maximal Lie subalgebra of D_n which is also a \mathbb{G}_n -invariant/ G_n -invariant Lie subalgebra.
- (Proposition 2.22) For $n \geq 2$, the G_n -module D_n/\mathfrak{div}_n^c is simple and infinite dimensional with $\text{End}_{G_n}(D_n/\mathfrak{div}_n^c) \simeq K$.

6Oct13

Theorem 1.3 For $n \geq 2$, the set of elements $\partial_1, x_2^2\partial_1, x_3^2\partial_1, \dots, x_n^2\partial_1, x_1^2\partial_2, x_1^2\partial_3, \dots, x_1^2\partial_n$ is a minimal set of generators for the Lie algebra \mathfrak{div}_n^0 .

A6Oct13

Theorem 1.4 For $n \geq 2$, the set of elements in Theorem 1.3 together with H_1 is a minimal set of generators for the Lie algebra \mathfrak{div}_n^c .

2 Proof of Theorems 1.1 and 1.2

P16AAA

In this section, proofs of Theorems 1.1 and 1.2 are given. In the first part of the section some useful results are proved that are used throughout the paper. The second part of the section can be seen as proofs of Theorem 1.1 and 1.2. The proofs are split into several statements that reflect ‘Structure of the proofs of Theorems 1.1 and 1.2’ given in the Introduction. As we have seen in the Introduction, Theorem 1.1 is the key point in the proof of Theorem 1.2.

The Lie algebra D_n is \mathbb{Z}^n -graded. The Lie algebra

xabdb

$$D_n = \bigoplus_{\alpha \in \mathbb{N}^n} \bigoplus_{i=1}^n Kx^\alpha \partial_i \quad (3)$$

is a \mathbb{Z}^n -graded Lie algebra

$$D_n = \bigoplus_{\beta \in \mathbb{Z}^n} D_{n,\beta} \quad \text{where} \quad D_{n,\beta} = \bigoplus_{\alpha - e_i = \beta} Kx^\alpha \partial_i,$$

i.e. $[D_{n,\alpha}, D_{n,\beta}] \subseteq D_{n,\alpha+\beta}$ for all $\alpha, \beta \in \mathbb{N}^n$ where $e_1 := (1, 0, \dots, 0), \dots, e_n := (0, \dots, 0, 1)$ is the canonical free basis for the free abelian group \mathbb{Z}^n . This follows from the commutation relations

xabdb1

$$[x^\alpha \partial_i, x^\beta \partial_j] = \beta_j x^{\alpha+\beta-e_i} \partial_j - \alpha_j x^{\alpha+\beta-e_j} \partial_i. \quad (4)$$

Clearly, for all $i, j = 1, \dots, n$ and $\alpha \in \mathbb{N}^n$,

xabdb2

$$[H_j, x^\alpha \partial_i] = \begin{cases} \alpha_j x^\alpha \partial_i & \text{if } j \neq i, \\ (\alpha_i - 1)x^\alpha \partial_i & \text{if } j = i, \end{cases} \quad (5)$$

xabdb3

$$[\partial_j, x^\alpha \partial_i] = \alpha_j x^{\alpha-e_j} \partial_i. \quad (6)$$

The *support* $\text{Supp}(D_n) := \{\beta \in \mathbb{Z}^n \mid D_{n,\beta} \neq 0\}$ is a submonoid of \mathbb{Z}^n . Let us find the support $\text{Supp}(D_n)$, the graded components $D_{n,\beta}$ and their dimensions $\dim_K D_{n,\beta}$. For each $i = 1, \dots, n$, let $\mathbb{N}^{n,i} := \{\alpha \in \mathbb{N}^n \mid \alpha_i = 0\}$ and $P_n^{\partial_i} := \ker_{P_n}(\partial_i)$. It follows from the decompositions $P_n = P_n^{\partial_i} \oplus P_n x_i$ for $i = 1, \dots, n$ that

Dnb2

$$D_n = \bigoplus_{i=1}^n (P_n^{\partial_i} \oplus P_n x_i) \partial_i = \bigoplus_{i=1}^n P_n^{\partial_i} \partial_i \oplus \bigoplus_{i=1}^n P_n H_i = \bigoplus_{i=1}^n P_n^{\partial_i} \partial_i \oplus \bigoplus_{\alpha \in \mathbb{N}^n} x^\alpha \mathcal{H}_n, \quad (7)$$

Therefore, any derivation $\partial = \sum_{i=1}^n a_i \partial_i \in D_n$ is the unique sum (where $a_i = b_i x_i + c_i$, $b_i \in P_n$ and $c_i \in P_n^{\partial_i}$) dus

$$\partial = \sum_{i=1}^n b_i H_i + \sum_{i=1}^n c_i \partial_i. \quad (8)$$

Hence, Dnb

$$\text{Supp}(D_n) = \prod_{i=1}^n (\mathbb{N}^{n,i} - e_i) \prod \mathbb{N}^n. \quad (9)$$

$$D_{n,\beta} = \begin{cases} Kx^\alpha \partial_i & \text{if } \beta = \alpha - e_i \in \mathbb{N}^{n,i} - e_i, \\ x^\beta \mathcal{H}_n & \text{if } \beta \in \mathbb{N}^n. \end{cases} \quad (10)$$

$$\dim_K D_{n,\beta} = \begin{cases} 1 & \text{if } \beta = \alpha - e_i \in \mathbb{N}^{n,i} - e_i, \\ n & \text{if } \beta \in \mathbb{N}^n. \end{cases}$$

Let \mathcal{G} be an abelian Lie algebra and $\mathcal{G}^* := \text{Hom}_K(\mathcal{G}, K)$. A \mathcal{G} -module M is called a *weight module* if Dnb1

$$M = \bigoplus_{\lambda \in \mathcal{G}^*} M_\lambda \text{ where } M_\lambda := \{m \in M \mid gm = \lambda(g)m \text{ for all } g \in \mathcal{G}\}.$$

The set $W(M) := \{\lambda \in \mathcal{G}^* \mid M_\lambda \neq 0\}$ is called the set of *weights* of M .

The direct sum $\mathfrak{div}_n^0 = \mathfrak{di}_n^0 \oplus \mathfrak{iv}_n^0$. Recall that $D'_n = \bigoplus_{\alpha \in \mathbb{N}^n} x^\alpha \mathcal{H}_n$. By (7), dus2

$$\mathfrak{div}_n^0 = \mathfrak{di}_n^0 \oplus \mathfrak{iv}_n^0 \text{ where } \mathfrak{di}_n^0 := \mathfrak{div}_n^0 \cap D'_n \text{ and } \mathfrak{iv}_n^0 := \bigoplus_{i=1}^n P_n^{\partial_i} \partial_i. \quad (11)$$

We will see that \mathfrak{di}_n^0 is a Lie subalgebra of \mathfrak{div}_n^0 but \mathfrak{iv}_n^0 is not for $n \geq 2$. Clearly, $\mathfrak{di}_1^0 = 0$ and $\mathfrak{div}_1^0 = \mathfrak{iv}_1^0 = K\partial_1$. There are inclusions

$$\begin{aligned} \mathfrak{iv}_1^0 &\subset \mathfrak{iv}_2^0 \subset \cdots \subset \mathfrak{iv}_n^0 \subset \cdots, \\ \mathfrak{di}_1^0 &\subset \mathfrak{di}_2^0 \subset \cdots \subset \mathfrak{di}_n^0 \subset \cdots, \\ \mathfrak{iv}_1^0 &\subset \mathfrak{iv}_2^0 \subset \cdots \subset \mathfrak{iv}_n^0 \subset \cdots. \end{aligned}$$

The K -linear maps $h_i = \partial_1 x_1, \dots, h_n = \partial_n x_n \in \text{End}_K(P_n)$ are bijections since for all $\alpha \in \mathbb{N}^n$ and $i = 1, \dots, n$, dus3

$$h_i(x^\alpha) = (\alpha_i + 1)x^\alpha. \quad (12)$$

The elements h_1, \dots, h_n commute, the polynomial algebra P_n is a weight \mathfrak{h} -module where $\mathfrak{h} := \bigoplus_{i=1}^n K h_i$ is an abelian Lie subalgebra of the Lie algebra $\text{End}_K(P_n)$ (where $[f, g] := fg - gf$) and the set $W(P_n)$ of weights of the \mathfrak{h} -module P_n is equal to $(1, \dots, 1) + \mathbb{N}^n$, i.e. $W(P_n) = \{\lambda = (\lambda_1, \dots, \lambda_n) \mid \lambda \in (1, \dots, 1) + \mathbb{N}^n\}$ where $\lambda(h_i) = \lambda_i$ for all i . For each derivation $\partial = \sum_{i=1}^n a_i H_i \in D'_n$, dus4

$$\text{div}(\partial) = \sum_{i=1}^n h_i(a_i). \quad (13)$$

K -bases for \mathfrak{div}_n^0 and \mathfrak{div}_n^c . For each pair $i \neq j$, the K -linear map dus5

$$\phi_{ij} : P_n \rightarrow \mathfrak{di}_n^0, \quad a \mapsto h_j(a)H_i - h_i(a)H_j, \quad (14)$$

is a (well-defined) injection: By (13), $\text{div}(\phi_{ij}(a)) = (h_i h_j - h_j h_i)(a) = 0$, and if $\phi_{ij}(a) = 0$ then $h_j(a)H_i = h_i(a)H_j$, and so $a = 0$ since the maps h_i and h_j are bijections. For all $\alpha \in \mathbb{N}^n$ and $i \neq j$, let

$$\theta_{ij}^\alpha := \phi_{ij}(x^\alpha) = x^\alpha((\alpha_j + 1)H_i - (\alpha_i + 1)H_j). \quad (15)$$

In particular, $\theta_{ij}^0 = H_i - H_j$. Then

$$[x^{\alpha - \alpha_i e_i} x_j \partial_i, x_i^{\alpha_i + 1} \partial_j] = \phi_{ji}(x^\alpha). \quad (16)$$

It is obvious that $\mathfrak{div}_1^0 = K\partial_1$ and $\mathfrak{div}_1^c = K\partial_1 + KH_1$.

Lemma 2.1 *Let $n \geq 2$. Then*

1. $\mathfrak{di}_n^0 = \bigoplus_{i=1}^{n-1} \phi_{i,i+1}(P_n)$.
2. The set of elements $\theta_i^\alpha := \phi_{i,i+1}(x^\alpha) = x^\alpha((\alpha_{i+1} + 1)H_i - (\alpha_i + 1)H_{i+1})$, where $i = 1, \dots, n-1$ and $\alpha \in \mathbb{N}^n$, is a K -basis for \mathfrak{di}_n^0 .
3. The set of elements θ_i^α in statement 2 and $x^\beta \partial_j$, where $x^\beta \in P_n^{\partial_j}$ and $j = 1, \dots, n$, is a K -basis for \mathfrak{div}_n^0 .
4. The set of elements in statement 3 and H_i , where i is any fixed index in the set $\{1, \dots, n\}$, is a K -basis for \mathfrak{div}_n^c .

Proof. 1. It is obvious that $R := \sum_{i=1}^{n-1} \phi_{i,i+1}(P_n) \subseteq \mathfrak{div}_n^0$, see (14). Recall that $\mathfrak{di}_n^0 = \mathfrak{div}_n^0 \cap D'_n$ and $D'_n = \bigoplus_{\alpha \in \mathbb{N}^n} x^\alpha \mathcal{H}_n$. By (14) and the fact that the K -linear maps h_1, \dots, h_n are invertible,

$$\mathfrak{di}_n^0 = R + \mathfrak{di}_n^0 \cap P_n H_n.$$

By (13), $\mathfrak{di}_n^0 \cap P_n H_n = 0$. Therefore, $\mathfrak{di}_n^0 = R$.

2. Statement 2 follows from statement 1.
3. Statement 3 follows from statement 2 and (11).
4. Statement 4 follows from statement 3 and the fact that $\mathfrak{div}_n^c = \mathfrak{div}_n^0 \oplus KH_i$, $i = 1, \dots, n$. \square

Let $\theta := x_1 \cdots x_n \in P_n$. Then $C_n := \bigoplus_{i \in \mathbb{N}} \theta^i \mathcal{H}'_n$ is an abelian Lie subalgebra of \mathfrak{div}_n^0 that contains \mathcal{H}'_n . We will see that C_n is a Cartan subalgebra of the Lie algebras \mathfrak{div}_n^0 and \mathfrak{div}_n^c (Lemma 2.3.(3,5)).

By Lemma 2.1.(2,3), for $n \geq 2$,

$$C_n = \bigoplus_{i=1}^{n-1} \bigoplus_{m \in \mathbb{N}} K \phi_{i,i+1}(\theta^m), \quad (17)$$

$$\mathfrak{div}_n^0 = \bigoplus_{i=1}^n \bigoplus_{\alpha \in \mathbb{N}^{n,i}} K x^\alpha \partial_i \oplus C_n \oplus \bigoplus_{i=1}^{n-1} \bigoplus_{m \in \mathbb{N}} \bigoplus_{\alpha \in \mathbb{N}_d^n \setminus \{0\}} K \phi_{i,i+1}(\theta^m x^\alpha), \quad (18)$$

where $\mathbb{N}_d^n := \bigcup_{i=1}^n \mathbb{N}^{n,i} = \{(\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n \mid \alpha_i = 0 \text{ for some } i\}$. We identify the vector space $\mathcal{H}'_n = \{\sum_{i=1}^n \lambda_i H_i \mid \sum_{i=1}^n \lambda_i = 0\}$ with its image in K^n under the K -linear injection $\mathcal{H}'_n \rightarrow K^n$, $\sum_{i=1}^n \lambda_i H_i \mapsto (\lambda_1, \dots, \lambda_n)$. So,

$$\mathcal{H}'_n = \{\lambda \in K^n \mid (\lambda, \bar{1}) = \sum_{i=1}^n \lambda_i = 0\}$$

where $\bar{1} := (1, 1, \dots, 1)$ and $(\lambda, \mu) := \sum_{i=1}^n \lambda_i \mu_i$ is the standard inner product on K^n . The dual space $\mathcal{H}'_n^* := \text{Hom}_K(\mathcal{H}'_n, K)$ can be identified with the factor space

$$K^n / K\bar{1} = \{[\mu] := \mu + K\bar{1} \mid \mu \in K^n\},$$

i.e. $[\mu](\sum_{i=1}^n \lambda_i H_i) = [\mu](\lambda) := (\mu, \lambda) = \sum_{i=1}^n \mu_i \lambda_i$. By (18), the \mathcal{H}'_n -module \mathfrak{div}_n^0 is a weight module and the summands in (18) are the (nonzero) weight vectors under the adjoint action of \mathcal{H}'_n on \mathfrak{div}_n^0 ,

$$\mathfrak{div}_n^0 = \bigoplus_{[\mu] \in W(\mathfrak{div}_n^0)} \mathfrak{div}_{n, [\mu]}^0$$

where $\mathfrak{div}_{n, [\mu]}^0 := \{\partial \in \mathfrak{div}_n^0 \mid [H, \partial] = [\mu](H)\partial \text{ for all } H \in \mathcal{H}'_n\}$ is the weight subspace of \mathfrak{div}_n^0 that corresponds to the weight $[\mu]$ and $W = W(\mathfrak{div}_n^0)$ is the set of weights for \mathfrak{div}_n^0 . To simplify the notation we identify the set \mathbb{N}_d^n with its isomorphic copy in the factor vector space $K^n/K\bar{1}$ via the map $K^n \rightarrow K^n/K\bar{1}$, $\lambda \mapsto K\bar{1}$. So,

WDnb

$$W(\mathfrak{div}_n^0) = \mathbb{N}_d^n. \quad (19)$$

Dnb1

$$\mathfrak{div}_{n, [\mu]}^0 = \begin{cases} Kx^\alpha \partial_i \oplus \bigoplus_{i=1}^{n-1} \bigoplus_{m \in \mathbb{N}} \phi_{i, i+1}(\theta^m x^{\alpha - e_i + \bar{1}}) & \text{if } [\mu] = [\alpha - e_i], \alpha \in \mathbb{N}^{n, i}, \\ C_n & \text{if } [\mu] = 0, \\ \bigoplus_{i=1}^{n-1} \bigoplus_{m \in \mathbb{N}} \phi_{i, i+1}(\theta^m x^\alpha) & \text{otherwise.} \end{cases} \quad (20)$$

$$\dim_K \mathfrak{div}_{n, [\mu]}^0 = \infty \text{ for all } [\mu] \in W(\mathfrak{div}_n^0).$$

The Lie algebra $\mathfrak{div}_n^0 = \mathfrak{iv}_n^0 \oplus \mathfrak{di}_n^0$ is the direct sum of its weight \mathcal{H}'_n -submodules with

Wiv

$$W(\mathfrak{iv}_n^0) = \prod_{i=1}^n (\mathbb{N}^{n, i} - e_i), \quad W(\mathfrak{di}_n^0) = \mathbb{N}_d^n. \quad (21)$$

For $H = \sum_{i=1}^n \lambda_i H_i \in \mathcal{H}_n$ and $\alpha \in K^n$, let

$$(H, \alpha) = (\alpha, H) = \sum_{i=1}^n \alpha_i \lambda_i.$$

Then, for all $\alpha \in \mathbb{N}^n$, $\text{div}(x^\alpha H) = (\alpha + \bar{1}, H)$. If, in addition, $H \in \mathcal{H}'_n$, that is $(H, \bar{1}) = 0$, then, for all $\alpha \in \mathbb{N}^n$, $\text{div}(x^\alpha H) = (\alpha + \bar{1}, H) = (\alpha, H)$. It follows that \mathfrak{di}_n^0 is the direct sum of vector spaces

Wiv1

$$\mathfrak{di}_n^0 = \bigoplus_{\alpha \in \mathbb{N}^n} \{Kx^\alpha H \mid (H, \alpha + \bar{1}) = 0, H \in \mathcal{H}_n\}. \quad (22)$$

Let

Wiv2

$$\mathfrak{di}_n^{\prime 0} := \bigoplus_{\alpha \in \mathbb{N}^n} \{Kx^\alpha H \mid (H, \alpha) = 0, H \in \mathcal{H}'_n\} = \bigoplus_{\alpha \in \mathbb{N}_d^n} \{K[\theta]x^\alpha H \mid (H, \alpha) = 0, H \in \mathcal{H}'_n\}. \quad (23)$$

Clearly, for $n \geq 2$, $\mathfrak{di}_n^{\prime 0} \subset \mathfrak{di}_n^0$ and the vector space $\mathfrak{di}_n^{\prime 0}$ is a left $K[\theta]$ -module. We will see shortly that $\mathfrak{di}_n^{\prime 0}$ is a non-Noetherian Lie algebra (Lemma 2.2). Notice that $\mathfrak{di}_n^{\prime 0} = C_2 = \bigoplus_{i \geq 0} K\theta^i (H_1 - H_2) = K[\theta](H_1 - H_2)$ where $\theta = x_1 x_2$.

The commutation relations of the weight vectors in \mathfrak{div}_n^0 . By (18) and (19), there are three types of commutation relations of elements from the weight spaces of the Lie algebra \mathfrak{div}_n^0 , see (24), (25) and (26). For all $x^\alpha \partial_i \in P_n^{\partial_i} \partial_i$ and $x^\beta \partial_j \in P_n^{\partial_j} \partial_j$,

xadxbd

$$[x^\alpha \partial_i, x^\beta \partial_j] = \begin{cases} \phi_{ji}(x^{\alpha+\beta-e_i-e_j}) & \text{if } \beta_i \neq 0, \alpha_j \neq 0 \\ \beta_i x^{\alpha+\beta-e_i} \partial_j & \text{if } \beta_i \neq 0, \alpha_j = 0 \\ -\alpha_j x^{\alpha+\beta-e_j} \partial_i & \text{if } \beta_i = 0, \alpha_j \neq 0 \\ 0 & \text{if } \beta_i = 0, \alpha_j = 0 \end{cases} \quad (24)$$

where $\phi_{ji}(x^{\alpha+\beta-e_i-e_j}) = x^{\alpha+\beta-e_i-e_j}(\beta_i H_j - \alpha_j H_i)$ (since $\alpha_i = 0$ and $\beta_j = 0$).

For all elements $x^\alpha H, x^{\alpha'} H' \in \mathfrak{di}_n^0$,

xadxbd1

$$[x^\alpha H, x^{\alpha'} H'] = x^{\alpha+\alpha'} ((H, \alpha')H' - (H', \alpha)H) \in \mathfrak{di}_n^0 \quad (25)$$

since $((H, \alpha')H' - (H', \alpha)H, \alpha + \alpha' + \bar{1}) = (H, \alpha')(H', \alpha' + \bar{1}) - (H', \alpha)(H, \alpha + \bar{1}) = 0$.

For all $x^\beta \in P_n^{\partial_i}$ and $x^\alpha H \in \mathfrak{di}_n^0$,

xadxbd2

$$[x^\beta \partial_i, x^\alpha H] = \alpha_i x^{\alpha+\beta-e_i} H - (H, \beta - e_i) x^{\alpha+\beta} \partial_i. \quad (26)$$

If, in addition $\alpha_i \neq 0$, then the equality (26) takes the form

xadxbd3

$$[x^\beta \partial_i, x^\alpha H] = x^{\alpha+\beta-e_i} (\alpha_i H - (H, \beta - e_i) H_i) \in \mathfrak{di}_n^0 \quad (27)$$

since $(\alpha_i H - (H, \beta - e_i) H_i, \alpha + \beta - e_i + \bar{1}) = -(H, \beta - e_i)(H_i, \beta - e_i + \bar{1}) = -(H, \beta - e_i) \cdot 0 = 0$. By (22) and (25), \mathfrak{di}_n^0 is a *Lie subalgebra* of \mathfrak{div}_n^0 which is not an ideal, by (26). By (24), \mathfrak{iv}_n^0 is not a Lie algebra for $n \geq 2$.

The Lie algebra \mathfrak{di}_n^0 is not Noetherian, $n \geq 2$. Let $n \geq 2$. The Lie algebra \mathfrak{di}_n^0 is a $K[\theta]$ -module, $K[\theta]\mathfrak{di}_n^0 \subseteq \mathfrak{di}_n^0$, and the Lie bracket on \mathfrak{di}_n^0 is a $k[\theta]$ -bilinear: for all $p, p' \in K[\theta]$ and $x^\alpha H, x^{\alpha'} H' \in \mathfrak{di}_n^0$,

ppHa

$$[px^\alpha H, p'x^{\alpha'} H'] = pp'[x^\alpha H, x^{\alpha'} H']. \quad (28)$$

So, the Lie algebra \mathfrak{di}_n^0 is not simple: $\theta^i \mathfrak{di}_n^0$, $i \in \mathbb{N}$ are distinct ideals of \mathfrak{di}_n^0 .

a3Apr13

Lemma 2.2 For $n \geq 2$, the Lie algebras \mathfrak{di}_n^0 and \mathfrak{di}_n^0 are not Noetherian.

Proof. If $n = 2$ then $\mathfrak{di}_2^0 = C_2$ is an infinite dimensional abelian Lie algebra, hence, non-Noetherian. Let $n \geq 3$. Let I be an ideal of the additive monoid \mathbb{N}^n ($I + \mathbb{N}^n \subseteq I$). Then $\mathfrak{a}'(I) := \bigoplus_{\alpha \in I} \{x^\alpha H \mid (\alpha, H) = 0, H \in \mathcal{H}'_n\}$ is an ideal of the Lie algebra \mathfrak{di}_n^0 , by (25). The map $I \mapsto \mathfrak{a}'(I)$ from the set $\mathcal{I}(\mathbb{N}^n)$ of ideals of \mathbb{N}^n to the set $\mathcal{I}(\mathfrak{di}_n^0)$ of ideals of the Lie algebra \mathfrak{di}_n^0 is an inclusion preserving injection ($I_1 \subsetneq I_2$ implies $\mathfrak{a}'(I_1) \subsetneq \mathfrak{a}'(I_2)$). The set $\mathcal{I}(\mathbb{N}^n)$ is not Noetherian (with respect to \subseteq) hence \mathfrak{di}_n^0 is not a Noetherian Lie algebra.

Similarly, by (25), for $n \geq 2$, the map $I \mapsto \mathfrak{a}(I) := \bigoplus_{\alpha \in I} \{x^\alpha H \mid (\alpha + \bar{1}, H) = 0, H \in \mathcal{H}_n\}$ from the set $\mathcal{I}(\mathbb{N}^n)$ to the set $\mathcal{I}(\mathfrak{di}_n^0)$ of ideals of the Lie algebra \mathfrak{di}_n^0 is an inclusion preserving injection. Therefore, the Lie algebra \mathfrak{di}_n^0 is non-Noetherian. \square

Let \mathcal{G} be a Lie algebra and \mathcal{H} be its Lie subalgebra. The *centralizer* $C_{\mathcal{G}}(\mathcal{H}) := \{x \in \mathcal{G} \mid [x, \mathcal{H}] = 0\}$ of \mathcal{H} in \mathcal{G} is a Lie subalgebra of \mathcal{G} . In particular, $Z(\mathcal{G}) := C_{\mathcal{G}}(\mathcal{G})$ is the *centre* of the Lie algebra \mathcal{G} . The *normalizer* $N_{\mathcal{G}}(\mathcal{H}) := \{x \in \mathcal{G} \mid [x, \mathcal{H}] \subseteq \mathcal{H}\}$ of \mathcal{H} in \mathcal{G} is a Lie subalgebra of \mathcal{G} , it is the largest Lie subalgebra of \mathcal{G} that contains \mathcal{H} as an ideal.

Let V be a vector space over K . A K -linear map $\delta : V \rightarrow V$ is called a *locally nilpotent map* if $V = \bigcup_{i \geq 1} \ker(\delta^i)$ or, equivalently, for every $v \in V$, $\delta^i(v) = 0$ for all $i \gg 1$. When δ is a locally nilpotent map in V we also say that δ *acts locally nilpotently* on V . Every *nilpotent* linear map δ , that is $\delta^n = 0$ for some $n \geq 1$, is a locally nilpotent map but not vice versa, in general. Let \mathcal{G} be a Lie algebra. Each element $a \in \mathcal{G}$ determines the derivation of the Lie algebra \mathcal{G} by the rule $\text{ad}(a) : \mathcal{G} \rightarrow \mathcal{G}$, $b \mapsto [a, b]$, which is called the *inner derivation* associated with a . The set $\text{Inn}(\mathcal{G})$ of all the inner derivations of the Lie algebra \mathcal{G} is a Lie subalgebra of the Lie algebra $(\text{End}_K(\mathcal{G}), [\cdot, \cdot])$ where $[f, g] := fg - gf$. There is the short exact sequence of Lie algebras

$$0 \rightarrow Z(\mathcal{G}) \rightarrow \mathcal{G} \xrightarrow{\text{ad}} \text{Inn}(\mathcal{G}) \rightarrow 0,$$

that is $\text{Inn}(\mathcal{G}) \simeq \mathcal{G}/Z(\mathcal{G})$ where $Z(\mathcal{G})$ is the *centre* of the Lie algebra \mathcal{G} and $\text{ad}([a, b]) = [\text{ad}(a), \text{ad}(b)]$ for all elements $a, b \in \mathcal{G}$. An element $a \in \mathcal{G}$ is called a *locally nilpotent element* (respectively, a *nilpotent element*) if so is the inner derivation $\text{ad}(a)$ of the Lie algebra \mathcal{G} .

The Cartan subalgebra C_n of \mathfrak{div}_n^0 . A nilpotent Lie subalgebra C of a Lie algebra \mathcal{G} such that $C = N_{\mathcal{G}}(C)$ is called a *Cartan subalgebra* of \mathcal{G} . We use often the following obvious observation: *An abelian Lie subalgebra that coincides with its centralizer is a maximal abelian Lie subalgebra.*

Example. \mathcal{H}_n is a Cartan subalgebra of D_n and $\mathcal{H}_n = C_{D_n}(\mathcal{H}_n)$ is a maximal abelian Lie subalgebra of D_n .

Lemma 2.3 *Let $n \geq 2$. Recall that $C_n = \bigoplus_{i \in \mathbb{N}} \theta^i \mathcal{H}'_n$ and $\theta = x_1 x_2 \cdots x_n$. Then*

1. $\mathcal{H}'_n = \mathcal{H}_n \cap \mathfrak{div}_n^0$.
2. $C_n = C_{\mathfrak{div}_n^0}(\mathcal{H}'_n)$ and $C_n = C_{\mathfrak{div}_n^0}(C_n)$ is a maximal abelian Lie subalgebra of \mathfrak{div}_n^0 .
3. C_n is a Cartan subalgebra of \mathfrak{div}_n^0 .
4. $C_n + \mathcal{H}_n = C_{\mathfrak{div}_n^c}(\mathcal{H}'_n)$ and $C_n = C_{\mathfrak{div}_n^c}(C_n)$ is a maximal abelian Lie subalgebra of \mathfrak{div}_n^c .
5. $N_{\mathfrak{div}_n^c}(C_n) = C_n + \mathcal{H}_n = N_{\mathfrak{div}_n^c}(C_n + \mathcal{H}_n)$ is a solvable but not nilpotent Lie algebra, and so $C_n + \mathcal{H}_n$ and C_n are not a Cartan subalgebra of \mathfrak{div}_n^c .
6. \mathcal{H}_n is a Cartan subalgebra of \mathfrak{div}_n^c and $\mathcal{H}_n = C_{\mathfrak{div}_n^c}(\mathcal{H}_n)$ is a maximal abelian Lie subalgebra of \mathfrak{div}_n^c .

Proof. 1. Let $H = \sum_{i=1}^n \lambda_i H_i \in \mathcal{H}_n$. Then $H \in \mathcal{H}_n \cap \mathfrak{div}_n^0$ iff $\sum_{i=1}^n \lambda_i = 0$ iff $H \in \mathcal{H}'_n$.

2. The fact that $C_n = C_{\mathfrak{div}_n^0}(\mathcal{H}'_n)$ follows from (20). By (25), C_n is an abelian Lie subalgebra of \mathfrak{div}_n^0 . Then the inclusion $\mathcal{H}'_n \subseteq C_n = C_{\mathfrak{div}_n^0}(\mathcal{H}'_n)$ implies the inclusions

$$C_n = C_{\mathfrak{div}_n^0}(\mathcal{H}'_n) \supseteq C_{\mathfrak{div}_n^0}(C_n) \supseteq C_n.$$

Therefore, $C_n = C_{\mathfrak{div}_n^0}(C_n)$ is a maximal abelian Lie subalgebra of \mathfrak{div}_n^0 .

3. Let $N := N_{\mathfrak{div}_n^0}(C_n)$. By statement 2, $C_n \subseteq N$. The \mathcal{H}'_n -module \mathfrak{div}_n^0 is weight and $C_n = C_{\mathfrak{div}_n^0}(\mathcal{H}'_n)$ is the zero weight component of \mathfrak{div}_n^0 . The inclusion $[N, \mathcal{H}'_n] \subseteq C_n$ implies $N \subseteq C_n$, and so $N = C_n$.

4. Notice that $\mathfrak{div}_n^c = \mathfrak{div}_n^0 \oplus KH_1$, $H_1 \in C_{\mathfrak{div}_n^c}(\mathcal{H}'_n)$ and $\mathcal{H}'_n \subseteq \mathfrak{div}_n^0$. By statement 2,

$$C_{\mathfrak{div}_n^c}(\mathcal{H}'_n) = C_{\mathfrak{div}_n^0}(\mathcal{H}'_n) \oplus KH_1 = C_n \oplus KH_1 = C_n + \mathcal{H}_n.$$

Now, the inclusion $\mathcal{H}'_n \subseteq C_n$ implies the inclusions

$$C_n \oplus KH_1 = C_{\mathfrak{div}_n^c}(\mathcal{H}'_n) \supseteq C_{\mathfrak{div}_n^c}(C_n) \supseteq C_n.$$

Hence $C_{\mathfrak{div}_n^c}(C_n) = C_n \oplus C_{\mathfrak{div}_n^c}(C_n) \cap KH_1 = C_n$ and so C_n is a maximal abelian Lie subalgebra of \mathfrak{div}_n^c .

5. Let $N^c := N_{\mathfrak{div}_n^c}(C_n)$. By statement 4, $C_n \subseteq N^c$. The \mathcal{H}'_n -module $\mathfrak{div}_n^c = \mathfrak{div}_n^0 \oplus KH_1$ is weight and $C_n \oplus KH_1 = C_{\mathfrak{div}_n^c}(\mathcal{H}'_n)$ is the zero weight component of \mathfrak{div}_n^c . The inclusion $[N^c, \mathcal{H}'_n] \subseteq C_n$ implies the inclusion $[N^c, \mathcal{H}'_n] \subseteq C_n \oplus KH_1 = C_{\mathfrak{div}_n^c}(\mathcal{H}'_n)$, and so $N^c \subseteq C_n \oplus KH_1$. Since $C_n \subseteq N^c$, it follows that $N^c = C_n \oplus N^c \cap KH_1 = C_n + KH_1$, i.e. C_n is not a Cartan subalgebra of \mathfrak{div}_n^c .

Clearly, $C_n + \mathcal{H}_n \subseteq \mathcal{N} := N_{\mathfrak{div}_n^c}(C_n + \mathcal{H}_n)$. Since $\mathcal{H}'_n \subseteq C_n + \mathcal{H}_n$, $[\mathcal{H}'_n, \mathcal{N}] \subseteq C_n + \mathcal{H}_n = C_{\mathfrak{div}_n^c}(\mathcal{H}'_n)$, hence $\mathcal{N} \subseteq C_n + \mathcal{H}_n$ (as $C_n + \mathcal{H}_n$ is the zero weight component of the weight \mathcal{H}'_n -module \mathfrak{div}_n^c), i.e. $\mathcal{N} = C_n + \mathcal{H}_n$. The Lie algebra $C_n + \mathcal{H}_n$ is solvable but not nilpotent, and so $C_n + \mathcal{H}_n$ is not a Cartan subalgebra of \mathfrak{div}_n^c .

6. Statement 6 follows from the Example above. \square

The Lie algebra \mathfrak{div}_n^0 is a weight \mathcal{H}'_n -module with respect to the adjoint action. In particular, the action of \mathcal{H}'_n on \mathfrak{div}_n^0 is *locally finite dimensional*, i.e. for any $\partial \in \mathfrak{div}_n^0$, $\dim_K(\sum_{i \in \mathbb{N}} \text{ad}(\mathcal{H}'_n)^i(\partial)) < \infty$. We can easily verify that the action of the Cartan subalgebra C_n of \mathfrak{div}_n^0 on \mathfrak{div}_n^0 is not locally finite dimensional, see (25) and (26).

P_n is a D_n -module. The polynomial algebra P_n is a (left) D_n -module: $D_n \times P_n \rightarrow P_n$, $(\partial, p) \mapsto \partial * p$. In more detail, if $\partial = \sum_{i=1}^n a_i \partial_i$ where $a_i \in P_n$ then

$$\partial * p = \sum_{i=1}^n a_i \frac{\partial p}{\partial x_i}.$$

The field K is a D_n -submodule of P_n and

IkerdiK

$$P_n^{D_n} := \bigcap_{i=1}^n \ker_{P_n}(\partial_i) = K. \quad (29)$$

Lemma 2.4 [2] *The D_n -module P_n/K is simple with $\text{End}_{D_n}(P_n/K) = \text{Kid}$ where id is the identity map.*

xa11Mar13

The G_n -module D_n . The Lie algebra D_n is a G_n -module,

$$G_n \times D_n \rightarrow D_n, \quad (\sigma, \partial) \mapsto \sigma(\partial) := \sigma \partial \sigma^{-1}.$$

Every automorphism $\sigma \in G_n$ is uniquely determined by the elements

$$x'_1 := \sigma(x_1), \dots, x'_n := \sigma(x_n).$$

Let $M_n(P_n)$ be the algebra of $n \times n$ matrices over P_n . The matrix $J(\sigma) := (J(\sigma)_{ij}) \in M_n(P_n)$, where $J(\sigma)_{ij} = \frac{\partial x'_j}{\partial x_i}$, is called the *Jacobian matrix* of the automorphism (endomorphism) σ and its determinant $\mathcal{J}(\sigma) := \det J(\sigma)$ is called the *Jacobian* of σ . So, the j 'th column of $J(\sigma)$ is the *gradient* $\text{grad } x'_j := \left(\frac{\partial x'_j}{\partial x_1}, \dots, \frac{\partial x'_j}{\partial x_n} \right)^T$ of the polynomial x'_j where T stands for the transposition. Then the derivations

$$\partial'_1 := \sigma \partial_1 \sigma^{-1}, \dots, \partial'_n := \sigma \partial_n \sigma^{-1}$$

are the partial derivatives of P_n with respect to the variables x'_1, \dots, x'_n ,

ddp=dx_i

$$\partial'_1 = \frac{\partial}{\partial x'_1}, \dots, \partial'_n = \frac{\partial}{\partial x'_n}. \quad (30)$$

Every derivation $\partial \in D_n$ is a unique sum $\partial = \sum_{i=1}^n a_i \partial_i$ where $a_i = \partial * x_i \in P_n$. Let $\partial := (\partial_1, \dots, \partial_n)^T$ and $\partial' := (\partial'_1, \dots, \partial'_n)^T$. Then

dp=Jnd

$$\partial' = J(\sigma)^{-1} \partial, \quad \text{i.e. } \partial'_i = \sum_{j=1}^n (J(\sigma)^{-1})_{ij} \partial_j \quad \text{for } i = 1, \dots, n. \quad (31)$$

In more detail, if $\partial' = A \partial$ where $A = (a_{ij}) \in M_n(P_n)$, i.e. $\partial'_i = \sum_{j=1}^n a_{ij} \partial_j$. Then for all $i, j = 1, \dots, n$,

$$\delta_{ij} = \partial'_i * x'_j = \sum_{k=1}^n a_{ik} \frac{\partial x'_j}{\partial x_k}$$

where δ_{ij} is the Kronecker delta function. The equalities above can be written in the matrix form as $AJ(\sigma) = 1$ where 1 is the identity matrix. Therefore, $A = J(\sigma)^{-1}$.

For all $\sigma, \tau \in G_n$,

Jst=JsJ

$$J(\sigma\tau) = J(\sigma) \cdot \sigma(J(\tau)). \quad (32)$$

By taking the determinants of both sides of (32), we have a similar equality of the Jacobians: for all $\sigma, \tau \in G_n$.

Jst=JsJ1

$$\mathcal{J}(\sigma\tau) = \mathcal{J}(\sigma) \cdot \sigma(\mathcal{J}(\tau)). \quad (33)$$

Properties of the divergence. Recall some of the properties of the divergence map

$$\operatorname{div} : D_n \rightarrow P_n, \quad \partial = \sum_{i=1}^n a_i \partial_i \mapsto \operatorname{div}(\partial) = \sum_{i=1}^n \frac{\partial a_i}{\partial x_i}.$$

(div-i) div is a K -linear map which is a surjection.

(div-ii) For all $a \in P_n$ and $\partial \in D_n$, $\operatorname{div}(a\partial) = a\operatorname{div}(\partial) + \partial(a)$.

(div-iii) For all $\partial, \delta \in D_n$, $\operatorname{div}([\partial, \delta]) = \partial(\operatorname{div}(\delta)) - \delta(\operatorname{div}(\partial))$.

(div-iv) Let $a_1, \dots, a_n \in P_n$; $\sigma : P_n \rightarrow P_n$, $x_i \mapsto a_i$, $i = 1, \dots, n$ and $\mathcal{J}(a_1, \dots, a_n) := \mathcal{J}(\sigma)$ be the Jacobian of σ . Then (Proposition 2.3.2, [6])

$$\partial * \mathcal{J}(a_1, \dots, a_n) = -\mathcal{J}(a_1, \dots, a_n)\operatorname{div}(\partial) + \sum_{i=1}^n \mathcal{J}(a_1, \dots, \partial * a_i, \dots, a_n).$$

(div-v) (Theorem 2.5.5, [6]) If $D_n = \bigoplus_{i=1}^n P_n \partial'_i$ and $\partial'_1, \dots, \partial'_n$ commute then

$$\operatorname{div}(\partial'_1) = \dots = \operatorname{div}(\partial'_n) = 0.$$

(div-vi) Let $\sigma : P_n \rightarrow P_n$, $x_i \mapsto x'_i$, $i = 1, \dots, n$, be an automorphism of P_n . Then $\partial'_1 := \sigma \partial_1 \sigma^{-1}, \dots, \partial'_n := \sigma \partial_n \sigma^{-1}$ are commuting derivations of P_n such that $D_n = \bigoplus_{i=1}^n P_n \partial'_i$ (by (31) and since $J(\sigma)^{-1} \in \operatorname{GL}_n(P_n)$). By (31), $\partial_i = \sum_{j=1}^n (J(\sigma)^{-1})_{ij} \partial_j$. By (div-v),

$$\sum_{j=1}^n \partial_j * (J(\sigma)^{-1})_{ij} = 0 \quad \text{for } i = 1, \dots, n.$$

The divergence commutes with polynomial automorphisms. The following known theorem shows that the divergence commutes with automorphisms, i.e. the divergence map $\operatorname{div} : D_n \rightarrow P_n$ is a G_n -module homomorphism. We give a short proof.

B16Mar13

Theorem 2.5 For all $\sigma \in G_n$ and $\partial \in D_n$,

$$\operatorname{div}(\sigma(\partial)) = \sigma(\operatorname{div}(\partial)).$$

Proof. Let $\partial = \sum_{i=1}^n a_i \partial_i$ where $a_i \in P_n$. Then $\partial' = \sigma \partial \sigma^{-1} = \sum_{i=1}^n \sigma(a_i) \partial'_i$ where, by (31), $\partial'_i = \sum_{j=1}^n (J(\sigma)^{-1})_{ij} \partial_j$. Now, by (div-vi),

$$\begin{aligned} \operatorname{div}(\partial') &= \sum_{i,j=1}^n \partial_j * ((J(\sigma)^{-1})_{ij} \sigma(a_i)) = \sum_{i=1}^n \left(\sum_{j=1}^n \partial_j * (J(\sigma)^{-1})_{ij} \right) \cdot \sigma(a_i) + \sum_{i=1}^n \sum_{j=1}^n (J(\sigma)^{-1})_{ij} \partial_j * \sigma(a_i) \\ &= \sum_{i=1}^n \partial'_i * \sigma(a_i) = \sum_{i=1}^n \sigma \partial_i \sigma^{-1} \sigma(a_i) = \sigma \left(\sum_{i=1}^n \partial_i(a_i) \right) = \sigma(\operatorname{div}(\partial)). \quad \square \end{aligned}$$

11Mar13

Theorem 2.6 $\mathbb{G}_n = G_n$.

The above theorem was announced in [8] where a sketch of a proof is given, it can also be deduced from [10]. A proof of the above theorem is given in [9] and in ([7], Theorem 3.6), a different approach and a short proof is given in [2]. Some generalizations are given in [4], [3] and [5].

a20Mar13

Corollary 2.7 For all $\sigma \in \mathbb{G}_n$ and $\partial \in D_n$, $\operatorname{div}(\sigma(\partial)) = \sigma(\operatorname{div}(\partial))$.

Proof. The statement follows from Theorem 2.6 and Theorem 2.5. \square

By (div-iii), \mathfrak{div}_n^0 and \mathfrak{div}_n^c are Lie subalgebras of D_n , \mathfrak{div}_n^0 is an ideal of \mathfrak{div}_n^c .

aB16Mar13

Corollary 2.8 *The Lie subalgebras \mathfrak{div}_n^0 and \mathfrak{div}_n^c of D_n are also G_n -submodules of D_n .*

Proof. This follows from Theorem 2.5. \square

By (div-iii),

$$0 \rightarrow \mathfrak{div}_n^0 \rightarrow D_n \xrightarrow{\text{div}} P_n \rightarrow 0 \quad (34)$$

divDnPn

is the short exact sequence of (left) \mathfrak{div}_n^0 -modules and \mathfrak{div}_n^c -modules, i.e.

divcom

$$\text{div}([\partial, \delta]) = \partial * \text{div}(\delta) \quad (35)$$

for all $\partial \in \mathfrak{div}_n^c$ and $\delta \in D_n$. So,

divDnPn1

$$0 \rightarrow \mathfrak{div}_n^0 \rightarrow \mathfrak{div}_n^c \xrightarrow{\text{div}} K \rightarrow 0 \quad (36)$$

is the short exact sequence of (left) \mathfrak{div}_n^0 -modules/ \mathfrak{div}_n^c -modules, and so

divDnPn2

$$\mathfrak{div}_n^c = \mathfrak{div}_n^0 \oplus KH_i \text{ for } i = 1, \dots, n, \quad (37)$$

since $\text{div}(H_i) = 1$ for all $i = 1, \dots, n$.

The maximal abelian Lie subalgebra \mathcal{D}_n of \mathfrak{div}_n^0 and \mathfrak{div}_n^c .

db11Mar13

Lemma 2.9 1. $C_{\mathfrak{div}_n^0}(\mathcal{D}_n) = C_{\mathfrak{div}_n^c}(\mathcal{D}_n) = \mathcal{D}_n$ and so \mathcal{D}_n is a maximal abelian Lie subalgebra of \mathfrak{div}_n^0 and \mathfrak{div}_n^c .

2. ([2]) $\text{Fix}_{G_n}(\mathcal{D}_n) = \text{Sh}_n$.

3. For $n \geq 2$, \mathfrak{div}_n^0 is a faithful G_n -module, i.e. the group homomorphism $G_n \rightarrow \mathbf{G}_n$, $\sigma \mapsto \sigma : \partial \mapsto \sigma \partial \sigma^{-1}$, is a monomorphism. For $n = 1$, the group Sh_1 is the kernel of the group homomorphism $G_1 \rightarrow \mathbf{G}_1$.

4. \mathfrak{div}_n^c is a faithful G_n -module, i.e. the group homomorphism $G_n \rightarrow \mathbf{G}_n^c$, $\sigma \mapsto \sigma : \partial \mapsto \sigma \partial \sigma^{-1}$, is a monomorphism.

5. $\text{Fix}_{G_n}(\mathcal{D}_n \oplus \mathcal{H}'_n) = \{e\}$ for $n \geq 2$.

Proof. 1. Since $C_{D_n}(\mathcal{D}_n) = \mathcal{D}_n$, [2], we must have $C_{\mathfrak{div}_n^0}(\mathcal{D}_n) = C_{\mathfrak{div}_n^c}(\mathcal{D}_n) = \mathcal{D}_n$, and so \mathcal{D}_n is a maximal abelian Lie subalgebra of \mathfrak{div}_n^0 and \mathfrak{div}_n^c .

3 and 5. By Corollary 2.8, \mathfrak{div}_n^0 is a G_n -submodule of \mathcal{D}_n . So, the group homomorphism in statement 3 is well-defined. The case $n = 1$ is obvious since $G_1 = \{x \mapsto \lambda x + \mu \mid \lambda \in K^*, \mu \in K\}$ and $\mathfrak{div}_1^0 = K\partial$.

So, let $n \geq 2$, and $\sigma \in \text{Fix}_{G_n}(\mathcal{D}_n \oplus \mathcal{H}'_n)$. Then $\sigma \in \text{Fix}_{G_n}(\mathcal{D}_n) = \text{Sh}_n$, by statement 2. So, $\sigma(x_1) = x_1 + \lambda_1, \dots, \sigma(x_n) = x_n + \lambda_n$ where $\lambda_i \in K$. Then for all $i \neq j$,

$$H_i - H_j = \sigma(H_i - H_j) = (x_i + \lambda_i)\partial_i - (x_j + \lambda_j)\partial_j = H_i - H_j + \lambda_i\partial_i - \lambda_j\partial_j.$$

So, $\lambda_1 = \dots = \lambda_n = 0$. This means that $\sigma = e$. So, $\text{Fix}_{G_n}(\mathcal{D}_n \oplus \mathcal{H}'_n) = \{e\}$ and \mathfrak{div}_n^0 is a faithful G_n -module.

4. By Corollary 2.8, \mathfrak{div}_n^c is a G_n -submodule of D_n . So, the group homomorphism in statement 4 is well-defined. Now, statement 4 follows from statement 3 for $n \geq 2$. For $n = 1$, statement 4 is obvious as $\mathfrak{div}_1^c = K\partial_1 + KH_1$ and $G_1 = \{x \mapsto \lambda x + \mu \mid \lambda \in K^*, \mu \in K\}$. \square

Lemma 2.10 $[\mathfrak{div}_n^c, \mathfrak{div}_n^c] = \mathfrak{div}_n^0$.

a16Mar13

Proof. The statement is obvious for $n = 1$ as $\mathfrak{div}_1^c = K\partial_1 + KH_1$ and $\mathfrak{div}_1^0 = K\partial_1$. So, let $n \geq 2$. By (div-iii), $\text{div}([\mathfrak{div}_n^c, \mathfrak{div}_n^c]) = 0$, and so $[\mathfrak{div}_n^c, \mathfrak{div}_n^c] \subseteq \mathfrak{div}_n^0$. Recall that $\mathfrak{div}_n^0 = \mathfrak{iv}_n^0 + \mathfrak{di}_n^0$ (see (11)), $\mathfrak{iv}_n^0 = \oplus_{i=1}^n P^{\partial_i} \partial_i$, $[x^\alpha \partial_i, H_i] = x^\alpha \partial_i$ for all $x^\alpha \in P_n^{\partial_i}$ and $i = 1, \dots, n$. Hence $\mathfrak{iv}_n^0 \subseteq [\mathfrak{div}_n^c, \mathfrak{div}_n^c]$.

Finally, for all $a \in P_n$ and $i \neq j$,

$$[H_k, h_j(a)H_i - h_i(a)H_j] = h_j(H_k * a)H_i - h_i(H_k * a)H_j.$$

Hence, $\{h_j(x^\alpha)H_i - h_i(x^\alpha)H_j \mid 0 \neq \alpha \in \mathbb{N}^n, i \neq j\} \subseteq [\mathfrak{div}_n^c, \mathfrak{div}_n^c]$. Finally, $H_i - H_j = [x_i \partial_j, x_j \partial_i] \in [\mathfrak{div}_n^c, \mathfrak{div}_n^c]$ for all $i \neq j$. Now, by Lemma 2.1.(3), $[\mathfrak{div}_n^c, \mathfrak{div}_n^c] = \mathfrak{div}_n^0$. \square

The following lemma is well-known and easy to prove.

c11Mar13

Lemma 2.11 1. D_n is a simple Lie algebra.

2. $Z(D_n) = \{0\}$.

3. $[D_n, D_n] = D_n$.

The next lemma is also known but we give a short elementary proof.

dc11Mar13

Lemma 2.12 1. \mathfrak{div}_n^0 is a simple Lie algebra.

2. $Z(\mathfrak{div}_n^0) = \begin{cases} \mathfrak{div}_n^0 & \text{if } n = 1, \\ 0 & \text{if } n \geq 2. \end{cases}$

3. $[\mathfrak{div}_n^0, \mathfrak{div}_n^0] = \begin{cases} 0 & \text{if } n = 1, \\ \mathfrak{div}_n^0 & \text{if } n \geq 2. \end{cases}$

Proof. All three statements are obvious if $n = 1$ since $\mathfrak{div}_1^0 = K\partial$. So, we assume that $n \geq 2$.

1. Let $0 \neq a \in \mathfrak{div}_n^0$ and $\mathfrak{a} = (a)$ be the ideal of the Lie algebra \mathfrak{div}_n^0 generated by the element a . We have to show that $\mathfrak{a} = \mathfrak{div}_n^0$. Using the inner derivations $\delta_1, \dots, \delta_n$ and $\text{ad}(H_i - H_j)$ (where $i \neq j$) of the Lie algebra \mathfrak{div}_n^0 we see that $\partial_i \in \mathfrak{a}$ for some i . Then $D_n \subseteq \mathfrak{a}$ since $\partial_j = [\partial_i, x_i \partial_j]$ for all $i \neq j$. Then $\mathfrak{iv}_n^0 \subseteq \mathfrak{a}$ since $x^\alpha \partial_i = [\partial_j, (\alpha_j + 1)^{-1} x^{\alpha + e_j} \partial_i]$ for all $i \neq j$ and $x^\alpha \in P_n^{\partial_i}$. Then $\mathcal{H}'_n \subseteq \mathfrak{a}$ since $[x_i \partial_j, x_j \partial_i] = H_i - H_j$ for all $i \neq j$. Using the commutation relations

$$[H_s - H_t, h_j(x^\alpha)H_i - h_i(x^\alpha)H_j] = (\alpha_s - \alpha_t)(h_j(x^\alpha)H_i - h_i(x^\alpha)H_j) \quad (38)$$

we see that $\{h_j(x^\alpha)H_i - h_i(x^\alpha)H_j \mid i \neq j, \alpha \in \mathbb{N}^n, \alpha \neq (p, p, \dots, p), p = 1, 2, \dots\} \subseteq \mathfrak{a}$. Let $\theta = x_1 \cdots x_n$. Finally, by Lemma 2.1.(3), $\mathfrak{a} = \mathfrak{div}_n^0$ since for all $p = 1, 2, \dots$ and $i \neq j$,

$$[x_j \partial_i, h_j(x_i x_j^{-1} \theta^p)H_i - h_i(x_i x_j^{-1} \theta^p)H_j] = (p+2)(h_j(\theta^p)H_i - h_i(\theta^p)H_j). \quad (39)$$

In more detail,

$$\begin{aligned} \text{LHS} &= [x_j \partial_i, p x_i x_j^{-1} \theta^p H_i - (p+2)x_i x_j^{-1} \theta^p H_j] = p(p+2)\theta^p H_i - (p+2)(p+1)\theta^p H_j \\ &+ (p+2)\theta^p H_i = (p+2)(p+1)(\theta^p H_i - \theta^p H_j) \\ &= (p+2)(h_j(\theta^p)H_i - h_i(\theta^p)H_j). \end{aligned}$$

2. Statement 2 follows from statement 1.

3. Statement 3 follows from statement 1. \square

For any $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$ and any $i \neq j$, we have $\alpha = \alpha_i e_i + \alpha_j e_j + \beta$ where $\beta \in \mathbb{N}^n$ with $\beta_i = \beta_j = 0$. Then

$$\theta_{ij}^\alpha = [x_i^{\alpha_i+1} \partial_j, x^{\alpha - \alpha_i e_i + \alpha_j e_j} \partial_i]. \quad (40)$$

taijx

Indeed, let c be the commutator. Then

$$c = x^\beta [x_i^{\alpha_i+1} \partial_j, x_j^{\alpha_j+1} \partial_i] = x^\beta \theta_{ij}^{\alpha_i e_i + \alpha_j e_j} = \theta_{ij}^\alpha.$$

By (40),

$$\mathfrak{d}\mathbf{i}_n^0 \subseteq [\mathbf{i}\mathbf{v}_n^0, \mathbf{i}\mathbf{v}_n^0]. \quad (41)$$

taijx1

The inner derivations $\delta_1 = \text{ad}(\partial_1), \dots, \delta_n = \text{ad}(\partial_n)$ of the Lie algebra D_n are commuting and locally nilpotent. The Lie algebra D_n is a union of vector spaces

$$D_n = \bigcup_{i \geq 0} D_{n,i} \text{ where } D_{n,i} := \{\partial \in D_n \mid \delta^\alpha(\partial) = 0 \text{ for all } \alpha \in \mathbb{N}^n \text{ such that } |\alpha| = i + 1\}$$

where $\delta^\alpha = \prod_{i=1}^n \delta_i^{\alpha_i}$. Clearly, $D_{n,i} = \oplus_{j=1}^n P_{n,i} \partial_j$ where $P_{n,i} := \{p \in P_n \mid \deg(p) \leq i\}$ and $[D_{n,i}, D_{n,j}] \subseteq D_{n,i+j-1}$ for all $i, j \geq 0$ where $D_{n,-1} := 0$. The inner derivations $\delta_1, \dots, \delta_n$ are also commuting locally nilpotent derivations of the Lie algebras $\mathfrak{d}\mathbf{i}\mathbf{v}_n^0$ and $\mathfrak{d}\mathbf{i}\mathbf{v}_n^c$. For each $i \in \mathbb{N}$, let $\mathfrak{d}\mathbf{i}\mathbf{v}_{n,i}^0 := \mathfrak{d}\mathbf{i}\mathbf{v}_n^0 \cap D_{n,i}$ and $\mathfrak{d}\mathbf{i}\mathbf{v}_{n,i}^c := \mathfrak{d}\mathbf{i}\mathbf{v}_n^c \cap D_{n,i}$. Then, for all $\alpha \in \mathbb{N}^n$ and $i = 1, \dots, n-1$,

tiadi

$$\theta_i^\alpha \in \mathfrak{d}\mathbf{i}\mathbf{v}_{n,|\alpha|+1}^0 \setminus \mathfrak{d}\mathbf{i}\mathbf{v}_{n,|\alpha|}^0. \quad (42)$$

The automorphisms $s_{i,i+1}$, $i = 1, \dots, n-1$ ($n \geq 2$). The automorphism $s = s_{i,i+1}$ of the polynomial algebra P_n that swaps the variables x_i and x_{i+1} leaving the rest of the variables untouched ($s(x_i) = x_{i+1}$ and $s(x_{i+1}) = x_i$) extends uniquely to an automorphism of the Lie algebra D_n . Clearly, s is also an automorphism of the Lie algebras $\mathfrak{d}\mathbf{i}\mathbf{v}_n^0$ and $\mathfrak{d}\mathbf{i}\mathbf{v}_n^c$ (Theorem 2.5). In particular,

$$s(\partial_i) = \partial_{i+1}, \quad s(\partial_{i+1}) = \partial_i, \quad s(H_i) = H_{i+1} \text{ and } s(H_{i+1}) = H_i.$$

Therefore, for all $\alpha \in \mathbb{N}^n$,

siia

$$s_{i,i+1}(\theta_i^\alpha) = -\theta_i^{s_{i,i+1}(\alpha)} \quad (43)$$

where, for $\alpha = (\alpha_1, \dots, \alpha_n)$, $s_{i,i+1}(\alpha) = (\alpha_1, \dots, \alpha_{i-1}, \alpha_{i+1}, \alpha_i, \alpha_{i+2}, \dots, \alpha_n)$. For $i = 1, \dots, n-1$; $j \in \mathbb{N}$ and $\alpha \in \mathbb{N}^n$ with $\alpha_i \geq 1$,

ditai

$$[x_{i+1}^j \partial_i, \theta_i^\alpha] = (\alpha_i + 1) \theta_i^{\alpha - e_i + j e_{i+1}}. \quad (44)$$

Proof. By (27),

$$\begin{aligned} \text{LHS} &= [x_{i+1}^j \partial_i, x^\alpha((\alpha_{i+1} + 1)H_i - (\alpha_i + 1)H_{i+1})] \\ &= x^{\alpha - e_i + j e_{i+1}} (\alpha_i((\alpha_{i+1} + 1)H_i - (\alpha_i + 1)H_{i+1}) - ((\alpha_{i+1} + 1)H_i - (\alpha_i + 1)H_{i+1}, j e_{i+1} - e_i)H_i) \\ &= (\alpha_i + 1) x^{\alpha - e_i + j e_{i+1}} ((\alpha_{i+1} + 1 + j)H_i - \alpha_i H_{i+1}) \\ &= (\alpha_i + 1) \theta_i^{\alpha - e_i + j e_{i+1}}. \quad \square \end{aligned}$$

For $i = 1, \dots, n-1$; $j \in \mathbb{N}$ and $\alpha \in \mathbb{N}^n$ with $\alpha_{i+1} \geq 1$,

ditai1

$$[x_i^j \partial_{i+1}, \theta_i^\alpha] = (\alpha_{i+1} + 1) \theta_i^{\alpha + j e_i - e_{i+1}}. \quad (45)$$

Proof. By applying the automorphism $s = s_{i,i+1}$ to the equality (44) and using (43) we obtain (45):

$$\begin{aligned} [x_i^j \partial_{i+1}, \theta_i^\alpha] &= -s([x_{i+1}^j \partial_i, \theta_i^{s(\alpha)}]) = -s((\alpha_{i+1} + 1) \theta_i^{s(\alpha) - e_i + j e_{i+1}}) \\ &= (\alpha_{i+1} + 1) \theta_i^{s(s(\alpha) - e_i + j e_{i+1})} \\ &= (\alpha_{i+1} + 1) \theta_i^{\alpha + j e_i - e_{i+1}}. \quad \square \end{aligned}$$

By (44) and (45), for $i = 1, \dots, n-1$; $j = 1, \dots, n$ and $\alpha \in \mathbb{N}^n$.

ditai4

$$[\partial_j, \theta_i^\alpha] = \begin{cases} (\alpha_j + 1) \theta_i^{\alpha - e_j} & \text{if } j = i, i+1 \text{ and } \alpha_j \geq 1, \\ (\alpha_{i+1} + 1) x^\alpha \partial_i & \text{if } j = i, \alpha_i = 0, \\ -(\alpha_i + 1) x^\alpha \partial_{i+1} & \text{if } j = i+1, \alpha_{i+1} = 0, \\ \alpha_j \theta_i^{\alpha - e_j} & \text{otherwise.} \end{cases} \quad (46)$$

Proposition 2.13 *Let $n \geq 2$. Then*

1. $\text{Fix}_{\mathbf{G}_n}(\mathcal{D}_n \oplus \mathcal{H}'_n) = \{e\}$.
2. Let $\sigma, \tau \in \mathbf{G}_n$. Then $\sigma = \tau$ iff $\sigma(\partial_i) = \tau(\partial_i)$ for $i = 1, \dots, n$ and $\sigma(H_j - H_{j+1}) = \tau(H_j - H_{j+1})$ for $j = 1, \dots, n-1$.
3. $\text{Fix}_{\mathbf{G}_n}(\partial_1, \dots, \partial_n) = \text{Sh}_n$.

Proof. 1. Let $\sigma \in F := \text{Fix}_{\mathbf{G}_n}(\mathcal{D}_n \oplus \mathcal{H}'_n)$. We have to show that $\sigma = e$. Let $\mathcal{F} := \text{Fix}_{\mathfrak{div}_n^0}(\sigma) := \{\partial \in \mathfrak{div}_n^0 \mid \sigma(\partial) = \partial\}$. We have to show that $\mathcal{F} = \mathfrak{div}_n^0$.

(i) $\mathcal{D}_n + \mathcal{H}'_n \subseteq \mathcal{F}$: Obvious.

(ii) $\mathfrak{iv}_n^0 \subseteq \mathcal{F}$: We have to show that $P_n^{\partial_i} \partial_i \subseteq \mathcal{F}$ for all $i = 1, \dots, n$. Fix i and $x^\alpha \partial_i \in P_n^{\partial_i} \partial_i$. If $\alpha = 0$ then $\partial_i \in \mathcal{D}_n \subseteq \mathcal{F}$, by (i). We use induction on $|\alpha| = \sum_{i=1}^n \alpha_i$ to prove the statement. The case $|\alpha| = 0$ is obvious. So, let $|\alpha| \geq 1$. The element $x^\alpha \partial_i$ has weight $[\alpha - e_i]$. The weight subspace $\mathfrak{div}_{n, [\alpha - e_i]}^0$ is σ -invariant ($\sigma(\mathfrak{div}_{n, [\alpha - e_i]}^0) = \mathfrak{div}_{n, [\alpha - e_i]}^0$) and the set $\{x^\alpha \partial_i, \theta_j^{\alpha - e_i + k\bar{1}} \mid j = 1, \dots, n-1; k = 1, 2, \dots\}$ is its K -basis, by (20). The element $x^\alpha \partial_i$ belongs to the σ -invariant subspace $N_{|\alpha|} := \mathfrak{div}_{n, |\alpha|}^0 (= \mathfrak{div}_n^0 \cap D_{n, |\alpha|})$. By (42) and (46),

$$V := N_{|\alpha|} \cap \mathfrak{div}_{n, [\alpha - e_i]}^0 = Kx^\alpha \partial_i$$

is a σ -invariant subspace. Therefore, $\sigma(x^\alpha \partial_i) = \lambda_{\alpha, i} x^\alpha \partial_i$ for all $x^\alpha \partial_i \in P_n^{\partial_i} \partial_i$ and $i = 1, \dots, n$ for some $\lambda_{\alpha, i} \in K$. Clearly, $\lambda_{0, i} = 1$ for all $i = 1, \dots, n$ since $\sigma(\partial_i) = \partial_i$. Applying the automorphism σ to the relations $[\partial_j, x^\alpha \partial_i] = \alpha_j x^{\alpha - e_j} \partial_i$ yields the equalities $\alpha_j (\lambda_{\alpha, i} - \lambda_{\alpha - e_j, i}) = 0$. If $\alpha_j \neq 0$ then $\lambda_{\alpha, i} = \lambda_{\alpha - e_j, i} = 1$ (by induction). Hence, $\mathfrak{iv}_n^0 \subseteq \mathcal{F}$.

(iii) $\mathfrak{di}_n^0 \subseteq \mathcal{F}$: By (ii) and (41).

Therefore, $\mathfrak{div}_n^0 = \mathfrak{di}_n^0 + \mathfrak{iv}_n^0 \subseteq \mathcal{F}$, and so $\mathfrak{div}_n^0 = \mathcal{F}$, as required.

2. Statement 2 follows from statement 1.

3. Clearly, $\text{Sh}_n \subseteq F := \text{Fix}_{\mathbf{G}_n}(\partial_1, \dots, \partial_n)$. Let $\sigma \in F$ and $H'_{i, j} := \sigma(H_{i, j})$ where $H_{i, j} := H_i - H_j$ for $i \neq j$. Then

$$[\partial_k, H'_{i, j} - H_{i, j}] = \sigma([\partial_k, H_{i, j}]) - [\partial_k, H_{i, j}] = [\partial_k, H_{i, j}] - [\partial_k, H_{i, j}] = 0$$

since $[\mathcal{D}_n, \mathcal{H}'_n] \subseteq \mathcal{D}_n$. Then $d_{ij} := H'_{i, j} - H_{i, j} \in C_{\mathfrak{div}_n^0}(\mathcal{D}_n) = \mathcal{D}_n$, and so $d_{ij} = \sum_{k=1}^n \lambda_{ij}^k \partial_k$ for some $\lambda_{ij}^k \in K$.

If $n = 2$ then $H'_{1, 2} = H_1 - H_2 + \lambda_1 \partial_1 - \lambda_2 \partial_2 = (x_1 + \lambda_1) \partial_1 - (x_2 + \lambda_2) \partial_2 = s_\lambda (H_1 - H_2)$ where $s_\lambda \in \text{Sh}_2$ and $\lambda = (\lambda_1, \lambda_2)$. Hence, $s_\lambda^{-1} \sigma(H_1 - H_2) = H_1 - H_2$, i.e. $s_\lambda^{-1} \sigma \in \text{Fix}_{\mathbf{G}_2}(\mathcal{D}_2 + \mathcal{H}'_2) = \{e\}$, and so $\sigma = s_\lambda \in \text{Sh}_2$.

Suppose that $n > 2$. If $n = 3$ then up to action of Sh_3 , we may assume that $H'_{1, 2} = H_{1, 2} + \lambda \partial_3$ and $H'_{2, 3} = H_{2, 3} + \mu \partial_1 + \nu \partial_2$ for some $\lambda, \mu, \nu \in K$. Then $0 = [H'_{1, 2}, H'_{2, 3}] = -\mu \partial_1 + \nu \partial_2 + \lambda \partial_3$ and so $\lambda = \mu = \nu = 0$.

If $n \geq 4$ then for any four distinct numbers $i, j, k, l \in \{1, 2, \dots, n\}$ we have the equality

$$0 = [H'_{i, j}, H'_{k, l}] = [H_{i, j}, d_{kl}] - [H_{k, l}, d_{ij}] = -\lambda_{kl}^i \partial_i + \lambda_{kl}^j \partial_j + \lambda_{ij}^k \partial_k - \lambda_{ij}^l \partial_l.$$

Therefore, $\lambda_{kl}^i = 0$ for all distinct i, j and k . Then using Sh_n , we may assume that

$$H'_{1, 2} = H_{1, 2}, \quad H'_{2, 3} = H_{2, 3} + \lambda_{23}^2 \partial_2, \quad H'_{3, 4} = H_{3, 4} + \lambda_{34}^3 \partial_3, \dots, \quad H'_{n-1, n} = H_{n-1, n} + \lambda_{n-1, n}^{n-1} \partial_{n-1}.$$

Then, for $i = 2, \dots, n-1$, $0 = [H'_{i-1, i}, H'_{i, i+1}] = \lambda_{i, i+1}^i \partial_i$, i.e. $\lambda_{i, i+1}^i = 0$ and $H'_{i, i+1} = H_{i, i+1}$. This means that $s_\lambda \sigma \in \text{Fix}_{\mathbf{G}_n}(\mathcal{D}_n + \mathcal{H}'_n) = \{e\}$ (statement 2) for some $s_\lambda \in \text{Sh}_n$, and so $\sigma = s_\lambda^{-1} \in \text{Sh}_n$. \square

Lemma 2.14 *Let $\sigma \in \mathbf{G}_n$, $\partial'_1 := \sigma(\partial_1), \dots, \partial'_n := \sigma(\partial_n)$; $\delta'_1 := \text{ad}(\partial'_1), \dots, \delta'_n := \text{ad}(\partial'_n)$ and \mathcal{D}' be the subalgebra of $\text{End}_K(P_n)$ generated by the linear maps $\partial'_1, \dots, \partial'_n$. Then*

1. $\partial'_1, \dots, \partial'_n$ are commuting, locally nilpotent derivations of P_n .
2. $\bigcap_{i=1}^n \ker_{P_n}(\partial'_i) = K$.
3. (Embedding trick) For $n \geq 2$, the short sequence of \mathcal{D}' -modules $0 \rightarrow K \rightarrow P_n \xrightarrow{\Delta'} \mathfrak{div}_n^0 \binom{n}{2}$ is exact where $\Delta'(p) := \prod_{i < j} (\partial'_i(p)\partial'_j - \partial'_j(p)\partial'_i)$. In particular, for all $p \in P_n$ and $\alpha \in \mathbb{N}^n$,

$$\Delta' \partial'^{\alpha}(p) = \delta'^{\alpha} \Delta'(p)$$

where $\partial'^{\alpha} := \prod_{i=1}^n \partial_i^{\alpha_i}$ and $\delta'^{\alpha} := \prod_{i=1}^n \delta_i^{\alpha_i}$.

4. Let $\Delta' : P_n \rightarrow D_n \binom{n}{2}$, $p \mapsto \prod_{i < j} (\partial'_i(p)\partial'_j - \partial'_j(p)\partial'_i)$. Then $\Delta' \in \text{Der}_{\mathcal{D}'}(P_n, D_n \binom{n}{2})$.

Proof. 2. Let $\lambda \in \bigcap_{i=1}^n \ker_{P_n}(\partial'_i)$. Then $\text{div}(\lambda \partial'_1) = \lambda \text{div}(\partial'_1) + \partial'_1 * \lambda = 0$, i.e. $\lambda \partial'_1 \in \mathfrak{div}_n^0$ and

$$\lambda \partial'_1 \in C_{\mathfrak{div}_n^0}(\partial'_1, \dots, \partial'_n) = \sigma(C_{\mathfrak{div}_n^0}(\partial_1, \dots, \partial_n)) = \sigma(C_{\mathfrak{div}_n^0}(\mathcal{D}_n)) = \sigma(\mathcal{D}_n) = \sigma\left(\bigoplus_{i=1}^n K \partial_i\right) = \bigoplus_{i=1}^n K \partial'_i,$$

since $C_{\mathfrak{div}_n^0}(\mathcal{D}_n) = \mathcal{D}_n$, Lemma 2.9.(1). Then $\lambda \in K$ since otherwise the infinite dimensional space $\bigoplus_{i \geq 0} K \lambda^i \partial'_1$ would be a subspace of the finite dimensional space $\sigma(\mathcal{D}_n)$.

3. The derivations $\partial'_1, \dots, \partial'_n$ commute since $\partial_1, \dots, \partial_n$ do. So, \mathcal{D}' is a commutative algebra. For all $p \in P_n$ and $\partial \in \mathfrak{div}_n^0$,

$$\text{div}(a\partial) = a\text{div}(\partial) + \partial(a) = \partial(a).$$

So, the map Δ' is well-defined: $\text{div}(\partial'_i(p)\partial'_j - \partial'_j(p)\partial'_i) = (\partial'_j\partial'_i - \partial'_i\partial'_j)(a) = 0$. Recall that the vector spaces P_n and \mathfrak{div}_n^0 are left \mathfrak{div}_n^0 -modules hence they are also left \mathcal{D}' -modules since $\partial'_1, \dots, \partial'_n \in \mathfrak{div}_n^0$. The map Δ' is a \mathcal{D}' -homomorphism since, for all $p \in P_n$,

$$\Delta' \partial'_i(p) = [\partial'_i, \Delta'(p)], \quad i = 1, \dots, n.$$

It remains to show that $\ker(\Delta') = K$. The inclusion $K \subseteq \ker(\Delta')$ is obvious.

- (i) $\ker(\Delta') = \{p \in P_n \mid \partial'_i(p)\partial'_j = \partial'_j(p)\partial'_i \text{ for all } i \neq j\}$: This is obvious. Let $P_n^{\partial'_i} := \ker_{P_n}(\partial'_i)$.
- (ii) $\ker(\Delta') \cap P_n^{\partial'_i} = K$ for $i = 1, \dots, n$: Given $p \in P_n$. By (i), $p \in \ker(\Delta') \cap P_n^{\partial'_i}$ iff $p \in \ker(\Delta') \cap P_n^{\partial'_j}$ for $j = 1, \dots, n$ ($0 = \partial'_i(p)\partial'_j = \partial'_j(p)\partial'_i \Rightarrow \partial'_j(p) = 0$) iff

$$p \in \ker(\Delta') \cap P_n^{\partial'_1, \dots, \partial'_n} = \ker(\Delta') \cap K = K,$$

by statement 2. Suppose that $K' := \ker(\Delta') \setminus K \neq \emptyset$, we seek a contradiction.

- (iii) If $p \in K'$ then $\partial'_1(p) \neq 0, \dots, \partial'_n(p) \neq 0$: This follows from (ii) ($K' = \ker(\Delta') \setminus K = \ker(\Delta') \setminus \ker(\Delta') \cap P_n^{\partial'_i} = \ker(\Delta') \setminus P_n^{\partial'_i}$).

- (iv) $P_n^{\partial'_1} = \dots = P_n^{\partial'_n}$: Fix $p \in K'$. By (iii), $\partial'_1(p) \neq 0, \dots, \partial'_n(p) \neq 0$. By (i),

$$\partial'_i(p)\partial'_j = \partial'_j(p)\partial'_i \text{ for all } i \neq j,$$

and so $P_n^{\partial'_1} = \dots = P_n^{\partial'_n}$.

- (v) $P_n^{\partial'_1} = \dots = P_n^{\partial'_n} = K$: This statement follows from (iv) and statement 2.

Suppose that $\ker(\Delta') \neq K$, we seek a contradiction. Fix an element $p \in \ker(\Delta') \setminus K$. By statement 2, $\partial'_j(p) \neq 0$ for some j . For all $i \neq j$, $\partial'_i(p)\partial'_j = \partial'_j(p)\partial'_i$ (since $p \in \ker(\Delta')$). Hence, $\partial'_i(p) \neq 0$ for all i . Therefore, for $i = 1, \dots, n$, $\partial'_i = f_i \partial$ for some $f_i \in P_n$ and a derivation $\partial = \sum_{s=1}^n p_s \partial_s \in D_n$ with $\text{gcd}(p_1, \dots, p_n) = 1$. For all elements $c_i \in C'_i := C_{\mathfrak{div}_n^0}(\partial'_i)$, $0 = [c_i, \partial'_i] = c_i(f_i \partial + f_i [c_i, \partial])$, and so $[c_i, \partial] \in P_n \partial$, by the choice of ∂ . By Theorem 1.3, the Lie algebra \mathfrak{div}_n^0 is generated by the set $C = \sum_{i=1}^n C_{\mathfrak{div}_n^0}(\partial_i)$ and also by the set $\sigma(C) = \sum_{i=1}^n C'_i$.

Hence, $[\mathfrak{div}_n^0, \partial] \subseteq P_n \partial$. In particular, for all i , $[\partial_i, \partial] = \sum_{j=1}^n \partial_i(p_j) \partial_j \in P_n \partial$. Hence, $\partial \in \mathcal{D}_n$, by degree argument. Then $P_n^\partial \neq K$, and so $P_n^{\partial_i} = P_n^\partial \neq K$. This contradicts to (v).

4. The map Δ' can be seen as the map

DpPb

$$\Delta' : P_n \rightarrow D_n^{\binom{n}{2}}, \quad p \mapsto \prod_{i < j} (\partial'_i(p) \partial'_j - \partial'_j(p) \partial'_i). \quad (47)$$

Notice that P_n and $D_n^{\binom{n}{2}}$ are left P_n -modules.

$\Delta' \in \text{Der}_{\mathcal{D}'}(P_n, D_n^{\binom{n}{2}})$, i.e. the map Δ' is a \mathcal{D}' -derivation from the polynomial algebra P_n to the left P_n -module $D_n^{\binom{n}{2}}$, i.e. for all $p, q \in P_n$,

$$\Delta'(pq) = q\Delta'(p) + p\Delta'(q) :$$

$$\begin{aligned} \Delta'(pq) &= \prod_{i < j} ((q\partial'_i(p) + p\partial'_i(q))\partial'_j - (q\partial'_j(p) + p\partial'_j(q))\partial'_i) \\ &= q \prod_{i < j} (\partial'_i(p)\partial'_j - \partial'_j(p)\partial'_i) + p \prod_{i < j} (\partial'_i(q)\partial'_j - \partial'_j(q)\partial'_i) \\ &= q\Delta'(p) + p\Delta'(q). \end{aligned}$$

1. The inner derivations $\delta_1, \dots, \delta_n$ of the Lie algebra \mathfrak{div}_n^0 are commuting and locally nilpotent. Hence so are the inner derivations $\delta_1, \dots, \delta'_n$, by statements 2 and 3. \square

A13Mar13

Theorem 2.15 [2] Let $\partial'_1, \dots, \partial'_n$ be commuting, locally nilpotent derivations of the polynomial algebra P_n such that $\bigcap_{i=1}^n \ker_{P_n}(\partial'_i) = K$. Then there exist polynomials $x'_1, \dots, x'_n \in P_n$ such that

con*

$$\partial'_i * x'_j = \delta_{ij} \quad \text{for } i, j = 1, \dots, n. \quad (48)$$

Moreover, the algebra homomorphism

$$\sigma : P_n \rightarrow P_n, \quad x_1 \mapsto x'_1, \dots, x_n \mapsto x'_n$$

is an automorphism such that $\partial'_i = \sigma \partial_i \sigma^{-1} = \frac{\partial}{\partial x'_i}$ for $i = 1, \dots, n$.

db13Mar13

Corollary 2.16 Let $\sigma \in \mathbf{G}_n$. Then $\tau\sigma \in \text{Fix}_{\mathbf{G}_n}(\partial_1, \dots, \partial_n)$ for some $\tau \in G_n$.

Proof. By Lemma 2.14, the elements $\partial'_1 := \sigma(\partial_1), \dots, \partial'_n := \sigma(\partial_n)$ satisfy the assumptions of Theorem 2.15. By Theorem 2.15, $\partial'_1 := \tau^{-1}(\partial_1), \dots, \partial'_n := \tau^{-1}(\partial_n)$ for some $\tau \in G_n$. Therefore, $\tau\sigma \in \text{Fix}_{\mathbf{G}_n}(\partial_1, \dots, \partial_n)$. \square

Proof of Theorem 1.1. If $n = 1$ then $\mathfrak{div}_1^0 = K\partial_1$, $\mathbf{G}_1 = \mathbb{T}^1 \simeq G_1/\text{Sh}_1$ since $G_1 = \mathbb{T}^1 \rtimes \text{Sh}_1$.

So, let $n \geq 2$. Let $\sigma \in \mathbf{G}_n$. By Corollary 2.16, $\tau\sigma \in \text{Fix}_{\mathbf{G}_n}(\partial_1, \dots, \partial_n) = \text{Sh}_n$ (Proposition 2.13.(3)). Therefore, $\sigma \in G_n$, i.e. $\mathbf{G}_n = G_n$. \square

a18Mar13

Lemma 2.17 $\text{Fix}_{\mathbf{G}_n^c}(\mathfrak{div}_n^0) = \begin{cases} \text{Sh}_1 & \text{if } n = 1, \\ \{e\} & \text{if } n \geq 2. \end{cases}$

Proof. If $n = 1$, $\sigma(\partial) = \partial$ for some $\sigma \in \mathbf{G}_1^c$ then applying σ to the equality $[\partial_1, H_1] = \partial_1$ yields $\sigma(H_1) = H_1 + \lambda\partial_1 = (x_1 + \lambda)\partial_1 = t_\lambda(H_1)$ for some $\lambda \in K$ where $t_\lambda \in \text{Sh}_1$. Hence $\text{Fix}_{\mathbf{G}_1^c}(\mathfrak{div}_1^0) = \text{Sh}_1$.

So, let $n \geq 2$. Let $\sigma \in F := \text{Fix}_{\mathbf{G}_n^c}(\mathfrak{div}_n^0)$, $H'_1 := \sigma(H_1), \dots, H'_n := \sigma(H_n)$. By (37), it suffices to show that $\sigma(H_i) = H_i$ for $i = 1, \dots, n$. For $i \neq j$, $\sigma(H_i - H_j) = H_i - H_j$, and so $d := H'_i - H_i = H'_j - H_j$. For all $i = 1, \dots, n$,

$$[\partial_i, d] = \sigma([\partial_i, H_i]) - [\partial_i, H_i] = \sigma(\partial_i) - \partial_i = \partial_i - \partial_i = 0.$$

So, $d \in C_{\mathfrak{div}_n^c}(\mathcal{D}_n) = \mathcal{D}_n$ (Lemma 2.9.(1)) and $d = \sum_{i=1}^n \lambda_i \partial_i$ for some $\lambda_i \in K$. The elements $H'_1 = H_1 + d, \dots, H'_n = H_n + d$ commute hence $d = 0$. Therefore, $\sigma = e$. \square

Proof of Theorem 1.2. If $n = 1$ then $\mathbf{G}_1^c \simeq \mathbb{T}^1 \ltimes \text{Sh}_1$. So, let $n \geq 2$. By (2) and Lemma 2.17, $\mathbf{G}_n^c = G_n$. \square

A17Mar13

Theorem 2.18 For all $\sigma \in G_n$,

$$\text{div}(\sigma(H_i) - H_i) = 0 \text{ for } i = 1, \dots, n.$$

Proof. For each $i = 1, \dots, n$, by (37), $\sigma(H_i) = \lambda_i(\sigma)H_i + d_i(\sigma)$ for some elements $\lambda_i(\sigma) \in K$ and $d_i(\sigma) \in \mathfrak{div}_n^0$. By Theorem 2.5,

$$1 = \text{div}(H_i) = \sigma(\text{div}(H_i)) = \text{div}(\sigma(H_i)) = \text{div}(\lambda_i(\sigma)H_i + d_i(\sigma)) = \lambda_i(\sigma). \quad \square$$

The automorphisms of the Lie algebra \mathfrak{div}_n^c preserve divergence.

b17Mar13

Corollary 2.19 1. For all $\sigma \in \mathbf{G}_n^c$ and $\partial \in \mathfrak{div}_n^c$, $\text{div}(\sigma(\partial)) = \text{div}(\partial)$.

2. Every automorphism of the Lie algebra \mathfrak{div}_n^0 is extendable to an automorphism of the Lie algebra \mathfrak{div}_n^c , and the extension is unique if $n \geq 2$.

Proof. 1. The result follows from Theorem 1.2 and Theorem 2.5.

2. Statement 2 follows from Theorem 1.1 and Theorem 1.2. \square

b2Apr13

Lemma 2.20 For all $n \geq 2$, P_n/K is a simple \mathfrak{div}_n^0 -module/ \mathfrak{div}_n^c -module; $\text{End}_{\mathfrak{div}_n^0}(P_n/K) = \text{End}_{\mathfrak{div}_n^c}(P_n/K) = \text{Kid}$ where id is the identity map. For $n = 1$, P_1/K is neither a simple \mathfrak{div}_1^0 -module nor a simple \mathfrak{div}_1^c -module.

Proof. If $n = 1$ then $\mathfrak{div}_1^0 = K\partial_1$, $\mathfrak{div}_1^c = K\partial_1 + KH_1$ and $\sum_{i \leq m} Kx_i^i \partial_i$ where $m \in \mathbb{N}$ are distinct $\mathfrak{div}_1^0/\mathfrak{div}_1^c$ -submodules of P_1 . Suppose that $n \geq 2$. It suffices to prove the statement for \mathfrak{div}_n^0 . Let M be a nonzero \mathfrak{div}_n^0 -submodule of P_n/K and $0 \neq m \in M$. Using the actions of $\partial_1, \dots, \partial_n \in \mathfrak{div}_n^0$ on m we obtain an element of M of the form $\lambda x_i + K$ for some $\lambda \in K^*$. Hence, $x_i + K \in M$. Then $x_j + K = x_j \partial_i * (x_i + K) \in M$ for all $j \neq i$. So, $x^\alpha + K \in M$ for all $\alpha \in \mathbb{N}^n$ with $|\alpha| = 1$. We use induction on $|\alpha|$ to show that all $x^\alpha + K \in M$. Suppose that $m := |\alpha| > 1$. If $x^\alpha + k = x_i^m + K$ for some $m \geq 2$ and i then $x_i^m + K = (m-1)^{-1} x_i (H_i - 2H_j) * (x_i^{m-1} + K)$. Then, by applying elements of the type $x_j \partial_i$ where $j \neq i$ to the element $x_i^m + K$ we obtain all the elements $x^\alpha + K$ with $|\alpha| = m$. Therefore, P_n/K is a simple \mathfrak{div}_n^0 -module/ \mathfrak{div}_n^c -module.

Let $f \in \text{End}_{\mathfrak{div}_n^0}(P_n/K)$. Then applying f to the equalities $\partial_i * (x_1 + K) = \delta_{i1}$ for $i = 1, \dots, n$, we obtain the equalities

$$\partial_i * f(x_1 + K) = \delta_{i1} \text{ for } i = 1, \dots, n.$$

Hence, $f(x_1 + K) \in \bigcap_{i=2}^n \ker_{P_n/K}(\partial_i) \cap \ker_{P_n/K}(\partial_1^2) = (K[x_1]/K) \cap \ker_{P_n/K}(\partial_1^2) = K(x_1 + K)$. So, $f(x_1 + K) = \lambda(x_1 + K)$ and so $f = \lambda \text{id}$, by the simplicity of the \mathfrak{div}_n^0 -module P_n/K . Therefore, $\text{End}_{\mathfrak{div}_n^0}(P_n/K) = \text{End}_{\mathfrak{div}_n^c}(P_n/K) = \text{Kid}$. \square

a2Apr13

Proposition 2.21 For $n \geq 2$, \mathfrak{div}_n^c is a maximal Lie subalgebra of D_n . For $n = 1$, \mathfrak{div}_1^c is not a maximal Lie subalgebra of D_1 . For each $n \geq 1$, \mathfrak{div}_n^c is a \mathbb{G}_n -invariant/ G_n -invariant Lie subalgebra of D_n .

Proof. For $n = 1$, $\mathfrak{div}_1^c = K\partial_1 + KH_1$ is contained in the Lie subalgebra $K\partial_1 + KH_1 + Kx_1H_1$ of D_1 . Suppose that $n \geq 2$. By (34) and (36),

divDnPn3

$$0 \rightarrow \mathfrak{div}_n^c \rightarrow D_n \xrightarrow{\text{div}} P_n/K \rightarrow 0 \quad (49)$$

is the short exact sequence of \mathfrak{div}_n^c -module. By Lemma 2.20, the \mathfrak{div}_n^c -module P_n/K is simple. Then, \mathfrak{div}_n^c is a maximal Lie subalgebra of D_n .

By Theorem 1.2, Theorem 2.6 and Theorem 2.5, \mathfrak{div}_n^c is a \mathbb{G}_n -invariant/ G_n -invariant Lie subalgebra of D_n . \square

Lemma 2.22 *Let $n \geq 2$. Then*

1. P_n/K is a simple G_n -module with $\text{End}_{G_n}(P_n/K) \simeq K$.
2. $D_n/\text{div}_n^c \simeq P_n/K$, an isomorphism of G_n -modules.
3. D_n/div_n^c is a simple G_n -module with $\text{End}_{G_n}(P_n/K) \simeq K$.

Proof. 1. Let M be a G_n -submodule of P_n properly containing K . We have to show that $M = P_n$, i.e. $x^\alpha \in M$ for all $\alpha \in \mathbb{N}^n$. The polynomial algebra $P_n = \bigoplus_{\alpha \in \mathbb{N}^n} Kx^\alpha$ is the direct sum of 1-dimensional non-isomorphic \mathbb{T}^n -modules. Hence M is a homogeneous submodule of P_n . Hence $x^\beta \in M$ for some $\beta \neq 0$. If $\beta_i \neq 0$ then using the automorphism $s_i : x_i \mapsto x_i + 1, x_j \mapsto x_j, j \neq i$, we see that $M \ni s_i(x^\beta) - x^\beta$ and $\deg_{x_i}(s_i(x^\beta) - x^\beta) = \beta_i - 1$. The module M is closed under the maps $s_i - 1$ for $i = 1, \dots, n$. Hence, $x^\gamma \in M$ for all $\gamma \leq \beta$ where $\gamma \leq \beta$ iff $\gamma_i \leq \beta_i$ for all i . In particular, all $x_1, \dots, x_n \in M$. Then applying the automorphism $\sigma_m : x_1 \mapsto x_1 + x_2^m, x_i \mapsto x_i$ for $i \neq 1$, to the element x_1 , we see that $M \ni (\sigma_m - 1)(x_1) = x_2^m$ for all $m \geq 1$. Then applying the automorphism $x_2 \mapsto \sum_{i=1}^n x_i, x_i \mapsto x_i$ for $i \neq 2$, we have $(x_1 + \dots + x_n)^m \in M$. This implies that all $x^\alpha \in M$, by the homogeneity of M .

Let $f \in \text{End}_{G_n}(P_n/K)$. Since f commutes with the action of the subgroup \mathbb{T}^n of G_n , we must have $f(x^\alpha + K) = \lambda_\alpha(x^\alpha + K)$ for all $\alpha \in \mathbb{N}^n$ and some $\lambda_\alpha \in K$. In particular, $f(x_1 + K) = \lambda(x_1 + K)$ for some $\lambda \in K$. Since f commutes with the action of the symmetric group S_n (which is obviously a subgroup of G_n), $f(x_i + K) = \lambda(x_i + K)$ for all $i = 1, \dots, n$. Now, we use induction on $|\alpha|$ show that $f(x^\alpha + K) = \lambda(x^\alpha + K)$. The initial case when $|\alpha| = 1$ has just been established. So, let $|\alpha| > 1$. Then $\alpha_i > 0$ for some i , and $\deg((s_i - 1)x^\alpha + K) < |\alpha|$. By induction, $f((s_i - 1)x^\alpha + K) = \lambda((s_i - 1)x^\alpha + K)$. Now, it follows from the equality

$$f(s_i(x^\alpha) + K) - f(x^\alpha + K) = f((s_i - 1)x^\alpha + K) = \lambda(s_i - 1)x^\alpha + K$$

that $\lambda_\alpha = \lambda$, and so $f = \lambda \text{id}$. Therefore, $\text{End}_{G_n}(P_n/K) = K \text{id}$.

2. Statement 2 follows from (49) and Theorem 2.5.
3. Statement 3 follows from statements 1 and 2. \square

a14Apr13

Lemma 2.23 *The Lie algebra div_n^0 is a \mathbb{G}_n -invariant/ G_n -invariant Lie subalgebra of D_n .*

Proof. The statement follows from Theorem 1.1, Theorem 2.6 and Theorem 2.5. \square

Conjecture: *Every nonzero homomorphism of the Lie algebra div_n^0 is an automorphism.*

3 Minimal set of generators for the Lie algebras div_n^0 and div_n^c

In this section, the proofs of Theorem 1.3 and Theorem 1.4 are given.

TMMSG
ditai2

$$[x_i^2 \partial_{i+1}, x_{i+1} \partial_i] = \theta_i^{e_i}. \quad (50)$$

In more detail, LHS = $x_i^2 \partial_i - 2x_i x_{i+1} \partial_{i+1} = x_i(H_i - 2H_{i+1}) = \theta_i^{e_i}$.

ditai3

$$[x_{i+1}^2 \partial_i, x_i \partial_{i+1}] = -\theta_i^{e_{i+1}}. \quad (51)$$

Similarly, LHS = $x_{i+1}^2 \partial_{i+1} - 2x_i x_{i+1} \partial_i = -x_{i+1}(2H_i - H_{i+1}) = -\theta_i^{e_{i+1}}$.

Proof of Theorem 1.3. The elements in Theorem 1.3 belong to div_n^0 and let \mathcal{G} be the Lie subalgebra of div_n^0 they generate.

(i) $\mathcal{G} = \mathfrak{div}_n^0$: To prove that the equality holds we use induction on n . Let $n = 2$. So, $\mathcal{G} = \langle \partial_1, x_2^2 \partial_1, x_1^2 \partial_2 \rangle$. Then $\partial_2, x_1 \partial_2, x_2 \partial_1 \in \mathcal{G}$:

$$x_1 \partial_2 = \frac{1}{2} [\partial_1, x_1^2 \partial_2], \quad \partial_2 = [\partial_1, x_1 \partial_2], \quad x_2 \partial_1 = \frac{1}{2} [\partial_2, x_2^2 \partial_1].$$

By Lemma 2.1.(3), we have to show that the elements $x_2^i \partial_1$, $x_1^i \partial_2$ and $\theta_1^\alpha = x^\alpha((\alpha_2 + 1)H_2 - (\alpha_1 + 1)H_1)$ belong to \mathcal{G} where $i \in \mathbb{N}$ and $\alpha \in \mathbb{N}^2$. By (50) and (51),

$$\theta_1^{e_1} = [x_1^2 \partial_2, x_2 \partial_1] \in \mathcal{G}, \quad \theta_1^{e_2} = -[x_2^2 \partial_1, x_1 \partial_2] \in \mathcal{G}.$$

Then using (44) and (45), we see that $\theta_1^\alpha \in \mathcal{G}$ for all $0 \neq \alpha \in \mathbb{N}^2$. For $\alpha = 0$, $\theta_1^0 = H_1 - H_2 = [x_1 \partial_2, x_2 \partial_1] \in \mathcal{G}$. By (26), for $j \geq 1$,

$$\mathcal{G} \ni [x_2 \partial_1, \theta_1^{j e_2}] = -((j+1)H_1 - H_2, -e_1 + e_2)x_2^{j+1} \partial_1 = (j+2)x_2^{j+1} \partial_1.$$

By symmetry, $\mathcal{G} \ni (j+2)x_1^{j+1} \partial_2$, i.e. $\mathcal{G} = \mathfrak{div}_n^0$.

Suppose that $n \geq 3$, and that the equality $\mathcal{G} = \mathfrak{div}_{n'}^0$ holds for all n' such that $2 \leq n' < n$.

Step 1. $\{\partial_i, x_j^\nu \partial_k \mid i, j, k = 1, \dots, n; j \neq k; \nu = 1, 2\} \subseteq \mathcal{G}$: For $i = 2, \dots, n$, $\partial_i = \frac{1}{2} [\partial_1, [\partial_1, x_1^2 \partial_i]] \in \mathcal{G}$ and $x_1 \partial_i = \frac{1}{2} [\partial_1, x_1^2 \partial_i] \in \mathcal{G}$. Then $x_i \partial_1 = \frac{1}{2} [\partial_i, x_i^2 \partial_1] \in \mathcal{G}$ for $i = 2, \dots, n$. For all $i \neq j$ such that $i \neq 1$ and $j \neq 1$, $x_i \partial_j = [x_i \partial_1, x_1 \partial_j] \in \mathcal{G}$ and $x_i^2 \partial_j = [x_i^2 \partial_1, x_1 \partial_j] \in \mathcal{G}$. For $i = 2, \dots, n$, fix an index j such that $j \neq 1, i$. Then $x_i \partial_1 = [x_i \partial_j, x_j \partial_1] \in \mathcal{G}$ and $x_i^2 \partial_1 = [x_i^2 \partial_j, x_j \partial_1] \in \mathcal{G}$. The proof of Step 1 is complete.

For $i = 2, \dots, n$, let $\mathfrak{div}_{n,i}^0$ be the Lie algebra \mathfrak{div}_{n-1}^0 for the polynomial algebra $K[x_1, \dots, \hat{x}_i, \dots, x_n]$ (x_i is missed).

Step 2. For $i = 1, \dots, n$, $\mathfrak{div}_{n,i}^0 \subseteq \mathcal{G}$: This follows from Step 1 and induction.

Step 3. $\mathfrak{di}_n^0 \subseteq \mathcal{G}$: This follows from (25) and Step 2.

Step 4. $\mathfrak{iv}_n^0 \subseteq \mathcal{G}$: This follows from (26) (where $\alpha_i = 0$) and Step 3.

(ii) *Minimality:*

(a) *The element ∂_1 cannot be dropped:* By Lemma 2.1.(3), \mathfrak{div}_n^0 is a \mathbb{Z} -graded Lie algebra which is determined by the degree \deg in the following way, for $x^\alpha \partial_i \in P_n^{\partial_i} \partial_i$, $\deg(x^\alpha \partial_i) = |\alpha| - 1$ and $\deg(\theta_i^\alpha) = |\alpha|$ for all $\alpha \in \mathbb{N}^n$ and $i = 1, \dots, n-1$. Clearly, $\deg(\partial_1) = -1$ (negative) and the degrees the rest of the generators are equal to 1 (positive). Therefore, the ∂_1 cannot be dropped.

(b) *The element $x_i^2 \partial_1$ ($i = 2, \dots, n$) cannot be dropped:* Since otherwise the Lie algebra generated by the remaining elements would belong to $\oplus_{j=1}^n K[x_1, \dots, \hat{x}_i, \dots, x_n] \partial_j$ (x_i is missed), a contradiction (see (i)).

(c) *The element $x_1^2 \partial_i$ ($i = 2, \dots, n$) cannot be dropped:* Since otherwise the Lie algebra generated by the remaining elements would belong to $\oplus_{j \neq i}^n P_n \partial_j$, a contradiction (see (i)). \square

Proof of Theorem 1.4. By Theorem 1.3, the elements in Theorem 1.4 generate the Lie algebra $\mathfrak{div}_n^c = \mathfrak{div}_n^0 \oplus KH_1$ and the element H_1 cannot be dropped (by Theorem 1.3).

(a) *The element ∂_1 cannot be dropped* by the same reason as in the proof of Theorem 1.3 as $\deg(H_1) = 0$.

(b) *The element $x_i^2 \partial_1$ ($i = 2, \dots, n$) cannot be dropped* by the same reason as in the proof of Theorem 1.3.

(c) *The element $x_1^2 \partial_i$ ($i = 2, \dots, n$) cannot be dropped* by the same reason as in the proof of Theorem 1.3. \square

Acknowledgements

The work is partly supported by the Royal Society and EPSRC.

References

- [1] V. V. Bavula, The inversion formula for automorphisms of the Weyl algebras and polynomial algebras, *J. Pure Appl. Algebra* **210** (2007) 147-159; arXiv:math.RA/0512215.

- [2] V. V. Bavula, The group of automorphisms of the Lie algebra of derivations of a polynomial algebra. Arxiv:math.RA:1304.3836.
- [3] D. Z. Dokovic and K. Zhao, Generalized Cartan type S Lie algebras in characteristic zero. *J. Algebra* **193** (1997) no. 1, 144–179.
- [4] J. Grabowski, Isomorphisms and ideals of the Lie algebras of vector fields. *Invent. Math.* **50** (1978/1979), no. 1, 13–33.
- [5] J. Grabowski and N. Poncin, Automorphisms of quantum and classical Poisson algebras. *Compos. Math.* **140** (2004), no. 2, 511–527.
- [6] A. Nowicki, Polynomial derivations and their rings of constants, Torun, 1994.
- [7] J. M. Osborn, Automorphisms of the Lie algebras W^* in characteristic zero, *Canad. J. Math.* **49** (1997) no. 1, 119–132.
- [8] A. N. Rudakov, Subalgebras and automorphisms of Lie algebras of Cartan type. *Funktsional. Anal. i Prilozhen.* **20** (1986) no. 1, 83–84.
- [9] T. Siebert, Lie algebras of derivations and affine algebraic geometry over fields of characteristic 0, *Math. Ann.* **305** (1996) 271–286.
- [10] S. M. Skryabin, Regular Lie rings of derivations. *Moscow Univ. Math. Bull.* **43** (1988) no. 3, 56–58.

Department of Pure Mathematics
 University of Sheffield
 Hicks Building
 Sheffield S3 7RH
 UK
 email: v.bavula@sheffield.ac.uk