



UNIVERSITY OF LEEDS

This is a repository copy of *Interventions Targeting Child Undernutrition in Developing Countries May Be Undermined by Dietary Exposure to Aflatoxin*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/124824/>

Version: Accepted Version

Article:

Watson, S, Gong, YY orcid.org/0000-0003-4927-5526 and Routledge, M (2017) Interventions Targeting Child Undernutrition in Developing Countries May Be Undermined by Dietary Exposure to Aflatoxin. *Critical Reviews in Food Science and Nutrition*, 57 (9). pp. 1963-1975. ISSN 1040-8398

<https://doi.org/10.1080/10408398.2015.1040869>

© 2017, Taylor & Francis Group, LLC . This is an author produced version of a paper published in *Critical Reviews in Food Science and Nutrition*. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Interventions targeting child undernutrition in developing countries may be undermined by dietary exposure to aflatoxin.

Sinead Watson, Yun Yun Gong, Michael N Routledge

Institute for Global Food Security, School of Biological Sciences, Queen's University Belfast, Northern Ireland (SW and YYG); Leeds Institute of Cardiovascular and Metabolic Medicine, School of Medicine, University of Leeds, UK (MNR)

Corresponding Author (reprints):

Michael Routledge, PhD

Leeds Institute of Genetics, Health and Therapeutics

LIGHT Building

Clarendon way

University of Leeds

Leeds LS2 9JT

Tel: 0113 343 7763

Fax 0113 3436300

Email: M.N.Routledge@leeds.ac.uk

Short running head: Aflatoxin exposure and child undernutrition

Abbreviations: AFB1, Aflatoxin B1; AFB2, aflatoxin B2; AFG1, aflatoxin G1; AFG2, aflatoxin G2; AFM1, aflatoxin M1; AF-alb, aflatoxin-albumin adduct; GM, geometric mean; HAZ, height-for-age z score; ID, Iron deficiency; IGF, insulin-like growth factor; LAZ, length-for-age z score; PEM, protein energy malnutrition; RCT, randomized control trial; SMD, standardized mean difference; VAD, vitamin A deficiency; WAZ, weight-for-age z score and WHZ, weight-for-height z score.

1 **Abstract**

2 Child undernutrition is a major adverse public health burden in developing countries,
3 specifically in sub-Saharan Africa and South Asia. Nutrition interventions such as
4 micronutrient supplementation, as well as complementary feeding targeting the major
5 micronutrient deficiencies have only reduced the burden of child undernutrition to a certain
6 extent, indicating that others factors may play a role. Aflatoxin exposure, which is also highly
7 prevalent in developing countries, may be considered to be an aggravating factor for child
8 undernutrition. Increasing evidence suggests that aflatoxin exposure can occur in any stage of
9 life including in utero through a trans-placental pathway and in early childhood (through
10 contaminated weaning food and family food). Early life exposure to aflatoxin is associated with
11 adverse effects on low birth weight, stunting, immune function suppression, and liver function
12 damage. The mechanisms underlying impaired growth and aflatoxin exposure are still unclear
13 but intestinal function damage, reduced immune function and alteration in the insulin-like
14 growth factor axis caused by liver damage, are suggested hypotheses. Given the fact that both
15 aflatoxin and child undernutrition are common in sub-Saharan Africa, effective interventions
16 aimed at reducing undernutrition cannot be satisfactorily achieved until the interactive
17 relationship between aflatoxin and child undernutrition is clearly understood and an aflatoxin
18 mitigation strategy has taken effect in those vulnerable mothers and young children.

19

20 **Keywords:** Aflatoxin, child undernutrition, stunting, kwashiorkor, micronutrient deficiencies

21

22 **Introduction**

23 Child undernutrition including stunting, wasting and micronutrient deficiencies is a major
24 public health problem for low-income countries. The short- and long-term health consequences
25 of child undernutrition can be severe and irreversible and include impaired cognitive

26 development, increased vulnerability to infectious diseases, and reduced educational outcomes
27 and economic productivity in adulthood (1). Furthermore, undernutrition is responsible for
28 approximately 3.1 million child deaths each year, with 45% of all child deaths in 2011 having
29 been attributed to this cause (1).

30

31 It is recognised that there is a window of opportunity for reducing the burden and the lasting
32 impact of child undernutrition, in particular impaired growth. This critical period is defined as
33 the first 1000 days of life from conception to 24 months of age (2, 3). Bhutta et al., (4) reviewed
34 the potential effect on child undernutrition outcomes of interventions such as breastfeeding
35 promotion, micronutrient supplementation and diversified complementary feeding during this
36 critical period and up to 36 months in the 36 countries with the highest burden of child stunting.
37 By modeling the survival and linear growth status of the annual birth cohort from birth to 36
38 months, these authors concluded that existing interventions could potentially reduce stunting at
39 36 months by 36%; mortality by 25% (from birth to 36 months); and stunting, wasting, fetal
40 growth restriction and micronutrient deficiencies disability-adjusted life-years by
41 approximately 25%. Although, these outcomes are encouraging, there are likely to be other
42 underlying determinants of undernutrition that need to be addressed.

43

44 There is increasing evidence that exposure to aflatoxin could be one of the underlying factors.
45 Aflatoxin is a mycotoxin produced by *Aspergillus flavus* and *Aspergillus parasiticus* that
46 contaminate staple crops in many of the countries where child stunting is also prevalent.
47 Although *Aspergillus* molds occur in soil across a wide geographic distribution, hot and humid
48 conditions are favourable for aflatoxin production, with stress to crops caused by drought
49 conditions promoting the contamination of susceptible crops (such as maize and groundnuts)
50 in the field (5). Further growth of the fungus and production of aflatoxin is enhanced by post-

51 harvest storage conditions that involve high humidity (6). There are four main types of
52 aflatoxin, namely aflatoxin B1 (AFB1), aflatoxin B2 (AFB2), aflatoxin G1 (AFG1) and
53 aflatoxin G2 (AFG2). AFB1 is the most potent toxin and is the most prevalent, accounting for
54 an average of 70% of the total aflatoxin content in food, although this may vary depending on
55 the strain of the fungus and local conditions. Aflatoxin M1 (AFM1) is a toxic metabolite of
56 aflatoxin B1, which can be found in milk of lactating mothers, and milk and meat of animals
57 exposed to aflatoxin.

58

59 Human exposure to contaminated food is highest in countries with high consumption of
60 susceptible staple crops grown and stored under optimal fungal growth conditions. Aflatoxin
61 exposure often causes acute outbreaks and sometimes fatal liver toxicity (7). Chronic exposure
62 can increase the risk of liver cancer (8), in particular through an interaction with the hepatitis B
63 virus. There is increasing evidence that aflatoxin plays a role in other health effects such as
64 hepatomegaly (9), immune suppression (10-12) and growth faltering in children (13, 14).
65 Chronic aflatoxin exposure is evident throughout life, including the critical first 1,000 days
66 (15).

67

68 With the increasing evidence that aflatoxin can exacerbate the effects of undernutrition, and
69 contribute to growth faltering, it is likely that aflatoxin exposure has inhibited the expected
70 growth improvement predicted for nutritional intervention programs. In this review we will
71 summarise the burden of childhood undernutrition and the current achievement of nutritional
72 specific interventions for improving child growth, review the evidence for aflatoxin exposure
73 exacerbating undernutrition and reflect on the necessity for considering aflatoxin exposure in
74 nutritional intervention programs.

75

76 **Child undernutrition and nutrition specific interventions in the developing world**

77

78 Protein energy malnutrition

79 Protein energy malnutrition (PEM), considered to be the leading form of childhood malnutrition
80 in developing countries, includes the disorders kwashiorkor, marasmus and marasmus-
81 kwashiorkor, which are differentiated by the balance between inadequate protein intake and
82 other energy sources (16). PEM is often a consequence of suboptimal breastfeeding, delayed
83 and/ or inadequate supplementation of appropriate complementary foods, lack of diet diversity
84 and infection that can lead to decreased absorption of essential nutrients.

85

86 In 2000, the WHO estimated that 26.7% of children < 5 years of age in developing countries
87 had PEM (17). There is a lack of recently conducted population based studies that have
88 investigated the prevalence of the different types of PEM in developing countries. Kwashiorkor,
89 oedematous malnutrition, has been included within the estimates for the prevalence of, and
90 deaths attributable to, severe acute malnutrition (SAM) (weight-for-height (WHZ) below -3,
91 according to WHO standards (18)). In 2011 the global prevalence of SAM in children < 5 years
92 was 3% (19 million) with higher percentages observed in central Africa (5.6%) and south-
93 central Asia (5.1%) (1).

94

95 A recent systematic review evaluated the effectiveness of inpatient management for SAM using
96 the WHO protocol, as well as community-based treatments in low- and middle-income settings
97 (19). The authors found that case fatality rates for inpatient management of SAM, following
98 the WHO protocol, which involves fluid management and micronutrient supplementation,
99 ranged from 3.4% to 35%. Only two studies reported nutrition recovery rates, which were
100 79.7% and 83.3%. For the community-based treatment of SAM that involves the use of ready-

101 to-use therapeutic food (RUTF), 51% of children were more likely to achieve nutritional
102 recovery than a standard care group. Although, this systematic review was limited in the
103 availability of high quality studies, the nutritional recovery rates of the interventions reviewed
104 were advantageous. The authors have concluded that future studies are warranted to compare
105 approaches to managing SAM and this includes identifying and tackling other aggravating
106 determinants of SAM.

107

108 Growth faltering

109 Stunting (height-for-age Z score (HAZ) < 2), wasting (weight-for-height Z score (WHZ) < 2)
110 and underweight (weight-for-age Z score (WAZ) < 2) (18) are major indicators of child
111 undernutrition. Severe undernutrition is considered when Z scores are < 3 . In 2011,
112 approximately 165 million (25.7%) children under the age of five years globally had stunted
113 growth, 52 million (8%) were classified as wasting and 100 million (16%) were underweight
114 (20). South-central Asia (36% stunted, 15% wasting and 30% underweight) as well as East
115 (42% stunted) and West Africa (36% stunted and 22% underweight) had the highest prevalence.
116 Growth faltering in early life is a predisposing risk factor for poor cognitive development,
117 reduced educational outcomes and economic productivity, as well as reduced survival in
118 adulthood (1). Micronutrient deficiencies alongside recurring infections are some of the well-
119 recognised causes of child growth faltering in developing countries. There are three
120 micronutrient deficiencies of public health concern in developing countries; vitamin A, Iron
121 and zinc deficiency. Interventions (supplementation) targeting these specific micronutrients and
122 their impact on growth outcomes are summarised in **table 1**.

123

124 Zinc deficiency

125 A recent analysis conducted by Wessells and Brown (21) estimated the global prevalence of
126 zinc deficiency (ZD) as 17% in 188 countries, using zinc intake obtained from FAO food
127 balance sheets, with zinc and phytate contents calculated using a nutrient composition database
128 (table 1). Low-income countries such as those in sub-Saharan Africa and South Asia were most
129 at risk with a ZD prevalence of over 25% (21). ZD is primarily caused by low intake of animal
130 products and exacerbated by persistent diarrhoea (22, 23). ZD can negatively impact the
131 immune system, thereby enhancing susceptibility to infectious diseases such as diarrhoea,
132 malaria and pneumonia, especially in children (22). It may also aggravate intestinal
133 permeability and chronic inflammation, both pathways that underlie environmental
134 enteropathy, which is a sub-clinical condition involving reduced intestinal function that can
135 affect micronutrient absorption (23). Zinc has a fundamental role in cell division and growth;
136 thus, it can result in decreased concentrations of circulatory Insulin-like Growth Factor 1 (IGF-
137 1), a possible pathway for slowed child growth in Zinc deficient children (24).

138
139 ZD in developing countries coincides with the high prevalence of stunted growth in children
140 observed in these countries (1, 21). In fact, assessing the number of children < 5 years old that
141 have stunted growth has been considered to be a proxy for zinc deficiency (21, 25), although
142 this is an indirect method of measuring ZD, and consequently is subject to confounding factors.
143 It would be expected, therefore, that zinc supplementation would have a positive effect on
144 growth. Four meta-analyses (26-29) have been identified that have investigated the impact of
145 zinc supplementation on growth indices in childhood (table 1). Three meta-analyses found that
146 zinc supplementation had a significant positive effect on linear growth (26, 27, 29) and two
147 found it had a positive effect on weight gain (26, 27). In contrast, Ramakrishnan et al. (28)
148 found no effect of zinc supplementation on linear growth or weight change but did find a
149 significant positive effect on change in WHZ score. Although it is apparent from the

150 aforementioned evidence that zinc can have a positive impact on growth, it is important to
151 highlight that its effect is only marginal.

152

153 Iron deficiency

154 Iron deficiency (ID) is the leading cause of anaemia (haemoglobin < 110g/L) and accounts for
155 ~50% of all cases (30). For this reason anaemia is typically used as a proxy for ID. Stevens et
156 al. (31) estimated the global prevalence of total and severe anaemia in three population groups
157 known to be most vulnerable to these conditions; women of child bearing age (15-49 years),
158 children (6-59 months) and pregnant women. Using representative population based data
159 collected from 107 countries, it was evident that anaemia is of epidemic proportions worldwide
160 (table 1). Regional analysis showed Central and West Africa as having the highest prevalence
161 of anaemia and severe anaemia in children aged < 5 years in 1995 (80% and 9.7%) and 2011
162 (71% and 4.9%). The high prevalence observed in developing parts of the world is mostly likely
163 due to diets low in iron rich foods alongside poor absorption and diets high in phytate
164 compounds that inhibit iron absorption (32). Parasite infections as well as tuberculosis and HIV
165 are also thought to be risk factors.

166

167 Poor growth and cognitive development during childhood have been suggested as major
168 consequences of iron deficiency; although, the evidence supporting these suggestions is
169 inconclusive. For example, several systematic reviews and meta-analyses of randomized
170 controlled trials (RCTs) have failed to discover a positive effect of iron supplementation on
171 different growth parameters in children (table 1) (33-36). However, a recent systematic review
172 and meta-analysis (37), found a small positive effect on growth (HAZ) in children that were
173 aged between 5 and 12 years. Likewise, systematic reviews have reported that iron
174 supplementation can have an impact on cognitive development especially in older children (37,

175 38) but appears to be ineffectual in young children and infants (35, 36, 38, 39). This evidence
176 indicates that iron supplementation may have more of an impact on growth performance and
177 cognitive development during mid-childhood. Of course, this may challenge the view that
178 interventions targeting growth should occur in the first 1,000 days of life (2, 3), as beyond this
179 timeframe interventions are considered to be ineffectual. Nevertheless, it is noticed that the
180 positive effect on growth reported in these studies (37) was only marginal, indicating that iron
181 supplementation targeting mid childhood may only have limited success as a public health
182 intervention.

183

184 Vitamin A deficiency

185 According to a WHO (40) report, vitamin A deficiency (VAD), defined as having serum
186 (plasma) retinol concentrations less than $< 0.70 \mu\text{mol/l}$ or having a history of night blindness in
187 more severe cases, is considered a major public health problem in developing countries,
188 specifically in Asia and sub-Saharan Africa. In that report, the global prevalence of VAD
189 measured between 1995 and 2005 in pregnant women was 15.3% and when stratified according
190 to WHO regions, Africa and Asia had the highest rates (14.3% and 18.4%). This trend was also
191 observed in children under 5 years old. Global prevalence was 33.3%, with Africa (41.6%) and
192 Asia (33.5%) having higher rates than other parts of the world.

193

194 The developing fetus and preschool aged children are considered to be at-risk populations,
195 owing to the rapid growth and subsequent increased nutritional requirements during these
196 stages of the life course. In developing countries these additional nutritional requirements are
197 frequently not met owing to the lack of diet diversity, as well as the affordability of foods high
198 in vitamin A such as animal products, citrus fruits and dark green vegetables.

199

200 Over the past decade, some observational studies have found that maternal VAD was associated
201 with lower birth weight (41, 42). In contrast, according to a recent systematic review and meta-
202 analysis vitamin A supplementation during pregnancy had no positive effect on birth weight
203 (43). Furthermore, vitamin A supplementation during childhood showed little or no effect on
204 growth performance in several RCTs (33, 44-47).

Table 1: Major micronutrient deficiencies of public health concern: prevalence, supplementation and growth outcomes in children

Micronutrient	Prevalence	Micronutrient supplementation on growth. Evidence from systematic reviews and meta-analyses	Effects on physical growth (95%CI)
Zinc	<p>Wessells and Brown (21) Using country specific FAO food balance sheets All ages (> 6months) Global: 17.3 ± 11.1% sub-Saharan Africa: 25.6 ± 12.2% South Asia: 29.6 ± 3.6% Prevalence of inadequate zinc intake was correlated with the prevalence of stunting in children < 5 years (r = 0.48; p <0.001)</p>	<p>Brown et al. (26) Meta-analysis of RCTs Children <12 years or prepubertal Zinc supplementation ≥ 8 weeks</p> <p>Brown et al. (27) Meta-analysis of RCTs Infants, pre-schooler and older pre-pubertal Zinc supplementation 2 weeks to 15 months</p> <p>Ramakrishnan et al. (28) Meta-analysis of RCTs Children ≤5 years Zinc supplementation ≥ 8 weeks</p>	<p>Zinc supplementation had a positive effect on change in height (effect size = 0.35; 95% CI: 0.19-0.51) and change in weight (effect size = 0.31; 95% CI: 0.18-0.44). There was no significant effect on WHZ.</p> <p>Zinc supplementation had a positive effect on change in height (effect size = 0.17; 95% CI: 0.08-0.26), change in weight (effect size = 0.12; 95% CI: 0.05-0.19) and a small marginal effect on change in WHZ score (effect size = 0.06; 95% CI: 0.00-0.12) compared with control groups.</p> <p>Zinc supplementation had no significant positive effect on change height or weight gain but did have a small positive effect on WHZ score (effect size = 0.06; 95% CI: 0.01-0.11) in comparison with placebo-controlled groups.</p>

		Imdad and Bhutta (29) Meta-analysis of RCTs Children < 5 years Zinc supplementation \geq 8 weeks	Zinc supplementation had a positive effect on linear growth (effect size = 0.19; 95% CI: 0.08-0.30) compared to placebo-controlled group.
Iron	Stevens et al., (31) Iron deficiency anaemia (haemoglobin <110 g/L) Children < 5 years Global: 43% (95% CI: 38-47) Central and West Africa: 71% (95% CI: 67-74) South Asia: 58% (95% CI: 44-69)	Ramakrishnan et al. (33) Meta-analysis of RCTs Children < 18 years Iron supplementation \geq 8 weeks	Iron supplementation had no significant effect on height or weight compared to a control group.
		Sachdev et al. (34) Meta-analysis of RCTs Children < 14 years Oral Iron supplementation duration 2 months to 12 months	Iron supplementation had no significant effect on WAZ, WHZ, HAZ, mid upper arm circumference, skinfold thickness or head circumference compared to control groups.
		Low et al. (37) Meta-analysis of RCTs Children 5 to 12 years Oral iron supplementation \geq 5 days per week	Iron supplementation had no significant effect on absolute height or absolute weight or WHZ score but did have a significant but small positive effect on HAZ score compared with a control group (effect size = 0.09; 95% CI: 0.01-0.17).
		Pasricha et al. (35) Meta-analysis of RCTs Children aged 4-23 months	Iron supplementation had no significant effect ($P > 0.05$) on final weight, WAZ scores, change in

		Daily oral iron supplementation	weight, final length, HAZ scores, change in length or weight for length z score in comparison with the control group.
		Thompson et al. (36) Meta-analysis of RCTs Children 2 to 5 years Oral iron supplementation \geq 5 days per week	Iron supplementation had no positive effect on final height, final weight, change in height and change in weight compared to a control group
Vitamin A	WHO (40) (Serum retinol <0.70 $\mu\text{mol/L}$) Children < 5 years Global: 33.3% (95% CI: 31.1-35.4) Africa: 44.4% (95% CI: 41.3-47.5) South East Asia: 49.9% (95% CI: 45.1-54.8)	Ramakrishnan et al. (33) Meta-analysis of RCTs Children < 18 years Vitamin A supplementation ≥ 8 weeks	Vitamin A supplementation had no positive effect on absolute height change or weight change

205 It is clear from the evidence above that supplementation interventions targeting the main
206 micronutrients of public health concern in developing countries are not entirely effective in
207 improving child growth. Vitamin A supplementation markedly has no impact on child growth,
208 whereas zinc and iron supplementation seem to have peripheral effects. This suggests that there
209 are other underlying determinants of child growth faltering that need to be addressed.

210

211 **Aflatoxin related undernutrition issues in the developing world**

212 Aflatoxin exposure and its relationship with growth faltering.

213 The development and application of the AF-alb biomarker has enabled a number of
214 epidemiology studies examining human health effects of aflatoxin exposure (48). This
215 biomarker, which is usually measured by an ELISA method (49) has shown a good correlation
216 with aflatoxin intake in adults through a groundnut based diet in The Gambia (50), as well as
217 in children through a maize-based weaning diet in Tanzania (51). Compared to other available
218 short term (for previous 1-2 days exposure) biomarkers such as the aflatoxin DNA adduct,
219 AFM1 and aflatoxin metabolites in urine, this biomarker reflects the previous 2-3 months
220 exposure at the individual level, and is therefore more appropriate for assessing chronic
221 exposure related health outcomes.

222

223 There is mounting evidence that aflatoxin exposure occurs from gestation onwards throughout
224 life (15). Exposure occurs in utero through the transfer of aflatoxins from the mother to the
225 foetus via the placenta. Several studies have investigated this route of exposure and have found
226 detectable levels of aflatoxin or AF-alb in cord blood samples (52-56). Only a few studies have
227 examined the impact of exposure in utero on birth weight (57-59). All have reported a
228 significant inverse relationship with higher exposure in utero corresponding to lower weight at

229 birth. A study by de Vries et al (57) conducted in rural Kenya, examined aflatoxin levels in
230 maternal and cord blood samples. Aflatoxin was detected in over half of the maternal samples
231 and 37% of the cord blood samples. Females born to aflatoxin positive mothers had a mean
232 birth weight that was 225g lower than those born to mothers free from aflatoxin exposure.
233 Similar results were observed in a study conducted in the Middle East by Abdulrazzaq et al.,
234 (58), where high aflatoxin levels in maternal and cord blood samples were significantly related
235 to lower birth weights ($r = -0.654$, $P = 0.0001$ and $r = -0.565$, $P = 0.001$, respectively). More
236 recently, a cross-sectional study of 785 pregnant Ghanaian women, after adjusting for socio-
237 demographic variables and other factors, found increased odds of delivering a baby with a low
238 birth weight in the highest quartile (59). The highest quartile represented the highest levels of
239 aflatoxin exposure measured in blood during pregnancy (OR, 2.09; 95% CI: 1.19–3.68).

240

241 Aflatoxin exposure in utero may also play a role in stunted growth in early childhood (up 24
242 months). Only one study to date has explored this temporal relationship (56), and found that
243 higher levels of AF-alb in maternal blood were significantly associated with lower weight ($P =$
244 0.012) and height ($P = 0.044$) gain, after adjusting for potential confounding factors.
245 Furthermore, the authors predicted that a reduction in maternal AF-alb level from 110 pg/mg
246 to 10 pg/mg would lead to a 2 cm increase in height and a 0.8 kg increase in weight within the
247 first 24 months of life.

248

249 Usually studies that have examined exposure in utero by measuring maternal blood only
250 obtained measurements at one point in time. A recent study conducted by Castelino et al., (60)
251 explored the effect of season and gestation stage on aflatoxin exposure in pregnant women from
252 Gambia. Results showed that mean AF-alb levels were higher during the dry season than the

253 rainy season. AF-alb levels increased marginally from early to later gestation during the dry
254 season (41.8 vs 34.5 pg/mg; $P < 0.05$). Although early pregnancy has been considered a period
255 when the foetus is most vulnerable, later pregnancy marks the fast growth period of the foetus,
256 which may exert a profound adverse impact on growth. Further research is warranted to
257 determine the longer term health effects of aflatoxin exposure during both early and late
258 pregnancy.

259

260 Weaning is the transition from breast milk to solid food, and typically commences between 3
261 and 6 months. It is often a period in developing countries when children are most susceptible
262 to PEM, specifically, kwashiorkor. Because weaning foods such as maize are prone to aflatoxin
263 contamination, there may also be high aflatoxin exposure during the weaning period. This was
264 evident in a study conducted by Gong et al., (61) in Benin and Togo, as children that were fully
265 weaned had approximately 2-fold higher mean AF-alb levels than children who were still
266 partially breastfed. Although breastfeeding is a period of lower aflatoxin exposure, there is still
267 some exposure from breast milk, with aflatoxin M1 having been found in breast milk samples
268 in many studies (62). Nevertheless, AFM1, which is the hydroxylated metabolite of aflatoxin
269 that is found in milk, is less toxic than AFB1 that is found in food; therefore extending the
270 breastfeeding period may help reduce the negative health impacts, such as growth faltering, that
271 are associated with aflatoxin exposure.

272

273 The impact of aflatoxin exposure on growth is considered the most prominent during the first
274 two years after birth. One of the first studies examining the association between aflatoxin
275 exposure and child growth performance was a cross-sectional study of 480 children from Benin
276 and Togo aged between 9-months and 5 years (13). Prevalence of aflatoxin was high in this

277 sample with 99% of the children having detectable levels and a reported geometric mean of
278 32.8 pg/mg. Undernutrition was also evident as 33%, 6 % and 29% of the children were
279 classified as having stunted growth (HAZ <-2), wasting (WHZ <-2) and being underweight
280 (WAZ <-2); respectively. Significant negative correlations between AF-alb and each of the
281 growth parameters were observed (P = 0.001 for stunting; P = 0.047 for wasting and P = 0.005
282 for underweight). Another cross-sectional study by Turner et al., (10) found that AF-alb levels
283 were weakly associated with wasting (P = 0.034) but not with stunting or underweight.

284

285 These earlier studies were the first in determining the association of aflatoxin dietary exposure
286 with growth impairment in human subjects, and generated hypotheses for further investigations.
287 Cross-sectional studies are the best way to measure prevalence (63); however, they do have
288 limitations, as they cannot be used to establish the temporal sequence of the relationship
289 observed. A subsequent study using a longitudinal design, examined the effects of aflatoxin
290 exposure on growth in a cohort of 200 children from Benin (16-37 months) followed up over
291 8-months (14). High prevalence of aflatoxin exposure was found across the cohort with almost
292 all samples being positive for aflatoxin at each time point and with mean AF-alb levels of 37.4
293 pg/mg (February), 38.7 pg/mg (June) and 86.8 pg/mg (October). Results showed that both AF-
294 alb levels measured in February and the mean AF-alb level from the three time points, were
295 inversely correlated with HAZ and WHZ growth parameters that were measured at the end of
296 the study. This relationship remained after adjusting for potential confounding factors such age,
297 sex, height, weaning status, SES and geographical location, although only for the HAZ growth
298 parameter (P < 0.001). Furthermore, there was a difference in height of 1.7 cm between the
299 highest and lowest AF-alb quartile over the 8 month period. This study has helped to show the
300 temporal relationship between aflatoxin exposure and impaired child growth. Although
301 additional longitudinal studies conducted in different geographical locations and populations

302 will strengthen the evidence on the likelihood of this effect being cause and effect. Furthermore,
303 plausible mechanisms that link aflatoxin exposure with impaired child growth should be
304 investigated.

305

306 Aflatoxin exposure and protein-energy malnutrition

307 It has been proposed that the development of kwashiorkor may be partly attributable to aflatoxin
308 exposure, although the evidence is circumstantial. Both aflatoxin exposure and kwashiorkor are
309 prevalent in hot and humid tropical countries where maize and rice are staples, both affect
310 children in early life and both are associated with impaired child growth (15, 64). In addition,
311 the clinical and metabolic manifestations of kwashiorkor are somewhat similar to those of
312 aflatoxin exposure, such as fatty liver and immunosuppression (65).

313

314 As shown in **table 2**, the association between the exposure to aflatoxin and kwashiorkor has
315 been investigated in a plethora of studies since the 1980's (65-78). The typical study designs
316 employed by the majority of these studies were case-control or cross-sectional, and involved
317 measuring the prevalence and concentration of aflatoxin in blood and urine samples. In most
318 studies it was found that aflatoxin was detected more frequently or concentrations were higher
319 in blood samples of children with kwashiorkor in comparison with children with marasmus,
320 and healthy children (65, 68, 69, 76-78). Furthermore, aflatoxin was detected more often in
321 liver specimens from children who had died from kwashiorkor compared to other diseases and
322 other protein malnutrition disorders (66).

323

324 Although evidence suggests that aflatoxin exposure may be related to kwashiorkor prevalence,
325 a causal relationship has not been established. Furthermore, most of the studies did not measure
326 AF-alb levels in serum of exposed children, which has been shown to be a more reliable
327 biomarker. A fundamental step in unravelling any link between aflatoxin and kwashiorkor is to
328 understand the possibility that the metabolic manifestations of kwashiorkor affect the way that
329 aflatoxins are metabolised and excreted from the body, or vice versa. Future studies,
330 undertaking a longitudinal design are required to determine if aflatoxin exposure plays an
331 aetiological role in the causation of kwashiorkor.

Table 2. The relationship between protein energy malnutrition and aflatoxin exposure

Study	Country/ study population	Aflatoxin Exposure		
		Blood – detection (%) and mean concentration	Urine – detection (%) and mean concentration	Other – detection (%) and mean concentration
Hendrickse et al. (65)	Country: Sudan 252 children K (n = 44) MK (n = 32) M (n = 70) AM controls (n = 106)	177 samples (total aflatoxin pg/ml). K (36.4%) (GM: 706) MK (21.9%) (GM: 412) M (19.3%) (GM: 211) AM controls (15.9%) (GM: 77) The difference between the groups approached significance (P = 0.05). Kwashiorkor group mean aflatoxin concentration was significantly higher than the control group (P = 0.01).	250 samples (total aflatoxin pg/ml). K (36.4%) (GM: 706) MK (21.9%) (GM: 412) M (19.3%) (GM: 211) AM controls (15.9%) (77) No significant differences between the groups identified.	
Lamplugh and Hendrickse, (66)	Country: Nigeria and South Africa 8 children (aged between 9 months and 24 months) K (n = 3) MK (n = 3) M (n = 1) Control (n = 1)			8 autopsy liver specimens. K (all three of the liver samples contained AFB1: 2000, 4900 and 1400 pg/g). MK (1 liver sample had no aflatoxins; one contained a small quantity of aflatoxin M1 (15 pg/g) and in the third sample aflatoxicol was found (8500 pg/g). M (no aflatoxins found) Control (no aflatoxin found)

Apeagyei et al., (67)	Country: Ghana 22 children (aged between 5 months and 48 months) K (n = 22)			22 autopsy liver specimens Aflatoxin B1 was detected in 20 of the samples (90.9%). The remaining 2 samples contained aflatoxicol (9.1%).
Coulter et al., (68)	Country: Sudan 584 children K (n = 141) MK(n = 152) M (n = 152) AM controls (n = 180)	457 samples (total aflatoxin pg/ml) K (37.7%) (GM: 154) MK (28.6%) (GM: 82) M (26.3%) (GM: 77) AM controls (21.3%) (GM: 81) Difference between the number of positive samples found in each group was significant (P <0.05). No differences between the groups in concentrations of aflatoxin identified.	463 samples (total aflatoxin pg/ml) K (27.2%) (GM: 308) MK(39.0%) (GM: 490) M (26.1%) (GM: 438) AM controls (28.4%) (GM:258) No significant difference between the numbers of positive samples found in each group. No significant differences between the groups in concentrations of aflatoxin identified.	
deVries et al., (69)	Country: Kenya 41 children K (n = 14) MK (n = 6) M (n = 11) Controls (n = 10)	39 samples (total aflatoxin (pg/ml) K (64%) (mean: 6666) MK (50%) (mean: 386) M (36%) (mean: 3412) Controls (30%) (mean: 759)	36 samples (total aflatoxin pg/ml) K (42%) (mean: 324) MK (60%) (mean: 1294) M (45%) (mean: 261) Controls (75%) (mean: 759) No differences in detection rates.	
de Vries et al., (70)	Country: Kenya 13 children K (n = 5)		K (4 out of 5 children excreted aflatoxin via urine).	K (all 5 of the children excreted aflatoxin in their stools).

	MK (n = 7) Underweight (n = 1)		MK (5 out of 7 children excreted aflatoxin via urine). The underweight child's urine samples tested negative for aflatoxin.	Total aflatoxin excreted (urine and stools) ranged from 0.08 ug/kg to 4 ug/kg body weight). MK (3 out of 7 children excreted aflatoxin in their stools). Total aflatoxin excreted (urine and stools) ranged from nil to 1.5 ug/kg body weight). The underweight child's stools tested negative for aflatoxin.
Househam and Hundt (71)	Country: South Africa 320 children (mean age of 38 months) K (n = 47) M (n = 17) Controls (n = 256)		448 urine samples Aflatoxin B1, B2, G1, G2 and aflatoxicol were not detected in any of the samples.	
Ramjee et al., (72)	Country: South Africa 109 children aged between 6 months and 2 years K (n = 45) M (n = 13) Underweight (n = 16) AM controls (n = 35)	109 samples K (56%) M (31%) Underweight (56%) AM controls (49%) No differences among the groups in the number of aflatoxin positive results.	50 samples K (16%) M (10%) Underweight (no samples tested) Age matched controls (25%) No differences among the groups in the number of aflatoxin positive results. The serum/ urine ratio was significantly higher in the kwashiorkor group than in the other groups (P = 0.001).	

Adhikari et al., (73)	Country: South Africa 36 children aged between 6 months and 2 years K (n=36)	36 samples Aflatoxin was detected in 21 samples (58%)	
Oyelami et al., (74)	Country: Nigeria 40 children (aged between 4 and 168 months) 20 children who died from kwashiorkor 20 children who died of other diseases		40 lung specimens K (90%) Other diseases (65%) No significant differences among the groups in the number of aflatoxin positive results.
Oyelami et al., (75)	Country: Nigeria 45 children 24 children who died from kwashiorkor (aged between 6 months and 72 months) 21 children who died of other diseases (aged between 4 months and 168 months)		45 kidney specimens (total aflatoxin pg/g) K (58%) (mean: 3851) Other diseases (62%) (mean: 1271) No significant differences among the groups in the number of aflatoxin positive results. No differences among the groups in mean concentrations of total aflatoxins.
Hatem et al., (76)	Country: Egypt 70 children (aged between 6 and 24 months) K (n = 30) M (n = 30) AM controls (n = 10)	30 samples (total aflatoxin ng/ml) K (80%) (mean: 70.58) Ma (46.7%) (mean: 25.21) AM controls (0) Aflatoxins were detected more frequently in blood samples of	30 samples (total aflatoxin ng/100ml) K (80%) (mean: M (46.7%) AM controls (0) Aflatoxins were detected more frequently in urine samples of

		the kwashiorkor group than the marasmus group (P = 0.007). Mean serum levels of total aflatoxin were significantly higher in the kwashiorkor group relative to the marasmus group (P <0.001).	the kwashiorkor group than the marasmus group (P = 0.007). Mean levels of total aflatoxin excreted in urine were significantly higher in the kwashiorkor group relative to the marasmus group (P = 0.052).
Tchana et al., (77)	Country: Cameroon 78 children (aged between 13 months and 12 years) K (n = 31) MK (n=11) AM controls (n = 36)		42 samples (aflatoxin B1) K (35.5%) MK (45.5%) AM controls (11.1%) Detection levels of AFB1 excreted in urine were significantly higher in the kwashiorkor and marasmus kwashiorkor groups relative to the control group (P <0.05).
Onyemelukwe et al. (78)	Country: Nigeria 111 children (aged between 7 months and 60 months) K (n = 36) MK (n = 29) M (n = 13) AM controls (n = 33)	111 samples (total aflatoxin ug/L) K (88.9%) (median: 165.6) MK (93.1%) (median: 228.4) M (76.9%) (median: 234.3) AM controls (63.6%) (median: 20.7) Median serum levels of total aflatoxin were significantly higher in each protein energy malnutrition group relative to	55 samples (total aflatoxin ug/L) K (84.6%) (median: 79) M (60%) (median: 43.8) Ma (81.8%) (median: 14.4) AM controls (90.9%) (median: 42.6) No differences among the groups in the number of aflatoxin positive results.

the control group (kwashiorkor vs. control P <0.001, marasmic kwashiorkor vs. control P <0.001, marasmus vs. control P = 0.031). There were no significant differences between the protein malnutrition groups.	Median total aflatoxin levels in urine samples were significantly higher in the kwashiorkor group relative to the marasmus group (P = 0.011). No other significant differences were identified between the groups.
---	--

Abbreviation: AM, age-matched; GM, geometric mean; K, kwashiorkor; M, marasmus; MK, Marasmic kwashiorkor

332 Aflatoxin exposure and micronutrient deficiencies

333 It has been hypothesized that aflatoxin exposure mediates intestinal damage resulting in
334 reduced nutrient absorption and increased intestinal permeability resulting in faltered growth
335 (79, 80). It is, therefore, possible that aflatoxin exposure exacerbates micronutrient deficiencies
336 and by reducing aflatoxin exposure the incidence of micronutrient deficiencies may be reduced
337 correspondingly. Previous research has established the relationship between aflatoxin exposure
338 and the effect on these micronutrients in feeding experiments in animal studies as reviewed by
339 Williams et al., (81). Increasing levels of aflatoxin in feed were significantly related to
340 decreasing concentrations of vitamin A in poultry (82); vitamin D concentrations in chickens
341 (83); vitamin A and E in swine (84) as well as zinc in piglets (85).

342

343 Owing to the species difference, it is difficult to directly apply these findings to humans. Only
344 a few studies have been identified that have examined the relationship between micronutrient
345 concentrations and aflatoxin exposure in humans. Two of these studies were conducted in
346 children (10, 14). As part of their investigation into the effect of aflatoxin exposure on immune
347 function in Gambian children aged between 6 and 9 years Turner et al. (10) investigated the
348 correlation between vitamins A (a- and b-carotene and lycopene) and C with AF-alb levels.
349 Vitamin C was the only micronutrient that demonstrated an inverse relationship with AF-alb (P
350 = 0.01). A study conducted by Gong et al. (14) that examined the relationship between aflatoxin
351 exposure during the post weaning period and growth faltering, measured vitamin A and zinc
352 levels to assess if they were potential confounding factors. No significant correlations between
353 vitamin A and zinc with AF-alb levels were observed. A more recent cross-sectional study (86)
354 of 147 Ghanaian adults found a significant negative correlation between AF-alb levels and
355 vitamin A concentrations in plasma samples (-0.20; $p < 0.05$). Participants with high AF-alb

356 levels (>0.80 pmol/mg albumin) had a 2.6-fold greater risk of having lower vitamin A levels
357 after adjusting for potential confounding factors (odds ratio = 2.61; CI = 1.03- 6.58; P = 0.04).
358 Tang et al., (87) found similar results in another sample of 507 Ghanaian adults. A correlation
359 analysis revealed significant negative correlations between AFB1-albumin adducts and vitamin
360 A ($r = -0.110$; $p = 0.013$) and vitamin E ($r = -0.149$; $p < 0.001$).

361

362 It is very difficult to draw specific conclusions based on the above evidence. Firstly, only a
363 small number of studies have been identified that have examined the relationship between
364 aflatoxin exposure and micronutrient deficiency in human subjects. Secondly, two studies
365 found no associations between vitamin A and AF-alb levels, whereas two studies did, indicating
366 that this relationship is not consistent across studies. Furthermore, the temporal relationship has
367 not yet been investigated as the above studies were cross-sectional; although Gong et al. (14)
368 was a longitudinal study, the micronutrients measured were only considered as potential
369 confounding factors for the relationship between aflatoxin exposure and impaired child growth,
370 and further explorations of these variables were not carried out. It is, consequently, still
371 unknown whether aflatoxin exposure exacerbates micronutrient deficiencies and if this
372 contributes to impaired child growth, which previous researchers have advocated (80). Future
373 studies opting for a longitudinal or experimental (RCT) design are warranted to help establish
374 whether a temporal relationship exists.

375

376 *Possible mechanisms for aflatoxin's effects on growth*

377 It has been hypothesized that aflatoxin may affect child growth through one or more of three
378 mechanisms; 1) by contributing to enteropathy, 2) immune suppression and 3) modulating the
379 insulin-like growth factor (IGF) pathway through liver toxicity (79, 80). Enteropathy is a

380 frequent condition observed in babies in Africa, and may be partly attributable to aflatoxin
381 related toxic damage to the intestine epithelium, which leads to further “leak” of nutrients, i.e.
382 aflatoxin exacerbates the reduction of nutrient uptake in an environment where undernutrition
383 is already rife. The immune suppression effect of aflatoxin, for which there is a lot of evidence
384 in animal species (88), and increasing evidence in humans (10-12), could enhance susceptibility
385 to infections such as those causing diarrhoea, which would reduce nutrient uptake. Liver
386 toxicity due to chronic aflatoxin exposure may damage the production of Insulin like Growth
387 Factor pathway proteins (IGFs) in the liver, leading to reduced IGFs in circulation and an
388 adverse impact on child growth. A recent in vitro study using human liver cells demonstrated
389 that aflatoxin down-regulated IGFs genes and protein levels in a dose-dependent manner (89).
390 In agreement with this result, both IGF1 and IGFBP3 levels were found to be inversely
391 correlated with AF-alb biomarker in Kenyan schoolchildren. Although the effect of aflatoxin
392 on IGFs only explained about 16% of total effect of aflatoxin on child growth, given the
393 complex causes of child stunting, the data provides preliminary evidence that aflatoxin-induced
394 changes in IGFs could contribute to growth impairment where aflatoxin exposure is high (89).

395

396 **Aflatoxin co-exposure with other mycotoxins on child undernutrition**

397 Many countries in sub-Saharan Africa have a largely maize-based diet for both weaning food
398 and family food. It has been noted that groundnuts, although often having higher incidence and
399 levels of aflatoxin contamination than maize, rarely cause aflatoxicosis. Major aflatoxicosis
400 often occurs in populations with high maize consumption. This is partly because maize is a
401 major component of the diet and is consumed in much larger amounts than groundnuts. Another
402 possibility is that another mycotoxin, fumonisin, often co-occurs with aflatoxin in maize in
403 these regions (90-92) and it is hypothesized that the co-exposure may greatly enhance aflatoxin

404 toxicity, both acute (aflatoxicosis), and chronic such as the childhood hepatomegaly reported
405 in Kenya (9).

406 Weaning food was found to be frequently co-contaminated with aflatoxin and fumonisin in
407 Tanzania, and fumonisin exposure by dietary assessment has been reported to be associated
408 with child stunting and linear growth in Tanzania (93). One hundred and sixty-six children
409 (aged 6-14 months) from representative regions in Tanzania were studied longitudinally over
410 one year to examine exposure to both mycotoxins and its impact on child growth. AF-alb levels
411 tripled during the first 6 months, and further doubled during the second 6 months, with mean
412 levels of 4.7, 12.9 and 23.5 pg/mg, respectively. Fumonisin exposure measured using urinary
413 FB1 biomarker was exceedingly high at both maize harvest seasons but with a lower level
414 observed at 6 months after harvest, reflecting a field mycotoxin contamination pattern (92).
415 Urinary FB1 at recruitment were negatively associated with HAZ at both 6 months and 12
416 months from recruitment. Mean levels of urinary FB1 had an inverse association with HAZ at
417 12 months from recruitment and length velocity. The negative association between AF-alb and
418 HAZ was not significant, possibly owing to study power limitation. These data show that
419 fumonisin may contribute to child growth impairment and highlight the potential role of co-
420 contamination with aflatoxin and fumonisin. More recently, Srey et al. (94) reported exposure
421 to dietary deoxynivalenol (DON), another mycotoxin with known growth inhibition in animals,
422 also occurs in these children, in agreement with food based exposure analysis in Tanzania (95).
423 This suggests that the children are frequently exposed to the three mycotoxins, all of which may
424 have an impact on growth faltering.

425

426 An increasing number of recent studies have reported multi-mycotoxin exposure in different
427 populations including some African groups (96-98). The methodology applied in these studies

428 typically involves simultaneous measurement of multiple mycotoxins using advanced LC-
429 MS/MS technique, and this offers great advantages as it gives useful data on multi-mycotoxin
430 exposure in a population. At present validation of the approach when applied to health outcomes
431 is in its infancy. It was evident from these studies that firstly, multiple mycotoxins co-exist in
432 staple foods such as maize and their by-products (96) and secondly, human populations in
433 Africa are co-exposed to proportionally high levels of multi-mycotoxins (97, 98). Ediage et al.
434 (97) cross-sectional study found no association between stunting, wasting or underweight in
435 children aged under five, although multiple mycotoxins were found in urine samples. Whilst
436 the multi-mycotoxin measurements require further validation, these studies provide a preview
437 of the co-exposure issue and with time more will be revealed, adding further complexity to the
438 health risk studies. How to assess the health outcomes associated with multiple toxins will thus
439 be a critical challenge ahead and this will lead to a new era of multiple toxins exposure
440 assessment methodology development.

441

442 **Conclusions**

443 Aflatoxin exposure is highly prevalent in developing countries; often this co-exists with
444 malnutrition, enteropathy, and infectious disease in young children. The fact that over 90% of
445 samples from young children from West Africa had detectable AF-alb, in contrast to less than
446 1% in the developed world clearly demonstrates a huge public health burden associated with
447 aflatoxin in sub-Saharan Africa. The greatest challenge ahead is not only to understand how
448 these problems may interactively impact on child stunting, but more importantly to explore the
449 most effective intervention method for child undernutrition, and eventually to reduce child
450 mortality. Many supplementation trials targeting the major micronutrient deficiencies aimed at
451 improving child growth have failed to produce a significant positive effect. We believe that the

452 high levels of aflatoxin exposure in these populations are likely to be exacerbating the problems
453 posed by child undernutrition and that future nutrition interventions should take aflatoxin
454 exposure into account. The most effective outcomes are likely to be produced by an attack on
455 two fronts- reduction of aflatoxin exposure and improvement in nutritional status.

Acknowledgments

The authors declare that there are no conflicts of interest. All authors were involved in conducting the literature search, writing and editing the article. All authors read and approved the final article.

References

- (1) Black RE, Victora CG, Walker SP, Bhutta ZA, Christian P, de Onis M, Ezzati M, Grantham-McGregor S, Katz J, Martorell R et al. Maternal and child undernutrition and overweight in low-income and middle-income countries. *Lancet* 2013;382:427-451. doi: 10.1016/S0140-6736(13)60937-X.
- (2) Shrimpton R, Victoria CG, de Onis M, Lima RC, Blossner M, Clugston G. Worldwide timing of growth faltering: Implications for nutritional interventions. *Pediatrics* 2001;107:E75.
- (3) Victora CG, de Onis M, Hallal PC, Blossner M, Shrimpton R. Worldwide timing of growth faltering: revisiting implications for interventions. *Pediatrics* 2010;125:e473-e480. doi: 10.1542/peds.2009-1519.
- (4) Bhutta ZA, Ahmed T, Black RE, Cousens S, Dewey K, Giugliani E, Haider BA, Kirkwood B, Morris SS, Sachdev HP et al. What works? Interventions for maternal and child undernutrition and survival. *Lancet* 2008;371:417-440. doi: 10.1016/S0140-6736(07)61693-6.
- (5) Pitt JI, Taniwaki MH, Cole MB. Mycotoxin production in major crops as influenced by growing, harvesting, storage and processing, with emphasis on the achievement of Food Safety Objectives. *Food Control* 2013;32:205-215. doi:10.1016/j.foodcont.2012.11.023
- (6) Mutegi CK, Wagacha JM, Christie ME, Kimani J, Karanja L. Effect of storage conditions on quality and aflatoxin contamination of peanuts (*Arachis hypogaea* L.). *Int J AgriScience* 2013;3:746-758.
- (7) Azziz-Baumgartner E, Lindblade K, Giesecker K, Rogers HS, Kieszak S, Njapau H, Schleicher R, McCoy LF, Misore A, DeCock K et al. Case-control study of an acute aflatoxicosis outbreak, Kenya, 2004. *Environ Health Perspect* 2005;113:1779-1783.
- (8) International Agency for Research on Cancer (IARC). Some traditional herbal medicines, some mycotoxins, naphthalene and styrene. IARC monographs on the evaluation of carcinogenic risks to humans 2002;82:171-300
- (9) Gong YY, Wilson S, Mwatha JK, Routledge MN, Castelino JM, Zhao B, Kimani G, Vennervald BJ, Dunne DW, Wild CP. Aflatoxin exposure may contribute to chronic hepatomegaly in Kenyan school children. *Environ Health Perspect* 2012;120:893-896. doi: 10.1289/ehp.1104357
- (10) Turner PC, Moore SE, Hall AJ, Prentice AM, Wild CP. Modification of immune function through exposure to dietary aflatoxin in Gambian children. *Environ Health Perspect* 2003;111:217-220.
- (11) Jiang Y, Jolly PE, Ellis WO, Wang JS, Phillips TD, Williams JH. Aflatoxin B1 albumin adduct levels and cellular immune status in Ghanaians. *Int Immunol* 2005;17:807-814.
- (12) Jiang Y, Jolly PE, Preko P, Wang JS, Ellis WO, Phillips TD, Williams JH. Aflatoxin-related immune dysfunction in health and in human immunodeficiency virus disease. *Clin Dev Immunol* 2008;790309. doi: 10.1155/2008/790309.

- (13) Gong YY, Cardwell K, Hounsa A, Egal S, Turner P, Hall AJ, Wild CP. Dietary aflatoxin exposure and impaired growth in young children from Benin and Togo: cross sectional study. *BMJ* 2002;325:20-21.
- (14) Gong YY, Hounsa A, Egal S, Turner PC, Sutcliffe AE, Hall AJ, Cardwell K, Wild CP. Postweaning exposure to aflatoxin results in impaired child growth: a longitudinal study in Benin, West Africa. *Environ Health Perspect* 2004;112:1334-1338.
- (15) Wild CP, Gong YY. Mycotoxins and human disease: a largely ignored global health issue. *Carcinogenesis* 2010;31:71-82. doi: 10.1093/carcin/bgp264.
- (16) Ahmed T, Hossain M, Sanin KI. Global burden of maternal and child undernutrition and micronutrient deficiencies. *Ann Nutr Metab* 2012;61:8-17. doi: 10.1159/000345165
- (17) World Health Organization. Nutrition for health and development: a global agenda for combating malnutrition. Geneva: World Health Organization, 2002.
- (18) World health Organization Working Group. Use and interpretation of anthropometric indicators of nutritional status. *Bull WHO* 1986; 64:929-941.
- (19) Lenters LM, Wazny K, Webb P, Ahmed T, Bhutta ZA. Treatment of severe and moderate acute malnutrition in low-and middle-income settings: a systematic review, meta-analysis and Delphi process. *BMC Public Health* 2013;13:S23. doi: 10.1186/1471-2458-13-S3-S23.
- (20) UNICEF, WHO, World Bank. Levels and trends in child malnutrition. Joint child malnutrition estimates. New York, NY: United Nations International Children's Fund; Geneva: World Health Organization; Washington DC: World Bank, 2012.
- (21) Wessells KR, Brown KH. Estimating the global prevalence of zinc deficiency: results based on zinc availability in national food supplies and the prevalence of stunting. *PLoS One* 2012;7:e50568. doi: 10.1371/journal.pone.0050568.
- (22) Ahmed T, Hossain M, Sanin KI. Global Burden of Maternal and Child Undernutrition and Micronutrient Deficiencies. *Ann Nutr Metab* 2012;61:8-17. doi: 10.1159/000345165.
- (23) Lindenmayer GW, Stoltzfus RJ, Prendergast AJ. Interactions between zinc deficiency and environmental enteropathy in developing countries. *Adv Nutr* 2014;5:1-6. doi: 10.3945/an.113.004838.
- (24) Prasad AS. Discovery of human zinc deficiency: its impact on human health and disease. *Adv Nutr* 2013;4:176-190. doi: 10.3945/an.112.003210.
- (25) de Benoist B, Darnton-Hill I, Davidsson L, Fontaine O, Hotz C. Conclusions of the joint WHO/UNICEF/IAEA/IZiNCG interagency meeting on zinc status indicators. *Food Nutr Bull* 2007;28:S480-S484.
- (26) Brown KH, Peerson JM, Rivera J, Allen LH. Effect of supplemental zinc on the growth and serum zinc concentrations of prepubertal children: a meta-analysis of randomized controlled trials. *Am J Clin Nutr* 2002;75:1062-1071.

- (27) Brown KH, Peerson JM, Baker SK, Hess SY. Preventive Zinc Supplementation among infants, preschoolers, and older prepubertal children. *Food Nutr Bull* 2009;30:12-40.
- (28) Ramakrishnan U, Nguyen P, Martorell R. Effects of micronutrients on growth of children under 5 y of age: meta-analyses of single and multiple nutrient interventions. *Am J Clin Nutr* 2009;89:191-203. doi: 10.3945/ajcn.2008.26862.
- (29) Imdad A, Bhutta ZA. Effect of preventive zinc supplementation on linear growth in children under 5 years of age in developing countries: a meta-analysis of studies for input to the lives saved tool. *BMC Public Health* 2011;11:S22. doi: 10.1186/1471-2458-11-S3-S22.
- (30) UNICEF, UNU, WHO. Iron Deficiency Anaemia. Assessment, prevention and control. A guide for programme managers. Geneva, World Health Organization, 2001.
- (31) Stevens GA, Finucane MM, De-Regil LM, Paciorek CJ, Flaxman SR, Branca F, Pena-Rosas JP, Bhutta ZA, Ezzati M, Nutrition Impact Model Study Group (Anaemia). Global, regional, and national trends in haemoglobin concentration and prevalence of total and severe anaemia in children and pregnant and non-pregnant women for 1995-2011: a systematic analysis of population-representative data. *Lancet Glob Health* 2013;1:e16-e25. doi: 10.1016/S2214-109X(13)70001-9.
- (32) Zimmermann MB, Hurrell RF. Nutritional iron deficiency. *Lancet* 2007;370:511-520.
- (33) Ramakrishnan U, Aburto N, McCabe G, Martorell R. Multimicronutrient interventions but not vitamin A or iron interventions alone improve child growth: results of 3 meta-analyses. *J Nutr* 2004;134:2592-2602.
- (34) Sachdev H, Gera T, Nestel P. Effect of iron supplementation on physical growth in children: systematic review of randomised controlled trials. *Public Health Nutr* 2006;9:904-920.
- (35) Pasricha SR, Hayes E, Kalumba K, Biggs BA. Effect of daily iron supplementation on health in children aged 4-23 months: a systematic review and meta-analysis of randomised controlled trials. *Lancet Glob Health* 2013;1:e77-e86. doi: 10.1016/S2214-109X(13)70046-9.
- (36) Thompson J, Biggs B, Pasricha SR. Effects of daily iron supplementation in 2- to 5-Year-Old Children: systematic review and meta-analysis. *Pediatrics* 2013;131:739-753. doi: 10.1542/peds.2012-2256.
- (37) Low M, Farrell A, Biggs BA, Pasricha SR. Effects of daily iron supplementation in primary-school-aged children: systematic review and meta-analysis of randomized controlled trials. *CMAJ* 2013;185:E791-E802. doi: 10.1503/cmaj.130628.
- (38) Sachdev H, Gera T, Nestel P. Effect of iron supplementation on mental and motor development in children: systematic review of randomised controlled trials. *Public Health Nutr* 2005;8:117-132.
- (39) Szajewska H, Rusczyński M, Chmielewska A. Effects of iron supplementation in nonanemic pregnant women, infants, and young children on the mental performance and

psychomotor development of children: a systematic review of randomized controlled trials. *Am J Clin Nutr* 2010;91:1684-1690. doi: 10.3945/ajcn.2010.29191.

(40) World Health Organization. Global prevalence of vitamin A deficiency in populations at risk 1995-2005: WHO global database on vitamin A deficiency. Geneva, World Health Organization, 2009.

(41) Gazala E, Sarov B, HersHKovitz E, Edvardson S, Sklan D, Katz M, Friger M, Gorodischer R. Retinol concentration in maternal and cord serum: its relation to birth weight in healthy mother-infant pairs. *Early Hum Dev* 2003;71(1):19-28.

(42) Tielsch JM, Rahmathullah L, Katz J, Thulasiraj RD, Coles C, Sheeladevi S, Prakash K. Maternal night blindness during pregnancy is associated with low birthweight, morbidity, and poor growth in south India. *J Nutr* 2008;138:787-792.

(43) Thorne-Lyman AL, Fawzi WW. Vitamin A and carotenoids during pregnancy and maternal, neonatal and infant health outcomes: a systematic review and meta-analysis. *Paediatr Perinat Epidemiol* 2012;26:36-54. doi: 10.1111/j.1365-3016.2012.01284.x.

(44) Rivera JA, Hotz C, Gonzalez-Cossio T, Neufeld L, Garcia-Guerra A. The effect of micronutrient deficiencies on child growth: a review of results from community-based supplementation trials. *J Nutr* 2003;133:4010S-4020S.

(45) Mahawithange ST, Kannanga KK, Wickremasinghe R, Chandrika UG, Jansz ER, Karunaweera ND, Wickremasinghe AR. Impact of vitamin A supplementation on health status, and absenteeism of school children in Sri Lanka. *Asia Pac J Clin Nutr* 2007;16:94-102.

(46) Chhagan MK, Van den Broeck J, Luabeya KA, Mpontshane N, Tomkins A, Bennis ML. Effect on longitudinal growth and anemia of zinc or multiple micronutrients added to vitamin A: a randomized controlled trial in children aged 6-24 months. *BMC Public Health* 2010;10:145. doi: 10.1186/1471-2458-10-145.

(47) Lima AA, Soares AM, Lima NL, Mota RM, Maciel BL, Kvalsund MP, Barret LJ, Fitzgerald RP, Blaner WS, Guerrant RL. Effects of vitamin A supplementation on intestinal barrier function, growth, total parasitic, and specific *Giardia* spp infections in Brazilian children: a prospective randomized, double-blind, placebo-controlled trial. *J Pediatr Gastroenterol Nutr* 2010;50:309-315. doi: 10.1097/MPG.0b013e3181a96489.

(48) Routledge MN, Gong YY. Developing biomarkers of human exposure to mycotoxins. In: De Saeger S, ed. Determining mycotoxins and mycotoxigenic fungi in food and feed. Cambridge, UK: Woodhead Publishing Limited, 2011:225-244.

(49) Chapot B, Wild CP. ELISA for quantification of aflatoxin-albumin adducts and their application to human exposure assessment. In: Warhol, M, van Velzen D, Bullock GR, eds. Techniques in diagnostic pathology. San Diego CA, USA: Academic Press 1991;2:135-155.

(50) Wild CP, Hudson GJ, Sabbioni G, Chapot B, Hall AJ, Wogan GN, Whittle H, Montesano R, Groopman JD. Dietary intake of aflatoxins and the level of albumin-bound aflatoxin in peripheral blood in the Gambia, West Africa. *Cancer Epidemiol Biomarkers Prev* 1992;1:229-234.

- (51) Routledge MN, Kimanya ME, Shirima CP, Wild CP, Gong YY. Quantitative correlation of aflatoxin biomarker with dietary intake of aflatoxin in Tanzanian children. *Biomarkers* 2014;19:430-435. doi: 10.3109/1354750X.2014.924998.
- (52) Maxwell SM, Apegyei F, De Vries HR, Mwanmut DD, Hendrickse RG. Aflatoxins in breast milk, neonatal cord blood and sera of pregnant women. *J Toxicol. Toxin Rev* 1989;8:19-29.
- (53) Denning DW, Allen R, Wilkinson AP, Morgan MR. Transplacental transfer of aflatoxin in humans. *Carcinogenesis* 1990;11:1033-1035.
- (54) Wild CP, Rasheed FN, Jawla MF, Hall AJ, Jansen LA, Montesano R. In-utero exposure to aflatoxin in West Africa. *Lancet* 1991;337:1602.
- (55) Jonsyn FE, Maxwell SM, Hendrickse RG. Human fetal exposure to ochratoxin A and aflatoxins. *Ann Trop Paediatr* 1995;15:3-9.
- (56) Turner PC, Collinson AC, Cheung YB, Gong YY, Hall AJ, Prentice AM, Wild CP. Aflatoxin exposure in utero causes growth faltering in Gambian infants. *Int J Epidemiol* 2007;36:1119-1125.
- (57) de Vries HR, Maxwell SM, Hendrickse RG. Foetal and neonatal exposure to Aflatoxins. *Acta Paediatr Scand* 1989;78:373-378.
- (58) Abdulrazzaq YM, Osman N, Yousif ZM, Trad O. Morbidity in neonates of mothers who have ingested aflatoxins. *Ann Trop Paediatr* 2004;24:145-151.
- (59) Shuaib FM, Jolly PE, Ehiri JE, Yatich N, Jiang Y, Funkhouser E, Person SD, Wilson C, Ellis WO, Wand JS et al. Association between birth outcomes and aflatoxin B-1 biomarker blood levels in pregnant women in Kumasi, Ghana. *Trop Med Int Health* 2010;15:160-167. doi: 10.1111/j.1365-3156.2009.02435.x.
- (60) Castelino JM, Dominguez-Salas P, Routledge MN, Prentice AM, Moore SE, Hennig BJ, Wild CP, Gong YY. Seasonal and gestation stage associated differences in aflatoxin exposure in pregnant Gambian women. *Trop Med Int Health* 2014;19:348-354. doi: 10.1111/tmi.12250.
- (61) Gong YY, Egal S, Hounsa A, Turner PC, Hall AJ, Cardwell KF, Wild CP. Determinants of aflatoxin exposure in young children from Benin and Togo, West Africa: the critical role of weaning. *Int J Epidemiol* 2003;32:556-562.
- (62) Khlangwiset P, Shephard GS, Wu F. Aflatoxins and growth impairment: a review. *Crit Rev Toxicol* 2011;41:740-755. doi: 10.3109/10408444.2011.575766
- (63) Mann CJ. Observational research methods. Research design II: cohort, cross sectional, and case-control studies. *Emerg Med J* 2003;20:54-60.
- (64) Hendrickse RG. Of sick turkeys, kwashiorkor, malaria, perinatal mortality, heroin addicts and food poisoning: research on the influence of aflatoxins on child health in the tropics. *Ann Trop Med Parasitol* 1997;91:787-793.

- (65) Hendrickse RG, Coulter JB, Lamplugh SM, Macfarlane SB, Williams TE, Omer MI, Suliman GI. Aflatoxins and kwashiorkor: a study in Sudanese children. *Br Med J* 1982;285:843-846.
- (66) Lamplugh SM, Hendrickse RG. Aflatoxins in the livers of children with kwashiorkor. *Ann Trop Paediatr* 1982;2:101-104.
- (67) Apeageyi F, Lamplugh SM, Hendrickse RG, Affram K, Lucas S. Aflatoxins in the livers of children with kwashiorkor in Ghana. *Trop Geogr Med* 1986;38:273-6.
- (68) Coulter J, Hendrickse RG, Lamplugh SM, Macfarlane SB, Moody JB, Omer MI, Suliman GI, Williams TE. Aflatoxins and kwashiorkor: clinical studies in Sudanese children. *Trans R Soc Trop Med Hyg* 1986;80:945-951.
- (69) de Vries HR, Lamplugh SM, Hendrickse RG. Aflatoxins and kwashiorkor in Kenya: a hospital based study in a rural area of Kenya. *Ann Trop Paediatr* 1987;7:249-251.
- (70) de Vries HR, Maxwell SM, Hendrickse RG. Aflatoxin excretion in children with kwashiorkor or marasmic kwashiorkor - a clinical investigation. *Mycopathologia* 1990;110:1-9.
- (71) Househam KC, Hundt HK. Aflatoxin exposure and its relationship to kwashiorkor in African children. *J Trop Pediatr* 1991;37:300-302.
- (72) Ramjee G, Berjak P, Adhikari M, Dutton MF. Aflatoxins and kwashiorkor in Durban, South Africa. *Ann Trop Paediatr* 1992;12:241-247.
- (73) Adhikari M, Ramjee G, Berjak P. Aflatoxin, kwashiorkor, and morbidity. *Nat Toxins* 1994;2:1-3.
- (74) Oyelami OA, Maxwell SM, Adelusola KA, Aladekoma TA, Oyelese AO. Aflatoxins in the lungs of children with kwashiorkor and children with miscellaneous diseases in Nigeria. *J Toxicol Environ Health* 1997;51:623-628.
- (75) Oyelami OA, Maxwell SM, Adelusola KA, Aladekoma TA, Oyelese AO. Aflatoxins in autopsy kidney specimens from children in Nigeria. *J Toxicol and Environ Health A* 1998;55:317-323.
- (76) Hatem NL, Hassab HM, Abd Al-Rahman EM, El-Deeb SA, El-Sayed Ahmed RL. Prevalence of aflatoxins in blood and urine of Egyptian infants with protein-energy malnutrition. *Food Nutr Bull* 2005;26:49-56.
- (77) Tchana AN, Moundipa PF, Tchouanguép FM. Aflatoxin contamination in food and body fluids in relation to malnutrition and cancer status in Cameroon. *Int J Environ Res Public Health* 2010;7:178-188. doi: 10.3390/ijerph7010178.
- (78) Onyemelukwe GC, Ogoina D, Ibiam GE, Ogbadu GH. Aflatoxins in body fluids and food of Nigerian children with protein-energy malnutrition. *AJFAND* 2012;12:6553-6566.

- (79) Gong YY, Turner PC, Hall AJ, Wild CP. Aflatoxin Exposure and Impaired Child Growth in West Africa: An Unexplored International Public Health Burden? In: Leslie JF, Bandyopadhyay R, Visconti A, eds. *Mycotoxins: Detection Methods, Management, Public Health, and Agricultural Trade*. Wallingford, UK: CABI, 2008:53-66.
- (80) Smith LE, Stoltzfus RJ, Prendergast A. Food chain mycotoxin exposure, gut health, and impaired growth: a conceptual framework. *Adv Nutr* 2012;3:526-531. doi: 10.3945/an.112.002188.
- (81) Williams JH, Phillips TD, Jolly PE, Stiles JK, Jolly CM, Aggarwal D. Human aflatoxicosis in developing countries: a review of toxicology, exposure, potential health consequences, and interventions. *Am J Clin Nutr* 2004;80:1106-1122.
- (82) Pimpukdee K, Kubena LF, Bailey CA, Huebner HJ, Afriyie-Gyawu E, Phillips TD. Aflatoxin-induced toxicity and depletion of hepatic vitamin A in young broiler chicks: protection of chicks in the presence of low levels of NovaSil PLUS in the diet. *Poult Sci* 2004;83:737-744.
- (83) Glahn RP, Beers KW, Bottje WG, Wideman RF, Huff WE, Thomas W. Aflatoxicosis alters avian renal function, calcium, and vitamin D metabolism. *J Toxicol Environ Health* 1991;34:309-321.
- (84) Harvey RB, Kubena LF, Elissalde MH. Influence of vitamin E on aflatoxicosis in growing swine. *Am J Vet Res* 1994;55:572-577.
- (85) Mocchegiani E, Corradi A, Santarelli L, Tibaldi A, DeAngelis E, Borghetti P, Bonomi A, Fabris N, Cabassi E. Zinc, thymic endocrine activity and mitogen responsiveness (PHA) in piglets exposed to maternal aflatoxicosis B1 and G1. *Vet Immunol Immunopathol* 1998;62:245-260.
- (86) Obuseh FA, Jolly PE, Kulczycki A, Ehiri J, Waterbor J, Desmond RA, Preko PO, Jiang Y, Piyathilake CJ. Aflatoxin levels, plasma vitamins A and E concentrations, and their association with HIV and hepatitis B virus infections in Ghanaians: a cross-sectional study. *J Int AIDS Soc* 2011;14:53. doi: 10.1186/1758-2652-14-53.
- (87) Tang L, Xu L, Afriyie-Gyawu E, Liu W, Wang P, Tang Y, Wang Y, Huebner HJ, Ankrah NA, Ofori-Adjei D et al. Aflatoxin-albumin adducts and correlation with decreased serum levels of vitamins A and E in an adult Ghanaian population. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess* 2009;26:108-118. doi: 10.1080/02652030802308472.
- (88) Bondy GS, Pestka JJ. Immunomodulation by fungal toxins. *J Toxicol and Environ Health B Crit Rev* 2000;3:109-143.
- (89) Castelino JM, Routledge MN, Wilson S, Dunne DW, Mwatha JK, Gachuhi K, Wild CP, Gong YY. Aflatoxin exposure is inversely associated with IGF1 and IGF1BP3 levels in vitro and in Kenyan schoolchildren. *Mol Nutr Food Res* 2014. doi: 10.1002/mnfr.201300619
- (90) Kimanya ME, De Meulenaer B, Tiisekwa B, Ndomondo-Sigonda M, Devlieghere F, Van Camp J, Kolsteren P. Co-occurrence of fumonisins with aflatoxins in home-stored maize for

human consumption in rural villages of Tanzania. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess* 2008;25:1353-1364. doi: 10.1080/02652030802112601.

(91) Kimanya ME, De Meulenaer B, Kolsteren PW. The Correlation between fumonisins exposure and growth retardation among infants in Rombo, Tanzania. *Ann Nutr Metab* 2009;55:614-614.

(92) Shirima CP, Kimanya ME, Kinabo JL, Routledge MN, Srey C, Wild CP, Gong YY. Dietary exposure to aflatoxin and fumonisin among Tanzanian children as determined using biomarkers of exposure. *Mol Nutr Food Res* 2013;57:1874-1881. doi: 10.1002/mnfr.201300116.

(93) Kimanya ME, De Meulenaer B, Roberfroid D, Lachat C, Kolsteren P. Fumonisin exposure through maize in complementary foods is inversely associated with linear growth of infants in Tanzania. *Mol Nutr Food Res* 2010;54:1659-1667. doi: 10.1002/mnfr.200900483.

(94) Srey C, Kimanya ME, Routledge MN, Shirima CP, Gong YY. Deoxynivalenol exposure assessment in young children in Tanzania. *Mol Nutr Food Res* 2014;58:1574-1580. doi: 10.1002/mnfr.201400012.

(95) Kimanya ME, Shirima CR, Magoha H, Shewiyo DH, De Meulenaer B, Kolsteren P, Gong YY. Co-exposures of aflatoxins with deoxynivalenol and fumonisins from maize based complementary foods in Rombo, Northern Tanzania. *Food Control* 2014;41:76-81.

(96) Abia WA, Warth B, Sulyok M, Krska R, Tchana A, Njobeh PB, Turner PC, Kouanfack C, Eyongetah M, Dutton M et al. Bio-monitoring of mycotoxin exposure in Cameroon using a urinary multi-biomarker approach. *Food Chem Toxicol* 2013;62:927-934. doi: 10.1016/j.fct.2013.10.003.

(97) Ediage EN, Di Mavungu JD, Song S, Sioen I, De Saeger S. Multimycotoxin analysis in urines to assess infant exposure: a case study in Cameroon. *Environ Int* 2013;57-58:50-59. doi: 10.1016/j.envint.2013.04.002.

(98) Shephard GS, Burger HM, Gambacorta L, Gong YY, Krska R, Rheeder JP, Solfrizzo M, Srey C, Sulyok M, Visconti A et al. Multiple mycotoxin exposure determined by urinary biomarkers in rural subsistence farmers in the former Transkei, South Africa. *Food Chem Toxicol* 2013;62:217-225. doi: 10.1016/j.fct.2013.08.040.

