



This is a repository copy of *Development of an amine-catalyzed regioselective synthesis of pyrroles*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/125781/>

Version: Accepted Version

Article:

Kakaawla, T.K.K. and Harrity, J.P.A. orcid.org/0000-0001-5038-5699 (2018) Development of an amine-catalyzed regioselective synthesis of pyrroles. *Organic Letters*, 20 (1). pp. 201-203. ISSN 1523-7060

<https://doi.org/10.1021/acs.orglett.7b03658>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.




eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Development of an Amine Catalyzed Regioselective Synthesis of Pyrroles

Taban K. K. Kakaawla, Joseph P. A. Harrity*

Department of Chemistry, University of Sheffield, Brook Hill, S3 7HF, UK.


Supporting Information Placeholder



ABSTRACT: A regioselective synthesis of pyrroles has been devised through the cycloaddition of 1,3-oxazolium-5-olates and enamines. Product regiochemistry is controlled by the enamine substitution pattern. Moreover, an amine catalyzed variant of this reaction allows aldehydes to be used directly as substrates for pyrrole synthesis.

Pyrroles are amongst the most important heteroaromatic compounds in the chemical sciences and they feature in a range of products, from light harvesting materials to pharmaceuticals.¹ The synthesis of pyrroles by traditional electrophilic aromatic substitution processes is complicated by their high reactivity, often rendering these processes difficult to control.² Cycloaddition reactions offer an appealing alternative to the synthesis of multi-substituted pyrroles as they can proceed under neutral conditions, often serving to regulate product regiochemistry. In this context, münchnones (1,3-oxazolium-5-olates) represent an interesting class of mesoionic heterocycles and they are known to furnish pyrroles

Scheme 1. Cycloadditions of alkynes and alkyne equivalents with münchnones




upon cycloaddition reactions with alkynes.³ These processes allow complex substitution patterns to be assembled very easily and the reactions can be quite regioselective, especially when arylacetylenes are employed.⁴ Indeed, the regiochemical insertion process favors the addition of the substituted alkyne carbon to the münchnone C4 position (Scheme 1, eq 1).

In contrast to the cycloaddition of alkynes, the corresponding reactions of münchnones and alkenes is more complicated as decarboxylation of the initial cycloadduct reveals a dipolar intermediate that can undergo several other reactions including proton transfer (with or without ring oxidation)⁵ or further cycloaddition,⁶ and these can be difficult to control.⁷ However, the use of alkenes bearing a potential leaving group LG could offer the opportunity to carry out in situ oxidation level adjustment to generate pyrroles directly. Moreover, if LG dictated reaction regiochemistry then this process would allow either regioisomer of the pyrrole to be accessed by simple choice of alkene substrate isomer, which would offer improved flexibility over the corresponding alkyne cycloadditions (Scheme 1, eq 2). The veracity of this idea has been demonstrated using β -nitrostyrenes, albeit with variable levels of regiocontrol.⁸ We report herein that enamines can offer high selectivities in münchnone cycloadditions, and that this approach successfully delivers a regiodivergent synthesis of pyrroles. Finally, we describe the development of this concept towards an amine-catalyzed cycloaddition of aldehydes.

We began our studies by exploring the reactions of enamines and münchnones, a cycloaddition that had not been previously investigated to the best of our knowledge. As münchnones are often unstable and require preparation and reaction in situ, we targeted stabilized analogs where an


Scheme 2. Regioselective cycloadditions of enamines with münchnones



electron acceptor is incorporated at C4 in order to allow us to focus explicitly on the efficiency of the cycloaddition step.⁹ As shown in Scheme 2, we were pleased to find that these substrates underwent relatively rapid cycloaddition with 1-substituted enamines to provide the corresponding pyrroles **2-5** in good yield, and as single regioisomers. Moreover, isomeric 2-substituted enamines also provided the corresponding pyrroles **6-9**, but with the opposite sense of regiochemical insertion. Indeed, examples **2** and **6**, **3** and **7** clearly highlight that pyrrole regiochemistry can be dictated by the substitution pattern of the enamine substrate, a distinct advantage over traditional alkyne cycloadditions which cannot currently be easily diverted from their innate regioselectivities.

We next opted to explore the relative reactivity of münchnones towards enamines and alkenes. We envisaged that 1-aminohexa-1,5-dienes could represent interesting substrates towards this end. Specifically, as shown in Scheme 3, a non-chemoselective cycloaddition of münchnones and 1-aminohexa-1,5-dienes could give rise to a range of products (formation of regioisomers is also possible although not explicitly shown).¹⁰ In the event, the reaction of substrates **1a,c** with 1-piperidyl hexa-1,5-diene provided pyrroles **10, 11** in reasonable yields, as single regioisomers. We noted the presence of several minor by-products in the crude material, however, products of cycloaddition with the alkene could not be identified. Therefore, although we have been unable to rule


Scheme 3. Relative reactivity of alkenes versus enamines



out the potential side reactions indicated in Scheme 3, it appears that they do not significantly compete with the enamine cycloaddition - elimination pathway.


The enamines employed in the cycloaddition reactions were accessed by simple condensation chemistry, and so the amine used at the start of the sequence was ultimately expelled during the cycloaddition as depicted in Scheme 4. This raised the possibility that an amine catalyzed variant of the cycloaddition could be devised whereby simple carbonyl compounds functioned as alkyne equivalents in the cycloaddition reaction.^{11,12} We decided to focus on acetaldehyde derivatives as these would be expected to rapidly form 2-substituted enamines in solution, thereby generating pyrroles with complementary regioselectivity to that of alkyne cycloadditions.

Scheme 4. Role of the amine in the overall transformation




Pleasingly, preliminary optimization studies showed that a range of secondary amines were effective for the catalytic cycloaddition of aldehydes, with dibenzylamine proving to be the most efficient. Exploring the scope of the reaction using stable 4-trifluoroacetyl-substituted münchnones highlighted that the process was compatible with aldehydes bearing a range of alkyl- and aryl-substituents, providing the corresponding pyrroles as single regioisomers (Scheme 5).¹³

Scheme 5. Amine catalyzed cycloaddition of aldehydes and münchnones



Extending this study to the imide substituted mesoionic reagents employed at the outset of our studies (c.f. Scheme 2) raised an interesting feature of the catalytic process. In these cases the N-tosylamide group that was essential for stabilization of the münchnone substrates was found to be cleaved during cycloaddition to provide the corresponding 1,2,3-trisubstituted pyrroles with complete regiocontrol (Scheme 6). Control reactions showed that pyrroles bearing N-tosylamides at C2 underwent conversion to the free pyrroles after heating in refluxing toluene overnight. The different outcomes highlighted in Schemes 2 and 6 are therefore due to the extended reaction times employed in the latter case.

Scheme 6. Amine catalyzed synthesis of 1,2,3-substituted pyrroles



In conclusion, we report the use of enamines in the cycloaddition - elimination reaction of münchnones for the synthesis of pyrroles. This method has several advantages over traditional alkyne cycloadditions, especially the ability of this approach to access complementary regioisomers. Moreover, this strategy allows acetaldehyde derivatives to formally function as substrates for cycloaddition via an amine catalyzed process, generating pyrroles with excellent regioselectivities. Moreover, this method access products with complementary regiochemistry to that of alkyne cycloadditions.

ASSOCIATED CONTENT

Supporting Information

Experimental procedures and characterization data. The Supporting Information is available free of charge on the ACS Publications website.

AUTHOR INFORMATION

Corresponding Author

* E-mail: j.harrity@sheffield.ac.uk

ACKNOWLEDGMENT

We are grateful to The Higher Committee for Education Development in Iraq for financial support.

REFERENCES

- (1) a) Domagala, A.; Jarosz, T.; Lapkowski, M. *Eur. J. Med. Chem.* **2015**, 100, 176; b) Walsh, C. T.; Garneau-Tsodikova, S.; Howard-Jones, A. R. *Nat. Prod. Rep.* **2006**, 23, 517.
- (2) a) Black, D. St. C. in *Comprehensive Heterocyclic Chemistry II*; Vol. 2 (Eds.: Katritzky, A. R.; Rees, C. W.; Scriven, E. F. V.), Pergamon: Oxford, 1996, p. 39; b) Joule, J. A.; Mills, K. *Heterocyclic Chemistry*, 4th ed.; Blackwell Science Ltd.: Oxford, 2000, p. 237.
- (3) Representative examples: a) Huisgen, R.; Gotthardt, H.; Bayer, H. O.; Schaefer, F. C. *Chem. Ber.* **1970**, 103, 2611; b) Reißig, H.-U.; Zimmer, R. *Angew. Chem. Int. Ed.* **2014**, 53, 9708; c) Arndtsen, B. A. *Chem. Eur. J.* **2009**, 15, 302; d) Lopchuk, J. M.; Gribble, G. W. *Tetrahedron Lett.* **2015**, 56, 3208; e) Dhawan, R.; Arndtsen, B. A. *J. Am. Chem. Soc.* **2004**, 126, 468.
- (4) Morin, M. S. T.; St-Cyr, D. J.; Arndtsen, B. A.; Krenske, E. H.; Houk, K. N. *J. Am. Chem. Soc.* **2013**, 135, 17349.
- (5) a) Peddibhotla, S.; Tepe, J. J. *J. Am. Chem. Soc.* **2004**, 126, 12776; b) Melhado, A. D.; Luparia, M.; Toste, F. D. *J. Am. Chem. Soc.* **2007**, 129, 12638.
- (6) a) Padwa, A.; Gingrich, H. L.; Lim, R. *J. Org. Chem.* **1982**, 47, 2447; b) Padwa, A.; Lim, R.; MacDonald, J. G.; Gingrich, H. L.; Kellar, S. M. *J. Org. Chem.* **1985**, 50, 3816.
- (7) Bélanger, G.; April, M.; Dauphin, É.; Roy, S. *J. Org. Chem.* **2007**, 72, 1104.
- (8) a) Lopchuk, J. M.; Hughes, R. P.; Gribble, G. W. *Org. Lett.* **2013**, 15, 5218; b) For a review, see: Gribble, G. W. in *The Chemistry of Heterocyclic Compounds, Oxazoles: Synthesis, Reactions, and Spectroscopy*; Taylor, E. C., Wipf, P., Eds.; John Wiley & Sons: Hoboken, NJ, 2003; Vol. 60, Part A, p 473.
- (9) Kakaawla, T. K. K.; Hartley, W. C.; Harrity, J. P. A. *Eur. J. Org. Chem.* **2016**, 2789.
- (10) The cage structure could arise from a cycloaddition with the alkene followed by loss of carbon dioxide and cycloaddition with the enamine. For a related process using cycloocta-1,5-diene, see: Gribble, G. W.; Sponholtz III, W. R.; Switzer, F. L.; D'Amato, F. J.; Byrne, M. P. *Chem. Commun.* **1997**, 993.
- (11) For an example of an amine catalyzed cycloaddition of azides and β -ketosulfones, see: Saraiva, M. T.; Costa, G. P.; Seus, N.; Schumacher, R. F.; Perin, G.; Paixão, M. W.; Luque, R.; Alves, D. *Org. Lett.* **2015**, 17, 6206.
- (12) For a complementary catalytic pyrrole synthesis, see: Donohoe, T. J.; Race, N. J.; Bower, J. F.; Callens, C. K. A. *Org. Lett.* **2010**, 12, 4094.
- (13) The regiochemistry of pyrroles **3**, **7**, **17** and **21** were assigned by nOe spectroscopy, and those of **19** and **20** were based on comparison with literature data. Other regiochemical assignments were made by inference.
