



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/125915/>

Version: Accepted Version

Article:

Kowal, J, Pressel, S, Duckett, JG et al. (2018) From rhizoids to roots? Experimental evidence of mutualism between liverworts and ascomycete fungi. *Annals Of Botany*, 121 (2). pp. 221-227. ISSN: 0305-7364

<https://doi.org/10.1093/aob/mcx126>

(c) 2018, The Author(s). Published by Oxford University Press on behalf of the Annals of Botany Company. All rights reserved. This is a pre-copyedited, author-produced version of an article accepted for publication in *Annals Of Botany* following peer review. The version of record Kowal, J, Pressel, S, Duckett, JG et al. (2018) From rhizoids to roots? Experimental evidence of mutualism between liverworts and ascomycete fungi. *Annals Of Botany*, 121 (2). pp. 221-227. is available online at: <https://doi.org/10.1093/aob/mcx126>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

1 **RESEARCH IN CONTEXT**

2 **From rhizoids to roots? Experimental evidence of mutualism between**
3 **liverworts and ascomycete fungi.**

4 Jill Kowal^{1,2,3}, Silvia Pressel³, Jeffrey G. Duckett³, Martin I. Bidartondo^{1,2}, Katie J.
5 Field⁴

6 ¹Imperial College London, London SW7 2AZ, UK; ²Royal Botanic Gardens, Kew,
7 Richmond TW9 3DS, UK; ³Natural History Museum, Cromwell Road, London SW7
8 5BD, UK; ⁴Centre for Plant Sciences, School of Biology, Faculty of Biological
9 Sciences, University of Leeds, Leeds LS2 9JT, UK

10 **Author for correspondence:** *Jill Kowal* - j.kowal@nhm.ac.uk

11 *Tel:* [+44 \(0\)20 7942 5276](tel:+442079425276)

12 Emails of co-authors: s.pressel@nhm.ac.uk; j.g.duckett@qmul.ac.uk;
13 m.bidartondo@imperial.ac.uk; k.field@leeds.ac.uk

14 Date of submission: May 6, 2017

15 Total word count (excluding summary, references and legends): 5873

16 Abstract: 208

17 Introduction: 881

18 Materials and Methods: 1449

19 Results and discussion: 1553

20 Acknowledgements: 36

21 No. of Figures: 3 (Fig. 1 and 2 in colour)

22 Supplemental data files: 5 (Table S1; Fig. S1,S2,S3; Methods S1)

1 **Abstract**

2 The rhizoids of leafy liverworts (Jungermanniales, Marchantiophyta) are
3 commonly colonized by the ascomycete fungus *Pezoloma ericae*. These associations
4 are hypothesized to be functionally analogous to the ericoid mycorrhizas (ErM)
5 formed by *P. ericae* with the roots of Ericaceae plants in terms of bi-directional
6 phosphorus-for-carbon exchange; however, this remains unproven. Here, we test
7 whether associations between the leafy liverwort *Cephalozia bicuspidata* and *P.*
8 *ericae* are mutualistic.

9 We measured movement of phosphorus and carbon between *C. bicuspidata*
10 and *P. ericae* using ³³P-orthophosphate and ¹⁴CO₂ isotope tracers in monoxenic
11 cultures. We also measured leafy liverwort growth, with and without *P. ericae*. We
12 present the first demonstration of nutritionally mutualistic symbiosis between a non-
13 vascular plant and an ErM-forming fungus, showing transfer of fungal-acquired P to
14 the liverwort and of liverwort-fixed C to the fungus alongside increased growth in
15 fungus-colonized liverworts.

16 Thus, this ascomycete-liverwort symbiosis can now be described as
17 mycorrhiza-like, providing further insights into ericoid mycorrhizal evolution and
18 adding Ascomycota fungi to mycorrhizal fungal groups engaging in mutualisms with
19 plants across the land plant phylogeny. As *P. ericae* also colonizes the rhizoids of
20 Schistochilaceae liverworts, which originated in the Triassic and are sister to all other
21 jungermannialean liverworts associated with fungi, our findings point toward an early
22 origin of ascomycete-liverwort symbioses, possibly predating their evolution in the
23 Ericales by some 150 MY.

24

1 **Key words:** *Cephalozia bicuspidata*; ericoid mycorrhizal fungi; liverwort; *Pezoloma*
2 *ericae*; symbiosis; carbon-for-nutrient exchange; mycorrhizas; mutualism

3

4 **Introduction**

5 Mycorrhizas are intimate symbioses formed between plant roots and soil fungi
6 that are prevalent across the globe in more than 80% of extant land plants (Smith and
7 Read, 2008). Through mycorrhizal associations, many plants engage in bi-directional
8 exchange of photosynthesis-derived plant carbon and fungal-acquired nutrients,
9 scavenged from sources beyond the root depletion zone or from soil pores too small
10 for roots to access (Smith and Read, 2008; Leake and Read, 2016). The role of
11 mycorrhizas in supplying extant land plants with nutrients, together with evidence of
12 mycorrhiza-like associations in Rhynie Chert plant fossils (Remy *et al.*, 1994) has led
13 to the hypothesis that mycorrhizal fungi likely facilitated the evolution of land plants
14 >470 Mya (Pirozynski *et al.*, 1975; Bidartondo *et al.*, 2011; Leake, 2015).
15 Additionally, recent studies have shown that the genes required for mycorrhization
16 are conserved across all land plant lineages (Wang *et al.*, 2010), including the earliest
17 diverging clade of liverworts – Haplomitriopsida (Fig. 1a, after Crandall-Stotler *et*
18 *al.*, 2009). The later-derived leafy liverworts (Jungermanniidae) have not been
19 incorporated in such analyses and a critical caveat of molecular studies is that the
20 presence of genes does not necessarily imply functional significance. Indeed, for
21 many groups of extant plants that form mycorrhizas and mycorrhiza-like associations
22 where roots are absent, knowledge regarding the physiological function of the
23 symbiosis has been severely limited (Read *et al.*, 2000, Field *et al.*, 2015a). To date,
24 amongst fungus-associated early-branching land plants (i.e. liverworts, hornworts and
25 lycophytes) nutritional mutualisms have only been demonstrated in a handful of early

1 diverging thalloid and Haplomitriopsida liverworts (Field *et al.*, 2012; 2015a; 2016),
2 (Fig. 1a), all which form symbioses with Mucoromycota fungi (Glomeromycotina
3 and/or Mucoromycotina) (Field *et al.* 2015b).

4 As far as can be established from published surveys and incidental
5 illustrations in floras (Pocock *et al.*,1984; Pocock & Duckett, 1985; Duckett *et*
6 *al.*,1991, Ligrone *et al.* 2007) rhizoidal ascomycete associations occur in far more
7 liverwort species than those containing basidiomycetes, glomeromycetes and
8 mucoromycetes (Pressel *et al.*, 2010). Following the accepted species names in the
9 Plant List (2013), we estimate that up to 1,000 late-diverging leafy liverworts are
10 likely to have ascomycete fungal symbionts, i.e. around 20% of liverwort species
11 worldwide compared to less than 100 with basidiomycetes, glomeromycetes and
12 mucoromycetes (see Table S2 for detailed breakdown). Reinforcing their symbiotic
13 rather than opportunistic status, the ascomycetes 1) induce swelling, branching and
14 septation of the rhizoids; and, 2) are ubiquitous rather than sporadic, in species where
15 they occur. In addition, transmission electron micrographs show healthy hyphae in
16 healthy host cells (Duckett *et al.*,1991; Pressel *et al.*, 2010).

17 There are several families in the leafy liverworts (Jungermanniidae),
18 including Schistochilaceae, Lepidoziaceae, Calypogeiaceae, Cephaloziaceae and
19 Cephaloziellaceae, which consistently associate with Ascomycota fungi (Fig. 1a)
20 (Pressel *et al.*, 2010). These fungal symbionts include *Pezoloma ericae* (D.J. Read)
21 Baral (syn. *Rhizoscyphus ericae* (D.J. Read) W.Y. Zhuang and Korf, 2004;
22 *Hymenoscyphus ericae* (D.J. Read) Korf and Kernan, 1983; and *Pezizella ericae* (D.J.
23 Read, 1974) (Duckett and Read, 1995; Read *et al.*, 2000; Pressel *et al.*, 2010)).
24 Notably, *Pezoloma ericae* is known to form ericoid mycorrhizas (ErM) with the roots
25 of Ericaceae plants and has previously been shown to provide nutrients to their

1 vascular plant hosts in exchange for fixed carbon (Read *et al.*, 2003; Smith and Read,
2 2008).

3 Ericaceous habitats are typically low in plant-available soil nutrients,
4 including N and/or P (Stribley and Read, 1974; Mitchell and Read, 1981; Leake *et al.*
5 1990; Bolan, 1991; Myers and Leake, 1996). The vascular plants inhabiting these
6 habitats, such as *Calluna*, *Erica*, *Rhododendron* and *Vaccinium*, grow together with
7 non-vascular plants, including the widespread leafy liverworts in the Cephaloziaceae
8 (Chambers *et al.*, 1999; Upson *et al.*, 2007) with which they ‘share’ fungal symbionts
9 (Duckett and Read, 1995; Read *et al.*, 2000). It was recently shown that that this
10 shared mycobiont can bring benefits in terms of establishment and survival to
11 ericaceous plants (Kowal *et al.*, 2015), and that the *P. ericae*-colonized liverwort may
12 serve as a source of fungal inoculum for vascular plants. It is possible that this effect
13 is driven by fungal-enhanced nutrition in liverworts and then in the vascular plant
14 species sharing the fungal symbiont, analogous to the nutritional role of arbuscular
15 mycorrhizal fungi (Glomeromycotina fungi) that associate with some thalloid
16 liverworts (Field *et al.*, 2012). It may be that *P. ericae* associates in leafy liverworts
17 play a similar role to the Glomeromycotina or Mucoromycotina fungal partners of
18 thalloid liverworts by supplementing plant phosphorus (P) assimilation (Field *et al.*,
19 2016).

20 Here, we aim to address the fundamental question of whether the ascomycete
21 fungus *P. ericae* forms mycorrhiza-like associations with leafy liverworts equivalent
22 to those formed by Glomeromycotina or Mucoromycotina fungi and thalloid
23 liverworts. We traced the movement of P from *P. ericae* fungal hyphae to *Cephalozia*
24 *bicuspidata* liverworts and the movement of carbon (C) from liverworts to the fungi
25 using isotope tracers. We determined liverwort growth responses to colonization by

1 *P. ericae* fungi by measuring the size and mass of liverworts, grown both with and
2 without *P. ericae* fungal symbionts.

3

4 **Materials and Methods**

5 We collected *Cephalozia bicuspidata* from Thursley Common, Surrey, in
6 autumn 2012 (OS grid reference SU900416). Mature sporophytes were harvested,
7 surface-sterilized and spores were cultured axenically (Duckett and Read, 1995) on
8 1.5% Phytigel™ (Sigma-Aldrich; ICP elemental analysis of Phytigel provided by
9 Sigma-Aldrich: 0.85% Ca, 0.35% Mg; 1.70% K, 0.15% P and 0.45% Na). No
10 additional nutrients were added to the culture medium. *Pezoloma ericae* was isolated
11 from the same liverwort collection and resynthesized with axenically-grown *C.*
12 *bicuspidata* using published methods (Kowal *et al.*, 2015), thus satisfying Koch's
13 postulates. Molecular identification of the fungal isolate as *P. ericae* was carried out
14 previously (Kowal *et al.*, 2015). Nomenclature for plants and fungi follows Hill *et al.*
15 (2008) and www.speciesfungorum.org, respectively.

16

17 *Fungus-to-plant phosphorus transfer*

18 We grew *C. bicuspidata* and *P. ericae* together in one compartment
19 ('liverwort and fungus') of 9 cm split-plate microcosms (Fig. S1) filled on both sides
20 of the divide ('B' in Fig. S1a) with 1.5% sterile Phytigel. Fragments of *P. ericae*
21 isolate (approximately 7 mm²) were inserted beneath the surface of the Phytigel and
22 an axenically-grown leafy liverwort stem (two per microcosm, c. 2 cm apart) was
23 gently pressed onto the surface, directly above the fungus fragment. Following
24 establishment of liverwort-fungal symbiosis, which is confined to the rhizoids (Fig.

1 1b), and growth of extraradical fungal hyphae (eleven weeks after planting), we
2 introduced 0.1 MBq $\text{H}_3^{33}\text{PO}_4$ (i.e. $0.03 \mu\text{g } ^{33}\text{P}$, specific activity $111 \text{ GBq mmol}^{-1}$;
3 Hartmann Analytics, Braunschweig, DE) into a well within the medium in the
4 contiguous compartment of the plate. Each ^{33}P -labelled well was then filled with
5 1.5% sterile Phytigel. The barrier dividing the microcosm prevented the liverwort
6 from encroaching into the compartment containing ^{33}P while fungal hyphae were able
7 to grow over the barrier and colonize the medium in both compartments (see Fig. S1).
8 We prepared a total of 16 microcosms with an additional fungus-free (control)
9 microcosm to measure non-fungal mediated diffusion of ^{33}P into liverwort tissue. We
10 also tested the effectiveness of the barriers in eight undivided microcosms containing
11 *C. bicuspidata* without fungus with ^{33}P -labelled wells placed the same distance from
12 plants as in divided microcosms. All microcosms were sealed with Parafilm 'M'
13 (Sigma), and placed in a controlled environment chamber (BDR16, Conviron,
14 Winnipeg, MB, Canada). The temperature regime was typical of late-spring/summer
15 for southeastern England with light intensity reflecting that at ground level beneath
16 canopy vegetation, similar to that experienced by the liverworts in their natural
17 environment (irradiance of $50 \mu\text{mol m}^{-2} \text{ s}^{-1}$, 12 h : 12 h, light : dark, $16^\circ \text{C} : 14^\circ \text{C}$
18 day : night, 80% RH and 440 ppm [CO_2]). After eight weeks, we removed the
19 liverworts from the microcosms, and freeze-dried all plant tissues and growth
20 medium containing fungal hyphae.

21 Between 10-30 mg of plant tissue and growth medium for each microcosm
22 was digested in 1 mL concentrated H_2SO_4 for two hours and then heated to 365°C for
23 15 minutes. After cooling, $100\mu\text{L}$ of hydrogen peroxide was added to each sample
24 before reheating to 365°C for two minutes, resulting in a clear solution. Samples
25 were diluted up to 10 mL with distilled water before 2 mL of the diluted digest

1 solution was mixed with 10 mL of the liquid scintillant Emulsify Safe (Perkin Elmer).
2 Activity of the samples was determined using liquid scintillation counting (Packard
3 TriCarb 3100, Isotech, Chesterfield, UK). The amount of ^{33}P transferred to the plants
4 by the fungus in each microcosm was calculated using equations from Cameron *et al.*
5 (2007). This figure was then adjusted for passive movement of ^{33}P from the substrate
6 via diffusion by subtracting the mean amount of ^{33}P measured in liverworts harvested
7 from fungus-free control microcosms.

8

9 *Plant-to-fungus carbon transfer*

10 We filled nine cube-shaped vessels (Sigma Magenta GA-7-3, 77 mm ×
11 77 mm × 97 mm) with Phytigel (1.5%) to a depth of 30 mm to prepare three replicate
12 microcosms with fungus only and six “complete” microcosms with liverwort and
13 fungus. A fragment of *P. ericae* isolate (ca. 7 mm²) was inserted beneath the surface
14 of the Phytigel for the fungus-only and complete microcosms. For the complete
15 microcosms, an axenically-grown leafy liverwort stem was gently pressed onto the
16 surface of the medium, directly above the fungus fragment. We maintained the plants
17 in the same controlled environment chamber and conditions as above. After six
18 months of growth, we labeled the microcosms with $^{14}\text{CO}_2$, separating the growth
19 medium from the chamber headspace with polythene (pre-cut with 25 mm diameter
20 holes for the liverworts) in order to minimize direct diffusion of $^{14}\text{CO}_2$ into the
21 substrate. The polythene was sealed with anhydrous lanolin where it met the
22 substrate and along the edges of the Magenta vessels (Fig. S2b). We generated 0.5
23 MBq of $^{14}\text{CO}_2$ gas by adding 6.8 μL ^{14}C -sodium bicarbonate to tubes in each
24 microcosm before introducing 500 μL 25% lactic acid. We allowed plants to fix
25 $^{14}\text{CO}_2$ for five hours in the middle of the day before introducing two Eppendorf tubes

1 containing 1 mL 2 M KOH into each microcosm for 30 min to trap any remaining
2 $^{14}\text{CO}_2$ from the headspace of each microcosm. We then harvested, separated and
3 freeze-dried the liverworts and Phytigel containing fungal hyphae (see Methods S1)
4 before weighing, homogenizing and determining ^{14}C content by sample oxidation
5 (Packard Sample Oxidiser) and liquid scintillation counting (Packard Tri-Carb 3100).

6 We calculated total carbon (^{12}C plus ^{14}C) fixed by the plant and transferred to
7 the fungus as a function of the total volume and CO_2 content of the vessel headspace
8 and the proportion of supplied $^{14}\text{CO}_2$ label fixed by the plants, using equations from
9 Cameron *et al.* (2008). The specific activity of the source was $2.04 \text{ TBq Mol}^{-1}$.

10 To account for carbon movement through passive diffusion, and to assess the
11 effectiveness of the polythene/lanolin barrier, total plant-fixed fungal carbon in the
12 complete microcosms was determined by subtracting the mean carbon measured in
13 the fungus from the fungus-only control microcosms from each of the complete
14 microcosms.

15

16 *Liverwort growth*

17 We measured *C. bicuspidata* growth both with and without fungal inoculation
18 looking first at changes in leafy liverwort surface area, and second at liverwort
19 biomass. In the first experiment, we introduced small fragments (c. 7 mm^2) of *P.*
20 *ericae* to 9 cm plates containing 1.5% Phytigel with axenically-grown leafy
21 liverworts (c. one year old). After fungal colonization of the liverwort rhizoids was
22 confirmed microscopically (Fig. 1b), we re-plated colonized liverworts and fungus-
23 free control liverworts individually using the same medium. Surface area
24 measurements of each liverwort stem (including leaves) were made before and after

1 six weeks using images taken with a Nikon Coolpix S10 digital camera and analyzed
2 using ImageJ software (Rasband, 1997-2012) (Fig. S3). For the second measurement
3 of the effect of *P. ericae* on *C. bicuspidata* growth, we freeze-dried and weighed the
4 liverworts from the complete microcosms used for the ¹⁴C transfer experiment
5 (above) after they were harvested at six months and compared the mean to that of
6 plants from liverwort-only microcosms (*n* = 6). This allowed us to test for
7 differences in mean plant biomass in liverwort growth with or without fungus over a
8 longer time period than the surface area growth experiment described above.
9 Liverwort stems were weighed before initial planting to ensure similarity in mass
10 between sample groups.

11

12 *Statistics*

13 After analyzing the data for normal distribution and homogeneity of variance,
14 we applied the appropriate statistical tests, i.e. parametric or non-parametric, using
15 GraphPad Prism (version 6.0h). The Mann-Whitney U test was used both for the dry
16 mass experiment and for testing barrier effect in the ³³P experiment, owing to the low
17 number of datapoints; Student's t-test was used for other datasets.

18

19 **Results and discussion**

20 *Leafy liverwort-Pezoloma ericae symbiosis is mutualistic and mycorrhiza-like*

21 In our radio-labeled liverwort microcosms, 1.786 ng g⁻¹ of ³³P was
22 assimilated from *P. ericae* into *C. bicuspidata* tissues (Fig. 2a, Table S1). There was
23 a significant difference in the liverwort ³³P content and concentration between plant-
24 only microcosms with barriers and in plant-only microcosms without barriers (Mann-

1 Whitney $U = 0$; $n_1=2$, $n_2=13$; $P = 0.019$ two-tailed (data not shown), further
2 demonstrating the efficacy of barriers in microcosms for preventing direct plant
3 access to ^{33}P -labeled wells. While we cannot exclude the possibility that the P
4 content of the Phytigel substrate used in our microcosms may have been directly
5 assimilated by the liverworts within the systems and therefore reduced plant demand
6 for fungal-acquired ^{33}P , our data unequivocally show movement of ^{33}P from the
7 fungus to the leafy liverwort.

8 The leafy liverworts transferred 0.019 ng (0.22 ng g^{-1}) of plant-fixed carbon
9 to their fungal partners (Fig. 2b); equivalent to 0.27% of the total amount of carbon
10 fixed during the labelling period (Table S1). By demonstrating unequivocal
11 exchange of fungal-acquired phosphorus and plant-fixed carbon between symbionts,
12 our results provide the first experimental evidence that associations between non-
13 vascular plants and ascomycete fungal symbionts are nutritionally mutualistic. This
14 confirms previous hypotheses based on cytological evidence (van der Heijden *et al.*
15 2015) and culturing experiments (Duckett and Read, 1991; Upson *et al.*, 2007; Kowal
16 *et al.* 2015). Additionally, liverworts resynthesized with *P. ericae* grew significantly
17 larger than liverworts grown under identical conditions without the fungus ($P < 0.01$
18 for both, Fig. 3a,b). Together with our evidence of carbon-for-nutrient exchange
19 between symbionts, it is clear that the symbiosis between non-vascular leafy
20 liverworts and their fungal partners is mutualistic and mycorrhiza-like. This lays the
21 foundation to investigate further functional differences between liverwort-ascomycete
22 partnerships and previously documented exchanges between *P. ericae* and vascular
23 plants (Pearson and Read, 1973; Read and Stribley, 1973; Upson *et al.* 2008, Kowal
24 *et al.*, 2015).

1 Our experiments aimed to uncover whether there is any movement of plant-
2 fixed carbon and fungal-acquired P between *C. bicuspidata* liverworts and *P. ericae*.
3 The evidence of ³³P transfer from fungus to plant in our experimental microcosm
4 (Fig. 2a) strongly supports a nutritional role for *P. ericae* that is particularly
5 significant given that phosphorus can be one of the most limiting nutrients in the
6 ericaceous habitats where *C. bicuspidata* grows. It is possible that the enhanced
7 growth we observed in liverworts with fungal symbionts compared to those without
8 (Fig. 3) was a result of fungal remineralization of organic compounds leached from
9 the liverwort, rather than a direct result of increased uptake of fungal-acquired P.
10 However, the provision of ³³P by the fungus, as shown here, supports the hypothesis
11 that the fungus plays a role in leafy liverwort P nutrition in natural environments.

12

13 *Evolutionary context*

14 Our demonstration that the association between a leafy liverwort and the
15 ericoid mycorrhizal fungus *P. ericae* can be reciprocally beneficial provides
16 additional clues about the evolution of ErM, and lends further weight to the idea that
17 mutualisms between leafy liverworts and ascomycete fungi pre-dated their formation
18 in the Ericales, estimated at less than 100 MYA (Brundrett, 2004). Following their
19 discovery that ascomycetes belonging to the *P. ericae* group colonize the rhizoids of
20 Schistochilaceae liverworts, thought to have originated in the Triassic (Heinrichs *et*
21 *al.*, 2007) as the sister group to all other fungus-containing lineages in the
22 Jungermanniales, Pressel *et al.* (2008) suggested that this association arose more than
23 250 MYA. Nonetheless, divergence time estimates for ascomycete lineages remain a
24 matter of debate particularly as the presumed recent origin of Leotiomycetes, the
25 clade containing *P. ericae*, is in the Cretaceous (James *et al.*, 2006; Prieto and Wedin,

1 2013). Until ever-improving phylogenetic methods integrating fossil evidence
2 (Beimforde *et al.*, 2014) provide more accurate date estimates, recent origins for the
3 association with multiple instances of host shifting between leafy liverworts and
4 Ericaceae plants, remain a plausible alternative scenario (Selosse, 2005; Pressel *et al.*,
5 2008).

6

7 *Observations on previously reported liverwort-fungal symbioses*

8 Although not directly comparable, given considerable differences in the
9 physiology of the liverwort hosts and in experimental design (i.e. axenic and edaphic
10 conditions), our finding of mycorrhizal-like associations between ascomycete fungi
11 and leafy liverworts now invites further comparisons with more ancient lineages of
12 liverworts, i.e. Haplomitriopsida (*Treubia* and *Haplomitrium*) and complex thalloid
13 liverworts (*Neohodgsonia*, *Allisonia* and *Marchantia*) and fungi - Mucoromycotina
14 and/or Glomeromycotina (Field *et al.* 2015a and 2016) (see Table S1; Fig. 1a). When
15 fungal-acquired ³³P uptake is normalized to biomass (Fig. 2a; Table S1), it appears *C.*
16 *bicuspidata* gains significantly less ³³P from its fungal partner than *Haplomitrium*,
17 *Treubia* (Mucoromycotina only), *Allisonia* and *Neohodgsonia* (Mucoromycotina and
18 Glomeromycotina), but a roughly similar amount to *Marchantia* (Glomeromycotina
19 only). *Cephalozia bicuspidata* carbon allocation (in terms of absolute amount, Fig.
20 2b) to its symbiotic fungus also points to a relatively low carbon demand by *P. ericae*
21 on its liverwort host, especially when compared to other liverwort lineages (0.27% of
22 plant-fixed carbon transferred to the symbiotic fungus vs. 2.2% - 14.2% - see Table
23 S1). This may be partially influenced by the facultative biotrophic nature of *P. ericae*
24 allowing it to gain at least some organic carbon from dead organic matter, and the
25 extent of colonisation (i.e. restricted to rhizoids vs. extensive through thallus). It is

1 possible that the fungus in our experiments was able to derive some C directly from
2 the Phytigel substrate (which contains glucose and trace levels of nutrients; see
3 materials and methods). Regardless of the phosphorus-for-carbon measurements
4 presented in Table S1, it is evident that the relative “cost” of maintaining mycorrhiza-
5 like symbioses varies between liverwort-fungal symbioses, with ErM-like
6 associations potentially requiring less liverwort investment in terms of photosynthate
7 allocation, and the liverwort-Mucoromycotina partnership requiring the most (Field *et*
8 *al.*, 2016).

9 Having established that there is a mycorrhizal-like nutritional exchange
10 between the leafy liverwort *C. bicuspidata* and its fungal symbiont *P. ericae*,
11 investigations are now needed which include fungus-to-plant N transfer and more
12 natural experimental conditions to provide a robust platform to investigate functional
13 variation in plant-fungal symbioses across evolutionary lineages and ecological
14 gradients.

15

16 *Future directions*

17 The diversity of fungal symbionts across the land plant phylogeny is
18 becoming increasingly apparent; however, there is still a relative dearth of
19 information regarding the function of plant-fungal symbioses in many clades,
20 particularly the bryophytes. The largest remaining functional knowledge gaps now
21 are the basidiomycete symbioses in thalloid Aneuraceae and leafy Scapaniaceae and
22 Arnelliaceae liverworts (Bidartondo and Duckett, 2010). In pteridophytes only two
23 pioneering studies to date have demonstrated the reciprocal exchange of plant-fixed C
24 for fungal-acquired N and P between green sporophytes of *Osmunda regalis* and

1 *Ophioglossum vulgatum* and their Glomeromycotina symbionts (Field *et al.*, 2012,
2 2015c; Pressel *et al.*, 2016).

3 Although we show transfer of C from *C. bicuspidata* to fungi *in vitro*, in
4 habitats where *C. bicuspidata* grows, often in deep shade under a canopy of
5 ericaceous plants, the fungal symbiont of *C. bicuspidata* may be getting a large
6 proportion of its organic C from surrounding vascular plants via a shared fungal
7 network, as demonstrated previously between *Betula* and the mycoheterotrophic
8 liverwort *Cryptothallus (Aneura) mirabilis* via a shared basidiomycete fungus (Read
9 *et al.*, 2000; Bidartondo *et al.*, 2003). How this affects ecosystem carbon and nutrient
10 budgets in terms of storage and cycling remains to be uncovered.

11 In conclusion, our findings provide a novel and important example of
12 mycorrhizal functioning of an additional fungal lineage with non-vascular plants.
13 Thus, our demonstration of symbiotic ErM-like functioning of *P. ericae* with a leafy
14 liverwort, now adds the Ascomycota to the list of fungal groups engaging in
15 mutualistic carbon-for-nutrient exchange across the land plant phylogeny. It has been
16 proposed that *C. bicuspidata* liverworts with ErM fungal symbionts can be used to
17 restore threatened heathlands by promoting establishment of native ericaceous plants
18 (Kowal *et al.*, 2015) that are limited by ErM inoculum availability (Diaz *et al.* 2006).
19 Our results suggest that at least one of the mechanisms underpinning the potential
20 role of *C. bicuspidata* in habitat restoration strategies is likely to be nutritional. The
21 potential ecological applications of pioneer non-vascular plants as mycorrhizal
22 reservoirs and vectors, particularly in habitat restoration, are only just becoming
23 apparent. Evidence of a mutualism between a widespread non-vascular plant and
24 ascomycete fungi that form mycorrhizas with dominant vascular plants should now
25 encourage further ecological and physiological experiments.

1

2

3 **Acknowledgements:**

4 J.K. thanks BSBI for funding for laboratory work and the University of Sheffield for

5 use of growth and analytical facilities. K.F. is supported by a BBSRC Translational

6 Fellowship (BB/M026825/1). All authors are supported by NERC (NE/N009665/1).

7 We thank the editor and reviewers for their constructive comments on an earlier

8 version of this manuscript.

Literature Cited

- Bajwa R, Read DJ. 1985.** The biology of mycorrhiza in the Ericaceae IX. Peptides as nitrogen sources for the ericoid endophyte and for mycorrhizal and non-mycorrhizal plants. *New Phytologist* **101**: 459-467.
- Bajwa R, Read DJ. 1986.** Utilization of mineral and amino N sources by the ericoid mycorrhizal endophyte *Hymenoscyphus ericae* and by mycorrhizal and non-mycorrhizal seedlings of *Vaccinium*. *Transactions of the British Mycological Society* **87**: 269-277.
- Beimforde C, Feldberg K, Nylinder S, Rikkinen J, Tuovila H, Dörfelt H, Gube M, Jackson DJ, Reitner J, Seyfullah LJ, Schmidt AR. 2014.** Estimating the Phanerozoic history of the Ascomycota lineages: combining fossil and molecular data. *Molecular Phylogenetics and Evolution* **78**: 386-398.
- Bidartondo MI, Bruns TD, Weiss M, Sergio S, Read DJ. 2003.** Specialized cheating of the ectomycorrhizal symbiosis by an epiparasitic liverwort. *Proceedings of the Royal Society B-Biological Sciences* **270**: 835-842.
- Bidartondo MI, Duckett JG. 2010.** Conservative ecological and evolutionary patterns in liverwort-fungal symbioses. *Proceedings of the Royal Society B-Biological Sciences* **277**: 485-492.
- Bidartondo MI, Read DJ, Trappe JM, Merckx V, Ligrone R, Duckett JG. 2011.** The dawn of symbiosis between plants and fungi. *Biology Letters* **7**: 574-577.
- Bolan NS. 1991.** A critical review on the role of mycorrhizal fungi in the uptake of phosphorus by plants. *Plant and Soil* **134**: 189-207.
- Brundrett MC. 2004.** Diversity and classification of mycorrhizal associations. *Biological Reviews* **79**: 473-495.
- Cameron DD, Leake JR, Read DJ. 2006.** Mutualistic mycorrhiza in orchids: evidence from plant–fungus carbon and nitrogen transfers in the green-leaved terrestrial orchid *Goodyera repens*. *New Phytologist* **171**: 405-416.
- Cameron DD, Johnson I, Leake JR, Read DJ. 2007.** Mycorrhizal acquisition of inorganic phosphorus by the green-leaved terrestrial orchid *Goodyera repens*. *Annals of Botany* **99**: 831-834.
- Cameron DD, Johnson I, Read DJ, Leake JR. 2008.** Giving and receiving: measuring the carbon cost of mycorrhizas in the green orchid, *Goodyera repens*. *New Phytologist* **180**: 176-184.
- Chambers SM, Williams PG, Seppelt RD, Cairney JWG. 1999.** Molecular identification of *Hymenoscyphus* sp. from rhizoids of the leafy liverwort *Cephaloziella exiliflora* in Australia and Antarctica. *Mycological Research* **103**: 286-288.
- Crandall-Stotler B, Stotler RE, Long, DG. 2009.** Phylogeny and classification of the Marchantiophyta. *Edinburgh Journal of Botany* **66**: 155–198.
- Diaz A, Green I, Benvenuto M, Tibbett M. 2006.** Are ericoid mycorrhizas a factor in the success of *Calluna vulgaris* heathland restoration? *Restoration Ecology* **14**: 187-195.

- Duckett JG, Read DJ. 1991.** The use of the fluorescent dye, 3,3'-dihexyloxycarbocyanine iodide, for selective staining of ascomycete fungi associated with liverwort rhizoids and ericoid mycorrhizal roots. *New Phytologist* **118**: 259-272.
- Duckett JG, Renzaglia KS & Pell K. 1991.** A light and electron microscope study of rhizoid-ascomycete associations and flagelliform axes in British hepatics: with observations on the effects of the fungus on host morphology. *New Phytologist*, **118**: 233-257.
- Duckett JG, Read DJ. 1995.** Ericoid mycorrhizas and rhizoid-ascomycete associations in liverworts share the same mycobiont - isolation of the partners and resynthesis of the associations *in-vitro*. *New Phytologist* **129**: 439-477.
- Field KJ, Cameron DD, Leake JR, Tille S, Bidartondo MI, Beerling DJ. 2012.** Contrasting arbuscular mycorrhizal responses of vascular and non-vascular plants to a simulated Palaeozoic CO₂ decline. *Nature Communication* **3**: 1-8.
- Field KJ, Pressel S, Rimington WR, Duckett JG, Bidartondo MI. 2015a.** Symbiotic options for the conquest of land. *Trends in Ecology and Evolution* **30**: 477-486.
- Field KJ, Rimington WR, Bidartondo MI, Allinson KE, Beerling DJ, Cameron DD, Duckett JG, Pressel S. 2015b.** First evidence of mutualism between ancient plant lineages (Haplomitriopsida liverworts) and Mucoromycotina fungi and its response to simulated Palaeozoic changes in atmospheric CO₂. *New Phytologist* **205**: 743-756.
- Field, KJ, Leake JR, Tille S, Allinson KE, Rimington WR, Bidartondo MI, Beerling DJ, Cameron DD. 2015c.** From mycoheterotrophy to mutualism: mycorrhizal specificity and functioning in *Ophioglossum vulgatum* sporophytes. *New Phytologist* **205**: 1492-1502.
- Field KJ, Rimington WR, Bidartondo MI, Allinson KE, Beerling DJ, Cameron DD, Duckett JG, Leake JR, Pressel S. 2016.** Functional analysis of liverworts in dual symbiosis with Glomeromycota and Mucoromycotina fungi under a simulated Palaeozoic CO₂ decline. *The ISME Journal* **10**: 1514-1526.
- Heinrichs J, Hentschel J, Wilson R, Feldberg K, Schneider H. 2007.** Evolution of leafy liverworts (Jungermanniidae, Marchantiophyta): estimates divergence times from chloroplast DNA sequences using penalized likelihood with integrated fossil evidence. *Taxon* **56**: 31-44.
- Hendry GAF, Grime JP. 1993.** *Methods in Comparative Plant Ecology: A Laboratory Manual*. London: Chapman and Hall.
- Hill MO, Blackstock TH, Long DG, Rothero GP. 2008.** *A checklist and census catalogue of British and Irish bryophytes*: Cheshire: British Bryological Society.
- Humphreys CP, Franks PJ, Rees M, Bidartondo MI, Leake JR, Beerling DJ. 2010.** Mutualistic mycorrhiza-like symbiosis in the most ancient group of land plants. *Nature Communication* **1**: 1-7.
- James TY, Kauff F, Schoch CL et al. 2006.** Reconstructing the early evolution of fungi using a six-gene phylogeny. *Nature* **443**: 818-822.

- Kerley SJ, Read DJ. 1997.** The biology of mycorrhiza in the Ericaceae .19. Fungal mycelium as a nitrogen source for the ericoid mycorrhizal fungus *Hymenoscyphus ericae* and its host plants. *New Phytologist* **136**: 691-701.
- Kowal J, Pressel S, Duckett JG, Bidartondo MI. 2015.** Liverworts to the rescue: an investigation of their efficacy as mycorrhizal inoculum for vascular plants. *Functional Ecology* **30**: 1014–1023.
- Leake JR, Shaw G, Read DJ. 1990.** The Role of Ericoid Mycorrhizas in the Ecology of Ericaceous Plants. *Agriculture Ecosystems and Environment* **29**: 237-250.
- Leake JR. 2015.** Box 12.1 Biogeochemical effects of coevolution of plants and fungi. In: Bronstein JL ed. *Mutualism*. Oxford, UK: Oxford University Press, 2.
- Ligrone R, Carafa A, Lumini E, Bianciotto V, Bonfante P, Duckett JG. 2007.** Glomeromycotean associations in liverworts: A molecular cellular and taxonomic analysis. *American Journal of Botany* **94**: 1756-1777.
- Mitchell DT, Read DJ. 1981.** Utilization of organic and inorganic phosphates by the mycorrhizal endophytes of *Vaccinium macrocarpon* and *Rhododendron ponticum*. *Transactions of the British Mycological Society* **76**: 255-260.
- Myers MD, Leake JR. 1996.** Phosphodiesterases as mycorrhizal P sources II. Ericoid mycorrhiza and the utilization of nucleic acids as a phosphorus and nitrogen source by *Vaccinium macrocarpon*. *New Phytologist* **132**: 445-451.
- Perotto S, Martino E, Abba S, Vallino M. 2012.** Genetic diversity and functional aspects of ericoid mycorrhizal fungi. In: Hock B ed. *Fungal associations, 2nd Edition*. Berlin: Springer-Verlag.
- Pearson V, Read DJ. 1973.** The biology of mycorrhiza in the Ericaceae. *New Phytologist* **72**: 1325-1331.
- Pietro M, Wedin M. 2013.** Dating the diversification of the major lineages of Ascomycota (Fungi). *PLOSone* **8**: e65576.
- Pirozynski KA, Malloch DW. 1973.** The origin of land plants: a matter of mycotropism. *Biosystems* **6**, 153–164.
- The Plant List. 2013.** Version 1.1. Published on the Internet; <http://www.theplantlist.org/> (accessed 1st January).
- Pocock K, Duckett JG, Grolle R, Mohamed MAH, Pang WC. 1984.** Branched and swollen rhizoids in hepatics from montane rain forest in Peninsular Malaya. *Journal of Bryology* **13**: 241-246.
- Pocock K, & Duckett JG. 1985.** On their occurrence of branched and swollen rhizoids in British hepatics: their relationships with the substratum and associations with fungi. *New Phytologist* **99**: 281-304.
- Pressel S, Ligrone R, Duckett JD, Davis, EC. 2008.** A novel ascomycetous endophytic association in the rhizoids of the leafy liverwort family, Schistochilaceae (Jungermanniidae, Hepaticopsida). *American Journal of Botany* **95**: 531-541.

- Pressel S, Bidartondo MI, Ligrone R, Duckett JG. 2010.** Fungal symbioses in bryophytes: New insights in the Twenty First Century. *Phytotaxa* **9**: 238-253.
- Pressel S, Bidartondo MI, Field KJ, Rimington WR, Duckett JG. 2016.** Pteridophyte fungal associations: Current knowledge and future perspectives. *Journal of Systematics and Evolution* **54**: 666-678.
- Rasband WS 1997-2012.** *ImageJ*. Maryland, USA: U. S. National Institutes of Health.
- Read DJ. 1993.** Mycorrhiza in plant communities. *Advances in Plant Pathology* **9**: 1-31.
- Read DJ, Stribley DP. 1973.** Effect of mycorrhizal infection on nitrogen and phosphorus nutrition of ericaceous plants. *Nature New Biology* **244**: 81-82.
- Read DJ, Duckett JG, Francis R, Ligrone R, Russell AJ. 2000.** Symbiotic fungal associations in 'lower' land plants. *Philosophical Transactions of the Royal Society B* **355**: 815-830.
- Read DJ, Leake JR, Perez-Moreno J. 2004.** Mycorrhizal fungi as drivers of ecosystem processes in heathland and boreal forest biomes. *Canadian Journal of Botany* **82**: 1243-1263.
- Remy W, Taylor TN, Hass H, Kerp H. 1994.** Four hundred million year old vesicular arbuscular mycorrhizae. *Proceedings of the National Academy of Sciences, USA* **91**: 11841-11843.
- Selosse M-A. 2005.** Are liverworts initiating mycorrhizas? *New Phytologist* **165**: 345-349.
- Smith SE, Read DJ. 2008.** *Mycorrhizal symbiosis*. Cambridge, UK: Academic Press.
- Stribley, DP, Read, DJ. 1974.** Biology of Mycorrhiza in Ericaceae IV. Effect of mycorrhizal infection on uptake of N-15 from labelled soil by *Vaccinium macrocarpon* Ait. *New Phytologist*, **73**, 1149-1155.
- Stribley DP, Read DJ. 1980.** The biology of mycorrhiza in the Ericaceae VII. The relationship between mycorrhizal infection and the capacity to utilize simple and complex organic nitrogen-sources. *New Phytologist* **86**: 365-371.
- Stevens PF. 2001.** *Angiosperm Phylogeny Website. Version 12, July 2012* [and more or less continuously updated since]. [WWW document] URL <http://www.mobot.org/MOBOT/research/APweb/>. [accessed 1 November 2016].
- Upton R, Read DJ, Newsham KK. 2007.** Widespread association between the ericoid mycorrhizal fungus *Rhizoscyphus ericae* and a leafy liverwort in the maritime and sub-Antarctic. *New Phytologist* **176**: 460-471.
- Walder F, van der Heijden MGA. 2015.** Regulation of resource exchange in the arbuscular mycorrhizal symbiosis. *Nature Plants* **1**: 15159.
- Wang B, Yeun LH, Xue JY, Liu Y, Ane JM, Qiu YL. 2010.** Presence of three mycorrhizal genes in the common ancestor of land plants suggests a key role of mycorrhizas in the colonization of land by plants. *New Phytologist* **186**: 514-525.

Figures

Fig. 1 (a) Liverwort phylogram showing diversity in liverwort-fungal associations (after Crandall-Stotler *et al.*, 2009) based on extensive sampling of nearly all the fungus-containing liverwort genera worldwide (Ligrone *et al.* 2007). (b) *Cephalozia bicuspidata* rhizoid resynthesized with *Pezoloma ericae* fungus (arrowed) in axenic culture.

Fig. 2 (a) Concentration of fungal-acquired ^{33}P in *Cephalozia bicuspidata* tissue (ng g^{-1}). (b) Total plant-derived carbon present in extraradical fungal mycelium after a five-hour labeling period (ng). Error bars denote 1 s.e. in both panels.

Fig. 3 (a) Change in surface area of leafy liverwort after six weeks growth with and without *Pezoloma ericae* fungal symbiont ($n = 16, 22$), $P < 0.01^{**}$ (using *t*-test). (b) Total biomass of liverworts (dry weight; g) after six months of growth with and without *P. ericae* (Mann-Whitney $U = 2$; $P = 0.0087^{**}$; $n_{1,2} = 6$ two-tailed). Error bars denote 1 s.e. in both panels.

Supplementary Figures and Data

Fig. S1 Diagrams and photographs of ^{33}P transfer experiment showing split plate experimental design set up before (a,b) and after (c,d) the fungus crossed the barrier. L – liverwort; H – hyphae; W – well and B – barrier.

Fig. S2 Diagram showing microcosm design for carbon transfer experiments. (a) *Cephalozia bicuspidata* and *Pezoloma ericae* growing on 1.5% Phytigel (b) $^{14}\text{CO}_2$ released into chamber headspace, fixed by plant then transferred into

fungal hyphae (blue). Dotted black line shows placement of polythene with lanolin seal in yellow. Polythene barrier minimizes $^{14}\text{CO}_2$ diffusion into Phytigel.

Fig. S3 Example of digitized pictures used to measure perimeter area of liverworts. (a) Control without fungus (b) Inoculated with fungus (arrow).

Methods S1 Methods for harvesting plants and fungus for ^{33}P acid digestion, liquid scintillation and nutrient budgeting.

Table S1 Summary of carbon-for-nutrient exchange between *Cephalozia bicuspidata* and *Pezoloma ericae*, alongside data from previous studies of early-diverging liverwort lineages and their fungal partners (below dotted line) (Field *et al.*, 2015a).

Table S2 Estimates of likely species numbers of liverworts worldwide with fungal symbionts.

A





