



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/126699/>

Version: Accepted Version

---

**Article:**

Akkerman, M, Kristensen, LS, Jespersen, L et al. (2017) Interaction between sodium chloride and texture in semi-hard Danish cheese as affected by brining time, dl -starter culture, chymosin type and cheese ripening. *International Dairy Journal*, 70. pp. 34-45. ISSN: 0958-6946

<https://doi.org/10.1016/j.idairyj.2016.10.011>

---

© 2016 Elsevier Ltd. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <http://creativecommons.org/licenses/by-nc-nd/4.0/>

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.


**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# Accepted Manuscript

Interaction between sodium chloride and texture in semi-hard Danish cheese as affected by brining time, DL-starter culture, chymosin type and cheese ripening

Marije Akkerman, Lise Søndergaard Kristensen, Lene Jespersen, Mia Balling Ryssel, Alan Mackie, Nick Nørreby Larsen, Ulf Andersen, Maria Kümpel Nørgaard, Mette Marie Løkke, Jean Robert Møller, Line Ahm Mielby, Barbara Vad Andersen, Ulla Kidmose, Marianne Hammershøj



PII: S0958-6946(16)30327-2

DOI: [10.1016/j.idairyj.2016.10.011](https://doi.org/10.1016/j.idairyj.2016.10.011)

Reference: INDA 4098

To appear in: *International Dairy Journal*

Received Date: 9 June 2016

Revised Date: 28 October 2016

Accepted Date: 29 October 2016

Please cite this article as: Akkerman, M., Kristensen, L.S., Jespersen, L., Ryssel, M.B., Mackie, A., Larsen, N.N., Andersen, U., Nørgaard, M.K., Løkke, M.M., Møller, J.R., Mielby, L.A., Andersen, B.V., Kidmose, U., Hammershøj, M., Interaction between sodium chloride and texture in semi-hard Danish cheese as affected by brining time, DL-starter culture, chymosin type and cheese ripening, *International Dairy Journal* (2016), doi: 10.1016/j.idairyj.2016.10.011.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Interaction between sodium chloride and texture in semi-hard Danish cheese as affected by brining**  
2 **time, DL-starter culture, chymosin type and cheese ripening**

3

4

5 Marije Akkerman<sup>a</sup>, Lise Søndergaard Kristensen<sup>b</sup>, Lene Jespersen<sup>b</sup>, Mia Balling Ryssel<sup>c</sup>, Alan Mackie<sup>d†</sup>,  
6 Nick Nørreby Larsen<sup>c</sup>, Ulf Andersen<sup>f</sup>, Maria Kümpel Nørgaard<sup>f</sup>, Mette Marie Løkke<sup>a</sup>, Jean Robert  
7 Møller<sup>a‡</sup>, Line Ahm Mielby<sup>g</sup>, Barbara Vad Andersen<sup>g</sup>, Ulla Kidmose<sup>g</sup>, Marianne Hammershøj<sup>a\*</sup>

8

9

10 <sup>a</sup> Department of Food Science, Aarhus University, Blichers Allé 20, DK-8830 Tjele, Denmark

11 <sup>b</sup> Department of Food Science, University of Copenhagen, Rolighedsvej 30, DK-1958 Frederiksberg C,  
12 Denmark

13 <sup>c</sup> Chr. Hansen A/S, Bøge allé 10-12, DK-2970 Hørsholm, Denmark

14 <sup>d</sup> Institute of Food Research, Norwich Research Park, Colney, Norwich NR47UA, United Kingdom

15 <sup>e</sup> Thise Dairy, Sundsørevej 62, DK-7870 Roslev, Denmark

16 <sup>f</sup> Arla Foods R&D, Rørdrumvej 2, DK-8220 Brabrand, Denmark

17 <sup>g</sup> Department of Food Science, Aarhus University, Kirstinebjergvej 10, DK-5792 Aarslev, Denmark

18

19

20 \* Corresponding author. Tel.: +45 8715 7974

21 E-mail address: [marianne.hammershoj@food.au.dk](mailto:marianne.hammershoj@food.au.dk) (M. Hammershøj)

22

23

24 † Present address: University of Leeds, School of Food Science & Nutrition, Leeds LS2 9JT, United  
25 Kingdom

26 ‡ Present address: Lindholmvej 13, 1tv, 8200 Aarhus N, Denmark

27

28 ABSTRACT

29

30 Reduced NaCl in semi-hard cheeses greatly affects textural and sensory properties. The interaction  
31 between cheese NaCl concentration and texture was affected by brining time (0–28 h), DL-starter cultures  
32 (C1, C2, and C3), chymosin type (bovine or camel), and ripening time (1–12 weeks). Cheese NaCl levels  
33 ranged from <0.15 to 1.90% (w/w). NaCl distribution changed during ripening; migration from cheese  
34 edge to core led to a more homogeneous NaCl distribution after 12 weeks. As ripening time increased,  
35 cheese firmness decreased. Cheeses with reduced NaCl were less firm and more compressible. Cheeses  
36 produced with C2 were significantly firmer than those produced with C1; cheeses produced with C3 had  
37 higher firmness and compressibility. In NaCl reduced cheese, use of camel chymosin as coagulant  
38 resulted in significantly higher firmness than that given using bovine chymosin. Overall, cheese NaCl  
39 content is reducible without significant textural impact using well-defined starter cultures and camel  
40 chymosin.

41

42

## 43 1. Introduction

44

45 Dietary sodium, which is typically consumed as sodium chloride (NaCl), is an important  
46 ingredient, especially in pre-processed food, contributing to flavour and acting as a preserving agent  
47 (Mattes & Donnelly, 1991). The daily recommendation of NaCl intake for an adult is around 6 g per day  
48 (WHO, 2012), but the average daily intake in many European countries varies from 9 to 13 g NaCl per  
49 day (European Union, 2012). This elevated intake of NaCl can promote negative health consequences,  
50 such as hypertension, cardiovascular diseases and kidney failure (Appel et al., 2012; Frisoli, Schmieder,  
51 Grodzicki, & Messerli, 2012). Hence, there is a growing pressure for reducing the sodium content in  
52 processed foods. Within the dairy industry, cheese is an evident dairy product with potential for reduction  
53 in its sodium content. The salt content in cheese differs markedly with variety, from 0.5% (w/w) in  
54 cottage cheese to 4–6% (w/w) in feta cheese (Fox, Guinee, Cogan, & McSweeney, 2000).

55 Salt is a key ingredient in cheese. It is the major preservative, as it controls the water activity and  
56 thereby the microbial growth, protein hydration, enzymatic activity, but it also contributes to flavour  
57 formation and the textural properties of the cheese (Fox et al., 2000; Guinee, 2004; Pastorino, Hansen, &  
58 McMahon, 2003). Reducing the NaCl content in semi-hard cheeses, like Cheddar cheese, results in  
59 increased bitterness and unpleasant aftertaste together with decrease in salty taste and firmness (Johnson,  
60 Kapoor, McMahon, McCoy, & Narasimmon, 2009; Rulikowska et al., 2013; Schroeder, Bodyfelt, Wyatt,  
61 & McDaniel, 1988).

62 The majority of the cheeses produced in Denmark belong to the Danish semi-hard cheese types,  
63 e.g., Danbo and Samsøe, which unlike, for example, Cheddar cheese, are brined cheeses. These semi-hard  
64 cheeses contain, on average, 1.7–1.8% (w/w) NaCl, have a few round holes and are smear-ripened  
65 (Madsen & Ardö, 2001; Sørensen & Benfeldt, 2001). The cheeses are traditionally produced using  
66 bovine chymosin to coagulate the cheese milk along with mesophilic DL-starter cultures, i.e., containing  
67 *Lactococcus lactis* ssp. *lactis*, *Lactococcus lactis* ssp. *cremoris*, citrate-positive strains of *Lactococcus*  
68 *lactis*, and *Leuconostoc* ssp. (Daly, 1983; Jacob, Jaros, & Rohm, 2011). The salting of these Danish semi-

69 hard cheese types is done after pressing by immersion of the cheese into a saturated NaCl-solution for up  
70 to 28 h, i.e., brining. This differs from the salting process of Cheddar cheese, where the NaCl is added to  
71 the milled curd, mixed and then pressed (Grummer, Karalus, Zhang, Vickers, & Schoenfuss, 2012). The  
72 majority of studies on salt-reduced semi-hard cheese are on Cheddar and Gouda cheese. Research on  
73 textural properties of salt reduced brined semi-hard cheeses of Danbo type is lacking, and the literature on  
74 effect of brining mainly deals with mozzarella, halloumi, feta and soft cheese types (Ayyash & Shah,  
75 2011; Hynes, Delacroix, Meinardi, & Zalazar, 1999; Katsiari, Voutsinas, Alichanidis, & Roussis, 1997;  
76 Thibaudeau, Roy, & St-Gelais, 2015). Hence, we need new knowledge on the effect of brining for salt  
77 uptake and salt distribution to be able to directly correlate salt-reduction to the cheese texture of such  
78 brined cheeses.

79 Both the starter cultures and coagulation enzyme affect texture and flavour formation in the  
80 cheese. The composition of the starter culture used in cheese production varies according to cheese type  
81 and is often a mixture of various strains to achieve the desired properties of the cheese (Beresford,  
82 Fitzsimons, Brennan, & Cogan, 2001). In Cheddar cheese, NaCl can influence the viability of the starter  
83 culture and the enzymatic activities, and the proteolysis of the caseins has been found to increase as NaCl  
84 content decreases (Mistry & Kasperon, 1998; Møller, Rattray, Høier, & Ardö, 2012). Autolysis of *Lc.*  
85 *lactis* is favoured by low NaCl content (0.17 M) and acidic pH (pH 5.4) (Ramírez-Nuñez, Romero-  
86 Medrano, Nevárez-Moorillón, & Gutiérrez-Méndez, 2011). We have previously shown that the NaCl  
87 content in Danish semi-hard cheese affects both the viability and autolysis of lactic acid bacteria, which  
88 depends to a high degree on the specific DL-starter culture (Søndergaard et al., 2015). Hence, to  
89 compensate for the environmental changes in the cheese caused by reducing NaCl, a DL-starter culture  
90 combining more specific bacteria strains could be thought to retain some of the traditional semi-hard  
91 cheese properties, e.g., texture and flavour.

92 The presence and the amount of chymosin in the cheese is reported to increase the proteolysis  
93 during ripening (Hynes et al., 2001), and this proteolysis is affected by NaCl in different ways when  $\alpha_{S1}$ -  
94 casein and  $\beta$ -casein are considered (Noomen, 1978), which may impact the texture of the mature cheese.

95 Furthermore, camel chymosin has been shown to be an alternative to the traditional bovine chymosin.  
96 Camel chymosin has a 70% higher clotting activity towards bovine milk compared with bovine chymosin  
97 (Bansal et al., 2009; Jensen et al., 2015; Kappeler et al., 2006).

98 Previous studies comparing camel and bovine chymosin in Cheddar cheese showed firmer and  
99 less bitter cheeses when using camel chymosin (Bansal et al., 2009). Moynihan et al. (2014) also found  
100 less proteolysis occurring in mozzarella made using camel chymosin. The use of camel chymosin is a new  
101 approach to the aim of providing texture in reduced-salt cheese. It is hypothesised to result in cheeses  
102 with similar textural properties to those of cheeses made with bovine chymosin and normal NaCl level.

103 Many of the studies in this area vary according to cheese type, production methods, starter culture  
104 and chymosin type, which makes comparisons complicated. To our knowledge, no previous studies have  
105 evaluated the effect of NaCl content in brined Danish semi-hard cheeses in relation to cheese texture, and  
106 the present study brings novelty into the understanding this relationship.

107 The aim of this study was therefore to study the effect of NaCl reduction in Danish semi-hard  
108 cheese in relation to chemical composition and textural properties during cheese ripening. Additionally,  
109 three different DL-starter cultures and two different types of chymosin were used to investigate whether  
110 the DL-starter culture and/or the rennet type could counteract the consequences of reducing the NaCl  
111 content during processing of brined semi-hard Danish cheeses.

112

## 113 2. Materials and methods

114

### 115 2.1. Starter cultures and chymosin

116

117 Three different commercially available DL-starter cultures (C1, C2 and C3) were used (Chr.  
118 Hansen, Hørsholm, Denmark). All three starter cultures comprised of strains of *Lc. lactis* subsp. *lactis*,  
119 *Lc. lactis* subsp. *cremoris*, citrate-positive strains of *Lc. lactis* and *Leuconostoc* spp. The DL-starter culture  
120 C1 was a traditional DL-starter culture, propagated and produced as mixed-strain containing all these

121 organisms. The DL-starter culture C2 was composed of defined strains of the above-mentioned organisms,  
122 where strains have been isolated from a traditional DL-starter culture, grouped and grown separately,  
123 before combined into the final starter culture. Detailed description of these two DL-starter cultures can be  
124 found in Søndergaard et al. (2015). The DL-starter culture C3 was produced by isolating selected strains  
125 from the DL-starter culture C2. These selected strains had been grown separately before combined into the  
126 final DL-starter culture C3. The main difference between the DL-starter C2 and C3 was an increased level  
127 of *Lc. lactis* subsp. *lactis* in the DL-starter C3 (Chr. Hansen). Two commercially available chymosins  
128 (CHY-MAX M® and CHY-MAX plus®) were used (Chr. Hansen). The chymosins differ from each  
129 other according to their origin; CHY-MAX plus® contains bovine chymosin (BC), while CHY-MAX  
130 M® contains camel chymosin (CC).

131

## 132 2.2. Cheese manufacture and sampling

133

134 In total, four cheese experiments were performed to produce the described semi-hard Danish  
135 cheese types. Experiment 1 and 2 were performed to test the effect of starter culture, and experiments 3  
136 and 4 to test the effect of the chymosin type. An overview of the experiments is shown in Table 1.

137 Milk for all cheese productions was fresh, pasteurised under high-temperature-short-time  
138 conditions (72 °C, 15 s) bovine milk standardised to a fat to protein ratio of 0.5 (1.88% fat and 3.75%  
139 protein).

140 In experiment 1, semi-hard Danish cheeses (type Samsøe, 30+) were manufactured at the Arla  
141 Foods R&D (Brabrand, Denmark), with brining times of 0, 6, 12, and 24 h in saturated NaCl solution.  
142 The manufacture of these cheeses is described in detail in Søndergaard et al. (2015). Sampling for all  
143 brining times was performed at 1, 2, 7 and 12 weeks of ripening with 3 cheese replicates of each  
144 treatment.

145 In experiment 2, semi-hard Danish cheeses (type Samsøe, 30+) were produced at Thise Dairy  
146 plant (Thise Dairy, Roslev, Denmark), with brining times of 0, 12, and 24 h, respectively. DL-starter

147 cultures C1 and C2 were used in both experiment 1 and 2. The manufacture of these cheeses is described  
148 in detail in Søndergaard et al. (2015). Sampling for all brining times was performed at 1, 2, 7 and 12  
149 weeks of ripening with 2 cheese replicates of each treatment.

150 In experiment 3, semi-hard Danish cheeses (type Danbo, 30+) were manufactured at Arla Foods  
151 R&D (Brabrand, Denmark), with brining times of 6, 12, and 24 h. The procedure was similar to  
152 Søndergaard et al. (2015), with some modifications. Two batches, each of 1000 L milk per day for two  
153 days, were used for cheese production. Each day, the milk for one batch was coagulated by use of CHY-  
154 MAX Plus® at a rate of 0.03% (w/w) (Chr. Hansen, Hørsholm, Denmark), and one batch was coagulated  
155 using CHYMAX-M® at a rate of 0.01% (w/w) (Chr. Hansen, Hørsholm, Denmark). These levels of  
156 chymosin correspond to equal levels of international milk clotting unit (IMCU) per L milk. The DL-starter  
157 culture C3 used in this experiment was added at a rate of 0.008 % (w/w) together with 0.005% (w/w)  
158 CaCl<sub>2</sub>. Cheeses were placed in a saturated NaCl solution (23.3%, w/v) with 0.25% (w/v) CaCl<sub>2</sub> at 11.5 °C  
159 for 6 h, 12 h and 24 h, respectively. The cheeses were smeared at the surface, cut in half (approximately  
160 15 × 30 × 15 cm), vacuum packed and ripened at 13 °C. Sampling for all brining times was performed  
161 after 1 and 12 weeks of ripening with 2 cheese replicates of each treatment.

162 In experiment 4, semi-hard Danish cheeses (type Danbo, 30+) were manufactured at Arla Foods  
163 Dairy plant (Taulov, Denmark). The procedure was similar to Søndergaard et al. (2015), with some  
164 modifications. Three batches each of 21,350 kg milk were used for production. The milk for one batch  
165 was coagulated using CHY-MAX Plus® at a rate of 0.03% (w/w) (Chr. Hansen). Cheeses from this batch  
166 were placed in a saturated NaCl solution (23.3%, w/v) with 0.25% (w/v) CaCl<sub>2</sub> at 11.5 °C for 28 h except  
167 for the 0 h brining treatment, where this step was omitted. Two batches obtained coagulation using  
168 CHYMAX-M® at a rate of 0.02% (w/w) (Chr. Hansen). The levels of chymosin correspond to equal  
169 levels of IMCU. Cheeses from these batches were split in two parts which were either placed in a  
170 saturated NaCl solution (23.3%, w/v) with 0.25% (w/v) CaCl<sub>2</sub> at 11.5 °C for 28 h or in the saturated NaCl  
171 solution for either 10 h or 15 h (see Table 1 for overview). Due to practical issues, another commercial  
172 DL-starter culture, the standard at Taulov Dairy, was used in the final trials, proven to give the same rate

173 of acidification, flavour development, and eye formation. The cheeses were produced in dimensions of 38  
174  $\times 76 \times 8.5$  cm. The longitudes of the cheeses were smeared, and stored for 4 weeks at  $\sim 15$  °C and relative  
175 humidity of 92–97%, then washed, coated with paraffin and further stored 3 weeks at  $\sim 8$ –9 °C. The  
176 cheeses were then cut into pieces of  $15 \times 9 \times 4.25$  cm, vacuum packed separately and stored at 3.5 °C  
177 until analysis after 12 weeks of ripening with 3 cheese replicates of each treatment.

178

### 179 2.3. *Texture analysis by uniaxial compression*

180

181 Textural properties of the cheeses were analysed by uniaxial compression analysis. Cheeses at the  
182 desired ripening time were stored for 24 h at 4°C before analysis. A lubricated cork borer, dipped in oil to  
183 minimize friction between cork borer and cheese, was slowly pressed through the cheese vertically to  
184 create cylindrical cheese pieces. Care was taken not to disrupt the cheese structure when using the cork  
185 borer. Cylindrical cheese samples with height (h) = 15 mm and diameter (d) = 15 mm from various  
186 locations (in experiments 1, 2, and 3, a total of 12 locations) in the cheese were used for textural analysis  
187 by uniaxial compression, and in experiment 4, as a result of the previous experiments only 2 locations  
188 were used for sampling (edge and core). The edge samples were taken at a position 1 cm from the surface  
189 of the cheese, while the core samples were taken in the centre of the cheese. The cheese cylinders were  
190 analysed immediately after cutting. Compression was performed until fracture of the cheese, or to a  
191 maximum distance of 12.5 mm, using a TA HDi Texture Analyzer (Stable Micro Systems, Godalming,  
192 UK) with a 100 kg load cell, 1 mN detection range, 75 mm diameter flat stainless steel plate and  
193 compression speed of  $0.8 \text{ mm s}^{-1}$ . Data were organised and initial data analysis was made in Texture  
194 Expert Exceed Version 2.63 (Stable Micro Systems, Godalming, UK). Recordings of force (N) and  
195 displacement (m) were converted into true axial stress  $\sigma$  (Equation 1) representing firmness of the cheese  
196 and Hencky strain  $\varepsilon$  (Equation 2) representing compressibility of the cheese according to the following:

$$197 \quad \sigma = \frac{F}{A} \frac{H}{H_1} \text{ [Pa]} \quad (1)$$

198 
$$\varepsilon = -\ln \frac{H}{H_i} [-] \quad (2)$$

199 where  $F$  = force (N),  $A$  = initial end area of sample ( $\text{m}^2$ ),  $H_i$  = initial sample height (m) and  $H$  = height  
200 (m) (Hammershoj, Larsen, Ipsen, & Qvist, 2001). The  $\sigma_f$  and  $\varepsilon_f$  were obtained as sample stress and sample  
201 strain, respectively, at the fracture point and used for further statistical analysis. For cheese samples,  
202 which did not obtain a breakage point within the distance of analysis, the force at maximum distance was  
203 used for further calculations.

204

#### 205 2.4. *Cheese composition*

206

207 At all sampling times, composition of analysis (%) was performed according to International  
208 Organization for Standardisation (ISO) methods for dry matter (ISO5534 and IDF004; ISO, 2004b), fat  
209 (ISO1735 and IDF005; ISO, 2004a), protein (6.38×N; ISO8968-1 and IDF020-1; ISO, 2014), and NaCl  
210 content (ISO5943 and IDF088; ISO, 2006). The pH of the grated cheese was measured by potentiometry.  
211 All analyses were conducted at Eurofins Steins Laboratory (Holstebro, Denmark) on a mixture of grated  
212 cheese from one whole cheese of each cheese replicate for all sampling times.

213

#### 214 2.5. *Determination of NaCl content*

215

216 The compressed cylindrical cheese samples from texture analysis were collected afterwards and  
217 used for analysis of the NaCl content. This approach was chosen to get a data set from the same position  
218 in the cheese to correlate textural properties to NaCl concentration. This was determined using a  
219 modification of the ISO method (ISO5943 & IDF088; ISO, 2006) based on potentiometric  $\text{AgNO}_3$   
220 titration of  $\text{Cl}^-$  ions. A mass of 1–1.5 g cheese from the compressed cylindrical cheese samples was  
221 collected in a 100 mL tube and added 30 mL 55 °C milliQ water along with 20 mL 1 M sodium citrate.  
222 The mixture was blended for 45 s with an Ultra Turrax homogeniser (IKA-Labortechnik, Janke & Kunkel

223 GmbH & Co., Staufen, Germany) at a speed of 10,000 rpm. To ensure that no sample mixture remained  
224 onto the homogenizer, 20 mL 55 °C milliQ water was used to wash off sample from the Ultra Turrax. The  
225 samples were then incubated at room temperature for 1 h after which time 10 mL 4 M HNO<sub>3</sub> was added.  
226 A Metrohm 862 compact titrosampler (Metrohm AG, Herisau, Switzerland) was used to determine the  
227 NaCl content potentiometrically using 0.1 M AgNO<sub>3</sub> solution as titrant. At the equivalence point of the  
228 titration, the amount of added silver nitrate was noted and used to calculate the concentration of NaCl in  
229 the sample solution (Equation 3):

$$230 \quad NaCl(\%) = \frac{v(AgNO_3) \cdot c(AgNO_3) \cdot M(NaCl) \cdot 0.1}{m(sample)} \quad (3)$$

231 Two cheese cylinders from the textural analysis, one from the core and one from the edge, were  
232 analysed for salt content for all cheeses.

233

#### 234 2.6. *Microstructure by scanning electron microscopy*

235

236 The network structure of cheeses after 12 weeks of ripening was studied by scanning electron  
237 microscopy (SEM). The procedure is described in detail in Søndergaard et al. (2015). A 1 mm<sup>3</sup> cube of  
238 cheese was fixed with 2.5% glutaraldehyde in 0.1 M piperazine-N,N'-bis(2-ethanesulfonic acid). The  
239 cheese sample was dehydrated by washing in ethanol in a series of stepwise increasing 10%  
240 concentrations from 10–100% ethanol each for 15 min. For the 20% ethanol step and onwards, the cheese  
241 sample was transferred to a critical point drying (CPD) capsule. The sample was washed in 100% dry  
242 ethanol for 2 × 15 min and stored at  $T = 4$  °C for at least 1 h before CPD. The CPD procedure was  
243 performed with liquid CO<sub>2</sub> using a Leica CPD300 (Leica Microsystems, Heidelberg, Germany), and the  
244 dried samples were stored in a sealed container at room temperature until analysis. Before analysis, the  
245 sample was secured at an aluminium SEM stub with Ag paint and fractured in the horizontal plane. The  
246 free-break surface was thereby facing upwards, and the surface was covered with a thin layer of Au using  
247 an agar high resolution sputter-coater (Agar Scientific, Stansted, UK). The prepared sample was observed

248 at 3 kV with a Zeiss Supra 55VP FEG Scanning Electron Microscope (Carl Zeiss, Oberkochen,  
249 Germany), at a working distance of ~5 mm at magnifications ranging from 1000 to 95,000 ×. Several  
250 pictures were captured for each cheese sample. Each sample was analysed in duplicates and samples were  
251 taken from the core and edge of the cheeses.

252

### 253 2.7. *Statistical analysis*

254

255 Two-way ANOVA and three-way ANOVA were performed to determine significant differences  
256 ( $P < 0.05$ ) among cheeses at different brining times, ripening time, depending on DL-starter culture and  
257 chymosin type. Differences were classified by the Ryan-Einot-Gabriel-Welsch multiple range test (SAS,  
258 version 9.3, SAS Institute Inc., Cary, NC). See Table 1 for the different variables and replicates.

259

## 260 3. **Results and discussion**

261

### 262 3.1. *Chemical composition during ripening*

263

264 Samples of all cheeses were collected during ripening for chemical compositional profiling. Fig. 1  
265 shows the chemical composition for the semi-hard Danish cheeses produced in experiment 1, with DL-  
266 starter culture C1, during ripening. Similar trends were observed for the production of the cheeses in  
267 experiment 2, 3 and 4 and are therefore not shown. As expected, a significant increase ( $P < 0.001$ ) in the  
268 total NaCl content was found with increased brining time (Fig. 1A). During the ripening period, the total  
269 NaCl content did not change significantly, which was also as expected. A small amount of  $\text{Cl}^-$  was  
270 detected in the non-brined cheeses. This is due to naturally occurring  $\text{Na}^+$  and  $\text{Cl}^-$  ions present in the milk  
271 before cheese making (Belitz, Grosch, & Schieberle, 2004). Results shown in Fig. 1 are derived from  
272 cheeses produced from the same batch of milk, and the only difference among the cheeses was the brining  
273 time. Changes in the chemical composition are therefore caused by differences in the NaCl content. The

274 protein content as a function of brining time and ripening time is shown in Fig.1B. No significant  
275 difference in the protein content was observed between brining times after 1, 2, and 7 weeks of ripening.

276 As ripening time increased to 12 weeks, a significant decrease in the protein content for the non-  
277 brined cheeses was observed ( $P < 0.05$ ). It is known that the water activity is higher and bacterial activity  
278 is increased, when the NaCl content is lowered, which might increase proteolysis (Guinee, 2004). It  
279 correlates with our previous results (Søndergaard et al., 2015), where the microbial activity in cheeses  
280 from experiments 1 and 2 were analysed. The cheese made using DL-starter culture C1 was found to have  
281 significantly higher number of colony forming units (cfu) per gram in the non-brined cheeses after 1 and  
282 2 weeks of ripening (58 and 71%, respectively) compared with the cheeses treated 24 h in brine (21 and  
283 22%, respectively), hence the proteolytic activity could likely be higher as the NaCl content decreased.

284 Proteolysis in non-brined cheeses is primarily due to primary proteolysis as NaCl-reduction  
285 accelerates the degradation of casein due to higher chymosin activity (Møller et al., 2012). With respect to  
286 non-starter lactic acid bacteria (NS-LAB), these were investigated and described in detail for cheeses  
287 produced in experiment 1 (Søndergaard et al., 2015). Here, the results for cheeses produced with C1 and  
288 C2 showed no significant influence of the NaCl concentration on the NS-LAB counts during the ripening  
289 period. However, the normal-salted cheeses had slightly lower NS-LAB counts after 2 weeks of ripening  
290 in cheeses produced with C1. Based on this we do not expect that the NS-LAB population will have a  
291 major impact on secondary proteolysis in this study.

292 The soluble peptides and free amino acids can to some extent be released from the protein matrix  
293 in the cheese and diffuse into the soluble fraction of the cheese, which may reduce the protein content in  
294 the cheese (Fox et al., 2000); however, as the protein content here was based on nitrogen analysis the  
295 peptides and amino acids still counted in the protein content (ISO8968-1 & IDF020-1, 2014). A more  
296 likely explanation could be the higher moisture content of non-brined cheeses (Fig. 1C), where the  
297 decrease in dry matter content with ripening time was more apparent for the non-brined cheeses ( $P <$   
298 0.05).

299 The content of dry matter, Fig. 1C, was found to increase as the brining time increased, which  
300 was expected ( $P < 0.001$ ). This is caused by NaCl migrating from the rind into the cheeses during  
301 ripening and water was expelled from the cheese (Guinee, 2004).

302 Fig. 1D shows the development in pH during ripening. The pH after 1 week was lower than pH of  
303 the cheese ripened for 2–12 weeks. This is related to the degradation of lactose by the lactic acid bacteria  
304 of the DL-starter culture (McSweeney & Fox, 2004). Ripening times above 1 week resulted in  
305 significantly increase in pH with only small changes in pH from ripening weeks 2–12. This has been  
306 shown to be caused by rebalancing of calcium phosphate equilibrium in the cheeses, proteolysis of  
307 proteins and degradation of lactic acid (Hassan, Johnson, & Lucey, 2004; McMahon et al., 2014).  
308 Cheeses with 24 h of brining time had generally lower pH during ripening (Fig. 1D), which seems to be  
309 related to increased syneresis (Nielsen, 2006). Furthermore, a difference between starter cultures is  
310 reported, as for C1 the pH was lower with increased salt content, whereas for C2 the pH did not vary  
311 significantly as function of salt content (Søndergaard et al., 2015). This also contributes to the explanation  
312 of the interlinked effect between the specific DL-starter culture and its activity, the protein content and the  
313 resulting pH, as the accumulation of organic acids inhibits the growth of microorganisms, i.e., inevitably  
314 also the starter culture (Beresford et al., 2001).

315 Overall, these findings in brined semi-hard Danish cheese are in agreement with previous findings  
316 in dry-salted Cheddar cheese by Schroeder et al. (1988) and Rulikowska et al. (2013), who analysed the  
317 chemical changes in Cheddar cheese with reduced NaCl content during ripening.

318

### 319 3.2. *NaCl distribution in the cheese*

320

321 The Cl<sup>-</sup> content in the cheeses was measured at various positions from the edge to the core of the  
322 cheeses as representative of the NaCl distribution in the cheeses during ripening. Fig. 2 shows the NaCl  
323 content in the cheese samples from the edge and core as a function of brining time for ripening times of 2,  
324 7 and 12 weeks in experiment 2. Similar observations were found in experiments 1 and 3 (data not

325 shown). At 2 weeks of ripening, (Fig. 2A) a significant ( $P < 0.01$ ) difference in the NaCl content between  
326 edge and core of brine treated cheese samples was observed, with samples from the edge having the  
327 highest content of up to 3% (w/w) NaCl with a gradient to the core of ~1.5% (w/w) for the 24 h brined  
328 cheese. At 7 weeks of ripening (Fig. 2B) a significant ( $P < 0.05$ ) increase in NaCl concentration was  
329 found in the core, while the NaCl content in the edge decreased (NS), compared with 2 weeks of ripening.

330 For brining at both 12 h and 24 h, the difference in NaCl content between edge and core was still  
331 significant ( $P < 0.05$ ) after 7 weeks of ripening. After 12 weeks (Fig. 2C), NaCl was equally distributed  
332 between edge and core of the cheeses with 12 h brining, but not for the 24 h brined cheeses. This  
333 diffusion of NaCl from edge to core of the cheese is driven by the concentration gradient (Geurts,  
334 Walstra, & Mulder, 1980). The time to reach NaCl equilibrium depends on cheese type, size and shape of  
335 the cheeses and ripening temperature. Sutherland (2002) observed similar results for 10 kg Gouda  
336 cheeses.

337

### 338 3.3. NaCl and DL-starter culture

339

340 The cheeses in experiments 1, 2 and 3 with three different commercial DL-starter cultures (C1, C2  
341 and C3) were analysed to evaluate the effect of the DL-starter culture on the chemical composition and  
342 textural properties of semi-hard cheeses. The mean NaCl contents of the cheeses with the three different  
343 DL-starter cultures are shown in Table 2 as a function of brining time after 12 weeks of ripening. The  
344 NaCl contents in the cheeses were analysed as an average of the entire cheese in contrast to the positional  
345 analysis, shown in Fig. 2. The DL-starter cultures did not affect the NaCl content of the cheeses  
346 significantly. Other factors may contribute to the final NaCl contents such as dairy factory, milk batch,  
347 pressing of the cheese, pore size and structure, brine saturation, etc. Furthermore, the cheeses produced  
348 with C3 had a tendency towards higher dry matter content, while no other differences in the chemical  
349 composition were observed (data not shown).

350 The most efficient brining was achieved during the first 6 h (experiments conducted with DL-  
351 starter cultures C1 and C2) with a rate of  $0.138 \pm 0.006\% \text{ NaCl h}^{-1}$ , while thereafter it decreased to  
352  $0.053 \pm 0.003\% \text{ NaCl h}^{-1}$  from 6–12 h and finally during the last 12 h of brining the rate was lowered to  
353  $0.036 \pm 0.002\% \text{ NaCl h}^{-1}$  for all starter cultures. This suggests that the brining process and NaCl uptake for  
354 these semi-hard Danish cheeses occurred in a very consistent way, regardless of the above-mentioned  
355 differences between the dairies, milk batches and DL-starter cultures.

356

#### 357 3.4. Cheese textural change during ripening

358

359 The cheese firmness, illustrated as axial stress, and compressibility, illustrated as Hencky strain,  
360 for experiment 2 with DL-starter culture C1 and C2 according to ripening time are shown in Fig. 3. The  
361 firmest cheeses were found with ripening of 1 week for all brining times and starter cultures, Fig.3A. As  
362 the ripening time increased, the firmness decreased significantly ( $P < 0.01$ ) for all cheeses. These results  
363 are in agreement with Murtaza et al. (2014), who followed the texture profile in Cheddar cheeses with  
364 various NaCl content during ripening. The firmness is correlated to the proteolysis, i.e., increased  
365 proteolysis during ripening results in decreased firmness of the cheeses (Fox, 1989; McSweeney, 2004).  
366 The cheese network of caseins is weakened by the proteolytic degradation into peptides, and as a result,  
367 the texture becomes softer over time.

368 The decrease in firmness was most pronounced for C2 with a brining time of 24 h while cheeses  
369 subjected to 0 h and 12 h of brining showed similar decreases in firmness. For both DL-starter cultures,  
370 the relative loss in cheese firmness during 12 weeks of ripening was highest for the non-brined cheeses  
371 with 68–72% loss relative to week 1. During the same period, the 24 h brined cheeses had a textural loss  
372 of 36–49%. It is noteworthy, that this was mainly caused by differences in the initial cheese firmness, as  
373 the actual decrease in stress was  $19.4 \pm 0.5 \text{ kPa}$  (all cheeses produced with C1) and  $27.0 \pm 3.3 \text{ kPa}$  (all  
374 cheeses produced with C2) during ripening regardless of brining time. This suggests a very similar  
375 development in cheese structure and therefore firmness during ripening.

376 Overall, the use of DL-starter culture C2 generally resulted in significantly ( $P < 0.05$ ) firmer  
377 cheeses compared with C1, regardless of brining time. From the standard deviation bars of Fig. 3, it is  
378 clear that large variations between samples were observed, especially in week 1 of the ripening period,  
379 while the variation between samples decreased as ripening time increased. This is due to that the mean  
380 value was generated from samples from both edge and core. As shown in Fig. 2 there were large  
381 variations in NaCl content among samples from edge and core of the cheese, until final ripening stage was  
382 reached, which resulted in variations in firmness.

383 The compressibility is given as Hencky strain as function of brining time, ripening and starter  
384 culture (Fig. 3B). For the present semi-hard Danish cheeses, a high Hencky strain value indicated a highly  
385 compressible or elastic cheese, while a low Hencky strain correlated with a cheese that fractured at low  
386 compression distance and was observed as more brittle. This is consistent with previous findings for  
387 Gouda cheese (Luyten, 1988). Throughout the ripening time, small variations for all salted cheeses  
388 occurred, but these were not significant. The non-brined cheeses increased in compressibility for ripening  
389 times of 2–7 weeks ( $P < 0.01$ ). For these cheeses, there was often not detected a fracture point of the  
390 cheese cylinder during the textural compression analysis. The samples were very elastic and could be  
391 compressed >83% without breaking during the analysis. These non-brined cheeses were also more prone  
392 to temperature, which made them lose their cylindrical structure very quickly, while all salted cheeses  
393 retained their shape at room temperature.

394 Generally, it is found that the compressibility decreases during ripening for cheeses like Cheddar  
395 and Gouda (Luyten, 1988; Zoon, 1993). Furthermore, Watkinson et al. (2001) observed an increase in  
396 Hencky strain during ripening of Gouda cheeses. In this study, the compressibility appeared unaffected by  
397 the changes occurring in the cheese during ripening of salted cheeses.

398 In comparison, the stress at fracture and Hencky strain values of 7 week ripened Danbo (30+)  
399 cheeses is reported to be 92 kPa and 1.10 (-), respectively, by Madsen and Ardö (2001), which is  
400 somewhat higher in stress at fracture than observed in the present study, where 7 week ripened cheeses  
401 had values of ~45 kPa (Fig. 3). Their compressibility levels are, however, comparable with levels

402 presented in Fig. 3B. The cheese firmness may be affected by a range of processing parameters, although  
403 the dry matter content was ~47% in both studies. This is illustrated for the textural analysis in experiment  
404 3, which resulted in much firmer reference cheeses (24 h brining, culture C3, bovine chymosin) after 12  
405 weeks ripening with fracture stress values of 100 kPa and Hencky strain of 1.09 (-) (data not shown).

406 Søndergaard et al. (2015) analysed the number of viable lactic acid bacteria (LAB), the extent of  
407 autolysis and also determined free amino acids of the cheeses as used in experiment 1 and 2. For the DL-  
408 starter culture C1, growth was found to be more affected by the NaCl concentration as compared with the  
409 DL-starter culture C2. Elevated levels of free amino acids have previously been found to increase stress  
410 and decrease strain due to binding of water to peptide bonds in the cheese matrix (Børsting et al., 2012;  
411 McSweeney, 2004), which can relate to the observed variation in texture between C1 and C2.

412 SEM micrographs of cheeses from experiment 2 (DL-starter cultures C1 and C2) brined for either  
413 0 h or 24 h after 12 weeks of ripening are shown in Fig. 4. The holes in the protein matrix originate from  
414 fat and water, which were removed during sample preparation. Variations in the number and size of voids  
415 in the cheese matrix can be observed. Fig. 4B and Fig. 4D show cheeses with 24h brining time. These had  
416 a more clearly structured protein matrix with many and smaller voids than in Fig. 4A and Fig. 4C, which  
417 are micrographs of non-brined cheeses. The protein matrix of the non-brined cheeses appeared less  
418 defined, which was seen by fewer and slightly larger voids. Comparing DL-starter culture C1 and C2,  
419 there was a tendency towards a more defined protein matrix when using C1. However, this was not  
420 confirmed with certainty by the SEM analysis. These microstructural observations support the chemical  
421 and textural results as the less defined protein matrix structure visualized by the SEM would be expected  
422 to result in softer and more compressible cheese texture as observed.

423

### 424 3.5. *Textural change as an effect of NaCl*

425

426 Cheese samples used for textural analysis were also analysed for NaCl content to explore the  
427 correlation between NaCl content and textural properties. Fig. 5 shows the correlation between textural

428 properties and NaCl content for cheeses from experiment 2, for both DL-starter cultures C1 and C2,  
429 during ripening of cheese samples from both edge and core. For the non-brined cheeses, there was not  
430 always a detectable fracture point of the cheese cylinders, when performing the texture analysis. These  
431 samples were so elastic that they could be compressed without breaking, and they were therefore not  
432 included in Fig. 5.

433 The firmness, given as axial stress, as a function of NaCl content is shown in Fig. 5A–C. An  
434 increase in firmness was observed with increasing NaCl content. This was expected, as NaCl is a major  
435 contributor to the formation of a strong gel network (Guinee, 2004; Mistry & Kasperson, 1998; Schroeder  
436 et al., 1988). After 2 weeks of ripening (Fig. 5A), large variations in texture were found between samples.  
437 During further ripening, (Fig. 5B,C), these variations became less pronounced and after 12 weeks of  
438 ripening there was a linear correlation with a regression coefficient of  $R^2 = 0.75$ . These observations  
439 could be related to the results shown in Fig. 2, which showed large differences in NaCl between edge and  
440 core in early ripening, while this became less pronounced during ripening.

441 The compressibility of the cheese, given as Hencky strain, (Fig. 5D–F) decreased linearly as the  
442 NaCl content increased. The fracture point of the cheese sample thereby occurred at a shorter distance in  
443 the textural compression analysis, which means that the samples became less elastic and more brittle.  
444 As for the firmness, the compressibility showed large variations among samples after 2 weeks of ripening  
445 and this became less during ripening. The correlation coefficients were generally low with  $R^2$ -values  
446 between 0.12-0.47, and the fit was poorest for the 2 weeks ripened cheeses. Especially for low salt  
447 content cheeses, the variations in Hencky strain at 7 and 12 weeks of ripening were very high. In  
448 perspective, a cheese with 0.5-1% (w/w) NaCl could have been useful to include to complete the picture.

449 The higher number of samples depicted in Fig. 5 revealed novel information on the texture in  
450 cheese core and cheese edge when salt migrated during ripening. At the very beginning of ripening (Fig.  
451 5A,D), the core and edge of brined cheese were clustered based on the textural properties, while increased  
452 ripening time resulted in more textural uniformity between the edge and core samples (Fig. 5C,F).

453 The relationship between firmness and compressibility and NaCl content for cheeses produced  
454 with DL-starter cultures C1, C2 and C3 and ripened for 12 weeks are shown in Table 2. Experiment 3  
455 cheeses with DL-starter culture C3 were produced at same dairy plant but at a different time, compared  
456 with cheeses produced with the C1 and C2 DL-starter cultures from experiment 1.

457 Comparing the DL-starter cultures, all cultures had similar tendencies to increase firmness with  
458 increased NaCl content (Table 2). However, the DL-starter culture C3 in experiment 3 produced much  
459 firmer cheeses at comparable brining hours than C1 and C2 in experiment 1, which resulted in 2-fold  
460 higher axial stress values for C3 compared with C1, regardless of NaCl content. The usage of starter  
461 culture C3 showed higher variations in firmness, which was found relating to variations between  
462 replicates. The variations were relatively lower for the DL-starter cultures C1 and C2. However, as  
463 explained earlier it is noted that the experimental set-up did vary for the cheeses produced with C1 and C2  
464 as compared with the cheeses produced with C3.

465 The compressibility, given as Hencky strain, for cheeses produced with C2 tended to be lower  
466 than cheeses produced with C3, while cheeses produced with C1 had the highest compressibility, i.e., they  
467 were more elastic.

468 C1 resulted in cheeses with lower firmness and higher compressibility compared with C2 and C3  
469 at similar brining times resulting in NaCl concentrations within a range of 0.11% at 6 h, 0.13% at 12 h,  
470 and 0.20% at 24 h brining time (Table 2). This indicates that the more defined DL-starter cultures,  
471 represented by C2 and especially C3, might result in firmer and more brittle cheeses. C3 resulted in the  
472 most firm cheeses, but these cheeses had also larger compressibility compared with the cheeses produced  
473 with C2. This indicates that the casein network of cheeses produced with C2 was more compact and  
474 therefore broke more easily.

475 Scientific studies on the relationship between NaCl content and cheese texture for brined semi-  
476 hard Danish cheeses are not available. However, for Cheddar cheese made from buffalo milk a reduction  
477 of NaCl content from 2.5% to 0.5% (w/w) resulted in lower hardness and crumbliness of the cheese  
478 textual properties (Murtaza et al., 2014). In another study, NaCl in Cheddar cheese was reduced from

479 2.3% to 0.9% (w/w); however, by maintaining an equal moisture content of  $37.6 \pm 0.1\%$ , the textural  
480 properties of the cheeses in the range from 0.9–1.7% (w/w) NaCl were kept similar (Møller, Rattray,  
481 Bredie, Høier, & Ardö, 2013). Also, replacing NaCl partly by other salts like KCl,  $MgCl_2$  and  $CaCl_2$  is  
482 reported to alter the hardness of Cheddar cheese in ways of both increased hardness and decreased  
483 hardness, even though the salt-to-moisture relationship and water activity was maintained at the same  
484 level (Grummer et al., 2012). The general trend of reducing NaCl in Cheddar cheese is a parallel change  
485 in textural properties (Floury et al., 2009; Rulikowska et al., 2013; Saint-Eve, Lauverjat, Magnan, Délérís,  
486 & Souchon, 2009), unless the NaCl reduction is substituted with other salts and/or moisture management  
487 is addressed.

488

### 489 3.6. Cheese textural effects of chymosin type

490

491 Cheeses of experiment 3 and 4 were analysed with regard to the effect of the origin of chymosin,  
492 camel (CC) or bovine (BC) on the chemical composition and textural properties of the cheese. However,  
493 as the choice of DL-starter culture varied between the cheese productions, the experiments cannot be  
494 compared directly. Table 3 shows the chemical composition and textural properties of the cheeses made  
495 with either chymosin type CC or BC for both experiments 3 and 4 after 12 weeks of ripening. Again, a  
496 significant increase in NaCl content was observed as the brining time increased ( $P < 0.05$ ), but no  
497 differences were found when comparing the chymosin types at equal brining times. The NaCl uptake in  
498 the cheeses was thus apparently not affected by the chymosin type. The dry matter content increased as  
499 brining time increased, but no significant differences between BC and CC cheeses were observed. The  
500 firmness of the cheeses with 6 h of brining for CC cheeses produced significantly ( $P < 0.05$ ) firmer  
501 cheeses compared with BC cheeses at equal brining times (Table 3). This is in agreement with Elagamy  
502 (2000), who observed that CC activity was less affected by low NaCl concentrations, while at high NaCl  
503 concentration both chymosin types were more equally affected. At brining times of  $< 12$  h, there was a  
504 significant textural effect of CC resulting in firmer cheeses than BC (experiment 3, Table 3), whereas at

505 brining times longer than 10 h, a tendency towards firmer cheeses with CC compared with BC was  
506 observed; however, this effect was not significant. In experiment 4, the CC renneted cheese brined for 15  
507 h had an axial stress level comparable with the BC renneted cheese brined for 28 h (Table 3) even at a  
508 NaCl content that was reduced by 18%. Firmer cheeses are generally found when using CC compared  
509 with BC (Bansal et al., 2009; Børsting et al., 2012; Govindasamy-Lucey, Lu, Jaeggi, Johnson, & Lucey,  
510 2010; Moynihan et al., 2014). It was therefore expected that the CC would result in firmer cheeses, as the  
511 amount of enzymes added corresponded to equal enzymatic activities (IMCU per mL milk). Different  
512 results among studies are most likely caused by variations in cheese type, DL-starter culture and amount  
513 of chymosin added.

514         The compressibility decreased as the NaCl content increased. For experiment 3, no significant  
515 differences in compressibility were found between chymosin types. In experiment 4, a significantly lower  
516 ( $P < 0.001$ ) compressibility was observed for CC compared with BC at comparable brining times of 28 h.  
517 Furthermore, for practical reasons, it was decided to not include a control treatment (0 h brining) in  
518 experiment 3, and only for the BC treatment in experiment 4. Basic knowledge on non-brined cheeses  
519 textural properties was obtained in experiments 1 and 2, and since the perspective for the Danish dairies is  
520 to reduce salt in cheese rather than avoiding salt in cheese, it was prioritised to include more treatments  
521 with reduced salt rather than with no salt in experiments 3 and 4.

522         SEM micrographs of cheeses from experiment 4 with 28 h of brining and ripened for 12 weeks  
523 are shown in Fig. 6. The structure of CC cheese (Fig. 6B) appears finer stranded and more compact than  
524 the BC cheese (Fig. 6A) and contains many small pores, while the BC cheese appears to contain more  
525 open network of relatively larger pores. Since this is the first time SEM images of salt reduced semi-hard  
526 brined cheeses are reported, we cannot compare to other studies. The structure show some agreement with  
527 Weijers, van de Velde, Stijnman, van de Pijpekamp, and Visschers (2006), who observed that gels  
528 composed of relatively thin network strands and small homogeneous pores are more brittle and would  
529 fracture at low strain values, while gels that fracture at high strain values are composed of thicker strands  
530 and relatively larger homogeneous pores.

531

**532 4. Conclusions**

533

534 Overall, this study has provided new knowledge on the effect of NaCl, DL-starter culture and  
535 chymosin type on the textural properties and chemical composition of Danish semi-hard cheeses. Shorter  
536 brining time reduced the NaCl content with a significant influence on firmness, compressibility and  
537 chemical composition of the cheeses. Cheese firmness increased and compressibility decreased linearly as  
538 the NaCl content increased. The three different DL-starter cultures influenced the textural properties of the  
539 cheeses. The most defined DL-starter culture, i.e., C3, produced significantly firmer cheeses while  
540 retaining a relative compressible cheese structure. The firmness was higher for cheeses made using camel  
541 chymosin at low NaCl content than for cheeses renneted with bovine chymosin. The compressibility of  
542 the cheeses was not significantly affected by chymosin type. However, the DL-starter culture may interact  
543 with the chymosin type in relation to cheese textural compressibility.

544 It therefore seems possible to reduce the NaCl content in semi-hard cheeses without  
545 compromising the textural properties by use of well-defined DL-starter cultures and camel chymosin. The  
546 cheese experiments performed at industrial scale provided novel insight into controlling cheese texture by  
547 brining under conditions that are readily applicable by the dairy industry. As the NaCl content also has an  
548 effect on the activity of the DL-starter cultures and the flavour formation, it is of importance to obtain  
549 knowledge on these parameters.

550

**551 Acknowledgements**

552

553 The authors thank for financial supported by the Danish Dairy Research Foundation, Arla Foods  
554 amba and the Future Food Innovation consortium of the region Mid-Jutland in Denmark. Laboratory  
555 technician Gitte Hald Kristiansen is acknowledged for performing the chloride analysis.

556

557 **References**

558

559 Appel, L. J., Angell, S. Y., Cobb, L. K., Limper, H. M., Nelson, D. E., Samet, J. M., et al. (2012).

560 Population-wide sodium reduction: The bumpy road from evidence to policy. *Annals of*  
561 *Epidemiology*, 22, 417–425.562 Ayyash, M. M., & Shah, N. P. (2011). Effect of partial substitution of NaCl with KCl on proteolysis of  
563 halloumi cheese. *Journal of Food Science*, 76, C31–C37.564 Bansal, N., Drake, M. A., Piraino, P., Broe, M. L., Harboe, M., Fox, P. F., et al. (2009). Suitability of  
565 recombinant camel (*Camelus dromedarius*) chymosin as a coagulant for Cheddar cheese.566 *International Dairy Journal*, 19, 510–517.567 Belitz, H. D., Grosch, W., & Schieberle, P. (2004). *Food chemistry* (3<sup>rd</sup> edn.). Berlin, Germany: Springer.568 Beresford, T. P., Fitzsimons, N. A., Brennan, N. L., & Cogan, T. M. (2001). Recent advances in cheese  
569 microbiology. *International Dairy Journal*, 11, 259–274.570 Børsting, M., Qvist, K., Rasmussen, M., Vindeløv, J., Vogensen, F., & Ardö, Y. (2012). Impact of  
571 selected coagulants and starters on primary proteolysis and amino acid release related to572 bitterness and structure of reduced-fat Cheddar cheese. *Dairy Science and Technology*, 92, 593–  
573 612.574 Daly, C. (1983). The use of mesophilic cultures in the dairy industry. *Antonie van Leeuwenhoek*, 49, 297–  
575 312.576 Elagamy, E. I. (2000). Physicochemical, molecular and immunological characterization of camel calf  
577 rennet: a comparison with buffalo rennet. *Journal of Dairy Research*, 67, 73v81.578 Floury, J., Camier, B., Rousseau, F., Lopez, C., Tissier, J.-P., & Famelart, M.-H. (2009). Reducing salt  
579 level in food: Part 1. Factors affecting the manufacture of model cheese systems and their  
580 structure–texture relationships. *LWT - Food Science and Technology*, 42, 1611–1620.581 Fox, P. F. (1989). Proteolysis during cheese manufacture and ripening. *Journal of Dairy Science*, 72,  
582 1379–1400.

- 583 Fox, P. F., Guinee, T. P., Cogan, T. M., & McSweeney, P. L. H. (2000). *Fundamentals of cheese science*.  
584 Gaithersberg, MD, USA: Aspen Publishers.
- 585 Frisoli, T. M., Schmieder, R. E., Grodzicki, T., & Messerli, F. H. (2012). Salt and hypertension: Is salt  
586 dietary reduction worth the effort? *American Journal of Medicine*, *125*, 433–439.
- 587 Geurts, T. J., Walstra, P., & Mulder, H. (1980). Transport of salt and water during salting of cheese. 2.  
588 Quantities of salt taken up and of moisture lost. *Netherlands Milk and Dairy Journal*, *34*, 229–  
589 254.
- 590 Govindasamy-Lucey, S., Lu, Y., Jaeggi, J. J., Johnson, M. E., & Lucey, J. A. (2010). Impact of camel  
591 chymosin on the texture and sensory properties of low-fat Cheddar cheese. *Australian Journal of*  
592 *Dairy Technology*, *65*, 139–142.
- 593 Grummer, J., Karalus, M., Zhang, K., Vickers, Z., & Schoenfuss, T. C. (2012). Manufacture of reduced-  
594 sodium Cheddar-style cheese with mineral salt replacers. *Journal of Dairy Science*, *95*, 2830–  
595 2839.
- 596 Guinee, T. P. (2004). Salting and the role of salt in cheese. *International Journal of Dairy Technology*,  
597 *57*, 99–109.
- 598 Hammershoj, M., Larsen, L. B., Ipsen, R. H., & Qvist, K. B. (2001). Effect of hen egg production and  
599 protein composition on textural properties of egg albumen gels. *Journal of Texture Studies*, *32*,  
600 105–129.
- 601 Hassan, A., Johnson, M. E., & Lucey, J. A. (2004). Changes in the proportions of soluble and insoluble  
602 calcium during the ripening of Cheddar cheese. *Journal of Dairy Science*, *87*, 854–862.
- 603 Hynes, E., Delacroix, B. A., Meinardi, C. A., & Zalazar, C. A. (1999). Relation between pH, degree of  
604 proteolysis and consistency in soft cheeses. *Australian Journal of Dairy Technology*, *54*, 24–27.
- 605 Hynes, E. R., Meinardi, C. A., Sabbag, N., Cattaneo, T., Candiotti, M. C., & Zalazar, C. A. (2001).  
606 Influence of milk-clotting enzyme concentration on the  $\alpha_{S1}$ -casein hydrolysis during soft cheeses  
607 ripening. *Journal of Dairy Science*, *84*, 1335–1340.

- 608 ISO. (2004a). *Cheese and processed cheese products - Determination of fat content - Gravimetric method*  
609 *(ISO1735 & IDF005 reference method)*. Geneva, Switzerland: International Standardisation  
610 Organisation.
- 611 ISO. (2004a). *Cheese and processed cheese - Determination of the total solids content (ISO5534 &*  
612 *IDF004 reference method)*. Geneva, Switzerland: International Standardisation Organisation.
- 613 ISO. (2006). *Cheese and processed cheese products - Determination of chloride content - Potentiometric*  
614 *titration method (ISO5943 & IDF088 reference method)*. Geneva, Switzerland: International  
615 Standardisation Organisation.
- 616 ISO. (2014). *Milk and milk products - Determination of nitrogen content - Part 1: Kjeldahl principle and*  
617 *crude protein calculation (ISO8968-1 & IDF020-1 reference method)*. Geneva, Switzerland:  
618 International Standardisation Organisation.
- 619 Jacob, M., Jaros, D., & Rohm, H. (2011). Recent advances in milk clotting enzymes. *International*  
620 *Journal of Dairy Technology*, 64, 14–33.
- 621 Jensen, J. L., Jacobsen, J., Moss, M. L., Rasmussen, F., Qvist, K. B., Larsen, S., et al. (2015). The  
622 function of the milk-clotting enzymes bovine and camel chymosin studied by a fluorescence  
623 resonance energy transfer assay<sup>1</sup>. *Journal of Dairy Science*, 98, 2853–2860.
- 624 Johnson, M. E., Kapoor, R., McMahon, D. J., McCoy, D. R., & Narasimmon, R. G. (2009). Reduction of  
625 sodium and fat levels in natural and processed cheeses: Scientific and technological aspects.  
626 *Comprehensive Reviews in Food Science and Food Safety*, 8, 252–268.
- 627 Kappeler, S. R., van den Brink, H. M., Rahbek-Nielsen, H., Farah, Z., Puhan, Z., Hansen, E. B., et al.  
628 (2006). Characterization of recombinant camel chymosin reveals superior properties for the  
629 coagulation of bovine and camel milk. *Biochemical and Biophysical Research Communications*,  
630 342, 647–654.
- 631 Katsiari, M. C., Voutsinas, L. P., Alichanidis, E., & Roussis, I. G. (1997). Reduction of sodium content in  
632 Feta cheese by partial substitution of NaCl by KCl. *International Dairy Journal*, 7, 465–472.

- 633 Luyten, H. (1988). *The rheological and fracture properties of Gouda cheese*. PhD Thesis. Wageningen,  
634 The Netherlands: University of Wageningen.
- 635 Madsen, J. S., & Ardö, Y. (2001). Exploratory study of proteolysis, rheology and sensory properties of  
636 Danbo cheese with different fat contents. *Cheese Ripening and Technology*, *11*, 423–431.
- 637 Mattes, R. D., & Donnelly, D. (1991). Relative contributions of dietary sodium sources. *Journal of the*  
638 *American College of Nutrition*, *10*, 383–393.
- 639 McMahon, D. J., Oberg, C. J., Drake, M. A., Farkye, N., Moyes, L. V., Arnold, M. R., et al. (2014).  
640 Effect of sodium, potassium, magnesium, and calcium salt cations on pH, proteolysis, organic  
641 acids, and microbial populations during storage of full-fat Cheddar cheese. *Journal of Dairy*  
642 *Science*, *97*, 4780–4798.
- 643 McSweeney, P. L. H. (2004). Biochemistry of cheese ripening. *International Journal of Dairy*  
644 *Technology*, *57*, 127–144.
- 645 McSweeney, P. L. H., & Fox, P. F. (2004). Metabolism of residual lactose and of lactate and citrate. In P.  
646 F. Fox, P. L. H. McSweeney, T. M. Cogan & T. P. Guinee (Eds.), *Cheese: Chemistry, physics*  
647 *and microbiology* (Vol. 1, pp. 361–371). New York, NY, USA: Academic Press.
- 648 Mistry, V. V., & Kasperson, K. M. (1998). Influence of salt on the quality of reduced fat Cheddar cheese.  
649 *Journal of Dairy Science*, *81*, 1214–1221.
- 650 Møller, K., Rattray, F., Høier, E., & Ardö, Y. (2012). Manufacture and biochemical characteristics during  
651 ripening of Cheddar cheese with variable NaCl and equal moisture content. *Dairy Science and*  
652 *Technology*, *92*, 515–540.
- 653 Møller, K. K., Rattray, F. P., Bredie, W. L. P., Høier, E., & Ardö, Y. (2013). Physicochemical and  
654 sensory characterization of Cheddar cheese with variable NaCl levels and equal moisture content.  
655 *Journal of Dairy Science*, *96*, 1953–1971.
- 656 Moynihan, A. C., Govindasamy-Lucey, S., Jaeggi, J. J., Johnson, M. E., Lucey, J. A., & McSweeney, P.  
657 L. H. (2014). Effect of camel chymosin on the texture, functionality, and sensory properties of  
658 low-moisture, part-skim Mozzarella cheese. *Journal of Dairy Science*, *97*, 85–96.

- 659 Murtaza, M. A., Huma, N., Sameen, A., Murtaza, M. S., Mahmood, S., Mueen-ud-Din, G., et al. (2014).  
660 Texture, flavor, and sensory quality of buffalo milk Cheddar cheese as influenced by reducing  
661 sodium salt content. *Journal of Dairy Science*, *97*, 6700–6707.
- 662 Nielsen, E. W. (2006). Principles of production of cheese. In Y. H. Hui (Ed.), *Handbook of food science,*  
663 *technology, and engineering* (Vol. 2, pp. 61–68). Cambridge, UK: Taylor & Francis Inc.
- 664 Noomen, A. (1978). Activity of proteolytic enzymes in simulated soft cheeses (Meshanger type). 2.  
665 Activity of calf rennet. *Netherlands Milk and Dairy Journal*, *32*, 49–68.
- 666 Pastorino, A. J., Hansen, C. L., & McMahon, D. J. (2003). Effect of salt on structure-function  
667 relationships of cheese. 1. *Journal of Dairy Science*, *86*, 60–69.
- 668 Ramírez-Nuñez, J., Romero-Medrano, R., Nevárez-Moorillón, G. V., & Gutiérrez-Méndez, N. (2011).  
669 Effect of pH and salt gradient on the autolysis of *Lactococcus lactis* strains. *Brazilian Journal of*  
670 *Microbiology*, *42*, 1495–1499.
- 671 Rulikowska, A., Kilcawley, K. N., Doolan, I. A., Alonso-Gomez, M., Nongonierma, A. B., Hannon, J. A.,  
672 et al. (2013). The impact of reduced sodium chloride content on Cheddar cheese quality.  
673 *International Dairy Journal*, *28*, 45–55.
- 674 Saint-Eve, A., Laverjat, C., Magnan, C., Déléris, I., & Souchon, I. (2009). Reducing salt and fat content:  
675 Impact of composition, texture and cognitive interactions on the perception of flavoured model  
676 cheeses. *Food Chemistry*, *116*, 167–175.
- 677 Schroeder, C. L., Bodyfelt, F. W., Wyatt, C. J., & McDaniel, M. R. (1988). Reduction of sodium chloride  
678 in Cheddar cheese: Effect on sensory, microbiological, and chemical properties. *Journal of Dairy*  
679 *Science*, *71*, 2010–2020.
- 680 Søndergaard, L., Ryssel, M., Svendsen, C., Høier, E., Andersen, U., Hammershøj, M., et al. (2015).  
681 Impact of NaCl reduction in Danish semi-hard Samsøe cheeses on proliferation and autolysis of  
682 DL-starter cultures. *International Journal of Food Microbiology*, *213*, 59–70.
- 683 Sørensen, J., & Benfeldt, C. (2001). Comparison of ripening characteristics of Danbo cheeses from two  
684 dairies. *International Dairy Journal*, *11*, 355–362.

- 685 Sutherland, B. J. (2002). Salting of cheese. In H. Roginski, J. W. Fuquay & P. F. Fox (Eds.),  
686 *Encyclopedia of dairy sciences* (pp. 293–300). Oxford, UK: Elsevier.
- 687 Thibaudeau, E., Roy, D., & St-Gelais, D. (2015). Production of brine-salted Mozzarella cheese with  
688 different ratios of NaCl/KCl. *International Dairy Journal*, *40*, 54–61.
- 689 Watkinson, P., Coker, C., Crawford, R., Dodds, C., Johnston, K., McKenna, A., et al. (2001). Effect of  
690 cheese pH and ripening time on model cheese textural properties and proteolysis. *International*  
691 *Dairy Journal*, *11*, 455–464.
- 692 Weijers, M., van de Velde, F., Stijnman, A., van de Pijpekamp, A., & Visschers, R. W. (2006). Structure  
693 and rheological properties of acid-induced egg white protein gels. *Food Hydrocolloids*, *20*, 146–  
694 159.
- 695 WHO. (2012). *Guideline: Sodium intake for adults and children*. Geneva, Switzerland: World Health  
696 Organisation.
- 697 Zoon, P. (1993). *Physical properties of cheese*. Madison, WI, USA: Proceedings of the Cheese Research  
698 and Technology Conference.

1 **Figure legends**

2

3 **Fig. 1.** Composition of semi-hard Danish cheeses from experiment 1 with DL-starter culture C1. A: NaCl  
4 content, B: protein content, C: dry matter and D: pH are shown as a function of ripening times (week) for  
5 brining times of 0 h (black), 6 h (dark grey), 12 h (grey) and 24 h (white). The results are shown as the mean  
6  $\pm$  SD,  $n = 3$ . Bars with different letters above differ significantly ( $P < 0.05$ ).

7

8 **Fig. 2.** NaCl distribution in semi-hard Danish cheeses from experiment 2 during ripening as a function of  
9 brining time, for samples from core (open circles) and edge (filled circles), with ripening times of 2 weeks  
10 (A), 7 weeks (B) and 12 weeks (C). Values are shown as mean  $\pm$  SD,  $n = 6$ ; values with different letters above  
11 are significantly different by a 3-factor interaction ( $P < 0.05$ ).

12

13 **Fig. 3.** Textural parameters Axial stress (A) and Hencky strain (B) for cheeses from experiment 2, produced  
14 with DL-starter culture C1 (■, ■, ■) and C2 (■, ■, □) in combination with brining times of 0 h (■, ■), 12 h  
15 (■, ■), and 24 h (■, □) according to ripening times (week). Values are shown as mean  $\pm$  SD,  $n = 24$ .

16

17 **Fig. 4.** Experiment 2. Scanning electron microscopy images (2000  $\times$  magnification) of semi-hard Danish  
18 cheeses after 11 weeks of ripening. A) DL-starter culture C1 with 0 h brining, B) DL-starter culture C1 with 24  
19 h brining, C) DL-starter culture C2 with 0 h brining and D) DL-starter culture C2 with 24 h brining.

20

21 **Fig. 5.** Correlations from experiment 2 of semi-hard Danish cheese NaCl content with textural properties;  
22 Axial stress (A+B+C) and Hencky strain (D+E+F) at fracture for ripening periods of 2 weeks (A+D), 7 weeks  
23 (B+E), and 12 weeks (C+F) for samples from core (open circles) and edge (filled circles). Linear regressions  
24 and the corresponding regression coefficient are given.

25

26 **Fig. 6.** Scanning electron microscopy images (5000 × magnification) of semi-hard Danish cheeses from  
27 experiment 4 receiving 28 h of brining and 12 weeks of ripening, using bovine chymosin (A) or camel  
28 chymosin (B).

29

ACCEPTED MANUSCRIPT

1 **Table 1**

2 Schematic overview of the four cheese experiments along with the main parameters studied, time of  
 3 analysis during ripening and number of replicates. <sup>a</sup>

Experiment	Chymosin type	DL-starter culture	Brining time (h)	Ripening time (weeks)	Cheese replicates	Positions for texture sampling
1	BC	C1	0, 6, 12, 24	1, 2, 7, 12	3	12
		C2	0, 6, 12, 24	1, 2, 7, 12	3	12
2	BC	C1	0, 12, 24	1, 2, 7, 12	2	12
		C2	0, 12, 24	1, 2, 7, 12	2	12
3	BC	C3	6, 12, 24	1, 12	2	12
	CC	C3	6, 12, 24	1, 12	2	12
4	BC	Commercial used at Taulov dairy	0, 28	12	12	2
	CC		10, 15, 28	12	12	2

4

5 <sup>a</sup> Abbreviations are: BC, bovine chymosin; CC, camel chymosin; C1, C2 and C3, DL-starter cultures  
 6 originating from Chr. Hansen A/S.

7

8

9 **Table 2**

10 Content of NaCl (%) and textural properties by axial stress (kPa) and Hencky strain (-) of semi-hard  
 11 Danish cheeses after 12 weeks of ripening from experiment 1 and 3. <sup>a</sup>

12

Parameters	Brining time (h)				F-test
	0	6	12	24	
C1 - experiment 1					P-value
NaCl (%)	0.21±0.03 <sup>d</sup>	1.00±0.09 <sup>c</sup>	1.30±0.15 <sup>b</sup>	1.70±0.10 <sup>a</sup>	<0.001
Stress (kPa)	-*	25.9±9.1 <sup>b</sup>	33.3±1.8 <sup>b</sup>	48.6±18.3 <sup>a</sup>	<0.001
Strain (-)	-*	1.34±0.17 <sup>a</sup>	1.26±0.18 <sup>a</sup>	1.11±0.14 <sup>b</sup>	<0.05
C2 - experiment 1					
NaCl (%)	0.19±0.03 <sup>d</sup>	1.05±0.05 <sup>c</sup>	1.39±0.07 <sup>b</sup>	1.83±0.14 <sup>a</sup>	<0.001
Stress (kPa)	19.8±4.8 <sup>c</sup>	36.5±11.9 <sup>b</sup>	48.4±16.1 <sup>ab</sup>	53.6±18.1 <sup>a</sup>	<0.001
Strain (-)	1.37±0.34 <sup>a</sup>	1.24±0.13 <sup>b</sup>	1.19±0.06 <sup>b</sup>	1.06±0.10 <sup>c</sup>	<0.001
C3 - experiment 3					
NaCl (%)	-**	1.11±0.04 <sup>c</sup>	1.43±0.13 <sup>b</sup>	1.90±0.18 <sup>a</sup>	<0.05
Stress (kPa)	-**	54.5±18.4 <sup>b</sup>	82.5±31.8 <sup>a</sup>	95.9±23.9 <sup>a</sup>	<0.001
Strain (-)	-**	1.30±0.10 <sup>a</sup>	1.25±0.07 <sup>a</sup>	1.07±0.11 <sup>b</sup>	<0.001

13

14 <sup>a</sup> The cheeses were produced with bovine chymosin and DL-starter cultures C1, C2 and C3. Values are  
 15 means ± standard deviation, n=6 (NaCl content), n=36 (textural analysis exp. 1), and n=24 (textural  
 16 analysis exp. 3); values within a row with different superscript letters differ significantly at the level of  
 17 given *P*-value. A single asterisk indicates no textural analysis was performed; a double asterisk indicates  
 18 no non-brined cheeses were produced using DL-starter culture C3.

19

20 **Table 3**

21 Experiment 3 and 4, effect of chymosin type and brining time used for semi-hard Danish cheeses on  
 22 NaCl content, dry matter, pH, and textural properties after 12 weeks of ripening. <sup>a</sup>

23

Chymosin type	Brining time (h)	NaCl (% w/w)	Dry matter (% w/w)	pH	Axial stress (kPa)	Hencky strain (-)
Experiment 3						
BC	6	1.11±0.03 <sup>c</sup>	45.9±0.5 <sup>b</sup>	5.51±0.04 <sup>a</sup>	54.5±18.4 <sup>c</sup>	1.31±0.10 <sup>a</sup>
BC	12	1.43±0.12 <sup>ab</sup>	48.5±1.1 <sup>a</sup>	5.51±0.05 <sup>a</sup>	82.5±31.8 <sup>b</sup>	1.25±0.07 <sup>a</sup>
BC	24	1.90±0.18 <sup>a</sup>	49.2±0.1 <sup>a</sup>	5.37±0.06 <sup>a</sup>	95.9±23.9 <sup>a</sup>	1.08±0.11 <sup>b</sup>
CC	6	1.17±0.04 <sup>bc</sup>	47.6±0.1 <sup>ab</sup>	5.53±0.02 <sup>a</sup>	77.7±25.9 <sup>b</sup>	1.33±0.07 <sup>a</sup>
CC	12	1.48±0.70 <sup>ab</sup>	48.7±0.1 <sup>a</sup>	5.48±0.02 <sup>a</sup>	88.0±35.5 <sup>b</sup>	1.25±0.10 <sup>a</sup>
CC	24	1.79±0.21 <sup>a</sup>	49.6±0.8 <sup>a</sup>	5.47±0.10 <sup>a</sup>	91.6±36.3 <sup>ab</sup>	1.05±0.12 <sup>b</sup>
Experiment 4						
BC	0	0.08±0.05 <sup>c</sup>	46.3±0.1 <sup>a</sup>	5.41±0.02 <sup>b</sup>	28.5±4.5 <sup>c</sup>	1.65±0.12 <sup>a</sup>
BC	28	1.51±0.11 <sup>a</sup>	47.8±0.1 <sup>a</sup>	5.52±0.02 <sup>a</sup>	66.0±16.3 <sup>a</sup>	1.23±0.05 <sup>b</sup>
CC	10	1.20±0.08 <sup>b</sup>	46.5±0.1 <sup>a</sup>	5.55±0.02 <sup>a</sup>	54.0±8.1 <sup>b</sup>	1.16±0.08 <sup>c</sup>
CC	15	1.23±0.11 <sup>b</sup>	47.7±0.1 <sup>a</sup>	5.58±0.02 <sup>a</sup>	62.8±7.6 <sup>a</sup>	1.11±0.08 <sup>c</sup>
CC	28	1.53±0.12 <sup>a</sup>	47.4±0.0 <sup>a</sup>	5.53±0.02 <sup>a</sup>	70.6±10.8 <sup>a</sup>	1.09±0.07 <sup>c</sup>

24

25 <sup>a</sup> Abbreviations are: BC, bovine chymosin; CC, camel chymosin. Values are least squares-means ±  
 26 standard deviation (n = 4, chemical analysis; n = 24, textural analysis); values within a column with  
 27 different superscript letters differ significantly ( $P < 0.05$ )

28

Fig. 1




Fig. 2




Fig. 3





Fig. 4



ACCEPTED

Fig. 5





Fig. 6



ACCEPTED MANUSCRIPT