



UNIVERSITY OF LEEDS

This is a repository copy of *Quantum principle of sensing gravitational waves: From the zero-point fluctuations to the cosmological stochastic background of spacetime*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/127249/>

Version: Accepted Version

Article:

Quiñones, DA orcid.org/0000-0001-5844-3312, Oniga, T, Varcoe, BTH orcid.org/0000-0001-7056-7238 et al. (1 more author) (2017) Quantum principle of sensing gravitational waves: From the zero-point fluctuations to the cosmological stochastic background of spacetime. *Physical Review D*, 96 (4). 044018. ISSN 2470-0010

<https://doi.org/10.1103/PhysRevD.96.044018>

© 2017 American Physical Society. This is an author produced version of a paper published in *Physical Review D*. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Quantum principle of sensing gravitational waves: From the zero-point fluctuations to the cosmological stochastic background of spacetime

Diego A. Quiñones,^{1,*} Teodora Oniga,^{2,†} Benjamin T. H. Varcoe,^{1,‡} and Charles H.-T. Wang^{2,§}

¹*School of Physics and Astronomy, University of Leeds, Leeds LS2 9JT, United Kingdom*

²*Department of Physics, University of Aberdeen, Aberdeen AB24 3UE, United Kingdom*

We carry out a theoretical investigation on the collective dynamics of an ensemble of correlated atoms, subject to both vacuum fluctuations of spacetime and stochastic gravitational waves. A general approach is taken with the derivation of a quantum master equation capable of describing arbitrary confined nonrelativistic matter systems in an open quantum gravitational environment. It enables us to relate the spectral function for gravitational waves and the distribution function for quantum gravitational fluctuations and to indeed introduce a new spectral function for the zero-point fluctuations of spacetime. The formulation is applied to two-level identical bosonic atoms in an off-resonant high- Q cavity that effectively inhibits undesirable electromagnetic delays, leading to a gravitational transition mechanism through certain quadrupole moment operators. The overall relaxation rate before reaching equilibrium is found to generally scale collectively with the number N of atoms. However, we are also able to identify certain states of which the decay and excitation rates with stochastic gravitational waves and vacuum spacetime fluctuations amplify more significantly with a factor of N^2 . Using such favourable states as a means of measuring both conventional stochastic gravitational waves and novel zero-point spacetime fluctuations, we determine the theoretical lower bounds for the respective spectral functions. Finally, we discuss the implications of our findings on future observations of gravitational waves of a wider spectral window than currently accessible. Especially, the possible sensing of the zero-point fluctuations of spacetime could provide an opportunity to generate initial evidence and further guidance of quantum gravity.

I. INTRODUCTION

A. Background

Recently, there has been considerable interest in experimenting with gravity using quantum instrument [1, 2], to better explore its theory, predictions, and applications in uncontested regimes [3–5]. This trend has been established through laboratory measurements of the gravitational acceleration by ultracold atom interferometers with accuracies exceeding that achievable by conventional gravimeters [6]. The realization of such quantum gravity and gravity gradient sensors as portable devices would herald a new level of quantum applications in engineering and technology [7].

For physicists, it has been hoped that major progress across a range of areas could be enabled by the availability of precision quantum gravity sensors. These include testing of the equivalence principle as a foundation of general relativity in the quantum domain [8], confronting macroscopic quantum behaviour with a microgravity environment [9], and assessing a long range effect such as

gravitational time dilation on quantum coherence [10]. Above all, quantum experiments may promise fundamental breakthroughs by offering a unique opportunity to access phenomena from quantum gravity and Planck-scale physics that would otherwise have required an inconceivable 10^{19} GeV Grand Unified Theory (GUT)-scale particle collider [11–14].

There are a number of counterintuitive effects unavailable in the classical domain to justify novelties and advantages of quantum measurements. For Bell’s inequality related experiments [15, 16], quantum entanglement and nonlocality are the key. More relevant to precision measurements, in the case of atom interferometry metrology, the short matter wavelength underpins a fine scale resolution. When atom interferometry is applied to gravity measurements including gradiometry and spacetime fluctuations, the slowness of the motion for cold atoms contributes to a longer interaction time to build up effects in terms of phase shift due to acceleration and decoherence due to granulation [8].

For quantum systems with a large number of correlated particles, nontrivial collective behaviours can arise [17, 18]. An early remarkable example shows up in Hanbury Brown and Twiss’s intensity interferometry [19]. Collective behaviours can give rise to significant amplifications of quantum effects associated with a large number of correlated particles, as exemplified in Dicke’s seminal superradiance [20]. In this paper, we will analyze previously unexploited collective quantum interactions

* pydaqo@leeds.ac.uk

† t.oniga@abdn.ac.uk

‡ b.varcoe@leeds.ac.uk

§ Corresponding author, c.wang@abdn.ac.uk

between a large number of particles with gravitational waves that may be relevant for their detections beyond the existing observation windows e.g. in the frequency domain of fundamental, astronomical and cosmological interest [21–24].

In particular, we consider stochastic gravitational waves of frequencies higher than the existing detection range, that may have a primordial origin and carry imprints of postinflation structure formation processes [21, 25–27]. The nature and possible observation of such stochastic gravitational waves have attracted significant recent attention since the groundbreaking detection of gravitational waves by the LIGO and Virgo collaborations [28] with the anticipation of broader gravitational wave astronomy and cosmology [23, 29, 30]. In what follows, we present a theoretical analysis with Gaussian stochastic gravitational waves [31] having a generic distribution function $N_{\text{gw}}(\omega)$ and the equivalent spectral function $\Omega_{\text{gw}}(f)$. The atoms are assumed to be bosonic so that the collective quantum dynamics of a large number of atoms occupying a common state is possible.

The effects of spacetime curvature on individual atoms have been considered in a number of works [32–36], showing slight shifts in the energy spectrum and weak transitions between the energy levels of the atoms induced by gravitational interactions. Gravity has been proposed to induce decoherence in atomic states [5, 37, 38], making the analysis of the evolution of atomic states a potential method for the detection of gravitational fluctuations [39–42]. However, these effects have also been found to be generally too small to be measured in practice [33].

It can be noted that Rydberg atoms [43, 44] have received particular attention for atom-gravity interactions due to a number of their distinct physical characteristics they possess: Their large principal quantum number n means that the effective mass quadrupole moment, which provides external gravitational wave coupling and scales with n^4 , can be greatly enhanced. These atoms are very stable having extremely long lifetimes of the internal transitions up to the order of 1 second allowing us to neglect them in many practical situations [45]. Furthermore, increasingly large numbers of Rydberg atoms with high principal quantum numbers $n > 10^2$ have been produced in laboratories [46]. The investigations of Rydberg atoms as a many-body system have often featured in the recent literature [47] with their collective quantum dynamics receiving current research interest both theoretically [48–51] and experimentally [52–58].

B. Summary

Rather than considering gravitational interactions with individual atoms, here we investigate the coherent quantum interactions for a large number of correlated atoms with gravitational waves. We find that

the resulting collective interactions can lead to amplified quadrupole transition rates of atomic states in a fashion akin to superradiance [20].

This collective amplification could be compared and combined with the application of Rydberg atoms, which are more sensitive as individual atoms to spacetime perturbations [42, 59]. However, their larger size can also decrease the number density and increase dipole-dipole interactions of the atoms. Consequently, we find that smaller atoms can provide more collective strength for gravitational wave interactions as described below, with specific numerical examples shown in Fig. 1, using an ensemble of heliumlike atoms for detecting high frequency stochastic gravitational waves and spacetime fluctuations. Our investigations may form a basis for a possible future detection scheme for gravitational waves with a wider spectral window, especially towards higher frequencies than currently accessible, which may be of fundamental and cosmological importance.

In the present theoretical investigation, we hypothesize for simplicity a setup of N identical two-level atoms with a quadrupole moment transition frequency $f = \omega/2\pi$ used to evaluate the distribution function $N_{\text{gw}}(\omega)$ and the equivalent spectral function $\Omega_{\text{gw}}(f)$ for the stochastic gravitational waves. Further refinement is possible to reduce the present level of idealization but will be deferred to future work. The system of atoms is considered to be placed in an off-resonant high- Q cavity with controlled boundary conditions that effectively inhibits undesirable electromagnetic delays [60–62]. The presence of the high- Q cavity can also change the quantum electrodynamics of the atom to modify the interaction between the nucleus and the electron so as to increase the quadrupole interaction while suppressing the dipole-dipole interactions between the atoms [63].

In Sec. II, we further the theoretical foundation of how general confined nonrelativistic matter interacts with an open gravitational environment, relevant to, for example, a system of atoms in a cavity attracting much recent attention. Specifically, we provide a general master equation through Eq. (14) with negligible self-interactions to focus on the collective interactions between such a matter system and gravitational waves in the environment due to both vacuum and stochastic fluctuations.

In Sec. III, the theoretical formulation is applied to a two-level atom that allows us to derive specific transition mechanisms through the quadrupole moment operator with respect to the two atomic states given by Eq. (28), satisfying a set of selection rules (31). In particular, we derive a formula for the gravitational transition rate Γ_0 in Eq. (29) for such a two-level atom. Unsurprisingly, the transition rate Γ_0 for a single atom is too small to be measured under typical laboratory conditions, as has been pointed out even for Rydberg atoms [33].

In Sec. IV, we invoke the generality of the formalism developed in Sec. II to extend the interaction between a




FIG. 1. Theoretical lower bounds for the spectral functions of stochastic gravitational waves $\Omega(f)$ determined by Eq. (66) that could be detected using an ensemble of correlated two-level atoms contained in an off-resonant high- Q cavity of volume L^3 and measurement time τ . Specifically, we consider heliumlike atoms with atomic quantum numbers $n_1 = n_2 = 2$, $l_1 = l_2 = 1$ and $-m_1 = m_2 = 1$, with a transition frequency induced by a Zeeman-type shift. For conventional stochastic gravitational waves, we have $\Omega(f) = \Omega_{\text{gw}}(f)$ with $\tau = \tau_{\text{gw}}$, whereas for zero-point, i.e. vacuum, metric fluctuations, we have $\Omega(f) = \Omega_{\text{vac}}(f)$ with $\tau = \tau_{\text{vac}}$ introduced in this work by Eqs. (62) and (64). In both cases, we choose $\tau = 1$ sec, min, and h as possible (future) transition times for the atoms with $L = 1$ and 10 m in generating the above diagram. It suggests, for example, measuring the zero-point fluctuations of spacetime would require a 1 m^3 cavity with a measurement time of 1 sec at $f = 1$ THz approximately. Note that regions where the values of $\Omega_{\text{gw}}(f)$ are below that of $\Omega_{\text{vac}}(f)$ would present a less stochastic gravitational wave signal to vacuum fluctuations “noise” ratio and hence the detection of the latter would be more likely using the discussed quantum methods. For comparison, we have superimposed the expected spectral functions of stochastic gravitational waves from various sources such as inflation, up to frequencies dominated by spacetime fluctuations, that may be detected by different methods such as LIGO from Ref. [21] and its references.

two-level atom and gravity to accommodate a correlated ensemble of N such atoms. In this case, as illustrated in Fig. 2, we find an overall factor of N collective amplifi-




FIG. 2. Simulations with N atoms using the master equation (1) with the quantum dissipator (37) for the correlated two-level atoms and negligible matter interaction H_{int} . From Eq. (45) the relaxation time is approximately given by $\tau_N = (N\Gamma_0)^{-1}$. The initial state of the forms (32) and (33) at $t = 0$ is given by $|0, N\rangle = |p = N + 1\rangle$ with all N atoms occupying the excited atomic level. During the dynamical evolution, the density matrix remains diagonal of the form (39) and relaxes into an inverse exponential profile represented by Eq. (48), yielding the asymptotic height of $\rho_{p=1} = [N_{\text{gw}}(\omega_0)(1 - e^{-N/N_{\text{gw}}(\omega_0)})]^{-1}$. The stochastic gravitational wave distribution function evaluated at the atomic transition frequency ω_0 is chosen to be $N_{\text{gw}}(\omega_0) = 1$. Accordingly, in plot (a) with $N = 100$ atoms, the initial state corresponding to the spike of height 1 at $(p, \Gamma_0 t) = (101, 0)$ relaxes into an inverse exponential profile peaked at $(p, \Gamma_0 t) = (1, 0.1)$ on time scale of $\tau_N = 0.01/\Gamma_0$. Similarly, in plot (b) with $N = 200$ atoms, the initial state corresponding to the spike of height 1 at $(p, \Gamma_0 t) = (201, 0)$ relaxes into an inverse exponential profile similarly peaked at $(p, \Gamma_0 t) = (1, 0.1)$ but on a halved time scale of $\tau_N = 0.005/\Gamma_0$.

cation of the transition rate from any initial state to the final equilibrium state satisfying the distribution in the large- N limit (48).

In Sec. V, we establish a relation between the spectral function $\Omega_{\text{gw}}(f)$ for gravitational waves and distribution function $N_{\text{gw}}(\omega)$ for open quantum gravitational systems and introduce a new spectral function $\Omega_{\text{vac}}(f)$




FIG. 3. Simulations of short-time collective spontaneous decays due to vacuum fluctuations of spacetime without additional stochastic gravitational waves so that $N_{\text{gw}}(\omega_0) = 0$. In plot (a) with $N = 100$ atoms, the initial state corresponds to the spike of height 1 at $(p, t) = (50, 0)$. As shown, it decays into adjacent lower states with a relaxation time $\tau_{\text{vac}} = 4/(N^2\Gamma_0) = 0.0004/\Gamma_0$ according to Eqs. (61) and (62). In plot (b) with $N = 200$ atoms, the initial state corresponds to the spike of height 1 at $(p, t) = (100, 0)$. For a doubled N , this state decays into adjacent lower states with a quarter of the preceding relaxation time with $\tau_{\text{vac}} = 4/(N^2\Gamma_0) = 0.0001/\Gamma_0$.

for the zero-point, i.e. quantum vacuum, fluctuations of spacetime. Although we have introduced a spectral function for vacuum fluctuations of spacetime $\Omega_{\text{vac}}(f)$ in Eq. (64), it should be emphasized that the detection of zero-point fluctuations of spacetime requires a quantum detector through decoherence as classical detectors have already decohered and so can no longer provide this function.

Indeed, the possibility of sensing the zero-point fluctuations of spacetime provides a unique physical characteristics for the collective quantum process which could lead to initial evidence and further guidance of quantum gravity. For optimal measurements, we have identified favourable states of which the decay and excitation rates with gravitational waves and spacetime fluctuations scale amplify with a factor of N^2 , as described by Eqs. (60) and (61) and illustrated in Figs. 2 and 3.

We then determine the theoretical lower bounds given by Eq. (66) for the spectral functions of gravitational

waves that could in principle be detected using an ensemble of identical two-level atoms given their excitation states and the size of the cavity as well as measurement time. The lower bound relation (66) applies to both conventional stochastic gravitational waves and novel zero-point spacetime fluctuations. Importantly, we find that, although as individual atoms with larger quadrupole moments, Rydberg atoms may have stronger interactions with gravitational waves, smaller atoms can nonetheless provide more collective strength for such interactions thanks to a higher possible number density of the atoms in a cavity. Specific numerical examples using an ensemble of heliumlike atoms are studied, leading to new prospects for detecting high frequency stochastic gravitational waves and spacetime fluctuations as shown in Fig. 1.

Finally, in Sec. VI, we conclude this work with an additional discussion.

II. GRAVITATIONAL MASTER EQUATION FOR CONFINED NONRELATIVISTIC SYSTEMS

In the interaction picture with relativistic units where $c = 1$, the general gravitational master equation for matter that may be interacting and relativistic has recently been obtained in Ref. [5] to be

$$\dot{\rho} = -\frac{i}{\hbar}[H_{\text{int}}, \rho] + \mathcal{D}\rho \quad (1)$$

with H_{int} describing matter interaction with itself or other fields not part of the gravitational environment, and the non-Markovian quantum dissipator

$$\begin{aligned} \mathcal{D}\rho = & -\frac{8\pi G}{\hbar} \int \frac{d^3k}{2(2\pi)^3k} \times \\ & \left\{ \int_0^t dt' e^{-ik(t-t')} ([\tau_{ij}^\dagger(\mathbf{k}, t), \tau_{ij}(\mathbf{k}, t')\rho] \right. \\ & \left. + N_{\text{gw}}(\omega_k) [\tau_{ij}^\dagger(\mathbf{k}, t), [\tau_{ij}(\mathbf{k}, t'), \rho]] + \text{H.c.} \right\} \quad (2) \end{aligned}$$

describing in general nonunitary statistical quantum evolution of the matter system as a result of dissipative interactions with the environment [5, 27]. Here, $N_{\text{gw}}(\omega)$ is a general distribution function for the fluctuating gravitational environment assumed to be Gaussian. If this environment is in thermal equilibrium at temperature T then $N_{\text{gw}}(\omega)$ is given by the Planck distribution

$$N_{\text{gw}}(\omega) = \frac{1}{e^{\hbar\omega/k_B T} - 1}. \quad (3)$$

However, we will be considering generic stochastic gravitational waves described by $N_{\text{gw}}(\omega)$ not restricted to the Planck distribution(3).

Since typical compact nonrelativistic systems have spatial extensions much smaller than the effectively coupled

gravitational wavelength [64, 65], we have $e^{-i\mathbf{k}\cdot\mathbf{r}} \approx 1$ and therefore

$$\begin{aligned}\tau_{ij}(\mathbf{k}, t) &= \int \tau_{ij}(\mathbf{r}, t) e^{-i\mathbf{k}\cdot\mathbf{r}} d^3x \\ &\approx \frac{1}{2} \ddot{\mathbf{F}}_{ij}^{\text{TT}}(\mathbf{k}, t)\end{aligned}\quad (4)$$

where

$$\mathbf{F}_{ij}^{\text{TT}}(\mathbf{k}, t) = P_{ijkl}(\mathbf{k}) \mathbf{F}_{kl}(t) \quad (5)$$

and

$$\mathbf{F}_{ij}(t) = \frac{1}{3} \int d^3x Q_{ij}(\mathbf{r}) T^{00}(\mathbf{r}, t) \quad (6)$$

in terms of the transverse-traceless (TT) projection operator P_{ijkl} [5, 64] and the quadrupole moment tensor

$$Q_{ij} = 3x^i x^j - \delta_{ij} r^2. \quad (7)$$

It is useful to introduce the frequency domain reduced quadrupole moments $\mathbf{F}_{ij}(\omega)$ for some positive frequencies ω by writing

$$\mathbf{F}_{ij}(t) = \sum_{\omega} [\mathbf{F}_{ij}(\omega) e^{-i\omega t} + \mathbf{F}_{ij}^{\dagger}(\omega) e^{i\omega t}]. \quad (8)$$

Let us consider the following integral in Eq. (2),

$$\begin{aligned}&\int_0^t dt' e^{-ik(t-t')} \tau_{ij}(\mathbf{k}, t') \\ &= -\sum_{\omega} \frac{\omega^2}{2} \mathbf{F}_{ij}^{\text{TT}}(\omega, \mathbf{k}) \int_0^t dt' e^{-ik(t-t')} e^{-i\omega t'} \\ &\quad - \sum_{\omega} \frac{\omega^2}{2} \mathbf{F}_{ij}^{\text{TT}\dagger}(\omega, \mathbf{k}) \int_0^t dt' e^{-ik(t-t')} e^{i\omega t'}\end{aligned}\quad (9)$$

which is nonlocal in time representing the non-Markov memory effect.

To apply the Markov approximation, valid for physical time scales where short-term transient memories are damped away [66], we change the time variable from t' to s using $t' = t - s$ in the above to get

$$\begin{aligned}\int_0^t dt' e^{-ik(t-t')} e^{\pm i\omega t'} &\approx e^{\pm i\omega t} \int_0^{\infty} ds e^{-i(k \pm \omega)s} \\ &= e^{\pm i\omega t} \left[\pi \delta(k \pm \omega) - i \mathcal{P} \frac{1}{k \pm \omega} \right]\end{aligned}$$

where the Sokhotski-Plemelj theorem [see Eq. (A15) below] has been used in the last step.

Using the above and neglecting the the Cauchy principal values, as they do not contribute to quantum deco-

herence and dissipation, we see that Eq. (9) becomes

$$\begin{aligned}&\int_0^t dt' e^{-ik(t-t')} \tau_{ij}(\mathbf{k}, t') \\ &= -\sum_{\omega} \frac{\pi \omega^2}{2} \mathbf{F}_{ij}^{\text{TT}}(\omega, \mathbf{k}) e^{-i\omega t} \delta(k - \omega) \\ &\quad - \sum_{\omega} \frac{\pi \omega^2}{2} \mathbf{F}_{ij}^{\text{TT}\dagger}(\omega, \mathbf{k}) e^{i\omega t} \delta(k + \omega)\end{aligned}\quad (10)$$

where the second term does not contribute as $k \geq 0$ and $\omega > 0$. Using Eqs. (4) and (8), we have

$$\begin{aligned}\tau_{ij}^{\dagger}(\mathbf{k}, t) &\approx \frac{1}{2} \ddot{\mathbf{F}}_{ij}^{\text{TT}\dagger}(\mathbf{k}, t) \\ &= -\frac{\omega^2}{2} [\mathbf{F}_{ij}^{\text{TT}}(\omega, \mathbf{k}) e^{-i\omega t} + \mathbf{F}_{ij}^{\text{TT}\dagger}(\omega, \mathbf{k}) e^{i\omega t}].\end{aligned}\quad (11)$$

Substituting Eqs. (10) and (11) into the dissipator (2) and applying the rotating wave approximation we have

$$\begin{aligned}\mathcal{D}\rho &= -\frac{G\omega^5}{8\pi\hbar} \int d\Omega(\mathbf{k}) \left\{ ([\mathbf{F}_{ij}^{\text{TT}}(\omega, \mathbf{k}), \mathbf{F}_{ij}^{\text{TT}}(\omega, \mathbf{k})\rho] \right. \\ &\quad \left. + N_{\text{gw}}(\omega) [\mathbf{F}_{ij}^{\text{TT}\dagger}(\omega, \mathbf{k}), [\mathbf{F}_{ij}^{\text{TT}}(\omega, \mathbf{k}), \rho]]) + \text{H.c.} \right\} \\ &= \frac{G\omega^5}{4\pi\hbar} \int d\Omega(\mathbf{k}) P_{ijkl}(\mathbf{k}) \times \\ &\quad \left\{ (N_{\text{gw}}(\omega) + 1) [\mathbf{F}_{ij}(\omega) \rho \mathbf{F}_{kl}^{\dagger}(\omega) - \frac{1}{2} \{ \mathbf{F}_{ij}^{\dagger}(\omega) \mathbf{F}_{kl}(\omega), \rho \}] \right. \\ &\quad \left. + N_{\text{gw}}(\omega) [\mathbf{F}_{ij}^{\dagger}(\omega) \rho \mathbf{F}_{kl}(\omega) - \frac{1}{2} \{ \mathbf{F}_{ij}(\omega) \mathbf{F}_{kl}^{\dagger}(\omega), \rho \}] \right\}\end{aligned}\quad (12)$$

summing over $\omega > 0$ and $i, j, k, l = 1, 2, 3$. With the aid of the formulae [65]

$$\int k_i k_j d\Omega = \frac{4\pi}{3} k^2 \delta_{ij}$$

$$\int k_i k_j k_k k_l d\Omega = \frac{4\pi}{15} k^2 (\delta_{ij} \delta_{kl} + \delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk})$$

we can calculate that

$$\int P_{ijkl}(\mathbf{k}) d\Omega = \frac{4\pi}{15} (3\delta_{ik} \delta_{jl} + 3\delta_{il} \delta_{jk} - 2\delta_{ij} \delta_{kl}). \quad (13)$$

Substituting Eq. (13) into Eq. (12) and using the tracelessness of \mathbf{F}_{ij} , we obtain the following general dissipator

$$\begin{aligned}\mathcal{D}\rho &= \frac{2G\omega^5}{5\hbar} \times \\ &\quad \left\{ (N_{\text{gw}}(\omega) + 1) [\mathbf{F}_{ij}(\omega) \rho \mathbf{F}_{ij}^{\dagger}(\omega) - \frac{1}{2} \{ \mathbf{F}_{ij}^{\dagger}(\omega) \mathbf{F}_{ij}(\omega), \rho \}] \right. \\ &\quad \left. + N_{\text{gw}}(\omega) [\mathbf{F}_{ij}^{\dagger}(\omega) \rho \mathbf{F}_{ij}(\omega) - \frac{1}{2} \{ \mathbf{F}_{ij}(\omega) \mathbf{F}_{ij}^{\dagger}(\omega), \rho \}] \right\}\end{aligned}\quad (14)$$

of the Lindblad form for any nonrelativistic matter systems with frequency domain reduced quadrupole moments $\mathbf{F}_{ij}(\omega)$. For particles in an isotropic harmonic potential with $N_{\text{gw}}(\omega) = 0$, Eq. (14) recovers the corresponding dissipator discussed in Ref. [27].

III. GRAVITATIONALLY INDUCED TRANSITIONS OF QUANTUM STATES

The theoretical framework established above is applied in this section to the quantum gravitational decoherence and radiation of a real scalar field ϕ with mass m and the associated inverse reduced Compton wavelength $\mu = m/\hbar$, subject to an external potential $\nu(\mathbf{r})$ described by the Lagrangian density

$$\mathcal{L} = -\frac{1}{2}g^{\alpha\beta}\phi_{,\alpha}\phi_{,\beta} - \left(\frac{1}{2} + \nu\right)\mu^2\phi^2. \quad (15)$$

To consider the nonrelativistic dynamics of the scalar field representing nearly Newtonian particles we assume the potential energy to be much less than the rest mass energy so that $\nu \ll 1$ and the total energy density is given approximately by

$$T^{00} = \frac{1}{2}(\dot{\phi}^2 + \mu^2\phi^2). \quad (16)$$

The unperturbed part of Eq. (15) satisfies quantum field equation

$$\ddot{\phi} = \nabla^2\phi - (1 + 2\nu)\mu^2\phi \quad (17)$$

having solutions of the convenient form

$$\phi = \sqrt{\frac{\hbar}{2\omega_n}} a_n \psi_n(\mathbf{r}) e^{-i\omega_n t} + \text{H.c.} \quad (18)$$

with some operators a_n and orthogonal complex functions $\psi_n(\mathbf{r})$. Hence Eq. (17) reduces formally to the time-independent Schrödinger equation

$$-\frac{\hbar^2}{2m}\nabla^2\psi_n + V\psi_n = E_n\psi_n \quad (19)$$

where $V = m\nu(\mathbf{r})$ and

$$E_n = \frac{1}{2m}(\hbar^2\omega_n^2 - m^2) \quad (20)$$

represent the potential and eigenenergies respectively.

Let us consider the atom potential energy $V(r)$ In the nonrelativistic domain, Eq. (18) becomes

$$\phi = \frac{\hbar}{\sqrt{2m}} (a_n e^{-i\omega_n t} + a_n^\dagger e^{i\omega_n t}) \psi_n(\mathbf{r})$$

using the multiple indices $\mathbf{n} = (n, l, m)$ and the atom wave functions

$$\psi_n(\mathbf{r}) = R_{nl}(r)Y_l^m(\theta, \phi)$$

with

$$\omega_n = \mu + E_n/\hbar$$

arising from the nonrelativistic limit of Eq. (20), where the corresponding ladder operators a_n and a_n^\dagger are annihilation and creation operators respectively. From Eqs. (6) and (16) we have

$$\mathcal{I}_{ij} = \frac{m}{3} \sum_{\substack{\mathbf{n}, \mathbf{n}' \\ (\omega_n > \omega_{n'})}} (\mathbf{n}'|Q_{ij}|\mathbf{n}) a_n^\dagger a_n e^{-i(\omega_n - \omega_{n'})t} + \text{H.c.} \quad (21)$$

where any irrelevant time-independent parts with $\omega_n = \omega_{n'}$ have been left out and for simplicity we have adopted the quantum mechanics styled notation

$$(\mathbf{n}'|Q_{ij}|\mathbf{n}) = \int d^3x Q_{ij} \psi_{n'}(\mathbf{r}) \psi_n(\mathbf{r}) \quad (22)$$

where Dirac's brackets are denoted by $(\cdot|\cdot)$ and $|\cdot\rangle$, as opposed to $\langle\cdot|$ and $|\cdot\rangle$ reserved for the main quantum states of our present discussion based on quantum field theory. Comparing Eqs. (8) and (21) we see that

$$\mathcal{I}_{ij}(\omega_n - \omega_{n'}) = \frac{m}{3} (\mathbf{n}'|Q_{ij}|\mathbf{n}) a_n^\dagger a_n \quad (23)$$

$$\mathcal{I}_{ij}^\dagger(\omega_n - \omega_{n'}) = \frac{m}{3} (\mathbf{n}'|Q_{ij}|\mathbf{n})^* a_n^\dagger a_n \quad (24)$$

for $\omega_n > \omega_{n'}$.

Let us apply the above general setup to a two-level atom model with states $|1\rangle = |n_1, l_1, m_1\rangle$ and $|2\rangle = |n_2, l_2, m_2\rangle$ having respective eigen energies $E_1 = \hbar\omega_1$ and $E_2 = \hbar\omega_2$ and the transition frequency $\omega_0 = \omega_2 - \omega_1 > 0$. It is useful to denote these two states by

$$|1\rangle = a_1^\dagger|0\rangle, \quad |2\rangle = a_2^\dagger|0\rangle$$

with the ladder operators

$$\sigma^+ = |2\rangle\langle 1| = a_2^\dagger a_1, \quad \sigma^- = |1\rangle\langle 2| = a_1^\dagger a_2.$$

Introducing the transition quadrupole moment

$$q_{ij} = \langle 1|Q_{ij}|2\rangle \quad (25)$$

we can express Eqs. (23) and (24) as

$$\mathcal{I}_{ij}(\omega_0) = \frac{m}{3} q_{ij} a_1^\dagger a_2 = \frac{m}{3} q_{ij} \sigma^- \quad (26)$$

$$\mathcal{I}_{ij}^\dagger(\omega_0) = \frac{m}{3} q_{ij} a_2^\dagger a_1 = \frac{m}{3} q_{ij}^* \sigma^+. \quad (27)$$

Substituting Eqs. (26) and (27) into Eq. (14) we obtain the dissipator for the two-level atom to be

$$\begin{aligned} \mathcal{D}\rho = & \Gamma_0(N_{\text{gw}}(\omega_0) + 1) \left[\sigma^- \rho \sigma^+ - \frac{1}{2} \{ \sigma^+ \sigma^-, \rho \} \right] \\ & + \Gamma_0 N_{\text{gw}}(\omega_0) \left[\sigma^+ \rho \sigma^- - \frac{1}{2} \{ \sigma^- \sigma^+, \rho \} \right] \end{aligned} \quad (28)$$

where we have introduced the gravitational spontaneous emission rate

$$\Gamma_0 = \frac{2Gm^2\omega_0^5 q^2}{45\hbar} \quad (29)$$

in terms of the coupling coefficient related to the squared modulus of the quadrupole moment of the two-level atom

$$q^2 = q_{ij}^* q_{ij} \quad (30)$$

subject to the following selection rules [33, 67],

$$l_1 + l_2 \geq 2, \quad \Delta l = 0, \pm 2, \quad \Delta m = 0, \pm 1, \pm 2 \quad (31)$$

where $\Delta l = l_2 - l_1$, $\Delta m = m_2 - m_1$, outside of which we have $q^2 = 0$.

An alternative derivation of Eq. (28) based on a more conventional quantum mechanical treatment is given in appendix A. Quantum field theory as the main approach of this work has the advantage of describing an arbitrary number of bosonic and fermionic particles and their collective behaviours more naturally and effectively.

IV. GRAVITATIONAL WAVE INTERACTIONS WITH CORRELATED TWO-LEVEL ATOMS

An N -particle two-level state is given by

$$|N_1, N_2\rangle = \frac{a_1^{\dagger N_1} a_2^{\dagger N_2}}{\sqrt{N_1!} \sqrt{N_2!}} |0\rangle \quad (32)$$

with $N_1 + N_2 = N$. There are $N + 1$ such states, which can be represented with

$$|p\rangle = |N - p + 1, p - 1\rangle \quad (33)$$

for $p = 1, 2, \dots, N + 1$, so that $|1\rangle = |N, 0\rangle, |2\rangle = |N - 1, 1\rangle, \dots, |N + 1\rangle = |0, N\rangle$. It follows from

$$\begin{aligned} \langle N'_1, N'_2 | a_1^\dagger a_2 | N_1, N_2 \rangle &= \sqrt{(N_1 + 1) N_2} \delta_{N'_1, N_1 + 1} \delta_{N'_2, N_2 - 1} \\ \langle N'_1, N'_2 | a_2^\dagger a_1 | N_1, N_2 \rangle &= \sqrt{N_1 (N_2 + 1)} \delta_{N'_1, N_1 - 1} \delta_{N'_2, N_2 + 1} \end{aligned}$$

that

$$\begin{aligned} \langle p' | a_1^\dagger a_2 | p \rangle &= \sqrt{(p - 1)(N - p + 2)} \delta_{p', p - 1} \\ \langle p' | a_2^\dagger a_1 | p \rangle &= \sqrt{p(N - p + 1)} \delta_{p', p + 1}. \end{aligned}$$

Using the above relations, we see that Eqs. (23) and (24) become

$$\mathbf{f}_{ij}(\omega_0) = \frac{m}{3} q_{ij} \Sigma^- \quad (34)$$

$$\mathbf{f}_{ij}^\dagger(\omega_0) = \frac{m}{3} q_{ij}^* \Sigma^+ \quad (35)$$

using the $(N + 1)$ -dimensional ladder operators

$$\Sigma_{p'p}^+ = \Sigma_{pp'}^- = \sqrt{p(N - p + 1)} \delta_{p', p + 1}. \quad (36)$$

Substituting Eqs. (34) and (35) into Eq. (14), we obtain the dissipator for the correlated N two-level atoms

$$\begin{aligned} \mathcal{D}\rho &= \Gamma_0 (N_{\text{gw}}(\omega_0) + 1) [\Sigma^- \rho \Sigma^+ - \frac{1}{2} \{\Sigma^+ \Sigma^-, \rho\}] \\ &\quad + \Gamma_0 N_{\text{gw}}(\omega_0) [\Sigma^+ \rho \Sigma^- - \frac{1}{2} \{\Sigma^- \Sigma^+, \rho\}]. \end{aligned} \quad (37)$$

For $N = 1$, we have the reduction $\Sigma^\pm = \sigma^\pm$. However, in general, we see that

$$\Sigma_{p\pm 1p}^\pm \approx \frac{N}{4}, \quad \text{for } p \approx \frac{N}{2} \quad (38)$$

with a large N , providing an N^2 amplification for Eq. (37) compared with Eq. (28).

Consider a diagonal density matrix

$$\rho_{pq} = \rho_p \delta_{p,q} \quad (39)$$

with the normalization

$$\sum_{p=1}^{N+1} \rho_p = 1 \quad (40)$$

from which we can calculate the following:

$$\begin{aligned} (\Sigma^+ \Sigma^- \rho)_{pq} &= (p - 1)(N - p + 2) \rho_p \delta_{p,q} \\ (\Sigma^- \Sigma^+ \rho)_{pq} &= p(N - p + 1) \rho_p \delta_{p,q} \\ (\Sigma^- \rho \Sigma^+)_{pq} &= p(N - p + 1) \rho_{p+1} \delta_{p,q} \\ (\Sigma^+ \rho \Sigma^-)_{pq} &= (p - 1)(N - p + 2) \rho_{p-1} \delta_{p,q} \\ (\rho \Sigma^- \Sigma^+)_{pq} &= p(N - p + 1) \rho_p \delta_{p,q} \\ (\rho \Sigma^+ \Sigma^-)_{pq} &= (p - 1)(N - p + 2) \rho_p \delta_{p,q}. \end{aligned}$$

Substituting the above relations into Eq. (37), we have

$$(\mathcal{D}\rho)_{pq} = \Gamma_0 D_p \delta_{p,q} \quad (41)$$

where

$$\begin{aligned} D_p &= (1 + N_{\text{gw}}(\omega_0)) p(N - p + 1) \rho_{p+1} \\ &\quad - (1 + N_{\text{gw}}(\omega_0)) (p - 1)(N - p + 2) \rho_p \\ &\quad + N_{\text{gw}}(\omega_0) (p - 1)(N - p + 2) \rho_{p-1} \\ &\quad - N_{\text{gw}}(\omega_0) p(N - p + 1) \rho_p. \end{aligned} \quad (42)$$

The trace of Eq. (41) vanishes

$$\text{Tr}(\mathcal{D}\rho) = \sum_{p=1}^{N+1} (\mathcal{D}\rho)_{pp} = 0 \quad (43)$$

as required for the consistency of Eq. (40) under time evolution. Denoting by $\rho(s) = N\rho_p$, $D(s) = ND_p$ using

$s = (p - 1)/N$ and $\epsilon = 1/N$ with $0 \leq s \leq 1$, we see that Eq. (42) becomes

$$\begin{aligned} D(s + \epsilon) &= [(s + \epsilon)(1 - s)\rho(s + 2\epsilon) \\ &\quad - s(1 - (s - \epsilon))\rho(s + \epsilon)]/\epsilon^2 \\ &\quad + N_{\text{gw}}(\omega_0)[(s + \epsilon)(1 - s)(\rho(s + 2\epsilon) - \rho(s + \epsilon)) \\ &\quad - s(1 - (s - \epsilon))(\rho(s + \epsilon) - \rho(s))]/\epsilon^2. \end{aligned}$$

For a large particle number $N \gg 1$ and hence $\epsilon \ll 1$, the above tends to the continuous limit

$$D(s) = \{s(1 - s)[N\rho(s) + N_{\text{gw}}(\omega_0)\rho'(s)]\}' \quad (44)$$

where $(\prime) = \partial_s$. Then the discrete dissipator (37) becomes the following continuous ‘‘diffuser,’’

$$\mathcal{D}\rho(s) = \Gamma_0 \{s(1 - s)[N\rho(s) + N_{\text{gw}}(\omega_0)\rho'(s)]\}' \quad (45)$$

with possible collective amplification by a factor up to N^2 represented by $(\prime) \sim N$ and $(\prime\prime) \sim N^2$. In terms of Eq. (44), the traceless condition (43) then becomes

$$\int_0^1 D(s) ds = 0. \quad (46)$$

For a stationary finite solution with $\mathcal{D}\rho = 0$ and $N_{\text{gw}}(\omega_0) > 0$, Eq. (45) reduces to

$$N_{\text{gw}}(\omega_0)\rho'(s) = -N\rho(s) \quad (47)$$

yielding

$$\rho(s) = \frac{e^{-s/w}}{w(1 - e^{-1/w})} \quad (48)$$

where $w = N_{\text{gw}}(\omega_0)/N$. See Fig. 3 for illustrative numerical examples of dynamical relaxations into the stationary states described above.

V. OPTIMAL GRAVITATIONAL WAVE INTERACTIONS WITH CORRELATED ATOMS

In this section, we would like to relate the above theoretical description to physically relevant values. To this end, let us first relate the distribution function $N_{\text{gw}}(\omega)$ to the spectral function $\Omega_{\text{gw}}(f)$ for gravitational waves.

The gravitational wave energy density

$$\rho_{\text{gw}} = \frac{1}{32\pi G} \langle \dot{h}_{ij} \dot{h}_{ij} \rangle$$

is related to $N_{\text{gw}}(k)$ by [68]:

$$\rho_{\text{gw}} = \frac{\hbar}{\pi^2} \int_0^\infty d\omega \omega^3 N_{\text{gw}}(\omega).$$

On the other hand, the energy density spectral function $\rho_{\text{gw}}(\omega)$ of the gravitational waves is also given by

$$d\rho_{\text{gw}} = \rho_{\text{gw}}(\omega) d\omega.$$

Comparing the above two relations, we see that

$$\rho_{\text{gw}}(\omega) = \frac{\hbar}{\pi^2} \omega^3 N_{\text{gw}}(\omega).$$

Using the wave frequency $f = \omega/2\pi$, Planck constant $\hbar = 2\pi\hbar$, and $d\rho_{\text{gw}} = \rho_{\text{gw}}(f) df$, we can write the above as

$$\rho_{\text{gw}}(f) = 8\pi\hbar f^3 N_{\text{gw}}(\omega).$$

Then, by using the critical energy density ρ_c in cosmology, we see that the dimensionless spectral function $\Omega_{\text{gw}}(f)$ for gravitational waves is given by

$$\Omega_{\text{gw}}(f) = \frac{1}{\rho_c} \frac{d\rho_{\text{gw}}}{d \ln f} = \frac{8\pi\hbar}{\rho_c} f^4 N_{\text{gw}}(\omega). \quad (49)$$

To recover $N_{\text{gw}}(\omega)$ from given $\Omega_{\text{gw}}(f)$, we can invert Eq. (49) to get

$$N_{\text{gw}}(\omega) = \frac{\rho_c}{8\pi\hbar} \frac{\Omega_{\text{gw}}(f)}{f^4} \quad (50)$$

which is the sought relation between the distribution function $N_{\text{gw}}(\omega)$ and spectral function $\Omega_{\text{gw}}(f)$.

Let us now consider an estimate of the single atom transition rate Γ_0 with states $|1\rangle = |n_1, l_1, m_1\rangle$ and $|2\rangle = |n_2, l_2, m_2\rangle$ where $n_2 \geq n_1$

In terms of the Bohr radius a_0 , we can make the following approximations

$$\begin{aligned} (R_{n_1, l_1} |r^2| R_{n_1, l_1}) &\approx n_1^2 a_0^2 \\ (R_{n_2, l_2} |r^2| R_{n_2, l_2}) &\approx n_2^2 a_0^2 \end{aligned}$$

and hence [33]

$$(R_{n_1, l_1} |r^2| R_{n_2, l_2})^2 \approx n_1^2 n_2^2 a_0^4. \quad (51)$$

Substituting Eq. (51) into Eq. (30) and using the related 3- j symbols described in appendix B we can therefore approximate

$$q^2 \approx n_1^2 n_2^2 a_0^4 \quad (52)$$

subject to the selection rules (31). Then the gravitational spontaneous emission rate for a single atom given by Eq. (29), with the speed of light c restored, reads

$$\Gamma_0 = \frac{2 G m^2 n_1^2 n_2^2 a_0^4 \omega_0^5}{45 \hbar c^5} \quad (53)$$

also subject to the selection rules (31).

To take advantage of an N^2 amplification factor for the transition rate discussed in Sec. IV, we will take as initial state a spikelike profile specified by

$$\rho_p = \delta_{p, p_0} \quad (54)$$

with $1 \leq p_0 \leq N + 1$; then, Eq. (42) becomes

$$\begin{aligned} D_p &= (1 + N_{\text{gw}}(\omega_0))p(N - p + 1)\delta_{p+1,p_0} \\ &\quad - (1 + N_{\text{gw}}(\omega_0))(p - 1)(N - p + 2)\delta_{p,p_0} \\ &\quad + N_{\text{gw}}(\omega_0)(p - 1)(N - p + 2)\delta_{p-1,p_0} \\ &\quad - N_{\text{gw}}(\omega_0)p(N - p + 1)\delta_{p,p_0}. \end{aligned} \quad (55)$$

The transition rate of the initial state follows as

$$\begin{aligned} D_{p_0} &= -(1 + N_{\text{gw}}(\omega_0))(p_0 - 1)(N - p_0 + 2) \\ &\quad - N_{\text{gw}}(\omega_0)p_0(N - p_0 + 1) \end{aligned} \quad (56)$$

whereas the transition rate of the state above the initial state follows as

$$D_{p_0+1} = N_{\text{gw}}(\omega_0)p_0(N - p_0 + 1). \quad (57)$$

For a large N , to maximize the absolute values of Eqs. (56) and (57), we choose $p_0 \approx N/2$, and then we have

$$D_{p_0} = -\frac{N^2}{4}(1 + 2N_{\text{gw}}(\omega_0)) \quad (58)$$

and

$$D_{p_0+1} = \frac{N^2}{4}N_{\text{gw}}(\omega_0) \quad (59)$$

to the leading order in N . The short-time evolutions of such ‘‘favourable states’’ under gravitational waves and spacetime fluctuations are illustrated in Figs. 3 and 4.

Using Eqs. (41), (53), (58), and (59), it is convenient to introduce the effective scale transition rates

$$\Gamma_{\text{gw}} = \frac{N^2}{4}\Gamma_0 N_{\text{gw}}(\omega_0) \quad (60)$$

$$\Gamma_{\text{vac}} = \frac{N^2}{4}\Gamma_0 N_{\text{vac}}(\omega_0) \quad (61)$$

and the associated relaxation times

$$\tau_{\text{gw}} = \frac{1}{\Gamma_{\text{gw}}}, \quad \tau_{\text{vac}} = \frac{1}{\Gamma_{\text{vac}}} \quad (62)$$

due respectively to stochastic gravitational waves and vacuum fluctuations, in terms of the constant vacuum distribution function

$$N_{\text{vac}}(\omega) = 1. \quad (63)$$

Eqs. (49) and (63) allow us to give an analogous spectral function for vacuum fluctuations as a quartic function of frequency to be

$$\Omega_{\text{vac}}(f) = \frac{8\pi\hbar f^4}{c^3\rho_c} \quad (64)$$

with the speed of light c restored.




FIG. 4. Simulations of short-time collective transitions due to vacuum fluctuations of spacetime and additional stochastic gravitational waves with a chosen $N_{\text{gw}}(\omega_0) = 1$. In plot (a) with $N = 100$ atoms, the initial state corresponds to the spike of height 1 at $(p, t) = (50, 0)$. While this state decays similarly to the descriptions in Fig. 4, its adjacent higher state with $p = 51$ is excited with an initial time scale of $\tau_{\text{gw}} = 4/(N^2\Gamma_0 N_{\text{gw}}(\omega_0)) = 0.0004/\Gamma_0$ according to Eqs. (60) and (62). In plot (b) with $N = 200$ atoms, the initial state corresponds to the spike of height 1 at $(p, t) = (100, 0)$. For a doubled N , its adjacent higher state with $p = 101$ is excited with a quarter of the preceding initial time scale with $\tau_{\text{gw}} = 4/(N^2\Gamma_0 N_{\text{gw}}(\omega_0)) = 0.0001/\Gamma_0$.

If we envisage an off-resonant high- Q cavity of volume L^3 , then the number of contained atoms N is limited by

$$N \lesssim \frac{L^3}{4\pi(n_2^2 a_0)^3/3}. \quad (65)$$

Substituting Eqs. (53), (60), (62), and (65) into Eq. (49),

we have

$$\Omega(f_0) \gtrsim \frac{160 \hbar^2 c^2 a_0^2 n_2^{10}}{GL^6 \rho_c m^2 \tau \omega_0 n_1^2} \quad (66)$$

which applies to $\Omega = \Omega_{\text{gw}}$ and Ω_{vac} for given $\tau = \tau_{\text{gw}}$ and τ_{vac} for the stochastic gravitational waves and vacuum spacetime fluctuations respectively. Considering the transition frequency given by

$$\omega_0 = \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} + \epsilon \right) \frac{E_R}{\hbar} \quad (67)$$

where ϵ indicates an energy shift due to, e.g., quantum defects or Lamb/Zeeman shifts, it is clear that the lower bound of the spectral function Eq. (66) is minimized for $f_0 \lesssim \text{THz}$ with $\epsilon \ll 1$ by *small* principal quantum numbers $n_2 = n$ which is as close to 1 as possible, with n_1 equal to or as close to n as possible.

Subject to further experimental constraints to be investigated further, Eq. (66) provides theoretical lower bounds on their spectral functions for detection, as precluded in Sec. I with Fig. 1, where specific numerical examples using an ensemble of heliumlike atoms are studied with $n_1 = n_2 = n = 2$, $l_1 = l_2 = 1$, $-m_1 = m_2 = 1$, satisfying the selection rules (31) with $l_1 + l_2 = 2$, $\Delta l = 0$, $\Delta m = 2$, and $0 < \epsilon \lesssim 0.003$, which could be induced by a Zeeman-type shift.

VI. CONCLUSION AND DISCUSSION

Recent developments in gravitational and quantum physics have not only enriched applications with emerging prospects but also escalated the quest for a unified theory of quantum gravity that in turn has important bearing on the structure and evolution of the Universe. The growing availabilities for quantum matter exemplified by ultracold atoms with higher excitations, more correlations, better controls, and in larger quantities promise valuable tools in this scientific endeavour.

In this work, we have carried out new theoretical analysis describing how a large number of correlated atoms could interact and sense gravitational waves of fundamental and cosmological origins to go beyond the physical capacities of conventional detectors. We have reported theoretical results within generic physical constraints, but not fully taken onboard experimental implementations, which are a subject of future work.

Nonetheless, apart from extending the conceptual framework, building up theoretical tools and guiding possible viable detection of gravitational waves outside existing windows including zero-point fluctuations, our findings can already be used to rule out potential claims of effects that can be mapped to below our theoretical lower bounds on measurable gravitational wave spectral function values using quantum techniques, as noted in the

caption of Fig. 1, unless new inputs in addition to considerations adopted here are supplied.

In going forward, it would also be worth exploring the effects of the interaction Hamiltonian H_{int} in Eq. (1), which we neglect at present for simplicity, to accommodate atom-atom interactions, external control of atoms, and interactions with regular and nonstochastic gravitational waves for their quantum sensing based our framework.

ACKNOWLEDGMENTS

The authors are grateful for financial support to the National Council for Science and Technology (CONA-CyT) (D.Q.), the Carnegie Trust for the Universities of Scotland (T.O.), and the Cruickshank Trust and EPSRC GG-Top Project (C.W.).

Appendix A: Alternative derivation of the quantum dissipator of the two-level atom

The quantum mechanical interaction Hamiltonian between a mass m and gravitational waves [33, 39] is given by

$$H_I = \frac{1}{2} m R_{0i0j} r_i r_j. \quad (A1)$$

Using the weak gravitational wave relation [64]

$$R_{0i0j} = -\frac{1}{2} \ddot{h}_{ij}^{\text{TT}}$$

we see that Eq. (A1) becomes

$$H_I = -\frac{1}{12} m Q_{ij} \ddot{h}_{ij}^{\text{TT}} \quad (A2)$$

where Q_{ij} is the quadrupole moment tensor given by Eq. (7).

It is useful to write Eq. (A2) in the standard form of the interaction Hamiltonian for open quantum systems [66]

$$H_I = \sum_{\alpha} A_{\alpha} B_{\alpha} \quad (A3)$$

with the following identifications:

$$\alpha \rightarrow (i, j) \quad (A4)$$

$$A_{\alpha} \rightarrow A_{ij} = m Q_{ij} \quad (A5)$$

$$B_{\alpha} \rightarrow B_{ij} = -\frac{1}{12} \ddot{h}_{ij}^{\text{TT}} \quad (A6)$$

yielding in our two-level case the following Lindblad operators

$$A_{ij}(\omega) = |1\rangle\langle 1|A_{ij}|2\rangle\langle 2| = m q_{ij} \sigma^- \quad (\text{A7})$$

$$A_{ij}(-\omega) = |2\rangle\langle 2|A_{ij}|1\rangle\langle 1| = m q_{ij}^* \sigma^+ \quad (\text{A8})$$

using Eq. (25) and

$$\sigma^+ = |2\rangle\langle 1|, \quad \sigma^- = |1\rangle\langle 2|.$$

Clearly, $A_{ij}(\omega)$ satisfies

$$A_{ij}^\dagger(\omega) = A_{ij}(-\omega), \quad A_{kk}(\omega) = 0 \quad (\text{A9})$$

for $\omega \rightarrow \pm\omega$.

Using the rotating wave approximation, we have the following dissipator

$$\mathcal{D}\rho = \sum_{\omega \rightarrow \pm\omega} \sum_{\alpha, \alpha'} \Gamma_{\alpha, \alpha'}(\omega) \times \\ \left[A_\alpha(\omega) \rho A_{\alpha'}^\dagger(\omega) - A_{\alpha'}^\dagger(\omega) A_\alpha(\omega) \rho \right] + \text{H.c.} \quad (\text{A10})$$

with the spectral correlation tensor

$$\Gamma_{\alpha, \alpha'}(\omega) = \frac{1}{\hbar^2} \int_0^\infty ds e^{i\omega s} \langle B_\alpha^\dagger(s) B_{\alpha'}(0) \rangle \quad (\text{A11})$$

using a Gaussian stationary bath of gravitational waves. This expression can be evaluated by using Eq. (A6) with the field expansion for h_{ij}^{TT} . To evaluate Eq. (A11) consider the wave expansion for the gravitons

$$h_{ij}^{\text{TT}} = \int d^3k \sqrt{\frac{G\hbar}{\pi^2 k}} e_{ij}^\lambda(\mathbf{k}) a_{\mathbf{k}}^\lambda e^{ikx} + \text{H.c.} \quad (\text{A12})$$

with the relations

$$[a_{\mathbf{k}}^\lambda, a_{\mathbf{k}'}^{\lambda'\dagger}] = \delta_{\lambda\lambda'} \delta(\mathbf{k}, \mathbf{k}')$$

$$[a_{\mathbf{k}}^{\lambda\dagger}, a_{\mathbf{k}'}^{\lambda'\dagger}] = [a_{\mathbf{k}}^\lambda, a_{\mathbf{k}'}^{\lambda'}] = 0.$$

To proceed, consider the discrete momentum \mathbf{k} of the atom in a box of volume V expressed in terms of the Cartesian wave numbers $n_i = 1, 2, \dots$ as follows:

$$\mathbf{k} = \left(\frac{2\pi n_1}{V^{1/3}}, \frac{2\pi n_2}{V^{1/3}}, \frac{2\pi n_3}{V^{1/3}} \right)$$

In order to write the field in this box, we use

$$\int d^3k \leftrightarrow \frac{(2\pi)^3}{V} \sum_{\mathbf{k}} \quad (\text{A13})$$

with the modes inside the box normalized such that

$$\int_V d^3x e^{i\mathbf{k}\mathbf{r}} e^{-i\mathbf{k}'\mathbf{r}} = V \delta_{\mathbf{k}\mathbf{k}'}$$

Along with the following representation for the delta function in a finite volume,

$$\delta(\mathbf{r} - \mathbf{r}') = \frac{1}{V} \sum_{\mathbf{k}} e^{i\mathbf{k}(\mathbf{r}-\mathbf{r}')}$$

The Dirac delta function maps to the Kronecker delta as follows:

$$\delta(\mathbf{k} - \mathbf{k}') \leftrightarrow \frac{V}{(2\pi)^3} \delta_{\mathbf{k}\mathbf{k}'}$$

Ladder operators for free fields are related by

$$a_{\mathbf{k}} \leftrightarrow \sqrt{\frac{V}{(2\pi)^3}} a_{\mathbf{k}}$$

$$a_{\mathbf{k}}^\dagger \leftrightarrow \sqrt{\frac{V}{(2\pi)^3}} a_{\mathbf{k}}^\dagger$$

so that

$$[a_{\mathbf{k}}, a_{\mathbf{k}'}^\dagger] = \delta(\mathbf{k} - \mathbf{k}')$$

$$[a_{\mathbf{k}}, a_{\mathbf{k}}^\dagger] = \delta_{\mathbf{k}\mathbf{k}'}$$

Then, the discrete version of (A12) takes the form

$$h_{ij}^{\text{TT}} = \sum_{\mathbf{k}} \sqrt{\frac{8\pi G\hbar}{Vk}} e_{ij}^\lambda(\mathbf{k}) a_{\mathbf{k}}^\lambda e^{ikx} + \text{H.c.} \quad (\text{A14})$$

For an equilibrium environment [66], we have

$$\langle a_{\mathbf{k}}^{\lambda\dagger} a_{\mathbf{k}'}^{\lambda'} \rangle = \delta_{\lambda\lambda'} \delta_{\mathbf{k}, \mathbf{k}'} N_{\text{gw}}(k)$$

$$\langle a_{\mathbf{k}}^\lambda a_{\mathbf{k}'}^{\lambda'\dagger} \rangle = \delta_{\lambda\lambda'} \delta_{\mathbf{k}, \mathbf{k}'} (1 + N_{\text{gw}}(k))$$

$$\langle a_{\mathbf{k}}^{\lambda\dagger} a_{\mathbf{k}'}^{\lambda'\dagger} \rangle = \langle a_{\mathbf{k}}^\lambda a_{\mathbf{k}'}^{\lambda'} \rangle = 0$$

for some distribution function $N_{\text{gw}}(k)$ with the quantum ensemble average $\langle \cdot \rangle$. Then using Eqs. (A6), (A14), and the above, Eq. (A11) becomes

$$\Gamma_{ij,kl}(\omega) = \frac{\pi G k^3}{9\hbar V} \sum_{\mathbf{k}} P_{ijkl}(\mathbf{k}) \times \\ \int_0^\infty ds [(1 + N_{\text{gw}}(k)) e^{-i(k-\omega)s} + N_{\text{gw}}(k) e^{i(k+\omega)s}]$$

Then by using Eq. (A13), the above returns to the continuous momentum case as follows

$$\Gamma_{ij,kl}(\omega) = \frac{G}{72\pi^2 \hbar} \int_0^\infty dk k^5 \int d\Omega P_{ijkl}(\mathbf{k}) \times \\ \int_0^\infty ds [(1 + N_{\text{gw}}(k)) e^{-i(k-\omega)s} + N_{\text{gw}}(k) e^{i(k+\omega)s}].$$

Substituting the relation (13) into the above, we have

$$\Gamma_{ij,kl}(\omega) = \frac{G(3\delta_{ik}\delta_{jl} + 3\delta_{il}\delta_{jk} - 2\delta_{ij}\delta_{kl})}{270\pi\hbar} \int_0^\infty dk k^5 \times \\ \int_0^\infty ds [(1 + N_{\text{gw}}(k)) e^{i(\omega-k)s} + N_{\text{gw}}(k) e^{i(\omega+k)s}]$$

Using the the Sokhotski-Plemelj theorem [66]

$$\int_0^\infty ds e^{-i\epsilon s} = \pi\delta(\epsilon) - iP\frac{1}{\epsilon} \quad (\text{A15})$$

where P denotes the Cauchy principal value, the above becomes

$$\Gamma_{ij,kl}(\omega) = \frac{G(3\delta_{ik}\delta_{jl} + 3\delta_{il}\delta_{jk} - 2\delta_{ij}\delta_{kl})}{270\pi\hbar} \times \int_0^\infty dk k^5 \left[(1 + N_{\text{gw}}(k)) \left(\pi\delta(\omega - k) + iP \frac{1}{\omega - k} \right) + N_{\text{gw}}(k) \left(\pi\delta(\omega + k) + iP \frac{1}{\omega + k} \right) \right].$$

Neglecting P terms as they do not contribute to quantum decoherence and dissipation, we see that the above reduces to

$$\Gamma_{ij,kl}(\omega) = \frac{G(3\delta_{ik}\delta_{jl} + 3\delta_{il}\delta_{jk} - 2\delta_{ij}\delta_{kl})}{270\hbar} \int_0^\infty dk \times k^5 \left[(1 + N_{\text{gw}}(k))\delta(\omega - k) + N_{\text{gw}}(k)\delta(\omega + k) \right]$$

yielding

$$\Gamma_{ij,kl}(\omega) = \frac{G\omega^5}{270\hbar} \times (3\delta_{ik}\delta_{jl} + 3\delta_{il}\delta_{jk} - 2\delta_{ij}\delta_{kl})(N_{\text{gw}}(\omega) + 1)$$

$$\Gamma_{ij,kl}(-\omega) = \frac{G\omega^5}{270\hbar} (3\delta_{ik}\delta_{jl} + 3\delta_{il}\delta_{jk} - 2\delta_{ij}\delta_{kl})N_{\text{gw}}(\omega).$$

Substituting these two relations into the dissipator (A10) and using (A9), we have

$$\begin{aligned} \mathcal{D}\rho = & \frac{G\omega^5}{90\hbar} \sum_{ij,kl} (\delta_{ik}\delta_{jl} + \delta_{il}\delta_{jk}) \times \\ & \left\{ (N_{\text{gw}}(\omega) + 1) [A_{ij}(\omega)\rho A_{kl}^\dagger(\omega) - A_{kl}^\dagger(\omega)A_{ij}(\omega)\rho \right. \\ & + A_{kl}(\omega)\rho A_{ij}^\dagger(\omega) - \rho A_{ij}^\dagger(\omega)A_{kl}(\omega)] \\ & + N_{\text{gw}}(\omega) [A_{ij}^\dagger(\omega)\rho A_{kl}(\omega) - A_{kl}(\omega)A_{ij}^\dagger(\omega)\rho \\ & \left. + A_{kl}^\dagger(\omega)\rho A_{ij}(\omega) - \rho A_{ij}(\omega)A_{kl}^\dagger(\omega)] \right\} \end{aligned}$$

which yields the following dissipator of the Lindblad form:

$$\begin{aligned} \mathcal{D}\rho = & \frac{2G\omega^5}{45\hbar} \sum_{ij} \left\{ (N_{\text{gw}}(\omega) + 1) \times \right. \\ & \left[A_{ij}(\omega)\rho A_{ij}^\dagger(\omega) - \frac{1}{2} \{ A_{ij}^\dagger(\omega)A_{ij}(\omega), \rho \} \right] \\ & \left. + N_{\text{gw}}(\omega) [A_{ij}^\dagger(\omega)\rho A_{ij}(\omega) - \frac{1}{2} \{ A_{ij}(\omega)A_{ij}^\dagger(\omega), \rho \}] \right\}. \end{aligned}$$

Finally, by using Eqs. (A7) and (A8), we can further simplify the above to the same form as Eq. (28).

Appendix B: Evaluation of the transition quadrupole moment

To evaluate q_{ij} in Eq. (25), let us consider the components of Q_{ij} given by Eq. (7), noting that Q_{ij} has five independent components for being traceless. For $l = 2$, we can express Q_{ij} in terms of the spherical harmonics $Y_{l=2}^m(\theta, \phi)$ as follows:

$$Q_{11} = \sqrt{\frac{6\pi}{5}} r^2 (Y_2^{-2} + Y_2^2) - \sqrt{\frac{4\pi}{5}} r^2 Y_2^0 \quad (\text{B1})$$

$$Q_{22} = -\sqrt{\frac{6\pi}{5}} r^2 (Y_2^{-2} + Y_2^2) - \sqrt{\frac{4\pi}{5}} r^2 Y_2^0 \quad (\text{B2})$$

$$Q_{33} = \sqrt{\frac{16\pi}{5}} r^2 Y_2^0 \quad (\text{B3})$$

$$Q_{12} = i\sqrt{\frac{6\pi}{5}} r^2 (Y_2^{-2} - Y_2^2) \quad (\text{B4})$$

$$Q_{13} = \sqrt{\frac{6\pi}{5}} r^2 (Y_2^{-1} - Y_2^1) \quad (\text{B5})$$

$$Q_{23} = i\sqrt{\frac{6\pi}{5}} r^2 (Y_2^{-1} + Y_2^1). \quad (\text{B6})$$

Denoting by

$$(R_{n_1, l_1} | f(r) | R_{n_2, l_2}) = \int_0^\infty dr r^2 f(r) R_{n_1, l_1}(r) R_{n_2, l_2}(r)$$

we can evaluate the matrix elements of the above Q_{ij} in terms of Wigner's 3- j symbols through the relation

$$\begin{aligned} & (n_1, l_1, m_1 | r^2 Y_{l=2}^m(\theta, \phi) | n_2, l_2, m_2) \\ & = (R_{n_1, l_1} | r^2 | R_{n_2, l_2}) (-1)^{m_1} \sqrt{\frac{(2l_1 + 1)(2l + 1)(2l_2 + 1)}{4\pi}} \\ & \quad \times \begin{pmatrix} l_1 & l & l_2 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l_1 & l & l_2 \\ -m_1 & m & m_2 \end{pmatrix} \quad (\text{B7}) \end{aligned}$$

with $l = 2$.

-
- [1] G. Amelino-Camelia, Quantum-Spacetime Phenomenology, *Living Rev. Relativity* **16**, 5 (2013).
- [2] C. Pfister *et al.*, A universal test for gravitational decoherence, *Nat. Commun.* **7** 13022 (2016).
- [3] C. H.-T. Wang, New “phase” of quantum gravity, *Phil. Trans. R. Soc. A* **364**, 3375 (2006).
- [4] C. Kiefer, *Quantum Gravity* (Oxford University Press, Oxford, England, 2013).
- [5] T. Oniga and C. H.-T. Wang, Quantum gravitational decoherence of light and matter, *Phys. Rev. D* **93**, 044027 (2016).
- [6] H. Müller, S. W. Chiow, S. Herrmann, S. Chu, and K.-Y. Chung, Atom-Interferometry Tests of the Isotropy of Post-Newtonian Gravity, *Phys. Rev. Lett.* **100**, 031101 (2008).
- [7] O. Carraz *et al.*, A spaceborne gravity gradiometer concept based on cold atom interferometers for measuring Earth’s gravity field, *Microgravity Sci. Technol.* **26**, 139 (2014).
- [8] G. Amelino-Camelia *et al.*, GAUGE: The GrAnd Unification Gravity Explorer, *Experimental Astron.* **23**, 549 (2009).
- [9] H. Müntinga *et al.* Interferometry with Bose-Einstein Condensates in Microgravity, *Phys. Rev. Lett.* **110**, 093602 (2013).
- [10] I. Pikovski, M. Zych, F. Costa, and C. Brukner, Universal decoherence due to gravitational time dilation, *Nat. Phys.* **11**, 668 (2015).
- [11] G. Amelino-Camelia, Gravity-wave interferometers as probes of a low-energy effective quantum gravity, *Phys. Rev. D* **62**, 024015 (2000).
- [12] C. H.-T. Wang, R. Bingham, and J. T. Mendonca, Quantum gravitational decoherence of matter waves, *Classical Quantum Gravity* **23**, L59 (2006).
- [13] B. Lamine, R. Herve, A. Lambrecht, and S. Reynaud, Ultimate Decoherence Border for Matter-Wave Interferometry, *Phys. Rev. Lett.* **96**, 050405 (2006).
- [14] M. S. Everitt, M. L. Jones, and B. T. Varcoe, Dephasing of entangled atoms as an improved test of quantized space time, *J. Phys. B* **46**, 224003, (2013).
- [15] A. Peres, Separability Criterion for Density Matrices, *Phys. Rev. Lett.* **77**, 1413 (1996).
- [16] J. Handsteiner *et al.*, Cosmic Bell Test: Measurement Settings from Milky Way Stars, *Phys. Rev. Lett.* **118**, 060401 (2017).
- [17] T. Oniga and C. H.-T. Wang, Spacetime foam induced collective bundling of intense fields, *Phys. Rev. D* **94**, 061501(R) (2016).
- [18] T. Oniga and C. H.-T. Wang, Quantum dynamics of bound states under spacetime fluctuations, *J. Phys. Conf. Ser.* **845**, 012020 (2017).
- [19] R. Hanbury Brown and R. Q. Twiss, Correlation between photons in two coherent beams of light, *Nature (London)* **177**, 27 (1956).
- [20] R. H. Dicke, Coherence in spontaneous radiation processes, *Phys. Rev.* **93**, 99 (1954).
- [21] LIGO Scientific and Virgo Collaborations, An upper limit on the stochastic gravitational-wave background of cosmological origin, *Nature (London)* **460**, 990 (2009) and references therein.
- [22] B. P. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), GW150914: Implications for the Stochastic Gravitational-Wave Background from Binary Black Hole, *Phys. Rev. Lett.* **116**, 131102 (2016).
- [23] P. D. Lasky *et al.*, Gravitational-wave cosmology across 29 decades in frequency, *Phys. Rev. X* **6**, 011035 (2016).
- [24] D.-G. Wang, Y. Zhang, and J.-W. Chen, Vacuum and gravitons of relic gravitational waves and the regularization of the spectrum and energy-momentum tensor, *Phys. Rev. D* **94**, 044033 (2016).
- [25] C. Kiefer, D. Polarski, and A. A. Starobinsky, Entropy of gravitons produced in the early universe, *Phys. Rev. D* **62**, 043518 (2000).
- [26] R. Easther and E. A. Lim, Stochastic gravitational wave production after inflation, *J. Cosmol. Astropart. Phys.* **2006**, 10 (2006).
- [27] T. Oniga and C. H.-T. Wang, arXiv:1701.04122.
- [28] B. P. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), Observation of Gravitational Waves from a Binary Black Hole Merger, *Phys. Rev. Lett.* **116**, 061102 (2016).
- [29] B. F. Schutz, Gravitational wave astronomy, *Classical Quantum Gravity* **16**, A131 (1999).
- [30] A. Sesana, Prospects for Multiband Gravitational-Wave Astronomy after GW150914, *Phys. Rev. Lett.* **116**, 231102 (2016).
- [31] B. Allen, *Astrophysical Sources of Gravitational Waves*, Proceedings of the Les Houches Summer School, edited by J.-A. Marck and J.-P. Lasota (Cambridge University Press, Cambridge, England, 1996); arXiv:gr-qc/9604033.
- [32] E. Fischbach, B. S. Freeman, and W.-K. Cheng General-relativistic effects in hydrogenic systems, *Phys. Rev. D* **23**, 2157 (1981).
- [33] U. Fischer, Transition probabilities for a Rydberg atom in the field of a gravitational wave, *Classical Quantum Gravity* **11**, 463 (1994).
- [34] G. A. Marques and V. B. Bezerra, Hydrogen atom in the gravitational fields of topological defects, *Phys. Rev. D* **66**, 105011 (2002).
- [35] G. J. Olmo, Hydrogen atom in Palatini theories of gravity, *Phys. Rev. D* **77**, 084021 (2008).
- [36] Z.-H. Zhao, Y.-X. Liu, and X.-G. Li, Gravitational corrections to energy-levels of a hydrogen atom, *Commun. in Theor. Phys.* **47**, 4 (2007).
- [37] M. P. Blencowe, Effective Field Theory Approach to Gravitationally Induced Decoherence, *Phys. Rev. Lett.* **111**, 021302 (2013).
- [38] D. A. Quiñones and B. Varcoe, Decoherence in excited atoms by low-energy scattering, *Atoms* **2016**, 28 (2016).
- [39] L. Parker, One-electron atom as a probe of spacetime curvature, *Phys. Rev. D* **22**, 1922 (1980).
- [40] L. Parker and L. O. Pimentel, Gravitational perturbation of the hydrogen spectrum, *Phys. Rev. D* **25**, 3180, (1982).
- [41] T. K. Leen, L. Parker, and L. O. Pimentel, Remote quantum mechanical detection of gravitational radiation, *General Relativ. Gravit.* **15**, 761 (1983).
- [42] F. Pinto, Rydberg atoms as gravitational-wave antennas, *General Relativ. Gravit.* **27**, 3839 (1995).

- [43] T. F. Gallagher, *Rydberg Atoms* (Cambridge University Press, Cambridge, England 2008).
- [44] M. Saffman, T. G. Walker, and K. Molmer, Quantum information with Rydberg atoms, *Rev. Mod. Phys.* **82**, 2313 (2010).
- [45] A. Kortyna *et al.*, Radiative lifetime measurements of rubidium Rydberg states, *J. Phys. B* **43**, 015002, (2010).
- [46] S. Ye, X. Zhang, T. C. Killian, F. B. Dunning, M. Hiller, S. Yoshida, S. Nagele, and J. Burgdörfer, Production of very-high- n strontium Rydberg atoms, *Phys. Rev. A* **88**, 043430 (2013).
- [47] A. Browaeys and T. Lahaye, Interacting Cold Rydberg Atoms: a Toy Many-Body System, *Séminaire Poincaré* **17**, 125 (2013) and references therein.
- [48] D. Jaksch *et al.*, Fast Quantum Gates for Neutral Atoms, *Phys. Rev. Lett.* **85**, 2208 (2000).
- [49] M. D. Lukin *et al.*, Dipole Blockade and Quantum Information in Mesoscopic Atomic Ensembles, *Phys. Rev. Lett.* **87**, 037901 (2001).
- [50] B. Olmos, R. González-Férez, and I. Lesanovsky, Creating collective many-body states with highly excited atoms, *Phys. Rev. A* **81**, 023604 (2010).
- [51] T. E. Lee, H. Häffner, and M. C. Cross, Collective Quantum Jumps of Rydberg Atoms, *Phys. Rev. Lett.* **108**, 023602 (2012).
- [52] A. Gaetan *et al.*, Observation of collective excitation of two individual atoms in the Rydberg blockade regime, *Nat. Phys.* **5**, 115 (2009).
- [53] T. Wilk *et al.*, Entanglement of Two Individual Neutral Atoms Using Rydberg Blockade, *Phys. Rev. Lett.* **104**, 010502 (2010).
- [54] R. Heidemann *et al.*, Evidence for Collective Rydberg Excitation in the Strong Blockade Regime, *Phys. Rev. Lett.* **99**, 163601 (2007).
- [55] Y.O. Dudin, L. Li, F. Bariani, and A. Kuzmich, Observation of coherent many-body Rabi oscillations, *Nat. Phys.* **8**, 790 (2012).
- [56] I. Mourachko *et al.*, Many-body effects in a frozen Rydberg gas, *Phys. Rev. Lett.* **80**, 253 (1998).
- [57] P. Schauss *et al.*, Observation of spatially ordered structures in a two-dimensional Rydberg gas, *Nature (London)* **491**, 87 (2012).
- [58] S. Kumar *et al.*, Collective state synthesis in an optical cavity using Rydberg atom dipole blockade, *J. Phys. B* **49**, 064014 (2016).
- [59] F. Pinto, Rydberg Atoms in Curved Space-Time, *Phys. Rev. Lett.* **70**, 25, (1993).
- [60] D. Kleppner, Inhibited Spontaneous Emission, *Phys. Rev. Lett.* **47**, 233 (1981)
- [61] R. G. Hulet, E. S. Hilfer, and D. Kleppner, Inhibited Spontaneous Emission by a Rydberg Atom, *Phys. Rev. Lett.* **55**, 2137 (1985)
- [62] H. Walther, B. T. H. Varcoe, B.-G. Englert, and T. Becker, Cavity quantum electrodynamics, *Rep. Prog. Phys.* **69**, 1325 (2006).
- [63] M. Marrocco, M. Weidinger, R. T. Sang, and H. Walther, Quantum Electrodynamical Shifts of Rydberg Energy Levels between Parallel Metal Plates, *Phys. Rev. Lett.* **81**, 5784 (1998).
- [64] E. E. Flanagan and S. A. Hughes, The basics of gravitational wave theory, *New J. Phys.* **7**, 204 (2005).
- [65] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation* (Freeman, New York, 1973).
- [66] H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, New York, 2002).
- [67] L. C. Biedenhorn, *Encyclopaedia of Mathematics and its Application* (Addison-Wesley, Reading, MA, 1981).
- [68] G. Schafer, Gravitational radiation resistance, radiation damping and field fluctuations, *J. Phys. A* **14**, 677 (1981).