

This is a repository copy of *Additive energy and the metric Poissonian property*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/127930/>

Version: Accepted Version

Article:

Bloom, Thomas F., Chow, Samuel Khai Ho, Gafni, Ayla et al. (1 more author) (2018) Additive energy and the metric Poissonian property. *Mathematika*. pp. 679-700. ISSN 2041-7942

<https://doi.org/10.1112/S0025579318000207>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

ADDITIVE ENERGY AND THE METRIC POISSONIAN PROPERTY

THOMAS F. BLOOM, SAM CHOW, AYLA GAFNI, AND ALED WALKER

ABSTRACT. Let A be a set of natural numbers. Recent work has suggested a strong link between the additive energy of A (the number of solutions to $a_1 + a_2 = a_3 + a_4$ with $a_i \in A$) and the metric Poissonian property, which is a fine-scale equidistribution property for dilates of A modulo 1. There appears to be reasonable evidence to speculate a sharp Khintchine-type threshold, that is, to speculate that the metric Poissonian property should be completely determined by whether or not a certain sum of additive energies is convergent or divergent. In this article, we primarily address the convergence theory, in other words the extent to which having a low additive energy forces a set to be metric Poissonian.

1. INTRODUCTION

The metric Poissonian property is a refined notion of the equidistribution of certain sequences on the unit circle. Initially studied in [11] for its connections to the Berry–Tabor conjecture in quantum mechanics, the property has recently received renewed interest, owing to the fundamental work of Aistleitner–Larcher–Lewko [2] who, for the first time, revealed an intimate quantitative connection between this property and the combinatorial notion of *additive energy*. We continue this quantitative investigation in one of our main theorems (Theorem 1.4).

In order to state the main result of [2], let us formally define the relevant notions. Let $\mathcal{A} \subseteq \mathbb{N}$ be an infinite set. For $X \rightarrow \infty$ a parameter, put

$$A = \mathcal{A} \cap [1, X], \quad N = \#A, \quad \delta = \delta(X) = N/X.$$

For $\alpha \in [0, 1)$, and $s > 0$ some fixed parameter, we define the pair correlation function

$$F(\alpha, s, X, \mathcal{A}) := N^{-1} \sum_{\substack{a, b \in A, a \neq b \\ \|\alpha(a-b)\| \leq s/N}} 1.$$

Here and in the rest of the paper we will use $\|x\| := \min_{n \in \mathbb{Z}} |x - n|$. The parameters s , X , and the underlying set \mathcal{A} will often be suppressed, with $F(\alpha)$ used to denote the above.

Considering αA as a subset of \mathbb{R}/\mathbb{Z} , the average gap length between consecutive elements is $1/N$, at least when $\alpha \notin \mathbb{Q}$. One can view the function $F(\alpha)$

2010 *Mathematics Subject Classification*. 11J71, 11J83, 05B10, 11B30, 60F10.

Key words and phrases. Pair correlations, distribution modulo 1, metric diophantine approximation, additive combinatorics, large deviations.

therefore as measuring the proportion of pairs $(\alpha a, \alpha b)$ of distinct elements such that the difference $\alpha a - \alpha b \pmod 1$ is on the scale of this average gap length. This scale is determined by the parameter s .

The set \mathcal{A} is said to be *metric Poissonian* if for almost all α we have

$$F(\alpha) \rightarrow 2s \quad (N \rightarrow \infty)$$

for all $s > 0$.

If \mathcal{A} is metric Poissonian, then for almost all α the set $\alpha\mathcal{A} \pmod 1$ mimics a certain statistic which holds almost surely for sequences (x_1, x_2, \dots) drawn uniformly and independently at random from $[0, 1]$. Indeed, if $I \subseteq \mathbb{R}/\mathbb{Z}$ is a uniformly chosen random interval of length $2s/N$, let Y_I denote the random variable $|I \cap \{x_1, \dots, x_N\}|$. For fixed k and large N one has

$$\begin{aligned} \mathbb{P}(Y_I = k) &= \frac{N!}{k!(N-k)!} (2s/N)^k (1 - 2s/N)^{N-k} \\ &= (1 + o_{k,s}(1)) \cdot \exp(-2s) \cdot (2s)^k / k!, \end{aligned}$$

so the limiting distribution of Y_I is Poisson with parameter $2s$. In particular its variance tends to $2s$ as N tends to infinity.

When the points x_1, x_2, \dots, x_N are deterministic, calculating the variance of Y_I is more or less equivalent to calculating the pair correlations

$$N^{-1} \sum_{\substack{x_i, x_j \\ x_i \neq x_j \\ \|x_i - x_j\| \leq t/N}} 1.$$

If for **all** $t > 0$ this tends to $2t$ as $N \rightarrow \infty$, then one may conclude that $\text{Var}(Y_I) \rightarrow 2s$. The metric Poissonian property, therefore, is the statement that for almost all α the set $\alpha\mathcal{A} \pmod 1$ matches the second order statistics of a random sequence in \mathbb{R}/\mathbb{Z} , regarding distribution in short intervals of length $O(1/N)$. The statistics of a random sequence are Poisson, in the large N limit, thus giving some explanation for the name ‘metric Poissonian’. For more details about this interpretation and connection, we direct the reader to the second section of [10].

When \mathcal{A} is the set of squares, or more generally the set of perfect k^{th} powers for $k \geq 2$, the set \mathcal{A} is metric Poissonian, as was shown by Rudnick and Sarnak in [11]. General lacunary sequences are also metric Poissonian (a result of [12]). Considerations of continued fractions show that $\mathcal{A} = \mathbb{N}$ is *not* metric Poissonian, however. As described at the start of the paper, the situation was greatly clarified in [2], by the consideration of *additive energy*

$$E = E(X) = \#\{(a, b, c, d) \in A^4 : a + b = c + d\}.$$

Indeed, the authors of [2] proved the following theorem.

Theorem 1.1. *Assume that $E \ll_{\xi} N^{3-\xi}$ for some $\xi > 0$. Then \mathcal{A} is metric Poissonian.*

We leave it to the reader to verify that this theorem implies the aforementioned results for perfect powers and lacunary sequences. We remark here that the trivial upper bound on E is N^3 , so the required hypothesis here is in some sense rather weak.

This theorem is neatly complemented by the following result of Bourgain, given in the appendix of [2].

Theorem 1.2. *Suppose*

$$\limsup_{N \rightarrow \infty} \frac{E}{N^3} > 0.$$

Then \mathcal{A} is not metric Poissonian.

These two theorems suggest a natural route for further investigation, namely to establish whether there is some threshold for the additive energy E which completely determines whether or not \mathcal{A} is metric Poissonian. Bourgain, in the same appendix of [2], gave an example of a set \mathcal{A} with $E = o(N^3)$ which was not metric Poissonian. This demonstrates that, if such a threshold exists, it is not at order N^3 .

The next development was made by the fourth named author in [14], answering a question of Nair concerning the primes.

Theorem 1.3. *The set of prime numbers is not metric Poissonian.*

In this instance, one has $E \sim N^3(\log N)^{-1}$. The proof of this theorem made use of the divergent part of Khintchine's theorem on Diophantine approximation, and rested on the divergence of the sum $\sum (N \log N)^{-1}$. At this point, we speculate that if $E \sim N^3\psi(N)$, for some decreasing function ψ , then \mathcal{A} is metric Poissonian if and only if $\sum \psi(N)/N$ converges.

Lachmann and Technau [8] have recently given examples of sets which are not metric Poissonian, and which come within an arbitrary finitely iterated logarithm of this putative threshold. For $r \in \mathbb{N}$, they showed that there exists a set \mathcal{A} which is not metric Poissonian, and which satisfies

$$E \asymp \frac{N^3}{\log(N) \log_2(N) \cdots \log_r(N)}.$$

Here $\log_0(N) = N$ and $\log_t(N) = \log(\log_{t-1}(N))$ for $t \in \mathbb{N}$. Here and throughout, we assume that N is large enough so that any iterated logarithms appearing are well-defined and positive.

We now come to the matter of the present paper. Firstly, we offer a large improvement over the results of [2], albeit under an extra hypothesis. Indeed, a key feature of Lachmann and Technau's work is that their constructed set is extremely sparse. Assuming instead that \mathcal{A} is quite dense, we show that if $E \ll_{\xi} N^3(\log N)^{-2-\xi}$ then \mathcal{A} is metric Poissonian. This can be considered a complement to Lachmann and Technau's construction.

Theorem 1.4. *Assume that*

$$E \ll_{\xi} \frac{N^3}{(\log N)^{2+\xi}}, \quad \delta \gg_{\xi} \frac{1}{(\log N)^{2+2\xi}}, \quad (1.1)$$

for some constant $\xi > 0$. Then \mathcal{A} is metric Poissonian.

If \mathcal{A} were ‘random’, e.g. each element of $\{1, 2, \dots, X\}$ chosen independently at random with probability δ , we would expect the energy to be approximately δN^3 . In the above theorem we permit δ to be smaller, by a power of a logarithm. Thus the theorem holds for all sets of energy $E \ll_{\xi} N^3(\log N)^{-2-\xi}$ bar those which are unexpectedly sparse.

For genuinely random sets \mathcal{A} , certain technical obstructions in the proof may be removed, and we obtain the following quantitatively stronger result.

Theorem 1.5. *Let $C > 2$. Let \mathcal{A} be a random set of natural numbers defined by choosing $x \in \mathcal{A}$ independently at random with probability*

$$\begin{cases} 1, & \text{if } x \leq 20, \\ (\log x)^{-1}(\log \log x)^{-C}, & \text{if } x > 20. \end{cases}$$

Then, with probability 1, the set \mathcal{A} is metric Poissonian.

In this theorem we have conceded a factor of roughly $\log \log N$ from the ‘Khintchine threshold’. Indeed, we verify in Appendix A that

$$E \asymp \frac{N^3}{(\log N) \cdot (\log \log N)^C}$$

with probability 1.

In the random setting we can also tackle the divergence side of the problem.

Theorem 1.6. *Let $0 \leq C \leq 1$. Let \mathcal{A} be a random set of natural numbers defined by choosing $x \in \mathcal{A}$ independently at random with probability*

$$\begin{cases} 1, & \text{if } x \leq 20, \\ (\log x)^{-1}(\log \log x)^{-C}, & \text{if } x > 20. \end{cases}$$

*Then, with probability 1, the set \mathcal{A} is **not** metric Poissonian.*

This complements Theorem 1.5. In a way, Theorem 1.6 demonstrates that the examples provided in [8] behave as one should expect.

We return to our fundamental motivating question: is there a universal ‘additive energy threshold’ for the metric Poissonian property?

Fundamental Question 1.7. *Is it true that if $E(N) \sim N^3\psi(N)$, for some weakly decreasing function $\psi : \mathbb{N} \rightarrow [0, 1]$, then \mathcal{A} is metric Poissonian if and only if $\sum \psi(N)/N$ converges?*

Theorem 1.5 shows that there are metric Poissonian sets with energy

$$\Theta\left(\frac{N^3}{(\log N) \cdot (\log \log N)^{2+\varepsilon}}\right),$$

whilst Lachmann and Technau [8, Corollary 1] construct sets of energy

$$\Theta\left(\frac{N^3}{(\log N) \cdot (\log \log N)}\right)$$

that are not metric Poissonian. In a way, we are within a doubly logarithmic factor of answering the question. Moreover, there is now strong evidence that the Khintchine threshold is the correct one, if such a threshold exists.

The paper is organised as follows. In section 2 we describe our method, introducing an idea of Schmidt which lies at the heart of the approach. The subsequent two sections are devoted to proving the key estimates. In section 5 we demonstrate how these estimates may be improved in the setting of Theorem 1.5. In section 6, we use a sandwiching idea we learnt from [2] to finish the proofs of Theorems 1.4 and 1.5. Then, in section 7, we use a simplified version of Walker’s method [14] to provide an abridged proof of Theorem 1.6. Finally, in Appendix A, we compute the additive energy in the random setting.

Regarding notation, throughout μ denotes Lebesgue measure. For $n \in \mathbb{N}$ write $[n] = \{1, 2, \dots, n\}$. We use Vinogradov and Bachmann–Landau notation, and we put $\mathbb{T} = [0, 1]$. It is convenient to introduce the notation \tilde{E} for the normalised additive energy E/N^3 , so that $\tilde{E} \in (0, 1]$.

Acknowledgements

This material is based upon work partially supported by the National Science Foundation under Grant No. DMS-1440140 while three of the authors were in residence at the Mathematical Sciences Research Institute in Berkeley, California, during the Spring 2017 semester. The second and third authors were MSRI Postdocs, and the fourth author a Program Associate, during the Analytic Number Theory Program. The first named author was visiting the Simons Institute for the Theory of Computing, Berkeley, for the Pseudorandomness Program 2017. All four authors are grateful for the working conditions provided by both institutions, at which this work was initiated. The second named author was also supported by EPSRC Programme Grant EP/J018260/1, and the fourth named author by EPSRC grant no. EP/M50659X/1. We thank Ben J. Green and Niclas Technau for helpful conversations. Finally, we thank an anonymous referee for a helpful and detailed report.

2. SCHMIDT’S TRICK

The arguments of Aistleitner, Larcher, and Lewko in [2] rest upon the bounds of Bondarenko and Seip on GCD-sums [4], which reach their natural limit once \tilde{E} is larger than $\exp((\log N)^{-1/2+\xi})$. To move above this threshold, we utilise a device of Schmidt from [13], which we learnt from Chapter 4 of Harman’s book [7]. The idea is that, when counting integer pairs (n, m) such that $|\alpha n - m|$ is small, one can profitably split into two cases, depending on the size of greatest common divisor of m and n . The resulting sum enjoys much better L^2 information than was available in [2], at a small L^1 cost.

We note at the outset that we will only consider values of $s > 0$ for which $s \asymp 1$. Let

$$r(n) = \#\{(a, b) \in A^2 : a - b = n, a \neq b\}$$

and

$$\mathcal{E}_n = \bigcup_{\substack{u \leq n \\ (u, n) \leq T}} \left(\frac{u - s/N}{n}, \frac{u + s/N}{n} \right) \bmod 1,$$

where $T \geq 2$ is a threshold to be specified in due course. Observe that

$$F(\alpha) = \frac{2}{N} \sum_{\substack{n \leq X \\ \|n\alpha\| \leq s/N}} r(n).$$

The idea is to replace $F(\alpha)$ by

$$F^*(\alpha) = F^*(\alpha, s, X, \mathcal{A}) = \frac{2}{N} \sum_{\substack{n \leq X \\ \alpha \in \mathcal{E}_n}} r(n).$$

We follow the exposition of Harman very closely, though the inclusion of the weight $r(n)$ means we need to redo much of the argument from Chapter 4 of [7]. Another difference is that the threshold T , as well as the parameter $2s/N$ of the intervals, is uniform, depending on N rather than n . This will actually simplify the proof of several lemmas.

We first establish two useful lemmas on the size of the sets \mathcal{E}_n . To this end, we introduce the quantity

$$\Phi(n) = \#\{u \in [n] : (u, n) \leq T\}$$

and observe that

$$\mu(\mathcal{E}_n) = \frac{2s}{N} \cdot \frac{\Phi(n)}{n}. \tag{2.1}$$

The second moment of $\Phi(n)$ will be particularly useful here, but in section 5 we will also have use for the first moment, when considering the random case Theorem 1.5.

Lemma 2.1. *We have*

$$\sum_{n \leq X} \left(1 - \frac{\Phi(n)}{n} \right) \ll \frac{X}{T}.$$

Proof. We begin by noting that

$$0 \leq n - \Phi(n) = \sum_{\substack{u \leq n: \\ (u, n) > T}} 1 = \sum_{\substack{d|n: \\ d > T}} \phi\left(\frac{n}{d}\right) \leq \sum_{\substack{d|n: \\ d > T}} \frac{n}{d},$$

where $m \mapsto \phi(m)$ is the Euler totient function. It follows that

$$\begin{aligned} 0 \leq \sum_{n \leq X} \left(1 - \frac{\Phi(n)}{n}\right) &\leq \sum_{n \leq X} \sum_{\substack{d|n: \\ d > T}} \frac{1}{d} = \sum_{T < d \leq X} \frac{1}{d} \sum_{\substack{n \leq X: \\ d|n}} 1 \\ &\leq X \sum_{T < d \leq X} d^{-2} \ll \frac{X}{T}. \end{aligned}$$

□

Lemma 2.2. *We have*

$$\sum_{n \leq X} \left(1 - \frac{\Phi(n)}{n}\right)^2 \ll \frac{X \log T}{T^2}.$$

Proof. Using the same initial manoeuvre as above, we have

$$\begin{aligned} \sum_{n \leq X} \left(1 - \frac{\Phi(n)}{n}\right)^2 &\leq \sum_{n \leq X} \sum_{\substack{d,e|n \\ d,e > T}} \frac{1}{de} = \sum_{T < d,e \leq X} \frac{1}{de} \sum_{\substack{n \leq X: \\ [d,e]|n}} 1 \\ &\leq X \sum_{T < d,e \leq X} \frac{(d,e)}{d^2 e^2}. \end{aligned}$$

Putting

$$g = (d, e), \quad d = gx, \quad e = gy$$

gives

$$\begin{aligned} \sum_{n \leq X} \left(1 - \frac{\Phi(n)}{n}\right)^2 &\leq X \sum_{g \leq X} g^{-3} \sum_{x,y \geq T/g} (xy)^{-2} \\ &\ll X \sum_{g \leq T} g^{-1} T^{-2} + X \sum_{g > T} g^{-3} \ll \frac{X \log T}{T^2}. \end{aligned}$$

□

With these estimates done, we move on to the main goal of this section, which is to establish the following L^1 estimate. We recall the use of \tilde{E} to denote the normalised additive energy E/N^3 .

Proposition 2.3. *We have*

$$\int_{\mathbb{T}} |F(\alpha) - F^*(\alpha)| d\alpha \ll \frac{\sqrt{\log T}}{T} (\tilde{E} \delta^{-1})^{1/2}.$$

Proof. Observe that

$$0 \leq F^*(\alpha) \leq F(\alpha),$$

and that

$$\int_{\mathbb{T}} F(\alpha) d\alpha = \frac{2}{N} \sum_{n \leq X} r(n) \frac{2s}{N}, \quad \int_{\mathbb{T}} F^*(\alpha) d\alpha = \frac{2}{N} \sum_{n \leq X} r(n) \mu(\mathcal{E}_n).$$

Therefore

$$\int_{\mathbb{T}} |F(\alpha) - F^*(\alpha)| d\alpha = \frac{2}{N} \sum_{n \leq X} r(n) \left(\frac{2s}{N} - \mu(\mathcal{E}_n) \right).$$

Using (2.1) it follows that

$$\begin{aligned} \int_{\mathbb{T}} |F(\alpha) - F^*(\alpha)| d\alpha &= \frac{2}{N} \sum_{n \leq X} r(n) \frac{2s}{N} \left(1 - \frac{\Phi(n)}{n} \right) \\ &\ll_s N^{-2} \sum_{n \leq X} r(n) \left(1 - \frac{\Phi(n)}{n} \right). \end{aligned}$$

Finally, Cauchy's inequality and Lemma 2.2 yield the desired inequality, since

$$E = N^2 + 2 \sum_{n \leq X} r(n)^2. \quad (2.2)$$

□

3. AN OVERLAP ESTIMATE

In this section, preparing for an L^2 argument, we bound the Lebesgue measure of the overlap $\mathcal{E}_n \cap \mathcal{E}_m$. For $m, n \in \mathbb{N}$, define

$$A(n, m) = \sum_{\substack{u \leq n, v \leq m \\ \frac{u}{n} = \frac{v}{m} \\ (u, n), (v, m) \leq T}} 1.$$

Note by symmetry that $A(n, m) = A(m, n)$. The following lemma is a minor adaptation of Lemma 4.4 of [7], in order to work with uniform cut-offs.

Lemma 3.1. *For $n \geq m \geq 1$ we have*

$$\mu(\mathcal{E}_n \cap \mathcal{E}_m) \leq \frac{4s^2}{N^2} + \frac{2s}{N} \frac{A(n, m)}{n}.$$

The constant 4 will be important, so we need to be quite precise here.

Proof. Considering separately the contributions from pairs of intervals for which $u/n = v/m$, it is easy to see that

$$\mu(\mathcal{E}_n \cap \mathcal{E}_m) \leq B_1 + B_2,$$

where

$$B_1 = \sum_{\substack{u \leq n, v \leq m \\ (u, n), (v, m) \leq T \\ u/n = v/m}} \frac{2s}{Nn} = \frac{2s}{N} \frac{A(n, m)}{n}$$

and

$$B_2 = \sum_{\substack{u \leq n, v \leq m \\ (u, n), (v, m) \leq T \\ u/n \neq v/m}} \mu \left(\left(\frac{u - s/N}{n}, \frac{u + s/N}{n} \right) \cap \left(\frac{v - s/N}{m}, \frac{v + s/N}{m} \right) \right).$$

It therefore remains to prove that $B_2 \leq 4s^2/N^2$.

If $u/n \neq v/m$ and the intervals $\left(\frac{u-s/N}{n}, \frac{u+s/N}{n}\right)$ and $\left(\frac{v-s/N}{m}, \frac{v+s/N}{m}\right)$ intersect then for some integer h we have

$$0 < |h| < (n+m)\frac{s}{N}, \quad \frac{u}{n} - \frac{v}{m} = \frac{h}{mn}.$$

The size of the overlap is bounded above by

$$\min\left(\frac{2s}{Nn}, \frac{s}{N}\left(\frac{1}{n} + \frac{1}{m}\right) - \frac{|h|}{mn}\right).$$

The number of positive integer solutions to $um - vn = h$ with $u \leq n$ and $v \leq m$ is bounded above by

$$\begin{cases} (m, n) & \text{if } (m, n) \mid h \\ 0, & \text{else.} \end{cases}$$

With $y = (n+m)s/N$, we now have

$$\begin{aligned} B_2 &\leq \sum_{\substack{0 < |h| < y \\ (m, n) \mid h}} (m, n) \min\left(\frac{2s}{Nn}, \frac{s}{N}\left(\frac{1}{n} + \frac{1}{m}\right) - \frac{|h|}{mn}\right) \\ &= \sum_{0 < |t| < y/(m, n)} (m, n) \min\left(\frac{2s}{Nn}, \frac{s}{N}\left(\frac{1}{n} + \frac{1}{m}\right) - \frac{(m, n)|t|}{mn}\right). \end{aligned}$$

The summand is non-increasing in $|t|$, so

$$\begin{aligned} B_2 &\leq 2 \int_0^{y/(m, n)} (m, n) \min\left(\frac{2s}{Nn}, \frac{s}{N}\left(\frac{1}{n} + \frac{1}{m}\right) - \frac{(m, n)t}{mn}\right) dt \\ &= 2 \int_0^y \min\left(\frac{2s}{Nn}, \frac{s}{N}\left(\frac{1}{n} + \frac{1}{m}\right) - \frac{h}{mn}\right) dh. \end{aligned}$$

Now

$$\begin{aligned} B_2 &\leq 2 \int_0^{(n-m)s/N} \frac{2s}{Nn} dh + 2 \int_{(n-m)s/N}^{(n+m)s/N} \left(\frac{s}{N}\left(\frac{1}{n} + \frac{1}{m}\right) - \frac{h}{mn}\right) dh \\ &= \frac{4s^2}{N^2} \frac{n-m}{n} + \frac{4s^2}{N^2} m \left(\frac{1}{n} + \frac{1}{m}\right) - \frac{1}{mn} \frac{s^2}{N^2} ((n+m)^2 - (n-m)^2), \end{aligned}$$

and finally we obtain

$$B_2 \leq \frac{4s^2}{N^2} \left(1 - \frac{m}{n} + \frac{m}{n} + 1 - 1\right) = \frac{4s^2}{N^2}.$$

□

We shall also need to bound the average overlap.

Lemma 3.2. *We have*

$$\sum_{n \leq X} \sum_{m \leq n} \frac{A(m, n)}{n} \ll X \log T.$$

Proof. We upper bound $\sum_{m \leq n} A(m, n)$ by neglecting one of the greatest common divisor constraints, i.e.

$$\sum_{m \leq n} A(m, n) \leq \sum_{\substack{1 \leq m, u, v \leq n \\ \frac{u}{m} = \frac{v}{n} \\ (v, n) \leq T}} 1.$$

Now, for fixed v , there are unique coprime positive integers a and b such that $\frac{v}{n} = \frac{a}{b}$. The denominator b is a divisor of n , and $a \leq b$. Further, with such an a and b fixed, the number of solutions to

$$\frac{a}{b} = \frac{u}{m}, \quad 1 \leq u, m \leq n$$

is exactly $\frac{n}{b}$.

We now consider how many times a particular divisor b of n can occur, as v ranges up to n . Indeed, since $b = \frac{n}{(v, n)}$, we see that b occurs exactly the number of times that $(v, n) = \frac{n}{b}$. If $\frac{n}{b} \leq T$, then this is at most b times; if $\frac{n}{b} > T$, then the greatest common divisor condition $(v, n) \leq T$ precludes this from ever occurring.

Therefore

$$\sum_{m \leq n} A(m, n) \leq \sum_{\substack{b|n \\ \frac{n}{b} \leq T}} b \frac{n}{b} = \sum_{\substack{c|n \\ c \leq T}} n.$$

It follows that

$$\sum_{n \leq X} \sum_{m \leq n} \frac{A(m, n)}{n} \leq \sum_{n \leq X} \sum_{\substack{c|n \\ c \leq T}} 1 \leq \sum_{c \leq T} \frac{X}{c} \ll X \log T$$

as claimed. \square

4. THE VARIANCE ESTIMATE

Like so much work on metric properties, we aim to show a result for almost all α by bounding the variance (α considered as a uniform random variable on \mathbb{T}). Rather than working directly with F , however, the objective of this section is instead to establish the following bound on the variance of F^* .

Proposition 4.1. *The variance of F^* satisfies*

$$\text{Var}(F^*) = \int_{\mathbb{T}} |F^*(\alpha) - \mathbb{E}F^*|^2 d\alpha \ll \frac{\sqrt{\log T}}{T} (\tilde{E}\delta^{-1})^{1/2} + \tilde{E}T \log T.$$

Proof. For brevity, we introduce the temporary notation

$$\rho = \frac{\sqrt{\log T}}{T} (\tilde{E}\delta^{-1})^{1/2}.$$

We will eventually choose T in such a way that $N^{-1} \leq \rho \leq 1$.

We begin by replacing $\mathbb{E}F^*$ with a simpler expression. Indeed, by Proposition 2.3 and the fact that $F^*(\alpha) \leq F(\alpha)$, we have

$$\mathbb{E}F^* = \mathbb{E}F + O(\rho).$$

Since

$$\mathbb{E}F = \int_{\mathbb{T}} F(\alpha) d\alpha = \frac{1}{N} \sum_{n \in \mathbb{Z} \setminus \{0\}} r(n) \frac{2s}{N} = 2s(1 - 1/N),$$

we must have

$$\mathbb{E}F^* = 2s + O(\rho). \quad (4.1)$$

A short calculation then tells us that

$$\text{Var}(F^*) = \int_{\mathbb{T}} (F^*(\alpha) - 2s)^2 d\alpha + O(\rho^2).$$

Expanding the main term gives

$$\frac{4}{N^2} \sum_{1 \leq n, m \leq X} r(n)r(m)\mu(\mathcal{E}_n \cap \mathcal{E}_m) - \frac{8s}{N} \sum_{1 \leq n \leq X} r(n)\mu(\mathcal{E}_n) + 4s^2.$$

By (2.1) and Lemma 3.1 this is at most

$$\begin{aligned} & \frac{16s^2}{N^4} \sum_{n, m \leq X} r(n)r(m) + \frac{16s}{N^3} \sum_{m \leq n \leq X} r(n)r(m) \frac{A(n, m)}{n} \\ & - \frac{16s^2}{N^2} \sum_{n \leq X} r(n) \frac{\Phi(n)}{n} + 4s^2. \end{aligned} \quad (4.2)$$

Now, as

$$\sum_{1 \leq n \leq X} r(n) = \frac{N^2}{2} + O(N),$$

we have

$$\frac{16s^2}{N^4} \sum_{1 \leq n, m \leq X} r(n)r(m) + 4s^2 = \frac{16s^2}{N^2} \sum_{1 \leq n \leq X} r(n) + O(N^{-1}).$$

Substituting this into (4.2) and bounding $\frac{1}{n}$ by $\frac{1}{\sqrt{mn}}$ when $m \leq n$ yields

$$\text{Var}(F^*) \ll S_1 + S_2 + N^{-1} + \rho^2, \quad (4.3)$$

where

$$S_1 = \frac{1}{N^2} \sum_{n \leq X} r(n) \left(1 - \frac{\Phi(n)}{n} \right)$$

and

$$S_2 = N^{-3} \sum_{m, n \leq X} r(m)r(n) \frac{A(m, n)}{\sqrt{mn}}.$$

Observe that the quantity S_1 was studied in the proof of Proposition 2.3, and in view of that calculation we have $S_1 \ll \rho$.

For S_2 , a more sensitive treatment of $A(m, n)$ than Lemma 3.2 will be required. We recall that $A(m, n)$ counts solutions $(u, v) \in [m] \times [n]$ to

$$un = vm, \quad (u, m), (v, n) \leq T.$$

The solutions are

$$u = \lambda \frac{m}{(m, n)}, \quad v = \lambda \frac{n}{(m, n)}$$

for $1 \leq \lambda \leq (m, n)$, subject to the further restriction given by T . As

$$(u, m) = \frac{m}{(m, n)}(\lambda, m, n), \quad (v, n) = \frac{n}{(m, n)}(\lambda, m, n),$$

we have

$$A(m, n) \leq \sum_{\substack{\lambda \leq (m, n): \\ (\lambda, m, n) \leq \frac{(m, n)}{\max\{m, n\}} T}} 1.$$

From this we extract two key pieces of information: that

$$A(m, n) \leq (m, n), \tag{4.4}$$

and that if $A(m, n) \neq 0$ then

$$\frac{m}{(m, n)}, \frac{n}{(m, n)} \leq T. \tag{4.5}$$

Just using (4.4) along with the gcd sum bounds of [4], which entirely removes the influence of T , we recover an analogous estimate to [2, Lemma 3] (see Remark 4.2 below). However, if we also use (4.5) and Cauchy–Schwarz, we obtain

$$\begin{aligned} S_2 &\leq \frac{1}{N^3} \sum_{\substack{m, n \leq X \\ \frac{m}{(m, n)}, \frac{n}{(m, n)} \leq T}} r(m)r(n) \frac{(m, n)}{\sqrt{mn}} \leq \frac{1}{N^3} \sum_{g \leq X} \left(\sum_{y \leq T} \frac{r(gy)}{\sqrt{y}} \right)^2 \\ &\leq \frac{1}{N^3} \sum_{g \leq X} \left(\sum_{y \leq T} \frac{1}{y} \right) \cdot \sum_{y \leq T} r(gy)^2 \ll \frac{\log T}{N^3} \sum_{n \in \mathbb{N}} r(n)^2 \sum_{\substack{y|n \\ y \leq T}} 1 \ll \tilde{E} T \log T. \end{aligned}$$

In view of (4.3), this completes the proof of the proposition. \square

Remark 4.2. There is an alternative approach to bounding S_2 , based on the general theory of gcd sums of the form

$$\sum_{k, \ell=1}^M \frac{(n_k, n_\ell)^{2\alpha}}{(n_k n_\ell)^\alpha}.$$

At $\alpha = 1/2$ we have already mentioned the essentially optimal bounds by Bondarenko and Seip [4, 5], which were employed in [2, Lemma 3]. We can pass to $\alpha = 1$ by Rankin’s trick, using (4.5); at this exponent there is Gál’s [6] prize-winning upper bound $O(M(\log \log M)^2)$. By [1, Lemma 4] and the discussion immediately following it, one can attach real weights to this at little cost, the idea being to apply this with weights $r(n)$. The sharpest version is

a recent breakthrough by Lewko and Radziwiłł [9, Theorem 2], by which we obtain the L^2 estimate

$$\sum_{k,\ell=1}^M c_k c_\ell \frac{(n_k, n_\ell)^2}{n_k n_\ell} \ll (\log \log M)^2 \sum_k c_k^2.$$

There are at most $M = N^2$ values of n for which $r(n) \neq 0$, so this approach ultimately yields

$$\text{Var}(F^*) \ll \frac{\sqrt{\log T}}{T} (\tilde{E} \delta^{-1})^{1/2} + T \tilde{E} (\log \log N)^2.$$

As it happens, for our eventual choice of T there is little quantitative difference between the two approaches, and certainly either is sufficient for Theorem 1.4.

5. IMPROVED ESTIMATES FOR THE RANDOM CASE

In this section we consider the setting of Theorem 1.5, and revisit the variance estimate from the preceding section. By a standard application of large deviation inequalities, we can assume that the representation function $r(n)$ is essentially constant. This leads to an improvement.

We begin with some easy bounds on N and $r(n)$.

Lemma 5.1. *Let $C > 2$, and let*

$$\psi(x) = \begin{cases} 1, & \text{if } x \leq 20, \\ (\log x)^{-1} (\log \log x)^{-C}, & \text{if } x > 20. \end{cases}$$

Let \mathcal{A} be as defined in the statement of Theorem 1.5, i.e. the random set of natural numbers defined by choosing $x \in \mathcal{A}$ independently at random with probability $\psi(x)$. Let $\varepsilon > 0$, and fix an integer X large enough in terms of ε . Consider the following properties:

(1) *N satisfies*

$$1 - \varepsilon \leq \frac{N}{X(\log X)^{-1}(\log \log X)^{-C}} \leq 1 + \varepsilon.$$

(2) *For all positive integers $n \leq X$, we have*

$$r(n) \leq (1 + \varepsilon) X (\log X)^{-2} (\log \log X)^{-2C}.$$

Then there is a constant $c_\varepsilon > 0$ such that property (1) holds with probability at least $1 - O(\exp(-c_\varepsilon X (\log X)^{-1} (\log \log X)^{-C}))$ and property (2) holds with probability at least $1 - O(X \exp(-c_\varepsilon X (\log X)^{-5}))$.

This is a quantitative version of a lemma which appears in Bourgain's appendix to [2], and is no doubt obvious to experts. Yet to keep the paper as self-contained as possible, we feel it is appropriate to provide the full details, particularly for part (2).

Proof. For each $x \in \mathbb{N}$, let ξ_x denote the Bernoulli random variable such that

$$\mathbb{P}(\xi_x = 1) = \psi(x).$$

Then $N = \sum_{x \leq X} \xi_x$ is a random variable with expectation $\sum_{x \leq X} \psi(x)$, which is asymptotic to $X(\log X)^{-1}(\log \log X)^{-C}$. Since the ξ_x are independent by assumption, one may settle part (1) immediately by applying the Chernoff bound, for instance by applying Corollary A.1.14 of [3].

For part (2), we first consider each $n \leq X - 1$ separately. Indeed, for x in the range $1 \leq x \leq X - n$, let $\omega_{x,n}$ denote the random variable $\omega_{x,n} = \xi_x \xi_{n+x}$. We have

$$r(n) = \sum_{x \leq X-n} \omega_{x,n},$$

which is a random variable with expectation $\sum_{x \leq X-n} \psi(x)\psi(n+x)$. Suppose first that $n \geq X - (1 + \varepsilon)X(\log X)^{-2}(\log \log X)^{-2C}$. Then by the trivial triangle inequality bound we have $r(n) \leq (1 + \varepsilon)X(\log X)^{-2}(\log \log X)^{-2C}$.

It remains to consider the case $n \leq X - (1 + \varepsilon)X(\log X)^{-2}(\log \log X)^{-2C}$. We wish to apply concentration of measure results for sums of independent random variables, and the family of random variables $\{\omega_{x,n} : x \leq X - n\}$ is very close to being independent. Methods of splitting this family into groups of genuinely independent random variables are alluded to in the discussion in the appendix of [2]. Here we describe an extremely coarse decomposition which nonetheless is strong enough for our purposes.

Split $[X - n]$ into two sets, \mathcal{S}_0 and \mathcal{S}_1 , constructed as follows. If $n \leq X/3$, let $x \in \mathcal{S}_j$ if $\lfloor x/n \rfloor \equiv j \pmod{2}$. If $n > X/3$, instead let

$$\mathcal{S}_0 = \left\{x \in \mathbb{N} : x \leq \frac{X - n}{2}\right\}, \quad \mathcal{S}_1 = \left\{x \in \mathbb{N} : \frac{X - n}{2} < x \leq X - n\right\}.$$

For each $j \in \{0, 1\}$ the family $\{\omega_{x,n} : x \in \mathcal{S}_j\}$ is independent, as no two indices differ by n . Applying the union bound and then Corollary A.1.14 of [3] once more, we have

$$\begin{aligned} & \mathbb{P}\left(r(n) \geq (1 + \varepsilon) \sum_{x \leq X-n} \psi(x)\psi(n+x)\right) \\ & \ll \mathbb{P}\left(\sum_{x \in \mathcal{S}_0} \omega_{x,n} \geq (1 + \varepsilon) \sum_{x \in \mathcal{S}_0} \psi(x)\psi(n+x)\right) \\ & \quad + \mathbb{P}\left(\sum_{x \in \mathcal{S}_1} \omega_{x,n} \geq (1 + \varepsilon) \sum_{x \in \mathcal{S}_1} \psi(x)\psi(n+x)\right) \\ & \ll \exp\left(-c_\varepsilon \sum_{x \in \mathcal{S}_0} \psi(x)\psi(n+x)\right) + \exp\left(-c_\varepsilon \sum_{x \in \mathcal{S}_1} \psi(x)\psi(n+x)\right). \end{aligned} \quad (5.1)$$

It remains to estimate these final quantities. Note that by construction we have $\min(|\mathcal{S}_0|, |\mathcal{S}_1|) \geq (X - n)/4$. So, since $\psi(x)$ is weakly decreasing, we

deduce from (5.1) that

$$\begin{aligned} \mathbb{P}\left(r(n) \geq (1 + \varepsilon) \sum_{x \leq X-n} \psi(x)\psi(n+x)\right) &\ll \exp\left(-c_\varepsilon \sum_{X - \frac{X-n}{4} < x \leq X} \psi(x)^2\right) \\ &\ll \exp(-c_\varepsilon X(\log X)^{-5}) \end{aligned}$$

by a simple calculation, reducing the quantity c_ε as necessary.

By a similar monotonicity principle, $\mathbb{P}(r(n) \geq (1 + \varepsilon) \sum_{x \leq X-n} \psi(x)\psi(n+x))$ is at least $\mathbb{P}(r(n) \geq (1 + \varepsilon) \sum_{x \leq X} \psi(x)^2)$. Since

$$\sum_{x \leq X} \psi(x)^2 \sim X(\log X)^{-2}(\log \log X)^{-2C},$$

and X large enough in terms of ε , we conclude that

$$r(n) \leq (1 + \varepsilon)X(\log X)^{-2}(\log \log X)^{-2C}$$

with probability greater than $1 - O(\exp(-c_\varepsilon X(\log X)^{-5}))$.

All these calculations were done for a single n . But applying a crude union bound over all $n \leq X - 1$, and noting that $r(X) = 0$, part (2) is proved. \square

Corollary 5.2. *With probability 1 there exists some X_0 such that properties (1) and (2) hold for all $X \geq X_0$.*

Proof. The sums

$$\sum_{X \geq 1} \exp(-c_\varepsilon X(\log X)^{-1}(\log \log X)^{-C})$$

and

$$\sum_{X \geq 1} X \exp(-c_\varepsilon X(\log X)^{-5})$$

are both convergent. So by the first Borel–Cantelli lemma, the corollary follows. \square

For the rest of this paper, whenever we consider the random case we restrict to the probability 1 event from Corollary 5.2.

To continue this section, we use the above work to get an improved version of Proposition 4.1, in the random case. First we improve on Proposition 2.3.

Proposition 5.3. *In the setting of Theorem 1.5, we have*

$$\int_{\mathbb{T}} |F(\alpha) - F^*(\alpha)| \, d\alpha \ll T^{-1}.$$

Proof. Following Proposition 2.3, we establish that

$$\int_{\mathbb{T}} |F(\alpha) - F^*(\alpha)| \, d\alpha = 2 \sum_{n \leq X} \frac{r(n)}{N} \frac{2s}{N} \left(1 - \frac{\Phi(n)}{n}\right) \ll X^{-1} \sum_{n \leq X} \left(1 - \frac{\Phi(n)}{n}\right).$$

The final inequality follows from Corollary 5.2. Then we apply Lemma 2.1, and the proposition follows. \square

We now go on to improve Proposition 4.1.

Proposition 5.4. *Let \mathcal{A} be as in Theorem 1.5. Then the variance of F^* satisfies*

$$\mathrm{Var}(F^*) \ll T^{-1} + (\log X)^{-1}(\log \log X)^{-C} \log T.$$

Proof. We will eventually choose T in such a way that $N^{-1} \leq T^{-1} \leq 1$. Following the proof of Proposition 4.1, we thus derive

$$\mathrm{Var}(F^*) \ll S_1 + S_3 + N^{-1} + T^{-2},$$

where

$$S_3 = N^{-3} \sum_{m \leq n \leq X} r(n)r(m) \frac{A(m, n)}{n}.$$

Observe that S_1 is the same expression considered in Proposition 5.3, and so $S_1 \ll T^{-1}$. It remains to bound S_3 , for which we already have all the necessary lemmas in place. Indeed, by Corollary 5.2, and by Lemma 3.2, we have

$$\begin{aligned} S_3 &\ll X^{-1}(\log X)^{-1}(\log \log X)^{-C} \sum_{m \leq n \leq X} \frac{A(m, n)}{n} \\ &\ll (\log X)^{-1}(\log \log X)^{-C} \log T. \end{aligned}$$

Combining everything together yields the proposition. \square

6. SANDWICHING, CONCENTRATION, AND BOREL–CANTELLI

Here we conclude the proofs of Theorems 1.4 and 1.5. We begin by proving the following assertion, which we will see implies that \mathcal{A} is metric Poissonian.

Proposition 6.1. *Let $\varepsilon, s > 0$. Let \mathcal{A} satisfy either the conditions of Theorem 1.4 or 1.5. Then there is a full measure set $\Omega_{\varepsilon, s}$ such that if $\alpha \in \Omega_{\varepsilon, s}$ then there exists $X_0 = X_0(\alpha, \varepsilon, s)$ such that if $X \geq X_0$ then $|F(\alpha) - 2s| \leq \varepsilon$.*

First, let us describe the argument in the deterministic setting of Theorem 1.4.

Proof. Define the sequence $N_j = \lfloor 2^{j^{1-\eta}} \rfloor$, where $\eta > 0$ is small in terms of ξ . Note that $N_{j+1}/N_j \rightarrow 1$. For $j \in \mathbb{N}$, define $X_j \in \mathbb{N}$ to be minimal such that $N_j = |\mathcal{A} \cap [X_j]|$. Let X be large in terms of ε and s , and suppose $X_j \leq X < X_{j+1}$. We begin with the inequalities noted in [2], namely

$$N_j F(\alpha, s \frac{N_j}{N_{j+1}}, X_j) \leq NF(\alpha, s, X) \leq N_{j+1} F(\alpha, s \frac{N_{j+1}}{N_j}, X_{j+1}),$$

which follow immediately from the definitions. Note that

$$s \frac{N_j}{N_{j+1}}, s \frac{N_{j+1}}{N_j} \asymp 1, \quad s \frac{N_j}{N_{j+1}}, s \frac{N_{j+1}}{N_j} \rightarrow s.$$

Motivated by Proposition 4.1, we choose $T = (\tilde{E}\delta)^{-1/4} + 1 \geq 2$. By Proposition 2.3, we thus have

$$\int_{\mathbb{T}} |F(\alpha) - F^*(\alpha)| d\alpha \ll (\log T)^{1/2} \tilde{E}^{3/4} \delta^{-1/4}.$$

At this stage we invoke our hypotheses (1.1): with $y := \tilde{E}^{-1} \gg (\log N)^{2+\xi}$, we have

$$(\log T)^{1/2} \tilde{E}^{3/4} \ll \frac{\sqrt{\log y} + \sqrt{\log \log N}}{y^{3/4}}.$$

The right hand side is decreasing in y for $y \geq 100$, so by our initial assumptions (1.1) we obtain

$$\int_{\mathbb{T}} |F(\alpha) - F^*(\alpha)| d\alpha \ll \frac{1}{(\log N)^{1+\xi/5}}. \quad (6.1)$$

The same calculation also shows that $N^{-1} \leq \rho \leq 1$, which was assumed in the proof of Proposition 4.1. Applying Proposition 4.1, therefore, we have

$$\text{Var}(F^*) \ll \tilde{E}^{3/4} \delta^{-1/4} \log T \ll \frac{1}{(\log N)^{1+\xi/5}}. \quad (6.2)$$

The L^1 estimate (6.1) implies that if $s \asymp 1$ and X is large then

$$|F(\alpha, s, X) - F^*(\alpha, s, X)| \leq \varepsilon/4 \quad (\alpha \notin E_{X,s}),$$

for some exceptional set $E_{X,s}$ with

$$\mu(E_{X,s}) \ll_{\varepsilon} \frac{1}{(\log N)^{1+\xi/5}}.$$

In particular

$$\sum_{j=1}^{\infty} \mu(E_{X_j, s \frac{N_j}{N_{j+1}}}) \ll \sum_{j=1}^{\infty} j^{(\eta-1)(1+\xi/5)} < \infty,$$

provided that η is chosen small enough depending on ξ . Now the Borel–Cantelli lemma tells us that for almost all $\alpha \in \mathbb{T}$ we have

$$\left| F(\alpha, s \frac{N_j}{N_{j+1}}, X_j) - F^*(\alpha, s \frac{N_j}{N_{j+1}}, X_j) \right| \leq \varepsilon/4 \quad (j \geq j_1(\alpha, \varepsilon)),$$

and similarly

$$\left| F(\alpha, s \frac{N_{j+1}}{N_j}, X_{j+1}) - F^*(\alpha, s \frac{N_{j+1}}{N_j}, X_{j+1}) \right| \leq \varepsilon/4 \quad (j \geq j_2(\alpha, \varepsilon)).$$

Recalling (4.1), the variance estimate (6.2) on F^* combined with Chebyshev's inequality implies that

$$\mu\left(\left\{\alpha \in \mathbb{T} : \left| F^*(\alpha, s \frac{N_j}{N_{j+1}}, X_j) - 2s \frac{N_j}{N_{j+1}} \right| > \varepsilon/4 \right\}\right) \ll_{\varepsilon} (\log N_j)^{-1-\xi/5}.$$

We again apply the Borel–Cantelli lemma, and find that for almost all $\alpha \in \mathbb{T}$ we have

$$\left| F^*(\alpha, s \frac{N_j}{N_{j+1}}, X_j) - 2s \frac{N_j}{N_{j+1}} \right| \leq \varepsilon/4 \quad (j \geq j_3(\alpha, \varepsilon)),$$

and similarly

$$\left| F^*(\alpha, s \frac{N_{j+1}}{N_j}, X_{j+1}) - 2s \frac{N_{j+1}}{N_j} \right| \leq \varepsilon/4 \quad (j \geq j_4(\alpha, \varepsilon)).$$

Now the triangle inequality gives, for almost all $\alpha \in \mathbb{T}$ and all $j \geq j_5(\alpha, \varepsilon)$,

$$F(\alpha, s \frac{N_j}{N_{j+1}}, X_j) \geq 2s \frac{N_j}{N_{j+1}} - \varepsilon/2$$

and

$$F(\alpha, s \frac{N_{j+1}}{N_j}, X_{j+1}) \leq 2s \frac{N_{j+1}}{N_j} + \varepsilon/2.$$

Since

$$F(\alpha, s \frac{N_j}{N_{j+1}}, X_j) \leq \frac{N}{N_j} F(\alpha, s, X)$$

and

$$F(\alpha, s \frac{N_{j+1}}{N_j}, X_{j+1}) \geq \frac{N}{N_{j+1}} F(\alpha, s, X),$$

we obtain

$$\frac{N_j}{N} \left(2s \frac{N_j}{N_{j+1}} - \varepsilon/2 \right) \leq F(\alpha, s, X) \leq \frac{N_{j+1}}{N} \left(2s \frac{N_{j+1}}{N_j} + \varepsilon/2 \right).$$

As j is large and $N_{j+1}/N_j \rightarrow 1$, we conclude as claimed that

$$|F(\alpha, s, X) - 2s| \leq \varepsilon.$$

□

Now let us prove Proposition 6.1 in the random setting of Theorem 1.5.

Proof. For some small $\eta > 0$, ultimately depending on ε and s , define the sequence $X_j = \lfloor 2^{j\eta} \rfloor$. Let X be large in terms of ε and s , and suppose that $X_j \leq X < X_{j+1}$. We note again the sandwiching inequalities

$$N_j F(\alpha, s \frac{N_j}{N_{j+1}}, X_j) \leq N F(\alpha, s, X) \leq N_{j+1} F(\alpha, s \frac{N_{j+1}}{N_j}, X_{j+1}).$$

Observe that

$$s \frac{N_j}{N_{j+1}}, s \frac{N_{j+1}}{N_j} \asymp 1,$$

and that for j large enough in terms of η we have

$$\frac{N_j}{N_{j+1}} \geq 2^{-2\eta}, \quad \frac{N_{j+1}}{N_j} \leq 2^{2\eta}.$$

Motivated by Proposition 5.4, we choose

$$T = \log^2 X.$$

By Proposition 5.3, we find that if $s \asymp 1$ and X is large in terms of η then

$$|F(\alpha, s, X) - F^*(\alpha, s, X)| \leq \eta/4 \quad (\alpha \notin E_{X,s}),$$

for some exceptional set $E_{X,s}$ with

$$\mu(E_{X,s}) \ll_\eta (\log X)^{-2}.$$

This means that

$$\sum_{j=1}^{\infty} \mu(E_{X_j, s \frac{N_j}{N_{j+1}}}) \ll_\eta \sum_{j=1}^{\infty} j^{-2} < \infty.$$

We apply the first Borel–Cantelli lemma as in the previous proof, concluding that for almost all $\alpha \in \mathbb{T}$ we have

$$\left| F(\alpha, s \frac{N_j}{N_{j+1}}, X_j) - F^*(\alpha, s \frac{N_j}{N_{j+1}}, X_j) \right| \leq \eta/4 \quad (j \geq j_1(\alpha, \eta)),$$

and similarly

$$\left| F(\alpha, s \frac{N_{j+1}}{N_j}, X_{j+1}) - F^*(\alpha, s \frac{N_{j+1}}{N_j}, X_{j+1}) \right| \leq \eta/4 \quad (j \geq j_2(\alpha, \eta)).$$

By Proposition 5.4, the variance of F^* is bounded above by a constant times

$$(\log X)^{-2} + (\log X)^{-1}(\log \log X)^{-C+1}.$$

We absorb the first term, and have by Chebychev's inequality that

$$\begin{aligned} \mu\left(\left\{\alpha \in \mathbb{T} : \left| F^*(\alpha, s \frac{N_j}{N_{j+1}}, X_j) - 2s \frac{N_j}{N_{j+1}} \right| > \eta/4 \right\}\right) \\ \ll_{\eta} (\log X_j)^{-1} (\log \log X_j)^{-C+1}. \end{aligned}$$

We again apply the Borel–Cantelli lemma, using the fact that

$$\sum_{j=1}^{\infty} (\log X_j)^{-1} (\log \log X_j)^{-C+1} \ll \eta^{-1} \sum_{j=1}^{\infty} j^{-1} (\log \eta j)^{-C+1} < \infty$$

since $C > 2$. Thus we again find that for almost all $\alpha \in \mathbb{T}$ we have

$$\left| F^*(\alpha, s \frac{N_j}{N_{j+1}}, X_j) - 2s \frac{N_j}{N_{j+1}} \right| \leq \eta/4 \quad (j \geq j_3(\alpha, \eta)),$$

and similarly

$$\left| F^*(\alpha, s \frac{N_{j+1}}{N_j}, X_{j+1}) - 2s \frac{N_{j+1}}{N_j} \right| \leq \eta/4 \quad (j \geq j_4(\alpha, \eta)).$$

The rest of the proof proceeds as in the deterministic case, reaching the expression

$$\frac{N_j}{N} \left(2s \frac{N_j}{N_{j+1}} - \eta/2 \right) \leq F(\alpha, s, X) \leq \frac{N_{j+1}}{N} \left(2s \frac{N_{j+1}}{N_j} + \eta/2 \right).$$

It is not true in this setting that $N_j/N_{j+1} \rightarrow 1$, but by our earlier observations we may establish

$$2^{-2\eta} \left(2s \cdot 2^{-2\eta} - \frac{\eta}{2} \right) \leq F(\alpha, s, X) \leq 2^{2\eta} \left(2s \cdot 2^{2\eta} + \frac{\eta}{2} \right),$$

and therefore

$$F(\alpha, s, X) = 2s + O_s(\eta).$$

Choosing η small enough, this error is at most ε , thus proving the proposition. \square

To complete the proofs of Theorems 1.4 and 1.5, it remains to deduce the metric Poissonian property from Proposition 6.1. Apply Proposition 6.1 to all $s \in \mathbb{Q}_{>0}$ simultaneously, with an exceptional set of measure zero. Now let s lie in a short interval (s_1, s_2) , where $s_1, s_2 \in \mathbb{Q}$, and the desired conclusion follows.

7. THE RANDOM DIVERGENCE THEORY

In this section we prove Theorem 1.6. We keep this brief, as the proof is straightforward by combining section 5 with the crux of [14].

By Khintchine's theorem [7, Theorem 2.2], we know that there is a full measure set Ω such that for all $\alpha \in \Omega$ there exist arbitrarily large positive integers M such that

$$\|M\alpha\| < (M\mathcal{L}(M))^{-1}, \quad (7.1)$$

where

$$\mathcal{L}(M) = \prod_{i \leq 5} \log_i(M).$$

We remind the reader that here $\log_0(M) = M$ and $\log_t(M) = \log(\log_{t-1}(M))$ for $t \in \mathbb{N}$.

Now let \mathcal{A} be as in Theorem 1.6. With negligible alteration, the proofs of Lemma 5.1 and Corollary 5.2 demonstrate that the associated quantities δ and $r(n)$ (for $1 \leq n \leq X/3$ with X large) have the expected order of magnitude. We conclude specifically that, with probability 1, there exists X_0 such that the following hold.

(1) If $X \geq X_0$ then

$$\frac{1}{2}(\log X)^{-1}(\log \log X)^{-C} \leq \delta \leq 2(\log X)^{-1}(\log \log X)^{-C}.$$

(2) If $X \geq X_0$ and $1 \leq n \leq X/3$ then $r(n) \geq \frac{1}{8}\delta N$.

Now let α be a member of Ω , and suppose that there exists an X_0 satisfying (1) and (2) above. Let $s > 0$ be constant, let M be a large positive integer satisfying (7.1), and put

$$N = \lfloor M \log_3(M) \rfloor, \quad K = \log M (\log \log M)^C \log_4 M.$$

Note that property (1) implies that $X \gg M(\log M)(\log \log M)^C(\log_3 M)$, and in particular that $KM \leq X/3$ (for large enough M).

By considering those n of the form $n = kM$ ($1 \leq k \leq K$), and using property (2), we have that

$$F(N) \geq N^{-1} \sum_{\substack{n \leq X/3 \\ \|n\alpha\| < s/N}} r(n) \gg \delta K \gg \log_4(M).$$

In particular, with probability 1, for all $\alpha \in \Omega$ there exist arbitrarily large N such that $F(N) > 3s$. Hence, with probability 1, \mathcal{A} is not metric Poissonian.

APPENDIX A. THE ENERGY IN THE RANDOM SETTING

In this appendix we include the computation of the additive energy in the random setting of Theorem 1.5. Recall (2.2). By Cauchy–Schwarz and Corollary 5.2, we have $\log N \asymp \log X$ and

$$E \geq \sum_{n \leq X} r(n)^2 \geq X^{-1} \left(\sum_{n \leq X} r(n) \right)^2 \asymp \frac{N^4}{X} \asymp \frac{N^3}{(\log N) \cdot (\log \log N)^C}$$

almost surely. Moreover, Corollary 5.2 also yields

$$E \ll N^2 + \sum_{n \leq X} (N^2/X)^2 \ll \frac{N^4}{X} \asymp \frac{N^3}{(\log N) \cdot (\log \log N)^C}$$

almost surely. We thus conclude that

$$E \asymp \frac{N^3}{(\log N) \cdot (\log \log N)^C}$$

with probability 1.

REFERENCES

- [1] C. Aistleitner, I. Berkes and K. Seip, *GCD sums from Poisson integrals and systems of dilated functions*, J. Eur. Math. Soc. (JEMS) **017** (2015), 1517–1546.
- [2] C. Aistleitner, G. Larcher and M. Lewko, *Additive energy and the Hausdorff dimension of the exceptional set in metric pair correlation problems*, with an appendix by J. Bourgain, Israel J. Math. **222** (2017), 463–485.
- [3] N. Alon and J. Spencer, *The Probabilistic Method*, third edition, John Wiley & Sons, Inc., Hoboken, NJ, 2008.
- [4] A. Bondarenko and K. Seip, *GCD sums and complete sets of square-free numbers*, Bull. London Math. Soc. **47** (2015), 29–41.
- [5] A. Bondarenko and K. Seip, *Large greatest common divisor sums and extreme values of the Riemann zeta function*, Duke Math. J. **166** (2017), 1685–1701.
- [6] I. S. Gál, *A theorem concerning Diophantine approximations*, Nieuw Arch. Wiskunde **23** (1949), 13–38.
- [7] G. Harman, *Metric number theory*, London Mathematical Society Monographs. New Series, 18. The Clarendon Press, Oxford University Press, New York, 1998.
- [8] T. Lachmann and N. Technau, *On Exceptional Sets in the Metric Poissonian Pair Correlations Problem*, arXiv:1708.08599.
- [9] M. Lewko and M. Radziwiłł, *Refinements of Gál’s theorem and applications*, Adv. Math. **305** (2017), 280–297.
- [10] J. Marklof, *Distribution modulo one and Ratner’s theorem*, Equidistribution in number theory, an introduction, 217–244, NATO Sci. Ser. II Math. Phys. Chem., 237, Springer, Dordrecht, 2007.
- [11] Z. Rudnick and P. Sarnak, *The pair correlation function of fractional parts of polynomials*, Comm. Math. Phys. **194** (1998), 61–70.
- [12] Z. Rudnick and A. Zaharescu, *A metric result on the pair correlation of fractional parts of sequences*, Acta Arith. **89** (1999), 283–293.
- [13] W. Schmidt, *A metrical theorem in diophantine approximation*, Canad. J. Math. **12** (1960), 619–631.
- [14] A. Walker, *The primes are not metric Poissonian*, Mathematika, to appear.

HEILBRONN INSTITUTE FOR MATHEMATICAL RESEARCH, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRISTOL, UNIVERSITY WALK, BRISTOL BS8 1TW, UNITED KINGDOM

E-mail address: `matfb@bristol.ac.uk`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, HESLINGTON, YORK YO10 5DD, UNITED KINGDOM

E-mail address: `sam.chow@york.ac.uk`

DEPARTMENT OF MATHEMATICS, 915 Hylan Building, UNIVERSITY OF ROCHESTER, ROCHESTER, NY 14627

E-mail address: `agafni@ur.rochester.edu`

ANDREW WILES BUILDING, UNIVERSITY OF OXFORD, RADCLIFFE OBSERVATORY QUARTER, WOODSTOCK RD, OXFORD OX2 6GG, UNITED KINGDOM

E-mail address: `walker@maths.ox.ac.uk`