



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/127991/>

Version: Accepted Version

Article:

MacVicar, S, Berrang-Ford, L, Harper, S et al. (2017) How seasonality and weather affect perinatal health: Comparing the experiences of indigenous and non-indigenous mothers in Kanungu District, Uganda. *Social Science & Medicine*, 187. pp. 39-48. ISSN: 0277-9536

<https://doi.org/10.1016/j.socscimed.2017.06.021>

© 2017 Published by Elsevier Ltd. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0>.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

1 **How seasonality and weather affect perinatal health: comparing the experiences of**
2 **Indigenous and non-Indigenous mothers in Kanungu District, Uganda**

3

4 **Abstract**

5 Maternal and newborn health disparities and the health impacts of climate change present grand
6 challenges for global health equity, and there remain knowledge gaps in our understanding of how
7 these challenges intersect. This study examines the pathways through which mothers are affected by
8 seasonal and meteorological factors in sub-Saharan Africa in general, and Kanungu District
9 (Uganda), in particular. We conducted a community-based study consisting of focus group
10 discussions with mothers and interviews with health care workers in Kanungu District. Using *a priori*
11 and *a posteriori* coding, we found a diversity of perspectives on the impacts of seasonal and weather
12 exposures, with reporting of more food available in the rainy season. The rainy season was also
13 identified as the period in which women performed physical labour for longer time periods, while
14 work conditions in the dry season were reported to be more difficult due to heat. The causal
15 pathways through which weather and seasonality may be affecting size at birth as reported by
16 Kanungu mothers were consistent with those most frequently reported in the literature elsewhere,
17 including maternal energy balance (nutritional intake and physical exertion output) and seasonal
18 illness. While both Indigenous and non-Indigenous mothers described similar pathways, however,
19 the *severity* of these experiences differed. Non-Indigenous mothers frequently relied on livestock
20 assets or opportunities for less taxing physical work than Indigenous women, who had fewer
21 options when facing food shortages or transport costs. Findings point to specific entry points for
22 intervention including increased nutritional support in dry season periods of food scarcity, increased
23 diversification of wage labour opportunities, and increased access to contraception. Interventions

24 should be particularly targeted towards Indigenous mothers as they face greater food insecurity, may
25 have fewer sources of income, and face greater overall deprivation than non-Indigenous mothers.

26 **Keywords:** Uganda; perinatal health; climate change; weather; season; Indigenous health

27

28 **1 Introduction**

29 Climate change will have substantial impacts for populations worldwide (Watts et al. 2015, Smith et
30 al. 2014). In response, research investigating climate impacts on health has grown rapidly in the past
31 two decades. Research has, however, primarily focused on health outcomes with relatively direct or
32 proximal relationships with climate (e.g., infectious disease, heat-related illness, mortality due to
33 extreme events (Smith et al. 2014)). Less attention has been paid to health outcomes with more
34 indirect or distal links to climate or weather, despite predictions that most climate impacts are
35 mediated through indirect social and ecological factors (Watts et al. 2015). Research has shown, for
36 example, an empirical relationship between weather and/or season and size at birth in multiple
37 regions and contexts (Strand, Barnett and Tong 2011, Laaidi et al. 2011, Beltran, Wu and Laurent
38 2014).

39 While the effects of climate change will have health impacts worldwide (Costello et al. 2009,
40 Few 2007, Haines and Patz 2004, McMichael, Woodruff and Hales 2006, Watts et al. 2015, Smith et
41 al. 2014), Indigenous populations will face a disproportionate burden of these negative impacts due
42 to their traditional reliance on biophysical resources and existing health inequalities. (Ford 2012).

43 The holistic definition of health in many Indigenous populations means environmental and
44 individual health are seen as inextricably linked, thus the effects of climate change on the
45 environmental are also felt at a personal level in such communities (Berry et al. 2010).

46 Connectedness to the land (both physical and spiritual) is seen as one of the essential components to
47 Indigenous health, and therefore a critical consideration in the planning of health adaptations to

48 climate change in these communities (Green and Minchin 2014). Much of the literature on
49 Indigenous health comes from North America, Australia, and New Zealand, and to a lesser extent,
50 South and Central America. There is a limited body of literature regarding the health of Indigenous
51 populations in Africa, where the existing burden of ill-health is high and investigation of differential
52 vulnerability between ethnic groups has been negligible (Ohenjo et al. 2006).

53 The burden of climate change on maternal and infant health will be inequitably distributed.
54 The groups already facing the greatest vulnerability—women, the poor, and Indigenous
55 populations—have been identified as being particularly at risk for adverse health impacts due to
56 climate change (Watts et al. 2015, McMichael et al. 2006, Smith et al. 2014). As Busby et al. (2013)
57 write, ethnicity may prove a key determinant in differential vulnerability to climate change. Already,
58 Indigenous mothers in remote areas often face inequities in perinatal health and are at risk of poorer
59 perinatal outcomes than non-Indigenous mothers (Graham et al. 2007, Gracey and King 2009); poor
60 women in both industrialized and low-resource settings tend to face disparities when compared to
61 their wealthier counterparts (Kramer et al. 2000, Lawn et al. 2009). There is a double-burden of
62 maternal and infant health inequity in populations reliant on subsistence agriculture: subsistence-
63 based Indigenous women are among the most vulnerable populations in the world due to persistent
64 health inequality and reliance on fluctuating food sources.

65 In both developed and developing settings, low birth weight is considered an important
66 determinant of infant mortality (Kramer 1987, Lawn, Cousens and Zupan 2005, McCormick 1985).
67 More than 80% of neonatal deaths in sub-Saharan Africa and south Asia occur in small babies as a
68 result of both preterm births and intrauterine growth restriction (IUGR) (Lawn et al. 2014). The
69 effects of being born small can persist throughout infancy and childhood (Lawn et al. 2014, Paneth
70 1995). A number of studies indicate that IUGR may increase risk of a range of adult-onset
71 conditions (Botero and Lifshitz 1999, Barker et al. 2002, Harding 2001, Kajantie et al. 2005, Low et

72 al. 1992). A combination of birth weight, gestational age, and Apgar scores are the recommended
73 predictors for infant mortality (Ma and Finch 2010), and are often the outcomes of interest when
74 examining in utero exposures (Chou et al. 2014, Porpora et al. 2013).

75 Maternal energy balance (i.e., food intake versus physical activity output) and seasonal
76 patterns in infectious disease (particularly malaria) are theorized in the literature as the predominant
77 pathways through which season and weather affect pregnant women in low-income country settings
78 (Rayco-Solon, Fulford and Prentice 2005, Laaidi et al. 2011, Beltran et al. 2014). For populations
79 reliant on subsistence agriculture, seasonal food shortages and seasonal trends in agricultural labour
80 activities are affected by weather and seasonality. Though research is mixed, most literature has
81 reported that increased incidence of lower birth weights coincides with periods of increased energy
82 expenditure, particularly when these periods coincide with food shortages. Variation in patterns of
83 malaria and other infectious diseases have also been associated with both weather and birth
84 outcomes, with malaria believed to be a key pathway for low birth weights resulting from preterm
85 births (Kinabo 1993; Rayco-Solon et al. 2005). Grace et al. (2015) investigate the role of weather on
86 birth outcomes across Africa, characterizing the variation in the relationships between weather and
87 birth weight across different livelihood zones . What remains unclear, however, is the extent to
88 which these aggregated results mask heterogeneity in the effect of weather on birth outcomes in
89 different countries, contexts, and *within* livelihood zones.

90 Climate, agricultural practices (Grace et al. 2015), birthing and pregnancy cultures (Brighton
91 et al. 2013, Magadi, Madise and Rodrigues 2000), and malaria transmission patterns (Noor et al.
92 2014) differ regionally, and the direction, magnitude, and nature of weather and/or climatic
93 determinants of infant health vary in diverse contexts (Strand et al. 2011, Laaidi et al. 2011, Beltran
94 et al. 2014, Carolan-Olah and Frankowska 2014, Chodick et al. 2009). Further, inequities in maternal
95 and infant health are multifactorial in origin, and arise based on different factors in different

96 contexts (Wirth et al. 2006). This regional diversity necessitates local studies characterizing the place-
97 specific experience of weather and season during pregnancy. Where regional differences do exist,
98 we do not yet have a complete picture as to how and why they differ—do the pathways through
99 which weather and pregnancy experiences may be linked differ in different regions? To date, there
100 has been limited qualitative investigation characterizing the pathways linking weather to birth
101 outcomes, and whether these are heterogeneous across and within populations. With the exception
102 of one paper among nomadic Turkana pastoralists (Pike 2000), we are aware of no studies exploring
103 the pathways by which weather influences perinatal health using empirical results from fieldwork
104 and qualitative analysis.

105 We address this research gap by bringing a qualitative lens to a problem that has primarily
106 benefitted from quantitative examination. This paper contributes to our understanding of how and
107 why season and weather influence pregnant mothers and newborns in a rural east African setting,
108 specifically among Indigenous and non-Indigenous subsistence-based populations in rural Uganda
109 where we have previously identified associations between *in utero* rainfall and temperature exposures
110 and birth weight [Anonymous Under review A]. Despite significant progress through Millennium
111 Development Goals initiatives, maternal mortality in Uganda remains 325 per 100 000 (Kassebaum
112 et al. 2014) and infant mortality 22 per 1000 live births (United Nations Inter-agency for Child
113 Mortality Estimation (UN IGME) 2014). We focus on some of the most vulnerable mothers within
114 this context (Indigenous women whose babies are born on average 295.5 g smaller than the general
115 population) [Anonymous under review A (on request)], and identify the most pressing needs of
116 these mothers and develop initiatives to enhance maternal resilience. The unique context in
117 Kanungu, a region in which Indigenous and non-Indigenous populations live in the same
118 environment with the same health services, permits study of the social modifiers of women’s
119 experiences during pregnancy. In this context, study objectives were to: 1) qualitatively characterize

120 how mothers and health care worker key informants perceive seasonal and weather exposures to
121 influence pregnancy and birth in Kanungu District, Uganda, and 2) assess compare the experiences
122 of these pathways between Indigenous and non-Indigenous mothers.

123

124 **2 Methods**

125 **2.1 Theoretical approach**

126 This study is grounded in the theories and methods of health geography, and is guided by the
127 integral consideration within health geography of *place* and *space* as important predictors of health
128 (Jones and Moon 1993, Macintyre, Ellaway and Cummins 2002). By situating this research as a
129 geographic inquiry, there is a freedom to not only assess whether or not a relationship exists
130 between environmental conditions and birth outcomes, but also to investigate *how* and *why* it exists.
131 One of the key elements in the shift from medical geography to health geography is the emphasis on
132 the use of critical theory to understand health disparities as a product of larger-scale forces and
133 systematic inequities (Cutchin 2007). Our inquiry is driven by the vulnerability approach (Adger
134 2006) which recognizes that climate vulnerability will manifest through existing social gradients. We
135 tackle the question of environmental impacts on pregnancy from a critical realist perspective (Yeung
136 1997), an approach focused on translating this knowledge into social change (Ng and Muntaner
137 2014) and emphasizing the policy relevance of findings (Fletcher 2017).

138 **2.2 Study population**

139 Kanungu District is located in southwestern Uganda, bordered by the Democratic Republic of the
140 Congo to the west and by protected park lands to the north (Kigezi Game Reserve) and south
141 (Bwindi Impenetrable National Park). In addition to subsistence agricultural activities, there is
142 considerable influence from the tourism industry in the region. Industrial tea and coffee production
143 also play a role in local livelihoods. The adult literacy rate in the region is comparable (71.9%)

144 (Kanungu District Local Government 2013) to the national rate (73%) (UNICEF 2014). The region
145 experiences bimodal seasonality, with rainy seasons from October to December and again from
146 March to May, and low average temperatures relative to the rest of Uganda (typically below 20°C)
147 (McSweeney, New and Lizcano 2010). Climate change projections indicate that the region will likely
148 experience increases in annual mean temperature and frequency of heavy rain events (Anyah and
149 Qiu 2012, Christensen et al. 2013).

150 [INSERT FIGURE 1 HERE]

151 **Figure 1. Map of Kanungu District**

152 The Bakiga ethnic group, a traditionally agrarian society, make up the majority of the
153 population of Kanungu District. One percent of the population (approximately 800 inhabitants) are
154 members of the Indigenous Batwa ethnic group. The Batwa [Anonymous 2012] and Bakiga
155 [Anonymous 2015 A] have both been identified as being highly vulnerable to the health impacts of
156 climate change. Perinatal health indicators for the region are also below national averages—only
157 about 40% of births occur in health facilities (Uganda Bureau of Statistics (UBOS) and ICF
158 International Inc 2012) as opposed to 57% nation-wide (UNICEF 2014). Though 59% of Ugandan
159 infants are delivered by a skilled provider, skilled providers are only present at 42% of births in the
160 Southwest Region (Uganda Bureau of Statistics (UBOS) and ICF International Inc 2012). In a
161 sample of newborns delivered at Bwindi Community hospital, the prevalence of low birth weight
162 births (<2500g) was 7.2% and the prevalence of preterm births was 8.1% [Anonymous Under
163 review A].

164 The Batwa ethnic group are a subgroup of the Central African Pygmy population and the
165 Indigenous residents of the Bwindi Impenetrable Forest [Anonymous 2012]. The Batwa have faced
166 historic oppression and marginalization, not unlike other Indigenous populations worldwide, and
167 they face a greater burden of climate change vulnerability [Anonymous 2012, Anonymous 2016 B,

168 Anonymous Under review B]. Evicted from their homes when conservationists created the National
 169 Park in the early 1990s, the Batwa were forcibly resettled in agrarian communities despite their
 170 history as traditional hunter-gatherers (Ohenjo et al. 2006, Jackson 2006) [Anonymous 2012].

171 There are deep health disparities between the Batwa and both the Ugandan population at
 172 large and the neighbouring Bakiga ethnic group (which has historically existed in settled agriculture-
 173 based communities), despite the fact that both groups live in the same biophysical environment with
 174 identical health services [Anonymous 2012]. The Batwa have a higher prevalence of malaria (9.4%
 175 compared to 4.5% in the Bakiga population [Anonymous 2016 A]) and acute gastrointestinal illness
 176 (compared to East Africa [Anonymous 2015 B]). The Batwa also face extreme food insecurity
 177 [Anonymous Under review B]. The prevalence of HIV among the Batwa population is lower than
 178 that in the Bakiga population (Birungi 2010). The two ethnic groups have some interaction, primarily
 179 when Batwa women perform wage labour for Bakiga farmers.

180 **Table 1. Indicators of socioeconomic status among Batwa and Bakiga communities.**
 181 **(Adapted from [Anonymous Under review A])**

Measure (variable descriptor)	Batwa (proportion of the population)	Bakiga (proportion of the sample)	Source
Moderate acute malnutrition among adult women (classified as moderately malnourished according to Uganda Ministry of Health Integrated Management of Acute Malnutrition Guidelines)	45.86%	0.42%	[Anonymous 2017]
Household mosquito net use (did not have nets)	70.99%	53.56%	[Anonymous 2016 A] (by request)
Assets (did not have any assets)	62.12%	19.01%	[Anonymous 2016 A] (by request)
Access to handwashing facilities (did not have access to handwashing)	73.85%	56.40%	[Anonymous 2016 A] (by request)
Access to soap (did not have access to soap)**	75.38%	62.06%	[Anonymous 2016 A] (by request)

182 **Only asked of people that had access to hand washing facility, for example for the Batwa, 32 or
 183 94% of the households that had access to handwashing had access to soap

184 In 2003, the Bwindi Community Hospital (BCH) was established by American medical
185 missionaries as an outreach clinic for the Batwa. It has since expanded to an inpatient hospital with a
186 catchment area population of 100 000 (BCH n.d.). BCH's antenatal clinic sees approximately 250
187 mothers per month and over 1000 deliveries are performed at the hospital annually (BCH 2014).
188 The hospital also operates a Waiting Mothers Hostel, where women who live far away can stay
189 during the weeks leading up to their deliveries (BCH 2009), and Village Health Teams from the
190 hospital provide outreach to both Indigenous and non-Indigenous communities through the region
191 on a monthly basis (Haven Nahabwe, Public Health Officer, Bwindi Community Hospital, 2015,
192 conversation). All of the healthcare workers at the hospital are non-Indigenous.

193 **2.3 Indigeneity and birth weight in Kanungu**

194 Defining Indigeneity remains a global and domestic challenge, and there is no universal consensus
195 on its definition and criteria (Stephens et al. 2006). The question of what it means to be Indigenous
196 in Africa is particularly contentious, and while there is little formal recognition of Indigenous
197 peoples by national governments, it is estimated that there are 14.2 million self-identifying
198 Indigenous people in Africa (Ohenjo et al. 2006). The Pygmy peoples of Central Africa, of which
199 the Batwa are a subgroup, number an estimated 920 000, a small proportion of whom live in Uganda
200 (Olivero et al. 2016). Though Indigeneity is often presumed to be a function of being tied to specific
201 geography, self-identification and the relationship of the group to a dominant state body have
202 become more salient qualifiers of Indigeneity (Maybury-Lewis 2002). In the case of the Batwa, their
203 historic marginalization and discrimination relative to the non-Indigenous ethnic majority have
204 translated into existing and persistent health and socioeconomic inequities (Table 1). Despite
205 contested definitions, the Batwa history (Jackson 2003) and current experiences [Anonymous 2012]
206 are consistent with most constructs of Indigeneity, including long-standing reliance on existing lands
207 and resources (Stephens et al. 2006, Shaw, Herman and Dobbs 2006), dispossession from traditional

208 resources, local stigmatization as ‘other’ compared to neighbouring populations, ongoing
209 discrimination and inequity based on their ethnicity (Maybury-Lewis 2002), and importantly, self-
210 identification as ‘Indigenous’ (Martinez Cobo 1981).

211 We sought to characterize how the pathways identified in existing literature linking weather
212 to birth outcomes were applicable to, manifested within, and differed between Indigenous Batwa
213 and non-Indigenous neighbouring Bakiga. A quantitative relationship between meteorological
214 factors and birth outcomes has already been established among Batwa and Bakiga, finding that there
215 is not only a difference in *magnitude* of the effect of weather on birth weight by ethnicity (effect
216 modification), but that the high-risk period of gestation and the meteorological variables of
217 significance differ as well [Anonymous Under review A]. This prior work revealed a significant
218 relationship between exposure to more days of precipitation and higher average temperatures in the
219 third trimester and birth weight. This relationship was different for the two ethnic groups: in the
220 non-Indigenous population, only exposures in the third trimester were associated with birth weight,
221 while the birth weights among infants from the Indigenous Batwa ethnic group were associated with
222 average temperature exposure throughout pregnancy. In this paper, we thus sought to qualitatively
223 characterize the pathways underpinning these previously established relationships, herein conducting
224 a qualitative, community-based study with Batwa and Bakiga mothers and health care workers in the
225 region.

226 **2.4 Data collection**

227 The research was guided by a community-based participatory research approach (Wallerstein and
228 Duran 2006, O’Fallon and Dearry 2002), building on existing partnerships with both Batwa and
229 Bakiga communities, the Bwindi Community Hospital, and the Batwa Development Program. The
230 research is underpinned by a phenomenological philosophy that is used to understand the
231 “common or shared experiences of a phenomenon” of several individuals (Creswell 2013, 81). We

232 sought to understand the individual and group perspectives of how mothers experienced the effects
233 of weather and season during their pregnancies. The study was conducted from June-August 2015 in
234 Kanungu District, Uganda. Qualitative data collection took two forms: focus group discussions
235 (FGDs) in both Batwa and Bakiga communities (n=16 FGDs, 8 Batwa and 8 Bakiga), and key
236 informant interviews (KIIs) with community members and hospital employees (8 KIIs, total n=10
237 individuals).

238 We sought permission from the village chairperson in advance of each focus group and held
239 each gathering at a designated communal gathering area within each settlement. Each group
240 consisted of five female participants, selected from all village women available on the day of the
241 focus group. A diverse range of ages was selected for each discussion to invite historical perspectives
242 from elder community members. Focus groups included both primagravidae and multigravidae
243 mothers. Discussions were conducted in Rukiga, the local language, with translation through two
244 local research assistants known to the communities. One research assistant facilitated the discussions
245 and provided translation after each response, while the other assisted with the logistics of gathering
246 the women and provided simultaneous translation as needed (Esposito 2001). This early stage
247 translation approach (Santos, Black and Sandelowski 2015) offered flexibility to English-speaking
248 researchers to redirect lines of questioning as appropriate and to better gauge participant engagement
249 (MacKenzie 2016, Esposito 2001). The discussion guide was designed to elicit storytelling and open-
250 ended responses, and was structured around three topic areas: personal background, pregnancy
251 experiences and childbirth experiences. Questions around environmental exposures during
252 pregnancy were open-ended, but included specific questions around diet, physical work, and
253 patterns of illness based on the three primary exposure pathways we had observed in the literature.
254 The total recorded focus group discussion time was 661 minutes and the average length per

255 discussion was 41 minutes. All participants were remunerated in a manner determined in
256 consultation with local partners (with community lunches or gifts of soap for individual mothers).

257 Semi-structured interviews were conducted with key informants (Brown and Durrheim 2009,
258 Fylan 2005) according to the interview guide included in our Supplemental Materials. All informants
259 were given the choice of language of interview; with the exception of one key informant, all
260 interviews were conducted in English (the official common language of Uganda). All key informants
261 were staff affiliated with the Bwindi Community Hospital except for one community informant. All
262 participants received a modest token of appreciation (e.g., pen, key chain) but no monetary
263 compensation from the researchers, as per hospital policy. The total recorded interview time with
264 key informants was 344 minutes, with an average interview length of 43 minutes.

265 The research team conducting focus groups included two outside female researchers
266 ([removed for blind review]) with unmistakable identifiers of privilege (both being white North
267 American researchers). The history of unethical research conducted on Indigenous and other
268 vulnerable populations (Smith 1999) was a constant consideration as we tried to minimize power
269 imbalances and create an environment where women felt safe discussing personal experiences of
270 pregnancy and childbirth. Confidentiality was stressed in each interview and focus group facilitators
271 explained that there were no ‘right’ or ‘wrong’ answers. Despite measures taken to ensure that
272 participants felt safe and comfortable sharing information with researchers, participants were
273 sometimes reluctant to divulge information about the things they perceived as frowned upon by
274 healthcare workers (e.g., use of traditional medicines). We acknowledge the unequal power
275 relationships that persist in spite of our efforts to balance power differentials in focus group and
276 interview settings (Nunkoosing 2005).

277 We obtained ethics approval for this research protocol from the [institution names removed
278 for blinded review] University Research Ethics Board, as well as from the Bwindi Community

279 Hospital administration. Informed consent was obtained prior to all research activities, and
280 participants were informed that they could end the interview at any time. To protect confidentiality
281 outside of the group and beyond the focus group, no demographic information was sought from
282 focus group participants apart from the number of children they had and the number of pregnancies
283 they had experienced. Key informants were offered the choice of being named or anonymous in
284 research notes and any subsequent publications.

285 **2.5 Data analysis**

286 All interviews were audio recorded with consent from participants. The recordings were
287 subsequently transcribed and verified for accuracy 1-2 times prior to coding. The recordings of all
288 focus groups conducted in Rukiga were translated by a translator who was present at all focus
289 groups but was not the discussion facilitator. This meant that the research team had a second
290 opportunity to verify the translations that were recorded in the initial discussions, allowing us to seek
291 clarification on any terms or cultural concepts that were unfamiliar.

292 As with data collection, data analysis was guided by a phenomenological approach. In this
293 study, the phenomenon of interest was how women experienced seasonal and weather exposures
294 during pregnancy and the health effects they attributed to these exposures. Throughout the data
295 collection process, memoing (Birks, Chapman and Francis 2008) was used to capture our initial
296 impressions during the interview and focus group processes. Through memoing, we generated a list
297 of significant statements that were later used to guide the coding process (Creswell 2013). We drew
298 on the dominant pathways identified in the systematic literature review to guide *a priori* deductive
299 coding (Crabtree and Miller 1999) while also examining transcripts for any *a posteriori* inductive
300 themes that could emerge (Pope, Ziebland and Mays 2000, Fereday and Muir-Cochrane 2006). The
301 framework of the interview guides allowed researchers to hone in on topics relevant to the analysis

302 of pathways through which seasonal or weather exposures may be affecting perinatal health, and to
303 code any mentions of concepts related to maternal nutrition, physical labour, or infectious disease.

304

305 **3 Results**

306 **3.1 Lived experience of pregnancy and seasonality**

307 A nine-month gestation means that all mothers experience the effects of both rainy and dry seasons
308 at various points in their pregnancies. Mothers and key informants reported that they observed
309 seasonal differences in the experiences of pregnancy. Respondents identified seasonal variation in
310 food supply and strenuous labour, as well as seasonal patterns in illness, as the primary ways they
311 were affected by season or weather during pregnancy. Focus group participants connected these
312 factors to the health of their babies: stated ways to have big healthy babies included “*not working so*
313 *hard,*” “*having peace,*” and “*not becoming sick, and even eating well*” (Batwa FGD 7B). The dry season was
314 identified as the period of greatest food scarcity, though many also reported environmental
315 challenges during the rainy season as well. Many women highlighted maternal nutrition as the most
316 important factor affecting the health of their infants, and associated lack of food with the most
317 severe outcomes, stating: “*we don’t have those foods [in the dry season], that’s why we produce smaller babies,*”
318 (Batwa FGD 6B) and “*that’s the reason why our babies die in our stomach, because we don’t have food*” (Batwa
319 FGD 7A). These statements reinforce prior research demonstrating high levels of food insecurity in
320 the region, particularly among the Batwa [Anonymous Under review B].

321 Few FGD participants perceived a difference in the health and size of their babies based on
322 the season of their birth. One focus group participant stated: “*dry season, or rainy season, the babies are all*
323 *the same*” (Batwa FGD 2A). Individual mothers did not associate seasonal effects on birth outcomes
324 when considered in the broader context of their livelihoods and other health and prenatal stressors.
325 In contrast, a local traditional birth attendant working in the area for over forty years, who has

326 experienced the birth of a large number of babies, reported that “*the baby of the rainy season is always*
327 *big. It’s because of eating well*” (KII 6).

328 Both mothers and hospital key informants emphasized access to health services as one of
329 the predominant challenges faced by pregnant women in the region. Respondents expressed that it
330 was difficult for mothers to leave seasonal work opportunities that sustained their families and to
331 leave other children at home to attend antenatal care or to come to the hospital for delivery.
332 Transport to health services was a critical barrier, both financially in times when paid labour was
333 scarce, and physically in the rainy season when road conditions worsened:

334 *And also it would become the dry season, where you find most of the jobs... these mothers... they*
335 *have no money... most of them go work for others, to get some little money, get some food... So [in*
336 *the] dry season, where are most of the [mothers coming to the hospital]? (KII 4)*

337 However, especially when it came to their desire for what they considered to be a “*good*” or “*safer*”
338 (*Batwa FGD 2B*) delivery at a health care facility, many women shared an attitude of resilience,
339 explaining that you find a way to the hospital however you can (e.g., by taxi, motorcycle, or walking
340 up to ten hours) because delivering at home is too great a risk to take—as one participant described,
341 if she cannot make it to the hospital, “*I have to deliver [from home] and wait to die.*” (*Batwa FGD 1B*).

342 **3.2 Reported pathways between meteorological conditions and birth outcomes**

343 **3.2.1 Maternal nutrition**

344 Eating well was identified in nearly all focus groups (n=15) as a crucial determinant of a healthy
345 pregnancy and healthy baby. Mothers also identified this as a factor that varied seasonally, with all
346 but one focus group confirming that there was less food available in the dry season:

347 *In the rainy season, you find you go to dig somewhere and you get food. You eat, and the baby*
348 *inside is receiving that food. And you find that you even got some vitamins for the baby. But in the*

349 *dry season, you find you have no energy, you stay home, you have not gone to look for food, even the*
350 *baby inside will not have energy. (Batwa FGD 6A)*

351 Key informants also reported the dry season as the period of greater food scarcity. A traditional
352 birth attendant explained, *“In the rainy season, that is when the food is available, like greens and some other*
353 *foods. So you find the woman is healthy in the rainy season” (KII 6)*. She explained the effects of this lack of
354 healthy food on the babies of malnourished mothers, stating that *“the babies [born in] the rainy season*
355 *are not the same as the dry season. The baby of the rainy season is produced healthy and the one of the dry season is*
356 *not healthy” (KII 6)*. Cost of food increases in the dry season as a result of scarcity: *“in the dry season,*
357 *their food is expensive... in the rainy season it’s in the harvesting time, and their food is available. But in the dry*
358 *season, there is no food” (Bakiga FGD 8A)*. Quality of food available in the dry season was also of
359 concern: *“in dry season, the greens are very few,”* stated one focus group participant (Bakiga FGD 4B).
360 One key informant who had expressed concern at how the seasonal trends in food insecurity
361 affected mothers also highlighted that changes in weather patterns have begun to affect agriculture
362 in the region:

363 *Sometimes, you may expect rainfall in a certain season... you may find... it’s time for rainy*
364 *season, winter, but [the sun] is shining heavily, every day. When it should be dry season, it’s*
365 *raining. So maybe you go and plant crops expecting rain and then the rain ceases for a month,*
366 *then you make a loss, so this is difficult. Or, you have planted crops, and because you get a lot of*
367 *rainfall, then they can’t grow, or it’s erosion, you find all of the crops planted, they are swept off by*
368 *moving water. (KII 9)*

369 **3.2.2 Physical labour**

370 Energy expended through physical work was described by participants as being season-dependent.
371 Participants described physical labour as being stratified across different livelihood activities. Rainy
372 season work included weeding, planting, mulching, and digging in the gardens. These activities were

373 described as being more difficult than most dry season work: harvesting, preparing the gardens,
374 drying crops, and weaving mats and baskets. Though digging was one of the predominant activities
375 in both seasons, participants stated that they spent longer hours digging in the rainy season than in
376 the dry. As one key informant described it:

377 *We usually have a lot of work in the rainy season because most of the activities are being done,*
378 *compared to the dry season; in the dry season, there is harvesting of some crops and preparation of*
379 *crops...when it rains, you have to prepare and wait for rain... then you start growing crops,*
380 *planting crops. (KII 9)*

381 Several women stated that being pregnant in the dry season was advantageous because it meant they
382 would have already planted their crops and would not need to go back to the gardens to dig. The
383 heat and hardness of the soil in the dry season were listed as intensifying strenuous work during the
384 dry periods. Women were aware of the potential for adverse outcomes due to overexerting
385 themselves with physical work during pregnancy: “[there are] some challenges, like digging, and you dig a lot
386 and you find you have some abdominal pain and you even can end up getting some abortion or miscarriages” (Batwa
387 FGD 7B). However, their experiences of physical work during pregnancy varied. Some women
388 described working up until the time of delivery (even to the extent of needing to deliver in the
389 garden), while others stopped working as early as the first trimester if they were feeling ill.

390 The ability to choose not to work was not available to all mothers. Some were motivated to
391 work by cravings for more expensive foods (“*When my heart wants meat, I have to go and dig for the money,*
392 *then I buy meat.*” – Batwa FGD 2B). Others worked to ensure the immediate survival of their families.

393 Stories from key informants highlighted the high levels of poverty among the population:

394 *Mothers work a lot in their pregnancy. Not only to earn money to attend the facility, but also*
395 *because they are the basic unit of the household's survival, so, to some mothers, it is the least of*
396 *their worries: money that will take them to the facility. Because they are still at day-to-day*

397 *survival. That is shown in how those mothers who present to us... She presents with mud on the*
398 *feet, meaning that labour started when she was in the field. ... (KII 1)*

399 **3.2.3 Seasonal illness**

400 Mothers told us that illness strikes in all seasons, but that there were seasonal differences in the types
401 of illnesses prevalent in the area. Focus group participants stated that in the rainy season “*there is a lot*
402 *of malaria and coldness, compared to the dry season*” (Batwa FGD 7A). Women were acutely aware of the
403 dangers of contracting malaria during pregnancy, associating it with pregnancies ending in
404 miscarriage. Mothers were well informed of prophylactic interventions providing protection from
405 malaria during pregnancy and many reported use of malarial prophylactics in prior pregnancies. In
406 the dry season, focus group participants identified hunger and fatigue as the primary threats to their
407 wellbeing. Feeling a lack of energy in the dry season was expressed by many mothers and confirmed
408 by a traditional birth attendant key informant, who stated “*In the rainy season they [pregnant mothers] have*
409 *energy, but in the dry season they have no energy*” (KII 6).

410 Maternal nutritional status plays an important role in seasonal variation in anemia, a
411 phenomenon highlighted by both participants and key informants alike. A participant stated: “*during*
412 *the dry season, we find we have no energy and there is a lot of sweating. We find there are no greens for eating. When*
413 *we go to the hospital, we are told that we are lacking blood in our bodies*” (Batwa FGD 1B). One key informant
414 expressed that cases of severe anemia in pregnancy were somewhat rare, since pregnant women
415 would prioritize their own nutritional needs and sell assets if needed to safeguard the health of their
416 foetus. However, the risks remain high, as several stated that the reason a woman might miscarry is
417 because she lacks the good foods (especially green leafy vegetables) that will help prevent anemia.
418 One key informant linked anemia to meteorological factors:

419 ...in the past, like half a year, we had a very dry period from December through January, up to...
420 end of March, that's when a bit of rain started coming. And in that period, we had two pregnant
421 mothers coming in with very severe anemia. (KII 1)

422 Mothers expressed concern at the effects of anemia on their infants: “The way you are in the dry season,
423 that is how the baby will be when it is produced. When you are low on energy in the dry season, the baby produced,
424 will also have low energy and low strength.” (Batwa FGD 1A).

425 **3.3 Comparing experiences of Batwa and Bakiga mothers**

426 When asked if there were differences between the challenges faced by Batwa and Bakiga mothers, a
427 hospital administrator key informant stated, “the challenges faced by Batwa and Bakiga mothers, they're the
428 same challenges. They face the same challenges. They live in the same environment... so they face the same issues” (KII
429 10). Batwa and Bakiga mothers reported similar experiences of seasonal and weather exposures
430 during pregnancy, however, the extent or *magnitude* to which these exposures influence their health
431 differed. Bakiga mothers spoke of selling off livestock to pay for transport to access health services
432 but no Batwa women mentioned such an option. Batwa mothers consistently reported that they
433 were unable to earn enough money for transport, and would deliver from home. Bakiga women
434 were more likely to report stopping physical work when such work became untenable with their
435 pregnancy. Batwa mothers rarely reporting reducing physical work. One Batwa mother stated: “If I
436 sit at home, who will go and dig for me? So I have to go” (Batwa FGD 6B). A key informant who has worked
437 among the Batwa for five years described the differences in the challenges facing Batwa and Bakiga
438 mothers:

439 All those challenges... for the Batwa, it's much, much, much, much worse. ...it's very hard, it's
440 cost of food... they don't have food at all. I may call it severe. (KII 9)

441 Last year we lost one Mutwa [Batwa singular], and their total population is fewer than a thousand. And for
442 women of reproductive age... when you lose one woman out of 200, you feel it. (KII 2)

443 Older Batwa mothers recalled the foods they would eat in the forest prior to eviction,
444 expressing frustration with being restricted from accessing the forest herbs that were used for
445 medicinal purposes. Some Batwa women stated that they had larger babies when living in the forest:
446 “Because we used to eat honey, and used to get fish from the water, that’s the reason why we used to produce healthy
447 babies” (Batwa FGD 6B). Despite these losses following their eviction from the forest, many Batwa
448 women praised the positive impact the hospital has had on their health during pregnancy,
449 mentioning in particular the availability of antenatal care that allows them to monitor the health of
450 their babies throughout pregnancy. Some of the older Batwa mothers reported that rates of infant
451 mortality were higher when they lived in the forest due to harsher living conditions and lack of
452 formal healthcare (Batwa FGD 1A).

453

454 **4 Discussion**

455 Prentice et al. (1987) were among the first to examine the effects of seasonality on birth weight and
456 posited that this relationship manifested through malaria morbidity, food shortages, and hard
457 physical work experienced during pregnancy. These same drivers were identified and validated by
458 the mothers and key informants we spoke to in Uganda.

459 Mothers generally stated that the babies were the same regardless of their season of birth but
460 expressed differing opinions as to whether they had found pregnancy more challenging in the rainy
461 season or in the dry season. This finding may indicate that there is not a clearly defined risk period,
462 but rather that pathways of risk differ under different seasonal conditions. Key informants offered
463 breadth of experience balancing the depth of individual perspectives, with many describing the
464 increased availability of food in the rainy season. This finding is consistent with our other work
465 [Anonymous Under review A], which found that exposure to more days of precipitation in the third
466 trimester was associated with increased birth weight (3.1g increase in birth weight per additional day

467 of exposure). Several prior studies from other parts of sub-Saharan Africa also identified the dry
468 season as a period in which food shortages and/or increases in physical labour occurred and lower
469 birth weights were recorded (Neufeld, Pelletier and Haas 1999, Onyiriuka 2006, Enquoselassie 2000,
470 Friis et al. 2004). Mothers expressed concern about sickness during both seasons, but malaria and
471 cold-related illness were of greater concern during the rainy season. Difficulties in accessing the
472 hospital and antenatal care services during the rainy season were mentioned by several mothers and
473 key informants. These access issues appeared to be driven by a lack of resources to pay for transport
474 to service providers and being unable to sacrifice time off work or to find child care to attend
475 appointments. The difference in the nature of the concerns by season may indicate that mothers
476 experience the effects of weather through different pathways in different seasons, with implications
477 for seasonally-sensitive interventions based on a woman's gestational stage.

478 Interviews and focus groups revealed that while both Batwa and Bakiga mothers experienced
479 the effects of weather and seasons via the same pathways, the extent of their vulnerability differed.
480 Batwa mothers had fewer assets to sell to pay for transport to the hospital, and did not have the
481 same level of flexibility in choosing when to stop doing strenuous work during their pregnancies.
482 Other prior research in the region has shown that Bakiga are considerably better equipped to cope
483 with shocks threatening food security, often by selling off livestock [Anonymous 2016 B]. Existing
484 disparities in social determinants of health between Batwa and Bakiga mothers appear to leave
485 Batwa mothers more vulnerable to the effects of weather and seasonal variation, magnifying the
486 adverse effects of these exposures on their pregnancies and on the health of their newborns.

487 This study sought to understand the pathways through which weather and season may affect
488 birth outcomes in low-resource settings. It is these pathways that might be modifiable to help
489 improve birth outcomes. Studies conducted in more developed settings where food security is a less
490 acute concern, where physical work may not vary seasonally, and where seasonal infectious diseases

491 are less common suggest different pathways through which meteorological factors may affect birth
492 weight (e.g. vitamin D exposure) (McGrath, Burne and Eyles 2005), and highlight interaction effects
493 with air pollution (Beltran et al. 2014, Chodick et al. 2009, Laaidi et al. 2011)). These pathways may
494 also be at work in Kanungu District, but the magnitude of their effects is likely to be marginal when
495 compared to the dominant pathways proposed in this paper (Strand et al. 2011). Consideration of
496 black carbon and other air pollutants—for which data were not available for this study—would also
497 be appropriate in this context given the extensive use of biofuels such as wood for cooking.

498 In the 5th Assessment Report of the Intergovernmental Panel on Climate Change, Smith et
499 al. (2014, 741) note that “[g]iven the increase globally in life expectancies, many babies born this
500 decade will be alive at the end of the century, and will be personally affected by the climate that is in
501 place in 2100”. However, climate change will begin affecting these babies far sooner than 2100—
502 unpredictable and intensified effects of season and weather will begin to affect their health in utero
503 (Rylander, Odland and Sandanger 2013, Grace et al. 2015). As climate change intensifies, the effects
504 of season and weather on birth outcomes will likely be amplified (Grace et al. 2015, Rylander et al.
505 2013). Understanding the new dimension climate change adds to existing disparities in perinatal
506 health should help illuminate interventions aimed at eliminating these inequities. Understanding the
507 context-dependent pathways means that we have intervention points around the prevention of
508 predictable food shortages, protecting women from excess physical exertion and ensuing best
509 practices for infectious disease prevention. The health effects of climate change and maternal and
510 newborn health disparities will be magnified by existing social gradients. Interventions supporting
511 those at the lower end of these social gradients have the potential for double benefit by addressing
512 two grand challenges in global health: maternal/child health and climate change adaptation.

513 Our findings suggest focused adaptation strategies targeting the pathways through which
514 mothers and developing foetuses are exposed to the effects of weather and season: maternal energy

515 balance and risk of seasonal illness. These pathways are relatively consistent across and within
516 populations, meaning they could be entry points to interventions in other subsistence agriculture-
517 based contexts. However, further place-based qualitative inquiry characterizing the nature and extent
518 of these pathways is warranted, particularly in communities where alternative livelihoods
519 predominate. These findings are of use to collaborators at Bwindi Community Hospital and the
520 Ugandan Ministry of Health as they develop strategies specific to the regional needs of the women
521 of Kanungu District. BCH has made it a priority in their strategic plan to reduce maternal and child
522 mortality by 25% by 2019, and the results of this study suggest placing a particular emphasis on
523 supporting Batwa mothers as a vulnerable population group to reach this goal. Hospital and ministry
524 planners may also take into consideration the need for more nutritional interventions in the dry
525 season and continued education around climate change and agriculture, as well as education
526 initiatives geared towards other income-generating activities. The need for better access to transport
527 in the rainy season (Caulfield et al. 2016) and opportunities to leverage the knowledge and access of
528 traditional birth attendants might also be considered in policy development (Rishworth et al. 2016,
529 Sarker et al. 2016). Promotion of family planning to time births according to the most optimal
530 seasonal conditions for pregnancy might also be also be considered as an initiative to enhance
531 perinatal health in the region.

532

533

534

535

536

537

538 **References**

- 539 [Anonymous 2012] Details omitted for double-blind reviewing.
540 [Anonymous 2015 A] Details omitted for double-blind reviewing.
541 [Anonymous 2015 B] Details omitted for double-blind reviewing.
542 [Anonymous 2016 A] Details omitted for double-blind reviewing.
543 [Anonymous 2016 B] Details omitted for double-blind reviewing.
544 [Anonymous 2017] Details omitted for double-blind reviewing.
545 [Anonymous Under review A] Details omitted for double-blind reviewing.
546 [Anonymous Under review B] Details omitted for double-blind reviewing.
547
548 Adger, W. N. (2006) Vulnerability. *Global environmental change*, 16, 268-281.
549 Anyah, R. O. & W. Qiu (2012) Characteristic 20th and 21st century precipitation and temperature
550 patterns and changes over the Greater Horn of Africa. *International Journal of Climatology*, 32,
551 347-363.
552 Barker, D., J. Eriksson, T. Forsén & C. Osmond (2002) Fetal origins of adult disease: strength of
553 effects and biological basis. *International Journal of Epidemiology*, 31, 1235-1239.
554 BCH, B. C. H. (2009) Safe deliveries in Bwindi. *Bwindi Community Hospital Newsletter*, 2, 2.
555 --- (2014) HMIS Annual Report 2013-2014 (unpublished data).
556 --- (n.d.) Our History.
557 Beltran, A. J., J. Wu & O. Laurent (2014) Associations of meteorology with adverse pregnancy
558 outcomes: A systematic review of preeclampsia, preterm birth and birth weight. *International*
559 *Journal of Environmental Research and Public Health*, 11, 91-172.
560 Berry, H. L., J. R. A. Butler, C. P. Burgess, U. G. King, K. Tsey, Y. L. Cadet-James, C. W. Rigby &
561 B. Raphael (2010) Mind, body, spirit: co-benefits for mental health from climate change
562 adaptation and caring for country in remote Aboriginal Australian communities. *New South*
563 *Wales Public Health Bulletin*, 21, 139-145.
564 Birks, M., Y. Chapman & K. Francis (2008) Memoing in qualitative research: Probing data and
565 processes. *Journal of Research in Nursing*, 13, 68-75.
566 Birungi, M. R. E. 2010. A comparative study of HIV risk factors and health care utilization between
567 the Batwa pygmies and surrounding communities of Kanungu District, South West Uganda.
568 Bwindi, Uganda: Bwindi Community Hospital.
569 Botero, D. & F. Lifshitz (1999) Intrauterine growth retardation and long-term effects on growth.
570 *Curr Opin Pediatr*, 11, 340-7.
571 Brighton, A., R. D'Arcy, S. Kirtley & S. Kennedy (2013) Perceptions of prenatal and obstetric care in
572 Sub-Saharan Africa. *International Journal of Gynecology & Obstetrics*, 120, 224-227.
573 Brown, L. & K. Durrheim (2009) Different Kinds of Knowing: Generating Qualitative Data
574 Through Mobile Interviewing. *Qualitative Inquiry*.
575 Busby, J. W., T. G. Smith, K. L. White & S. M. Strange (2013) Climate change and insecurity:
576 mapping vulnerability in Africa. *International Security*, 37, 132-172.
577 Carolan-Olah, M. & D. Frankowska (2014) High environmental temperature and preterm birth: A
578 review of the evidence. *Midwifery*, 30, 50-59.
579 Caulfield, T., P. Onyo, A. Byrne, J. Nduba, J. Nyagero, A. Morgan & M. Kermode (2016) Factors
580 influencing place of delivery for pastoralist women in Kenya: a qualitative study. *BMC*
581 *Women's Health*, 16, 52.
582 Chodick, G., S. Flash, Y. Deoitch & V. Shalev (2009) Seasonality in birth weight: Review of global
583 patterns and potential causes. *Human Biology*, 81, 463-477.

584 Chou, W.-C., Y.-T. Chung, H.-Y. Chen, C.-J. Wang, T.-H. Ying, C.-Y. Chuang, Y.-C. Tseng & S.-L.
585 Wang (2014) Maternal Arsenic Exposure and DNA Damage Biomarkers, and the
586 Associations with Birth Outcomes in a General Population from Taiwan. *PLoS ONE*, 9,
587 e86398.

588 Christensen, J. H., K. Krishna Kumar, E. Aldrian, S.-I. An, I. F. A. Cavalcanti, M. de Castro, W.
589 Dong, A. Goswami, A. Hall, J. K. Kanyanga, A. Kitoh, J. Kossin, N.-C. Lau, J. Renwick, D.
590 B. Stephenson, S. P. Xie & T. Zhou. 2013. Climate Phenomena and their Relevance for
591 Future Regional Climate Change. In *Climate Change 2013: The Physical Science Basis. Contribution*
592 *of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*,
593 eds. T. F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y.
594 Xia, V. Bex & P. M. Midgley. Cambridge, UK and New York, NY, USA.

595 Costello, A., M. Abbas, A. Allen, S. Ball, S. Bell, R. Bellamy, S. Friel, N. Groce, A. Johnson, M. Kett,
596 M. Lee, C. Levy, M. Maslin, D. McCoy, B. McGuire, H. Montgomery, D. Napier, C. Pagel, J.
597 Patel, J. A. de Oliveira, N. Redclift, H. Rees, D. Rogger, J. Scott, J. Stephenson, J. Twigg, J.
598 Wolff & C. Patterson (2009) Managing the health effects of climate change: Lancet and
599 University College London Institute for Global Health Commission. *Lancet*, 373, 1693-733.

600 Crabtree, B. F. & W. L. Miller. 1999. *Doing Qualitative Research*. SAGE Publications.

601 Creswell, J. W. 2013. *Qualitative inquiry and research design: Choosing among five approaches*. Thousand
602 Oaks, USA: SAGE Publications.

603 Cutchin, M. P. (2007) The need for the “new health geography” in epidemiologic studies of
604 environment and health. *Health & Place*, 13, 725-742.

605 Enquoselassie, F., Minyilshewa, A. (2000) Changes in birthweight-weight of Hospital-delivered
606 neonates in Addis Ababa. *Ethiopian Journal of health development*, 14, 169-176.

607 Esposito, N. (2001) From Meaning to Meaning: The Influence of Translation Techniques on Non-
608 English Focus Group Research. *Qualitative Health Research*, 11, 568-579.

609 Fereday, J. & E. Muir-Cochrane (2006) Demonstrating Rigor Using Thematic Analysis: A Hybrid
610 Approach of Inductive and Deductive Coding and Theme Development. *International Journal*
611 *of Qualitative Methods*, 5, 80-92.

612 Few, R. (2007) Health and climatic hazards: Framing social research on vulnerability, response and
613 adaptation. *Global Environmental Change*, 17, 281-295.

614 Fletcher, A. J. (2017) Applying critical realism in qualitative research: methodology meets method.
615 *International Journal of Social Research Methodology*, 20, 181-194.

616 Ford, J. D. (2012) Indigenous Health and Climate Change. *American Journal of Public Health*, 102,
617 1260-1266.

618 Friis, H., E. Gomo, N. Nyazema, P. Ndhlovu, H. Krarup, P. Kæstel & K. F. Michaelsen (2004)
619 Maternal body composition, HIV infection and other predictors of gestation length and
620 birth size in Zimbabwe. *British Journal of Nutrition*, 92, 833-840.

621 Fylan, F. (2005) Semi structured interviewing. *A handbook of research methods for clinical and health*
622 *psychology*, 65-78.

623 Grace, K., F. Davenport, H. Hanson, C. Funk & S. Shukla (2015) Linking climate change and health
624 outcomes: Examining the relationship between temperature, precipitation and birth weight
625 in Africa. *Global Environmental Change*, 35, 125-137.

626 Gracey, M. & M. King (2009) Indigenous health part 1: determinants and disease patterns. *The*
627 *Lancet*, 374, 65-75.

628 Graham, S., L. R. Pulver, Y. A. Wang, P. M. Kelly, P. J. Laws, N. Grayson & E. A. Sullivan (2007)
629 The urban-remote divide for Indigenous perinatal outcomes. *Med J Aust*, 186, 509-12.

630 Green, D. & L. Minchin (2014) Living on Climate-Changed Country: Indigenous Health, Well-Being
631 and Climate Change in Remote Australian Communities. *EcoHealth*, 11, 263-272.

632 Haines, A. & J. A. Patz (2004) HEalth effects of climate change. *JAMA*, 291, 99-103.

633 Harding, J. (2001) The nutritional basis of the fetal origins of adult disease. *International Journal of*
634 *Epidemiology*, 30, 15-23.

635 Jackson, D. 2003. Twa Women, Twa Rights in the Great Lakes Region of Africa. ed. M. R. G.
636 International. London, UK.

637 ---. 2006. The health situation of women and children in central African Pygmy peoples. ed. T. F. P.
638 Programme. Moreton-in-Marsh, UK.

639 Jones, K. & G. Moon (1993) Medical geography: taking space seriously. *Progress in Human Geography*,
640 17, 515-515.

641 Kajantie, E., C. Osmond, D. J. Barker, T. Forsen, D. I. Phillips & J. G. Eriksson (2005) Size at birth
642 as a predictor of mortality in adulthood: a follow-up of 350 000 person-years. *Int J Epidemiol*,
643 34, 655-63.

644 Kanungu District Local Government. 2013. Statistical Abstract 2012/13. ed. U. B. o. S. (UBOS).
645 Kanungu, Uganda.

646 Kassebaum, N. J., A. Bertozzi-Villa, M. S. Coggeshall, K. A. Shackelford, C. Steiner, K. R. Heuton,
647 D. Gonzalez-Medina, R. Barber, C. Huynh & D. Dicker (2014) Global, regional, and
648 national levels and causes of maternal mortality during 1990–2013: a systematic analysis for
649 the Global Burden of Disease Study 2013. *The Lancet*, 384, 980-1004.

650 Kramer, M. S. (1987) Determinants of low birth weight: methodological assessment and meta-
651 analysis. *Bulletin of the World Health Organization*, 65, 663-737.

652 Kramer, M. S., L. Séguin, J. Lydon & L. Goulet (2000) Socio-economic disparities in pregnancy
653 outcome: why do the poor fare so poorly? *Paediatric and Perinatal Epidemiology*, 14, 194-210.

654 Laaidi, M., A. Boumendil, T. C. Tran, H. Kaba, P. Rozenberg & P. Aegerter (2011) Effects of
655 meteorological conditions on pregnancy outcome: Literature review. *Environnement, Risques et*
656 *Sante*, 10, 128-141.

657 Lawn, J. E., H. Blencowe, S. Oza, D. You, A. C. C. Lee, P. Waiswa, M. Lalli, Z. Bhutta, A. J. D.
658 Barros, P. Christian, C. Mathers & S. N. Cousens (2014) Every Newborn: progress,
659 priorities, and potential beyond survival. *The Lancet*, 384, 189-205.

660 Lawn, J. E., S. Cousens & J. Zupan (2005) 4 million neonatal deaths: When? Where? Why? *The*
661 *Lancet*, 365, 891-900.

662 Lawn, J. E., K. Kerber, C. Enweronu-Laryea & O. Masee Bateman (2009) Newborn survival in low
663 resource settings—are we delivering? *BJOG: An International Journal of Obstetrics & Gynaecology*,
664 116, 49-59.

665 Low, J. A., M. H. Handley-Derry, S. O. Burke, R. D. Peters, E. A. Pater, H. L. Killen & E. J. Derrick
666 (1992) Association of intrauterine fetal growth retardation and learning deficits at age 9 to 11
667 years. *Am J Obstet Gynecol*, 167, 1499-505.

668 Ma, S. & B. K. Finch (2010) Birth Outcome Measures and Infant Mortality. *Population Research and*
669 *Policy Review*, 29, 865-891.

670 Macintyre, S., A. Ellaway & S. Cummins (2002) Place effects on health: how can we conceptualise,
671 operationalise and measure them? *Social science & medicine*, 55, 125-139.

672 MacKenzie, C. A. (2016) Filtered meaning: appreciating linguistic skill, social position and
673 subjectivity of interpreters in cross-language research. *Qualitative Research*, 16, 167-182.

674 Magadi, M. A., N. J. Madise & R. N. Rodrigues (2000) Frequency and timing of antenatal care in
675 Kenya: explaining the variations between women of different communities. *Social science &*
676 *medicine*, 51, 551-561.

677 Martinez Cobo, J. 1981. Study of the Problem of Discrimination Against Indigenous Populations:
678 Final report ed. U. N. D. f. S. P. a. D. I. Peoples. Geneva, Switzerland.

679 Maybury-Lewis, D. 2002. *Indigenous peoples, ethnic groups, and the state*. Needham, Massachusetts, USA:
680 Allyn & Baker.

681 McCormick, M. C. (1985) The Contribution of Low Birth Weight to Infant Mortality and Childhood
682 Morbidity. *New England Journal of Medicine*, 312, 82-90.

683 McGrath, J. J., T. H. Burne & D. W. Eyles (2005) Seasonal variation in birth weight. *Canadian Medical*
684 *Association Journal*, 173, 733-733.

685 McMichael, A. J., R. E. Woodruff & S. Hales (2006) Climate change and human health: Present and
686 future risks. *Lancet*, 367, 859-869.

687 McSweeney, C., M. New & G. Lizcano. 2010. UNDP Climate Change Country Profiles: Uganda.

688 Nahabwe, H. 2015. Public Health Officer at Bwindi Community Hospital. ed. S. MacVicar.

689 Neufeld, L., D. L. Pelletier & J. D. Haas (1999) The timing hypothesis and body proportionality of
690 the intra-uterine growth retarded infant. *American Journal of Human Biology*, 11, 638-646.

691 Ng, E. & C. Muntaner (2014) A Critical Approach to Macrosocial Determinants of Population
692 Health: Engaging Scientific Realism and Incorporating Social Conflict. *Current Epidemiology*
693 *Reports*, 1, 27-37.

694 Noor, A. M., D. K. Kinyoki, C. W. Mundia, C. W. Kabaria, J. W. Mutua, V. A. Alegana, I. S. Fall &
695 R. W. Snow (2014) The changing risk of Plasmodium falciparum malaria infection in Africa:
696 2000–10: a spatial and temporal analysis of transmission intensity. *The Lancet*, 383, 1739-
697 1747.

698 Nunkoosing, K. (2005) The Problems With Interviews. *Qualitative Health Research*, 15, 698-706.

699 O'Fallon, L. R. & A. Deary (2002) Community-based participatory research as a tool to advance
700 environmental health sciences. *Environmental Health Perspectives*, 110, 155-159.

701 Ohenjo, N., R. Willis, D. Jackson, C. Nettleton, K. Good & B. Mugarura (2006) Health of
702 Indigenous people in Africa. *Lancet*, 367, 1937-1946.

703 Olivero, J., J. E. Fa, M. A. Farfán, J. Lewis, B. Hewlett, T. Breuer, G. M. Carpaneto, M. Fernández,
704 F. Germi & S. Hattori (2016) Distribution and numbers of Pygmies in Central African
705 forests. *PloS one*, 11, e0144499.

706 Onyiriuka, A. N. (2006) Trends in incidence of delivery of low birth weight infants in Benin City,
707 southern Nigeria. *Niger Postgrad Med J*, 13, 189-94.

708 Paneth, N. S. (1995) The Problem of Low Birth Weight. *The Future of Children*, 5, 19-34.

709 Pike, I. L. (2000) Pregnancy outcome for nomadic Turkana pastoralists of Kenya. *American Journal of*
710 *Physical Anthropology*, 113, 31-45.

711 Pope, C., S. Ziebland & N. Mays (2000) Analysing qualitative data. *BMJ*, 320, 114-116.

712 Porpora, M. G., R. Lucchini, A. Abballe, A. M. Ingelido, S. Valentini, E. Fuggetta, V. Cardi, A.
713 Ticino, V. Marra, A. R. Fulgenzi & E. D. Felip (2013) Placental Transfer of Persistent
714 Organic Pollutants: A Preliminary Study on Mother-Newborn Pairs. *International Journal of*
715 *Environmental Research and Public Health*, 10, 699-711.

716 Prentice, A. M., T. J. Cole, F. A. Foord, W. H. Lamb & R. G. Whitehead (1987) Increased
717 birthweight after prenatal dietary supplementation of rural African women. *The American*
718 *Journal of Clinical Nutrition*, 46, 912-25.

719 Rayco-Solon, P., A. J. Fulford & A. M. Prentice (2005) Differential effects of seasonality on preterm
720 birth and intrauterine growth restriction in rural Africans. *The American Journal of Clinical*
721 *Nutrition*, 81, 134-139.

722 Rishworth, A., J. Dixon, I. Luginaah, P. Mkandawire & C. T. Prince (2016) "I was on the way to the
723 hospital but delivered in the bush": Maternal health in Ghana's Upper West Region in the
724 context of a traditional birth attendants' ban. *Social Science & Medicine*, 148, 8-17.

- 725 Rylander, C., J. O. Odland & T. M. Sandanger (2013) Climate change and the potential effects on
726 maternal and pregnancy outcomes: An assessment of the most vulnerable - the mother,
727 fetus, and newborn child. *Global Health Action*, 6.
- 728 Santos, H. P. O., A. M. Black & M. Sandelowski (2015) Timing of Translation in Cross-Language
729 Qualitative Research. *Qualitative Health Research*, 25, 134-144.
- 730 Sarker, B. K., M. Rahman, T. Rahman, J. Hossain, L. Reichenbach & D. K. Mitra (2016) Reasons for
731 Preference of Home Delivery with Traditional Birth Attendants (TBAs) in Rural Bangladesh:
732 A Qualitative Exploration. *PLoS one*, 11, e0146161.
- 733 Shaw, W. S., R. D. K. Herman & G. R. Dobbs (2006) Encountering indigeneity: Re-imagining and
734 decolonizing geography. *Geografiska Annaler: Series B, Human Geography*, 88, 267-276.
- 735 Smith, K. R., A. Woodward, D. Campbell-Lendrum, D. D. Chadee, Y. Honda, Q. Liu, J. M.
736 Olwoch, B. Revich & R. Sauerborn. 2014. Human health: impacts, adaptation, and co-
737 benefits. In *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and*
738 *Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the*
739 *Intergovernmental Panel of Climate Change*, eds. C. B. Field, V. R. Barros, D. J. Dokken, K. J.
740 Mach, M. D. Mastrandrea, T. E. Bilir, M. Chatterjee, K. L. Ebi, Y. O. Estrada, R. C. Genova,
741 B. Girma, E. S. Kissel, A. N. Levy, S. MacCracken, P. R. Mastrandrea & L. L. White, 709-
742 754. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press.
- 743 Smith, L. T. 1999. *Decolonizing methodologies: Research and indigenous peoples*. London, UK and New York,
744 USA: Zed Books.
- 745 Stephens, C., J. Porter, C. Nettleton & R. Willis (2006) Disappearing, displaced, and undervalued: a
746 call to action for Indigenous health worldwide. *The lancet*, 367, 2019-2028.
- 747 Strand, L. B., A. G. Barnett & S. Tong (2011) The influence of season and ambient temperature on
748 birth outcomes: A review of the epidemiological literature. *Environmental Research*, 111, 451-
749 462.
- 750 Uganda Bureau of Statistics (UBOS) and ICF International Inc. 2012. Uganda Demographic and
751 Health Survey 2011. Kampala, Uganda: UBOS and Calverton, MD, USA: ICF International
752 Inc.
- 753 UNICEF. 2014. State of the World's Children 2015: Reimagine the Future. ed. U. N. C. s. Fund.
754 New York, NY: UNICEF.
- 755 United Nations Inter-agency for Child Mortality Estimation (UN IGME). 2014. Levels & Trends in
756 Child Mortality. ed. W. UNICEF, World Bank, UNDP. New York, USA.
- 757 Wallerstein, N. B. & B. Duran (2006) Using Community-Based Participatory Research to Address
758 Health Disparities. *Health Promotion Practice*, 7, 312-323.
- 759 Watts, N., W. N. Adger, P. Agnolucci, J. Blackstock, P. Byass, W. Cai, S. Chaytor, T. Colbourn, M.
760 Collins, A. Cooper, P. M. Cox, J. Depledge, P. Drummond, P. Ekins, V. Galaz, D. Grace, H.
761 Graham, M. Grubb, A. Haines, I. Hamilton, A. Hunter, X. Jiang, M. Li, I. Kelman, L. Liang,
762 M. Lott, R. Lowe, Y. Luo, G. Mace, M. Maslin, M. Nilsson, T. Oreszczyn, S. Pye, T. Quinn,
763 M. Svensdotter, S. Venevsky, K. Warner, B. Xu, J. Yang, Y. Yin, C. Yu, Q. Zhang, P. Gong,
764 H. Montgomery & A. Costello (2015) Health and climate change: policy responses to protect
765 public health. *The Lancet*, 386, 1861-1914.
- 766 Wirth, M. E., D. Balk, E. Delamonica, A. Storeygard, E. Sacks & A. Minujin (2006) Setting the stage
767 for equity-sensitive monitoring of the maternal and child health Millennium Development
768 Goals. *Bulletin of the World Health Organization*, 84, 519-527.
- 769 Yeung, H. W.-c. (1997) Critical realism and realist research in human geography: a method or a
770 philosophy in search of a method? *Progress in Human Geography*, 21, 51-74.