

This is a repository copy of *Global temperature definition affects achievement of long-term climate goals*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/128103/>

Version: Accepted Version

---

**Article:**

Richardson, Mark, Cowtan, Kevin [orcid.org/0000-0002-0189-1437](https://orcid.org/0000-0002-0189-1437) and Millar, Richard J. (2018) Global temperature definition affects achievement of long-term climate goals. *Environmental Research Letters*. 054004. pp. 1-8. ISSN: 1748-9326

<https://doi.org/10.1088/1748-9326/aab305>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

ACCEPTED MANUSCRIPT • OPEN ACCESS

# Global temperature definition affects achievement of long-term climate goals

To cite this article before publication: Mark Richardson *et al* 2018 *Environ. Res. Lett.* in press <https://doi.org/10.1088/1748-9326/aab305>

## Manuscript version: Accepted Manuscript

Accepted Manuscript is “the version of the article accepted for publication including all changes made as a result of the peer review process, and which may also include the addition to the article by IOP Publishing of a header, an article ID, a cover sheet and/or an ‘Accepted Manuscript’ watermark, but excluding any other editing, typesetting or other changes made by IOP Publishing and/or its licensors”

This Accepted Manuscript is **US Government**.

As the Version of Record of this article is going to be / has been published on a gold open access basis under a CC BY 3.0 licence, this Accepted Manuscript is available for reuse under a CC BY 3.0 licence immediately.

Everyone is permitted to use all or part of the original content in this article, provided that they adhere to all the terms of the licence <https://creativecommons.org/licenses/by/3.0>

Although reasonable endeavours have been taken to obtain all necessary permissions from third parties to include their copyrighted content within this article, their full citation and copyright line may not be present in this Accepted Manuscript version. Before using any content from this article, please refer to the Version of Record on IOPscience once published for full citation and copyright details, as permissions may be required. All third party content is fully copyright protected and is not published on a gold open access basis under a CC BY licence, unless that is specifically stated in the figure caption in the Version of Record.

View the [article online](#) for updates and enhancements.

1 Global temperature definition affects achievement of long-term  
2 climate goals

3  
4 Mark Richardson<sup>a</sup>, Kevin Cowtan<sup>b</sup>, Richard J. Millar<sup>c</sup>

5  
6 <sup>a</sup>Corresponding author: Jet Propulsion Laboratory, California Institute of Technology. email:  
7 [markr@jpl.nasa.gov](mailto:markr@jpl.nasa.gov) Phone: (+1) 818-354-4270 Mail: Jet Propulsion Laboratory, M/S 233-300,  
8 4800 Oak Grove Drive, Pasadena, CA 91101, U.S.A.

9 <sup>b</sup>Department of Chemistry, University of York, Heslington, York, YO10 5DD, UK

10 <sup>c</sup>Environmental Change Institute, Oxford University Centre for the Environment, South Parks Road,  
11 Oxford, OX1 3QY, UK

12  
13 **Keywords:** Climate change, Global Warming, Paris Agreement, Carbon Budgets, Observational  
14 Temperature Records

©2017. All rights reserved.

**Abstract**

The Paris Agreement on climate change aims to limit “global average temperature” rise to “well below 2 °C” but reported temperature depends on choices about how to blend air and water temperature data, handle changes in sea ice and account for regions with missing data. Here we use CMIP5 climate model simulations to estimate how these choices affect reported warming and carbon budgets consistent with the Paris Agreement. By the 2090s, under a low-emissions scenario, modelled global near-surface air temperature rise is 15% higher (5—95% range 6—21 %) than that estimated by an approach similar to the HadCRUT4 observational record. The difference reduces to 8% with global data coverage, or 4% with additional removal of a bias associated with changing sea-ice cover. Comparison of observational datasets with different data sources or infilling techniques supports our model results regarding incomplete coverage. From high-emission simulations, we find that a HadCRUT4-like definition means higher carbon budgets and later exceedance of temperature thresholds, relative to global near-surface air temperature. 2 °C warming is delayed by seven years on average, to 2048 (2035—2060), and CO<sub>2</sub> emissions budget for a >50% chance of <2 °C warming increases by 67 GtC (246 GtCO<sub>2</sub>).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86

**1. Introduction**

Reflecting the 90—100 % consensus among relevant research(1, 2), the 5<sup>th</sup> Assessment Report of the Intergovernmental Panel on Climate Change (IPCC AR5) stated that “warming of the climate system is unequivocal” and “It is extremely [95—100 %] likely that human influence has been the dominant cause of the observed warming since the mid-20<sup>th</sup> century”.(3) Such scientific findings can inform policy responses in concert with other factors such as risk aversion, discounting of the future and assessments of the severity of future climate impacts. The Paris Agreement of the United Nations Framework Convention on Climate Change (UNFCCC), whose Article 2.1(a) expresses a long-term goal of:

“Holding the increase in the global average temperature to well below 2°C above pre-industrial levels and pursuing efforts to limit the temperature increase to 1.5°C above pre-industrial levels, recognizing that this would significantly reduce the risks and impact of climate change”.

However, “global average temperature” is not precisely defined, and achievement of the Agreement’s goal may depend on possible different definitions and available measurement techniques. A related concept is that of a carbon budget, the allowable cumulative carbon dioxide (CO<sub>2</sub>) emissions consistent with a specified level of peak warming with a particular probability(4–6).

The IPCC 5<sup>th</sup> Assessment Report (AR5) assessed carbon budgets for various levels of warming in billions of tonnes of carbon (GtC) or of carbon dioxide (GtCO<sub>2</sub>) based on projections of global near-surface air temperature change, which we refer to as “global-tas”, where tas means “temperature, air, at surface”, from complex Earth System Models (ESMs). In general, climate modelling studies use global-tas, whereas observational records typically combine non-global coverage of near-surface air temperature over land with sea-surface temperature (SST) over oceans into a single timeseries. As it is likely that stakeholders may have diverse interpretations as to what global average temperature refers, here we provide carbon budgets for different definitions of global average temperature, including definitions consistent with current observational products. Three main factors contribute to differences in “global average temperature” change between global-tas and observational records. Firstly, there are regions with missing data that may not warm at the global-mean rate. For example, the Arctic is now rapidly becoming warmer and wetter(7), but much of it is commonly excluded due to lack of long-term data(8). Secondly, under CO<sub>2</sub>-driven global warming, modelled near-surface air temperatures warm more than SSTs(9). Finally, data providers must decide how to account for changes in sea ice. There may be a change from reporting estimated near-surface air temperatures to SSTs where ice has retreated. In the HadCRUT4 dataset(10) this approach probably

©2017. All rights reserved.

results in an artificially low reported warming compared with the true air warming due to features of the normalisation procedure.

We refer to issues related to missing data as being due to “masking”, and the other two factors together as “blending”, specifically “air-sea blending” and “sea-ice blending”.

One early study accounted for the masking and air-sea blending issues(11), and some studies have accounted for masking but this is not universal. Recently, it was shown that over 1861—1880 to 2000—2009, modelled global-tas increased 24 % more than a HadCRUT4-like blended-masked estimate(12). Current observed temperature records should therefore exceed 2°C later than global-tas, implying a larger carbon budget if compliance were assessed using one of them. Here we extend this prior work by (i) reporting results to 2099, (ii) calculating carbon budgets using IPCC techniques, (iii) accounting for realistic potential future data coverage and (iv) applying blending and masking to a low-emission scenario. In particular, the addition of a low-emission scenario allows us to determine to what extent temperature definitions matter if policymakers choose to take strong mitigation action.

Future blending-masking biases may change relative to the past because of increased modern data coverage: indeed, the blending-masking bias under transient warming with 2000—2009 data coverage was estimated to be 15 % instead of 24 %(12). Furthermore, with strong mitigation sea-ice cover would be expected to stabilise before 2100, suppressing the future sea-ice blending bias(13). In addition, the long-term warming pattern may differ from the historical pattern, leading to a different effect of coverage bias(14–16).

## 2. Methods

We consider two emission scenarios from the Coupled Model Intercomparison Project, phase 5 (CMIP5): the low emissions Representative Concentration Pathway 2.6 (RCP2.6(17, 18)) and the high emissions RCP8.5 (19). Among CMIP5 scenarios, only RCP2.6 has a substantial probability of <2 °C warming so we use it as representative of a world of strong mitigation. This allows us to estimate shifts in the probability of compliance with Paris targets in such a world, and to determine whether the magnitude of blending and masking biases should change substantially in the future. Meanwhile, RCP8.5 is used to estimate carbon budgets in a manner that is comparable with a set reported by the IPCC 5<sup>th</sup> Assessment Report. Note that we report decadal temperature changes relative to 1861—1880 to include simulations beginning in 1861 and avoid major volcanic eruptions. Supplementary Figures 1 & 2 further justify the choice of these reference periods.

We process CMIP5 simulations on a  $1 \times 1^\circ$  lat-lon grid using the Cowtan et al. (2015)(20) algorithm and assuming that 2005—2014 geographic data coverage is maintained in future. This is done by downsampling the HadCRUT4 historical coverage up to December 2014 to  $1 \times 1^\circ$  and extending this coverage to December 2099 in the following fashion. For each calendar month, coverage is allowed if data are reported at that location for that calendar month in more than 5 years from 2005—2014 inclusive. Mapping at  $1 \times 1^\circ$  instead of  $5 \times 5^\circ$  doesn't affect reported global temperature but keeps spatial information that may be useful in future. We area weight all reporting cells, whereas HadCRUT4 calculates hemispheres separately then averages those: this introduces a minor 1.9 % difference in 1861—2016 warming (Supplementary Figure 3).

We calculate 4 temperature series for each simulation beginning with the widely used “global-tas”, and then add the effect of SST blending by mixing air temperatures and SSTs before calculating the anomalies, which we call “air-sea blended”. Next, we add the effect of sea-ice blending by calculating the anomalies in air and ocean temperatures separately before combining them, and call this “fully blended”. Finally we restrict coverage to follow the historical or assumed future HadCRUT4-like data availability and call this “blended-masked”.

We select all CMIP5 simulations that have continuous historical and RCP2.6 or RCP8.5 runs from 1861—2099 inclusive and for which we could obtain the required output fields. These fields are Near-surface Air Temperature (short name “tas”), Sea Surface Temperature (SST, “tos”), Sea Ice Concentration (“sic”) and Sea Area Fraction (“sftof”, see the CMIP5 Standard Output description at [http://cmip-pcmdi.llnl.gov/cmip5/data\\_description.html](http://cmip-pcmdi.llnl.gov/cmip5/data_description.html) ). Simulations are listed in Supplementary Tables 1 & 2 and model configurations can be found in Table 9.A.1 of AR5 (21). Each simulation was processed using the Cowtan et al. (2015) code and our updated future coverage mask. Blended temperature at the  $i,j^{th}$  grid point,  $T_{blend,i,j}$  are obtained using:

$$T_{blend,i,j} = w_{air,i,j} T_{air,i,j} + (1 - w_{air,i,j}) T_{ocean,i,j} \quad (1)$$

Where  $w_{air,i,j}$  is the fraction of the grid cell from which near-surface air temperatures are taken,  $T_{air,i,j}$  refers to the local air temperature “tas” and  $T_{ocean,i,j}$  the local SST “tos”. Each of these is converted into temperature anomaly relative to the local baseline of the same type (i.e. air or water). After the local anomalies are calculated, the grid points are then averaged with a spherical Earth area weighting. For global-tas,  $w_{air} = 1$  always, while for blended series  $w_{air,i,j}$  is the fraction of land plus sea ice within the grid cell. For air-sea blended, a grid cell's  $w_{air,i,j}$  is fixed based on the initial sea ice extent whereas for fully blended the sea-ice fraction changes depending on the monthly sea ice concentration.

©2017. All rights reserved.

Carbon Threshold Exceedance Budgets (TEBs) are calculated as in the Technical Summary of IPCC AR5(22). Linear interpolation between decadal means are used to compute the diagnosed cumulative CO<sub>2</sub> emissions since 1870 to the point that warming exceeds a given temperature threshold. Unlike in ref. (22), only complex ESMs are included in the analysis with Earth System Models of Intermediate Complexity (EMICs) excluded. Reported percentiles correspond to percentiles of the distribution of ESM TEBs for that warming threshold. ESMs (models that can interactively diagnose compatible CO<sub>2</sub> emissions with a prescribed concentration pathway) considered here are identified with an asterisk in Supplementary Table 2.

### 3. Results

#### 3.1 Effect of temperature definition under low emissions.

Figure 1(a) shows the CMIP5 historical-RCP2.6 and historical-RCP8.5 ensemble time series of global-tas. Figure 1(b) shows the blending-masking differences and Figure 1(c) the decadal averages of these differences as a function of global-tas for historical-RCP2.6 and panels (d) and (e) the same for historical-RCP8.5. All results shown here use a single simulation from each model, labelled “r1i1p1” in CMIP5 nomenclature. Results are not sensitive to including the full ensemble (Supplementary Table 3 and Supplementary Figure 4).

In RCP2.6 the air-sea blending bias stabilises and begins to decrease in the last ~70 years of the simulations while Figure 1(c) shows that the sea-ice-blending and masking biases increase with global-tas throughout the series, but at a much slower rate than under RCP8.5. This suggests that the temperature stabilisation reduces sea-ice loss and its contribution to reported temperature bias. Similarly, the errors bars in Figure 1(b) show that uncertainty introduced by sea ice change is smaller under RCP2.6.

However, temperature bias still continues to grow with time in RCP2.6, and Figure 2 demonstrates that the masking bias component is likely dominated by the warming at high northern latitudes, which tend to warm much more than the global average and are poorly sampled.

Table 1 contains the ensemble median and 5—95 % range for RCP2.6 and RCP8.5 temperature changes over periods spanning the past (1861—1880), present (2007—2016) and future (2090—2099). Under RCP2.6, the percentage of simulations consistent with 2 °C warming increases from 75 % for air-sea blended (the same as for global-tas) to 90 % for blended-masked. Percentage blended-masked bias is calculated separately for each simulation and the median and ranges of these



percentages are reported: the 16 % bias for 1861—2016 differs from the 24 % reported previously(12) due to the changed time period, available RCP2.6 runs, and because this result doesn't use the same HadCRUT4 hemisphere-weighting.

The ensemble suggests a decrease in air-sea blending bias in future, with global-tas warming from 2007—2016 to 2090—2099 just 1.9 % (0.5—3.9 %, all bracketed values 5—95 % ensemble range) greater than the air-sea blended value. In addition, improved geographical data coverage relative to most of the historical period reduces the masking bias, although the ice-blending issue remains at a similar magnitude. Overall, 21<sup>st</sup> century global-tas warming is 10.6 (1.2—29.7) % greater than the blended-masked estimate. The full-period blending-masking bias from 1861—1880 is approximately 14.9 (5.7—20.6) %.

As masking contributes the most to our blending-masking biases we assess whether our model-based estimates are realistic by considering observational data records that handle land data in different ways and have different masking biases. These datasets are HadCRUT4(10), Cowtan and Way(8) and Berkeley Earth,(23, 24) all of which combine land air temperature data with the HadSST3 ocean product(25, 26) and extend over our full period. HadCRUT4 is subject to the full blending-masking bias while Cowtan & Way follows the HadCRUT4 method except that missing regions are infilled by kriging, a statistical method that accounts for spatial covariance in the field and more heavily weights nearby data. Berkeley Earth uses a similar approach, but handles the raw station data in a different manner.

From 1861—1880 to 2007—2016 the global warming in HadCRUT4 is 0.84 °C, in Cowtan & Way is 0.94 °C and in Berkeley Earth is 0.99 °C. The two records which do not assume that regions of missing data warm at the global-average rate show 12—18 % more warming, the same order of magnitude as the masking biases inferred from CMIP5 simulations, although particularly poor historical coverage over the Antarctic and Southern Ocean means that infilling from neighbouring regions may be inadequate.

3.2 Carbon budgets and temperature thresholds under higher emissions.

*IPCC AR5 carbon budgets correspond to cumulative emissions compatible with thresholds of modelled global-tas warming. Carbon budgets for a 1.5 °C or 2 °C warming in any form of blended or blended-masked estimate will therefore be higher than the corresponding IPCC AR5 budget. Budgets given in*

$(\Delta T_{global-tas} - \Delta T_{blended})/\Delta T_{global-tas}$ (%)						
Period:	historical-RCP2.6			historical-RCP8.5		
	Air-sea blended	Fully blended	Blended-masked	Air-sea blended	Fully blended	Blended-masked

©2017. All rights reserved.

1861—1880 to 2007—2016	5.8 (3.2-7.3)	8.7 (6.0-10.9)	16.2 (5.2-28.7)	5.8 (3.9-7.4)	8.9 (6.2-11.7)	17.9 (5.5-27.1)
2007—2016 to 2090—2099	1.9 (0.5-3.9)	6.9 (3.9-12.0)	10.6 (1.2-29.7)	4.0 (2.5-6.1)	10.1 (8.4-11.7)	11.7 (8.2-15.9)
1861—1880 to 2090—2099	4.2 (2.3-6.3)	8.0 (5.4-9.8)	14.9 (5.7-20.6)	4.3 (2.8-6.3)	9.7 (7.3-11.4)	12.8 (7.4-18.1)
Cases with $\Delta T < 2^{\circ}\text{C}$	15/20 (75 %)	15/20 (75 %)	18/20 (90 %)	N/A	N/A	N/A

Table 2 correspond to cumulative CO<sub>2</sub> emissions since 1870 until the point of exceeding 1.5 °C or 2 °C warming (a threshold exceedance budget or TEB(22, 27)) under RCP8.5 (see Methods). The IPCC AR5 results are also included for comparison, and differ somewhat since they include EMIC runs and were reported to the nearest 50 GtCO<sub>2</sub>, or approximately 13.6 GtC.

For the blended-masked timeseries the 1.5 °C and 2 °C thresholds are reached a median 7—8 years later than for global-tas under this high-emission scenario. This has implications for carbon budgets, with the TEB for which 50 % of the ESMs have warming below 1.5 °C increasing by 53 GtC (194 GtCO<sub>2</sub>) and 67 GtC (246 GtCO<sub>2</sub>) for the 2 °C threshold.

The IPCC carbon budgets were reported relative to 1870, but policymakers require up-to-date guidance to inform discussions related to the Paris Agreement. We therefore also calculate the remaining post-2015 carbon budget based on the ESM ensemble after adjusting for observed warming through 2015 following the approach of Millar et al. (2017, (28)). For example, given that HadCRUT4 shows approximately 0.9 °C human-induced warming to 2015, another 0.6 °C results in a total of 1.5 °C. In our ESM simulations, the remaining blended-masked carbon budget with a >66 % chance of <0.6 °C warming post-2015 is 246 GtC. However, if Berkeley Earth were to be used, then historical human-induced warming is greater. It would also likely show greater future warming for a given quantity of CO<sub>2</sub> emissions too as Berkeley better approximates air-sea blended temperatures rather than the blended-masked approach of HadCRUT4. We estimate the remaining 1.5 °C budget at near 161 GtC in that case (see Supplementary Table 4 and related discussion).

#### 4. Discussion

Here we have shown that achievement of the Paris Agreement's long-term goals could depend on the definition of "global average temperature". The scientific background to the Paris Agreement was informed directly by the Structured Expert Dialogue(29), which used non-infilled datasets to track warming to date. Our results indicate the potential impact of choosing different types of

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

236 observational product in the future to measure global temperatures in the context of the  
237 Agreement. As it is unlikely that estimates of global mean air temperature, which inherently rely on  
238 climate models, will be used, we show how the use of “blended” observational products would  
239 increase the policy-relevant carbon budgets for 2 °C relative to the global air-temperature budgets  
240 given by IPCC-AR5 (see Table 2).

241 A recent study estimated the post-2015 carbon budget with a >66 % chance of achieving a 1.5 °C  
242 target at 204 GtC, rather than the 70 GtC implied by IPCC AR5, suggesting that the 1.5 °C target is  
243 “not yet a geophysical impossibility”, but likely requires “strengthened pledges for 2030 followed by  
244 challengingly deep and rapid mitigation”, i.e. cuts in net anthropogenic emissions (Millar et al., 2017,  
245 (28)). The Millar et al. value differs from IPCC-AR5 budgets as it updated these calculations using  
246 observed warming and emissions from a 2010—2019 reference period using CMIP5-consistent  
247 relationships between future warming and future CO<sub>2</sub> emissions. A fraction of this difference was  
248 due to the IPCC carbon budgets being calculated for global-tas, whereas Millar et al. used human-  
249 induced warming estimated from the blended-masked HadCRUT4 dataset, consistent with the  
250 Structured Expert Dialogue, to define the remaining warming between the present-decade and  
251 1.5°C.

252 If an alternative observational dataset were used to monitor global temperature in the context of  
253 the Paris Agreement then estimates of human-induced warming and compatible carbon budgets  
254 would change. For example, the Berkeley Earth product uses infilling techniques with more data  
255 sources and a different sea-ice algorithm which should reduce differences with global-tas. It shows  
256 almost 20 % more human-induced warming than HadCRUT4 through 2015, and hence would reduce  
257 post-2015 carbon budgets by around 80 GtC.

258 Biases associated with incomplete data coverage and the blending of air and water data both  
259 suppress reported warming relative to global near-surface air temperatures. Our analysis and results  
260 in Table 1 indicate that these biases will tend to be smaller in future provided that the improved  
261 data coverage of recent decades is maintained. Furthermore, under a scenario of strong mitigation,  
262 the differences introduced by the retreat of sea-ice are marginally smaller than under high emissions  
263 where sea ice retreat is more pronounced.

264 **5. Conclusion**

265 We have demonstrated here the importance of a clear understanding of different definitions of  
266 global mean temperature with regards to carbon budgets and achievement of long-term climate  
267 goals under the Paris Agreement. We propose that the definition of global mean temperature should

©2017. All rights reserved.

be physically based, transparent and verifiable in order for stakeholders to have confidence in its value. For a timeseries to be truly “global”, it must account for the incomplete spatial coverage of direct observations, requiring techniques such as those used in Cowtan & Way or Berkeley Earth. It is key that policy-makers unambiguously elucidate how they intend to measure global temperatures in the context of the Paris Agreement to enable the most useful mitigation advice to be provide by the scientific community. If pure observation-based timeseries are used then further efforts for data-recovery in data-sparse regions would help, as would more long-term stations at high latitudes. In addition, the sea-ice blending effect is a non-physical artefact of algorithm design and it should be possible to account for this in future datasets. However, the long-term air-sea blending effect is difficult to verify due to the lack of robust, homogenised and long-term collocated air-SST ocean data, and its lack of measurability may justify the definition of global-average temperature as being an air-sea blended value. Under this definition, potential blending biases are reduced to an equivalent of an apparent 2—3 year delay in exceeding temperature targets, instead of the 7—8 years for a fully blended-masked series.

## Acknowledgements

MR's contribution was carried out at the Jet Propulsion Laboratory, California Institute of Technology under contract with the National Aeronautics and Space Administration. RJM was financially supported by the Oxford Martin Net Zero Carbon Investment Initiative.

## Author contributions

M.R. generated the main figures, generated the RCP time series, calculated the non-carbon-budget results, and contributed to the text. K.C. produced & supports the blending-masking code, did the supplementary observation analysis and contributed to the text. R.J.M. did the carbon budget analysis and contributed to the text.

## References

1. Cook J, et al. (2013) Quantifying the consensus on anthropogenic global warming in the scientific literature. *Environ Res Lett* 8(2):24024.

©2017. All rights reserved.

- 298 2. Cook J, et al. (2016) Consensus on consensus: a synthesis of consensus estimates on human-  
299 caused global warming. *Environ Res Lett* 11(4):48002.
- 300 3. Stocker T, et al. (2013) *IPCC, 2013: Summary for Policymakers. In: Climate Change 2013: The*  
301 *Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the*  
302 *Intergovernmental Panel on Climate Change.* (Cambridge, United Kingdom and New York, NY,  
303 USA).
- 304 4. Meinshausen M, et al. (2009) Greenhouse-gas emission targets for limiting global warming to  
305 2 °C. *Nature* 458(7242):1158–1162.
- 306 5. Allen MR, et al. (2009) Warming caused by cumulative carbon emissions towards the  
307 trillionth tonne. *Nature* 458(7242):1163–1166.
- 308 6. Matthews HD, Caldeira K (2008) Stabilizing climate requires near-zero emissions. *Geophys Res*  
309 *Lett* 35(4):L04705.
- 310 7. Boisvert LN, Stroeve JC (2015) The Arctic is becoming warmer and wetter as revealed by the  
311 Atmospheric Infrared Sounder. *Geophys Res Lett* 42(11):4439–4446.
- 312 8. Cowtan K, Way RG (2014) Coverage bias in the HadCRUT4 temperature series and its impact  
313 on recent temperature trends. *Q J R Meteorol Soc* 140(683):1935–1944.
- 314 9. Richter I, Xie S-P (2008) Muted precipitation increase in global warming simulations: A surface  
315 evaporation perspective. *J Geophys Res* 113(D24):D24118.
- 316 10. Morice CP, Kennedy JJ, Rayner NA, Jones PD (2012) Quantifying uncertainties in global and  
317 regional temperature change using an ensemble of observational estimates: The HadCRUT4  
318 data set. *J Geophys Res Atmos* 117(D8):n/a-n/a.
- 319 11. Santer BD (2000) Interpreting Differential Temperature Trends at the Surface and in the  
320 Lower Troposphere. *Science* (80- ) 287(5456):1227–1232.
- 321 12. Richardson M, Cowtan K, Hawkins E, Stolpe MB (2016) Reconciled climate response estimates  
322 from climate models and the energy budget of Earth. *Nat Clim Chang*.  
323 doi:10.1038/nclimate3066.
- 324 13. Swart NC, Fyfe JC, Hawkins E, Kay JE, Jahn A (2015) Influence of internal variability on Arctic  
325 sea-ice trends. *Nat Clim Chang* 5(2):86–89.
- 326 14. Armour KC, Bitz CM, Roe GH (2013) Time-Varying Climate Sensitivity from Regional  
327 Feedbacks. *J Clim* 26(13):4518–4534.

©2017. All rights reserved.

15. Andrews T, Gregory JM, Webb MJ (2015) The Dependence of Radiative Forcing and Feedback on Evolving Patterns of Surface Temperature Change in Climate Models. *J Clim* 28(4):1630–1648.
16. Held IM, et al. (2010) Probing the Fast and Slow Components of Global Warming by Returning Abruptly to Preindustrial Forcing. *J Clim* 23(9):2418–2427.
17. Taylor KE, Stouffer RJ, Meehl GA (2012) An Overview of CMIP5 and the Experiment Design. *Bull Am Meteorol Soc* 93(4):485–498.
18. van Vuuren DP, et al. (2011) RCP2.6: exploring the possibility to keep global mean temperature increase below 2°C. *Clim Change* 109(1–2):95–116.
19. Riahi K, et al. (2011) RCP 8.5—A scenario of comparatively high greenhouse gas emissions. *Clim Change* 109(1–2):33–57.
20. Cowtan K, et al. (2015) Robust comparison of climate models with observations using blended land air and ocean sea surface temperatures. *Geophys Res Lett* 42(15):6526–6534.
21. Flato G, et al. (2013) *Evaluation of Climate Models in: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (Cambridge, United Kingdom and New York, NY, USA).
22. Stocker T (2013) *IPCC, 2013: Technical Summary. In: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.* (Cambridge, United Kingdom and New York, NY, USA).
23. A. Muller R, Rohde R, Jacobsen R, Muller E, Wickham C (2013) A New Estimate of the Average Earth Surface Land Temperature Spanning 1753 to 2011. *Geoinformatics Geostatistics An Overv* 1(1). doi:10.4172/2327-4581.1000101.
24. Rohde R, Muller R, Jacobsen R, Perlmutter S, Mosher S (2013) Berkeley Earth Temperature Averaging Process. *Geoinformatics Geostatistics An Overv* 1(2). doi:10.4172/2327-4581.1000103.
25. Kennedy JJ, Rayner NA, Smith RO, Parker DE, Saunby M (2011) Reassessing biases and other uncertainties in sea surface temperature observations measured in situ since 1850: 1. Measurement and sampling uncertainties. *J Geophys Res* 116(D14):D14103.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

26. Kennedy JJ, Rayner NA, Smith RO, Parker DE, Saunby M (2011) Reassessing biases and other uncertainties in sea surface temperature observations measured in situ since 1850: 2. Biases and homogenization. *J Geophys Res* 116(D14):D14104.

27. Rogelj J, et al. (2016) Differences between carbon budget estimates unravelled. *Nat Clim Chang* 6(3):245–252.

28. Millar RJ, et al. (2017) Emission budgets and pathways consistent with limiting warming to 1.5 °C. *Nat Geosci* 10(10):741–747.

29. “Subsidiary Body for Scientific and Technological Advice”, Implementation" "Subsidiary Body for (2015) *Report on the structured expert dialogue on the 2013–2015 review* Available at: [http://unfccc.int/science/workstreams/the\\_2013-2015\\_review/items/7521.php](http://unfccc.int/science/workstreams/the_2013-2015_review/items/7521.php).

30. Dodd EMA, Merchant CJ, Rayner NA, Morice CP (2015) An Investigation into the Impact of using Various Techniques to Estimate Arctic Surface Air Temperature Anomalies\*. *J Clim* 28(5):1743–1763.

©2017. All rights reserved.




Figure 1 – (a) CMIP5 global near-surface air temperature change under RCP2.6 and RCP8.5 relative to 1861–1880, with the ensemble median as a line and the shaded area representing the 5–95 % ensemble range (r1i1p1 simulations only). (b,d) The difference for each scenario (as labelled) between the CMIP5 ensemble median blended-masked temperature change and the global tas-only, shown as blended-masked minus tas-only. Each line represents one extra blending or masking factor: blue is ocean-blend only, orange is ocean-blend plus sea-ice blend, and green is both blends plus masking for data coverage. On the right of the figure, each point and bar represents the ensemble median and 5–95 % range of each difference for the final decade. (c,e) decadal means of the differences from (b,d) plotted as a function of the global tas-only temperature change relative to 1861–1880 for the labelled scenario.






Figure 2 – Time series of mean CMIP5 near-surface air temperature change from 1861–1880 under historical-RCP2.6 scenario split into latitude bands as labelled in the legend. The solid black line shows the global-tas value, undersampling of any region that shows warming above this black line will lead to a cooling masking bias, and vice versa. The modelled bias is dominated by undersampling of the northern high latitudes where by far the greatest warming occurs.

©2017. All rights reserved.

**Table 1 – Percentage increase in observed temperature change between selected periods when considering global-tas relative to the blended or blended-masked version. “Fully blended” includes the sea-ice change effect in addition to air-water warming differences. CMIP5 historical-RCP2.6 ensemble median reported with 5–95 % range in brackets. Bottom row shows the number of simulations and percentage of ensemble that show <2 °C difference between 1861–1880 and 2090–2099.**

Period:	$(\Delta T_{\text{global-tas}} - \Delta T_{\text{blended}}) / \Delta T_{\text{global-tas}} (\%)$			$(\Delta T_{\text{global-tas}} - \Delta T_{\text{blended}}) / \Delta T_{\text{global-tas}} (\%)$		
	historical-RCP2.6			historical-RCP8.5		
	Air-sea blended	Fully blended	Blended-masked	Air-sea blended	Fully blended	Blended-masked
1861–1880 to 2007–2016	5.8 (3.2-7.3)	8.7 (6.0-10.9)	16.2 (5.2-28.7)	5.8 (3.9-7.4)	8.9 (6.2-11.7)	17.9 (5.5-27.1)
2007–2016 to 2090–2099	1.9 (0.5-3.9)	6.9 (3.9-12.0)	10.6 (1.2-29.7)	4.0 (2.5-6.1)	10.1 (8.4-11.7)	11.7 (8.2-15.9)
1861–1880 to 2090–2099	4.2 (2.3-6.3)	8.0 (5.4-9.8)	14.9 (5.7-20.6)	4.3 (2.8-6.3)	9.7 (7.3-11.4)	12.8 (7.4-18.1)
Cases with $\Delta T < 2^\circ\text{C}$	15/20 (75 %)	15/20 (75 %)	18/20 (90 %)	N/A	N/A	N/A

**Table 2 – Estimated carbon budgets expressed in GtC for various percentiles of the ESM distribution for 1.5 °C or 2 °C global warming thresholds and different definitions of “global average temperature”. The median and 5–95 % ensemble range of exceedance years are also shown and correspond to the full set of RCP8.5 CMIP5 simulations and not just the ESM subset.**

Percentile of ESM distribution	1.5 °C budget (GtC)			2 °C budget (GtC)			Year $\Delta T$ exceeded	
	>33 %	>50 %	>66 %	>33 %	>50 %	>66 %	1.5 °C	2 °C
IPCC since 1870	695	614	614	900	818	791	-	-
global-tas	703	667	588	931	852	794	2027 (2015–2039)	2041 (2028–2053)
air-sea blended	731	692	627	951	889	831	2030 (2018–2041)	2043 (2031–2056)
fully-blended	749	708	645	987	916	849	2031 (2018–2041)	2044 (2032–2058)
blended-masked	787	720	638	1053	919	855	2035 (2022–2044)	2048 (2035–2060)