

This is a repository copy of *Disruption of the GDP-mannose synthesis pathway in Streptomyces coelicolor results in antibiotic hyper-susceptible phenotypes.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/128230/>

Version: Accepted Version

---

**Article:**

Howlett, Robert, Anttonen, Katri, Read, Nicholas et al. (1 more author) (2018) Disruption of the GDP-mannose synthesis pathway in *Streptomyces coelicolor* results in antibiotic hyper-susceptible phenotypes. *Microbiology* (Reading, England). ISSN: 1465-2080

<https://doi.org/10.1099/mic.0.000636>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# Microbiology

## Disruption of the GDP-mannose synthesis pathway in *Streptomyces coelicolor* results in antibiotic hyper-susceptible phenotypes

--Manuscript Draft--

Manuscript Number:	MIC-D-18-00002R1
Full Title:	Disruption of the GDP-mannose synthesis pathway in <i>Streptomyces coelicolor</i> results in antibiotic hyper-susceptible phenotypes
Article Type:	Research Article
Section/Category:	Physiology and metabolism
Corresponding Author:	Margaret CM Smith, PhD University of Leeds Leeds, West Yorkshire UNITED KINGDOM
First Author:	Robert Howlett, PhD
Order of Authors:	Robert Howlett, PhD Katri Anttonen, PhD Nicholas Read, PhD Margaret CM Smith, PhD
Abstract:	<p>Actinomycete bacteria use polyprenol phosphate mannose as a lipid linked sugar donor for extra-cytoplasmic glycosyl transferases that transfer mannose to cell envelope polymers, including glycoproteins and glycolipids. We showed recently that strains of <i>Streptomyces coelicolor</i> with mutations in the <i>ppm1</i> gene, which encodes polyprenol phosphate mannose synthase, were both resistant to phage <math>\phi</math>C31 and have greatly increased susceptibility to antibiotics that mostly act on cell wall biogenesis. Here we show that mutations in the genes encoding enzymes that act upstream of Ppm1 in the polyprenol phosphate mannose synthesis pathway can also confer phage resistance and antibiotic hyper-susceptibility. GDP-mannose is a substrate for Ppm1 and is synthesized by GDP-mannose pyrophosphorylase (GMP;ManC) which uses GTP and mannose-1-phosphate as substrates. Phosphomannomutase (PMM;ManB) converts mannose-6-phosphate to mannose-1 phosphate. <i>S. coelicolor</i> strains with knocked down GMP activity or with a mutation in <i>sco3028</i> encoding PMM acquire phenotypes that resemble those of the <i>ppm1</i>- mutants i.e. <math>\phi</math>C31 resistant and susceptible to antibiotics. Differences in the phenotypes of the strains were observed, however. While the <i>ppm1</i>- strains have a small colony phenotype, the <i>sco3028::Tn5062</i> mutants had an extremely small colony phenotype indicative of an even greater growth defect. Moreover we were unable to generate a strain in which GMP activity encoded by <i>sco3039</i> and <i>sco4238</i> is completely knocked out, indicating that GMP is also an important enzyme for growth. Possibly GDP-mannose is at a metabolic branch point where it is modified to supply alternative nucleotide sugar donors.</p>

**Disruption of the GDP-mannose synthesis pathway in *Streptomyces coelicolor*  
results in antibiotic hyper-susceptible phenotypes**

Robert Howlett<sup>a</sup>, Katri Anttonen<sup>b</sup>, Nicholas Read<sup>a</sup>, and Margaret C.M Smith<sup>a,b, #</sup>

<sup>a</sup> Department of Biology, University of York, York, United Kingdom;

<sup>b</sup> Institute of Medical Sciences, University of Aberdeen, Aberdeen, United Kingdom;

#Address correspondence to Margaret C.M. Smith, [Maggie.smith@york.ac.uk](mailto:Maggie.smith@york.ac.uk)

Running Title: *S. coelicolor* strains disrupted in the synthesis of GDP-mannose

Abbreviations; Ppm1; polyprenol phosphate mannose synthase, PMM;  
Phosphomannomutase, GMP; GDP-mannose pyrophosphorylase, PGM;  
phosphoglucomutase, MESG; 2-amino-6-mercapto-7-methylpurine ribonucleoside,  
PNP; purine nucleoside phosphorylase, SMMS; supplemented minimal medium  
solid.

18

19 **Abstract**

20 Actinomycete bacteria use polyprenol phosphate mannose as a lipid linked sugar  
21 donor for extra-cytoplasmic glycosyl transferases that transfer mannose to cell  
22 envelope polymers, including glycoproteins and glycolipids. We showed recently that  
23 strains of *Streptomyces coelicolor* with mutations in the gene *ppm1* encoding  
24 polyprenol phosphate mannose synthase were both resistant to phage  $\phi$ C31 and  
25 have greatly increased susceptibility to antibiotics that mostly act on cell wall  
26 biogenesis. Here we show that mutations in the genes encoding enzymes that act  
27 upstream of Ppm1 in the polyprenol phosphate mannose synthesis pathway can also  
28 confer phage resistance and antibiotic hyper-susceptibility. GDP-mannose is a  
29 substrate for Ppm1 and is synthesized by GDP-mannose pyrophosphorylase  
30 (GMP;ManC) which uses GTP and mannose-1-phosphate as substrates.  
31 Phosphomannomutase (PMM; ManB) converts mannose-6-phosphate to mannose-1  
32 phosphate. *S. coelicolor* strains with knocked down GMP activity or with a mutation in  
33 *sco3028* encoding PMM acquire phenotypes that resemble those of the *ppm1*<sup>-</sup>  
34 mutants i.e.  $\phi$ C31 resistant and susceptible to antibiotics. Differences in the  
35 phenotypes of the strains were observed, however. While the *ppm1*<sup>-</sup> strains have a  
36 small colony phenotype, the *sco3028::Tn5062* mutants had an extremely small  
37 colony phenotype indicative of an even greater growth defect. Moreover we were  
38 unable to generate a strain in which GMP activity encoded by *sco3039* and *sco4238*  
39 is completely knocked out, indicating that GMP is also an important enzyme for  
40 growth. Possibly GDP-mannose is at a metabolic branch point where it is modified to  
41 supply alternative nucleotide sugar donors.

42

## Introduction

*Streptomyces* spp. are prolific producers of secondary metabolites, many with potent antibiotic activity. In nature *Streptomyces* spp. produce antibiotics either to inhibit competitors thus providing the producer with a growth advantage or as signalling molecules in microbial communities [1, 2]. Either way *Streptomyces* bacteria are constantly exposed to antibiotics produced by other soil microorganisms and consequently have evolved resistance mechanisms [3]. As such *Streptomyces* spp. are a model system to study how the mechanisms of antibiotic resistance evolve in an environmental organism.

We recently showed that strains of *S. coelicolor* lacking the ability to synthesise polyprenol phosphate mannose due to mutations in polyprenol phosphate mannose synthase (Ppm1) were hyper-sensitive to multiple antibiotics (Howlett et al, submitted). We used RNA-seq and Raman spectroscopy to demonstrate that the strains had undergone changes to the membrane phospholipids, with possible subsequent changes to membrane functions. Polyprenol phosphate mannose synthase, Ppm1, transfers mannose from GDP-mannose to polyprenol phosphate (Fig. 1). Previously we demonstrated that the synthesis of polyprenol phosphate mannose was entirely dependent on membrane associated Ppm1 [4].

Polyprenol phosphate mannose is the mannose donor for extracytoplasmic glycosyl transferases. One of these is a protein mannosyl transferase (Pmt), which glycosylates periplasmic and membrane proteins in *Streptomyces* [4, 5]. Pmt defective strains also show increased antibiotic susceptibility compared to the parent strain, but to fewer antibiotics and to a lower level than the *ppm1*<sup>-</sup> mutants (Howlett et

al, submitted). Loss of protein glycosylation is therefore likely to contribute in part to the antibiotic hyper susceptible phenotype of the *ppm1<sup>-</sup>* mutants. In addition both *ppm1<sup>-</sup>* strains and the *pmt* strains are resistant to the phage  $\phi$ C31, most likely through loss of the receptor, although the exact nature of the phage receptor is still unknown [5, 6].

Polyprenol phosphate mannose is likely to be a mannose donor for other cell envelope macromolecules with one of these likely to be phosphoinositol mannosides (PIMs) [7, 8]. In other Actinobacteria including *Mycobacterium* and *Corynebacterium* spp. PIMs are precursors for the synthesis of lipoarabinomannan and lipomannan [9], but neither of these polymers have been reported in *Streptomyces*. Ppm1 is an essential enzyme in mycobacteria and a *ppm1<sup>-</sup>* strain of *Corynebacterium* is growth retarded indicating the central role polyprenol phosphate mannose has in both organisms [10, 11]. The protein O-glycosylation pathway is present in most Actinobacteria and Pmt in Mtb has been shown to be important for virulence [12, 13]. In *Streptomyces coelicolor* other putative glycosyl transferases are also likely to use polyprenol phosphate mannose as a sugar donor and some of these are described in Howlett et al (submitted).

The role of polyprenol phosphate mannose in antibiotic resistance and the pathway leading to its synthesis is addressed further in this paper (Fig. 1). D-mannose is either taken up from the medium and converted by hexokinase to D-mannose-6-phosphate or the latter can be produced from D-fructose-6-phosphate by phosphomannoisomerase (ManA). Phosphomannomutase (ManB: PMM) then converts D-mannose-6-phosphate to D-mannose-1-phosphate which is a substrate

for GDP mannose pyrophosphorylase (ManC:GMP). In *Corynebacterium glutamicum* deletion of the *manC* homologue (NCgl0710) conferred retarded growth and loss of nearly all mannoglycans from the envelope [14]. This phenotype resembles that of the *ppm1* mutant of *C. glutamicum* [10] and suggests that the ManB, ManC pathway is responsible for the synthesis of GDP-mannose. We hypothesized that strains containing blocks in the pathway leading to the synthesis of GDP-mannose ought to be phenotypically similar to the *ppm1*<sup>-</sup> strains as they too will be deficient in polyprenol phosphate mannose. Here we analyzed the roles of three putative *manC* genes in the *S. coelicolor* genome and a *manB* gene. We show that both a GMP depleted strain and a strain lacking PMM do indeed have phenotypes reminiscent of the *ppm1*<sup>-</sup> mutants. The phenotype of the *S. coelicolor manB*<sup>-</sup> strains constructed here varied from that reported previously for a *manB*<sup>-</sup> strain [15, 16]. We conclude that GMP activity in *S. coelicolor* is provided by expression of two genes, *sco3039* and *sco4238*. Moreover both GMP and PMM activities are part of the same metabolic pathway leading to the synthesis of polyprenol phosphate mannose and ultimately to glycoprotein biosynthesis in *S. coelicolor*.

## Methods

### *DNA manipulations*

Chemically competent *E. coli* cells were prepared, stored and used in the transformation procedure as described previously [17]. Plasmid DNA extraction from *E. coli* was performed using a Spin Miniprep Kit following the protocol supplied by the manufacturer (QIAGEN). Cosmids were manipulated as described [18]. Restriction enzymes and T4 ligase were obtained from New England Biolabs (NEB) and used according to the manufacturer's instruction. Phusion<sup>®</sup> High-Fidelity DNA Polymerase

(NEB) was employed for PCR amplification. Primers used in the present study are listed (Table S1). In-fusion® HD cloning kit (Clontech) was used according to the protocol supplied by the manufacturer. DNA sequencing (Sanger) was outsourced to Source Bioscience.

#### *Plasmid, cosmid and strain constructions*

A list of plasmids and cosmids used in this work is provided (Table 1). Plasmid pRH01 was produced by cloning of the PCR amplified product from primers RH11 and RH12 and J1929 genomic DNA as template, into EcoRV digested vector pAV11b [19-21]. Plasmid pRH12 was produced by cloning the PCR product from primers RH91 and RH92 and J1929 template into NdeI digested pIJ10257. Plasmids pRH11 and pRH14 were produced similar to pRH12 but using *E.coli* DH5  $\alpha$  genomic DNA as template and primer pairs RH93/RH94, and RH140/RH141, respectively. Expression plasmids for *sco3039* (pRH06) and *sco4238* (pRH07) were produced through the ligation of XhoI and NdeI digested PCR products from primers pairs RH71/RH72, and RH73/RH74, respectively, and *S. coelicolor* J1929 template DNA, into XhoI and NdeI digested pET21a fusing both ORFs to an inframe C-terminal hexa-histidine tag. All constructs were confirmed as correct through Sanger sequencing performed by Source Bioscience.

The apramycin resistance markers within the Tn5062 transposon of StD8A.2.D12 and St1A8A.1.B09 were replaced with spectinomycin and hygromycin markers, respectively, using the REDIRECT methodology [18]. Cosmids were introduced into *S. coelicolor* J1929 by conjugation and resistant exconjugants were selected according to the marker on Tn5062 (apramycin, spectinomycin or hygromycin resistance). Those that had undergone a double cross over recombination event were identified initially as they lost the marker (kanamycin resistance) on the cosmid



vector backbone. Presence of the interrupted allele and loss of the wild type allele was confirmed by PCR and Southern blotting.

#### *Phage sensitivity assays*

Plaque assays were performed as described [22]. Briefly Difco nutrient agar supplemented with 10 mM MgSO<sub>4</sub> and 8 mM Ca (NO<sub>3</sub>)<sub>2</sub> were inoculated with dilutions of  $\phi$ C31  $\Delta c25$  (clear plaque) phage [23] and then overlaid with soft nutrient agar containing approximately  $1 \times 10^7$  spores of the desired test strain. The streak plate assay was performed using square 10 cm plates containing Difco nutrient agar (10 mM MgSO<sub>4</sub> and 8 mM Ca (NO<sub>3</sub>)<sub>2</sub>). One half of the plate was inoculated with 100  $\mu$ l of  $\phi$ C31  $\Delta c25$  (approx  $1 \times 10^8$  pfu/ml) and a single streak of the test spore preparation was inoculated across the plate beginning on the phage-free region. Plates were incubated at 30 °C.

#### *Protein expression*

An overnight culture of *E. coli* BL21DE3 (pRH07) in LB containing ampicillin was grown at 37 °C and used to inoculate 2YT, which was grown to OD 0.6. IPTG (0.15 mM) was then added to induce expression and the culture was further incubated (22 °C for 22 hours). The bacteria were harvested by centrifugation and resuspended in binding buffer (30 ml; 20 mM Tris HCl pH 7.4, 0.5M NaCl, 30 mM imidazole) and sonicated. The cell lysate was cleared by centrifugation (4 °C, 5 minutes, 10,000g) to remove unlysed cells and debris and then the supernatant was loaded onto a HiTrap Ni<sup>2+</sup> affinity column (AKTA Purifier). After washing with 2 column volumes of binding buffer the bound protein was eluted with a gradient of increasing imidazole concentration using the elution buffer (20 mM Tris HCl pH 7.4, 0.5M NaCl, 500 mM

imidazole). Pooled fractions were then loaded onto a desalting column to remove imidazole and eluted in 20 mM Tris HCl pH 7.4, 0.5M NaCl. The protein was concentrated using Vivaspin (GE healthcare) spin columns to approximately 10 mg/ml. Glycerol was added to a final concentration of 50% and aliquots were stored at -80 °C. Protein concentration was assayed using the BioRad protein assay solution and is based on the Bradford assay [24].

#### *GDP mannose pyrophosphorylase assays*

Activity was measured by monitoring the release of pyrophosphate using the EnzChek® pyrophosphate assay kit (Thermofisher). Briefly the kit includes a pyrophosphatase that catalyses the conversion of the pyrophosphate released from the GMP activity to two equivalents of phosphate, which is then used as a substrate in a reaction with 2-amino-6-mercapto-7-methylpurine ribonucleoside (MESG) and purine nucleoside phosphorylase (PNP) to release ribose 1-phosphate and 2-amino-6-mercapto-7-methyl-purine. The latter compound was detected spectrophotometrically by absorbance at 360 nm. Assays were performed according to the manufacturers instructions except that they were scaled down to enable use of a 96 well plate reader (200 µl assay volume per well). GMP activity rates were obtained using different nucleotides (1 mM ATP, GTP, CTP or dTTP) and sugars, (1 mM mannose-1-phosphate or mannose-6-phosphate) as substrates. Initial rates were calculated and plotted against substrate concentration using SIGMAplot.

## **Results**

### *Identification of putative GDP-mannose pyrophosphorylases*

GDP-mannose, a substrate for Ppm1, is synthesized by GDP-mannose pyrophosphorylase (GMP) encoded by *manC* (Fig. 1). BLAST searches of the *Streptomyces coelicolor* genome with the characterised *Corynebacterium glutamicum* GMP/ManC (encoded by NCgl0710) and *Mycobacterium tuberculosis* GMP/ManC (encoded by Rv3264c) identified SCO1388, SCO3039 and SCO4238 as putative GMP candidates (Fig. 2) [14]. The nucleotidyl transferase domains of all three *Streptomyces* GMP/ManC candidates contain the GXGXR<sub>n</sub>K signature motif of phosphorylases, and variations on the F(V)EKP motif characteristic of the GMP active site (Fig. 2) [25, 26]. SCO3039 and SCO1388 have protein domains in addition to the nucleotidyl transferase domain; SCO1388 in particular appears to be a bifunctional enzyme with both GMP and phosphomannomutase (ManB) activity (Fig. 2).

#### *SCO3039 and SCO4238 have overlapping functions*

The *S. coelicolor* genes *sco1388*, *sco3039* and *sco4238* were disrupted by allelic exchange with cosmids containing Tn5062 in the gene of interest to produce single insertion mutants RH501 (*sco1388::Tn5062*), SKA211 (*sco3039::Tn5062*) and SKA311 (*sco4238::Tn5062*). The cosmids were obtained from the transposon insertion cosmid library [27] (Table 1). Validated mutants were tested for  $\phi$ C31 resistance as a potential indicator for a loss of GMP/ManC activity due to a lack of protein O-glycosylation [5, 6], but all of the mutants were still sensitive to phage infection. A mild increase in blue pigment production was seen in the mutant strains SKA311 (*sco4238::Tn5062*) and SKA211 (*sco3039::Tn5062*) when grown on supplemented minimal media (SMM) (not shown).

217  
218 Double mutants were created to assess whether there is redundancy in gene  
219 function between *sco3039*, *sco4238* and *sco1388*. The cosmid StD8A.2.D12<sup>spec</sup>  
220 (*sco4238::Tn5062<sup>spec</sup>*) was introduced into J1929 by conjugation to create the  
221 spectinomycin resistant *sco4238* insertion mutant, RH25. The cosmid StE34.1.G05  
222 (*sco3039::Tn5062*) was then introduced into RH25 by conjugation, selecting for  
223 apramycin resistance. Only 8 exconjugants from several hundred that were screened  
224 had the spectinomycin-resistant, apramycin-resistant, kanamycin-sensitive  
225 phenotype indicative of a *sco4238::Tn5062<sup>spec</sup>*, *sco3039::Tn5062* double mutant.  
226 However subsequent analysis by polymerase chain reaction (PCR) to amplify the  
227 genomic region containing *sco3039* showed that this gene was uninterrupted in all  
228 eight candidate double mutant strains and mutations must have occurred elsewhere  
229 to confer resistance to apramycin. Thus we were unable to create a simple double  
230 mutant containing *Tn5062* insertions in both *sco3039* and *sco4238*, suggesting that  
231 these genes share an important function for growth. Multiple *sco4238::Tn5062<sup>spec</sup>*,  
232 *sco1388::Tn5062* and *sco3039::Tn5062*, *sco1388::Tn5062<sup>hyg</sup>* double mutant strains  
233 were produced and confirmed through kanamycin sensitivity. The phenotypes of  
234 these strains were no different from the individual *sco4238::Tn5062* and  
235 *sco3039::Tn5062* mutants, SKA311 and SKA211, respectively. The product of  
236 *sco1388* therefore probably contributes little to the total GMP activity in *S. coelicolor*.  
237

238 We were able to create a strain containing both *sco4238::Tn5062<sup>spec</sup>* and  
239 *sco3039::Tn5062* insertions in the presence of a conditionally expressed *sco4238*.  
240 Plasmid pRH01, encoding *sco4238* under the control of the anhydrotetracycline  
241 (ATC) inducible promoter, *tcp830* [20], was introduced into RH25 to create strain

242 RH221 (*sco4238::Tn5062<sup>spec</sup>*, *tcp830-sco4238*, *hyg*). Conjugation of StE34.1.G05  
243 (*sco3039::Tn5062*) into RH221 in the presence of ATC resulted in multiple *spec<sup>R</sup>*,  
244 *apra<sup>R</sup>*, *hyg<sup>R</sup>*, *kan<sup>S</sup>* exconjugants (RH2213) that were subsequently confirmed as  
245 *sco4238::Tn5062<sup>spec</sup>*, *sco3039::Tn5062* double mutants through PCR. Surprisingly  
246 RH2213 could grow in the absence of ATC, an observation that was at odds with our  
247 inability to isolate the transposon double mutants in the absence of pRH01. Colony  
248 sizes of the RH2213 strains in the absence of ATC were indistinguishable from the  
249 wild type parent strain, J1929, but a significant increase in blue pigments were  
250 observed compared to the single mutants RH25 (*sco4238::Tn5062<sup>spec</sup>*) and SKA211  
251 (*sco3039::Tn5062*)(Fig. 2B). The *tcp830* promoter has been shown by others to be  
252 incompletely turned off in the absence of ATC and we propose that this is also the  
253 case in our experiments [19]. It seems likely that RH2213 grown in the absence of  
254 ATC has a depleted level of GMP compared to the parent strain and compared to  
255 RH2213 grown in the presence of ATC.

256

257

258 *Strains depleted in the putative GMPs SCO3039 and SCO4238 are hyper*  
259 *susceptible to antibiotics and partially resistant to  $\phi$ C31*

260

261 Ppm1 uses GDP-mannose as a substrate and we therefore hypothesized that  
262 inability to synthesize GDP-mannose, for example through GMP depletion, should  
263 result in a similar phenotype to those strains deficient in Ppm1. RH2213 isolates  
264 (*sco4239::Tn5062<sup>spec</sup>*, *sco3039::Tn5062*, pRH01 encoding inducible *sco4238*) were  
265 still able to support  $\phi$ C31 plaque formation but displayed resistance to  $\phi$ C31 on a  
266 streak assay in the absence of ATC (Fig 3A).

267

268 We then tested the putative GMP deficient strains for their susceptibilities to  
269 antibiotics, notably those to which the *ppm1* and *pmt* mutants were particularly  
270 sensitive. *S. coelicolor* strains SKA211 and RH25 containing Tn5062 insertions in  
271 *sco3039* and *sco4238*, respectively, had the same antibiotic resistances as the  
272 parent strain J1929. However RH2213, with depleted levels of GMP in the absence  
273 of ATC, was highly susceptible to antibiotics, strongly resembling the phenotypes of  
274 the *ppm1* mutants (Fig. 3B). The phenotypes of the GMP depleted mutants indicate  
275 that *sco3039* and *sco4238* provide the majority of the GMP activity in *S. coelicolor*.

276

277 *SCO4238 encodes a highly specific GDP-mannose pyrophosphorylase activity*

278

279 To confirm the phenotypes mentioned above were due to a depletion of GMP activity  
280 in RH2213, *sco4238* and *sco3039* were overexpressed in *E. coli* in order to assay  
281 GMP activity on purified proteins. Overproduced SCO4238 showed high GMP activity  
282 (Fig. 4). The enzyme was highly specific for GTP and D-mannose-1-phosphate  
283 substrates, with no or very low rates achieved with CTP, ATP and dTTP (not shown).  
284 Approximately 50% activity was observed with D-mannose-6-phosphate and GTP,  
285 with the Hill coefficient showing a loss of the cooperativity seen with D-mannose-1-  
286 phosphate. In *Mycobacterium tuberculosis* the essential enzyme, RmlA, catalyses  
287 the synthesis of dTDP-glucose, an intermediate in dTDP-rhamnose biosynthesis  
288 required for the integrity of the cell wall [28]. Given the apparent essentiality of GMP  
289 in *S. coelicolor* we tested whether SCO4238 had activity on glucose-1-phosphate in  
290 combination with any nucleotide, including dTTP but no activity was detected.

291 Attempts to obtain soluble, active SCO3039 from several overexpression constructs  
292 in *E. coli* failed (not shown).

293

294 *Strains with a mutation in the manB gene, sco3028 are also phenotypically similar to*  
295 *the ppm1 mutants*

296 Previous work has shown that SCO3028 is a dual functioning enzyme capable of  
297 phosphomannomutase (PMM, mannose-6-phosphate to mannose-1-phosphate) and  
298 phosphoglucomutase (PGM, glucose-6-phosphate to glucose-1-phosphate) activity  
299 [16]. The authors constructed a *manB* deletion mutant,  $\Delta manB$ , which had increased  
300 actinorhodin production and had lost chloramphenicol resistance but displayed  
301 apparently similar growth to the parent strain, M145 [15, 16]. Both phenotypes of the  
302  $\Delta manB$  strain were complemented when wild type *E. coli manB* but not *S. coelicolor*  
303 *pgm* gene (*sco7443*) were introduced. Thus PMM activity was shown to be solely  
304 responsible for an increase in chloramphenicol sensitivity and actinorhodin  
305 production in *S. coelicolor* M145. If SCO3028 is the sole PMM enzyme in *S.*  
306 *coelicolor* we would expect a similar phenotype in the *sco3028* mutant as we see for  
307 the GMP depleted strains. However, Yang *et al* did not detect increased susceptibility  
308 of their  $\Delta manB$  strain to vancomycin, bacitracin or ampicillin [16].

309

310 In order to assess the phenotype of an *sco3028* mutant in our  $\phi$ C31 sensitive strain  
311 *S. coelicolor* J1929, a *pglY*<sup>-</sup> derivative of M145 [29], the cosmid StE34.1.B03  
312 (*sco3028::Tn5062*) was introduced into J1929 by conjugation. Exconjugants that had  
313 undergone a double crossover (RHB42 strains, validated by PCR) were isolated at  
314 low frequency and had an extreme small (XS) colony phenotype, even smaller than  
315 the colony size seen in the *ppm1* mutant DT3017 (Fig. 5). The XS colony phenotype

in RHB42 could be fully restored to wild type through complementation with *S. coelicolor sco3028 (manB)*, and *Escherichia coli manB (cpsG)* as observed in strains RHB4212 and RHB4211, respectively. RHB42 containing *Escherichia coli pgm*, encoding phosphoglucomutase, was capable of partially restoring colony size (RHB4214), suggesting it is the loss of both PMM and PGM activity that had resulted in the XS colony phenotype in RHB42.

RHB42 was highly resistant to  $\phi$ C31 infection in a plaque assay, resembling phage resistance in the *pmt* and *ppm1* mutants (Fig. 5A). RHB42 was also highly susceptible to a number of cell wall acting antibiotics, as well as the RNA polymerase targeting antibiotic, rifampicin (Fig. 5B). Phage sensitivity and antibiotic resistance were restored to wild type in RHB4212 (*sco3028*<sup>+</sup>) and RHB4211 (*cpsG*<sup>+</sup>) but not RHB4214 (*pgm*<sup>+</sup>). No difference in chloramphenicol resistance between RHB42 and J1929 was observed. These phenotypes are consistent with SCO3028 being the primary PMM in *S. coelicolor* and in the same metabolic pathway that synthesizes polyprenol phosphate mannose.

An increase in pigment production was recorded in RHB42, similar to that seen previously following *sco3028* deletion [16] and that seen in the GMP depleted strain, RH2213 (Fig. 5C). The production of blue pigment was reduced to wild type level in RHB4212 (*sco3028*<sup>+</sup>) and RHB4211 (*cpsG*<sup>+</sup>) but not RHB4214 (*pgm*<sup>+</sup>) (Fig. 5). To further validate our observations (as they differ from those of Yang et al, [15, 16]), we created two more sets of *sco3028::Tn5062* mutants: First we used a different Tn5062 insertion in J1929 using cosmid, STE34.2.D03, generating strain JD182 and second we generated derivatives of M145 containing the Tn5062 insertions from both



STE34.1.B03 and STE34.2.D03 to generate strains MD202 and MB92, respectively.  
All three strains had an identical phenotype to RHB42 (Figs S1 and S2).

## Discussion

Mannose is a component of cell envelope polymers including mannlipids, phosphoinositol mannosides (PIMs) and glycoproteins in many bacteria [9, 30-32]. Extracytoplasmic glycosyl transferases use polyprenol phosphate mannose as the lipid linked sugar donor in the biosynthesis of mannose containing polymers [4, 33]. The synthesis of polyprenol phosphate mannose by Ppm1 is therefore an important activity and *ppm1* mutants are considerably less fit than the parent strains [10, 11]. In the case of *S. coelicolor*, *ppm1*<sup>-</sup> mutants have a small colony growth phenotype and are hyper-susceptible to multiple antibiotics, most of which inhibit cell wall biogenesis suggesting that these mutants are pleiotropically deficient in membrane and/or periplasmic function (Howlett et al, submitted). Mutants lacking Ppm1 or Pmt are also resistant to phage infection and we have proposed that  $\phi$ C31 uses a glycoprotein(s) as its receptor [5, 6]. We show here that depletion of enzymes in the mannose metabolism pathway prior to Ppm1 display a phenotype that resembles that of the *ppm1*<sup>-</sup> mutants. We conclude that synthesis of polyprenol phosphate mannose and its subsequent role as a mannose donor in the periplasm is required for a wild type antibiotic resistant phenotype.

Although the overall phenotypes of the GDP-mannose pyrophosphorylase (GMP) deficient and the *manB* mutant strains resembled the *ppm1*<sup>-</sup> strain there were some minor differences. We had difficulty in generating a GMP deficient strain. *S. coelicolor* has three candidate genes that could express GMP activity and we could only obtain

a double *sco3039*<sup>-</sup>, *sco4238*<sup>-</sup> mutant if *sco4238* was expressed conditionally using the anhydrotetracycline-inducible *tcp830* promoter. While this is not absolute proof that the GMP activity is essential in *S. coelicolor*, it would seem that some low level of activity, possibly that provided by the leakiness of the repressed *tcp830* promoter reading into an integrated copy of *sco4238*, is required for the simultaneous interruption of both *sco3039* and *sco4238* by Tn5062. Similarly the insertion in *sco3028* (*manB*) was obtained at very low frequency and the colonies were extremely small, indicative of a requirement for both phosphomannomutase (PMM) and phosphoglucomutase (PGM) activities encoded by this gene. If GMP activity is essential then we would also expect PMM activity to be essential, but there may be sufficient PMM activity from other closely related enzymes (such as other PGM paralogues; *sco7443* or *sco4916* a possible alternative phosphomannomutase) to allow growth. The more severe phenotypes of the GMP depletion mutant and the *manB* mutant also suggest that GDP-mannose could be located at a metabolic branch point i.e. GDP-mannose is required for polyprenol phosphate mannose synthesis but also perhaps for modification into other nucleotide sugars. For example, the *S. coelicolor* genome encodes a GDP-mannose dehydrogenase (SCO0382) that is predicted to make GDP-mannuronate, one of the building blocks in the synthesis of alginates in *Pseudomonas*. *Sco0382* lies within an operon *sco381* to *sco386* that has features of an extracellular polysaccharide biosynthesis gene cluster including a polyprenol dependent glycosyl transferase and various other membrane proteins.

Enzyme assays with purified SCO4238 showed it to be a monofunctional GMP (ManC) with a slim substrate tolerance similar to Rv3264 (previously miss-annotated

as *rmlA*) of *Mycobacterium* [28, 34] and in contrast to the more promiscuous ManC enzymes of *E. coli* and *P. furiosus* [35, 36]. In *Mycobacterium* and in *Corynebacterium* the ManC enzymes (Rv3264 and NCgl0710, respectively) provide essential supplies of GDP-mannose for phosphatidyl inositol mannoside (PIM) biosynthesis and lipoglycans [14].

The phenotype described here for RHB42 (*sco3028:Tn5062*, *manB*) has differences and similarities to a  $\Delta manB$  strain of *S. coelicolor* M145 that was described previously [15, 16]. A notable difference is the sensitivity to antibiotics of RHB42 as Yang *et al* did not detect an increase in susceptibility of their  $\Delta manB$  strain to vancomycin, bacitracin or ampicillin [16]. Whilst we cannot explain these differences in phenotypes, both RHB42 and the  $\Delta manB$  of Yang *et al* have increased pigment production. Pigment production is also greatly increased in the ManC deficient strains. The increase in pigment production could be indicative of the activation of several stress pathways or, as discussed by Yang *et al*, could be due to the increase in carbon flux through glycolysis as the pathway to GDP-mannose is blocked. The *ppm1<sup>-</sup>* strain DT3017 has a mild pigment overproduction phenotype (data not shown). Neither Yang *et al*, or Rajesh *et al* could test phage sensitivity in their  $\Delta manB$  strain as they used a Pgl<sup>+</sup> strain of *S. coelicolor*, which confers  $\phi$ C31 resistance.

Mannose is used in other *Streptomyces spp* in the biosynthesis of antibiotics eg mannopeptimycins and amphotericin [37, 38]. ManB and ManC activities are required in *S. nodosus* for the glycosylation of amphotericin [38]. The identification of the *manC* genes and the construction of the *manC* deficient strain could be useful in

heterologous expression and combinatorial biosynthesis of several antibiotic pathways in *S. coelicolor*.

## Acknowledgements

We are grateful to Professor Mervyn Bibb for providing plasmid pIJ10257 and to Professor Paul Dyson for the provision of mutagenized cosmids.

## Funding Information

This work was funded by project grants\_BB/J016691 and BBS/B/05990 from the Biotechnology and Biological Sciences Research Council, UK.

## Conflicts of Interest and Ethics

None

## References

1. **Wang W, Ji J, Li X, Wang J, Li S et al.** Angucyclines as signals modulate the behaviors of *Streptomyces coelicolor*. *Proc Natl Acad Sci U S A* 2014;111(15):5688-5693.
2. **Nesme J, Simonet P.** The soil resistome: a critical review on antibiotic resistance origins, ecology and dissemination potential in telluric bacteria. *Environmental microbiology* 2015;17(4):913-930.
3. **D'Costa VM, McGrann KM, Hughes DW, Wright GD.** Sampling the antibiotic resistome. *Science* 2006;311(5759):374-377.
4. **Wehmeier S, Varghese AS, Gurcha SS, Tissot B, Panico M et al.** Glycosylation of the phosphate binding protein, PstS, in *Streptomyces coelicolor* by a pathway that resembles protein O-mannosylation in eukaryotes. *Mol Microbiol* 2009;71(2):421-433.
5. **Cowlshaw DA, Smith MCM.** Glycosylation of a *Streptomyces coelicolor* A3(2) cell envelope protein is required for infection by bacteriophage  $\phi$ C31. *Mol Microbiol* 2001;41(3):601-610.
6. **Cowlshaw DA, Smith MCM.** A gene encoding a homologue of dolichol phosphate-b-D mannose synthase is required for infection of *Streptomyces coelicolor* A3(2) by phage  $\phi$ C31. *J Bacteriol* 2002;184(21):6081-6083.

7. **Hoischen C, Gura K, Luge C, Gumpert J.** Lipid and fatty acid composition of cytoplasmic membranes from *Streptomyces hygroscopicus* and its stable protoplast-type L form. *J Bacteriol* 1997;179(11):3430-3436.
8. **Sandoval-Calderon M, Geiger O, Guan Z, Barona-Gomez F, Sohlenkamp C.** A eukaryote-like cardiolipin synthase is present in *Streptomyces coelicolor* and in most actinobacteria. *J Biol Chem* 2009;284(26):17383-17390.
9. **Guerin ME, Kordulakova J, Alzari PM, Brennan PJ, Jackson M.** Molecular basis of phosphatidyl-myo-inositol mannoside biosynthesis and regulation in mycobacteria. *J Biol Chem* 2010;285(44):33577-33583.
10. **Gibson KJ, Eggeling L, Maughan WN, Krumbach K, Gurucha SS et al.** Disruption of Cg-Ppm1, a polyprenyl monophosphomannose synthase, and the generation of lipoglycan-less mutants in *Corynebacterium glutamicum*. *J Biol Chem* 2003;278(42):40842-40850.
11. **Rana AK, Singh A, Gurucha SS, Cox LR, Bhatt A et al.** Ppm1-encoded polyprenyl monophosphomannose synthase activity is essential for lipoglycan synthesis and survival in mycobacteria. *PloS one* 2012;7(10):e48211.
12. **Liu CF, Tonini L, Malaga W, Beau M, Stella A et al.** Bacterial protein-O-mannosylating enzyme is crucial for virulence of *Mycobacterium tuberculosis*. *P Natl Acad Sci USA* 2013;110(16):6560-6565.
13. **Mahne M, Tauch A, Puhler A, Kalinowski J.** The *Corynebacterium glutamicum* gene pmt encoding a glycosyltransferase related to eukaryotic protein-O-mannosyltransferases is essential for glycosylation of the resuscitation promoting factor (Rpf2) and other secreted proteins. *FEMS Microbiol Lett* 2006;259(2):226-233.
14. **Mishra AK, Krumbach K, Rittmann D, Batt SM, Lee OY et al.** Deletion of manC in *Corynebacterium glutamicum* results in a phospho-myo-inositol mannoside- and lipoglycan-deficient mutant. *Microbiology* 2012;158(Pt 7):1908-1917.
15. **Rajesh T, Song E, Lee BR, Park SH, Jeon JM et al.** Increased sensitivity to chloramphenicol by inactivation of *manB* in *Streptomyces coelicolor*. *Journal of microbiology and biotechnology* 2012;22(10):1324-1329.
16. **Yang YH, Song E, Park SH, Kim JN, Lee K et al.** Loss of phosphomannomutase activity enhances actinorhodin production in *Streptomyces coelicolor*. *Applied microbiology and biotechnology* 2010;86(5):1485-1492.
17. **Green RM, Sambrook J.** *Molecular Cloning: A laboratory Manual*, Fourth Edition ed. Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press; 2012.
18. **Gust B, Challis GL, Fowler K, Kieser T, Chater KF.** PCR-targeted *Streptomyces* gene replacement identifies a protein domain needed for biosynthesis of the sesquiterpene soil odor geosmin. *Proc Natl Acad Sci U S A* 2003;100(4):1541-1546.
19. **Jyothikumar V, Klanbut K, Tiong J, Roxburgh JS, Hunter IS et al.** Cardiolipin synthase is required for *Streptomyces coelicolor* morphogenesis. *Mol Microbiol* 2012;84(1):181-197.
20. **Rodriguez-Garcia A, Combes P, Perez-Redondo R, Smith MCM.** Natural and synthetic tetracycline-inducible promoters for use in the antibiotic-producing bacteria *Streptomyces*. *Nucleic Acids Res* 2005;33(9):e87.
21. **Fayed B, Younger E, Taylor G, Smith MC.** A novel *Streptomyces* spp. integration vector derived from the *S. venezuelae* phage, SV1. *BMC biotechnology* 2014;14:51.
22. **Kieser T, Bibb MJ, Buttner MJ, Chater KF, Hopwood DA.** *Practical Streptomyces Genetics*. Norwich: The John Innes Foundation; 2000.

23. **Sinclair RB, Bibb MJ.** The repressor gene (c) of the *Streptomyces* temperate phage  $\phi$ C31: nucleotide sequence, analysis and functional cloning. *Mol Gen Genet* 1988;213(2-3):269-277.
24. **Bradford MM.** A rapid and sensitive method for the quantitation of microgram quantities of protein, utilising the principal of protein-dye binding. *Anal Biochem* 1976;72:248-254.
25. **Pelissier MC, Lesley SA, Kuhn P, Bourne Y.** Structural insights into the catalytic mechanism of bacterial guanosine-diphospho-D-mannose pyrophosphorylase and its regulation by divalent ions. *J Biol Chem* 2010;285(35):27468-27476.
26. **Sousa SA, Moreira LM, Leitao JH.** Functional analysis of the *Burkholderia cenocepacia* J2315 BceAJ protein with phosphomannose isomerase and GDP-D-mannose pyrophosphorylase activities. *Applied microbiology and biotechnology* 2008;80(6):1015-1022.
27. **Fernandez-Martinez LT, Del Sol R, Evans MC, Fielding S, Herron PR et al.** A transposon insertion single-gene knockout library and new ordered cosmid library for the model organism *Streptomyces coelicolor* A3(2). *Antonie Van Leeuwenhoek* 2011;99(3):515-522.
28. **Ma Y, Stern RJ, Scherman MS, Vissa VD, Yan W et al.** Drug targeting *Mycobacterium tuberculosis* cell wall synthesis: genetics of dTDP-rhamnose synthetic enzymes and development of a microtiter plate-based screen for inhibitors of conversion of dTDP-glucose to dTDP-rhamnose. *Antimicrob Agents Chemother* 2001;45(5):1407-1416.
29. **Bedford DJ, Laity C, Buttner MJ.** Two genes involved in the phase-variable  $\phi$ C31 resistance mechanism of *Streptomyces coelicolor* A3(2). *J Bacteriol* 1995;177(16):4681-4689.
30. **Dobos KM, Khoo KH, Swiderek KM, Brennan PJ, Belisle JT.** Definition of the full extent of glycosylation of the 45-kilodalton glycoprotein of *Mycobacterium tuberculosis*. *J Bacteriol* 1996;178(9):2498-2506.
31. **Michell SL, Whelan AO, Wheeler PR, Panico M, Easton RL et al.** The MPB83 antigen from *Mycobacterium bovis* contains O-linked mannose and (1 $\rightarrow$ 3)-mannobiose moieties. *J Biol Chem* 2003;278(18):16423-16432.
32. **Mishra AK, Driessen NN, Appelmek BJ, Besra GS.** Lipoarabinomannan and related glycoconjugates: structure, biogenesis and role in *Mycobacterium tuberculosis* physiology and host-pathogen interaction. *FEMS Microbiol Rev* 2011;35(6):1126-1157.
33. **Besra GS, Morehouse CB, Rittner CM, Waechter CJ, Brennan PJ.** Biosynthesis of mycobacterial lipoarabinomannan. *J Biol Chem* 1997;272(29):18460-18466.
34. **Ning B, Elbein AD.** Purification and properties of mycobacterial GDP-mannose pyrophosphorylase. *Arch Biochem Biophys* 1999;362(2):339-345.
35. **Mizanur RM, Pohl NL.** Phosphomannose isomerase/GDP-mannose pyrophosphorylase from *Pyrococcus furiosus*: a thermostable biocatalyst for the synthesis of guanidinediphosphate-activated and mannose-containing sugar nucleotides. *Organic & biomolecular chemistry* 2009;7(10):2135-2139.
36. **Yang YH, Kang YB, Lee KW, Lee TH, Park SS et al.** Characterization of GDP-mannose pyrophosphorylase from *Escherichia coli* O157 : H7 EDL933 and its broad substrate specificity. *J Mol Catal B-Enzym* 2005;37(1-6):1-8.
37. **Magarvey NA, Haltli B, He M, Greenstein M, Hucul JA.** Biosynthetic pathway for mannopeptimycins, lipoglycopeptide antibiotics active against drug-

resistant gram-positive pathogens. *Antimicrob Agents Chemother* 2006;50(6):2167-2177.

38. **Nic Lochlainn L, Caffrey P.** Phosphomannose isomerase and phosphomannomutase gene disruptions in *Streptomyces nodosus*: impact on amphotericin biosynthesis and implications for glycosylation engineering. *Metabolic engineering* 2009;11(1):40-47.

39. **Hong HJ, Hutchings MI, Hill LM, Buttner MJ.** The role of the novel Fem protein Vank in vancomycin resistance in *Streptomyces coelicolor*. *J Biol Chem* 2005;280(13):13055-13061.

40. **MacNeil DJ.** Characterization of a unique methyl-specific restriction system in *Streptomyces avermitilis*. *J Bacteriol* 1988;170(12):5607-5612.

## Legends to Figures

**Figure 1. The GDP-mannose biosynthesis pathway in *Streptomyces coelicolor*.**

**Figure 2. GDP-mannose pyrophosphorylases in *Streptomyces coelicolor* A.** Domain structures for *S. coelicolor* genes with putative GDP-mannose phosphorylase activity. B. Pigment overproduction in *sco4238*, *sco3039* double mutants. Strain RH2213 (*sco4239::Tn5062spec*, *sco3039::Tn5062*, pRH01 encoding inducible *sco4238*) overproduced blue pigment on supplemented minimal medium solid (SMMS) agar in the absence of anhydrotetracycline (ATC) but not in the presence of 0.5  $\mu$ g/ml ATC.

**Figure 3. *Streptomyces coelicolor* strains depleted in GDP mannose pyrophosphorylase activity are partially resistant to  $\phi$ C31 and are hyper-susceptible to some antibiotics A.** Spores of the indicated *S. coelicolor* strains were streaked from an area free from  $\phi$ C31 to an area inoculated with  $1 \times 10^7$  pfu  $\phi$ C31 on Difco nutrient agar plates with or with the supplementation of 0.5  $\mu$ g/ml anhydrotetracycline (ATC). RH2213 (*sco4239::Tn5062<sup>spec</sup>*, *sco3039::Tn5062*, pRH01

encoding inducible *sco4238*) showed conditional phage resistance growing only in the absence of ATC. For comparison the phage resistant phenotype of the *ppm1*<sup>-</sup> mutant (DT3017), the parent strain (J1929) and the strains with single mutations in the *manC* candidate genes, *sco3039* and *sco4238*, (SKA211 and RH25, respectively). B. RH2213 showed increased susceptibility to antibiotics in the absence of ATC but not in the presence of ATC. This phenotype is comparable to the antibiotic hyper-susceptible phenotype of the *ppm1*<sup>-</sup> mutant (DT3017). Results show the mean diameter of the disc diffusion inhibition zones from at least 3 replicates. Antibiotics were all used at 4  $\mu$  g/disc with the exception of ampicillin that was used at 40  $\mu$  g/disc.

**Figure 4. Kinetics of GDP-mannose pyrophosphorylase activity of SCO4238.**

Dependence of initial velocity of SCO4238 GMP activity with increasing concentration of mannose-1-phosphate (A) and mannose-6-phosphate (B). C. Kinetic parameters  $K_m$ ,  $n_H$  and  $V_{max}$  for the two sugar phosphate substrates.

**Figure 5. Phenotypes of the *manB* (*sco3028::Tn5062*) mutants.**

A. Extreme small colony phenotype (left) and resistance to phage  $\phi$ C31 (right) in the RHB42 (*sco3028::Tn5062*) mutant strains. The poor growth of RHB42 was complemented with an additional copy of *sco3028* (RHB4212) or the *E. coli* gene *cpsG* (*manB* homologue; RHB4211) but not *pgm* from *E. coli* (encoding phosphoglucomutase; RHB4214). The  $\phi$ C31 resistance in RHB42 reverted to phage sensitivity in the complemented strains RHB4211 and RHB4212. B. RHB42 has increased susceptibility to antibiotics resembling the phenotype of the *ppm1*<sup>-</sup> mutant (DT3017). Antibiotic susceptibility could be complemented with an additional copy of *sco3028*



605 and *E. coli cpsG* but not *pgm* from *E. coli*. C. Increased pigment production in RH42  
606 compared to the parental strain, J1929. This phenotype was complemented by  
607 additional copy of *sco3028* and *E. coli cpsG* but not *pgm* from *E. coli*.

608

609

610

611 Table 1. Bacteria, plasmids and cosmids

Plasmid Name	Description	References
pAVIIb	Integrating vector with <i>tcp830</i> promoter and <i>Tetris</i> cassette	[21]
pRH01	<i>sco4238</i> in pAV11b	This study
pDT16	<i>sco1423 (ppm1)</i> in pSET152	[6]
pDT10	<i>sco3154 (pmt)</i> in pSET152	[5]
pET21a	Overexpression vector containing HIS <sub>6</sub> -tag, T7 promotor	Merck Chemicals
pRH06	<i>sco3039</i> in pET21a	This study
pRH07	<i>sco4238</i> in pET21a	This study
pIJ10257	Integrating vector with constitutive promoter <i>ermEp*</i>	[39]
pRH11	<i>cpsG</i> in pIJ10257	This study
pRH12	<i>sco3028</i> in pIJ10257	This study
pRH14	<i>pgm</i> in pIJ10257	This study
Cosmid	Description	References
St1A8A.1.B09	SCO1388:: <i>Tn5062</i> at nt 572	[27]
StD8A.2.D12	SCO4238:: <i>Tn5062</i> at nt 69	[27]
StE34.1.G05	SCO3039:: <i>Tn5062</i> at nt 155	[27]
StE34.1.B03	SCO3028:: <i>Tn5062</i> at nt 590	[27]
StD8A.2.D12 <sup>spec</sup>	StD8A.2.D12 with <i>sco4238</i> :: <i>Tn5062</i> <sup>spec</sup>	This study
St1A8A.1.B09 <sup>hyg</sup>	St1A8A.1.B09 with <i>sco1388</i> :: <i>Tn5062</i> <sup>hyg</sup>	This study
<i>Streptomyces</i> strain	Genotype	References
M145	Prototroph	[22]
J1929	<i>pglY</i> mutant	[29]
DT3017	<i>ppm1</i> <sup>E218V</sup> mutant	[6]
DT1020	<i>ppm1</i> <sup>H116D</sup> mutant	[6]
DT1029	<i>ppm1</i> <sup>S163L</sup> mutant	[6]
DT1025	<i>pmt</i> , frameshift from A121	[5]
DT2008	<i>pmt</i>	[5]

SKA211	<i>sco3039::Tn5062</i>	This study
SKA311	<i>sco4238::Tn5062</i>	This study
RH501	<i>sco1388::Tn5062</i>	This study
RH25	<i>sco4238::Tn5062<sup>spec</sup></i>	This study
RH221	<i>sco4238::Tn5062<sup>spec</sup></i> , pRH01	This study
RH2213	<i>sco4238::Tn5062<sup>spec</sup> sco3039::Tn5062</i> , pRH01	This study
RHB42	<i>sco3039::Tn5062</i>	This study
RHB4211	<i>sco3028::Tn5062</i> , pRH11	This study
RHB4212	<i>sco3028::Tn5062</i> , pRH12	This study
RHB4214	<i>sco3028::Tn5062</i> , pRH14	This study
<b><i>E.coli</i> strain</b>	<b>Genotype</b>	<b>References</b>
ET12567(pUZ800 2)	<i>dam- dcm- hsdS</i> , RP4 transfer genes	[40]
BL21 (DE3)	<i>lon, ompT, gal, hsdS</i> , □DE3	[17]
DH5α	F- Φ80 <i>lacZ</i> Δ M15 Δ ( <i>lacZYA-argF</i> ) U169 <i>recA1 endA1 hsdR17</i> (rk-, mk+) <i>phoA supE44 λ -thi-1 gyrA96 relA1</i>	[17]

612

613




Figure 1





Figure 2

A



B




Figure 3




Figure 4




Figure 5






Figure S1




Figure S2

## Supplementary Figure Legends. Howlett *et al.*

**Figure S1. Extreme small colony phenotype of various *manB*<sup>-</sup> strains of *S. coelicolor*.** MD202 is a derivative of *S. coelicolor* M145 containing *sco3028::Tn5062* at nt 590 derived from cosmid StE34.1.B03 (1, 2). MB92 and JD182 are derivatives of *S. coelicolor* M145 and J1929, respectively, containing *sco3028::Tn5062* at nt 576 derived from cosmid StE34.2.D03 (1, 2). In each case a transposon insertion in *sco3028*, (encoding ManB) confers an extreme small colony phenotype. The phenotype can be fully complemented by introduction of a wild type copy of *sco3028* (*sco3028*<sup>+</sup> in pRH12), *E. coli cpsG* (encoding ManB in pRH11) but not by *E. coli pgm* (encoding phosphoglucomutase, pRH14).

**Figure S2. Antibiotic hyper-susceptible phenotype of various *manB*<sup>-</sup> strains of *S. coelicolor*.** Strains are as described in the legend to Figure S2. For comparison the antibiotic susceptibility profiles of the parent strains J1929 and M145 and that of the *ppm1*<sup>-</sup> strain DT3017 are also shown.

## References

1. Bishop A, Fielding S, Dyson P, Herron P. Systematic insertional mutagenesis of a streptomycete genome: a link between osmoadaptation and antibiotic production. *Genome Res.* 2004;14(5):893-900.
2. Fernandez-Martinez LT, Del Sol R, Evans MC, Fielding S, Herron PR, Chandra G, et al. A transposon insertion single-gene knockout library and new ordered cosmid library for the model organism *Streptomyces coelicolor* A3(2). *Antonie Van Leeuwenhoek.* 2011;99(3):515-22.

Table S1. Oligonucleotides for Howlett, Read *et al*,

Primer	Sequence	Use
RH11	TCTAGAGACGTCGATATCATGACCGATCCGAACGCCGCGTCC	<i>sco4238</i> F Infusion; pRH01
RH12	CATGCATGATCAGATATCTCAGCGGCCGGACAGGGCCG	<i>sco4238</i> R Infusion; pRH01
RH71	GACGCCCATATGACAGAAGCGATCCTCCTG	<i>sco3039</i> (NdeI); pRH06
RH72	CGTTACAAGCTTCGTGTCCGGTGAGAAGCG	<i>sco3039</i> (HindIII); pRH06
RH73	GGCGCCCATATGACCGATCCGAACGCCGCGTC	<i>sco4238</i> (NdeI); pRH07
RH74	GTACCTCGAGGCGGCCGGACAGGGCCG	<i>sco4238</i> (XhoI); pRH07
RH91	ACAGGAGGCCCCATATGGCTGCTGATCTGTGCGCAG	<i>sco3028</i> F Infusion; pRH12
RH92	ACTCGAGATCTCATATGTCACGCCCGGATGATCGCC	<i>sco3028</i> R Infusion; pRH12
RH93	ACAGGAGGCCCCATATGAAAAAATTAACCTGCTTT	<i>cpsG(manB)</i> F Infusion; pRH11
RH94	ACTCGAGATCTCATATGTTACTCGTTCAGCAACG	<i>cpsG(manB)</i> R Infusion; pRH11
RH140	ACAGGAGGCCCCATATGGCAATCCACAATCGTGC	<i>pgm</i> F Infusion; pRH14
RH141	ACTCGAGATCTCATATGTTACGCGTTTTTCAGAA	<i>pgm</i> R Infusion; pRH14