



This is a repository copy of *Associations Between Serum Bone Biomarkers in Early Breast Cancer and Development of Bone Metastasis: Results From the AZURE (BIG01/04) Trial*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/128376/>

Version: Supplemental Material

Article:

Brown, J., Rathbone, E., Hinsley, S. et al. (13 more authors) (2018) Associations Between Serum Bone Biomarkers in Early Breast Cancer and Development of Bone Metastasis: Results From the AZURE (BIG01/04) Trial. *JNCI: Journal of the National Cancer Institute*, 110 (8). pp. 871-879. ISSN 0027-8874

<https://doi.org/10.1093/jnci/djx280>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.


Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.




eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Supplementary Figure 1. Forest plot of Invasive Disease Free Survival (IDFS) treatment hazard ratios and 95% confidence intervals (CIs) for all patients in the AZURE study (black) and patients in the biomarker population (blue). P-values were calculated using the likelihood ratio χ^2 test statistic and tests were performed at the two-sided 5% significance level



Supplementary Figure 2. χ^2 values from adjusted Cox proportional hazards model, analysing bone metastasis at any time by CTX, with differing high vs. normal CTX cut-points. Optimum cut-point observed at 0.2 ng/ml with a corresponding p-value of 0.01. P-values were calculated using the likelihood χ^2 test statistic and tests were performed at the two-sided 5% significance level.



Supplementary Figure 3. χ^2 values from adjusted Cox proportional hazards model, analysing bone metastasis at any time by 1-CTP, with differing high vs. normal 1-CTP cut-points. Optimum cut-point observed at 3.7 ng/ml with a corresponding p-value of 0.01. P-values were calculated using the likelihood χ^2 test statistic and tests were performed at the two-sided 5% significance level.

