



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/128698/>

Version: Accepted Version

Article:

Chalendar, I, Esterle, J and Partington, JR (2018) Estimates near the origin for functional calculus on analytic semigroups. *Journal of Functional Analysis*, 275 (3). pp. 698-711.
ISSN: 0022-1236

<https://doi.org/10.1016/j.jfa.2018.03.012>

(c) 2018 Elsevier Inc. All rights reserved. Licensed under the Creative Commons Attribution-Non Commercial No Derivatives 4.0 International License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Estimates near the origin for functional calculus on analytic semigroups

I. Chalendar*, J. Esterle† and J.R. Partington‡

January 9, 2018

Abstract

This paper provides sharp lower estimates near the origin for the functional calculus $F(-uA)$ of a generator A of an operator semigroup defined on a sector; here F is given as the Fourier–Borel transform of an analytic functional. The results are linked to the existence of an identity element in the Banach algebra generated by the semigroup. Both the quasinilpotent and non-quasinilpotent cases are considered, and sharp results are proved extending many in the literature.

MATHEMATICS SUBJECT CLASSIFICATION (2000): Primary: 47D03, 46J40, 46H30 Secondary: 30A42, 47A60

KEYWORDS: strongly continuous semigroup, functional calculus, Fourier–Borel transform, analytic semigroup, maximum principle.

1 Introduction

The purpose of this paper is to prove results concerning norm estimates in analytic semigroups which complement the results proved in [3] for semigroups

*Université Paris Est Marne-la-Vallée, 5 bd Descartes, Champs-sur-Marne, 77454 Marne-la-Vallée, cedex 2, France isabelle.chalendar@u-pem.fr.

†IMB, UMR 5251, Université de Bordeaux, 351 cours de la Libération, 33405 Talence Cedex, France. jean.esterle@math.u-bordeaux1.fr.

‡School of Mathematics, University of Leeds, Leeds LS2 9JT, U.K. J.R.Partington@leeds.ac.uk.

defined on \mathbb{R}^+ . For good references on analytic semigroups we recommend the books [4, 11]: we note that analytic semigroups of operators are norm-continuous in the open sector of the plane on which they are defined, and thus also act as semigroups by multiplication on the Banach algebra that they generate. Thus we may easily pass from the language of operators to the language of Banach algebras.

In [1] the following result was proved for semigroups defined on the right-hand half-plane \mathbb{C}_+ . Here, ρ denotes the spectral radius of an operator, and Rad denotes the radical of an algebra.

Theorem 1.1. *Let $(T(t))_{t \in \mathbb{C}_+}$ be an analytic non-quasinilpotent semigroup in a Banach algebra. Let \mathcal{A}_T be the closed subalgebra generated by $(T(t))_{t \in \mathbb{C}_+}$ and let $\gamma > 0$. If there exists $t_0 > 0$ such that*

$$\sup_{t \in \mathbb{C}_+, |t| \leq t_0} \rho(T(t) - T((\gamma + 1)t)) < 2$$

then $\mathcal{A}_T / \text{Rad } \mathcal{A}_T$ is unital, and the generator of $(\pi(T(t)))_{t > 0}$ is bounded, where $\pi : \mathcal{A}_T \rightarrow \mathcal{A}_T / \text{Rad } \mathcal{A}_T$ denotes the canonical surjection.

This can be seen as a lower estimate for a functional calculus in \mathcal{A}_T , determined by $F(-A) = T(t) - T((\gamma + 1)t)$, where $F : s \rightarrow e^{-s} - e^{-(\gamma+1)s}$ is the Laplace transform of the atomic measure $\delta_1 - \delta_{\gamma+1}$.

This approach was taken in [3] for semigroups defined on \mathbb{R}_+ , and very general results were proved for both the quasinilpotent and non-quasinilpotent cases, providing extensions of results in [5, 6, 9] and elsewhere. For a detailed history of the subject, we refer to [2].

Fewer results are available for analytic semigroups, and virtually nothing involving a general functional calculus, although some dichotomy results are given in [2]. To prove more general results for analytic semigroups requires the notion of the Fourier–Borel transform of a distribution acting on analytic functions, and in Section 2 we define these transforms and the associated functional calculus.

In Section 3 we derive results in the non-quasinilpotent case, using the properties of the characters defined on the algebra \mathcal{A}_T , putting Theorem 1.1 in a much more general context. Note that the results we prove are sharp, as is shown in Example 3.6.

Finally, the more difficult case of quasinilpotent semigroups is treated in Section 4, adapting the complex variable methods introduced in [3]. Here there are additional technical difficulties involved in defining the functional calculus, since we now work with measures supported on compact subsets of \mathbb{C} .

2 Analytic semigroups and functional calculus

For $0 < \alpha < \pi/2$ let S_α denote the sector

$$S_\alpha := \{z \in \mathbb{C}_+ : |\arg z| < \alpha\}.$$

Let $H(S_\alpha)$ denote the Fréchet space of analytic functions on S_α , endowed with the topology of local uniform convergence; thus, if $(K_n)_{n \geq 1}$ is an increasing sequence of compact subsets of S_α with $\bigcup_{n \geq 1} K_n = S_\alpha$, we may specify the topology by the seminorms

$$\|f\|_n := \sup\{|f(z)| : z \in K_n\}.$$

Now let $\varphi : H(S_\alpha) \rightarrow \mathbb{C}$ be a continuous linear functional, in the sense that there is an index n and a constant $M > 0$ such that $|\langle f, \varphi \rangle| \leq M\|f\|_n$ for all $f \in H(S_\alpha)$. These are sometimes known as *analytic functionals* [8].

We define the *Fourier–Borel transform* of φ by

$$\mathcal{FB}(\varphi)(z) = \langle e_{-z}, \varphi \rangle,$$

for $z \in \mathbb{C}$, where $e_{-z}(\xi) = e^{-z\xi}$ for $\xi \in S_\alpha$. This is an analogue of the Laplace transform, and is given under that name in [8, Sec. 4.5]. We follow the terminology of [12].

If $\varphi \in H(S_\alpha)'$, as above, then by the Hahn–Banach theorem, it can be extended to a functional on $C(K_n)$, which we still write as φ , and is thus given by a Borel measure μ supported on K_n .

That is, we have

$$\langle f, \varphi \rangle = \int_{S_\alpha} f(\xi) d\mu(\xi),$$

where μ (which is not unique) is a compactly supported measure. For example, if $\langle f, \varphi \rangle = f'(1)$, then

$$\langle f, \varphi \rangle = \frac{1}{2\pi i} \int_C \frac{f(z) dz}{(z-1)^2}, \quad (1)$$

where C is any sufficiently small circle surrounding the point 1.

Note that

$$\mathcal{FB}(\varphi)(z) = \int_{K_n} e^{-z\xi} d\mu(\xi),$$

and thus it is an entire function of z satisfying $\sup_{\operatorname{Re} z > r} |\mathcal{FB}(\varphi)(z)| \rightarrow 0$ as $r \rightarrow \infty$. We shall sometimes find it convenient to use the alternative notation $\mathcal{FB}(\mu)$.

Now let $T := (T(t))_{t \in S_\alpha}$ be an analytic semigroup on a Banach space \mathcal{X} , with infinitesimal generator A . Let $\varphi \in H(S_\alpha)'$ and let $F = \mathcal{FB}(\varphi)$.

We may thus define, formally to start with,

$$F(-A) = \langle T, \varphi \rangle = \int_{S_\alpha} T(\xi) d\mu(\xi),$$

which is well-defined as a Bochner integral in \mathcal{A}_T . It is easy to verify that the definition is independent of the choice of μ representing φ .

Moreover, if $u \in S_{\alpha-\beta}$, where $\operatorname{supp} \mu \subset S_\beta$ and $0 < \beta < \alpha$, then we may also define

$$F(-uA) = \int_{S_\beta} T(u\xi) d\mu(\xi),$$

since $u\xi$ lies in S_α .

In the following, a symmetric measure is a measure such that $\mu(\overline{S}) = \overline{\mu(S)}$ for $S \subset S_\alpha$. A symmetric measure will have a Fourier–Borel transform F satisfying $F(z) = \tilde{F}(z) := \overline{F(\overline{z})}$ for all $z \in \mathbb{C}$.

3 The non-quasinilpotent case

For a semigroup $(T(t))_{t \in S_\alpha}$ with generator A , we write \mathcal{A}_T for the commutative Banach algebra generated by the elements of the semigroup, and $\widehat{\mathcal{A}}_T$ for its character space (Gelfand space).

The following result is proved for semigroups on \mathbb{R}_+ in [3, Lem. 3.1] (see also [5, Lem. 3.1] and [1, Lem. 3.1]). It enables us to regard A itself as an element of $C(\widehat{\mathcal{A}}_T)$ by defining an appropriate value $\chi(A) = -a_\chi$ for each $\chi \in \widehat{\mathcal{A}}_T$.

Lemma 3.1. *For a strongly continuous and eventually norm-continuous semigroup $(T(t))_{t>0}$ and a nontrivial character $\chi \in \widehat{\mathcal{A}}_T$ there is a unique $a_\chi \in \mathbb{C}$ such that $\chi(T(t)) = e^{-ta_\chi}$ for all $t > 0$. Moreover, the mapping $\chi \mapsto a_\chi$ is continuous, and $\chi(F(-uA)) = F(ua_\chi)$ in the case that F is the Laplace transform of a measure μ on \mathbb{R}_+ .*

A similar result holds for analytic semigroups, with the same proof, where now F is the Fourier–Borel transform of a distribution (defined above).

The following theorem extends [1, Thm. 3.6]

Theorem 3.2. *Let $0 < \beta < \alpha < \pi/2$. Let $\varphi \in H(S_\alpha)'$, induced by a symmetric measure $\mu \in M_c(S_\beta)$ such that $\int_{S_\beta} d\mu(z) = 0$, and let $F = \mathcal{FB}(\varphi)$. Let $(T(t))_{t \in S_\alpha} = (\exp(tA))_{t \in S_\alpha}$ be an analytic non-quasinilpotent semigroup and let \mathcal{A}_T be the subalgebra generated by $(T(t))_{t \in S_\alpha}$. If there exists $t_0 > 0$ such that*

$$\sup_{t \in S_{\alpha-\beta}, |t| \leq t_0} \rho(F(-tA)) < \sup_{z \in S_{\alpha-\beta}} |F(z)|, \quad (2)$$

then $\mathcal{A}_T / \text{Rad } \mathcal{A}_T$ is unital and the generator of $\pi(T(t))_{t \in S_\alpha}$ is bounded, where $\pi : \mathcal{A}_T \rightarrow \mathcal{A}_T / \text{Rad}(\mathcal{A}_T)$ denotes the canonical surjection.

Proof. By the maximum principle, for each $\theta \in (-\pi/2, \pi/2)$, F attains its maximum absolute value M_θ , say, on the ray $R_\theta = \{z : \arg z = \theta\}$ and M_θ is an increasing function of θ on $[0, \pi/2)$. Moreover, $M_\theta = M_{-\theta}$, since μ is symmetric.

Clearly there exists a $d > 0$ such that the maximum value of F on each ray R_θ is attained at a point z such that $\operatorname{Re} z \leq d$.

By Lemma 3.1, the hypotheses of the theorem, including non-quasinilpotency, imply that for each $\chi \in \widehat{\mathcal{A}}_T$ there exists $a_\chi \in \mathbb{C}$ such that $\chi(T(t)) = e^{-a_\chi t}$ for all $t \in S_\alpha$, and hence $\chi(F(-tA)) = F(a_\chi t)$. Moreover, we know from (2) that

$$|F(a_\chi t)| < \sup_{z \in S_{\alpha-\beta}} |F(z)|$$

for all $t \in S_{\alpha-\beta}$ with $|t| \leq t_0$.

If for any point t in the sector $\{t \in S_{\alpha-\beta} : |t| \leq t_0\}$ we have $\operatorname{Re} a_\chi t > d$, and $|\arg a_\chi t| \geq \alpha - \beta$, then

$$|F(\lambda a_\chi t)| \geq \sup_{z \in S_{\alpha-\beta}} |F(z)|$$

for some real λ between 0 and 1, giving a contradiction. In particular, if $|\arg a_\chi t_0| \geq \alpha - \beta$, then $\operatorname{Re} a_\chi t_0 \leq d$.

Now suppose that $0 \leq \gamma = \arg a_\chi t_0 < \alpha - \beta$ (the other case is similar); then we know that $\operatorname{Re}(a_\chi t_0 e^{i(\alpha-\beta-\gamma)}) \leq d$; writing $a_\chi t_0 = r e^{i\gamma}$ and $a_\chi t_0 e^{i(\alpha-\beta-\gamma)} = r e^{i(\alpha-\beta)}$, we deduce that $\operatorname{Re} a_\chi t_0 \leq d \cos \gamma / \cos(\alpha - \beta)$. Hence, in all cases, we obtain $\operatorname{Re} a_\chi t_0 \leq d / \cos(\alpha - \beta)$ and

$$|\chi(T(t_0))| \geq \exp(-d / \cos(\alpha - \beta)),$$

and this holds for every $\chi \in \widehat{\mathcal{A}}_T$. Hence $\widehat{\mathcal{A}}_T$ is compact. Now by [10, Thm. 3.6.3, 3.6.6], we conclude that $\mathcal{A}_T / \operatorname{Rad} \mathcal{A}_T$ is unital. Since the algebra generated by the norm-continuous semigroup $(\pi(T(t)))_{t>0}$ is dense in $\mathcal{A}_T / \operatorname{Rad} \mathcal{A}_T$, the generator of this semigroup is bounded. □

By considering the convolution of a functional $\varphi \in H(S_\alpha)'$, possessing the Fourier–Borel transform F , and the functional $\tilde{\varphi}$, possessing the Fourier–Borel transform \tilde{F} (defined by the convolution of defining measures), we obtain a functional whose transform is the product $F\tilde{F}$, and may deduce the following result.

Corollary 3.3. *Let $0 < \beta < \alpha < \pi/2$. Let $\varphi \in H(S_\alpha)'$, induced by a measure $\mu \in M_c(S_\beta)$ such that $\int_{S_\beta} d\mu(z) = 0$, and let $F = \mathcal{FB}(\varphi)$. Let $(T(t))_{t \in S_\alpha} = (\exp(tA))_{t \in S_\alpha}$ be an analytic non-quasinilpotent semigroup and let \mathcal{A}_T be the subalgebra generated by $(T(t))_{t \in S_\alpha}$. If there exists $t_0 > 0$ such that*

$$\sup_{t \in S_{\alpha-\beta}, |t| \leq t_0} \rho(F(-tA)\tilde{F}(-tA)) < \sup_{z \in S_{\alpha-\beta}} |F(z)| |\tilde{F}(z)|,$$

then $\mathcal{A}_T / \text{Rad } \mathcal{A}_T$ is unital and the generator of $\pi(T(t))_{t \in S_\alpha}$ is bounded, where $\pi : \mathcal{A}_T \rightarrow \mathcal{A}_T / \text{Rad}(\mathcal{A}_T)$ denotes the canonical surjection.

A similar proof gives the following result.

Theorem 3.4. *Let $0 < \alpha < \pi/2$. Let $\varphi \in H(S_\alpha)'$, induced by a symmetric measure $\mu \in M_c(S_\alpha)$ such that $\int_{S_\alpha} d\mu(z) = 0$, and let $F = \mathcal{FB}(\varphi)$. Let $(T(t))_{t \in S_\alpha} = (\exp(tA))_{t \in S_\alpha}$ be an analytic non-quasinilpotent semigroup and let \mathcal{A}_T be the subalgebra generated by $(T(t))_{t \in S_\alpha}$. If there exists $t_0 > 0$ such that*

$$\rho(F(-tA)) < \sup_{x > 0} |F(x)|,$$

for all $0 < t \leq t_0$, then $\mathcal{A}_T / \text{Rad } \mathcal{A}_T$ is unital and the generator of $\pi(T(t))_{t \in S_\alpha}$ is bounded, where $\pi : \mathcal{A}_T \rightarrow \mathcal{A}_T / \text{Rad}(\mathcal{A}_T)$ denotes the canonical surjection.

Proof. The proof is similar to that of Theorem 3.2, and we adopt the same notation. We cannot have $\text{Re } a_\chi t_0 > d$ for any $\chi \in \widehat{\mathcal{A}}_T$, as then there would be a $\lambda \in (0, 1)$ with

$$|\chi(F(-\lambda t_0 A))| = |F(\lambda a_\chi t_0)| \geq \sup_{x > 0} |F(x)|.$$

We conclude that $|\chi(T(t_0))| \geq \exp(-d)$ for all $\chi \in \widehat{\mathcal{A}}_T$. The proof is now concluded as before. □

Again there is an immediate corollary for general φ .

Corollary 3.5. *Let $0 < \alpha < \pi/2$. Let $\varphi \in H(S_\alpha)'$, induced by a measure $\mu \in M_c(S_\alpha)$ such that $\int_{S_\alpha} d\mu(z) = 0$, and let $F = \mathcal{FB}(\varphi)$. Let $(T(t))_{t \in S_\alpha} = (\exp(tA))_{t \in S_\alpha}$ be an analytic non-quasinilpotent semigroup and let \mathcal{A}_T be the subalgebra generated by $(T(t))_{t \in S_\alpha}$. If there exists $t_0 > 0$ such that*

$$\rho(F(-tA)\tilde{F}(-tA)) < \sup_{x>0} |F(x)|^2,$$

for all $0 < t \leq t_0$, then $\mathcal{A}_T/\text{Rad } \mathcal{A}_T$ is unital and the generator of $\pi(T(t))_{t \in S_\alpha}$ is bounded, where $\pi : \mathcal{A}_T \rightarrow \mathcal{A}_T/\text{Rad } (\mathcal{A}_T)$ denotes the canonical surjection.

Example 3.6. *In the Banach algebra $\mathcal{A} = C_0[0, 1]$ consider the semigroup $T(t) : x \mapsto x^t$ for $t \in S_\alpha$. Clearly there is no convergence as t approaches 0.*

For $x \in (0, 1]$ (which can be identified with the Gelfand space of \mathcal{A}) let $F = \mathcal{FB}(\mu)$ and

$$F(-tA)(x) = \int_{S_\alpha} x^{t\xi} d\mu(\xi) = \int_{S_\alpha} e^{t\xi \log x} d\mu(\xi),$$

where $\mu \in M_c(S_\alpha)$, supposing that $\int_{S_\alpha} d\mu(z) = 0$ and that F is real for $x > 0$.

Thus $F(-tA)(x) = F(-t \log x)$ and

$$\rho(F(-tA)) = \|F(-tA)\| = \sup_{x \in (0,1]} |F(-t \log x)| = \sup_{r>0} |F(tr)|.$$

Clearly

$$\sup_{t \in S_\alpha, |t| \leq t_0} \rho(F(-tA)) = \sup_{t \in S_\alpha} |F(z)|$$

for all $t_0 > 0$. This shows that the hypothesis (2) of Theorem 3.2 is sharp.

Remark 3.7. *In [3, Thm. 3.3] it is shown that if F is the Laplace transform of a real compactly-supported measure μ with $\int_0^\infty d\mu = 0$, then the condition*

$$\rho(F(-u_k A)) < \sup_{t>0} |F(u_k t)|$$

for a (real) sequence $u_k \rightarrow 0$ implies that the algebra \mathcal{A}_T possesses an exhaustive sequence of idempotents $(P_n)_{n \geq 1}$ (i.e., $P_n^2 = P_n P_{n+1} = P_n$ for all n

and for every $\chi \in \widehat{\mathcal{A}}_T$ there is a p such that $\chi(P_n) = 1$ for all $n \geq p$, such that each semigroup $(P_n T(t))_{t>0}$ has a bounded generator.

The directly analogous result for analytic semigroups does not hold. For example, we may consider a modification of Example 3.6, by taking $\mathcal{A} = C_0[0, 1]$ and the semigroup $T(t) : x \rightarrow x^t = \exp(t \log x)$ for $x \in (0, 1]$ and $t \in S_{\pi/2}$ with generator A , say. Let $F(z) = e^{-z} - e^{-2z} = \mathcal{FB}(\delta_1 - \delta_2)$. Now let $\omega = \exp(i\pi/6)$ and define $\widetilde{T}(t) = T(\omega t)$. Thus $\widetilde{T}(t)$ is an analytic semigroup in $S_{\pi/3}$, with generator $\widetilde{A} = \omega A$. Define $\widetilde{F}(z) = F(z/\omega)$, the Fourier–Borel transform of a measure supported in $S_{\pi/3}$. Now, for $u > 0$,

$$\rho(\widetilde{F}(-u\widetilde{A})) = \rho(F(-uA)) = \rho(x^u - x^{2u}) = 1/4;$$

however,

$$\sup_{t>0} |\widetilde{F}(tu)| = \sup\{|F(tu/\omega)| : t > 0\} > 1/4,$$

by the maximum principle applied to F on the sector $S_{\pi/6}$ (numerically, the supremum is about 0.29).

4 The quasinilpotent case

We suppose throughout this section that $(T(t))_{t \in S_\alpha}$ is a nontrivial strongly continuous semigroup of uniformly bounded quasinilpotent operators acting on a Banach space $(\mathcal{X}, \|\cdot\|)$.

4.1 Preliminary results

We shall need the following theorem from complex analysis, which was proved in [3, Thm. 2.2].

Theorem 4.1. *Let $f : \overline{\mathbb{C}_+} \rightarrow \mathbb{C}$ be a continuous bounded nonconstant function, analytic on \mathbb{C}_+ , such that $f([0, \infty)) \subset \mathbb{R}$, $f(0) = 0$, satisfying the condition $\lim_{x \rightarrow \infty, x \in \mathbb{R}} f(x) = 0$.*

Assume that $\sup_{t \in S_\beta \cap D(0,1)} \|T(t)\| < +\infty$ for every $\beta \in [0, \alpha)$, so that the Bochner integral

$$\int_K T(s\zeta u) e^{(s-1)\zeta\lambda} \zeta d\mu(\zeta)$$

is well-defined in $\mathcal{B}(\mathcal{X})$ for every $\zeta \in K$, $\lambda \in \mathbb{C}$, and $u \in \mathbb{C}$ with $uK \subset S_\alpha$. Then, for $F = \mathcal{FB}(\varphi)$ we have

$$F(-uA) - F(\lambda)I = (uA + \lambda I) \left(\int_0^1 \left[\int_K T(s\zeta u) e^{(s-1)\zeta\lambda} \zeta d\mu(\zeta) \right] ds \right).$$

Also, if $x \in D(A)$, we have

$$F(-uA)x - F(\lambda)x = \left(\int_0^1 \left[\int_K T(s\zeta u) e^{(s-1)\zeta\lambda} \zeta d\mu(\zeta) \right] ds \right) (uA + \lambda I)x.$$

Proof. Using Fubini's theorem and the fundamental theorem of the calculus, we have

$$\begin{aligned} \int_0^1 \left[\int_K T(s\zeta u) e^{(s-1)\zeta\lambda} \zeta d\mu(\zeta) \right] ds &= \int_K (uA + \lambda I)^{-1} (T(\zeta u) - e^{-\zeta\lambda} I) d\mu(\zeta) \\ &= (uA + \lambda I)^{-1} (F(-uA) - F(\lambda)I), \end{aligned}$$

from which the result is clear. \square

Corollary 4.3. *Let $(T(t) = (e^{tA}))_{t \in S_\alpha}$ be a bounded analytic semigroup on S_α . Then there exists a constant $C > 0$ such that*

$$\|(F(-uA) - F(\lambda)I)(uA + \lambda I)^{-1}\| \leq C \int_K |\zeta| d|\mu|(\zeta),$$

for all u with $uK \subset S_\alpha$ and $|\lambda| \leq 1$.

Proof. This follows immediately from Lemma 4.2 since $|e^{(s-1)\zeta\lambda}|$ is uniformly bounded for $\zeta \in K$ and $s \in [0, 1]$. \square

We shall require one further preliminary result, which is well-known for z in the closed left half-plane, and was proved in [3].

Lemma 4.4. *Let $(T(t))_{t \in S_\alpha}$ be bounded analytic quasinilpotent semigroup with generator A , and let $0 < \beta < \alpha$. Then $\|(A + zI)^{-1}\|$ is uniformly bounded on the set*

$$\left\{ z \in \mathbb{C} : \frac{\pi}{2} - (\alpha - \beta) \leq |\arg z| \leq \pi \right\}.$$

Proof. It is standard that $\|z(A + zI)^{-1}\|$ is bounded in the given set for any bounded analytic semigroup. Moreover, quasinilpotency implies that A is invertible and then $\|(A + zI)^{-1}\|$ is bounded near 0. \square

4.2 Estimates for quasinilpotent semigroups

The following theorem is a version of [3, Thm. 2.5] for analytic semigroups. The proof is based on the same ideas from complex analysis with several significant modifications. For $0 < \beta < \pi/2$ let

$$V_\beta := \{z \in \mathbb{C}_+ : |\arg z| \leq \beta\}.$$

Theorem 4.5. *Let $(T(t))_{t \in S_\alpha}$ be a nontrivial bounded semigroup of quasinilpotent operators and let F be the Fourier–Borel transform of a symmetric measure μ supported on a compact set $K \subset V_\beta$ for some β with $0 < \beta < \alpha$, such that $\int_K d\mu = 0$.*

Then there is an $\eta > 0$ such that

$$\|F(-uA)\| > \sup_{t>0} |F(t)|$$

for all $u \in S_{\alpha-\beta}$ with $|u| \leq \eta$.

Proof. Note that F is real on the positive real axis, since μ is symmetric. Let $b > 0$ be such that $|F(x)| \leq |F(b)|$ for all $x \geq 0$. By considering $-\mu$ instead of μ , if necessary, we may suppose that $F(b) > 0$.

By Corollary 4.3

$$\|F(-uA)(uA + \lambda I)^{-1}\| \geq \|F(\lambda)(uA + \lambda I)^{-1}\| - C \int_K |\zeta| d|\mu|(\zeta).$$

for $u \in S_{\alpha-\beta}$ and $|\lambda| \leq 1$, and hence

$$\|F(-uA)\| \geq |F(\lambda)| - \frac{C}{\|(uA + \lambda I)^{-1}\|} \int_K |\zeta| d|\mu|(\zeta).$$

Suppose that there exists a $u \in S_{\alpha-\beta}$ with $|u| < 1$ such that $\|F(-uA)\| \leq F(b)$, and consider the simple Jordan curve

$$\Gamma := [b, a_1] \cup \Gamma_1 \cup [a_2, a_3] \cup [a_3, \bar{a}_3] \cup [\bar{a}_3, \bar{a}_2] \cup \bar{\Gamma}_1 \cup [\bar{a}_1, b],$$

where Γ_1, a_1, a_2, a_3 are defined as in Theorem 4.1, taking $f = F$ (see Figure 1).

We now make various estimates of $\|(uA + \lambda I)^{-1}\|$ for λ on three different parts of Γ .

1) By (i) of Theorem 4.1, for $\lambda \in [b, a_1] \cup [\bar{a}_1, b]$ we have

$$\begin{aligned} F(b) \geq \|F(-uA)\| &\geq |F(\lambda)| - \frac{C}{\|(uA + \lambda I)^{-1}\|} \int_K |\zeta| d|\mu|(\zeta) \\ &\geq F(b) + \delta|\lambda - b|^m - \frac{C}{\|(uA + \lambda I)^{-1}\|} \int_K |\zeta| d|\mu|(\zeta). \end{aligned}$$

Hence we obtain

$$\|(uA + \lambda I)^{-1}\| \leq \frac{C}{\delta|\lambda - b|^m} \int_K |\zeta| d|\mu|(\zeta). \quad (3)$$

2) By (ii) of Theorem 4.1, for $\lambda \in \Gamma_1 \cup [a_2, a_3] \cup [\bar{a}_3, \bar{a}_2] \cup \bar{\Gamma}_1$ we have

$$\begin{aligned} F(b) \geq \|F(-uA)\| &\geq |F(\lambda)| - \frac{C}{\|(uA + \lambda I)^{-1}\|} \int_K |\zeta| d|\mu|(\zeta) \\ &\geq |F(a_0)| - \frac{C}{\|(uA + \lambda I)^{-1}\|} \int_K |\zeta| d|\mu|(\zeta). \end{aligned}$$

It follows that

$$\|(uA + \lambda I)^{-1}\| \leq \frac{C}{|F(a_0)| - F(b)} \int_K |\zeta| d|\mu|(\zeta). \quad (4)$$

3) By Lemma 4.4 there is a constant $C' > 0$ depending only on the semigroup such that

$$\left\| \left(A + \frac{\lambda}{u} I \right)^{-1} \right\| \leq C' \quad (5)$$

for $\lambda \in [a_3, \bar{a}_3] \subset i\mathbb{R}$.

We can now provide estimates for the quantity $\left\| (\lambda - b)^m \left(A + \frac{\lambda}{u} I \right)^{-1} \right\|$ for λ on Γ . Let $R = \max_{\lambda \in \Gamma} |\lambda - b|$.

By (3)

$$\left\| (\lambda - b)^m \left(A + \frac{\lambda}{u} I \right)^{-1} \right\| \leq \frac{C|u|}{\delta} \int_K |\zeta| d|\mu|(\zeta)$$

for all $\lambda \in [b, a_1] \cup [\bar{a}_1, b]$.

By (4)

$$\left\| (\lambda - b)^m \left(A + \frac{\lambda}{u} I \right)^{-1} \right\| \leq \frac{C|u|R^m}{|F(a_0)| - F(b)} \int_K |\zeta| d|\mu|(\zeta)$$

for all $\lambda \in \Gamma_1 \cup [a_2, a_3] \cup [\bar{a}_3, \bar{a}_2] \cup \bar{\Gamma}_1$.

By (5)

$$\left\| (\lambda - b)^m \left(A + \frac{\lambda}{u} I \right)^{-1} \right\| \leq C' R^m$$

for all $\lambda \in [a_3, \bar{a}_3]$.

Since $0 < |u| \leq 1$, for all $z \in \Gamma \cup \text{int } \Gamma$ we have

$$\left\| \left(A + \frac{z}{u} I \right)^{-1} \right\| \leq \frac{M}{|z - b|^m},$$

by the maximum modulus principle, where

$$M = \max \left(\frac{C}{\delta} \int_K |\zeta| d|\mu|(\zeta), \frac{C R^m}{|F(a_0)| - F(b)} \int_K |\zeta| d|\mu|(\zeta), C' R^m \right).$$

Since by hypothesis $F(0) = 0$, there is an $r \in (0, b)$ such that

$$\sup_{|z| \leq r} |F(z)| < F(b).$$

Taking r sufficiently small, we have $\overline{D(0, r)} \cap \Gamma \cap \mathbb{C}_+ = \emptyset$, and then $\overline{D(0, r)} \cap \overline{\mathbb{C}_+} \subset \Gamma \cup \text{int } \Gamma$.

Now if $z \in \overline{D(0, r)}$ with $\operatorname{Re} z > 0$, we have $|z - b| \geq b - r$, and thus we have

$$\left\| \left(A + \frac{z}{u} I \right)^{-1} \right\| \leq \frac{M}{|z - b|^m} \leq \frac{M}{(b - r)^m}. \quad (6)$$

Also, by Lemma 4.4 there is an $M' > 0$ such that

$$\sup_{z \in E} \|(A + zI)^{-1}\| \leq M',$$

where

$$E = \left\{ z \in \mathbb{C} : \frac{\pi}{2} - (\alpha - \beta) \leq |\arg z| \leq \pi \right\}.$$

Hence

$$\left\| \left(A + \frac{z}{v} I \right)^{-1} \right\| \leq M' \quad (7)$$

for $\operatorname{Re} z \leq 0$ and $v \in S_{\alpha - \beta}$.

Now, since by Liouville's theorem the function $\lambda \mapsto \|(A + \lambda I)^{-1}\|$ is unbounded on \mathbb{C} , and so there is a $\lambda \in \mathbb{C}$ with

$$\|(A + \lambda I)^{-1}\| > \max \left\{ M', \frac{M}{(b - r)^m} \right\}.$$

It follows that there is an $\eta > 0$ such that for all $u \in S_{\alpha - \beta}$ with $|u| \leq \eta$ we have

$$\left\| \left(A + \frac{z}{u} I \right)^{-1} \right\| > \max \left\{ M', \frac{M}{(b - r)^m} \right\}$$

for some $z = \lambda u \in \overline{D(0, r)}$. Note that $z \in \mathbb{C}_+$ by (7).

This contradicts (6), and it follows that

$$\|F(-uA)\| > F(b) \quad \text{for all } u \in S_{\alpha - \beta}, \quad |u| \leq \eta.$$

□

Example 4.6. *The analytic distribution $\varphi : f \mapsto f'(1)$ discussed earlier (cf. (1)) leads to $F(z) = -ze^{-z}$ and $F(-uA) = uAT(u)$. Theorem 4.5 now asserts that $\|uAT(u)\| \geq 1/e$ for sufficiently small $u \in S_{\alpha - \beta}$, where β corresponds to the support of φ and can therefore be taken arbitrarily small.*

This may be seen in the context of the result of Hille [6] (see also [7, Thm. 10.3.6]) that if $\limsup_{t \rightarrow 0^+} t \|AT(t)\| < 1/e$, then A is bounded, and hence the semigroup cannot be quasinilpotent. There is a further discussion of this example in [1, Thm. 3.12, Thm 3.13].

We have a corollary for measures that are not necessarily symmetric. If μ is now a complex compactly-supported measure, with Fourier–Borel transform F , then we write $\tilde{F}(z) = \overline{F(\bar{z})}$; then \tilde{F} is also an entire function, indeed, the Fourier–Borel transform of $\bar{\mu}$.

Corollary 4.7. *Let $(T(t))_{t \in S_\alpha}$ be a nontrivial bounded analytic semigroup of quasinilpotent operators and let F be the Fourier–Borel transform of a measure μ supported on a compact set $K \subset V_\beta$ for some β with $0 < \beta < \alpha$, such that $\int_K d\mu = 0$.*

Then there is an $\eta > 0$ such that

$$\|F(-uA)\tilde{F}(-uA)\| \geq \sup_{t>0} |F(t)|^2$$

for all $u \in S_{\alpha-\beta}$ with $|u| \leq \eta$.

Proof. The result follows on applying Theorem 4.5 to the real measure $\nu := \mu * \bar{\mu}$, whose Fourier–Borel transform satisfies

$$\mathcal{FB}(\nu)(s) = F(s)\tilde{F}(s).$$

□

Acknowledgements

The authors thank the Institut Camille Jordan of Université Lyon 1 and the IMB of the Université de Bordeaux for their hospitality and financial support. They are also grateful to the referee for many useful suggestions.

References

- [1] Z. Bendaoud, I. Chalendar, J. Esterle, and J. R. Partington. Distances between elements of a semigroup and estimates for derivatives. *Acta Math. Sin. (Engl. Ser.)*, 26 (2010), no. 12, 2239–2254.
- [2] I. Chalendar, J. Esterle and J.R. Partington, Dichotomy results for norm estimates in operator semigroups, *Operator Semigroups meet Complex Analysis, Harmonic Analysis and Mathematical Physics*. Ed. by W. Arendt, R. Chill and Y. Tomilov, Operator Theory: Advances and Applications, Vol. 250, Birkhäuser, 2015.
- [3] I. Chalendar, J. Esterle, and J. R. Partington. Lower estimates near the origin for functional calculus on operator semigroups. *J. Funct. Anal.* 269 (2015), no. 6, 1620–1635.
- [4] K.-J. Engel and R. Nagel, *One-parameter semigroups for linear evolution equations*. Graduate Texts in Mathematics, 194. Springer-Verlag, New York, 2000.
- [5] J. Esterle and A. Mokhtari. Distance entre éléments d’un semi-groupe dans une algèbre de Banach. *J. Funct. Anal.*, 195 (2002), no. 1, 167–189.
- [6] E. Hille, On the differentiability of semi-group operators. *Acta Sci. Math. (Szeged)*, 12 (1950), 19–24.
- [7] E. Hille and R. S. Phillips, *Functional analysis and semi-groups*. Third printing of the revised edition of 1957. American Mathematical Society Colloquium Publications, Vol. XXXI. American Mathematical Society, Providence, R. I., 1974.
- [8] L. Hörmander, *An introduction to complex analysis in several variables*. Third edition. North-Holland Mathematical Library, 7. North-Holland Publishing Co., Amsterdam, 1990.

- [9] N. Kalton, S. Montgomery-Smith, K. Oleszkiewicz and Y. Tomilov, Power-bounded operators and related norm estimates. *J. London Math. Soc.* (2) 70 (2004), no. 2, 463–478.
- [10] C. E. Rickart. *General theory of Banach algebras*. The University Series in Higher Mathematics. D. van Nostrand Co., Inc., Princeton, N.J.–Toronto–London–New York, 1960.
- [11] A. M. Sinclair, *Continuous semigroups in Banach algebras*. London Mathematical Society Lecture Note Series, 63. Cambridge University Press, Cambridge–New York, 1982.
- [12] D.C. Struppa, *Fourier–Borel transform*. Encyclopedia of Mathematics. http://www.encyclopediaofmath.org/index.php?title=Fourier-Borel_transform.