



This is a repository copy of *Diagnostic accuracy of biomarkers and imaging for bone turnover in renal osteodystrophy*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/128959/>

Version: Supplemental Material

---

**Article:**

Salam, S. [orcid.org/0000-0001-9236-3906](https://orcid.org/0000-0001-9236-3906), Gallagher, O., Gossiel, F. et al. (3 more authors) (2018) Diagnostic accuracy of biomarkers and imaging for bone turnover in renal osteodystrophy. *Journal of the American Society of Nephrology*, 29 (5). pp. 1557-1565. ISSN 1046-6673

<https://doi.org/10.1681/ASN.2017050584>

---

© 2018 The American Society of Nephrology. This is an author produced version of a paper subsequently published in *Journal of the American Society of Nephrology*. Uploaded in accordance with the publisher's self-archiving policy.

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# Table of Contents

**Supplementary methods..... 2-3**

**Supplementary tables ..... 4**

## Supplementary methods

**BTMs:** The inter-assay coefficient of variations (CVs) were  $\leq 9\%$  for iPTH, BALP and CTX;  $\leq 5\%$  for intact PINP;  $\leq 10\%$  for 25-hydrovitamin D and TRAP5b; and  $\leq 4\%$  for total PINP. Inter-assay CVs were  $\leq 5\%$  for all biochemistry measured on the same day as the blood sample collection using Roche Cobas c701/702 analyser (Roche Diagnostics, England).

**HR-pQCT:** The non-dominant side was scanned unless there was a previous fracture or an arterio-venous fistula present. The scans were acquired in the high-resolution mode (image matrix = 1,536 x 1,536) using a source potential of 60 kVp, a tube current of 900 mA, and an integration time of 100 ms. Each scan resulted in the acquisition of a total of 110 image slices (stack height = 9.02mm) at an isotropic resolution of 82  $\mu\text{m}$ . A maximum of one repeat scan at either or both anatomical sites was performed in the event of patient movement. All images were graded by a single operator (MP), and any images with an unacceptable degree of movement artefact were excluded from further analysis. We had to exclude images of 9 radii from controls, 12 radii from CKD, one tibia from control and two tibiae from CKD due to movement artefact. Microarchitectural parameters included trabecular number (Tb.N, 1/mm), trabecular thickness (Tb.Th, mm), trabecular separation (Tb.Sp, mm), bone volume/tissue volume (BV/TV, %) and cortical thickness (Ct.Th, mm). Extended cortical analysis for cortical microstructural bone properties, including cortical porosity (%) and cortical pore diameter (Ct.Po.Dm,  $\mu\text{m}$ ).

**DXA:** Lumbar spine scans were assessed by a single operator (MP) for vertebral fracture and other causes leading to falsely high BMD (such as degenerative changes and osteophytes). One lumbar spine scan was excluded due to multiple vertebral fractures and one patient did not have a forearm scan due to previous bilateral wrist fractures.

**Transiliac bone biopsy & histomorphometry:** To assess bone formation rate (BFR), patients received four days of tetracycline (250mg, four times a day) and after a 10-day intermission, a further two days of demeclocycline (300mg, twice a day). Trans-iliac bone biopsy was performed within 2-4 days of the last dose of demeclocycline. The undecalcified bone sample was placed in 70%, 80%, 90% and 100% ethanol in turn for at least one week in each solution while shielded from light at all time. It was then infiltrated and embedded in LR White medium grade resin. The resin block was sectioned at 6 $\mu$ m thickness and stained with Masson Goldner trichrome.

**Supplementary Table 1:** Correlation between all bone turnover markers in 43 advanced CKD patients with bone histomorphometry data.

	Intact PINP	Total PINP	bALP	tALP	CTX	TRAP5b
iPTH	0.619	0.523	0.621	0.508	0.567	0.504
Intact PINP		0.813	0.866	0.762	0.732	0.741
Total PINP			0.749	0.698	0.591	0.575
bALP				0.869	0.710	0.672
tALP					0.542	0.548
CTX						0.709

All Spearman's rho correlations are significant at  $\leq 0.001$  (2-tailed)

**Supplementary Table 2:** Diagnostic accuracy of DXA areal BMD Z-scores for identifying advanced CKD patients with low and high bone turnover (N=43).

Areal BMD Z-score	AUC (95% CI)
Low bone turnover	
Forearm	0.645 (0.483 – 0.786)
Total hip	0.598 (0.438 – 0.744)
Lumbar spine	0.554 (0.395 – 0.705)
High bone turnover	
Forearm	0.685 (0.523 – 0.819)
Total hip	0.580 (0.420 – 0.729)
Lumbar spine	0.533 (0.375 – 0.686)

**Supplementary Table 3:** Diagnostic accuracy of HR-pQCT Z-scores for identifying advanced CKD patients with high bone turnover (N=43).

HR-pQCT Z-scores	Distal radius AUC (95% CI)	Distal tibia AUC (95% CI)
Total vBMD	0.627 (0.450 – 0.782)	0.553 (0.392 – 0.706)
Cortical vBMD	0.569 (0.394 – 0.732)	0.579 (0.417 – 0.729)
Trabecular vBMD	0.625 (0.448 – 0.780)	0.541 (0.381 – 0.696)
Cortical thickness	0.552 (0.377 – 0.717)	0.525 (0.365 – 0.681)
Cortical porosity	0.627 (0.450 – 0.782)	0.548 (0.387 – 0.702)
Cortical BV/TV	0.628 (0.451 – 0.783)	0.576 (0.414 – 0.727)
Trabecular thickness	0.583 (0.407 – 0.744)	0.575 (0.413 – 0.726)
Trabecular number	0.603 (0.427 – 0.762)	0.541 (0.381 – 0.696)
Trabecular separation	0.614 (0.438 – 0.771)	0.533 (0.373 – 0.688)
Trabecular BV/TV	0.628 (0.451 – 0.783)	0.542 (0.382 – 0.697)