UNIVERSITY of York

This is a repository copy of Arbuscular Mycorrhizal Fungi and Plant Chemical Defence:Effects of Colonisation on Aboveground and Belowground Metabolomes.

White Rose Research Online URL for this paper: <u>https://eprints.whiterose.ac.uk/129480/</u>

Version: Accepted Version

# Article:

Hill, Elizabeth M, Robinson, Lynne A., Abdul-Sada, Ali et al. (3 more authors) (2018) Arbuscular Mycorrhizal Fungi and Plant Chemical Defence:Effects of Colonisation on Aboveground and Belowground Metabolomes. Journal of chemical ecology. pp. 198-208. ISSN 0098-0331

https://doi.org/10.1007/s10886-017-0921-1

# Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

# Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk https://eprints.whiterose.ac.uk/

# Journal of Chemical Ecology ARBUSCULAR MYCORRHIZAL FUNGI AND PLANT CHEMICAL DEFENCE: EFFECTS OF COLONISATION ON ABOVEGROUND AND BELOWGROUND METABOLOMES

--Manuscript Draft--

Manuscript Number:	JOCE-D-17-00277R		
Full Title:	ARBUSCULAR MYCORRHIZAL FUNGI AND PLANT CHEMICAL DEFENCE: EFFECTS OF COLONISATION ON ABOVEGROUND AND BELOWGROUND METABOLOMES		
Article Type:	Original Research		
Keywords:	Arbuscular mycorrhizal symbiosis, Above-belowground interactions, Metabolomics, Pyrrolizidine alkaloids, Blumenols, Senecio jacobaea, Rhizophagus irregularis.		
Corresponding Author:	Sue Hartley University of York UNITED KINGDOM		
Corresponding Author Secondary Information:			
Corresponding Author's Institution:	University of York		
Corresponding Author's Secondary Institution:			
First Author:	Elizabeth Hill		
First Author Secondary Information:			
Order of Authors:	Elizabeth Hill		
	Lynne Robinson		
	Ali Abdul-Sada		
	Adam Vanbergen		
	Angela Hodge		
	Sue Hartley		
Order of Authors Secondary Information:			
Funding Information:			
Abstract:	Arbuscular mycorrhizal fungal (AMF) colonisation of plant roots is one of the most ancient and widespread interactions in ecology, yet the systemic consequences for plant secondary chemistry remain unclear. We performed the first metabolomic investigation into the impact of AMF colonisation, by Rhizophagus irregularis on the chemical defences, spanning above- and below-ground tissues, in its host-plant ragwort (Senecio jacobaea). We used a non-targeted metabolomics approach to profile, and where possible identify, compounds induced by AMF colonisation in both roots and shoots. Metabolomics analyses revealed that 33 compounds were significantly increased in the root tissue of AMF-colonised plants, including seven blumenols, plant-derived compounds known to be associated with AMF colonisation. One of these was a novel structure conjugated with a malonyl-sugar and uronic acid moiety, hitherto an unreported combination. Such structural modifications of blumenols could be significant for their previously reported functional roles associated with the establishment and maintenance of AM colonisation. Pyrrolizidine alkaloids (PAs), key anti-herbivore defence compounds in ragwort, dominated the metabolomic profiles of root and shoot extracts. Analyses of the metabolomic profiles revealed an increase in four PAs in roots (but not shoots) of AMF colonised plants, with the potential to protect colonised plants from below-ground organisms.		

Dear Prof Hartley:

Your manuscript, JOCE-D-17-00277, "ARBUSCULAR MYCORRHIZAL FUNGI AND PLANT CHEMICAL DEFENCE: EFFECTS OF COLONISATION ON ABOVEGROUND AND BELOWGROUND METABOLOMES", that you submitted to the JOURNAL OF CHEMICAL ECOLOGY has been reviewed by the handling editor. Some minor edits are needed before final acceptance.

Please notify all co-authors of the decision.

To submit your revised manuscript, please log onto the following:

Your username is: SHartley-928 If you forgot your password, you can click the 'Send Login Details' link on the EM Login page at http://joce.edmgr.com/.

Please click "Author Login" to get into the site. Be sure to delete the old version of your paper before uploading the revised version.

Please also be sure to upload a letter in which you explain in detail how you have responded to the reviewers' concerns. Please find specific formatting instructions in the Author Instructions at:

http://www.springer.com/west/home/life+sci?SGWID=4-10027-70-356744610&detailsPage=contentItemPage&contentItemId= 143542&CIPageCounter=CI\_FOR\_AUTHORS\_AND\_EDITORS\_ PAGE0&SHORTCUT=www.springer.com/sgw/cda/frontpage/0,11855,4-10027-70-35674461-0,00.html

(they can also be found below the reviewer comments at the end of this email). Please read these over before you prepare the final manuscript and follow them precisely. Papers are not accepted by the Journal until they have been edited after revision. Papers requiring major revisions are subject to additional review. If you anticipate that it will take longer than 2 months to complete the revision, please let us know.

Please make sure to submit your editable source files (i. e. Word, TeX).

Thank you for submitting your work to the JOURNAL OF CHEMICAL ECOLOGY for appraisal. If you have questions or concerns, please do not hesitate to email me.

Sincerely, Nicole M. van Dam Associate Editor Journal of Chemical Ecology

Comments for the author:

The changes made this a great contribution to J Chem Ecology. It was a pleasure to read the revised MS. Thanks for taking the effort!

There is one minor detail that you need to revise: could you please align the floating points of all the values in the tables?

All the points have been aligned in the tables. Furthermore I think the word: Plants in line 309 should be removed,

# Removed

and in the legend of figure 1, it should be :Open and closed circles (move the word circles).

## Corrected.

The title "Table 3" (was 4) still contains a correction marking. Can you please make it final?

# Corrected.

Thanks again for submitting your work to JCE

1 2	1	ARBUSCULAR MYCORRHIZAL FUNGI AND PLANT CHEMICAL
3 4 5	2	DEFENCE: EFFECTS OF COLONISATION ON ABOVEGROUND AND
6 7 8	3	BELOWGROUND METABOLOMES
9 10 11	4	
12 13 14	5	ELIZABETH M. HILL <sup>1</sup> , LYNNE A. ROBINSON <sup>1,2</sup> , ALI ABDUL-SADA <sup>1</sup> ,
15 16 17	6	ADAM J. VANBERGEN <sup>2</sup> , ANGELA HODGE <sup>3</sup> AND SUE E. HARTLEY <sup>3, 4*</sup>
19 20 21	7 8	<sup>1</sup> School of Life Sciences, University of Sussex, Brighton BN1 9QG, UK.
22 23	9	<sup>2</sup> Centre for Ecology and Hydrology, Bush Estate, Penicuik, Midlothian EH26 0QB, UK.
24 25 26	10	<sup>3</sup> Department of Biology, University of York, Wentworth Way, York YO10 5DD, UK.
27 28	11	<sup>4</sup> York Environment and Sustainability Institute, Department of Biology, University of York,
29 30 21	12	Wentworth Way, York YO10 5DD, UK.
31 32 33	13	* Corresponding author: sue.hartley@york.ac.uk
34 35	14	
36 37 38	15	
39 40 41	16	
42 43	17	
44 45 46	18	
47 48 49		
50 51		
52 53		
54 55		
56 57		
58 59		
60 61		
ю⊥ 62		1
63 64		
65		

Abstract - Arbuscular mycorrhizal fungal (AMF) colonisation of plant roots is one of the most ancient and widespread interactions in ecology, yet the systemic consequences for plant secondary chemistry remain unclear. We performed the first metabolomic investigation into the impact of AMF colonisation, by *Rhizophagus irregularis* on the chemical defences, spanning above- and below-ground tissues, in its host-plant ragwort (Senecio jacobaea). We used a non-targeted metabolomics approach to profile, and where possible identify, compounds induced by AMF colonisation in both roots and shoots. Metabolomics analyses revealed that 33 compounds were significantly increased in the root tissue of AMF-colonised plants, including seven blumenols, plant-derived compounds known to be associated with AMF colonisation. One of these was a novel structure conjugated with a malonyl-sugar and uronic acid moiety, hitherto an unreported combination. Such structural modifications of blumenols could be significant for their previously reported functional roles associated with the establishment and maintenance of AM colonisation. Pyrrolizidine alkaloids (PAs), key anti-herbivore defence compounds in ragwort, dominated the metabolomic profiles of root and shoot extracts. Analyses of the metabolomic profiles revealed an increase in four PAs in roots (but not shoots) of AMF colonised plants, with the potential to protect colonised plants from below-ground organisms.

**Key Words** – Arbuscular mycorrhizal symbiosis, Above-belowground interactions, Metabolomics, Pyrrolizidine alkaloids, Blumenols, Senecio jacobaea, Rhizophagus irregularis.

**INTRODUCTION** 

Plant-mediated interactions between organisms above- and below-ground are increasingly recognised as a structuring force in ecology, though in many systems there is still a need for a more mechanistic understanding of the chemical basis of these effects (Johnson et al. 2012, 2016; Wardle et al. 2004). Microbial symbionts have significant impacts on the chemical composition of their host plant, so are potentially key drivers of these indirect interactions (Hartley and Gange 2009). One of the most ancient and widely occurring plant-symbiont relationships involves arbuscular mycorrhizal fungi (AMF) (Smith and Read 2008). AMF are obligate symbionts that, via specialised structures (arbuscules) within the roots of their hosts, enhance plant uptake of macronutrients, particularly phosphorus and nitrogen (Hodge 2016; Hodge and Storer 2015; Javot et al. 2007; Leigh et al. 2009; Thirkell et al. 2016). In return, AMF are wholly dependent on the host plant for their carbon provision, usually acquiring around 10% of host photosynthate (Bago et al. 2000). These nutrient and carbon exchanges modify the nutrient status of the host plant, but also affect the allocation to secondary metabolites and other resistance mechanisms (Cameron et al. 2013; Minton et al. 2016). For example, colonisation by AMF has been shown to increase plant resistance to below-ground antagonists such as parasitic plants (Li et al. 2012) and plant parasitic nematodes (Rodriguez-Echeverria et al. 2009), as well as to generalist below-ground herbivores (Vannette and Rasmann 2012). Plant interactions with above-ground herbivores can also be affected by AMF colonisation (van der Putten et al. 2001), although usually to a lesser degree than below-ground (Van Wees et al. 2008), and outcomes vary with AMF species identity, herbivore diet breadth and/or feeding mode (Koricheva et al. 2009).

Early studies examining the impact of AMF colonisation on the chemical protection of plant tissues usually targeted specific chemical groups, limiting their power to detect the full range of induced chemical responses to AMF presence (de Deyn et al. 2009; Eftekhari et al. 2012). More recently, non-targeted metabolomic approaches have been used to investigate the effects of AMF on a wide range of metabolites (the metabolome) in foliar

tissues (Fester et al. 2011; Schweiger et al. 2014; Schweiger and Muller 2015). Again, several of these did not measure effects on secondary metabolites, so much remains to be discovered about the potential mechanisms by which AMF colonisation of plant roots can affect above-ground plant-herbivore interactions (Gange et al. 2012; Schweiger and Muller 2015). Similarly, the impact of AMF colonisation on the root metabolome has been little studied, although Schliemann et al (2008a) used a metabolomic approach to show that both primary and secondary metabolites differed between the roots of Medicago truncatula L. individuals with and without AMF colonisation, and Saia et al (2015) studied impacts of colonisation on the amino acid content of wheat. However, neither of these studies examined above-ground foliar impacts, whilst none of the foliar studies (Fester et al. 2011; Schweiger et al. 2014; Schweiger and Muller 2015) examined root tissue. Therefore, hitherto there has been no metabolomic investigation into the concomitant above- and below-ground changes in plant secondary chemistry following AMF colonisation, an important omission given the growing interest in above-belowground links and systemic defence responses (Johnson et al. 2016). This study addresses this knowledge gap by using non-targeted metabolomics profiling techniques to examine the simultaneous effects of arbuscular mycorrhizal (AM) colonisation on plant secondary chemistry in foliar and root tissues of ragwort (Senecio jacobaea L.), a species which forms symbiotic associations with AMF (Gange et al. 2002; Reidinger et al. 2012). We used experimental and untargeted metabolomic approaches to test the effect of AMF (*Rhizophagus irregularis*) colonisation on the secondary chemistry of ragwort roots and shoots. We aimed to identify the chemicals induced or repressed in host root and shoot tissues as a result of AM colonisation. Effects of colonisation on host metabolites might be expected to differ between the two tissue types given the established differences in local vs systemic responses to AMF (Van Wees et al. 2008), and, whilst recent

research suggests high specificity in leaf metabolomic responses to AMF (Schweiger et al.

2014), this is as yet untested in roots. Hence, we address two questions:

1) How does colonisation by AMF affect the profiles of secondary metabolites in ragwort and are the impacts more pronounced in root vs. foliar tissue?

2) Are the metabolites induced by AM colonisation similar to those identified in other plant-fungal systems, or is there evidence of novel compounds, potentially indicating a degree of host specificity? 

# MATERIALS AND METHODS

### Microcosm experiment

Ragwort plants were grown from seeds collected from a single plant on the University of Sussex campus (Lat: 50°52'18.6366"N, Long: 000°04'59.6860"W), sterilised in a 5% bleach solution and germinated on double autoclaved vermiculite. Thirty-one pots (1.7 L) received a double autoclaved (121°C; 60 min) growth medium comprising a 1:1 mixture of silver sand:Terragreen® (an attapulgite clay; Turf-Pro Ltd, UK) and bone meal (0.25g/L<sup>-1</sup>), a complex N and P source to encourage mycorrhizal development (Hodge and Fitter 2010). After mixing the growth media, all pots were twice flushed to saturation with water, aiming 43 110 to remove the pulse of nutrients released following sterilisation (Troelstra et al. 2001). Seedlings were assigned to replicate pots and randomly assigned to AMF (n = 16) and control 48 112 (n = 15) treatments. Each replicate in the AMF treatment was inoculated with 50 g (dry weight) of granular Rhizophagus irregularis (see Redecker et al. 2013 for recent changes in 53 114 nomenclature; obtained from PlantWorks, Kent, UK). Those replicates assigned to the 55 115 control treatment also had 50 g of this granular inoculate added, but double autoclaved (121°C; 60 min) to kill the *R. irregularis*, to control for the input of organic material with the inoculation (see Hodge 2001). In addition, non-AM pots received 100 ml of filtered 60 117

washings of the R. irregularis inoculum, but with AMF propagules removed to limit initial differences among pots in starter microbial communities (as Atkin et al. 2009). To account for any differences caused by the addition of water, 100 ml of deionised water was applied to the AMF-treated microcosms. Each microcosm received a weekly dose of 50 ml of half-strength Rorison's solution (Hewitt 1966), but with the plant-available phosphate removed to encourage AMF colonisation. The experiment ran for 10 weeks in a greenhouse maintained at 15-25°C with supplementary lighting (400W, high pressure sodium lamps) on a 16:8 L:D photoperiod. Microcosms were watered with tap water ad libitum.

Plants were destructively harvested after 10 weeks in treatment. From each replicate three new leaves (fully mature closest to the centre of the rosette) and two old leaves (closest to the edge of the rosette with no visible signs of senescence) were excised, weighed, immediately snap frozen in liquid nitrogen to halt enzymatic processes and stored in a -80°C freezer for metabolomic analysis. Two representative root samples (each ~10% of the total root mass) per plant were weighed and used for: i) plant metabolomic analysis (snap frozen then stored at  $-80^{\circ}$ C) and ii) to measure AMF colonisation (stored at  $4^{\circ}$ C until processed). Root sections were cleared in a 10% potassium hydroxide solution, ink-stained (Vierheilig et al. 1998), slide mounted and the presence or absence of AMF structures was recorded for a minimum of 100 root intersections using a compound microscope (x 200 magnification). Numbers of arbuscules, vesicles and root length colonized (RLC; the percentage of total intercepts where hyphae or other AM fungal structures were present) were recorded for each intersection (as Hodge 2003). The remaining shoot and root material was weighed and then oven-dried (60°C, 72 hours) to obtain dry biomass.

A homogenised sub-sample  $(0.1 \pm 0.01g)$ , of the root and shoot tissue taken from each replicate was mixed with 2mL of a 3:3:2 solvent mix of isopropanol, acetonitrile and water using a ball mill (Pulverisette 23. Fritsch, Germany). This solvent mixture extracts both polar metabolites (carbohydrates, amino acids) and non-polar metabolites (lipids) (Sana et al. 2010). Samples were spiked with two deuterated internal standards; 17β-estradiol 2,4,16,16d<sub>4</sub> sodium 3-sulfate (E2-d<sub>4</sub>-S, >99% D atom) and progesterone-2,2,4,6,6,17α,21,21,21-d<sub>9</sub> (P-d<sub>9</sub>, 98% D atom) (Cambridge Isotope Laboratories Inc. MA and CDN isotopes, Quebec, Canada, respectively). The sample was vortexed (1 min) and after overnight extraction at -20°C, samples were centrifuged, the supernatant removed and the pellet extracted with 2 mL of 80% methanol for 12 hours. The supernatants were combined and a 3 mL aliquot was evaporated to dryness under vacuum and redissolved in 160  $\mu$ l of methanol:water (3:1, v/v). The extract was filtered  $(0.2 \,\mu\text{m})$  prior to MS analysis. All extraction chemicals used were purchased from Rathburn Chemicals Ltd, Walkerburn, UK. Metabolites were profiled using ultraperformance liquid chromatography quadrupole time-of-flight mass spectrometry (UPLC-QTOFMS) (Waters, Manchester, UK). Aliquots of plant extracts were injected on to an Acquity UPLC BEH C18 column (1.7 µm particle size, 2.1 x 100 mm, Waters, UK). Metabolites were separated using a water-formic acid and acetonitrile gradient as follows: 0-9.0 min, from 0 to 30.0% acetonitrile; 9.0-15.0 min, from 30.0 to 100% acetonitrile; 15.0 to 23.0 min, 100% acetonitrile. The flow rate was 0.2 mL

161 min<sup>-1</sup> and the column temperature was 30°C. The injection volume of plant extracts was 20 162  $\mu$ l.

Metabolites were detected in both positive and negative ESI modes using a Micromass TOF-MS system (Waters, Manchester, UK). The mass spectrometer was tuned to 9000 mass resolution and data collected in full scan mode from 100 to 1200 *m/z*. The

collision gas used was argon, a constant collision energy of 10eV was used for all experiments, and the TOF penning pressures ranged from  $4.63 \times 10^{-7}$  to  $4.83 \times 10^{-7}$  mbar. Capillary voltage was 2.6 in positive mode and -2.9 in negative mode. In positive mode, the cone voltage was set at 36V, and the multiplier voltage was set at 654V. In negative mode the cone and multiplier voltage were set at 35V and 550V respectively. Desolvation N2 gas flow was set at 401L h<sup>-1</sup> for both ionisation modes.

MarkerLynx software (V 4.1, Waters, Manchester, UK) was used to align, normalise and remove isotopic peaks from the metabolomic profiles. Each metabolite signal was the description of an analyte using its specific retention time (r.t.) and mass-to-charge ratio (m/z). Using SIMCA-P multivariate analysis software (Umetrics UK Ltd, Windsor, UK) the data were pareto scaled, log-transformed and modelled using partial least square-discriminate analysis (PLS-DA) for > 2 classes, or orthogonal partial least-square discriminate analysis (OPLS-DA) for comparison of just 2 classes followed by examination of the loading plots to detect MS signals associated with AMF colonisation (Liland 2011; Trygg and Wold 2002). For all multivariate analyses, the explained variation  $(R^2Y)$  and the predictive power of the model  $(Q^2)$  were examined to assess the performance of the models. 

Biochemical markers associated with AM colonisation were extracted from 'S'-plots derived from the OPLS-DA models (Wiklund et al. 2008). The identity of biochemical markers was determined from their accurate mass composition and isotopic fit using the elemental composition tool from the MassLynx software (V 4.1, Waters, Manchester, UK). Fragmentation data, obtained from collision induced dissociation (CID) using QTOFMS (collision energy between 20 and 50eV) was used to confirm the putative identity of the markers.

Analyses of pyrrolizidine alkaloids in metabolomic profiles of tissue extracts. 

Metabolomic profiles of root extracts acquired in positive electrospray ionization (ESI) mode were dominated by highly abundant pyrrolizidine alkaloids (PAs) saturating the detector response. PAs are the principal secondary metabolites in Senecio species, synthesised in the roots (Hartmann et al. 1989) before being transported to the shoots (Hartmann and Dierich 1998) where they are effective defences against generalist herbivores (Macel et al. 2005; Narberhaus et al. 2005; Thoden et al. 2009). In order to determine whether PA profiles changed as a result of AM colonisation, an additional analysis of a 0.5 µl injection of root extracts was undertaken to reduce the highly abundant PA signals. The chromatograms (of both root and shoot extracts) were manually searched, and the relative abundance of observed signals corresponding to PA metabolites was determined relative to the internal standard using MassLynx peak integration software. PAs were assigned putative identities as described above. Where possible, authenticated standards were purchased to confirm PA identity: a mixed standard of PAs seneciphylline and senecionine (Carl Roth, GmbH & Co, Karlsruhe, Germany) and a retrorsine N-oxide standard (PhytoLab GmbH & Co. KG, Nürnberg, Germany). Details of the MS fragmentation data used for the identification of each metabolite are outlined in the supporting information.

### Data analysis

Adjusted fresh weights ((total fresh weight/100)\*water content) of plant tissues were calculated to account for a significant between-treatment difference in root and shoot water content (water content = (fresh weight-dry weight)/(fresh weight)\*100). Statistical analyses of differences in plant biomass, biochemical markers and AM colonisation between treatments were performed in SPSS v11. Where the residuals of the data met the assumptions, either before or following transformations, of parametric models (normality, homogeneity of variance), then one-way ANOVA (for plant biomass) or t-tests (for

percentage RLC, metabolite and PA concentrations) were used. For non-parametric data, Mann-Whitney U tests were employed. Bonferroni corrections were used to account for the increased false discovery rate (FDR) associated with the multiple testing of metabolic datasets containing many thousands of metabolite signals (Broadhurst and Kell 2006). A less conservative Benjamini and Hochberg FDR correction was applied to the univariate tests of PA concentrations (Benjamini et al. 2001), as this was more appropriate for these much smaller datasets. The association between blumenol metabolites and percentage RLC were investigated using Spearman rank-order correlations, as the data did not meet the assumptions of parametric tests.

# 226 RESULTS

# 227 AMF colonisation and plant biomass

Mean percent RLC was  $47 \pm 4\%$  in the *R. irregularis* colonised plants. Frequency of arbuscules was  $16 \pm 1\%$  and vesicles  $13 \pm 2\%$  respectively. There was no AMF colonisation of the non-AM controls. Shoot and root water content were higher in control plants than in *R. irregularis* colonised plants, though plant biomass was unaffected by *R. irregularis* colonisation (Table S1).

234 Root metabolome

The metabolite profiles of root extracts from AM colonised and non-AM control plants were clearly separated, indicating a shift in chemical composition of plants in the different treatment groups; (see Figure 1a for the positive ESI mode dataset, and a similar scores plot was obtained for the negative ESI mode in Figure S1a). OPLS-DA of both datasets (positive and negative ESI modes) produced models that explained a large amount of the variation in metabolite profiles between treatments (negative:  $R^2Y=0.99$ , positive:  $R^2Y=0.98$ ) with good

predictability (negative:  $Q^2 = 0.53$ , positive:  $Q^2 = 0.48$ ). This separation between treatment classes was driven by increased concentrations of 33 metabolites associated with AM colonisation (threshold values after Bonferroni adjustment: positive mode:  $P < 5.20 \times 10^{-7}$ ; negative mode:  $P < 5.01 \times 10^{-7}$ ). The fold increase in concentration of the 33 metabolites associated with AM roots was between two to many thousands (up to 37,995-fold in one instance). Eight of these metabolites were assigned identities (Table 1); the remaining metabolites (n = 25) are listed in Table S2. Blumenol standards are not commercially available, so identifications are derived from comparisons with published mass spectra (cited in Strack and Fester 2006; Peipp et al. 1997; Schliemann et al. 2008b); see supporting information (Note 1, Figure S2 and Table S3) for further details. 

Seven of the identified metabolite signals were conjugates of C<sub>13</sub> cyclohexenones, i.e. blumenol apocarotenoids (metabolites 1-7, Table 1, Figure 2). Metabolites 1 and 4 were identified as blumenols with a 13-hydroxyblumenol C moiety ( $C_{13}H_{23}O_3$ ). Metabolites 2, 3, 5, 6 and 7 were identified as blumenols with a blumenol C moiety  $(C_{13}H_{23}O_2)$ . Whilst identified metabolites 1-7 share common blumenol structures, they differed in the nature of conjugation (Figure 2). Metabolites 1, 2 and 5 were conjugated with glycoside sugars, whereas metabolites 4 and 7 were conjugated with malonylglycoside sugars. Metabolites 3 and 6 were conjugated with an uronide group consistent with a glucuronide (Strack and Fester 2006), and so were identified as a glycosyl-glucuronide and a malonylglycoside-glucuronide conjugate respectively. Metabolite 6, blumenol C malonylglycosyl-glucuronide, is a hitherto unreported structure, whereas metabolites 1-5 and 7 have been found to be associated with AM colonisation in other species (Peipp et al. 1997; Schliemann et al. 2008b). All blumenols were present as their molecular ions, which were either [M+H]<sup>+</sup> or [M-H]<sup>-</sup> in positive or negative ESI, respectively. In positive ESI, some metabolites were also present as the sodium adduct of the molecular ion (Table S3). Of those blumenols identified,

two were observed in both positive and negative ESI modes (metabolites 1 & 6, Table 1).
The concentration changes of the identified blumenols corresponded to large differences,
with fold increases ranging from 250 to over 8000 in AM plants. The final identified
metabolite (number 8) was determined to be hexadecenoyl-glycero-phosphocholine, a
lysophospholipid.

All blumenol apocarotenoids were positively associated with increasing percentage RLC in the AM colonised plants (Spearman correlation coefficient 0.523-0.711,  $P \le 0.03$ -0.002, Table 2). Levels of hexadecenoyl-glycero-phosphocholine were not significantly correlated with RLC (P  $\ge$  0.11). A number of unidentified metabolites were detected that were associated with AM colonisation (Table S2) and these were likely to be conjugated phenolic compounds, two additional conjugated blumenols and (in one case) a phospholipid. Based on their empirical formula, none of the metabolites (except for the phospholipid) were present in the plant metabolite databases and so potentially represent new structures. However, their concentrations were too low to allow further structural characterisation. Changes in the concentrations of primary metabolites were not observed as a result of AM colonisation, but this was most likely due to the fact that some structures such as lipids and carotenoids were not have been detected by ESI and small organic molecules, including alkaloid precursors, would not have been retained on the HPLC column. 

285 Shoot metabolome

PLS-DA models of the shoot dataset from positive ESI analysis revealed good predictability ( $Q^2 = 0.66$ ) and a high degree of explained variation (total  $R^2Y = 0.99$  for six latent variables) (Figure 1b). PLS-DA recorded in negative ESI mode revealed similar models that explained a high level of variation in shoot chemistry ( $R^2Y = 0.62$ , 4 latent variables) with lower predictability ( $Q^2 = 0.36$ ) than in positive mode (Figure S1b). In both positive and negative

modes, the first latent variable separated the metabolomes of old and new leaves. The second, whilst separating AM and non-AM control groups, only explained a relatively small amount of the observed variation in metabolite profiles between these treatments. OPLS-DA models with the highest predictability ( $R^2Y=0.989$ ,  $Q^2=0.431$ ) were associated with the positive ESI profiles of old leaves from the different treatment groups. However, in all OPLS-DA models of shoot tissues, no metabolites were found to vary significantly in association with AMF colonisation of ragwort. 

# Concentrations of PA metabolites in the root and shoot metabolome.

Analysis of the UPLC-QTOFMS chromatograms from the metabolomics study revealed that signals of PA metabolites were saturated in samples of the root extracts. Analysis of lower volumes of samples revealed 19 PA signals that could be resolved in the metabolomic profiles of roots, however more PA structures (potentially up to 33 metabolites, Kostkeno et al. 2013) were likely to be present, which were not detected due to masking of their signals by the most abundant PA signals. The concentration of individual PAs within a species or genotype varies greatly in both roots and shoots (Cheng et al. 2011a). Of the detected PAs, four were significantly increased in AM plants (Table 3). The PAs that responded to AMF colonisation were identified as senecionine, jacoline N-oxide, jaconine N-oxide, and usaramine N-oxide. Of these four PAs, only levels of senecionine were significantly correlated with RLC (Spearman correlation coefficient -0.560,  $P \le 0.02$ ). Details of the identification of the different PA structures are given in supporting information. The relative abundance of PAs was also determined in the metabolomics datasets of the shoot extracts. A total of 17 abundant PA structures were resolved in the metabolomics profiles of the shoot extracts out of a potential of 29 PA metabolites (Kostkeno et al. 2013). However, there were no significant differences in concentrations (after FDR corrections) of

PA signals between control and AMF-colonised treatment groups which was consistent with
the lack of discriminating metabolites observed in the above study on the shoot metabolome
(Table S4).

320 DISCUSSION

This study, the first to use a non-targeted metabolomic approach to assess the effect of AMF colonisation on host plant secondary chemistry in both roots and shoots, showed that colonisation by AMF could increase the concentrations of 33 metabolites in roots by between two and many thousand fold. The extent of AMF colonisation was positively correlated with levels of compounds thought to be associated with establishing and maintaining AMF colonisation, specifically seven C<sub>13</sub> cyclohexenone blumenol structures, including one with a hitherto unidentified structure. In addition, AMF colonisation affected the levels of the main anti-herbivore defences in root tissues, with four PAs significantly increased in concentration. Despite these below-ground changes, no significant changes in the metabolome of above-ground tissues were detected in response to AMF colonisation of roots, although there were clear metabolomic differences between leaf tissues of different ages. This is noteworthy as had changes been found in aboveground tissues then it would indicate the AM fungus had a direct impact upon the plant. However, as differences were only found in the 'mycorrhizal' (literally, 'fungus-root') root tissue, an unknown fraction of the metabolomics response observed here, and in other studies (e.g. Rivero et al. 2015; Saia et al. 2015), may be due directly to the fungal partner. Experimentally distinguishing between responses driven by the AMF, the root and the mycorrhizal root is hampered by the fact that many of these fungal symbionts cannot be grown in the absence of a host plant (Hodge et al. 2010). Nevertheless, our results demonstrate that colonisation of roots by AMF results in clear metabolomics differences compared to the un-colonised condition, a significant finding given that c. two-

thirds of land plants form this type of close symbiotic association (Hodge and Fitter 2013;
Smith and Read 2008).

The process of initiating and maintaining the plant-AMF relationship involves a range of metabolic changes within the host (Fester et al. 2011; Schliemann et al. 2008a), and the bumenol metabolites we detected in our non-targeted analysis are of particular significance, as they are thought to be involved in the chemical signalling underpinning the plant-AMF interaction (Walter et al. 2010; Maier et al. 1995). It has been suggested that blumenols and C14 polyene apocarotenoids have a role in maintaining the plant-AMF interaction once formed (Fester et al. 2002). For instance, concentrations of root C14 polyenes and blumenols increase (Peipp et al. 1997; Schliemann et al. 2006; 2008b) as the plant-AMF interaction progresses, particularly in root cells hosting AMF structures (Fester et al. 2002). Mutant plants, where blumenol production is reduced, are associated with an increased amount of dead and degenerating mycorrhizal structures (Floss et al. 2008), providing further evidence that these blumenol C13 cyclohexenones may have a role in the persistence of AMF colonisation (Walter et al. 2010).

In support of this hypothesis, and in accordance with other studies (Maier et al. 2000; Schliemann et al. 2008a), the concentrations of the seven blumenols we identified were all positively related to levels of AMF colonisation. One of these induced blumenols (metabolite 6) was conjugated with a malonyl-sugar and glucuronic acid moiety, a combination that has never been reported before. Species-specific blumenol metabolites have been observed in a range of plant taxa (Fester et al. 2002, 2005; Maier et al. 2000; Peipp et al. 1997; Schliemann et al. 2006, 2008a), and have been suggested as a mechanism of species recognition between the AMF and its host (Strack and Fester 2006). This is a hypothesis we raise for further testing, but current data is sparse. So far, only a few other plant species, such as the legume Medicago truncatula colonised by Glomus intraradices (Schliemann et al. 2008a), tomato

colonised by Rhizophagus irregularis and Funneliformis mossae (Rivero et al. 2015) and wheat colonised by multiple AM species (Saia et al. 2015) have had the root tissue response to AMF colonisation been examined in an untargeted metabolomic approach. Until a wider variety of plant-AMF species combinations are tested, it is too early to conclude whether any species-specific signalling exists in AMF-plant interactions. It is notable, however, that one recent study (Schweiger et al. 2014) which compared the impacts of the AMF Rhizophagus irregularis on the leaf metabolome of five different plant species found that species-specific metabolic changes following colonisation far out-numbered more generic cross-species responses. Similarly, Rivero et al. (2015) detected fungus-specific aspects of the AM-associated changes in the tomato metabolome.

Despite substantial changes to the root metabolome induced by AMF colonisation, there were no concomitant shifts in the aboveground metabolome. Conceivably, changes in the shoot metabolome may have been too small or inconsistent to have been detected, but this seems unlikely given that our UPLC-QTOFMS methodology clearly discriminated between the metabolomes of old and new leaves (Figure 1b), and that we have used the same technique to successfully detect differences in the metabolomes of above-ground tissues in ragwort previously (Hartley et al. 2012). Previous studies have demonstrated that AMF-mediated changes occur in above-ground plant tissues, with effects on gene expression (Taylor and Harrier 2003) and on herbivore performance (Koricheva et al. 2009), but changes in foliar secondary metabolites in response to colonisation seem more inconsistent. It has been suggested that induction of chemical defences by beneficial microbes are localised rather than systemic (Van Wees et al. 2008), which may explain why the increases in PA concentration in AMF colonised ragwort we observed were confined to the roots. Increases in PA concentrations in the roots may even reduce PA levels in the shoots. Although our study did not detect such a reduction in foliar allocation to PAs, Reidinger et al. (2012) showed that

leaf concentrations of both total PAs and jacoline were negatively correlated with AMFcolonisation of ragwort in the field.

The increased PA levels that we found in AMF colonised roots could impact on trophic interactions. AMF colonisation has been shown to have a negative influence on generalist below-ground herbivores such as parasitic nematodes (Li et al. 2006; Rodriguez-Echeverria et al. 2009) and beetle larvae (Gange et al. 1994), but positive effects on the root-feeding larvae of specialist weevils (Currie et al. 2011). Overall, generalist herbivores, whether above or below ground, seem more affected by AMF colonisation of their host plants than specialists (Gange et al. 2012; Koricheva et al. 2009). In ragwort, the highly toxic PAs deter generalist herbivores (Joosten and van Veen 2011; Thoden et al. 2009), whilst specialist herbivores have the capacity to use PAs to locate and select hosts (Cheng et al. 2013), to detoxify them, and even sequester these compounds for their own defence against predators (Beuerle et al. 2007; Narberhaus et al. 2003). Most previous studies have demonstrated these effects on above-ground herbivores (e.g. Cheng et al. 2011b; Wei et al. 2015); far less is known about the impacts of PAs on root-feeding herbivores (Cheng et al. 2011a) and we have found no published studies on how AMF colonisation modifies those effects. It has been shown that root damage increases PA levels (Hol et al. 2004) and that PAs affect root fungi and the composition of the rhizosphere (Hol and van Veen 2002; Kowalchuk et al. 2006), so there is potential for significant ecological consequences deriving from the changes in PAs we observed in AM roots.

In summary, this study has shown that colonisation by the AMF species *R*. *irregularis* causes numerous and significant changes in the root metabolome of ragwort, but not in the shoot metabolome. In root tissues, the key changes were in groups of chemicals associated with AMF-plant signalling (blumenols) and anti-herbivore defence (PAs). We showed strong correlations between the levels of blumenols and AMF colonisation,

416 supporting the idea that these metabolites play a key role in the interactions between AMF 417 and their hosts. The discovery that one of these was a novel structure not previously reported 418 raises questions about the significance of structural modifications in species recognition and 419 signalling that would repay further study.

*Acknowledgements* - For advice and practical assistance with the metabolomic analysis we would like to thank Dr Raghad Al-Salhi and Ms Julia Horwood. Dr Stefan Reidinger and Dr Kate Storer provided mycorrhizal cultures and advice on experimental procedures. We also thank the University of Sussex and the Centre for Ecology and Hydrology (project NEC03605) for co-funding this project.

- 428 REFERENCES
- Atkin OK, Sherlock D, Fitter AH, Jarvis S, Hughes JK, Campbell C, Hurry V, Hodge A,
  (2009) Temperature dependence of respiration in roots colonized by arbuscular
  mycorrhizal fungi. New Phytol 182:188-199. doi:10.1111/j.1469-8137.2008.02727.x
  Bago B, Pfeffer PE, Shachar-Hill Y (2000) Carbon metabolism and transport in arbuscular
  mycorrhizas. Plant Physiol 124:949-957. doi:10.1104/pp.124.3.949
  Benjamini Y, Drai D, Elmer G, Kafkafi N, Golani I (2001) Controlling the false discovery
  rate in behavior genetics research. Behav Brain Res 125:279-284. doi:10.1016/s01664328(01)00297-2
  - Beuerle T, Theuring C, Klewer N, Schulz S, Hartmann T (2007) Absolute configuration of the creatonotines and callimorphines, two classes of arctiid-specific pyrrolizidine alkaloids. Insect Biochem Mol Biol 37:80-89. doi:10.1016/j.ibmb.2006.10.005

	440	Broadhurst DI, Kell DB (2006) Statistical strategies for avoiding false discoveries in
1 2 3	441	metabolomics and related experiments. Metabolomics 2:171-196.
4 5	442	doi:10.1007/s11306-006-0037-z
6 7 8	443	Cameron DD, Neal AL, Van Wees SCM, Ton J (2013) Mycorrhiza-induced resistance: more
9 10	444	than the sum of its parts? Trends Plant Sci 18:539-545.
11 12 13	445	doi:10.1016/j.tplants.2013.06.004
14 15	446	Cheng D, Kirk H Mulder PPJ Vrieling K Klinkhamer PGL (2011a) Pyrrolizidine alkaloid
16 17 18	447	variation in shoots and roots of segregating hybrids between Jacobaea vulgaris and
19 20	448	Jacobaea aquatica. New Phytol 192: 1010-1023 doi: 10.1111/j.1469-
21 22	449	8137.2011.03841.x
23 24 25	450	Cheng D, Kirk H, Vrieling K, Mulder PPJ, Klinkhamer PGL (2011b) The relationship
26 27	451	between structurally different pyrrolizidine alkaloids and western flower thrips
28 29 30	452	resistance in F-2 hybrids of Jacobaea vulgaris and Jacobaea aquatica. Journal of
31 32	453	Chemical Ecology 37:1071-1080. doi:10.1007/s10886-011-0021-6
33 34 35	454	Cheng D, van der Meijden E, Mulder PPJ, Vrieling K, Klinkhamer PGL (2013) Pyrrolizidine
36 37	455	alkaloid composition influences cinnabar moth oviposition preferences in Jacobaea
38 39 40	456	hybrids. Journal of Chemical Ecology 39:430-437. doi:10.1007/s10886-013-0257-4
41 42	457	Currie AF, Murray PJ, Gange AC (2011) Is a specialist root-feeding insect affected by
43 44	458	arbuscular mycorrhizal fungi? Appl Soil Ecol 47:77-83.
43 46 47	459	doi:10.1016/j.apsoil.2010.12.002
48 49	460	de Deyn GB, Biere A, van der Putten WH, Wagenaar R, Klironomos JN (2009) Chemical
50 51 52	461	defense, mycorrhizal colonization and growth responses in <i>Plantago lanceolata</i> L.
53 54	462	Oecologia 160:433-442. doi:10.1007/s00442-009-1312-2
55 56 57	463	Eftekhari M, Alizadeh M, Ebrahimi P (2012) Evaluation of the total phenolics and quercetin
58 59	464	content of foliage in mycorrhizal grape (Vitis vinifera L.) varieties and effect of
60 61		
ь2 63 64		19
65		

1	465	postharvest drying on quercetin yield. Ind Crop Prod 38:160-165.
1 2 3	466	doi:10.1016/j.indcrop.2012.01.022
4 5 6	467	Fester T, Hause B, Schmidt D, Halfmann K, Schmidt J, Wray V, Hanse G, Strack D (2002)
6 7 8	468	Occurrence and localization of apocarotenoids in arbuscular mycorrhizal plant roots.
9 10	469	Plant Cell Physiol 43:256-265. doi:10.1093/pcp/pcf029
11 12 13	470	Fester T, Fetzer I, Buchert S, Lucas R, Rillig MC, Hartig C (2011) Towards a systemic
14 15	471	metabolic signature of the arbuscular mycorrhizal interaction. Oecologia 167:913-
16 17 18	472	924. doi:10.1007/s00442-011-2037-6
19 20	473	Fester T, Wray V, Nimtz M, Strack D (2005) Is stimulation of carotenoid biosynthesis in
21 22 22	474	arbuscular mycorrhizal roots a general phenomenon? Phytochem 66:1781-1786.
23 24 25	475	doi:10.1016/j.phytochem.2005.05.009
26 27	476	Floss DS, Hause B, Lange PR, Kuster H, Strack D, Walter MH (2008) Knock-down of the
28 29 30	477	MEP pathway isogene 1-deoxy-D-xylulose 5-phosphate synthase 2 inhibits formation
31 32	478	of arbuscular mycorrhiza-induced apocarotenoids, and abolishes normal expression of
33 34 35	479	mycorrhiza-specific plant marker genes. Plant J 56:86-100. doi:10.1111/j.1365-
36 37	480	313X.2008.03575.x
38 39 40	481	Gange AC, Bower E, Brown VK (2002) Differential effects of insect herbivory on arbuscular
41 42	482	mycorrhizal colonization. Oecologia 131:103-112. doi:10.1007/s00442-001-0863-7
43 44	483	Gange AC, Brown VK, Sinclair GS (1994) Reduction of black vine weevil larval growth by
43 46 47	484	vesicular-arbuscular mycorrhizal infection. Entomol Exp Appl 70:115-119.
48 49	485	Gange AC, Eschen R, Schroeder V (2012) The soil microbial community and plant foliar
50 51 52	486	defences against insects. In: Iason GR, Dicke M and Hartley SE (eds) The ecology of
53 54	487	plant secondary metabolites: from genes to global processes. Cambridge University
55 56 57	488	Press, New York, pp 170-189
58 59		
60 61		
63 64		20
65		

1	489	Hartley SE, Gange AC (2009) Impacts of plant symbiotic fungi on insect herbivores:
2 3	490	mutualism in a multitrophic context. In: Annual Review of Entomology, vol 54, Palo
4 5 6	491	Alto, pp 323-342. doi:10.1146/annurev.ento.54.110807.090614
6 7 8	492	Hartley SE, Eschen R, Horwood JM, Robinson LA, Hill EM (2012) Plant secondary
9 10	493	metabolites and the interactions between plants and other organisms: the potential of a
11 12 13	494	metabolomic approach. In: Iason GR, Dicke M and Hartley SE (eds) The ecology of
14 15	495	plant secondary metabolites: from genes to global processes, Cambridge University
16 17 18	496	Press, New York, pp. 204-225
19 20	497	Hartmann T, Dierich B (1998) Chemical diversity and variation of pyrrolizidine alkaloids of
21 22 23	498	the senecionine type: biological need or coincidence? Planta 206:443-451.
23 24 25	499	doi:10.1007/s004250050420
26 27	500	Hartmann T, Ehmke A, Eilert U, Vonborstel K, Theuring C (1989) Sites of synthesis,
20 29 30	501	translocation and accumulation of pyrrolizidine alkaloid N-oxides in Senecio vulgaris
31 32	502	L. Planta 177:98-107. doi:10.1007/bf00392159
33 34 35	503	Hewitt EJ (1966) Sand and water culture methods used in the study of plant nutrition. CAB,
36 37	504	Farnham Royal, UK.
38 39 40	505	Hodge A (2001) Arbuscular mycorrhizal fungi influence decomposition of, but not plant
41 42 43 44	506	nutrient capture from, glycine patches in soil. New Phytol 151:725-734.
	507	doi:10.1046/j.0028-646x.2001.00200.x
45 46 47	508	Hodge A (2003) N capture by Plantago lanceolata and Brassica napus from organic
48 49	509	material: the influence of spatial dispersion, plant competition and an arbuscular
50 51 52	510	mycorrhizal fungus. J Exp Bot 54:2331-2342. doi:10.1093/jxb/erg249
53 54	511	Hodge A (2016) Accessibility of inorganic and organic nutrients for mycorrhizas. In: Johnson
55 56 57	512	NC, Gehring C and Jansa J (eds) Mycorrhizal mediation of soil: fertility, structure,
58 59	513	and carbon storage. Elsevier, Amsterdam, pp 129-148
60 61 62 63 64 65		21

1	514	Hodge A, Fitter AH (2010) Substantial nitrogen acquisition by arbuscular mycorrhizal fungi
2 3	515	from organic material has implications for N cycling. P Natl Acad Sci USA
4 5 6	516	107:13754-13759. doi:10.1073/pnas.1005874107
6 7 8	517	Hodge A, Fitter AH (2013) Microbial mediation of plant competition and community
9 10	518	structure. Funct Ecol 27:865-875. doi:10.1111/1365-2435.12002
11 12 13	519	Hodge A, Storer K (2015) Arbuscular mycorrhiza and nitrogen: implications for individual
14 15	520	plants through to ecosystems. Plant Soil 386:1-19. doi:10.1007/s11104-014-2162-1
16 17 18	521	Hodge A, Helgason T, Fitter AH (2010) Nutritional ecology of arbuscular mycorrhizal fungi.
19 20	522	Fungal Ecol 3:267-273. doi: 10.1016/j.funeco.2010.02.002
21 22 23	523	Hol WHG, Macela M, van Veen JA, van der Meijden E (2004) Root damage and
23 24 25	524	aboveground herbivory change concentration and composition of pyrrolizidine
26 27	525	alkaloids of Senecio jacobaea Basic Appl Ecol 5: 253–260. doi:
28 29 30	526	10.1016/j.baae.2003.12.002
31 32 33	527	Hol WHG, Van Veen JA (2002) Pyrrolizidine alkaloids from Senecio jacobaea affect fungal
34 35 36	528	growth. J Chem Ecol 28: 1763-1772. doi: 10.1023/A:1020557000707
37 38 39	529	Javot H, Pumplin N, Harrison MJ (2007) Phosphate in the arbuscular mycorrhizal symbiosis:
40 41	530	transport properties and regulatory roles. Plant Cell Environ 30:310-322.
42 43	531	doi:10.1111/j.1365-3040.2006.01617.x
45 46	532	Johnson SN, Clark KE, Hartley SE, Jones TH, McKenzie SW, Koricheva J (2012)
47 48	533	Aboveground-belowground herbivore interactions: a meta-analysis. Ecology 93:2208-
49 50 51	534	2215. doi: 10.1890/11-2272.1
52 53	535	Johnson SN, Erb M, Hartley SE (2016) Roots under attack: contrasting plant responses to
54 55 56	536	below- and aboveground insect herbivory. New Phytol 210:413-418.
57 58	537	doi:10.1111/nph.13807
59 60 61		
62 63		22
64 65		

1	538	Joosten L, van Veen JA (2011) Defensive properties of pyrrolizidine alkaloids against			
1 2 3	539	microorganisms. Phytochem Rev 10:127-136. doi:10.1007/s11101-010-9204-y			
4 5 6	540	Koricheva J, Gange AC, Jones T (2009) Effects of mycorrhizal fungi on insect herbivores: a			
7 8	541	meta-analysis. Ecology 90:2088-2097. doi:10.1890/08-1555.1			
9 10 11	542	Kostenko O, Mulder PJ, Martijn Bezemer T (2013) Effects of Root Herbivory on			
11 12 13	543	Pyrrolizidine Alkaloid Content and Aboveground Plant-Herbivore-Parasitoid			
14 15	544	Interactions in Jacobaea Vulgaris. J Chem Ecol 39:109-119. doi.org/10.1007/s10886-			
16 17 18	545	012-0234-3			
19 20	546	Kowalchuk GA, Hol WHG, van Veen JA (2006) Rhizosphere fungal communities are			
21 22 23	547	influenced by Senecio jacobaea pyrrolizidine alkaloid content and composition. Soil			
24 24 25	548	Biology & Biochemistry 38: 2852–2859 doi: 10.1016/j.soilbio.2006.04.043			
26 27 28	549	Leigh J, Hodge A, Fitter AH (2009) Arbuscular mycorrhizal fungi can transfer substantial			
29 30 31 32 33 34 35 36	550	amounts of nitrogen to their host plant from organic material. New Phytol 181:199-			
	551	207. doi:10.1111/j.1469-8137.2008.02630.x			
	552	Li AR, Smith SE, Smith FA, Guan KY (2012) Inoculation with arbuscular mycorrhizal fungi			
36 37 38	553	suppresses initiation of haustoria in the root hemiparasite Pedicularis tricolor. Ann			
39 40	554	Bot 109:1075-1080. doi:10.1093/aob/mcs028			
41 42 43	555	Li HY, Yang GD, Shu HR, Yang YT, Ye BX, Nishida I, Zheng CC (2006) Colonization by			
44 45	556	the arbuscular mycorrhizal fungus Glomus versiforme induces a defense response			
46 47 48	557	against the root-knot nematode Meloidogyne incognita in the grapevine (Vitis			
49 50	558	amurensis Rupr.), which includes transcriptional activation of the class III chitinase			
51 52	559	gene VCH3. Plant Cell Physiol 47:154-163. doi:10.1093/pcp/pci231			
53 54 55	560	Liland KH (2011) Multivariate methods in metabolomics - from pre-processing to dimension			
56 57	561	reduction and statistical analysis. Trends Anal Chem 30:827-841.			
58 59 60	562	doi:10.1016/j.trac.2011.02.007			
61 62 63 64 65		23			

1	563	Macel M, Bruinsma M, Dijkstra SM, Ooijendijk T, Niemeyer HM, Klinkhamer PGL (2005)
1 2 3	564	Differences in effects of pyrrolizidine alkaloids on five generalist insect herbivore
4 5 6	565	species. J Chem Ecol 31:1493-1508. doi:10.1007/s10886-005-5793-0
6 7 8	566	Maier W, Peipp H, Schmidt J, Wray V, Strack D (1995) Levels of a terpenoid glycoside
9 10	567	(blumenin) and cell wall-bound phenolics in some cereal mycorrhizas. Plant Physiol
11 12 13	568	109:465-470. doi:10.1104/pp.109.2.465
14 15	569	Maier W, Schmidt J, Nimtz M, Wray V, Strack D (2000) Secondary products in mycorrhizal
16 17 18	570	roots of tobacco and tomato. Phytochem 54:473-479. doi:10.1016/s0031-
19 20	571	9422(00)00047-9
21 22	572	Minton MM, Barber NA, Gordon LL (2016) Effects of arbuscular mycorrhizal fungi on
23 24 25	573	herbivory defense in two Solanum (Solanaceae) species. Plant Ecol Evol 149:157-
26 27	574	164. doi:10.5091/plecevo.2016.1176
28 29 30	575	Narberhaus I, Theuring C, Hartmann T, Dobler S (2003) Uptake and metabolism of
31 32	576	pyrrolizidine alkaloids in Longitarsus flea beetles (Coleoptera : Chrysomelidae)
33 34 35	577	adapted and non-adapted to alkaloid-containing host plants. J Comp Physiol B-
36 37	578	Biochem Syst Environ Physiol 173:483-491. doi:10.1007/s00360-003-0356-6
38 39 40	579	Narberhaus I, Zintgraf V, Dobler S (2005) Pyrrolizidine alkaloids on three trophic levels -
41 42	580	evidence for toxic and deterrent effects on phytophages and predators. Chemoecology
43 44	581	15:121-125. doi:10.1007/s00049-005-0302-z
43 46 47	582	Peipp H, Maier W, Schmidt J, Wray V, Strack D (1997) Arbuscular mycorrhizal fungus-
48 49	583	induced changes in the accumulation of secondary compounds in barley roots.
50 51 52	584	Phytochem 44:581-587. doi:10.1016/s0031-9422(96)00561-4
53 54	585	Redecker D, Schussler A, Stockinger H, Sturmer SL, Morton JB, Walker C (2013) An
55 56 57	586	evidence-based consensus for the classification of arbuscular mycorrhizal fungi
58 59	587	(Glomeromycota). Mycorrhiza 23:515-531. doi:10.1007/s00572-013-0486-y
60 61 62 63 64 65		24

1	588	Reidinger S, Eschen R, Gange AC, Finch P, Bezemer TM (2012) Arbuscular mycorrhizal
1 2 3	589	colonization, plant chemistry, and aboveground herbivory on Senecio jacobaea. Acta
4 5 6	590	Oecol 38:8-16. doi:10.1016/j.actao.2011.08.003
6 7 8	591	Rivero J, Gamir J, Aroca R, Pozo MJ, Flors V (2015) Metabolic transition in mycorrhizal
9 10	592	tomato roots. Front Microbiol 6: 598. doi: 10.3389/fmicb.2015.00598
11 12 13	593	Rodriguez-Echeverria S, de la Pena E, Moens M, Freitas H, van der Putten WH (2009) Can
14 15	594	root-feeders alter the composition of AMF communities? Experimental evidence from
16 17 18	595	the dune grass Ammophila arenaria. Basic Appl Ecol 10:131-140.
19 20	596	doi:10.1016/j.baae.2008.01.004
21 22	597	Saia S, Ruisi P, Fileccia V, Di Miceli G, Amato G, Martinelli F (2015) Metabolomics
23 24 25	598	suggests that soil inoculation with arbuscular mycorrhizal fungi decreased free amino
26 27	599	acid content in roots of durum wheat grown under N-limited, P-rich field conditions.
28 29 30	600	PLoS One 10:15. doi:10.1371/journal.pone.0129591
31 32	601	Sana TR, Fischer S, Wohlgemuth G, Katrekar A, Jung KH, Ronald PC, Fiehn O (2010)
33 34 35	602	Metabolomic and transcriptomic analysis of the rice response to the bacterial blight
36 37	603	pathogen Xanthomonas oryzae pv. oryzae. Metabolomics 6:451-465.
38 39 40	604	doi:10.1007/s11306-010-0218-7
41 42	605	Schliemann W, Ammer C, Strack D (2008a) Metabolite profiling of mycorrhizal roots of
43 44	606	Medicago truncatula. Phytochem 69:112-146. doi:10.1016/j.phytochem.2007.06.032
45 46 47	607	Schliemann W, Ammer C, Strack D (2008b) Metabolite profiling of mycorrhizal roots of
48 49	608	Medicago truncatula. Phytochem 69:1446-1447.
50 51 52	609	doi:10.1016/j.phytochem.2008.02.004
53 54	610	Schliemann W, Schmidt J, Nimtz M, Wray V, Thomas F, Strack D (2006) Accumulation of
55 56 57	611	apocarotenoids in mycorrhizal roots of Ornithogalum umbellatum. Phytochem
57 58 59	612	67:1196-1205. doi:10.1016/j.phytochem.2006.05.005
60 61 62 63 64 65		25

1	613	Schweiger R, Baier MC, Persicke M, Muller C (2014) High specificity in plant leaf metabolic
1 2 3 4 5 6	614	responses to arbuscular mycorrhiza. Nat Commun 5:11. doi:10.1038/ncomms4886
	615	Schweiger R, Muller C (2015) Leaf metabolome in arbuscular mycorrhizal symbiosis. Curr
7 8	616	Opin Plant Biol 26:120-126. doi:10.1016/j.pbi.2015.06.009
9 10 11	617	Smith SE, Read DJ (2008) Mycorrhizal symbiosis. Academic Press, San Diego
$12 \\ 13$	618	Strack D, Fester T (2006) Isoprenoid metabolism and plastid reorganization in arbuscular
14 15	619	mycorrhizal roots. New Phytol 172:22-34. doi:10.1111/j.1469-8137.2006.01837.x
16 17 18 19 20	620	Taylor J, Harrier LA (2003) Expression studies of plant genes differentially expressed in leaf
	621	and root tissues of tomato colonised by the arbuscular mycorrhizal fungus Glomus
21 22 23	622	mosseae. Plant MolBiol 51:619-629. doi:10.1023/a:1022341422133
23 24 25 26 27	623	Thirkell TJ, Cameron DD, Hodge A (2016) Resolving the 'nitrogen paradox' of arbuscular
26 27	624	mycorrhizas: fertilization with organic matter brings considerable benefits for plant
28 29 30 31 32	625	nutrition and growth. Plant Cell Environ 39:1683-1690. doi:10.1111/pce.12667
	626	Thoden TC, Boppre M, Hallmann J (2009) Effects of pyrrolizidine alkaloids on the
33 34 35	627	performance of plant-parasitic and free-living nematodes. Pest Manag Sci 65:823-
36 37	628	830. doi:10.1002/ps.1764
30 39 40 41 42 43 44 45 46 47	629	Troelstra SR, Wagenaar R, Smant W, Peters BAM (2001) Interpretation of bioassays in the
	630	study of interactions between soil organisms and plants: involvement of nutrient
	631	factors. New Phytol 150:697-706. doi:10.1046/j.1469-8137.2001.00133.x
	632	Trygg J, Wold S (2002) Orthogonal projections to latent structures (O-PLS). J Chemometr
48 49	633	16:119-128. doi:10.1002/cem.695
50 51 52	634	van der Putten WH, Vet LEM, Harvey JA, Wackers FL (2001) Linking above- and
53 54	635	belowground multitrophic interactions of plants, herbivores, pathogens, and their
55 56	636	antagonists. Trends Ecol Evol 16:547-554. doi:10.1016/s0169-5347(01)02265-0
58 59		
60 61		
62 63 64		26
65		

1	637	Van Wees SCM, Van der Ent S, Pieterse CMJ (2008) Plant immune responses triggered by
2 3	638	beneficial microbes. Curr Opin Plant Biol 11:443-448. doi:10.1016/j.pbi.2008.05.005
4 5	639	Vannette RL, Rasmann S (2012) Arbuscular mycorrhizal fungi mediate below-ground plant-
6 7 8	640	herbivore interactions: a phylogenetic study. Funct Ecol 26:1033-1042.
9 10	641	doi:10.1111/j.1365-2435.2012.02046.x
11 12 13	642	Vierheilig H, Coughlan AP, Wyss U, Piche Y (1998) Ink and vinegar, a simple staining
14 15	643	technique for arbuscular-mycorrhizal fungi. Appl Environ Microbiol 64:5004-5007.
16 17 18	644	Walter MH, Floss DS, Strack D (2010) Apocarotenoids: hormones, mycorrhizal metabolites
19 20	645	and aroma volatiles. Planta 232:1-17. doi:10.1007/s00425-010-1156-3
21 22 23	646	Wardle DA, Bardgett RD, Klironomos JN, Setala H, van der Putten WH, Wall DH (2004)
23 24 25	647	Ecological linkages between aboveground and belowground biota. Science 304:1629-
26 27	648	1633. doi:10.1126/science.1094875
28 29 30	649	Wei X, Vrieling K, Mulder PPJ, Klinkhamer PGL (2015) Testing the generalist-specialist
31 32	650	dilemma: the role of pyrrolizidine alkaloids in resistance to invertebrate herbivores in
33 34 35	651	Jacobaea species. J Chem Ecol 41:159- 167. doi: 10.1007/s10886-015-0551-4
36 37	652	Wiklund S, Johansson E, Sjostrom L, Mellerowicz EJ, Edlund U, Shockcor JP, Gottfries J,
38 39 40	653	Moritz T, Trygg J (2008) Visualization of GC/TOF-MS-based metabolomics data for
41 42	654	identification of biochemically interesting compounds using OPLS class models. Anal
43 44	655	Chem 80:115-122. doi:10.1021/ac0713510
45 46 47	656	
48	657	
49 50	658	
51	659	
52 53		
54		
55		
56		
58		
59		
60		
61 62		27
63		27
64		
65		

# **TABLE 1** METABOLITES IDENTIFIED IN RAGWORT ROOTS THAT WERE SIGNIFICANTLY INCREASED AFTER COLONISATION BY RHIZOPHAGUS IRREGULARIS

Metabolite	Observed	UPLC-	Putative	Theoretical	Putative identity	Fold	P-value <sup>b</sup>
number <sup>¶</sup>	ion $(m/z)$	TOFMS	formula	mass of ion		change <sup>a</sup>	
		r.t.					
1	389.2177	6.37	$C_{19}H_{33}O_8$	389.2175	13-hydroxyblumenol C glycoside [M+H] <sup>+</sup>	1151.2*	6.65 x 10 <sup>-9</sup>
1	387.2018	6.37	$C_{19}H_{33}O_8$	387.2018	13-hydroxyblumenol C glycoside [M-H] <sup>-</sup>	746.6*	6.65 x 10 <sup>-9</sup>
2	373.2228	7.23	$C_{19}H_{33}O_7$	373.2226	Blumenol C glycoside [M+H] <sup>+</sup>	8065.8	6.65 x 10 <sup>-9</sup>
3	547.2389	7.30	$C_{25}H_{39}O_{13}$	547.2391	Blumenol C glycosyl-glucuronide [M-H] <sup>-</sup>	1553.3	6.65 x 10 <sup>-9</sup>
4	475.2182	7.55	$C_{22}H_{35}O_{11}$	475.2179	13-hydroxyblumenol C malonylglycoside [M+H] <sup>+</sup>	489.7	2.99 x 10 <sup>-7</sup>
5	373.2222	8.20	$C_{19}H_{33}O_7$	373.2226	Blumenol C glycoside [M+H] <sup>+</sup>	5032.0	6.65 x 10 <sup>-9</sup>
6	635.2551	8.24	$C_{28}H_{43}O_{16}$	635.2551	Blumenol C malonylglycosyl-glucuronide [M+H] <sup>+</sup>	3101.3*	6.65 x 10 <sup>-9</sup>
6	633.2396	8.24	$C_{28}H_{41}O_{16}$	633.2395	Blumenol C malonylglycosyl-glucuronide [M-H] <sup>-</sup>	250.4*	6.65 x 10 <sup>-9</sup>
7	459.2232	9.38	$C_{22}H_{35}O_{10}$	459.2230	Blumenol C malonylglycoside [M+H] <sup>+</sup>	1109.2	6.65 x 10 <sup>-9</sup>
8	494.3247	14.20	$C_{24}H_{49}NO_7P$	494.3247	Hexadecenoyl-glycero-phosphocholine [M+H] <sup>+</sup>	7.86	2.00 x 10 <sup>-7c</sup>

<sup>¶</sup>Structures given in Figure 2.

r.t.= retention time

<sup>a</sup> Fold change indicates the concentration increase in roots colonised with AMF when compared to the concentrations observed in control plants.

<sup>b</sup> Significance determined using t-tests (<sup>c</sup>) or Mann-Whitney U tests (unmarked) after Bonferroni adjustments.

\* Differences in estimation of fold change between [M+H]<sup>+</sup> and [M-H]<sup>-</sup> signals for the same metabolite were due to presence of a Na adduct competing with the [M+H]<sup>+</sup> ion in positive ESI mode.

# **TABLE 2** SPEARMAN RANK-ORDER CORRELATIONS (N = 16) BETWEEN THE<br/>CONCENTRATION OF THE BLUMENOL METABOLITES MEASURED IN<br/>RAGWORT ROOTS AND RHIZOPHAGUS IRREGULARIS COLONISATION IN THE<br/>AMF-TREATED PLANTS

Metabolite structure	Putative identity	Correlation coefficient	P-value <sup>¶</sup>
(see Figure 3)			
1	13-hydroxyblumenol C glycoside	0.852	1.16 x 10 <sup>-9</sup>
2	Blumenol C glycoside	0.902	4.33 x 10 <sup>-12</sup>
3*	Blumenol C glycosyl-glucuronide	0.828	9.37 x 10 <sup>-9</sup>
4	13-hydroxyblumenol C	0.841	3.21 x 10 <sup>-9</sup>
	malonylglycoside		
5	Blumenol C glycoside	0.844	2.41 x 10 <sup>-9</sup>
6	Blumenol C malonylglycosyl-	0.832	6.39 x 10 <sup>-9</sup>
	glucuronide		
7	Blumenol C malonylglycoside	0.855	9.36 x 10 <sup>-10</sup>

\* Result from MS analysis in -ESI mode (required to detect metabolite 3); all other results from +ESI mode

<sup>¶</sup>All P-values significant after Bonferroni adjustment

Theoretical mass of ion	Putative formula	UPLC- TOFMS r.t.	Control	<i>G. intraradices</i> colonised	P-value <sup>a</sup>	Fold change <sup>b,c</sup>	Putative identity <sup>c</sup>
334.1654	C <sub>18</sub> H <sub>23</sub> NO <sub>5</sub>	6.00	49.7 (±5.3)	59.2 (±5.3)	0.188		
336.1811	C <sub>18</sub> H <sub>25</sub> NO <sub>5</sub>	6.86	211.7 (±17.7)	289.8 (±21.6)	3.63 x 10 <sup>-3</sup>	1.37	Senecionine
350.1604	$C_{18}H_{23}NO_6$	6.34	220.5 (±27.7)	223.2 (±21.6)	0.571		
	$C_{18}H_{23}NO_6$	9.01	24.5 (±3.2)	38.6 (±6.2)	0.054		
352.1760	$C_{18}H_{25}NO_6$	4.89	23.7 (±8.7)	52.0 (±14.2)	0.033		
	$C_{18}H_{25}NO_6$	5.82	76.1 (±11.3)	55.3 (±5.4)	0.216		
	$C_{18}H_{25}NO_6$	6.13	327.9 (±23.9)	303.8 (±17.9)	0.711		
	$C_{18}H_{25}NO_6$	6.99	48.8 (±2.5)	53.0 (±2.4)	0.029		
	$C_{18}H_{25}NO_6$	7.16	108.8 (±4.2)	113.0 (±3.5)	0.110		
366.1553	$C_{18}H_{23}NO_7$	4.29	193.2 (±46.1)	174.4 (±31.4)	0.598		
	$C_{18}H_{23}NO_7$	4.98	126.0 (±33.8)	141.9 (±39.5)	0.891		
368.1709	C <sub>18</sub> H <sub>25</sub> NO <sub>7</sub>	5.24	45.9 (±6.7)	65.2 (±10.2)	<b>4.81</b> x 10 <sup>-3</sup>	1.42	Usaramine N-oxide
	$C_{18}H_{25}NO_7$	5.62	93.8 (±20.2)	119.3 (±18.8)	0.231		
	$C_{18}H_{25}NO_7$	6.39	550.5 (±54.4)	530.8 (±38.6)	0.953		
370.1866	$C_{18}H_{27}NO_7$	5.94	24.0 (±2.2)	25.8 (±2.8)	0.379		
376.1760	$C_{20}H_{25}NO_{6}$	9.55	118.5 (±12.1)	118.4 (±14.3)	1.000		
386.1815	C <sub>18</sub> H <sub>27</sub> NO <sub>8</sub>	3.68	34.6 (±9.9)	105.6 (±16.0)	<b>2.01</b> x 10 <sup>-3</sup>	3.05	Jacoline N-oxide
392.1709	$C_{20}H_{25}NO_7$	9.77	196.5 (±26.17)	194.0 (±31.4	0.740		
404.1476	C <sub>18</sub> H <sub>26</sub> NO <sub>7</sub> Cl	5.22	39.3 (±9.73)	122.2 (±13.5)	5.87 x 10 <sup>-5</sup>	3.11	Jaconine N-oxide

# **TABLE 3** AVERAGE RELATIVE CONCENTRATIONS (±S.E.) OF THE PYRROLIZIDINE ALKALOID (PA) SIGNALS MEASURED IN THE POSITIVE ESI UPLC-TOFMS PROFILES OF CONTROL (N=15) AND *RHIZOPHAGUS IRREGULARIS* COLONISED (N=16) RAGWORT ROOTS

Relative concentrations were quantified as the ratio of the analyte signal: internal standard signal per 0.3 mg plant mass.

<sup>a</sup> Results that were significant after Benjamini and Hochberg corrections (Benjamini et al. 2001) are displayed in bold.

<sup>b</sup> Fold change indicates the concentration increase in roots colonised with AMF when compared to the concentrations observed in control plants.

<sup>c</sup> Only given for those PAs that differed significantly between treatment groups.

- FIGURE LEGENDS

4	FIG. 1 a) Orthogonal partial least square-discriminate analysis (OPLS-DA) scores
5	plot of the chemical profiles of root extracts from control and Rhizophagus irregularis
6	treated ragwort, b) Partial least square-discriminate analysis (PLS-DA) scores plot of
7	the chemical profiles of leaves from control and Rhizophagus irregularis treated
8	ragwort. Open and closed circles represent control and R. irregularis treated ragwort
9	plants, respectively. In b), blue symbols represent samples from new leaves and
10	orange symbols those from old leaves and the percentages of explained variation
11	$(R^2Y)$ modelled by the first two latent variables are displayed on the axes.
12	Both plots show datasets from samples profiled in positive ESI MS mode; data from
13	negative ESI mode revealed similar model characteristics (Figure S1).
14	
15	FIG. 2 Schematic structure of the identified blumenol apocarotenoids associated with
16	Rhizophagus irregularis colonisation of ragwort roots.
17	
18	
19	
20	
21	
22	





Metabolite	R1	R2
1	-CH <sub>2</sub> OH	-OH
2	-CH <sub>3</sub>	-OH
3	-CH <sub>3</sub>	$-C_6H_9O_7$
4	-CH <sub>2</sub> OH	-OCOCH <sub>2</sub> CO <sub>2</sub> H
5	-CH <sub>3</sub>	-OH
6	-CH <sub>3</sub>	-OCOCH <sub>2</sub> CO-C <sub>6</sub> H <sub>9</sub> O <sub>7</sub>
7	-CH <sub>3</sub>	-OCOCH <sub>2</sub> CO <sub>2</sub> H

Supplementary Material

Click here to access/download **Supplementary Material** Hill Hartley et al J Chem Ecol REVISED Supp Ino Resub.docx