



UNIVERSITY OF LEEDS

This is a repository copy of *Adversity-related injury in adolescence is associated with increased risk of suicide, alcohol and drug-related deaths in the decade following admission*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/129947/>

Version: Accepted Version

Article:

Friend, AJ orcid.org/0000-0001-9864-5605 (2018) Adversity-related injury in adolescence is associated with increased risk of suicide, alcohol and drug-related deaths in the decade following admission. *Archives of Disease in Childhood: Education and Practice Edition*, 103 (4). p. 223. ISSN 1743-0585

<https://doi.org/10.1136/archdischild-2017-313943>

© 2018, Article author(s) (or their employer(s) unless otherwise stated in the text of the article). All rights reserved. No commercial use is permitted unless otherwise expressly granted. This is an author produced version of a paper published in *Archives of Disease in Childhood: Education and Practice Edition*. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Declarative Title: Adversity-related Injury in Adolescence is Associated with Increased Risk of Suicide, Alcohol and Drug-Related Deaths in the Decade Following Admission

STUDY DESIGN

Design: Retrospective cohort study.

STUDY QUESTION

Setting: NHS hospitals in England.

Patients: 333,009 adolescents admitted following injury between 1 April 1997 and 31 March 2012.

Exposure: Adversity-related injury (defined as self-inflicted, drug related, alcohol related or violent injury) vs accident-related injury.

Outcomes: Risk and causes of death.

Follow-up period: Up to 10 years following initial admission.

Patient follow-up: Ascertained from Hospital Episode Statistics.

MAIN RESULTS:

Relative risk of death from various causes in the 10 years following initial admission was calculated for various subgroups. Results are summarised in table 1 below.

Gender	Subgroup	Adjusted HR (95% CI)	
		Suicide	Drug or alcohol-related death
Female	All adversity-related injury	4.54 (3.25 to 6.36)	4.71 (3.28 to 6.76)
	Self-inflicted injury	5.11 (3.61 to 7.23)	5.14 (3.50 to 7.55)
	Drug- or alcohol-related injury	4.55 (3.23 to 6.39)	4.52 (3.14 to 6.51)
	Violent injury	1.48 (0.73 to 2.98)	2.75 (1.47 to 5.17)
Male	All adversity-related injury	3.15 (2.73 to 3.63)	3.15 (2.73 to 3.63)
	Self-inflicted injury	6.20 (5.27 to 7.30)	5.91 (4.96 to 7.03)
	Drug- or alcohol-related injury	4.51 (3.89 to 5.24)	4.91 (4.24 to 5.73)
	Violent injury	1.43 (1.15 to 1.78)	1.78 (1.44 to 2.19)

Table 1: Relative risk of death by cause in adolescents admitted with adversity-related injury (all vs accident-related injury)

CONCLUSION: Risk of suicide was significantly increased in the 10 years following an adversity-related injury, and risk of drug- and alcohol-related death was similarly raised.

ABSTRACTED FROM: Herbert, A., Gilbert, R., Cottrell, D. and Li, L., 2017. Causes of death up to 10 years after admissions to hospitals for self-inflicted, drug-related or alcohol-related, or violent injury during adolescence: a retrospective, nationwide, cohort study. *The Lancet*

Adolescents admitted to hospital following episodes of self-inflicted, violent, drug-, or alcohol-related injury are well known to be at increased risk of death up to 10 years after presentation¹, however causes of death have not been previously explored in detail. Despite this, there has been a tendency for some treating doctors to be dismissive of these patients as "attention seekers" or "time wasters"².

This study describes the increased risk of suicide in patients who self-harm - which itself is not new or surprising³. The important findings of increased risk of drug- or alcohol-related death warrants further consideration and highlights the importance of discussing substance misuse when assessing young people admitted following episodes of self-harm. Equally, the high rates of suicide are described in patients who present following drug- or alcohol-related injury, suggesting that exploration of potential suicidal ideation may be indicated in this group as well as those who self-harm.

Those admitted following violent injury were reported to be at increased risk of both suicide and of drug- or alcohol-related death; a stark reminder of the effect of adversity on mental health.

The significant risk to future health, and indeed life, in adolescents presenting with adversity-related injury should concern all who are involved in the care of young people and prompt us to search for more effective harm reduction strategies.

1. Herbert, A., Gilbert, R., Gonzalez-Izquierdo, A., Pitman, A. and Li, L., 2015. 10-y risks of death and emergency re-admission in adolescents hospitalised with violent, drug-or alcohol-related, or self-inflicted injury: a population-based cohort study. *PLoS medicine*, 12(12), p.e1001931.
2. Saunders, K.E., Hawton, K., Fortune, S. and Farrell, S., 2012. Attitudes and knowledge of clinical staff regarding people who self-harm: a systematic review. *Journal of affective disorders*, 139(3), pp.205-216.

3. Hawton, K. and Harriss, L., 2007. Deliberate self-harm in young people: characteristics and subsequent mortality in a 20-year cohort of patients presenting to hospital. *The Journal of clinical psychiatry*.