

This is a repository copy of *Unravelling past impacts of climate change and land management on historic peatland development using proxy-based reconstruction, monitoring data and process modelling*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/130093/>

Version: Accepted Version

Article:

Heinemeyer, Andreas orcid.org/0000-0003-3151-2466 and Swindles, Graeme T. (2018) Unravelling past impacts of climate change and land management on historic peatland development using proxy-based reconstruction, monitoring data and process modelling. *Global Change Biology*. GCB-18-0338.R1. pp. 4131-4142. ISSN: 1354-1013

<https://doi.org/10.1111/gcb.14298>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Unravelling past impacts of climate change and land management on historic peatland development using proxy-based reconstruction, monitoring data and process modelling

Running head: Exploring historic peat development

Andreas Heinemeyer¹ & Graeme T. Swindles²

¹Stockholm Environment Institute, Environment Department, University of York, York, YO5NG, United Kingdom

²water@leeds, School of Geography, University of Leeds, LS2 9JT, United Kingdom

Corresponding author: Andreas Heinemeyer; xx44 1904 32 2991; andreas.heinemeyer@york.ac.uk

Keywords: water-table, testate amoebae, peatlands, MILLENNIA model, carbon budgets, carbon emissions, peat accumulation, peatland management

Type of paper: Primary Research Article

Abstract

Peatlands represent globally significant soil carbon stores that have been accumulating for millennia under water-logged conditions. However, deepening water-table depths (WTD) from climate change or human-induced drainage could stimulate decomposition resulting in peatlands turning from carbon sinks to carbon sources. Contemporary WTD ranges of testate amoebae (TA) are commonly used to predict past WTD in peatlands using quantitative transfer function models. Here we present, for the first time, a study comparing TA-based WTD reconstructions to instrumentally-monitored WTD and hydrological model predictions using the MILLENNIA peatland model to examine past peatland responses to climate change and land management. Although there was very good agreement between monitored and modelled WTD, TA-reconstructed water table was consistently deeper. Predictions from a larger European TA transfer function data set were wetter, but the overall directional fit to observed WTD was better for a TA transfer function based on data from northern England. We applied a regression-based offset correction to the reconstructed WTD for the validation period (1931-2010). We then predicted WTD using available climate records as MILLENNIA model input and compared the offset-corrected TA reconstruction to MILLENNIA WTD predictions over an extended period (1750-1931) with available climate reconstructions. Although the comparison revealed striking similarities in predicted overall WTD patterns, particularly for a recent drier period (1965-1995), there were clear periods when TA-based WTD predictions underestimated (i.e. drier during 1830-1930) and overestimated (i.e. wetter during 1760-1830) past WTD compared to MILLENNIA model predictions. Importantly, simulated grouse moor management scenarios may explain the drier TA WTD predictions, resulting in considerable model predicted carbon losses and reduced methane emissions, mainly due to drainage. This study demonstrates the value of a site-specific and combined data-model validation step towards using TA-derived moisture

conditions to understand past climate-driven peatland development and carbon budgets alongside modelling likely management impacts.

Introduction

Globally, peatlands contain ~30% of all soil organic carbon (SOC), despite covering only 3% of the land surface (Parish *et al.*, 2008). They occur mainly in the Northern Hemisphere circumpolar region (henceforth NHCP), where low temperatures, high soil moisture and slow decay rates of litter input via net primary production (NPP) allow peat to form (i.e. a long-term positive balance between NPP and litter decay), often under conditions of high water-table depth (WTD). Crucially, this slow decay preserves an archive of peatland development (i.e. TA and plant remains) that can be dated and used to reconstruct past drivers of peatland growth, such as WTD and vegetation composition, providing key information on how peatlands respond to changes in climate.

Blanket bogs are a globally rare peatland habitat with the UK containing about 15% of this habitat (Tallis, 1998) but mostly in a degraded state (Natural England, 2008), largely due to past environment (e.g. N deposition) and management (e.g. drainage) impacts. In the UK blanket bogs represent about 90% of all peatlands (Bain *et al.*, 2011), which are often managed for grazing and grouse shooting, commonly supported by draining the peat and regular burning of vegetation. The consequence is dominance of heather (*Calluna vulgaris*), very low overall plant biodiversity, suppressed cover of peat-forming *Sphagnum* mosses (Lindsay, 2010), and often drier and eroding peat. In fact, only about 12% of protected blanket bog sites are classified as in favourable condition (Natural England, 2008). However,

1 little is known about management impacts on the long-term SOC accumulation and soil C
2 emissions.

3
4 UK blanket bogs have accumulated peat since the start of the Holocene about 11.6 k years
5 ago under varying climate, but current bioclimatic models highlight the threat by climate
6 change to their natural range (Gallego-Salla *et al.*, 2010), suggesting that they might start to
7 degrade as the climate warms (Gallego-Sala & Prentice, 2013) resulting in water-table
8 drawdown and increased decomposition. However, existing bioclimatic models do not
9 consider extremely important autogenic negative feedbacks within peatlands that may
10 actually act as a 'buffer' to climate change (Swindles *et al.*, 2012). Such feedbacks include
11 ecohydrological linkages between changes in WTD and shifts in vegetation communities
12 (with different rooting depth and litter quality and thus affecting SOC inputs across depth and
13 peat decomposition rates). A better process-level understanding of climate-peatland SOC
14 feedbacks is clearly needed (Davidson & Janssens, 2006) since the mineralisation of peatland
15 soil organic matter (SOM) has the potential to release vast amounts of previously locked-up
16 C into the atmosphere. Depending on the water-table level, C emissions from decomposition
17 are either as CO₂ (oxic acrotelm) or CH₄ (anoxic catotelm), the latter particularly responsible
18 for exacerbating climate change and affecting the overall greenhouse gas (GHG) emissions.
19 However, a key limiting issue in more accurate predictions of future climate is still the
20 inadequate model representation of climate – terrestrial carbon (C) cycle feedbacks,
21 particularly of peatland soil organic carbon (SOC; Limpens *et al.*, 2008). Several peatland
22 models of varying complexity and feedback mechanisms have been developed (Clymo, 1984;
23 Gignac *et al.*, 1991; Bauer *et al.*, 2003; Frohking *et al.*, 2010; Heinemeyer *et al.*, 2010; Baird
24 *et al.*, 2012), which have often been compared to measured C stocks. However, there is a lack

of validating the C cycle underpinning hydrological model predictions against past palaeo records.

Peat archives from peat cores are also important for testing peatland development models, which predict peat layer accumulation and their associated chemical (e.g. carbon content) and physical (e.g. bulk density) properties, enabling comparison between the two. Data-model comparisons have revealed uncertainties in peat accumulation processes (Clark *et al.*, 2010), particularly considering hydrology, vegetation and NPP. Further, recent peat core studies (Charman *et al.*, 2013) indicate increased C accumulation during warmer periods due to increased NPP outweighing higher decomposition, which contradicts most global earth system model carbon cycle simulations (e.g. Friedlingstein *et al.*, 2006). Moreover, those global earth system models used by the Intergovernmental Panel on Climate Change (IPCC) do not yet adequately include peatland SOC dynamics, limiting global predictions on future climate C-cycle feedbacks and resulting GHG emissions. Latest attempts to include peatlands in global dynamic vegetation models (e.g. LPX; Spahni *et al.*, 2013) are promising, yet still show major process-level uncertainties.

Testate amoebae (TA) are commonly used to reconstruct peatland hydrological changes over the Holocene, as indicator species are aligned across a gradient of wet to dry conditions based on present-day associations (Amesbury *et al.*, 2016; Turner *et al.*, 2014). Peat cores provide a stratigraphic (i.e. temporal) archive of past TA species composition, thus allowing to predict past moisture (and likely water-table) conditions from understanding of contemporary ecology. Quantitative transfer functions are used to establish a relationship between present-day species composition and hydrological data and then applied to subfossil data from cores (Amesbury *et al.*, 2016).

Model predictions of peat carbon stock and flux changes rely on capturing seasonal and inter-annual WTD changes. Whereas in the short-term site measurements can be used for model validations, validation over longer-time scales relies on comparing model predictions to WTD reconstructions, for example based on TA taxa found within the peat core. In addition to other dating tools such as radiocarbon (Turner *et al.*, 2014), the use of Spheroidal Carbonaceous Particles (SCPs; Swindles, 2010) allows the generation of temporally-constrained records of palaeo-hydrological conditions in the recent past. Together with dating tools, palaeo-reconstructions thus offer crucial insights into peatland development, (i) understanding baseline trajectories in peatland development, and (ii) assessing long-term resilience and recovery rates of peatlands to climate or management impacts (Swindles *et al.*, 2016). TA based reconstructions of past WTDs can then be compared to process model predictions, providing an important hydrological model validation tool. A good fit between TA and model predicted WTDs supports applying model scenarios to explore past management impacts on peatland functioning and carbon storage. The MILLENNIA peatland model predicts peat hydrological conditions (i.e. water-table depth) based on climatic conditions and long-term peat column growth either for annual or monthly time steps. Here we used the annual MILLENNIA version (Heinemeyer *et al.*, 2010) for long-term peat accumulation during the Holocene until 1914, and then either the annual or monthly version (Carroll *et al.*, 2015) until 2012. This choice of model application reflected the availability of climate data. The WTDs predicted by the MILLENNIA model allowed comparison to TA-based WTD reconstructions from ca. 1760 onwards for a peat core at Moor House National Nature Reserve (NNR) using TA transfer functions. The WTDs from different management model scenarios were then compared to the TA-based WTD, and related to C accumulation and C emissions affecting the GHG balance.

Materials and Methods

Site location and environmental conditions

The study site, Moor House National Nature Reserve (NNR), covers about 75 km² of a typical UK blanket bog with much known about its ecology (Garnett *et al.*, 2001; Heal & Perkins, 1978) together with detailed meteorological (i.e. weather station) and hydrological (i.e. water-table) data collected by the Environmental Change Network (ECN). It is located in the northern Pennines across an elevation range of 290 - 850 m (a.s.l.) and is characterised by a sub-arctic–oceanic climate with an average long-term mean annual temperature (MAT) and precipitation (MAP) at 550 m altitude of about 5.1°C (1931-1997) and around 2000 mm (last 20 years), respectively (cf. Garnett, 1998). Peat formation at the study site started about 9,000 years ago (see Heinemeyer *et al.*, 2010 for more site information). This study focuses on a square kilometre around the ECN meteorological station at 550 m (NY757328; 54° 68' N, 2° 37' W), a site with mostly deep (>1 m) peat (histosol), supporting dominant vegetation of *Calluna vulgaris* L. [Hull] and *Eriophorum* spp. with some *Sphagnum* spp., and classified as a *Calluna vulgaris*–*Eriophorum* blanket mire (for more information on vegetation and peat depth see Garnett *et al.*, 2000 and Heinemeyer *et al.*, 2010). Typically the peatland site has a pH of generally between 3.0 – 4.2 and a high mean annual WTD of within 5 cm of the peat surface for >80% during the year (Evans *et al.*, 1999).

Peatland model predictions

The MILLENNIA peatland model (Heinemeyer *et al.*, 2010) predicts peat hydrological conditions (i.e. water-table) based on climatic conditions either for annual or monthly time steps. The hydrological conditions, in connection with internal factors (e.g. litter quality) and external modifiers (e.g. temperature and oxygen availability), then determine decomposition

1 rates of old and new carbon fractions (as in Bauer, 2004) across the peat profile based on
2 litter inputs via NPP as a function of potential evapotranspiration (PET) based on Lieth &
3 Box (1972). Detailed model explanations are provided in Heinemeyer *et al.* (2010; annual
4 model) and Carroll *et al.* (2015; monthly model). Here we used the annual version for the
5 long-term peat accumulation during the Holocene and for the period until 1914, and either the
6 annual or monthly version until 2012 (reflecting model application and climate data
7 availability). However, model outputs are only provided as annual averages in relation to
8 average TA predictions. The main model driver is climate with either monthly or annual
9 temperature and rainfall amounts as inputs. Both versions consider topography (e.g. slope
10 affecting temperature, runoff and erosion), vegetation (e.g. affecting NPP, litter quality and
11 transpiration losses) and proceeding hydrological conditions (e.g. high water-tables leading to
12 higher runoff) to predict a new water-table, vegetation (based on the preceding five-year
13 average water-table) and corresponding changes in soil carbon fluxes (i.e. CO₂ and CH₄ from
14 decomposition), peat depth increments (i.e. accumulation) and soil C budgets (change in soil
15 C stock in relation to input from NPP and losses from soil C fluxes and erosion).

16
17 Further changes were implemented to improve hydrological and C flux process
18 representation by calculating water filled pore space in the peat, bedrock drainage, plant-
19 mediated transport and methane oxidation (oxidation):

20
21 *The available pore space* in relation to the height above the water-table depth (WTD)
22 was based on data by Hayward & Clymo (1982; see Fig. 4, but ignoring the short
23 term hysteresis effect); an exponential relationship is assumed between the distance
24 to the water-table and the available pore space ($0.2 \cdot \text{EXP}(1.6 \cdot \text{WTD}^2)$), such that
25 available space increases with distance from the water-table. Total space is then

calculated by integrating the available pore space over the available unsaturated peat cohorts. By combining the water entering the system with the available space, a new WTD is calculated.

To simulate drainage of the peat column into the bedrock, two drainage factors are included, specific yield (SY) (0.1) and hydraulic conductivity (HC). These are set to default values of 0.02 (SY in %) and 0.1 (HC in cm/year) reflecting average values for clay reported by Johnson (1967) for SY (2%) and for unweathered clay based on Bear (1972) for HC (10-5 feet/day). However, SY and HC can be altered as a user input.

The plant functional type (vegetation) composition of shrub, sedge, rush, grass, herb, *Sphagnum*, other moss) is based on a moving average of five years of previous water-tables, allowing representing a more stable/resilient vegetation in the case of only a few very dry or wet years.

The anoxic ratio is set to 0.035, similar to values reported in previous literature (Bauer, 2004) ranging from 0.025 to 0.0625.

Methane oxidation is set to 0.05 g C/g/yr and reflects the range of the very scarce data available on methane oxidation in relation to dry peat and/or on a carbon (mass) basis, i.e. McDonald *et al.* (1996) provided incubation values at 20°C for UK peat of around 0.08 g C/g/yr, Watson *et al.* (1997) quoting 0.012 g C/g/yr (i.e. 0.001021 molC/g/yr equal to 1.021e-3 molC/g/yr), but Yrjala *et al.* (2010) quoted only 0.0009 g C/g/yr (0.2 µmol/gDW/d) and Whaalen & Reeburg (2010) measured around 0.002 g C/g/yr.

1
2 We used available reconstructed Holocene climate data (based on a combination of recent
3 instrumental data and a variety of existing multiproxy climate reconstructions, see Morris et
4 al., 2015) to model long-term peat accumulation, Met Office 5 km gridded data (Perry &
5 Hollis, 2015) for the recent past (1914-1991) and ECN data (ECN Data Centre:
6 <http://data.ecn.ac.uk>) for recent present periods (1992-2012). Met Office data were adjusted
7 for elevation for the Moor House site in order to achieve the same long-term average
8 temperature and rainfall amounts as the Moor House ECN data (see Carroll *et al.*, 2015).
9 Model predictions of monthly water-table depth could be compared to averaged ECN hourly
10 automated dipwell data (UK grid location: NY 75072 33425; missing data were gap-filled by
11 interpolation of manual data) for the period of 1999 - 2012. Water-table depth standard
12 deviation for a Moor House model evaluation (see supplementary data in Carroll *et al.*, 2015)
13 was predicted to within 0.32 cm (linear $R^2 = 0.57$).

14
15 Moor House was a formal shooting estate during 1842-1951 and grouse moor management
16 scenarios reflected available site information (Rob Rose, CEH; personal communication),
17 which indicated a 20 year burn rotation. The associated drainage was assumed to last from
18 1831, before intensification of grouse shooting (to enable enhanced heather growth and drier
19 access conditions for gamekeepers), until the late 1950s. Burning was anticipated to have
20 started in 1851 and to reduce NPP to 1% in the burnt year (and charcoal adding about 5% to
21 an inert carbon pool), subsequently recovering in a sigmoidal shape to 100% by either 5 or 10
22 years after burning (based on field observations by Heinemeyer et al., unpublished data).
23 Drainage was expected to reduce WTD on average by 5 cm, based on the field evidence of
24 Wilson et al. (2010). Reduced WTD were linked to enhanced decomposition and increased
25 the associated CO₂ but decreased CH₄ emissions similarly to modelled impacts of natural

WTD changes (Heinemeyer *et al.*, 2010). Drainage (grip) effectiveness was assumed to be at optimum for 25 years and declining to 60% over the subsequent 15 years (renewed once in 1871 and then maintained at optimum until 1905, reflecting intense grouse shooting), finally declining to 0% by 1955. Grazing pressure was anticipated to be insignificant above 450 m (i.e. no reduction in NPP at the modelled altitude of 550 m a.s.l.). Further model scenarios considered a no management option (no shoot).

Water-table reconstructions

A Russian peat core was taken from the top 50 cm of peat beside the Moor House ECN station (Lat. 54.695500; Lon. -2.387400) following De Vleeschouwer *et al.* (2010). The core was returned to the laboratory and stored at 4°C before analysis. The top 20 cm of the core was sampled at 0.5 cm resolution and testate amoebae were prepared using the standard method of Booth *et al.* (2010). We applied the Northern England and pan-European transfer functions to the subfossil TA data to reconstruct water-table depth (Turner *et al.*, 2013; Amesbury *et al.*, 2016). In both cases weighted-averaging tolerance downweighted regression with inverse deshrinking was used as it yielded the best performance statistics. The water-table reconstructions were standardised following Swindles *et al.* (2015). There is some variation in predicted water-tables between the two transfer functions which is caused by inter-model differences in WTD optima of key taxa. Some drier taxa (e.g. *Nebela militaris*, *Trigonopyxis arcuata* type) have drier optima in the Northern England model owing to the inclusion of some very dry peatlands affected by fire and drainage from this region. Sample-specific standard errors for the TA transfer function reconstructions were based on 1000 bootstrapping cycles (e.g. Amesbury *et al.*, 2016). The resulting WTD predictions could then

be paired with mean annual WTD predictions for years obtained from the MILLENNIA model based on establishing a peat age-depth profile based on a SCP profile.

Peat core age-depth profile

SCPs have been deemed to provide reliable age information from peatlands in Northern Britain and Ireland for the last ~150 years (Swindles, 2010; Swindles *et al.*, 2010; Turner *et al.*, 2014). Three unambiguous features can be used as age-equivalent stratigraphic markers: (i) the start of the record (c. 1850), (ii) the rapid increase in SCPs (c. 1950) and (iii) the peak (c. 1978). These represent (i) the start of high temperature combustion and power generation; (ii) the post-WW2 increase in power generation and fossil fuel burning and (iii) the peak of SCP production before reliance on other methods of power generation and the advent of clean fuel technologies (e.g. Rose *et al.*, 1995). We used the method of Swindles (2010) to prepare SCPs from the peat. Linear interpolation was used to generate a simple age-depth model between the three SCP age-equivalent stratigraphic markers and the date of core sampling (2011) as the uppermost time point; dates before the start of the SCPs are an extrapolation based on accumulation rate.

Results

Water table reconstructions

The testate amoebae data from Moor House (Fig. 1) along with the SCP concentrations and water-table reconstructions show periods of changes in wetness across the peat depth profile.




Fig. 1 Testate amoebae record (%) from the Moor House blanket bog, Northern Britain alongside the SCP concentration data with chronological assignments marked (i.e. 1978 at ~4.5 cm; 1950 at ~9.0 cm; 1850 at ~14.0 cm). The water-table depth reconstruction using the transfer function of Turner *et al.* (2014) is illustrated with standard errors derived from bootstrapping (1000 cycles). The standardised water-table reconstruction in relation to z-scores (following Swindles *et al.*, 2015) is also shown.

The earliest part of the record (before 1850) is very wet owing to the presence of wet-indicator species such as *Amphitrema wrightianum* and *Archerella flavum*. After this, the assemblage suggests quite a dry peatland surface with the abundance of *Nebela militaris*, *Cyclopyxis arcelloides* type (mostly comprised of *Phryganella acropodia*) and *Heleopera petricola*. All testate amoeba taxa identified in fossil samples were present in the modern training sets of the two transfer functions. The comparison of the Northern England and pan-European modelled water-tables (Fig. 2) illustrates that the WTD optima of TA species from the wider European area are wetter than those for Northern England (for the regression equation see Fig. 2 legend).




Fig. 2 Water-table depth (WTD) reconstructions for the Moor House blanket bog peat core based on alternative testate amoebae transfer functions: Europe, EU (grey) (Amesbury *et al.*, 2016) *versus* Northern England, NE (black) (Turner *et al.*, 2013). Note, a lower WTD value means wetter conditions. The linear regression for the reconstructed WTD comparison is $y_{[NE]} = 0.79x_{[EU]} + 7.58$; $R^2 = 0.65$.

TA-derived *versus* MILLENNIA model water-table predictions and site records

Water-table reconstructions for the Moor House core based on the Northern England transfer function (WTD_{TA_NE}) showed an overall good correlation with the MILLENNIA modelled WTD during the period of high quality climate records available for modelling (i.e. 1933-2004; see Fig. 3a) but TA-predictions were consistently drier ($WTD = 1.79 * WTD_{TA_NE} - 39.57$; $r^2 = 0.68$). A comparison with the EU TA transfer function (Fig. 3b) indicated slightly wetter conditions but with a less good fit to the measured data ($WTD = 0.714 * WTD_{TA_EU} - 10.613$; $R^2 = 0.47$). The reason behind this difference in TA-predicted WTD likely reflects

different WTD optima of key testate amoebae species in the different geographical areas. Therefore, we decided to use the WTD_{TA_NE} model for subsequent analysis because of the better directional fit in addition to closer geographical location of the TA records.




Fig. 3 Linear regression of MILLENNIA model predicted mean annual water-table depth (WTD in cm) for 19 years during 1933-2004 *versus* the corresponding years for the two testate amoebae (TA) reconstructions (see Fig. 2) based on the **A**) uncorrected transfer function derived for northern England (NE) as in Turner *et al.* (2013), and **B**) the uncorrected European (EU) transfer function (Amesbury *et al.*, 2016). The corresponding regression equations are shown.

The available ECN site climate data (1931-2010) for Moor House were used to make comparisons between annual mean WTD predicted by the MILLENNIA model to the paired years (± 1 year) with the offset corrected available TA-derived WTD_{TA_NE} from the Moor House core (Fig. 4). Not only did the WTD range predicted by the model agree with the TA-derived WTD, but it also reproduced the general pattern of dry *versus* wet years (MILLENNIA $WTD = WTD_{TA_NE} + 0.004$; $r^2 = 0.68$). Both WTD predictions indicated that the years between 1965 and 1995 were an extended period of fairly dry conditions compared

to the long-term WTD average of 4.4 ± 1.8 cm (based on annual averages of ECN dipwell data for 1995-2012).




Fig. 4 Selected years for mean annual water-table depth (WTD in cm) predictions (in the peat profile) between 1931-2004 from the MILLENNIA model (grey line) paired with years (triangles) of available testate amoebae (TA) based predictions (black line) using the offset regression (Fig. 3A). The dashed line indicates the long-term (1995-2012) mean annual WTD at Moor House (ECN data).

The annual MILLENNIA model prediction of WTD during 1931-2012 also showed very good agreement with the paired years of available offset corrected TA-derived WTD, capturing peaks and troughs as well as the average trend (Fig. 5). Moreover, the model predicted WTD (WTD_{Mod}) followed the available annual ECN site WTD (WTD_{ECN}) data very well ($WTD_{ECN} = WTD_{Mod} * 0.813 + 0.595$, $r^2 = 0.57$; see Carroll *et al.*, 2015).




Fig. 5 Annual water-table depth (WTD in cm) predictions (in the peat profile) since 1930 from the MILLENNIA model (dark grey lines, with 4-year average as dashed line) *versus* the 4-yearly predictions (black line) from testate amoebae (TA) based on the offset regression (Fig. 3A) and actual annual site WTD averages (white circles and light grey line) based on available continuous WTD data (1995-2012).

Comparison of predicted water-tables for management scenarios and site records

Use of the available long-term climate data for a nearby Northern England peatland site (Morris *et al.*, 2015), adjusted to the long-term mean temperature and total rainfall for Moor House, together with Moor House ECN climate data since 1931, provided a basis for a comparison of the WTD predicted by the MILLENNIA model over an extended time period (1750s till 2012), for which TA WTD reconstructions were available (Fig. 6). Although the general WTD patterns agreed very well between the two predictions, the (unadjusted) TA WTD predictions were much drier than the MILLENNIA predictions. Whilst MILLENNIA

model predictions indicated mean annual WTD conditions during 1760-1830 of between 2-6 cm, the unadjusted TA predictions of WTDs were around 15 to 25 cm. Adjusting for the 1931-2004 determined TA-WTD offset (Fig. 3a) increased the TA-based WTD predictions to above the peat surface (-2.92 ± 6.16 cm; data not shown). However, whereas MILLENNIA model predictions of the unmanaged (no shoot) scenario remained very wet during 1830-1940, unadjusted TA-predictions showed much lower WTD during that period, but overlapped again with the MILLENNIA model predictions from the 1940s onwards (Fig. 6).





Fig. 6 Water-table depth (WTD in cm) predictions from the MILLENNIA model (grey lines) versus 4-yearly testate amoebae (TA) based predictions (black line) and actual annual site WTD averages (white circles) based on available continuous site WTD data (1995-2012). Light dashed grey line indicates model scenario without any management, whereas dark solid grey line indicates a grouse moor management scenario during 1842-1951 (dashed arrow) with a period of active drainage (1831-1920). Note the different y-axis scales (i.e. unadjusted TA WTD predictions).

Predicted impacts on soil C emissions, peat C budgets and peat accumulation rates

The observed drier than average mean annual WTD predictions from the MILLENNIA model during 1965-1995 (Fig. 4) corresponded to $5.2 \text{ g C m}^{-2} \text{ yr}^{-1}$ lower net methane (CH_4) but $13.4 \text{ g C m}^{-2} \text{ yr}^{-1}$ higher CO_2 emissions (Fig. 7a) than during the remaining period of 1931-2012. This change in soil C fluxes to the atmosphere reflected the increased aerobic decomposition processes under a lower WTD, which resulted in near zero mean peat depth increment (reduced by 0.041 cm yr^{-1}) and a lower mean soil C budget (reduced by $6.2 \text{ g C m}^{-2} \text{ yr}^{-1}$) (Fig. 7b) during the drier 1965-1995 period.






Fig. 7 MILLENNIA model predictions (five year running means) of climatic impacts during 1931 to 2012 on A) annual water-table depth (WTD in cm; black line) and carbon (C) fluxes of CO₂ (dark grey line) and CH₄ (light grey line), and B) annual peat depth increments (grey line) and soil C budget (black line) with positive numbers representing C gains.

The MILLENNIA model scenario period of 1851-1950 (Fig. 8) showed a positive (C gain) annual soil C budget (Fig. 8b) of $12.7 \pm 22.7 \text{ g C m}^{-2}$ for the unmanaged scenario (with a mean WTD of $4.7 \pm 1.5 \text{ cm}$; Fig. 8a) compared to a negative (C loss) budget of $-38.6 \pm 110.1 \text{ g C m}^{-2}$ for the grouse managed scenario with drainage (mean WTD of $7.7 \pm 2.1 \text{ cm}$) assuming a 10-year NPP recovery time after burning (resulting in $51.3 \text{ g C m}^{-2} \text{ yr}^{-1}$ less soil carbon gain under grouse management). The corresponding peat depth increments predicted by the MILLENNIA model during 1851-1950 showed a mean peat accumulation of $0.03 \pm 0.59 \text{ cm}$ for the unmanaged (no shoot) *versus* a loss of $-0.06 \pm 0.61 \text{ cm}$ for the grouse managed scenario (i.e. 0.09 cm yr^{-1} less peat depth increment). Moreover, the 5-year NPP re-growth scenario (data not shown) resulted in about 45% lower mean annual soil C budget loss ($-21.6 \pm 93.7 \text{ g C m}^{-2}$) and half the peat increment loss ($-0.03 \pm 0.63 \text{ cm}$) compared to the

10-year NPP recovery scenario. In comparison to the default burn and drain scenario, the burn but no drain 10-year NPP re-growth scenario (data not shown) resulted in about 30% lower C and peat accumulation losses with a mean annual soil C budget of $-28.4 \pm 108.7 \text{ g C m}^{-2}$ and $-0.04 \pm 0.63 \text{ cm}$ mean annual peat depth increment (with a mean WTD of $4.8 \pm 1.9 \text{ cm}$). The burn but no drain 5-year NPP re-growth scenario reduced this further (i.e. 80% lower soil C and peat increment losses with a mean annual soil C budget of $-10.9 \pm 92.6 \text{ g C m}^{-2}$ and a mean annual peat depth increment of $-0.01 \pm 0.64 \text{ cm}$).

During 1831-1850, the period of drainage only (i.e. no burning), the managed scenario reduced the mean annual soil C budget by 27.8 g C m^{-2} to $-7.2 \pm 15.8 \text{ g C m}^{-2}$, which reflected an average reduction in mean annual WTD by 3.9 cm to 8.3 cm (Fig. 8a). These changes in soil C budget under drainage only corresponded to an annual peat depth increment reduction by 0.06 cm to $-0.02 \pm 0.63 \text{ cm}$ compared to the unmanaged scenario (Fig. 8b).

The grouse moor management (i.e. 1851-1950) not only impacted C dynamics via reduced water-tables from drainage (Fig. 8a), it also altered C inputs and thus C dynamics via reduced NPP following burning. Overall, drained and 10-year NPP recovery scenarios reduced both mean annual C losses from soil CO_2 fluxes ($308 \pm 106 \text{ g C}$) and annual CH_4 emissions ($4.9 \pm 8.3 \text{ g C}$) compared to the unmanaged scenario for which mean annual values were $417 \pm 60 \text{ g C}$ for CO_2 and $13.2 \pm 20.4 \text{ g C}$ for CH_4 net emissions (i.e. including methane oxidation, ebullition and plant-mediated transfer (PMT) processes via sedge leaves and stems). However, whereas the no drain 10-year NPP burn scenario decreased CO_2 ($292 \pm 110 \text{ g C}$) and increased CH_4 ($9.8 \pm 18.0 \text{ g C}$) emissions (Fig. 8a), the 5-year NPP scenario (data not shown) increased both CO_2 fluxes ($348 \pm 92 \text{ g C}$) and net CH_4 emissions ($11.2 \pm 18.8 \text{ g C}$) emissions, reflecting quicker vegetation re-growth and thus NPP and PMT recovery.





Fig. 8 MILLENNIA model predictions (five year running means) of management impacts during 1825 until 1945 on **A**) annual water-table depth (WTD; black line) and carbon (C) fluxes of CO₂ (dark grey line) and CH₄ (light grey line), and **B**) annual peat depth increments (grey line) and the soil C budget (black line) with positive numbers representing net C gains. Unbroken lines represent the unmanaged (no shoot) ‘natural state’, whereas dashed lines represent predictions for grouse moor management (1842-1951) including active drainage (1931-1920); drainage efficiency 25 years at maximum plus 15 years reduction to 60% and 0% by 1955) and burning (20 year cycles; with an exponential 10-year re-grow period to full net primary productivity) during 1850 till 1950.

Discussion

This study provided novel insights into ecological applications of using TA-derived water-table depth reconstructions in a site-specific model validation context. The findings highlight the value of combining palaeo-ecological records with process level modelling to allow better understanding of the effects of climate and management on peat development and C cycling. This combination shows promising potential in allowing to understand the contributions of past environmental (e.g. climate) and human-induced (e.g. grouse management) changes in peatland development over time, C stocks and corresponding C fluxes.

Although the EU transfer function predictions (as used in Amesbury *et al.*, 2016) showed wetter conditions overall (Fig. 2), the model fit to the MILLENNIA predictions was not improved, indicating a better overall fit using the geographically closer NE transfer function, possibly applying across the wider UK context, which has yet to be tested. Previous work comparing TA to recent short-term monitored WTD by Amesbury *et al.* (2016) was based around a collection of samples from several sites and did not compare TA-derived reconstructions to site-specific long-term modelled hydrological time series. Moreover, whereas Amesbury provided a WTD reconstruction based on comparing two different TA transfer functions, we identified and adjusted for a generic transfer function offset based on a site-specific validation period (Fig. 3) to allow a context-dependent (i.e. NE blanket bog at Moor House) model comparison over several centuries. The fairly constant offset could well reflect a TA sample bias of summer sampling and thus drier conditions overall (i.e. during contemporary sampling to obtain TA calibration data on moisture or WTD relationships); water-tables at Moor House frequently drop to 20 - 30 cm during July and August (Carroll *et al.*, 2015). With fairly constant water-table drawdown during drier periods (unpublished data

by Heinemeyer et al.), a general offset can be expected (i.e. wetter sites are always wetter and vice-versa).

Notwithstanding the overall good fit from the offset corrected TA-derived *versus* model predicted water-tables (Fig. 4), some resulting TA-derived water-table predictions such as the very wet period between 1730-1830 (as shown in Fig. 6) might seem questionable. However, the predicted standing water conditions of up to -10 cm when applying the offset adjusted TA functions (using the offset corrected NE transfer function as shown in Fig. 3a) fall within the tolerance and optima values (i.e. Turner et al., 2013) of two wet to aquatic TA indicator taxa (see Fig. 5 in Amesbury *et al.*, 2016), *Amphitrema wrightianum* type (2 ± 6 cm mean WTD) and *Archerella flavum* (8 ± 7 cm mean WTD). This coincides with documented wet bog conditions across Ireland (Swindles *et al.*, 2013) and Northern Britain (Mauquoy *et al.*, 2002) during this period (i.e. the Little Ice Age) with potentially standing water for most of the year or water pools in hollows. Such conditions of standing water are not specifically captured by either the model or TA WTD predictions, which are relatively insensitive to water-tables above the surface. Moreover, the climate data used in the model become less reliable before the 1850s, and this period might well have been wetter than the by Morris *et al.* (2015) reconstructed climate record suggests. Therefore, MILLENNIA predicted WTD during 1730-1830 might be under predicting the WTD in relation to uncertain climate input data.

Most interesting was the period between 1840s and 1940s, an unexpectedly drier period based on TA predicted hydrological conditions compared to the wetter conditions predicted by the MILLENNIA model based on climate only (Figure 6). Moor House was a formal shooting estate throughout exactly this period (i.e. 1842-1951), based on grouse bags and predator control information (ECN data provided by Rob Rose from the Centre for Ecology

and Hydrology (CEH) Lancaster, personal communication). Grazing intensity on true blanket bogs such as Moor House is low (0.5 sheep/ha according to Rawes & Heal, 1978), except perhaps for recently burnt areas; historically grazing was probably low overall because farmers were unable to grow sufficient feed to maintain their stock over the winter, particularly at altitudes above 500 m. Therefore, burning in connection with drainage seems to be the most likely factor explaining the water-table lowering. The available anecdotal evidence (R. Rose (CEH), personal communication) from some of the game keepers in the latter part of the period suggest that an average of 250 acres were burnt each year out of a total area of about 5000 acres (i.e. averaging a 20 year burn cycle) and drainage ditches were maintained to aid the heather management. Such intense grouse moor management would have most likely resulted in a lowering of the water-table by around 5 cm, as has been observed in a grip blocking study by Wilson *et al.* (2010). Interestingly, the MILLENNIA scenarios simulating such grouse moor management resulted in the modelled WTD capturing this drier period as inferred from the TA record, which in the model scenario was mainly a result of drainage.

Moreover, the managed model scenario indicated considerable losses in the soil C budget and decreased peat accumulation rates. Up-scaled to an intense grouse moor management period of 100 years these accumulated C losses (i.e. 1850-1950) relate to around 5 kg C m⁻² less soil carbon or 10 cm lower peat depth accumulation than predicted for the unmanaged scenario. Notably, burning and drainage contributed equally to the carbon loss, via reduced NPP (i.e. less litter input) and enhancing decomposition (i.e. lower WTD). Garnett *et al.* (2001) indicated a carbon loss and lower peat accumulation on burn management at Moor House (based on experimental burn plots) of around 73 g C m⁻² yr⁻¹, slightly higher but very similar to our model predictions of 51 g C m⁻² yr⁻¹. However, the Garnett *et al.* study contained some

1 methodological uncertainties in relation to calculating C stocks (Clay *et al.*, 2010) and the
2 MILLENNIA model did not account for any potential charcoal impacts on hydrology and
3 decomposition. In particular, impacts on peat bulk density (i.e. charcoal pieces entering peat
4 pores) with possibly changes in water holding capacity and charcoal inputs representing an
5 inert C pool (i.e. biomass partly by-passing decomposition; see Clay *et al.*, 2010) with
6 possible additional effects on decomposition rates (e.g. negative priming; see Lu *et al.*, 2014)
7 might need to be considered in future model developments. Notably, the Garnett *et al.* (2001)
8 study has been highlighted by Evans *et al.* (2014) as the only substantial study assessing
9 burning impacts on UK blanket bogs. Therefore, burn management implications on the long-
10 term peat C stock remain uncertain.

11
12 Whilst the estimated mean annual soil CO₂ fluxes of around 300-400 g C m⁻² yr⁻¹ (i.e. range
13 of managed vs. unmanaged scenario, respectively) are within the same order of magnitude
14 reported for a study at Moor House (Clay *et al.* 2010), the CH₄ emissions of around 5-10 g C
15 m⁻² yr⁻¹ were slightly higher compared to published values for the Flow Country (4.3 g C m⁻²
16 yr⁻¹; Levey & Gray, 2015) and Moor House (3.9 g C m⁻² yr⁻¹; Worrall *et al.*, 2007, and 6.4 g
17 C m⁻² yr⁻¹; Worrall *et al.*, 2009). However, the existing data on net CH₄ emissions from
18 peatlands are very uncertain, with many older studies using inadequate methodologies (i.e.
19 long chamber incubation periods and inaccurate analyser techniques such as gas
20 chromatographs). Moreover, recent data from Moor House during two wet and warm years
21 (2015-2016) measured much higher than previous average annual CH₄ emissions from peat
22 decomposition of 2 to 400 g C m⁻² yr⁻¹ (unpublished; pers. comm. Rob Rose at CEH).
23 However, the largest uncertainty for field measurements is likely related to manual chamber
24 measurements often not capturing plant-mediated transfer rates and ebullition, bypassing
25 methane oxidation and thus leading to an overall underestimation of 'true' methane fluxes.

Due to both model and measurement uncertainties, the presented CH₄ emissions should thus not be taken as real flux predictions but rather be seen as indicators of possible differences to be assessed by future monitoring.

Peat cores provide a valuable archive for reconstructing peat development in relation to past climate, particularly TA-based water-table reconstructions as a driver of peat and carbon accumulation. We have shown here that combining this peatland archive with actual site measurements and model scenarios of past water-tables can provide additional information on likely impacts of land management, which are not easily detectable otherwise, but are crucial for explaining observed peat accumulation not explained by climate alone.

Acknowledgements

The Environmental Change Network kindly provided access to ECN data (Environmental Change Network 2012. Dataset accessed 25/06/2013 at <http://data.ecn.ac.uk> - data citation code: 'ECN:AH2/12 & ECN: AH2/14'). Paul Morris at the University of Leeds (UK) kindly provided updated Holocene climate data. The MILLENNIA model was developed with funding from NERC from the CTCD and NCEO (NERC F14/G6/105) centres, UKPopNet (NERC R8-H12-01) and Defra (BD5104); we thank Simon Croft for help with previous model developments. We thank Gabriel Mason-Dixon for assistance in the laboratory and Manuel Gloor for field assistance. There are no conflicts of interests.

References

Amesbury, MJ, Swindles, GT, Bobrov, A, *et al.* (2016) Development of a new pan-European *testate amoeba* transfer function for reconstructing peatland palaeohydrology. Quaternary Science Reviews, **152**, 132–151.

Bain C, Bonn A, Stoneman R. *et al.* 2011. IUCN UK Commission of Inquiry on Peatlands. IUCN UK Peatland Programme, Edinburgh.

Baird AJ, Morris PJ, Belyea LR (2012) The DigiBog peatland development model 1: rationale, conceptual model, and hydrological basis. Ecohydrology, **5**, 242–255. doi:10.1002/eco.230.

Bauer IE, Gignac LD, Vitt DH (2003) Development of a peatland complex in boreal western Canada: lateral site expansion and local variability in vegetation succession and long-term peat accumulation. Canadian Journal of Botany, **81**, 833-847.

Bauer IE (2004) Modelling effects of litter quality and environment on peat accumulation over different time-scales. Journal of Ecology, **92**, 661-674.

Bear J (1972) Dynamics of Fluids in Porous Media, first ed. American Elsevier, New York.

Booth RK, Lamentowicz M, Charman DJ (2010) Preparation and analysis of testate amoebae in peatland paleoenvironmental studies. Mires and Peat, **7**(2), 1-7.

1 Clark JM, Gallego-Sala AV, Allott TEH *et al.* (2010) Assessing the vulnerability of
2 blanket peat to climate change using an ensemble of statistical bioclimatic envelope
3 models. *Climate Research*, **45**, 131–150.

4
5 Carroll M, Heinemeyer A, Pearce-Higgins JW *et al.* (2015) Hydrologically-driven ecosystem
6 processes determine the distribution and persistence of ecosystem-specialist predators under
7 climate change. *Nature Communication*, **7851**, doi:10.1038/ncomms8851.

8
9 Charman DJ, Beilman DW, Blaauw *et al.* (2013) Climate-related changes in peatland carbon
10 accumulation during the last millennium. *Biogeosciences*, **10**(2), 929-944; DOI: 10.5194/bg-
11 10-929-2013.

12
13 Clay GD, Worrall F, Rose R (2010) Carbon budgets of an upland blanket bog managed by
14 prescribed fire. *Journal of Geophysical Research-Biogeosciences*, **115**, G04037.

15
16 Clymo RS (1984) The Limits to Peat Bog Growth. *Philosophical Transactions Royal Soc*
17 *London, B*, **303**, 605-654.

18
19 Davidson EA, Janssens IA (2006) Temperature sensitivity of soil carbon decomposition and
20 feedbacks to climate change. *Nature*, **440**, 165–173.

21
22 De Vleeschouwer F, Chambers FM, Swindles GT (2010) Coring and sub-sampling of
23 peatlands for palaeoenvironmental research. *Mires and Peat*, **7**, 1-10.

1 Evans MG, Burt TP, Holden J, Adamson JK (1999) Runoff generation and water table
2 fluctuations in blanket peat: evidence from UK data spanning the dry summer of 1995.
3 Journal of Hydrology, **221**, 141-160.

4
5 Evans CD, Bonn A, Holden, J, *et al.* (2014) Relationships between anthropogenic pressures
6 and ecosystem functions in UK blanket bogs: Linking process understanding to ecosystem
7 service valuation. Ecosystem Services, **9**, 5–19.

8
9 Friedlingstein, P., Cox, P., Betts, *et al.* (2006) Climate-carbon cycle feedback analysis: results
10 from the C⁴MIP model intercomparison. Journal of Climate, **19**, 3337–3353.

11
12 Frohking S, Roulet NT, Tuittila E, Bubier JL, Quillet A, Talbot J, Richard PJH (2010) A new
13 model of Holocene peatland net primary production, decomposition, water balance, and peat
14 accumulation. Earth System Dynamics **1**(1): 1-21.

15
16 Gallego-Sala AV, Prentice IC (2013) Blanket peat biome endangered by climate change.
17 Nature Climate Change, **3**, 152–155.

18
19 Gallego-Sala AV, Clark JM, House JJ, *et al.* (2010) Bioclimatic envelope model of climate
20 change impacts on blanket peatland distribution in Great Britain. Climate Research, **45**(1),
21 151-162.

22
23 Garnett (1998) Carbon storage in Pennine moorland and response to change. PhD
24 dissertation, Department of Geography, University of Newcastle-Upon-Tyne.

- 1 Garnett MH, Ineson P, Stevenson AC (2000) Effects of burning and grazing on carbon
2 sequestration in a Pennine blanket bog, UK. *The Holocene*, **10**, 729–736.
- 3
- 4 Garnett MH, Ineson P, Stevenson AC, Howard DC (2001) Terrestrial organic carbon storage
5 in a British moorland. *Global Change Biology*, **7**, 375–388.
- 6
- 7 Gignac LD, Vitt DH, Bayley SE (1991) Bryophyte response surfaces along ecological and
8 climatic gradients. *Vegetatio*, **93**, 29-45.
- 9
- 10 Hayward PM, Clymo RS (1982) Profiles of Water Content and Pore Size in *Sphagnum* and
11 Peat, and their Relation to Peat Bog Ecology. *Proceedings of the Royal Society of London.*
12 *Series B, Biological Sciences*, **215**(1200), 299-325.
- 13
- 14 Heal OW, Perkins DF (1978) *Ecological Studies 27, Production Ecology of British Moors*
15 *and Mountain Grasslands*, Heal OW, Perkins DF (eds.).
- 16
- 17 Heinemeyer A, Croft S, Garnett MH, Gloor M, Holden J, Lomas MR & Ineson P (2010) The
18 MILLENNIA peat cohort model: predicting past, present and future soil carbon budgets and
19 fluxes under changing climates in peatlands. *Climate Research*, **45**, 207-226.
- 20
- 21 Johnson AI (1967) Specific yield: compilation of specific yields for various materials, *Water*
22 *Supply Paper*, **1662**-D.
- 23
- 24 Levey PE, Gray A (2015) Greenhouse gas balance of a semi-natural peatbog in northern
25 Scotland. *Environmental Research Letters*, **10**(9), 094019.

Lieth H, Box E (1972) Evapotranspiration and primary productivity: Thornthwaite Memorial Model. Publications in Climatology, **25**, 37-46.

Limpens J, Berendse F, Blodau C, *et al.* (2008) Peatlands and the carbon cycle: from local processes to global implications ? A synthesis. Biogeosciences Discussions, **5**, 1379–1419.

Lindsay R (2010) Peatbogs and carbon: a critical synthesis to inform policy development in oceanic peat bog conservation and restoration in the context of climate change. Environmental Research Group, University of East London, London.

Lu W, Ding W, Zhang J, Li Y, Luo J, Bolan N, Xie Z (2014) Biochar suppressed the decomposition of organic carbon on a cultivated sandy loam soil: a negative priming effect. Soil Biology and Biochemistry, **76**, 12-21.

Mauquoy D, van Geel B, Blaauw M, van der Plicht J (2002) Evidence from northwest European bogs shows ‘Little Ice Age’ climatic changes driven by variations in solar activity. The Holocene, **12**(1), 1–6; doi.org/10.1191/0959683602hl514rr.

McDonald IR, Hall GH, Pickup RW, Murrell JC (1996) Methane oxidation potential and preliminary analysis of metanotrophs in blanket bog peat using molecular ecology techniques. FEMS Microbiology Ecology, **21**, 197-211.

Morris PJ, Baird AJ, Young DM, Swindles GT (2015) Untangling climate signals from autogenic changes in long-term peatland development. *Geophysical Research Letters*, **42**, 10788–10797.

Natural England (2008) State of the Natural Environment; 3.8 Wetlands. Cat. Code: NE85. Available at: http://www.naturalengland.org.uk/Images/sone-section3.8_tcm6-4742.pdf [Accessed July 2017].

Parish F, Sirin A, Charman D, Joosten H, Minaeva T, Silvius M (eds.) (2008) Assessment on peatlands, biodiversity and climate change. Kuala Lumpur, Global Environment Centre and Wageningen, Wetlands International. 179 pp.

Perry M, Hollis D (2015) The generation of monthly gridded datasets for a range of climatic variables over the UK. *International Journal of Climatology*, **25**, 1041-1054.

Rawes M, Heal OW (1978) The Blanket bog as part of a Pennine moorland. In *Ecological Studies 27, Production Ecology of British Moors and Mountain Grasslands*, Heal OW, Perkins DF (eds).

Rose NL, Harlock S, Appleby PG, Battarbee RW (1995) The dating of recent lake sediments in the United Kingdom and Ireland using spheroidal carbonaceous particle concentration profiles. *The Holocene*, **5**, 328-335.

Spahni R, Joos F, Stocker BD, Steinacher M, Yu ZC (2013) Transient simulations of the carbon and nitrogen dynamics in northern peatlands: from the Last Glacial Maximum to the 21st century, *Climate of the Past*, **9**, 1287-1308.

Swindles GT (2010) Dating recent peat profiles using spheroidal carbonaceous particles (SCPs). *Mires and Peat*, **7**(3), 1-5.

Swindles GT, Blundell A, Roe HM, Hall VA (2010) A 4500-year proxy climate record from peatlands in the North of Ireland: the identification of widespread summer ‘drought phases’? *Quaternary Science Reviews*, **29**, 1577-1589.

Swindles GT, Morris PJ, Baird AJ, Blaauw M, Plunkett G (2012) Ecohydrological feedbacks confound peat-based climate reconstructions. *Geophysical Research Letters*, **39**, L11401.

Swindles GT, Lawson IT, Matthews IP, *et al.* (2013) Centennial-scale climate change in Ireland during the Holocene. *Earth Science Reviews*, **126**, 300-320.

Swindles GT, Holden J, Raby C, Turner TE, Blundell A, Charman DJ, Menberu MW, Kløve B (2015) Testing peatland water-table depth transfer functions using high-resolution hydrological monitoring data. *Quaternary Science Reviews*, **120**, 107-117.

Swindles GT, Morris PJ, Wheeler J, Smith M, Bacon KL, Turner TE, Headley A, Galloway JM (2016) Resilience of peatland ecosystem services over millennial timescales: evidence from a degraded British bog. *Journal of Ecology*, **104**, 621-636.

1 Tallis JH (1998) Growth and Degradation of British and Irish Blanket Mires. Environmental
2 Reviews, **6**, 81-122.

3
4 Turner TE, Swindles GT, Charman DJ, Blundell A (2013) Comparing regional and supra-
5 regional transfer functions for palaeohydrological reconstruction from Holocene peatlands.
6 Palaeogeography Palaeoclimatology, **369**, 395-408.

7
8 Turner TE, Swindles GT, Roucoux K (2014) Late Holocene ecohydrological and carbon
9 dynamics of a UK raised bog: human activity or climate change? Quaternary Science
10 Reviews, **84**, 65-85.

11
12 Watson A, Stephen KD, Nedwell DB, Arah JRM (1997) Oxidation of methane in peat:
13 kinetics of CH₄ and O₂ removal and the role of plant roots. Soil Biology and Biochemistry,
14 **29**, 1257-1267.

15
16 Whalen SC, Reeburgh WS (2000) Methane Oxidation, Production, and Emission at
17 Contrasting Sites in a Boreal Bog, Geomicrobiology Journal, **17**(3), 237-251.

18
19 Wilson L, Wilson J, Holden J, Johnstone I, Armstrong A, Morris M (2010) Recovery of
20 water tables in Welsh blanket bog after drain blocking: discharge rates, time scales and the
21 influence of local conditions. Journal of Hydrology, **391**, 377–386.

22
23 Worrall F, Armstrong A, Adamson JK (2007) The effects of burning and sheep-grazing on
24 water table depth and soil water quality in an upland peat. Journal of Hydrology, **339**, 1–14.

1 Worrall F, Burt TP, Rowson JG, Warburton J, Adamson JK (2009) The multi-annual carbon
2 budget of a peat-covered catchment. *Science of The Total Environment*, **407**, 4084–4094.

3

4 Yrjälä KIM, Tuomivirta T, Juottonen H, *et al.* (2011) CH₄ production and oxidation
5 processes in a boreal fen ecosystem after long-term water table drawdown. *Global Change*
6 *Biology*, **17**(3), 1311-1320.

7

8