



This is a repository copy of *Linear and Nonlinear Estimated GFR Slopes in ADPKD Patients Reaching ESRD*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/130268/>

Version: Supplemental Material

---

**Article:**

Neagu, M., Coca, D. [orcid.org/0000-0003-2878-2422](https://orcid.org/0000-0003-2878-2422) and Ong, A.C.M. (2018) Linear and Nonlinear Estimated GFR Slopes in ADPKD Patients Reaching ESRD. *American Journal of Kidney Diseases*, 71 (6). pp. 912-913. ISSN 0272-6386

<https://doi.org/10.1053/j.ajkd.2018.01.052>

---

**Reuse**







This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>



### **Figure legend**

a. Example of a linear model; b. Example of non-linear decline (two phase); c. Example of non-linear decline (negative exponential); d. Example of non-linear decline (prolonged stable phase); e. Patients with initial eGFR  $>100\text{ml}/\text{min}/1.73\text{m}^2$  (n=12); f. Patients with initial eGFR  $<100\text{ml}/\text{min}/1.73\text{m}^2$  (n=25).