UNIVERSITY of York

This is a repository copy of *Physiological and behavioral risk factors of type 2 diabetes mellitus in rural India*.

White Rose Research Online URL for this paper: <u>https://eprints.whiterose.ac.uk/130448/</u>

Version: Published Version

Article:

Barik, Anamitra, Mazumdar, Sumit, Chowdhury, Abhijit et al. (1 more author) (2016) Physiological and behavioral risk factors of type 2 diabetes mellitus in rural India. BMJ open diabetes research & care. e000255. ISSN 2052-4897

https://doi.org/10.1136/bmjdrc-2016-000255

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk https://eprints.whiterose.ac.uk/

BMJ Open Diabetes Research & Care

Physiological and behavioral risk factors of type 2 diabetes mellitus in rural India

Anamitra Barik,^{1,2} Sumit Mazumdar,³ Abhijit Chowdhury,^{1,4} Rajesh Kumar Rai¹

To cite: Barik A,

Mazumdar S, Chowdhury A, et al. Physiological and behavioral risk factors of type 2 diabetes mellitus in rural India. *BMJ Open Diabetes Research and Care* 2016;**4**: e000255. doi:10.1136/ bmidrc-2016-000255

Received 21 April 2016 Revised 15 June 2016 Accepted 5 July 2016



¹Society for Health and Demographic Surveillance, Suri, West Bengal, India ²Niramay TB Sanatorium and Chest Clinic, District Hospital, Suri, Birbhum, West Bengal, India

³Institute of Public Health Kalyani, Society for Health and Demographic Surveillance, Netaji Subhas Sanatorium (T.B. Hospital), Kalyani, West Bengal, India ⁴Department of Hepatology, School of Digestive and Liver Diseases, Institute of Post Graduate Medical Education and Research (IPGMER), Kolkata, West Bengal, India

Correspondence to

Rajesh Kumar Rai; rajesh.iips28@gmail.com

ABSTRACT

Background: The dynamics of physiological and behavioral risk factors of diabetes in rural India is poorly understood. Using data from a health and demographic surveillance site of Birbhum district in West Bengal, India, this study aims to assess the risk factors associated with type 2 diabetes mellitus. **Research design and methods:** A total of 7674 individuals aged \geq 18 years participated in a crosssectional study. Venous plasma glucose method was used for measuring and reporting glucose concentrations in blood, categorized as individuals with diabetes, pre-diabetes or impaired, and normoglycemic. Aside from a set of physiological and behavioral risk factors, a range of socioeconomic confounders of diabetes was computed. Bivariate

analysis with χ^2 test, and multivariate ordered logit regression methods were deployed to attain the study's objective.

Results: Overall 2.95% and 3.34% of study participants were diagnosed as individuals with diabetes and pre-diabetes or impaired, respectively. Compared to the poorest, the richest have higher probability (β : 0.730; 95% CI 0.378 to 1.083) of being diagnosed with diabetes. As compared to people with normal body mass index, overweight/obese people are more prone to being diagnosed with diabetes (β : 0.388; 95% CI 0.147 to 0.628). With a decreasing level of physical activity, people are more likely to be diagnosed with diabetes.

Conclusions: To curb the level of diabetes, this study recommends a culturally sensitive, focused intervention for the adoption of physical activity with more traditional dietary practices, to control the level of overweight/obesity. Attention should be paid to relatively older patients with diabetes or adults with pre-diabetes.

INTRODUCTION

As a signatory to the sustainable development goals (SDGs), India has committed to extend every possible effort to reduce one-third of premature mortality from non-communicable diseases (SDG: 3.4) by 2030.¹ In 2012, noncommunicable diseases (NCDs), also known as chronic diseases, killed over 38 million people in the world, and 1.5 million of all deaths were attributed to diabetes.² In 2011,

Key messages

- Being overweight or obese and low physical activity are the significant physiological and behavioral risk factors for diabetes in rural India.
- A focused lifestyle modification intervention to encourage physical activity and traditional dietary intake should be encouraged in rural India where the effect of tobacco and alcohol use did not appear to have any significant effect on diabetes.
- In future, study sites such as health and demographic surveillance system-Birbhum Population Project should aim to collect more information on diabetes, and its related comorbidity and mortality patterns among the sampled cohort population.

under the leadership of the WHO, over 190 countries endorsed a global mechanism to reduce the burden of diabetes called a 'Global Action Plan for the Prevention and Control of NCDs 2013-2020'.³ Globally, an estimated 422 million adults were living with diabetes in 2014, compared to 108 million in 1980.² Diabetes accounted for 12% of health expenditures in 2010, or at least \$376 billion -a figure expected to hit \$490 billion by 2030.⁴ With over 1.25 billion people, India has a little lower prevalence of diabetes (among adult population) than the global level (India: 7.8% vs worldwide: 8.5%).² Diabetes is growing alarmingly in India, with more than 65 million people having the disease, compared to 51 million in 2010.

Type 2 diabetes mellitus (formerly called non-insulin-dependent or adult-onset) which results from the body's ineffective use of insulin comprises 90% of diabetes cases.² Risk factors of diabetes can be categorized as 'metabolic or physiological risk factors' overweight or obesity, and hyperlipidemia (high levels of fat in the blood); and 'modifiable behavioral risk factors' –unhealthy diet, physical inactivity and the harmful use of tobacco and alcohol.^{2 6–10} In addition, socioeconomic factors act as potential confounders of diabetes, driven by forces that

BMJ

Epidemiology/health services research

include ageing, rapid unplanned urbanization and the globalization of unhealthy lifestyles.^{2 6 7}

Despite having a lower body mass index (BMI) on average, some Asian countries have similar or even higher prevalence of diabetes than Western countries,¹¹ and the risk of type 2 diabetes starts at a lower BMI for Asians than for Europeans.¹² In India, traditional dietary patterns are being lost as the population adapts to more industrialized and urban food environment.⁶ These changes enhance type 2 diabetes risks by increasing body weight and central adiposity.⁶ As lower income Asian countries shift away from agricultural labor toward a service sector economy, energy expenditure has declined dramatically, even in rural India. A metaanalysis revealed that current smokers had a 45% increased risk of developing diabetes compared with non-smokers.¹³ The accumulation of visceral adipose tissue among smokers may be due to increased plasma cortisol levels induced by the stimulation of sympathetic nervous system activity. Smoking decreases testosterone level leading to adiposity. High consumption of alcohol leads to excess caloric intake and obesity, which in turn accelerates propensity for diabetes.¹⁴

Although modification of physiological and behavioral risk factors are known to be essential for preventing and managing the type 2 diabetes, there is currently insufficient information available on type 2 diabetes and associated factors in Asian countries, especially for rural India. Using a cross-sectional data of 7674 individuals drawn from a health and demographic surveillance system (HDSS) site in rural India, this study assesses the physiological and behavioral risk factors associated with type 2 diabetes mellitus. It is hoped that this study will help understand strategies to halt the level of diabetes in a rural Indian context.

METHODS Data set

Data for the present study were drawn from the Birbhum Population Project (BIRPOP), a Health and Demographic Surveillance System (HDSS) site located in Birbhum district of West Bengal, a state located along the eastern boundary of India.¹⁵ The HDSS-BIRPOP site covers 4 of 19 blocks of Birbhum, namely Mohammad Bazar, Rajnagar, Sainthia, and Suri I. At its inception in 2008, the 2001 National Census was used as the sampling frame to select the study population. By adopting a multistage sampling design, 12 300 self-weighted households were sampled taking into account a 10% dropout and non-participation rate. The HDSS-BIRPOP collects extensive data on demographic processes, population health and epidemiology, and healthcare utilization, with a focus on anthropometric indices of cardiometabolic health, and cardiometabolic imaging data in a subgroup of individuals. More about the sampling procedure and the profile of HDSS-BIRPOP is available elsewhere.¹⁵

The data used in the study were collected between July 2012 and March 2013. It covered 12 557 households comprising a total of 54 585 rural Indians. Blood samples were collected from a subsample of 9886 individuals drawn from 36 176 adults (over \geq 18 years). At completion, 7674 individuals with blood tests were found eligible to be included in this study. The study flow chart is shown in figure 1. The participants have no clinical evidence of major cardiovascular disorders, which was defined as having a history of myocardial infarction, angina pectoris, prior revascularization, heart failure, atrial fibrillation, stroke, or peripheral arterial disease and end organ damage at enrolment. In addition, none of the participants were found to be suffering from type 1 diabetes mellitus.

The survey was conducted by 44 trained interviewers who had at least an undergraduate degree and some exposure to large scale sample surveys. All the interviewers (designated as the surveyors in HDSS-BIRPOP) were native speakers of at least one of the local languages—Bengali and Santhali (a tribal language). The follow-up protocols include study visits to each household every 3 weeks for assessment of follow-up for clinical events. Despite (small) effects of migration, there was very little loss in the follow-up within this cohort (1% per year maximally). Ethical clearance for collection and analysis of the blood samples was obtained through a formal review by the HDSS' institutional review board and signed informed consent from the participating individuals.¹⁵

Defining type 2 diabetes mellitus

To define type 2 diabetes mellitus, this study has adopted the guideline developed by the WHO and the definition of the International Diabetes Federation.¹⁶



Figure 1 Selection of participants included in this study.

People were diagnosed with diabetes if they recorded fasting plasma glucose (FPG) $\geq 126 \text{ mg/dL}$ or if they were taking antidiabetic drugs (oral hypoglycemic or insulin). The FPG cut-point for impaired fasting glucose ranged from 110 to 125 mg/dL; and individuals were categorized as normoglycemic (defined if individual has normal glucose level with low risk of developing diabetes or cardiovascular disease) if FPG was <110 mg/dL.¹⁶

Venous plasma glucose is the standard method for measuring and reporting glucose concentrations in blood. The study participants were instructed to fast for at least 12 hours prior to blood collection. Fasting blood samples were obtained by a trained phlebotomist in K2-EDTA tubes (3.0 mL); fluoride tubes (2.0 mL) and serum tubes (7.0 mL). Serum was prepared within 2 hours of blood collection. Plasma glucose was estimated by using the hexokinase/glucose-6-phosphate dehydrogenase method within 2 hours of blood collection.

Covariates

Guided by the existing literature, this study uses a set of potential physiological and behavioral risk factors of dia-The variables representing physiological betes. characteristics are BMI categorized as normal, underweight and overweight/obese; and physical activity level, categorized as high, medium, and low. BMI is defined as the ratio of the weight in kilograms to the square of the height in meters (kg/m^2) . A BMI of $<18.5 kg/m^2$ is widely used as a practical measure of being underweight, indicating chronic energy deficiency, whereas BMI ranges of 18.6–22.9 kg/m² and \geq 23 kg/m² are considered indicative of normal, and overweight/obesity, respectively.¹⁷ The level of physical activity was measured by asking how often (never, irregular/once or twice a day, or daily) an individual undertook certain types of physical activity on a typical day. The details of computing physical activity levels are discussed in appendix 1. Information obtained on tobacco use (both smoking and chewing), and alcohol consumption was used as the behavioral risk factors of diabetes. Respondents were investigated if they were currently (preceding 30 days of survey date) using tobacco or consuming alcohol, or if they have any history of tobacco use or alcohol consumption.

In addition, a list of potential confounders of diabetes was adapted to measure the risk factors of diabetes. The variables include completed age (categorized as 18 to 24, 25 to 34, 35 to 44, and \geq 45 years), sex (categorized as male or female), level of education (categorized as illiterate, grades 1 to 5, 6 to 10, and \geq 11), religion (Hindu, Islam or Others), social group (categorized as scheduled caste, scheduled tribe, other backward class, and others), occupation (categorized as unemployed, non-earning household chore, labor, and service/ pensioner/ retired), and wealth quintile (categorized as poorest, poorer, middle, richer, and richest). Scheduled Castes and Scheduled Tribes are identified by the government of India as socially and economically backward social groups and considered to be in need of protection from social injustice and exploitation, whereas other backward classes are a diverse collection of intermediate castes; others, or the so-called 'general' castes, are a residual group that enjoys higher status in the social group hierarchy. To compute a composite proxy indicator of income, a relative index of household wealth quintile was calculated using principal component analysis of the standard set of assets owned by the household, including ownership of consumer items and dwelling characteristics. Individuals were ranked on the basis of their household scores and divided into different quintiles, each representing 20% of the score, ranked between 1 (poorest) and 5 (richest).¹⁸

Statistical analysis

To attain the study's objectives, descriptive as well as multivariate analyses were deployed. χ^2 Test was used to understand the proportional difference in diabetes prevalence. Ordered logit regression model was applied to understand the net effects of the correlates of diabetes. The ordered logit model is used when the outcome variable is categorized on an ordinal scale, ordered by some conceptual or subjective criteria.¹⁹ It is the probability model used here for the diabetes score, which is ordered from 0 (categorized as normoglycemic) to 1 (impaired or pre-diabetes), to 2 (individuals with diabetes). The ordered logit model can be expressed as

 $\log[\gamma_j \chi/1 - \gamma_j \chi] = \theta_j - \beta^{\mathrm{T}} \chi$, where $j = 1, \dots, k-1$

for *k* categories of the response variable, where $\gamma_j = \Pr(Y \le \theta | \mathbf{x})$ is the cumulative probability up to and including *j*, for a covariate vector χ ; and θ_j is the cutpoint for the *j*th category. Exponentiating both sides of the equation gives the odds of falling into category *j* or lower versus falling into a category higher than *j* with a given set of covariates. The OR for a unit change in a particular covariate, say from $\chi = \mathbf{x}_1$ to $\chi = \mathbf{x}_2$, is given by

$$[\gamma_j \chi_1/(1-\gamma_j \chi_1)]/[\gamma_j \chi_{12}/(1-\gamma_j \chi_2)] = \exp(-\beta_{\chi}(\chi_1-\chi_2))$$

where β_{χ} is the coefficient of interest. A negative coefficient indicates that a higher value of the independent variable increases the odds of a lower value of the outcome. Since some of the selected predictor variables appear to have multicollinearity issues, a variance inflation factor (VIF) was estimated. All VIF values were much smaller than 5.0, suggesting that the possibility of high multicollinearity was low. The analyses were conducted using statistical software, Stata V.12. The variables found significant at p<0.05 level in the ordered logit regression model were brought to attention.

RESULTS

Table 1 represents the percentage distribution of diabetes status by selected background characteristics. The χ^2 test reveals that age group, education, social group, wealth quintile, body mass index, and physical activity level were measured to have a significant (p<0.05) difference in proportion within each group. Overall 2.9% and 3.3% of study participants were diagnosed with diabetes and pre-diabetes or impaired, respectively. The proportion of people diagnosed with diabetes was estimated high with increasing age, and almost 6% of those surveyed with education of 11th grade or above were diagnosed with diabetes. Almost 4% of people belonged to the 'others' social group, and nearly 7% of the richest respondents were diagnosed with diabetes. Nearly 7% of overweight/obese people were with diabetes, whereas close to 5% of people who had recorded low physical activity were with diabetes.

Table 2 shows the coefficient (β) with 95% CI estimated from ordered logit regression predicating the factors associated with diabetes. People who belonged to age group 15–44 years (β : 1.352; CI 0.802 to 1.902), and age group ≥45 years (β : 1.747; CI 1.203 to 2.291) have increasing risk of diabetes, as compared to people in the 18–24 years age group. Compared to the poorest, the richest have a higher probability (β : 0.730; CI 0.378 to 1.083) of being diagnosed with diabetes. As compared to people with normal BMI, people who are underweight tend to have low rate of diabetes (β : -0.419; CI -0.650 to 0.188); whereas overweight/obese people are more prone to being diagnosed with diabetes (β : 0.388; CI 0.147 to 0.628). With decreasing level of physical activity, people are more likely to be diagnosed with diabetes.

DISCUSSION

Using data of 7674 individuals from HDSS-BIRPOP, this study demonstrates the physiological and behavioral risk factors associated with the type 2 diabetes mellitus. While performing the analysis, the potential socioeconomic confounders were taken into consideration. Results indicate that age group, economic status, being overweight/obese, and low level of physical activity are the potential risk factors of diabetes. Although these risk factors appear as independent characteristics associated with diabetes, they are also interrelated.

Studies have demonstrated that the risk of type 2 diabetes tend to develop early in the life cycle among Indians, as compared to non-Indians. This study explains the higher probability of diabetes among people in their advanced age. This finding is consistent with the claim that the countries with the largest number of patients with diabetes are, and will be, India, China and the USA; in the first two countries diabetes occurs mostly in the age range of 45–64 years, in contrast with age >65 years in developed countries.⁹ A younger age at onset of diabetes has been noticed in Asian Indians in several studies, but this study focuses on an adult group aged ≥ 18 years only.

The high BMI for overweight/ obesity increases the probability of diabetes. Indians have, on average, a lower

BMI than those of European descent, and risk of diabetes starts to increase at very low levels of BMI for Indians.⁹ Higher percentage of body fat that is concentrated in the abdominal area is a key risk factor for developing diabetes. Also, irrespective of economic status, people living in rural India, increasingly refrain from labor-intensive activities leading to low physical activity.^{7 9 20} Efforts that focus on diet and physical activity have the potential to reduce the risk of obesity, the single most important risk factor for type 2 diabetes. Nutrition transition refers to a shift from consumption of simple, traditional foods to foods high in calories, sugar, and animal fat but low in vitamins and minerals derived from fruits and vegetables. Interactions between Westernized diets and lifestyle and genetic background may accelerate the growth of diabetes in the context of rapid nutrition transition. This food intake behavior is increasingly becoming common in rural India, which could lead to overweight/obesity, a prelude to diabetes.²¹ In earlier years there was a very low prevalence of diabetes among rural population. However, studies from Maharashtra²² and Andhra Pradesh²³ report very high prevalence of diabetes along with correlation with BMI. More commonly, rich and privileged groups have abandoned the traditional diet, and as their purchasing power is high, they are inclined to buy high calorie foods.

Early detection of individuals with diabetes or prediabetes and treatment in outpatient settings provides cost-saving ways of tackling the disease. Several randomized controlled trials have shown that lifestyle modification can reduce conversion from pre-diabetes to type 2 diabetes by 50%. Now the basic science needs to play the role for prevention and treatment for diabetes in new directions.²⁴ Integrating individual-level factors with environmental and policy factors to produce an interacting network are the need of the hour. An intervention could be designed to encourage physical activity, which in turn will maintain BMI and ultimately reduce the burden of diabetes. While designing the intervention, attention should be paid to aged people. Restoration of traditional diets in rural India should be encouraged, over food with high calorie concentration. Traditional Indian diets are largely based on polished white rice and refined wheat, having high glycemic index (GI) and glycemic load (GL) values.²⁵ Previously the metabolic effects of high GI or GL diets were nullified by high levels of physical activity.

In Brazil, Colombia, and Fiji, contextually relevant environmental interventions that were accompanied by national health education and promotional programs showed tremendous success in increasing physical activity at the population level.²⁶ Despite convincing evidence that type 2 diabetes is largely preventable through diet and lifestyle modifications, most current agricultural, nutrition, and urban planning policies are not conducive to nutritious eating and active living. Efforts to increase active travel in rural areas should be integral

Fable 1 Percentage distribution of diabetes status by select background characteristics						
	Normoglycemic	Impaired	Individuals with diabetes	Total	χ²	
Age group (years)					p<0.001	
18–24	98.31	1.35	0.34	890		
25–34	97.02	1.74	1.24	2013		
35–44	93.41	4.05	2.54	2048		
45 and above	90.01	4.63	5.36	2723		
Sex					p=0.077	
Male	93.05	3.61	3.35	3467		
Female	94.27	3.11	2.61	4207		
Education (grade)					p<0.001	
Illiterate	94.57	3.05	2.38	3277	P	
1–5	94.54	3.56	1.90	1629		
6–10	92.97	3.28	3 74	2163		
11 and above	89.59	4 46	5.95	605		
Religion	00.00		0.00	000	n=0.073	
Hindu	93.39	3 64	2 97	5353	p=0.070	
Islam and others	94 49	2.63	2.89	2321		
Social group	04.40	2.00	2.00	2021	n~0.001	
Scheduled caste	9/ 78	3 31	1 90	2625	p<0.001	
Scheduled tribe	94.70	2.01	1.90	646		
Other backward class	02 47	2.34	0.77	505		
Othera	93.47	3.70	2.11	2000		
Occupation	92.09	3.30	3.95	2090	n_0.962	
Unemployed	00.77	0.60	0.55	1050	p=0.863	
Nen eerrier beweeheld ebere	93.77	3.00	2.55	1059		
Non-earning nousenoid chore	94.06	3.03	2.90	3099		
Labor	93.38	3.48	3.13	3099		
Service/pensioner/retired	93.53	3.60	2.88	417	0.004	
wealth quintile	05 50	0.40	4.05	4000	p<0.001	
Poorest	95.53	3.12	1.35	1633		
Poorer	95.66	3.16	1.18	1359		
Middle	95.69	2.55	1.76	1530		
Richer	93.83	2.71	3.45	1622		
Richest	87.97	5.16	6.86	1530		
Body mass index					p<0.001	
Normal	93.54	3.57	2.89	3360		
Underweight	95.83	2.65	1.52	3095		
Overweight/obese	88.84	4.43	6.73	1219		
Physical activity level					p<0.001	
High	95.77	2.82	1.41	2911		
Medium	93.36	3.65	2.99	2575		
Low	91.41	3.66	4.94	2188		
Tobacco (smoking)					p=0.316	
No	93.97	3.25	2.78	5439		
Yes	93.11	3.53	3.36	2235		
Tobacco (chewing)					p=0.999	
No	93.72	3.33	2.95	5728		
Yes	93.73	3.34	2.93	1946		
Alcohol use					p=0.081	
No	93.74	3.20	3.05	6714		
Yes	93.54	4.27	2.19	960		
Total	93.72	3.34	2.95	7674		

to strategies to curb the level of diabetes, thus preventing NCDs in India.

Study limitations and way forward

Limitations of this study include the cross-sectional nature of data, which prevents us from drawing causal

inferences. Second, the study uses information on people who consume alcohol and tobacco users based on their recollection, which could generate recall errors. However, this study does provide physiological and behavioral risk factors of diabetes that can be used for future focused interventions. The study tool and the

Table 2 Coefficient (β) with 95% CI (estimated from ordered	logit regression pred	sion predicting the factors associated with diabetes		
	β	p Value	95% CI (lower bound, upper bound)		
Age group (years)					
18 to 24 (ref.)					
25 to 34	0.563	0.056	-0.014 to 1.141		
35 to 44	1.352	0.000	0.802 to 1.902		
45 and above	1.747	0.000	1.203 to 2.291		
Sex					
Male (ref.)					
Female	0.128	0.410	-0.177 to 0.434		
Education (grade)					
Illiterate (ref.)					
1–5	-0.040	0.785	-0.324 to 0.244		
6–10	0.012	0.937	-0.281 to 0.305		
11 and above	0.095	0.659	-0.326 to 0.515		
Religion					
Hindu (ref.)					
Islam and others	-0.178	0.197	-0.449 to 0.093		
Social group					
Scheduled caste (ref.)					
Scheduled tribe	-0.130	0.564	–0.571 to 0.311		
Other backward class	-0.254	0.245	-0.683 to 0.174		
Others	0.074	0.616	-0.216 to 0.365		
Occupation					
Unemployed (ref.)					
Non-earning household chore	-0.010	0.949	-0.306 to 0.287		
Labor	0.089	0.551	-0.204 to 0.382		
Service/pensioner/retired	0.039	0.870	-0.434 to 0.513		
Wealth quintile					
Poorest (ref.)	0.005	0.004			
Poorer	-0.025	0.891	-0.380 to 0.330		
Middle	-0.048	0.789	-0.396 to 0.301		
Richer	0.247	0.149			
	0.730	0.000	0.378 to 1.083		
Body mass index					
Normal (ret.)	0.410	0.000	0.050 to . 0.100		
Onderweight	-0.419	0.000			
Overweight/obese	0.388	0.002	0.147 10 0.628		
Physical activity level					
Madium	0.007	0.010	0.055 to 0.509		
	0.327	0.018	$0.055\ 10\ 0.596$		
LOW Tabaaaa (amaking)	0.594	0.000	0.273 10 0.916		
No (rof)					
No (lel.)	0.010	0.042	0.284 to 0.264		
Tebacco (chowing)	-0.010	0.942	-0.264 10 0.264		
No (ref.)					
Vos	0.015	0 000	0.212 to 0.241		
Alcohol use	0.015	0.900	-0.212 10 0.241		
No (ref.)					
	0 186	0.285	-0 155 to 0 527		
/cut1	4 575	0.205	3 855 to 5 296		
/cut2	5.306		4 668 to 6 125		
$Proh > v^2$	0.000		7.000 10 0.120		
l og likelihood	-1985 121				
n	7674				
	1017				

measure of physical activity patterns that we have used here is not a strictly standardized measure similar to Global Physical Activity Questionnaire (GPAQ) or other recent adaptations in India.²⁷ The scale used here is a crude measure based on initial exercises to understand the broad patterns of physical activity in the population, and identify the major activities according to their intensity levels. We are in the process of adapting the scale to

Epidemiology/health services research

the above measures in the ensuing future rounds of data collection. In future, the HDSS-BIRPOP also aims to collect further specific information on diabetes, and its related co-morbidity and mortality patterns among the sampled cohort population. In the present data set, we do not have any detailed information on food consumption or dietary intake, but investigation of dietary practices, an important behavioral factor that influences the risk of glucose intolerance is being included in an upcoming study on the study cohort. Additionally, in future studies every effort will be made to increase the participation rate.

Contributors AB, SM, AC and RKR. provided intellectual input through different stages of preparation of this manuscript equally.

Funding This work was supported by the Department of Health and Family Welfare, Government of West Bengal, India (Memo number: HF/O/MERT/ 1464/HSL (MISC)—35/2008).

Competing interests None declared.

Ethics approval This study was executed by the Birbhum Population Project (BIRPOP) of the Society for Health and Demographic Surveillance (SHDS), a Health and Demographic Surveillance System (HDSS). Ethical approval was obtained from the ethics review committee appointed by the chairperson of the governing body of HDSS-BIRPOP. This survey was conducted only after obtaining an informed and written consent from the participants. More information on the ethics review can be accessed from the BIRPOP's official website—http://www.shds.in/.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work noncommercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: http:// creativecommons.org/licenses/by-nc/4.0/

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work noncommercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: http:// creativecommons.org/licenses/by-nc/4.0/

REFERENCES

- United Nations (UN). Draft outcome document of the United Nations Summit for the adoption of the post-2015 development agenda. New York: UN, 2015.
- 2. World Health Organization (WHO). *Global Report on Diabetes*. Geneva: WHO, 2016.
- 3. World Health Organization (WHO). *Global action plan for the prevention and control of noncommunicable diseases 2013–2020.* Geneva: WHO, 2013.

APPENDIX 1

Our summary measure for individual physical activity is a combination of different types of physical activity and the reported frequency of such activity by individuals in a typical week. Information was collected on seven different types of physical activity: heavy manual work such as lifting loads, pulling carts and heavy agricultural work, was classified as 'heavy physical activity' (HPA); 'moderately high

- Zhang P, Zhang X, Brown J, et al. Global healthcare expenditure on diabetes for 2010 and 2030. *Diabetes Res Clin Pract* 2010;87:293–301.
- 5. International Diabetes Federation (IDF). *Diabetes atlas.* 6th edn, Brussels: IDF, 2013.
- Siegel K, Narayan KM, Kinra S. Finding a policy solution to India's diabetes epidemic. *Health Aff (Millwood)* 2008;27:1077–90.
- 7. Hu FB. Globalization of diabetes: the role of diet, lifestyle, and genes. *Diabetes Care* 2011;34:1249–57.
- Hu FB, Satija A, Manson JE. Curbing the diabetes pandemic: the need for global policy solutions. JAMA 2015;313:2319–20.
- Chan JC, Malik V, Jia W, *et al.* Diabetes in Asia: epidemiology, risk factors, and pathophysiology. *JAMA* 2009;301:2129–40.
- 10. Polonsky KS. The past 200 years in diabetes. *N Engl J Med* 2012;367:1332–40.
- 11. Yoon KH, Lee JH, Kim JW, *et al.* Epidemic obesity and type 2 diabetes in Asia. *Lancet* 2006;368:1681–8.
- Huxley R, James WP, Barzi F, et al, Obesity in Asia Collaboration. Ethnic comparisons of the cross-sectional relationships between measures of body size with diabetes and hypertension. Obes Rev 2008;9(Suppl 1):53–61.
- Willi C, Bodenmann P, Ghali WA, *et al.* Active smoking and the risk of type 2 diabetes: a systematic review and meta-analysis. *JAMA* 2007;298:2654–64.
- Koppes LL, Dekker JM, Hendriks HF, et al. Moderate alcohol consumption lowers the risk of type 2 diabetes: a meta-analysis of prospective observational studies. *Diabetes Care* 2005;28:719–25.
- Ghosh S, Barik A, Majumder S, *et al.* Health & demographic surveillance system profile: the Birbhum population project (Birbhum HDSS). *Int J Epidemiol* 2015;44:98–107.
- World Health Organization (WHO). *Definition and diagnosis of diabetes mellitus and intermediate hyperglycemia*. Geneva: WHO, 2006.
- WHO Expert Consultation. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet* 2004;363:157–63.
- 18. Rutstein SO, Johnson K. *The DHS Wealth Index*. DHS Comparative Reports No. 6. Calverton, MD: ORC Macro, 2004.
- McCullagh P, Nelder JA. Generalized linear models. Monographs on statistics and applied probability 37. London: Chapman and Hall, 1989.
- Ramachandran A, Snehalatha C, Kapur A, *et al*, Diabetes Epidemiology Study Group in India (DESI). High prevalence of diabetes and impaired glucose tolerance in India: National Urban Diabetes Survey. *Diabetologia* 2001;44:1094–101.
- Qi L, Cornelis MC, Zhang C, et al. Genetic predisposition, Western dietary pattern, and the risk of type 2 diabetes in men. Am J Clin Nutr 2009;89:1453–8.
- Deo SS, Zantye A, Mokal R, *et al.* To identify the risk factors for high prevalence of diabetes and impaired glucose tolerance in Indian rural population. *Int J Diab Dev Countries* 2006;26:19–23.
- Chow CK, Raju PK, Raju R, et al. The prevalence and management of diabetes in rural India. *Diabetes Care* 2006;29:1717–18.
- Ramachandran A, Snehalatha C, Mary S, *et al.*, Indian Diabetes Prevention Programme (IDPP). The Indian Diabetes Prevention Programme shows that lifestyle modification and metformin prevent type 2 diabetes in Asian Indian subjects with impaired glucose tolerance (IDPP-1). *Diabetologia* 2006;49:289–97.
- Popkin BM. The nutrition transition and obesity in the developing world. J Nutr 2001;131:871S–3S.
- Bauman A, Schoeppe S, Lewicka M, et al. Review of best practice in interventions to promote physical activity in developing countries: background document prepared for the WHO workshop on physical activity and public health. Beijing, China: World Health Organization, 2005.
- Matsuzaki M, Sullivan R, Ekelund U, et al. Development and evaluation of the Andhra Pradesh Children and Parent Study Physical Activity Questionnaire (APCAPS-PAQ): a cross-sectional study. BMC Public Health 2016;16:48.

physical activity' (MHPA) included activities such as carrying water, filling utensils with water, playing active games such as football, volleyball or swimming, and washing clothes; cooking and other normal household chores, washing utensils, dusting/mopping rooms and cleaning household premises were classified as 'moderate physical activity' (MPA); activities such as driving vehicles, tractors, light agricultural work (such as watering plants or spraying insecticides) were

Epidemiology/health services research

grouped into 'light physical activity' (LPA); finally, activities such as teaching, working in shops/vending wares in a stationary position were considered 'sedentary activity' (SA). Each of these component activities were coded on a scale of 0–2, where '0' denoted such activities were never conducted, '1' for occasional reporting of such activities, usually once or twice in a week, and '2' denoting usual or regular activity. Consequently, the scores for each of the five domains mentioned above were computed as a simple additive score of the component activity. Finally, these five physical activity domains were aggregated to derive a composite measure of physical activity as follows:

CPAS=(SA×0)+(LPA×2)+(MPA×4)+(MHPA×6)+(HPA×8); where CPAS denotes the Composite Physical Activity Score.ⁱ

The CPAS was then normalized into a scale of 0-100 to smoothen the distribution, and finally three equal-sized tercile classes were computed

from the normalized score variable. These three classes were subsequently used as ordered variables in subsequent multivariate analysis.

ⁱThe listing of specific component physical activities and their grouping into the five physical activity domains were based on an elaborate qualitative community-level ranking exercise, first involving a free-listing of the daily activities by selected groups of adult community members from both sexes, followed by pile-sorting of these activities into similar, hierarchical groups of physical activity in an increasing order of severity or involvement of physical stress or energy. The multipliers used in order to derive the aggregate score are arbitrary constants, but with the property that each higher-ordered domain is twice as stressful as its preceding domain. Using alternative scoring matrices such as equal (unitary) weights yielded results that were not significantly different.