



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/131619/>

Version: Accepted Version

Article:

de Lajudie, Philippe M and Young, J Peter W (2017) International Committee on Systematics of Prokaryotes Subcommittee for the Taxonomy of Rhizobium and Agrobacterium Minutes of the meeting, Budapest, 25 August 2016. *International Journal of Systematic and Evolutionary Microbiology*. pp. 2485-2494. ISSN: 1466-5034

<https://doi.org/10.1099/ijsem.0.002144>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

International Committee on Systematics of Prokaryotes
Subcommittee for the Taxonomy of *Rhizobium-Agrobacterium*
Minutes of the meeting, Budapest, August 25th, 2016

Philippe de Lajudie, *Secretary.*

J. Peter W. Young, *Chairperson.*

Minute 1. Call to order. The subcommittee closed meeting was called to order by Philippe de Lajudie (secretary) at 14:45 on July 25th, 2016 just before the 12th Nitrogen Fixation Conference in Budapest, Hungary.

Minute 2. Record of attendance.

Present: Philippe de Lajudie (Secretary, Institut de Recherche pour le Développement, Montpellier, France), J. Peter W. Young (University of York, UK), Kristina Lindström (University of Helsinki, Finland), Chang-Fu Tian (China Agricultural University), Julie Ardley (Murdoch University, Australia).

Apologies for absence: Bertrand Eardly (Penn State Berks College Reading, USA), Esperanza Martinez-Romero (Chairperson, Universidad Nacional Autónoma de Mexico, Cuernavaca, Mor., Mexico), Xavier Nesme (Institut National de la Recherche Agronomique, University of Lyon I, France), Pablo Vinuesa (Universidad Nacional Autónoma de Mexico, Cuernavaca, Mor., Mexico), Gehong Wei (China), Anne Willems (University of Gent, Belgium).

JPW Young was designated as the chairperson of the meeting and, in the name of the whole subcommittee, warmly welcomed Julie Ardley and Chang-Fu Tian for their first attendance as subcommittee members.

Minute 3. 2014 Minutes approval. Minutes of our last meeting (7 September 2014 and online) were unanimously approved. They have since been published (de Lajudie and Martinez Romero, 2017).

Minute 4. Resignations and thanks. Prof. Esperanza Martinez-Romero resigned as the chairperson of the subcommittee; the subcommittee is grateful to her for the many years since the 1990s that she held this position and for the accomplished work. The subcommittee assumes that Dr. Rosemary Sylvester-Bradley must have resigned long ago since she has not attended a subcommittee meeting since the late 1990s nor responded to messages. The subcommittee expressed their warm thanks to Prof. Wenxin Chen (China Agricultural University) and to Dr. Peter Van Berkum (USDA, Beltsville, USA) for their long and active years of membership with appreciated contributions. The subcommittee also remembers, with gratitude, Dr. Gisèle Laguerre (1957-2013, INRA France) and Prof. Brion Jarvis (1928-2015, New Zealand), who both actively participated in the subcommittee for many years with significant inputs.

Minute 5. Election of a new chairperson of the subcommittee. JPW Young was unanimously elected by vote of the members present as the new chairperson of the subcommittee.

Minute 6. New members. The subcommittee decided to encourage new members to join in order to face new challenges, to cover more items, diversity of rhizobial taxonomic groups, scopes, technical and geographical concerns, i.e. representatives from all continents, experts of genome-based taxonomic approaches, *Agrobacterium*, beta-rhizobia and new rhizobial groups, etc. Several names were suggested and the Secretary agreed to contact them.

Minute 7. The scope of the subcommittee. Originally restricted to the genera *Rhizobium* and *Agrobacterium*, the subcommittee concerns gradually extended to the taxonomy of « rhizobia », a general name covering all bacteria having the capacity to induce nodules on legumes where they fix dinitrogen. Therefore the subcommittee now deals with *Agrobacterium* and with all bacterial species nodulating legumes. These were all initially described as *Rhizobium* and later split in *Rhizobium sensu stricto*, *Azorhizobium*, *Bradyrhizobium*, *Sinorhizobium* (syn. *Ensifer*), *Mesorhizobium*, *Allorhizobium*, *Pararhizobium*, *Neorhizobium*, etc. Bacteria with this symbiotic ability have gradually also been discovered in other genera, i.e. *Methylobacterium*, *Devosia*, *Ciceribacter*, *Aminobacter*, *Bosea*, *Phyllobacterium*, *Ochrobactrum*, *Microvirga*, *Shinella*, *Labrys*, *Burkholderia*, *Paraburkholderia*, *Ralstonia*, *Cupriavidus*, *Herbaspirillum*. With « *Rhizobium* » having become “rhizobia”, including at least 21 genera in Alphaproteobacteria and Betaproteobacteria, the subcommittee discussed the opportunity to change its name accordingly. A longer name would not be convenient nor exhaustive either, and probably more rhizobia in more genera will be described in future, possibly in more proteobacterial classes. On the other hand, the general term “agrobacteria” covers all bacteria that form galls or root hypertrophies on plants by transfer of T-DNA, and are so far included in *Agrobacterium*, *Rhizobium*, *Pararhizobium* and *Allorhizobium*. Consequently, the subcommittee agreed to request a change in its name to the “**subcommittee for the taxonomy of rhizobia and agrobacteria**”, to reflect our growing knowledge of nitrogen-fixing microsymbionts of legumes and bacteria that transfer T-DNA to plants, consisting of diverse bacteria with similar accessory functions (nodule/gall forming ability), but the taxonomy of which follows the general bacterial rules, largely based on core genes/functions. There is precedent for a remit based on biological function rather than genera or families, such as the subcommittee for phototrophic bacteria. The subcommittee is aware that some rhizobia may fall within the taxonomic remit of another subcommittee, and agreed that species descriptions should have regard to the recommendations of that subcommittee as well as to any that are specific to rhizobia.

Thus the subcommittee understands its own main missions as:

1. To provide guidelines for describing new agrobacterial and rhizobial species,
2. To maintain information records about *Rhizobium*, *Agrobacterium*, agrobacterial and rhizobial species published,
3. To provide the list of taxa proposed since the last subcommittee meeting, and
4. To evaluate and comment on each genus/species validation, including rhizobia and agrobacteria described outside *Rhizobium* – *Agrobacterium*

Minute 8. New species. The number of rhizobial species increased exponentially in recent years. Table 1 lists the new taxa described since the last subcommittee meeting in September 2014, consisting of 2 new genera (*Neorhizobium*, *Pararhizobium*), 83 spp.

nov. (29 in *Rhizobium*, 24 in *Bradyrhizobium*, 12 in *Mesorhizobium*, 4 in *Ensifer*, 1 in *Pararhizobium*, 1 in *Agrobacterium*, 1 in *Bosea*, 1 in *Microvirga*, 1 in *Phyllobacterium*, 7 in *Burkholderia*, 2 in *Paraburkholderia*), 30 comb. nov. (3 in *Agrobacterium*, 6 in *Allorhizobium*, 3 in *Neorhizobium*, 4 in *Pararhizobium*, 12 in *Paraburkholderia*, 2 in *Ensifer*). *Rhizobium vignae* was found as a later synonym of (*Neo*)*rhizobium galegae* (Mousavi et al., 2014). *Rhizobium gei* (Shi et al., 2016) should be transferred as *Pararhizobium gei*. The rhizobial authentication of the 4 *Bosea* species (de Meyer and Willems, 2012 ; Safronova et al., 2015) may be regarded as pending as none of them was formally demonstrated to induce nodules on legumes so far, nor to harbor known nodulation genes (*nodA* and *nodC* could not be amplified) despite being isolated from nodules. As far as we know *Ciceribacter* is not confirmed to nodulate, despite having been isolated from the rhizosphere of chickpea. Authors publishing new species proposals outside IJSEM are encouraged to submit details of their papers to IJSEM for these species to appear in a validation list in that journal.

Minute 9. Type strains. The rule is that when a new species is validly described, the type strain should be deposited in two international collections (LMG, DSMZ, HAMBI, USDA, etc.) and available from them on request including a signed standard MTA (Material Transfer Agreement) form. Although the 'two collections rule' was not enforced rigorously in the past, recent examples show that it is being enforced: '*Allorhizobium taibaishanense*' comb. nov., '*Allorhizobium paknamense*' comb. nov. and '*Pararhizobium sphaerophysae*' comb. nov. (Mousavi et al., 2015) did not appear in IJSEM validation lists so far due to strain unavailability. In practice, some type strains may be difficult to obtain due to specific dissemination restrictions from authors in agreement with their specific national laws concerning biodiversity. In some countries the authorities do not allow any strain to be delivered abroad, but in general these restrictions vary from a simple additional declaration to the depositor, to the validation of an additional MTA by the depositor. Most of the restrictions imposed by MTAs, for example prohibitions on further distribution and commercial use, do not impede taxonomic study. However, a prohibition on the publication of the genome sequence of the type strain is problematic as it is now widely expected to have type strain genome sequences available, and the type strain genome sequence is likely to become mandatory for new species description quite soon. This issue may become a problem for taxonomists in countries with restrictions on genome sequencing to propose new species. Our subcommittee will initiate a discussion with the International Committee on Systematics of Prokaryotes on this sensitive case of national laws being problematic for taxonomists.

Minute 10. *Agrobacterium*. Recently, the number of agrobacterial species increased because of transferring the species *Rhizobium nepotum*, *Rhizobium pusense*, *Rhizobium skierniewicense* to the genus *Agrobacterium* (Mousavi et al, 2015; Table1) and also the description of a new species *Agrobacterium arsenijevicei* (Kuzmanović et al., 2016). *Agrobacterium tumefaciens* was found a later heterotypic synonym of *Agrobacterium radiobacter*, suggesting that *Ag. tumefaciens* should be amalgamated with *Ag. radiobacter* which has taxonomic priority over *Ag. tumefaciens* (Tindall, 2014). Even if treated as a later heterotypic synonym of *Ag. radiobacter*, the type species of the genus is *Ag. tumefaciens*, typified by the strain defined on the Approved Lists of Bacterial Names. This is the reason why it is recommended to retain ATCC 19358 (=LMG 140) as the type-

strain of *Ag. radiobacter* and B6 (=LMG 187=ATCC 23308) as the the type-strain of the genus *Agrobacterium*.

The genus *Agrobacterium* could be characterized by the presence of a large linear chromid [Ramirez-Bahena, 2014]. Based on the results of MLSA and genomic studies of the family *Rhizobiaceae* (Mousavi *et al.*, 2015 ; Ormeño-Orrillo *et al.*, 2015, unpublished data), the “*Rhizobium aggregatum* complex” consisting of the species *Agrobacterium albertimagni*, *R. rosettiformans*, *R. aggregatum*, *R. naphthalenivorans*, *R. selenitireducens*, and *R. daejeonense* should be considered as members of a separate genus, closely related to *Agrobacterium*. In addition, based on the results of MLSA (Mousavi *et al.*, 2014&2015), the species *Agrobacterium vitis* along with six *Rhizobium* species were transferred to the genus *Allorhizobium*. For other species that were named *Rhizobium* while genera were amalgamated the subcommittee recommends to use their original names (e.g. *Allorhizobium undicola*, *Agrobacterium larrymoorei* and *Ag. rubi*); *Rhizobium qilianshanense* (Xu *et al.*, 2013) should be transferred to *Allorhizobium*.

Minute 11. *Burkholderia*, *Paraburkholderia*, *Caballeronia*. The subcommittee discussed the particular case of *Burkholderia*. As currently circumscribed, *Burkholderia sensu lato* includes over 100 species, many of economic or clinical importance, that display remarkable diversity in their genome size, metabolism and ecological niches. Previous studies have established that *Burkholderia sensu lato* comprises three major clades: Clade I includes *Burkholderia cepacia* complex (Bcc) species, several plant pathogens and a group related to the animal pathogens *Burkholderia mallei* and *Burkholderia pseudomallei*; Clade II corresponds to the *Burkholderia glathei* clade; Clade III encompasses primarily environmental and plant-associated species, including rhizobial species and other members exhibiting beneficial plant growth promoting traits. In addition to the three major clades, *Burkholderia sensu lato* also includes several other deep-branching lineages, such as *Burkholderia rhizoxinica* and *Burkholderia endofungorum*; a group consisting of *Burkholderia caryophylli*, *Burkholderia symbiotica* and *Burkholderia soli*; and *Burkholderia andropogonis* (Depoorter *et al.*, 2016; Estrada-de los Santos *et al.*, 2016) and references therein). Informal proposals discussed splitting *Burkholderia sensu lato* into distinct genera, in particular to distinguish a plant-associated clade from a pathogen clade (Estrada-de los Santos *et al.*, 2013; Gyaneshwar *et al.*, 2011).

Sawana *et al.* (2014) were first to formally propose splitting *Burkholderia* into two genera, a revised *Burkholderia* retaining all pathogenic members and *Paraburkholderia* gen. nov. consisting of all plant-associated ones, with 62 new combinations. The genus *Paraburkholderia* was rapidly validated in IJSEM (Oren & Garrity, 2015b; Oren & Garrity, 2015c) and is now in use in databases.

In particular, symbiotic legume-nodulating *Burkholderia* species were transferred to *Paraburkholderia* with corresponding new combinations: the *Mimosa* symbionts as *Paraburkholderia tuberum*, *P. mimosarum*, *P. phymatum*, *P. nodosa*, *P. sabiae*, *P. symbiotica* and *P. diazotrophica*; *Lebeckia* symbionts as *P. sprentiae* comb. nov. and *P. dilworthii* comb. nov.; *Rhynchosia* symbionts as *P. rhynchosiae* comb. nov.; *Aspalathus* symbionts as *P. aspalathi* comb. nov.; *Phaseolus* symbionts as *P. caballeronis* comb. nov.. *Burkholderia sensu lato* was further split, with the concomitant publication of *Caballeronia* gen. nov. to accommodate twelve species of *Burkholderia* and *Paraburkholderia* (Dobritsa & Samadpour, 2016).

The taxonomic scheme of Sawana *et al.* (2014) was based on 16S rRNA phylogeny, as well as the percentage guanine plus cytosine content and conserved indels in whole

genome sequences of some 25 formally named *Burkholderia* species and several unclassified strains. However, Eberl and Vandamme (2016) criticized the lack of several type strains in the study by Sawana *et al.* and noted that the authors ignored phylogenetic heterogeneity, as revealed by 16S rRNA-based divergence and by differences in the distribution of 22 additional conserved sequence indels, in their simplistic inclusion of all species outside Clade I (*Burkholderia sensu stricto*) within the novel genus *Paraburkholderia*. According to this scheme, the deep-branching lineages of *B. rhizoxinica*, *B. endofungorum*, *B. caryophylli*, *B. symbiotica*, *B. soli* and *B. andropogonis* are all to be included within *Paraburkholderia*.

Similarly, Dobritsa & Samadpour (2016) proposed *Caballeronia* gen. nov. for species within the *B. glathei*-like Clade II, on the basis of 16S rRNA phylogeny and conserved indels, but have omitted a number of type strains in their study, including those of *B. caryophylli*, *B. soli* and *B. symbiotica*. We would further note a recent paper, which ascribed a strain to a novel species of *Burkholderia* on the basis of a phylogenetic analysis of 16S rRNA genes that did not include sequences from all type strains (Weber & King, 2017).

It is the opinion of this subcommittee that the studies describing *Caballeronia* gen. nov. and *Paraburkholderia* gen. nov. do not represent best taxonomic practice, and lead to confusion. Already, the phylogenetic classification of "*P. andropogonis*" within *Burkholderia* has been reassessed, with the result that it is now described as a new genus, *Robbsia andropogonis* gen. nov., comb. nov. (Lopes-Santos *et al.*, 2017).

While remaining within the Principles of the Bacterial Code (Parker *et al.*, 2015), in particular Principle 1(4) regarding freedom of taxonomic thought, we would note that implications exist outside prokaryotic taxonomy circles in the renaming of bacterial taxa, especially in the case of a genus of economic and clinical importance such as *Burkholderia*. In this regard, the following principles of the Code of Nomenclature of Prokaryotes should be considered:

Principle 1 - The essential points in nomenclature are as follows.

1. Aim at stability of names.
2. Avoid or reject the use of names which may cause error or confusion.
3. Avoid the useless creation of names.

Principle 9 - The name of a taxon should not be changed without sufficient reason based either on further taxonomic studies or on the necessity of giving up a nomenclature that is contrary to the Rules of this Code.

Notwithstanding that there are valid taxonomic reasons for splitting *Burkholderia sensu lato* into several genera, the members of this group cannot be described as pathogens or non-pathogens solely on the basis of their affiliation to a particular clade (Eberl & Vandamme, 2016). However, many strains, particularly rhizobial and plant beneficial strains from Clade III, show great potential as biofertilisers and bioremediants in sustainable agricultural systems. Research efforts directed towards robust characterisation and taxonomy of *Burkholderia sensu lato* species can help in realising this agricultural potential (Estrada-de los Santos *et al.*, 2016). Clearly, large-scale phylogenomic study is required for resolving these taxa.

Minute 12. The subcommittee home page. The sub-committee home page

(<http://edzna.ccg.unam.mx/rhizobial-taxonomy/node/2>) refers to LPSN

(<http://www.bacterio.net/>) for validly published bacterial species names and focuses on the taxonomy of rhizobia, their distribution and phylogenetic relationships with their close bacterial relatives, host plants, nodulation capacity and some features. The

subcommittee noticed the problem of obsolete genus and species affiliation in molecular sequence databases where the affiliation (genus, species) of the strain is forever retained as it was given at the time of its sequence deposit, with no further update of their later identification or genus/species transfers. The subcommittee will produce a list of the taxonomic transfer history and related sequence accession numbers for type strains to make this information available on its website.

Minute 13. Guidelines and Recommendations. The subcommittee recognizes the urgent need to update the last formal guidelines published for rhizobial species description 25 years ago (Graham et al., 1991) and will reactivate the process of publishing recommendations in IJSEM. There is a need of clarification regarding the minimal standards required to describe novel bacterial taxa and to better inform researchers and so seek to avoid errors in the classification of bacteria. It would appear, for example, that effective publication of a bacterial taxon does not have to be in print, and there is no longer a requirement for a new genus to be distinguishable by phenotypic characteristics.

Minute 14. Next meeting. Next meetings were planned to be held during the 20th International Congress on Nitrogen Fixation, Granada, Spain, 3-7 September 2017 and during the 13th European Nitrogen Fixation Conference (ENFC) in Stockholm, Sweden (2018).

Minute 15. Adjournment. The meeting was adjourned at 17:00 on August 25th 2016. The meeting continued online. Online discussion was adjourned on 18 April 2017.

Note Added in Preparation

Minute 16 (online) New members.

Several scientists were later contacted by the subcommittee secretary and Seyed Abdollah Mousavi (University of Helsinki, Finland), Alvaro Peix (Consejo Superior de Investigaciones Científicas, Instituto de Recursos Naturales y Agrobiología de Salamanca, Spain), Emma Steenkamp (University of Pretoria, South Africa), Jerri Zilli (Empresa Brasileira de Pesquisa Agropecuária, Seropédica, Brazil), Lionel Moulin (Institut de Recherche pour le Développement, Montpellier, France), Mitchell Andrews (Lincoln University, New Zealand), Ridha Mhamdi (Centre of Biotechnology of Borj-Cédria, Hammam-Lif, Tunisia) and Tomasz Stepkowski (Warsaw Life Sciences University, Warsaw, Poland) were further elected as members, and contributed in the online discussion to prepare the present minutes.

Minute 17. Current membership. The current members of the subcommittee are J.P.W. Young (UK, Chairperson), P. de Lajudie (France, Secretary), M. Andrews (New Zealand), J. Ardley (Australia), B. Eardly (USA), K. Lindström (Finland), R. Mhamdi (Tunisia), E. Martinez-Romero (Mexico), L. Moulin, (France), A. Mousavi, (Finland / Iran), X. Nesme (France), A. Peix, (Spain), E.T. Steenkamp (South Africa), T. Stepkowski (Poland), C.F. Tian (China), P. Vinuesa (Mexico), G. Wei (China), A. Willems (Belgium), J. Zilli, (Brazil).

References

Barauna AC, Rouws LF, Simoes-Araujo JL, Dos Reis Junior FB, Iannetta PP, Maluk M, Goi SR, Reis VM, James EK, Zilli JE. *Rhizobium altiplani* sp. nov., isolated from effective nodules on *Mimosa pudica* growing in untypically alkaline soil in central Brazil. *Int J Syst Evol Microbiol* 2016;66:4118-4124.

Bournaud C, Moulin L, Cnockaert M, de Faria SM, Prin Y, Severac D, Vandamme P. *Paraburkholderia piptadeniae* sp. nov. and *Paraburkholderia ribeironis* sp. nov., two root-nodulating symbiotic species of *Piptadenia gonoacantha* in Brazil. *Int J Syst Evol Microbiol* 2017;67:432-440.

Chahboune R, Carro L, Peix A, Ramírez-Bahena MH, Barrijal S, Velázquez E, Bedmar EJ. *Bradyrhizobium rifense* sp. nov. isolated from effective nodules of *Cytisus villosus* grown in the Moroccan Rif. *Syst Appl Microbiol* 2012;35:302-305.

Da Silva K, De Meyer SE, Rouws LFM, Farias ENC, Dos Santos MAO, O'hara G, Ardley JK, Willems A, Pitard RM, Zilli JE. *Bradyrhizobium ingae* sp. nov., isolated from effective nodules of *Inga laurina* grown in Cerrado soil. *Int J Syst Evol Microbiol* 2014;64:3395-3401.

Dall'agnol RF, Ribeiro RA, Delamuta JRMO, Ormeño-Orrillo E, Rogel MA, Andrade DS, Martínez-Romero E, Hungria M. *Rhizobium paranaense* sp. nov., an effective N₂-fixing symbiont of common bean (*Phaseolus vulgaris* L.) with broad geographical distribution in Brazil. *Int J Syst Evol Microbiol* 2014 ;64:3222-3229.

de Lajudie P, Martinez Romero E. International Committee on Systematics of Prokaryotes Subcommittee on the taxonomy of *Agrobacterium* and *Rhizobium*. Minutes of the meeting, 7 September 2014, Tenerife, Spain. *Int J Syst Evol Microbiol* 2017;67 :516-520.

De Meyer SE, Cnockaert M, Ardley JK, Maker G, Yates R, Howieson JG, Vandamme P. *Burkholderia sprentiae* sp. nov., isolated from *Lebeckia ambigua* root nodules. *Int J Syst Evol Microbiol* 2013;63:3950-3957.

De Meyer SE, Cnockaert M, Ardley JK, Trengove RD, Garau G, Howieson JG, Vandamme P. *Burkholderia rhynchosiae* sp. nov., isolated from *Rhynchosia ferulifolia* root nodules. *Int J Syst Evol Microbiol* 2013;63:3944-3949.

De Meyer SE, Cnockaert M, Ardley JK, Van Wyk BE, Vandamme PA, Howieson JG. *Burkholderia dilworthii* sp. nov., isolated from *Lebeckia ambigua* root nodules. *Int J Syst Evol Microbiol* 2014;64:1090-1095.

De Meyer SE, Tan HW, Andrews M, Heenan PB, Willems A. *Mesorhizobium calcicola* sp. nov., *Mesorhizobium waitakense* sp. nov., *Mesorhizobium sophorae* sp. nov., *Mesorhizobium newzealandense* sp. nov. and *Mesorhizobium kowhaii* sp. nov. isolated from *Sophora* root nodules. *Int J Syst Evol Microbiol* 2016;66:786-795.

De Meyer SE, Tan HW, Heenan PB, Andrews M, Willems A. *Mesorhizobium waimense* sp. nov. isolated from *Sophora longicarinata* root nodules and *Mesorhizobium cantuariense* sp. nov. isolated from *Sophora microphylla* root nodules *Int J Syst Evol Microbiol* 2015;65:3419-3426.

Delamuta JR, Ribeiro RA, Ormeno-Orrillo E, Parma MM, Melo IS, Martinez-Romero, E, Hungria M. *Bradyrhizobium tropiciagri* sp. nov. and *Bradyrhizobium embrapense* sp. nov., nitrogen-fixing symbionts of tropical forage legumes. *Int J Syst Evol Microbiol* 2015;65:4424-4433.

Depoorter E, Bull MJ, Peeters C, Coenye T, Vandamme P, Mahenthiralingam E. *Burkholderia*: an update on taxonomy and biotechnological potential as antibiotic producers. *Appl Microbiol Biotechnol* 2016;100:5215-5229.

Dobritsa AP, Samadpour M. Transfer of eleven *Burkholderia* species to the genus *Paraburkholderia* and proposal of *Caballeronia* gen. nov., a new genus to accommodate twelve species of *Burkholderia* and *Paraburkholderia*. *Int J Syst Evol Microbiol* 2016;66:2836-2846.

Durán D, Rey L, Mayo J, Zúñiga-Dávila D, Imperial J, Ruiz-Argüeso TS, Martínez-Romero E, Ormeño-Orrillo E. *Bradyrhizobium paxllaeri* sp. nov. and *Bradyrhizobium*

icense sp. nov., nitrogen-fixing rhizobial symbionts of Lima bean (*Phaseolus lunatus* L.) in Peru. *Int J Syst Evol Microbiol* 2014;64:2072-2078.

Durán D, Rey L, Navarro A, Busquets A, Imperial J, Ruiz-Argüeso T. *Bradyrhizobium valentinum* sp. nov., isolated from effective nodules of *Lupinus mariae-josephae*, a lupine endemic of basic-lime soils in Eastern Spain. *Syst Appl Microbiol.* 2014;37:336-341.

Eberl L, Vandamme P. Members of the genus *Burkholderia*: good and bad guys [version 1; referees: 3 approved]. *F1000Research* 2016, 5(F1000 Faculty Rev):1007.

Estrada-de los Santos, P., Vinuesa, P., Martínez-Aguilar, L., Hirsch, A. M. & Caballero-Mellado, J. Phylogenetic analysis of *Burkholderia* species by multilocus sequence analysis. *Current Microbiol* 2013;67:51-60.

Estrada-de los Santos P, Rojas-Rojas FU, Tapia-García EY, Vásquez-Murrieta MS, Hirsch AM. To split or not to split: an opinion on dividing the genus *Burkholderia*. *Annals Microbiol* 2016;66:1303-1314.

Graham, PH, Sadowsky MJ, Keyser HH, Barnett YM, Bradley RS, Cooper JE, De Ley D J, Jarvis BDW, Roslycky EB, Strijdom BW, Young JPW. Proposed Minimal Standards for the Description of New Genera and Species of Root- and Stem-Nodulating Bacteria. *Int J Syst Evol Microbiol* 1991;41:582-587.

Grisson CM, Jackson S, Merlot S, Dobson A, Grison C. *Rhizobium metallidurans* sp. nov., a symbiotic heavy metal resistant bacterium isolated from the *Anthyllis vulneraria* Zn-hyperaccumulator. *Int J Syst Evol Microbiol* 2015;65:1525-1530.

Grönemeyer JL, Chimwamurombe P, Reinhold-Hurek B. *Bradyrhizobium subterraneum* sp. nov., a symbiotic nitrogen-fixing bacterium from root nodules of groundnuts *Int J Syst Evol Microbiol* 2015;65:3241-3247.

Grönemeyer JL, Hurek T, Bünger W, Reinhold-Hurek B. *Bradyrhizobium vignae* sp. nov., a nitrogen-fixing symbiont isolated from effective nodules of *Vigna* and *Arachis*. *Int J Syst Evol Microbiol* 2016;66:62-69.

Gu, T, Sun LN, Zhang J, Sui XH, Li SP. *Rhizobium flavum* sp. nov., a triazophos-degrading bacterium isolated from soil under the long-term application of triazophos. *Int J Syst Evol Microbiol* 2014;64:2017-2022.

Guerrouj K, Ruiz-Diez B, Chahboune R, Ramirez-Bahena MH, Abdelmoumen H, Quinones MA, El Idrissi MM, Velazquez E, Fernandez-Pascual M, Bedmar EJ, Peix A. Definition of a novel symbiovar (sv. *retamae*) within *Bradyrhizobium retamae* sp. nov., nodulating *Retama sphaerocarpa* and *Retama monosperma*. *Syst Appl Microbiol* 2013;36: 218-223.

Gyaneshwar P, Hirsch AM, Moulin L, Chen WM, Elliott GN, Bontemps C, Estrada-de Los Santos P, Gross E, Dos Reis FB, Sprent JI, Young JP, James EK. Legume-nodulating betaproteobacteria: diversity, host range, and future prospects. *Mol Plant-Microbe Interact* 2011;24:1276-88.

Helene LC, Delamuta JR, Ribeiro RA, Ormeno-Orrillo E, Rogel MA, Martinez-Romero E, Hungria M. *Bradyrhizobium viridifuturi* sp. nov., encompassing nitrogen-fixing symbionts of legumes used for green manure and environmental services. *Int J Syst Evol Microbiol* 2015;65:4441-4448.

Jiao YS, Yan H, Ji ZJ, Liu YH, Sui XH, Wang ET, Guo BL, Chen WX, Chen WF. *Rhizobium sophorae* sp. nov. and *Rhizobium sophoriradicis* sp. nov., nitrogen-fixing rhizobial symbionts of the medicinal legume *Sophora flavescens*. *Int J Syst Evol Microbiol* 2015;65: 497-503.

Khalid R, Zhang YJ, Ali S, Sui XH, Zhang XX, Amara U, Chen WX, Hayat R. *Rhizobium pakistanensis* sp. nov., isolated from groundnut (*Arachis hypogaea*) nodules grown in rainfed Pothwar, Pakistan. *Antonie van Leeuwenhoek* 2015;107:281-290.

Kittiwongwattana C, Thawai C. *Rhizobium lemnae* sp. nov., a bacterial endophyte of *Lemna aequinoctialis*. *Int J Syst Evol Microbiol* 2014;64:2455-2460.

Kumar N, Lad G, Giuntini E, Kaye ME, Udomwong P, Shamsani NJ et al. Bacterial genospecies that are not ecologically coherent: population genomics of *Rhizobium leguminosarum*. *Open Biology* 2015;5:140133–140133.

Kuzmanović N, Puławska J, Prokića A, Ivanovića M, Zlatkovića N, Jones JB, Obradović A. *Agrobacterium arsenijevicei* sp. nov., isolated from crown gall tumors on raspberry and cherry plum. *Syst Appl Microbiol* 2016;38:373–378.

Grönemeyer JL, Hurek T, Reinhold-Hurek B. *Bradyrhizobium kavangense* sp. nov., a symbiotic nitrogen-fixing bacterium from root nodules of traditional Namibian pulses. *Int J Syst Evol Microbiol* 2015;65:4886-4894.

Le Quéré A, Tak N, Singh Gehlot H, Lavire C, Meyer T, Chapulliot D, Rathi S, Sakrouhi I, Rocha G, Rohmer M, Severac D, Filali-Maltouf A, Munive JA. Genomic characterization of *Ensifer aridi*, a proposed new species of nitrogen-fixing rhizobium recovered from Asian, African and American deserts. *BMC Genomics* 2017;18:85.

Li Y, Yan J, Yu B, Wang ET, Li X, Yan H, Liu W, Xie Z. *Ensifer alkalisoli* sp. nov. isolated from root nodules of *Sesbania cannabina* grown in saline-alkaline soils. *Int J Syst Evol Microbiol* 2016;66:5294-5300.

Li YH, Wang R, Zhang XX, Young JP, Wang ET, Sui XH, Chen WX. *Bradyrhizobium guangdongense* sp. nov. and *Bradyrhizobium guangxiense* sp. nov., isolated from effective nodules of peanut. *Int J Syst Evol Microbiol* 2015;65:4655-4661.

Lin SY, Hsu YH, Liu YC, Hung MH, Hameed A, Lai WA, Yen WS, Young CC. *Rhizobium straminoryzae* sp. nov., isolated from the surface of rice straw. *Int J Syst Evol Microbiol* 2014;64:2962-2968.

Lin SY, Hung MH, Hameed A, Liu YC, Hsu YH, Wen CZ, Arun AB, Busse HJ, Glaeser SP et al. *Rhizobium capsici* sp. nov., isolated from root tumor of a green bell pepper (*Capsicum annuum* var. *grossum*) plant. *Antonie van Leeuwenhoek* 2015;107:773-784.

Liu TY, Li Jr Y, Liu XX, Sui XH, Zhang XX, Wang ET, Chen WX, Chen WF, Puławska, J. *Rhizobium cauense* sp. nov., isolated from root nodules of the herbaceous legume *Kummerowia stipulacea* grown in campus lawn soil. *Syst Appl Microbiol* 2012;35:415-420.

Lopes-Santos L, Castro DBA, Ferreira-Tonin M, Corrêa DBA, Weir BS, Park D, Ottoboni LMM, Neto JR, Destéfano SAL. Reassessment of the taxonomic position of *Burkholderia andropogonis* and description of *Robbsia andropogonis* gen. nov., comb. nov. *Antonie van Leeuwenhoek* 2017;1-10.

Lu JK, Dou YJ, Zhu YJ, Wang SK, Sui XH, Kang LH. *Bradyrhizobium ganzhouense* sp. nov., an effective symbiotic bacterium isolated from *Acacia melanoxylon* R. Br. nodules. *Int J Syst Evol Microbiol* 2014;64:1900-1905.

Martínez-Aguilar L, Salazar-Salazar C, Méndez RD, Caballero-Mellado J, Hirsch AM, Vásquez-Murrieta MS, Estrada-De Los Santos P. *Burkholderia caballeronis* sp. nov., a nitrogen fixing species isolated from tomato (*Lycopersicon esculentum*) with the ability to effectively nodulate *Phaseolus vulgaris*. *Antonie van Leeuwenhoek*, 2014;104:1063-1071.

Martínez-Hidalgo P, Ramírez-Bahena MH, Flores-Felix JD, Rivas R, Igual JM, Mateos PF, Martínez-Molina E, León-Barrios M, Peix A, Velázquez E. Revision of the taxonomic status of type strains of *Mesorhizobium loti* and reclassification of strain USDA 3471^T as the type strain of *Mesorhizobium erdmanii* sp. nov. and ATCC 33669^T as the type strain of *Mesorhizobium jarvisii* sp. nov. *Int J Syst Evol Microbiol* 2015;65:1703-1708.

Martínez-Hidalgo P, Ramírez-Bahena MH, Flores-Felix JD, Igual JM, Sanjuán J, León-Barrios M, Peix A, Velázquez E. Reclassification of strains MAFF 303099T and R7A into *Mesorhizobium japonicum* sp. nov. *Int J Syst Evol Microbiol* 2016;66:4936-4941.

Mavengere NR, Ellis AG, Le Roux, JJ. *Burkholderia aspalathi* sp. nov., isolated from root nodules of the South African legume *Aspalathus abietina* Thunb. *Int J Syst Evol Microbiol* 2014;64:1906-1912.

Mnasri B, Liu TY, Saidi S, Chen WF, Chen WX, Zhang XX, Mhamdi R. *Rhizobium azibense* sp. nov., a nitrogen fixing bacterium isolated from root-nodules of *Phaseolus vulgaris*. *Int J Syst Evol Microbiol* 2014;64:1501-1506.

Mousavi SA, Österman J, Wahlberg N, Nesme X, Lavire C, Vial L, Paulin L, de Lajudie P, Lindström K. Phylogeny of the *Rhizobium-Allorhizobium-Agrobacterium* clade supports the delineation of *Neorhizobium* gen. nov. *Syst Appl Microbiol* 2014;37:208-215.

Mousavi SA, Willems A, Nesme X, de Lajudie P, Lindström K. Revised phylogeny of *Rhizobiaceae*: proposal of the delineation of *Pararhizobium* gen. nov., and 13 new species combinations. *Syst Appl Microbiol* 2015;38:84-90.

Nguyen TM, Pham VHT, Kim J. *Mesorhizobium soli* sp. nov., a novel species isolated from the rhizosphere of *Robinia pseudoacacia* L. in South Korea by using a modified culture method. *Antonie van Leeuwenhoek* 2015;108:301-310.

Oren A, Garrity GM. List of new names and new combinations previously effectively, but not validly, published. *Int J Syst Evol Microbiol* 2015a;65:741-744.

Oren A, Garrity GM. List of new names and new combinations previously effectively, but not validly, published. *Int J Syst Evol Microbiol* 2015b;65:2017-2025.

Oren A, Garrity GM. List of new names and new combinations previously effectively, but not validly, published. *Int J Syst Evol Microbiol* 2015c;65:2777-2783.

Oren A, Garrity GM. List of new names and new combinations previously effectively, but not validly, published. *Int J Syst Evol Microbiol* 2016;66:4299-4305.

Ormeño-Orrillo E, Martínez-Romero E. Phenotypic tests in *Rhizobium* species description: an opinion and (a sympatric speciation) hypothesis. *Syst Appl Microbiol* 2013;36:145-147.

Parag B, Sasikala Ch, Ramana ChV. Molecular and culture dependent characterization of endolithic bacteria in two beach sand samples and description of *Rhizobium endolithicum* sp. nov. *Antonie van Leeuwenhoek* 2014;104:1235-1244.

Parker CT, Tindall BJ, Garrity GM. International code of nomenclature of prokaryotes. *Int J Syst Evol Microbiol* 2015; doi: [10.1099/ijsem.0.000778](https://doi.org/10.1099/ijsem.0.000778)

Peix A, Ramírez-Bahena MH, Flores-Félix JD, Alonso De La Vega P, Rivas R, Mateos PF, Igual JM, Martínez-Molina E, Trujillo ME, Velázquez E. Revision of the taxonomic status of the species *Rhizobium lupini* and reclassification as *Bradyrhizobium lupini* comb. nov. *Int J Syst Evol Microbiol* 2015;65:1213-1219.

Puławska J, Kuzmanović N, Willems A, Pothier JF. *Pararhizobium polonicum* sp. nov. isolated from tumors on stone fruit rootstocks. *Syst Appl Microbiol* 2016;39:164-169.

Radl V, Simões-Araújo JL, Leite J, Passos SR, Martins LM, Xavier GR, Rumjanek NG, Baldani JJ, Zilli JE. *Microvirga vignae* sp. nov., a root nodule symbiotic bacterium isolated from cowpea grown in semi-arid Brazil. *Int J Syst Evol Microbiol* 2014;64:725-730.

Ramírez-Bahena MH, Flores-Félix JD, Chahboune R, Toro M, Velázquez E, Peix A. *Bradyrhizobium centrosemae* (symbiovar centrosemae) sp. nov., *Bradyrhizobium americanum* (symbiovar phaseolarum) sp. nov. and a new symbiovar (tropici) of

Bradyrhizobium viridifuturi establish symbiosis with *Centrosema* species native to America. *Syst Appl Microbiol* 2016;39:378-83.

Ramírez-Bahena MH, Vial L, Lassalle F, Diel D, Chapulliot D, Daubin V, Nesme X, Muller D. Single acquisition of protelomerase gave rise to speciation of a large and diverse clade within the *Agrobacterium/Rhizobium* supercluster characterized by the presence of a linear chromid. *Mol Phyl Evol* 2014;73:202-207.

Rashid MH, Young JPW, Overall I, Clercx P, Willems A, Santhosh Braun M, Wink M. Average nucleotide identity of genome sequences supports the description of *Rhizobium lentis* sp. nov., *Rhizobium bangladeshense* sp. nov. and *Rhizobium binae* sp. nov. from lentil (*Lens culinaris*) nodules. *Int J Syst Evol Microbiol* 2015;65:3037-3045.

Ribeiro RA, Martins TB, Ormeño-Orrillo E, Marcon Delamuta JR, Rogel MA, Martínez-Romero E, Hungria M. *Rhizobium ecuadorensis* sp. nov., an indigenous N₂-fixing symbiont of the Ecuadorian common bean (*Phaseolus vulgaris* L.) genetic pool. *Int J Syst Evol Microbiol* 2015;65:3162-3169.

Román-Ponce, B., Jing Zhang, Y., Soledad Vásquez-Murrieta, M., Hua Sui, X., Feng Chen, W., Carlos Alberto Padilla, J., Wu Guo, X., Lian Gao, J., Yan, J., Hong Wei, G. & Tao Wang, E. (2016). *Rhizobium acidisoli* sp. nov., isolated from root nodules of *Phaseolus vulgaris* in acid soils. *Int J Syst Evol Microbiol* 66, 398-406.

Rozahon M, Ismayil N, Hamood B, Erkin R, Abdurahman M, Mamtimin H, Abdukerim M, Lal R, Rahman E. *Rhizobium populi* sp. nov., an endophytic bacterium isolated from *Populus euphratica*. *Int J Syst Evol Microbiol* 2014;64:3215-3221.

Safronova VI, Kuznetsova IG, Sazanova AL, Kimeklis AK, Belimov AA, Andronov EE, Pinaev AG, Chizhevskaya EP, Pukhaev AR, Popov KP, Willems A, Tikhonovich IA. *Bosea vaviloviae* sp. nov., a new species of slow-growing rhizobia isolated from nodules of the relict species *Vavilovia formosa* (Stev.) Fed. *Antonie van Leeuwenhoek* 2015;107:911-920.

Saïdi S, Ramírez-Bahena MH, Santillana N, Zúñiga D, Álvarez-Martínez E, Peix A, Mhamdi R, Velázquez E. *Rhizobium laguerreae* sp. nov. nodulates *Vicia faba* on several continents. *Int J Syst Evol Microbiol* 2014;64:242-247.

Shamseldin A, Carro L, Peix A, Velázquez E, Moawad H, Sadowsky MJ. The symbiovar trifolii of *Rhizobium bangladeshense* and *Rhizobium aegyptiacum* sp. nov. nodulate *Trifolium alexandrinum* in Egypt. *Syst Appl Microbiol* 2016;39:275-279.

Sánchez M, Ramírez-Bahena MH, Peix A, Lorite MJ, Sanjuán J, Velázquez E, Monza J. *Phyllobacterium loti* sp. nov. isolated from nodules of *Lotus corniculatus*. *Int J Syst Evol Microbiol* 2014;64:781-786.

Sawana A, Adeolu M, Gupta RS. Molecular signatures and phylogenomic analysis of the genus *Burkholderia*: proposal for division of this genus into the emended genus *Burkholderia* containing pathogenic organisms and a new genus *Paraburkholderia* gen. nov. harboring environmental species. *Front Genet* 2014;5:429.

Sheu SY, Chou JH, Bontemps C, Elliott GN, Gross E, James EK, Sprent JI, Young JPW, Chen WM. *Burkholderia symbiotica* sp. nov., isolated from root nodules of *Mimosa* spp. native to north-east Brazil. *Int J Syst Evol Microbiol* 2012;62:2272-2278.

Sheu SY, Chou JH, Bontemps C, Elliott GN, Gross E, Dos Reis Junior FB, Melkonian R, Moulin L, James EK, Sprent JI, Young JPW, Chen WM. *Burkholderia diazotrophica* sp. nov., isolated from root nodules of *Mimosa* spp. *Int J Syst Evol Microbiol* 2013;63:435-441.

Sheu SY, Huang HW, Young CC, Chen WM. *Rhizobium alvei* sp. nov., isolated from a freshwater river. *Int J Syst Evol Microbiol* 2015;65:472-478.

Shi X, Li C, Zhao L, Si M, Zhu L, Xin K, Chen C, Wang Y, Shen X, Zhang L. *Rhizobium gei* sp. nov., a bacterial endophyte of *Geum aleppicum*. *Int J Syst Evol Microbiol* 2016;66:4282-4288.

Silva FV, De Meyer SE, Simões-Araújo JL, Da Costa Barbé T, Xavier GR, O'hara G, Ardley JK, Rumjanek NG, Willems A, Zilli, JE. *Bradyrhizobium manausense* sp. nov., isolated from effective nodules of *Vigna unguiculata* grown in Brazilian Amazonian rainforest soils. *Int J Syst Evol Microbiol* 2014;64:2358-2363.

Tan ZY, Kan F, Peng GX, Wang ET, Reinhold-Hurek B, Chen WX. *Rhizobium yanglingense* sp. nov., isolated from arid and semi-arid regions in China. *Int J Syst Evol Microbiol* 2001;51:909-914.

Tindall BJ. *Agrobacterium radiobacter* (Beijerinck and van Delden 1902) Conn 1942 has priority over *Agrobacterium tumefaciens* (Smith and Townsend 1907) Conn 1942 when the two are treated as members of the same species based on the principle of priority and Rule 23a, Note 1 as applied to the corresponding specific epithets. *Int J Syst Evol Microbiol* 2014;64:3590–3592.

Torres Tejerizo G, Rogel MA, Ormeño-Orrillo E, Althabegoiti MJ, Nilsson JF, Niehaus K, Schlüter A, Pühler A, Del Papa MF, Lagares A, Martínez-Romero E, Pistorio M. *Rhizobium favelukesii* sp. nov., isolated from the root nodules of alfalfa (*Medicago sativa* L). *Int J Syst Evol Microbiol* 2016;66:4451-4457.

Wang R, Chang YL, Zheng WT, Zhang D, Zhang XX, Sui XH, Wang ET, Hu JQ, Zhang LY, Chen WX. *Bradyrhizobium arachidis* sp. nov., isolated from effective nodules of *Arachis hypogaea* grown in China. *Syst Appl Microbiol* 2013;36:101-105.

Wang YC, Wang F, Hou BC, Wang ET, Chen WF, Sui XH, Chen WX, Li Y, Zhang YB. Proposal of *Ensifer psoraleae* sp. nov., *Ensifer sesbaniae* sp. nov., *Ensifer morelense* comb. nov. and *Ensifer americanum* comb. nov. *Syst Appl Microbiol* 2013;36: 467-473.

Weber CF, King GM. Volcanic soils as sources of novel CO-oxidizing *Paraburkholderia* and *Burkholderia*: *Paraburkholderia hiiakae* sp. nov., *Paraburkholderia metrosideri* sp. nov., *Paraburkholderia paradisi* sp. nov., *Paraburkholderia peleae* sp. nov., and *Burkholderia alpina* sp. nov. a member of the *Burkholderia cepacia* complex. *Front Microbiol* 2017;8: Article 207. doi.org/10.3389/fmicb.2017.00207.

Xu L, Zhang Y, Deng ZS, Zhao L, Wei XL, Wei GH. *Rhizobium qilianshanense* sp. nov., a novel species isolated from root nodule of *Oxytropis ochrocephala* Bunge in China. *Antonie Van Leeuwenhoek* 2013;103:559-65.

Yan H, Yan J, Sui XH, Wang ET, Chen WX, Zhang XX, Chen WF. *Ensifer glycinis* sp. nov., a rhizobial species associated with species of the genus *Glycine*. *Int J Syst Evol Microbiol* 2016;66:2910-2916.

Yan J, Yan H, Liu LX, Chen WF, Zhang XX, Verastegui-Valdes MM, Wang ET, Han XZ. *Rhizobium hidalgonense* sp. nov., a nodule endophytic bacterium of *Phaseolus vulgaris* in acid soil. *Arch Microbiol* 2017;199:97-104.

Yao Y, Sui XH, Zhang XX, Wang ET, Chen WX. *Bradyrhizobium erythrophlei* sp. nov. and *Bradyrhizobium ferriligni* sp. nov., isolated from effective nodules of *Erythrophleum fordii*. *Int J Syst Evol Microbiol* 2015;65:1831-1837.

Yu X, Cloutier S, Tambong JT, Bromfield ESP. *Bradyrhizobium ottawaense* sp. nov., a symbiotic nitrogen fixing bacterium from root nodules of soybeans in Canada. *Int J Syst Evol Microbiol* 2014;64:3202-3207.

Zhang YJ, Zheng WT, Everall I, Young JP, Zhang XX, Tian CF, Sui XH, Wang ET, Chen WX. *Rhizobium anhuiense* sp. nov. isolated from effective nodules of *Vicia faba* and *Pisum sativum*. *Int J Syst Evol Microbiol* 2015;65:2960-2967.

Zhang XX, Gao JS, Cao YH, Sheirdil RA, Wang XC, Zhang L. *Rhizobium oryzicola* sp. nov., potential plant-growth-promoting endophytic bacteria isolated from rice roots. *Int J Syst Evol Microbiol* 2015;65:2931-2936.

Zhang XX, Tang X, Sheirdil RA, Sun L, Ma XT. *Rhizobium rhizoryzae* sp. nov., isolated from rice roots *Int J Syst Evol Microbiol* 2014;64:1373-1377.

Zhang YJ, Zheng WT, Everall I, Young JP, Zhang XX, Tian CF, Sui XH, Wang ET, Chen WX. *Rhizobium anhuiense* sp. nov., isolated from effective nodules of *Vicia faba* and *Pisum sativum*. *Int J Syst Evol Microbiol* 2015;65:2960-2967.

Zhu YJ, Kun J, Chen YL, Wang SK, Sui XH, Kang LH. *Mesorhizobium acaciae* sp. nov., isolated from root nodules of *Acacia melanoxylon* R.Br. *Int J Syst Evol Microbiol* 2015;65:3558-3563.

Zilli JE, Baraúna AC, Da Silva K, De Meyer SE, Farias ENC, Kaminski PE, Da Costa IB, Ardley JK, Willems A, Camacho NN, Dourado FDS, O'hara G. *Bradyrhizobium neotropicale* sp. nov., isolated from effective nodules of *Centrolobium paraense*. *Int J Syst Evol Microbiol* 2014;64:3950-3957.