

1 **Thymic B cell-mediated attack of thymic stroma precedes Type 1**
2 **Diabetes Development.**

3 Ana Isabel Pinto, Jennifer Smith, Miriam R. Kissack, Karen Hogg and E. Allison Green*

4 Centre for Immunology and Infection, Hull York Medical School and the Department of
5 Biology, University of York, Wentworth Way, York YO10 5DD.

6 *`Address for correspondence: Dr. E. Allison Green; allison.green@york.ac.uk

7

8 **Abstract**

9 Type 1 diabetes results from a co-ordinated autoimmune attack of insulin producing beta
10 cells in the pancreas by the innate and adaptive immune systems, beta cell death being
11 predominantly T cell-mediated. In addition to T cells, peripheral B cells are important in type
12 1 diabetes progression. The thymus of mice and man also contain B cells, and lately they
13 have been linked to central tolerance of T cells. The role of thymic B cells in type 1 diabetes
14 is undefined. Here we show there are abnormalities in the thymic B cell compartment prior to
15 beta cell destruction and type 1 diabetes manifestation.

16 Using non-obese diabetic (NOD) mice, we document that preceding type 1 diabetes
17 development, there is significant accumulation of thymic B cells-partly through *in situ*
18 development- and the putative formation of ectopic germinal centres. In addition, in NOD
19 mice we quantify thymic plasma cells and observe *in situ* binding of immunoglobulins to
20 undefined antigens on a significant proportion of medullary thymic epithelial cells. In
21 contrast, no ectopic germinal centres, or pronounced intrathymic autoantibodies are
22 detectable in animals not genetically predisposed to developing type 1 diabetes. Binding of
23 autoantibodies to thymic stroma correlates with apoptosis of medullary thymic epithelial
24 cells, including insulin-expressing cells. In contrast, apoptosis of medullary thymic epithelial
25 cells was decreased by 50% in B cell deficient NOD mice suggesting intrathymic
26 autoantibodies may selectively target certain medullary thymic epithelial cells for destruction.
27 Furthermore, we observe that these thymic B cell-associated events correlated with an
28 increased prevalence of premature thymic emigration of T cells.

29 Together our data suggests that the thymus may be a principal autoimmune target in type 1
30 diabetes and contributes to disease progression.

Introduction

The thymus is a primary lymphoid organ involved in shaping the T cell repertoire. Sequential compartmentalisation of developing T cells into the cortical region of the thymus, and subsequently the medulla enable the effective positive and negative selection events, respectively, that are integral in generating an immature T cell repertoire enriched to respond to pathogens but not self-tissue- termed central tolerance [1]. Central to this role for the thymus, are the medullary thymic epithelial cells (mTECS) [2; 3]; capable of autoimmune regulator (AIRE) driven expression of peripheral tissue specific antigens (TSAs) [4] and presentation in the context of MHC class I or class II molecules, they trigger events that lead to apoptosis of developing T cells bearing high affinity receptors for self-peptides.

Although extensive studies have documented the importance of mTECs for negative selection of autoreactive T cells [5], other antigen presenting cell (APC) populations within the thymus have also been shown to participate in T cell negative selection, particularly dendritic cells [6]. A newer member of this family of APCs involved in negative selection are the thymic B cells [7; 8], although it is still not clear how significant their role is in negative selection with respect to that of mTECs and thymic DCs [9]. Thymic B cells are present both in man and mice; constituting a minor population of the thymic cellular pool, they are detectable in foetal through to adult mammals [10; 11]. Thymic B cells have a similar phenotype to peripheral B2 cells [12; 13], and their thymic frequency is stable from birth onwards. Interestingly expansion of thymic B cells in Myasthenia and Systemic Lupus Erythematosus (SLE) patients [14; 15], or animal models of SLE have been linked to disease progression, suggesting thymic B cells may have a potential role in breakdown of central tolerance [16].

Type 1 Diabetes (T1D) is an autoimmune condition where insulin secreting β cells in the islets of Langerhans are destroyed through co-ordinated attack by both the innate and

adaptive immune systems; the final assault being perpetuated by CD8⁺ cytotoxic T cells [17; 18; 19]. Defects in central tolerance is linked to emergence of a β cell-specific T cell repertoire [20], yet definitive understanding of the mechanisms underlying defective central tolerance are unclear. Much of our understanding of the immunological events leading to β cell pathology has been derived from the non-obese diabetic (NOD) mouse, a murine model that spontaneously develops T1D with many similarities to those seen in man [21]. Studies in NOD mice show T1D is a progressive condition, with priming of the T cell repertoire to β cell antigens in early life followed by infiltration of islets with immune cells (termed insulitis), a period of regulation of the autoreactive response, but ultimately an aggressive and sustained attack on the β cells. It is not clear what immunological event triggers this final stage of the disease.

B cells, too, are known to be important in the T1D process both in man and in NOD mice; abnormally high numbers of islet-infiltrating B cells is linked to rapid progression to T1D in young children [22], and increasing diversity of serum antibodies for β cell antigens increase substantially the risk factor of developing T1D genetically-predisposed children [23]. In NOD mice, genetic or immunological ablation of B cells protects against T1D development [24; 25], and in both diabetic NOD mice and diabetic patients, depletion of B cells can resolve the condition albeit transiently [26; 27]. To date, the role for B cells in T1D progression has been linked to their peripheral APC function- their ability to present β cell antigens to β -reactive CD4⁺ T cells [28] enhances CD4⁺ T helper cell activation of CD8⁺ T cells, and in islets B cells provide survival signals for activated CD8⁺ T cells enabling a sustained cytotoxic attack on β cells [29].

Here we provide that the thymus of diabetes-prone NOD mice displays evidence of autoreactivity prior to type 1 diabetes development. . We show that the post-insulitic/pre-diabetic phase is characterised by abnormally high thymic B cell development, B cell

81 accumulation in follicles at the cortical-medullary junction and the emergence of ectopic
82 germinal centres. Intrathymic autoantibodies bind to undefined antigens on selective mTECs.
83 Subsequently increased mTECs apoptosis, including insulin-expressing mTECs occurs.
84 These events correlate with increased levels of peripheral T cells that have a RAG-GFP
85 phenotype akin to thymocytes that have yet to undergo negative selection, suggesting in NOD
86 mice thymic B cells may contribute to decreased efficacy of negative selection of
87 autoreactive T cells.

88 Our data provides new insights into thymic abnormalities that precede β cell destruction and
89 highlight the importance of focusing research on these unique thymic B cells as mediators of
90 this chronic condition.

Methods

Mice

C57BL/6 (B6), FVB.RAGp2-GFP reporter mice [30] and NOD. μ MT^{-/-} mice [25] have been described elsewhere. FVB.RAGp2-GFP reporter mice were backcrossed 20 generations to either non-obese diabetic (NOD) mice (NOD.RAGp2-GFP) or NOD. μ MT^{-/-} mice (NOD. μ MT^{-/-}.RAGp2-GFP mice). All mice used in this study were maintained under specific-pathogen free conditions with a 12 hour light-dark cycle and fed normal chow. All animal experimental procedures were carried out in accordance with the Animals and Scientific Procedures Act 1986 were approved by the University of York Animal Welfare and Ethics Review Board and conducted under UK Home Office License approval conforming to ARRIVE guidelines (<https://www.nc3rs.org.uk/arrive-guidelines>). Diabetes development was determined by assessing urine glucose levels using Diastix (Bayer, Inc). All animals used in this study were not diabetic. In addition, only female mice were used.

Antibodies and Flow Cytometry

All antibodies, unless otherwise stated, were purchased from e-Bioscience. Single-cell suspensions were incubated with antibodies against CD16/32 unconjugated (93), CD3 FITC (145-2C11), CD3 BV421 (17A2; Biolegend), CD4 eFluor450 (RM4-5), CD4 PE (RM4-5), CD4 BV650 (GK1.5; Biolegend), CD8 α FITC (53-6.7), CD8 β PE-Cy7 (H35-17.2), CD19 eFluor450 (6D5), CD19 PE (6D5), CD19 BV421 (1D3; Biolegend), CD21/CD35 PE (4E3), CD23 PE-Cy7 (B3B4), IgM APC (II/41), IgD eFluor450 (11-26c), IgE BV650 (R35-72; BD-Biosciences), IgG PerCP-eFluor 710 (Polyclonal), IgA PE (11-44-2), biotinylated-insulin (ibtsystems), CD45 PerCPCy5 (30-F11), CD45 BV510 (30-F11; Biolegend); PD1 (29F.1A12;

Biolegend), ICOS APC (C398.4A), CD138 BV650 (281-2; Biolegend), CD11b FITC (M1/70),
 , CD11c PE (N418), B220 eFluor450 (RA3-6B2), BCL-6 PerCP-eFluor 710 (BCL-DWN),
 CXCR5 PE (SPRCL5), IL-21 PE (mhalx21) and Ki67 PE (B5; BD Biosciences). Intracellular
 labelling of Ki67 was performed using eBioscience kit following manufacturer's guidelines
 (catalogue number 00-5523-00). Cells were acquired using a BD LSR Fortessa X-20 (BD
 Biosciences) and data analysed using FlowJo software® (Tree Star). Doublets were excluded
 using forward light-scatter gating (FSC-A versus FSC-W) followed by gating on cells based
 on FSC-SSC. Dead cells were excluded by gating on LIVE/DEAD® Fixable Dead Cell
 Staining (ThermoFisher) negative cells. The gating strategies are described in the paper in the
 main figures and supplementary figures, explicit in the axis or described in detail in figures
 legends. The gates were defined using fluorescence minus one and isotype controls: Rat IgG2a
 eF450 (eBR2a), Rat IgG2a FITC (eBR2a), Rat IgG2a PE (eBR2a), Rat IgG2a PE-Cy7
 (eBR2a), Rat IgG2a APC (eBR2a), Rat IgG2a BV421 (RTK2758, Biolegend), Rat IgG2a
 BV650 (RTK2758, Biolegend), Rat IgG2a PerCP-eFluor 710 (eBR2a), Rat IgG2b PE (10H5),
 Armenian Hamster IgG APC (eBio299Arm), Rat IgG1 Biotin (eBRG1) and Rat IgG1 BV650
 (RTK2071; Biolegend).

Detection of thymic B cells bearing insulin-specific receptors.

The detection of B cells with receptors that bind insulin has been previously described [31].
 Briefly single cell suspensions isolated from the thymus were incubated overnight at 4°C in
 PBS supplemented with 1% foetal bovine serum, 1% anti-CD16/32 antibodies (eBiosciences)
 and biotinylated insulin (0.1 µg/10⁶ cells, ibtsystems). Bound insulin was detected with
 fluorochrome-labelled streptavidin Alexa6470(Invitrogen) for 30 minutes at 4°C. The cells
 were subsequently incubated with anti- CD19 PE (6D5; eBiosciences), B220 eFluor450 (RA3-
 6B2; eBiosciences), - CD4 BV650 (GK1.5; Biolegend), CD8β PE-Cy7 (H35-17.2;
 eBiosciences), -CD45 PerCPCy5.5 (30-F11; eBiosciences) antibodies and LIVE/DEAD

Fixable Dead Cell Stains (Thermo Fisher Scientific) for 30 min at 4°C, following which the cells were analysed by flow cytometry. B cell gates were defined following exclusion of dead cells and T cells (dump channel). All samples were stained with insulin-biotin followed by streptavidin or with streptavidin only, frequencies of B cells insulin⁺ were calculated subtracting the background calculated in sample-matched streptavidin only control.

Soluble tissue extracts and Enzyme Linked Immunosorbent Assay (ELISA)

Cell-free supernatants from thymic and splenic tissue were prepared as described [32]. Briefly, single cell suspension were centrifuged at 300g for 10 mins, 4 °C then 15 mins, 4 °C at 3000g. Cell-free supernatants were collected and stored at -20°C until analysis. IL-2 and IL-21 cytokines were detected using mouse IL-2 ELISA Ready-SET-Go and mouse IL-21 ELISA Ready-SET-Go ELISA kits following manufacturer guidelines (eBioscience). Isotype classification of immunoglobulins in thymic cell-free supernatants or serum was achieved using a rapid ELISA Mouse mAb isotyping kit (ThermoFisher) following manufacturer's guidelines.

Cultures

Bone marrow derived dendritic cells (BM-DCs) were prepared from the appropriate mice by standard methodology. Immature DCs were pulsed for 16 hours with whole insulin (Sigma; 5 µg/ml), LPS (Sigma; 10ng/ml), or pro-insulin peptide pB15-23 peptide (Thermo Fisher; p4878-1; 5 µg/ml). Thymocytes were prepared from mice described in the results and 1x10⁶ thymocytes were co-cultured in complete RPMI media (10% FCS, 50µmol/L β-mercaptoethanol, L-glutamine, 50 units/ml penicillin and streptomycin (Life-Sciences)) with 3x10⁴ BM-DCs only, or 3x10⁴ BM-DCs pulsed with insulin or 3x10⁴ BM-DCs pulsed with B15-23 peptide, or anti-CD3 (5µg/ml) (eBioscience) and anti-CD28 (2.5µg/ml) antibodies (eBioscience). The co-cultures were incubated at 37°C, 5% CO₂ for 72 hrs, following which cell proliferation was assessed by flow cytometry. The stimulation index was calculated

dividing the frequency of T cells in active proliferation (Ki67⁺) in cells following antigen stimulation by the frequency of T cells in active proliferation in paired non-stimulated culture (background).

For the detection of IL-21, single cell suspensions from the appropriate tissues were prepared placed in RPMI media (as above) supplemented with 50ng/ml PMA and 1µg/ml ionomycin for a total of 5 hours at 37°C, 5% CO₂. Brefeldin A (SIGMA) was added to the cultures at a concentration of 0.4mg/ml 1 hr after the initiation of the culture.

Immunofluorescence Analysis.

Thymi frozen in OCT compound were sectioned (~8 µm) on a cryostat. Sections were fixed in 4% paraformaldehyde or ice-cold acetone then blocked in PBS supplemented with 0.5% BSA. The sections were incubated with unconjugated primary antibodies rabbit-anti mouse IgG (Abcam), rabbit anti-mouse insulin (Abcam) or rabbit anti-mouse cytokeratin V (Abcam) overnight at 4 °C. Detection of bound antibody was achieved with goat anti-rabbit IgG-Alexa 647 or goat anti-rabbit Ig-Alexa488 (Invitrogen) or goat anti-rat IgG Alexa 488 (Invitrogen). Anti-B220 directly conjugated with Alexa 647 was incubated for 45 minutes at room temperature. For detection of apoptosis, following incubation with the secondary antibody an *in situ* apoptosis kit was used (Click-iT™ Plus TUNEL Assay, Alexa Fluor™ 647 dye; Thermofisher) according with the manufacturer instructions. Sections were counterstained with DAPI (Molecular Probes) and mounted in Prolong Gold anti-fade or Prolong Diamond (Invitrogen). Confocal microscopy was undertaken using Zen software on a Zeiss LSM 710 fitted on an Axioimager using a 63x (1.4) planApochromat or 20x (0.6) Neofluor. Binding of autoreactive immunoglobulin and TUNEL in microscopy images was quantified using StrataQuest V64 software. Individual nuclei were counted and the data was presented as

scatterplots of mean fluorescence intensity of DAPI versus mean fluorescence intensity of Ig or TUNEL positive cells.

RNA Isolation and real-time RT-PCR analysis

Thymic tissues were stored at -80°C in RLT. Samples were allowed to thaw and RNA were carried out using the RNeasy mini kits (Qiagen, Manchester, UK), according to the manufacturer's instructions. On-column DNase digestion was carried out to remove any contaminating genomic DNA using the RNase-free DNase set (Qiagen, Manchester, UK) according to the manufacturer's instructions. The cDNA syntheses were performed with the Superscript II reverse transcriptase system (Invitrogen), according to manufacture's instructions. The qRT-PCR of *aicda* mRNA expression (*AID* gene) in total thymus was performed with the Taqman qPCR Kit (Applied Biosystems, Warrington, UK)). mRNA expression levels were normalized to *HPRT1* housekeeping gene using $\Delta\Delta C_t$ calculations. Mean relative mRNA expression levels between control and experimental groups were calculated using the $2^{-\Delta\Delta C_t}$ calculations.

Statistical analysis

Statistical analyses were performed by parametric or non-parametric tests, selected based on the distribution of the raw data. The comparisons between experimental groups were performed using student unpaired t-test, Mann-Whitney and one-way ANOVA as appropriate. The statistical analyses for fold-changes were performed using Wilcoxon signed-rank test. All analyses were conducted using GraphPad InStat (version 5) software (GraphPad).

Results

T1D progression correlates with increased intrathymic B cell numbers in NOD mice

Thymic B cells normally constitute a small population of cells within the murine and human thymus in normal individuals. Abnormality in thymic B cell numbers has been linked to certain autoimmune conditions [14; 15]. To determine whether thymic B cell populations differ between diabetes-prone or non-prone mice, we performed time-course flow cytometric studies of age-matched, sex-match NOD and control C57BL/6 (B6) mice.

Diabetes incidence in our female NOD mouse colony is 95%, approximately 3% of mice develop T1D at 12-14 weeks of age, 85% at 18-20 weeks of age with the remaining 7% of females progressing to T1D by 23 weeks of age. Animals not diabetic by 23 weeks of age rarely develop T1D. The data is based on a cohort of 200 animals (Supplementary Figure 1a). This diabetes incidence, combined with the insulinitis score- that is the degree of immune cell infiltration of islets and degree of β cell destruction- as mice age (Supplementary Figure 1b) highlight that 12-14 weeks of age in our colony represents late insulinitic-preultimate diabetic stage, a critical time when immunoregulation of the autoreactive response starts to breakdown. Thus, in our initial studies we focused on two major time points; the pre-early insulinitic phase (4-6 weeks) and the post-insulinitic/pre-diabetic phase (12-14 weeks) to assess the presence of CD19⁺ thymic B cells. Representative flow cytometry plots for the respective mice are shown in Figure 1a. Although absolute numbers of CD19⁺ B cells remained relatively static in the thymi of control B6 mice at the time points investigated (Figure 1b), with perhaps a slight increase at 12-14 weeks of age, the absolute numbers of CD19⁺ B cells increased significantly in the later age group of NOD mice in comparison to numbers seen either at 4-6 week old NOD mice or 12-14 week old B6 mice. Importantly, the number of thymic B cells at 4-6 weeks of age was comparable between NOD and control B6 mice.

This increased number of thymic B cells in 12-14 week old NOD mice was not related to increased B cell development in the bone marrow, as frequencies of CD19⁺ B cells in this primary lymphoid tissue was comparable between the two strains of mice at both time points investigated (data not shown). These data show that inappropriate accumulation of thymic B cells precedes the overt β cell destruction phase of T1D.

Intrathymic signals trigger enhanced B cell development in NOD mice.

Although previous studies have documented the ability of the thymic environment to enable B cell development in non-autoimmune-prone mice, other reports suggest thymic B cells accumulate via periphery B cell migration to the thymus [16; 33]. To determine whether the NOD mouse thymus promotes B cell development we used recombination activation gene green fluorescent protein (RAG2p-GFP) reporter mice on a non-T1D-prone FVB background (hereafter called FVB-RAG-GFP), or on the NOD background (hereafter called NOD-RAG-GFP). In RAG2p-GFP reporter mice, highest GFP expression occurs when RAG genes are active [30]. Once recombination of the B cell receptors and T cell receptors is complete and RAG activity is silenced, GFP expression decreases over a 54 hour period [30]. As such, newly developed B cells can be identified from thymic resident/recirculatory B cells based on the expression of GFP.

Since our control RAG2p-GFP transgenic mice are on a FVB background, we compared thymic B cell frequencies and numbers of this alternative control murine strain to control B6 mice or NOD mice. Although frequencies and absolute numbers of thymic B cells in the FVB strain were higher than the B6 strain, the NOD strain demonstrated significantly greater thymic B cell frequencies and numbers to the FVB strain (Supplementary Figure 2a-b).

We performed time-course, flow cytometry studies of the two strains of mice at the ages shown in Figure 1c, and quantified the number of GFP^{hi} B cells. Representative flow cytometry plots showing the gating strategy for CD19⁺GFP^{hi} B cells is shown in Supplementary Figure 1c. Recently developed CD19⁺GFP^{hi} B cells were readily detectable in both strains of mice at all time points analysed (Figure 1c). In control FVB-GFP mice, there was no significant changes in B cell development as mice aged. In the NOD strain, although there was no significant change in B cell development when the two age groups were compared, it was clear that thymic B cell development is enhanced as mice enter the late insulinitic-prediabetic phase of the T1D pathway.

In light of evidence that the late insulinitic-prediabetic phase is characterised by increased B cell development, we asked if homeostatic proliferation of thymic B cells is also affected as mice enter the late insulinitic-prediabetic phase. We performed comparative flow cytometric studies between NOD and control B6 mice, assessing for Ki67 expression as a marker for homeostatic proliferation. Interestingly for both strains of mice, the highest level of homeostatic proliferation of thymic B cells is an early event, with CD19⁺Ki67⁺ B cell frequencies higher in younger mice when compared to older mice (Figure 1d). Further, this decrease in homeostatic proliferation in the 12-14 week old group was more pronounced in NOD mice, although the decrease was not significant.

The NOD thymic environment has ectopic germinal centre formation potentiality.

To further investigate the phenotype of thymic B cells in NOD mice we assessed their surface markers. B cells undergo a series of transitions from the immature stage developing follicular or marginal zone properties. Thus, we qualified the phenotype of thymic B cells assessing for follicular (IgM^{lo}IgD⁺CD21/35⁺,CD23⁺) versus marginal zone (IgM⁺IgD^{lo}CD23⁻CD21/35⁺).

285 We focused our studies on 11-14 week old mice due to the evidence that at this age B cell
286 development is enhanced as are thymic B cell numbers in NOD mice when compared to
287 control mice. Representative flow cytometric plots for our gating strategies are shown in
288 Supplementary Figure 2d.

289 As shown in Figure 2, the frequency of follicular B cells within the thymic B cell pool was
290 significantly higher in NOD mice compared to control B6 mice (Figure 2a). This
291 enhancement in follicular B cells in the NOD mouse thymus was recapitulated when absolute
292 number of follicular B cells was calculated (Figure 2b). In contrast, although the frequency of
293 B cells with a marginal zone phenotype were significantly decreased in the NOD mouse
294 thymus compared to control B6 mouse thymus, the absolute numbers of these cells was
295 similar between the two strains of mice.

296 The increased numbers of FO B cells in NOD mice with respect to B6 control mice led us to
297 investigate whether the thymic B cell form follicle-like structures. Immunohistochemical
298 studies revealed B cell follicle-like structures form only in NOD mice (Figure 2c). Initially B
299 cells are detectable at the cortical-medullary junction at 9 weeks of age in NOD mice (not
300 shown) with pronounced accumulation of B cells into follicle-like structures in this location
301 by 11 weeks of age. The presence and location of B cell follicle-like structures was identical
302 irrespective of whether we used anti-B220 or anti-CD19 antibodies to identify B cells (not
303 shown) confirming the accumulating B220⁺ cells are not plasmacytoid DCs. We quantified
304 the number of B cell follicle-like structures in the thymus of 9-11 week old NOD mice; of 15
305 individual sections assessed, 90% contained 1 follicle, 5% two follicles and 5% no follicles.

306 The presence of follicle-like structures in the thymus of late insulitic-pre-diabetic NOD mice,
307 but not control B6 mice, lead us to ask if the thymic environment could support germinal
308 centre formation. Of interest was the relationship between IL-2 and IL-21, the latter being a

key mediator of germinal centre formation; the cytokine promotes B cell somatic hypermutation and class switching, and the development and maintenance of T follicular helper cells (T_{fh} cells) [34]. In NOD mice IL-21 has been associated with T1D progression [35; 36] and CD4⁺CD45R⁻ T cells isolated from T1D patients secrete greater quantities of IL-21 than quantified from normal individuals [37]. We prepared cell-free supernatants [32] from the thymi of NOD and control B6 mice at the ages shown in Figure 3a and performed ELISA assays. As a comparison, we analysed cell-supernatants from the spleens of the same mice. The results were tabulated as ratio of IL-21:IL-2. No differences were seen in IL-21:IL-2 ratios in splenic preparations from the two strains of mice. However, the NOD mouse thymus had a significant bias in IL-21 concentrations in comparison to B6 mice.

In light of this IL-21 bias, we quantified the frequency and absolute numbers of CD4SP cells that expressed a T_{fh} cell phenotype in the thymus of the two strains of mice. As shown in Figure 3b-c, NOD mice exhibited a significant increase in frequencies and absolute numbers of CD4SPPD1^{hi}ICOS⁺ T cells, and these cells also expressed transcription factor Bcl-6 (Figure 3d) and CxCR5 (Supplementary Figure 3a). In addition, approximately 5% of NOD putative thymic T_{fh} cells secreted IL-21, a frequency that was comparable to that seen for splenic T_{fh} cells from the same mice (Supplementary Figure 3b). Furthermore, this increase in thymic T_{fh} cells in NOD mice in comparison to B6 control mice correlated with an increased number of CD4⁻CD8⁻B220^{low/-}CD138⁺ plasma cells in the NOD mouse thymus, although this increased number was not significant (Figure 3e, Supplementary Figure 3c). Together, these data suggested that ectopic germinal centres could be present in the NOD mouse thymus, but absent in control B6 mouse thymus. To support this hypothesis we looked for a bone-fide germinal centre marker; the enzyme activation-induced cytidine deaminase (AID). RNA was prepared from thymi isolated from NOD mice or control B6 mice and quantitative real-time RT-PCR performed. As an additional control we included thymic

mRNA isolated from age-matched, sex-matched NOD- μ MT^{-/-} mice. The relative expression of transcripts for AID in NOD mice was normalised to control B6 mice. As shown in Figure 3f, the NOD mouse thymus has enhanced AID expression in comparison to control B6 mice. Thus, ectopic germinal centre formation is likely a feature of the NOD thymus and precedes the preultimate β cell destruction phase of T1D.

Thymic immunoglobulins binding selective mTECs correlates with mTEC apoptosis.

The presence of AID and enhanced plasma cell frequencies in the NOD thymus with respect to control B6 mice, made us query the immunoglobulin isotype of the thymic B cells and secreted antibodies. Since we previously had investigated the IgM⁺ B cell thymic subtype (Figure 2), this time we focused on class-switched IgM⁻ cells. The number of IgM⁻IgD⁻IgA⁺ and IgM⁻IgD⁻IgG⁺ B cells were similar in the thymus of both NOD and control B6 mice as determined by flow cytometry (Figure 4a). In contrast, the number of IgM⁻IgD⁻IgE⁺ B cells were significantly increased in the NOD mouse thymus with respect to control mice. Interesting, a unique population of IgM⁻IgD⁺ B cells (similar to those reported in T1D patients [31]) was detectable in the thymic tissue. These IgM⁻IgD⁺ B cells dually expressed IgG, IgE or IgA with the number of dual expressing IgD⁺IgA⁺ and IgD⁺IgG⁺ B cells being significantly higher in the NOD mouse thymus than the B6 control mouse thymus, the most significant being the IgD⁺IgG⁺ isotype (Figure 4b, Supplementary Figure 4a). In contrast, no differences in IgD⁺IgE⁺ B cell numbers were seen between the two strains of mice. We next assessed the isotype of soluble thymic immunoglobulin by ELISA, in comparison to serum immunoglobulin. Only the IgG1 and IgA immunoglobulin isotypes were enhanced in the thymus of NOD mice in comparison to control B6 mice (Figures 4c,e). In contrast, IgG2a, IgG2b and IgM antibody levels were similar in both strains of mice, with IgG3 antibody

358 levels being slightly lower in NOD mice than in the thymus of B6 control mice. Interestingly,
359 in NOD mice thymus, B cells predominately used the kappa light chain, there being a
360 significant decrease in the presence of lambda light chains when compared to B6 control
361 mice (Supplementary Figure 4c). In addition, this isotype pattern documented in the NOD
362 mouse thymus seemed unique for this tissue, as similar ELISA-based isotyping of
363 immunoglobulins in the serum of the two strains of mice revealed little difference in levels of
364 each isotype assessed (Figure 4d, f). However, similar to the thymus, in the serum there was a
365 significant decrease in lambda light chain usage in NOD mice in comparison to B6 controls
366 (Supplementary Figure 4b). Quantification of the thymic Ig isotypes supported the data that
367 IgG1 and IgA are significantly greater in the thymus of NOD mice with respect to control B6
368 mice (Supplementary Figure 5a).

369 We decided to explore further these thymic B cells to determine whether they harboured
370 receptors specific for islet autoantigens, focusing on their specificity for insulin [31].
371 Representative gating strategy for identifying insulin-reactive B cells is shown in
372 Supplementary Figure 5b. Although the frequency of cells bearing receptors specific for
373 insulin is significantly less in the thymus of NOD mice with respect to control B6 mice
374 within the B cell fraction, absolute numbers of insulin reactive B cells was similar in NOD
375 mice and B6 control mice (Supplementary Figure 5c). Thus, insulin-reactive B cell numbers
376 do not correlate with T1D susceptibility at this time point. Due to this finding, we decided to
377 ask whether thymic B cells produce antibodies that target, as yet, undefined antigens on
378 thymic stroma. Thymic tissue sections from 11 week old NOD mice were incubated with
379 anti-mouse antibodies that would detect any mouse immunoglobulin bound to thymic stroma
380 *in situ* and bound antibodies were detected by confocal microscopy. To qualify whether any
381 immunoglobulins that bound to thymic stroma interacted with mTECs, we included mTEC-
382 binding anti-cytokeratin V antibodies in the assay. As shown in Figure 5a, there was

detectable binding of murine immunoglobulins to thymic stroma, suggesting these cells had murine immunoglobulins bound to them *in situ*. Interestingly, intrathymic immunoglobulins were bound almost exclusively to cytokeratin V⁺ mTECs and it appeared that only a proportion of mTECs were being targeted by the immunoglobulins. In contrast to NOD mice, there was substantially less intrathymic immunoglobulin in control B6 mice interacted with thymic stroma, particularly cytokeratin V⁺ mTECs (Figure 5b). Further, there was no evidence of intrathymic immunoglobulins bound to thymic stroma, including cytokeratin V⁺ mTECs in B cell deficient NOD- μ MT^{-/-} mice confirming the specificity of the anti-mouse antibodies for mouse immunoglobulins (Supplementary Figure 6). We quantified the frequency of cells with murine immunoglobulin bound in the thymus of 11-14 week old NOD and B6 mice. We selected images that had comparative frequencies of cytokeratin V⁺ mTECs and counted $2-3 \times 10^4$ DAPI⁺ cells/mm². As shown in Figure 5c, approximately 7% of cells had bound murine immunoglobulin (Ig) in the NOD mouse thymus. In contrast, the frequency of cells bound by murine immunoglobulins in B6 mice was so low as to be undetectable.

Finally we queried the significance of *in situ* binding of thymic stroma by immunoglobulins, particularly the potential that a selective number of mTECs underwent apoptosis. In this regard, we incubated thymic tissue sections from 11 week old, female NOD mice with antibodies specific to cytokeratin V⁺ and assessed for apoptosis by confocal microscopy following TUNEL staining (Figure 6a). As controls, we similarly analysed thymic tissue sections from control B6 mice and NOD- μ MT^{-/-} mice. The inclusion of NOD- μ MT^{-/-} mice was important to determine whether the diabetes-associated MHC class II molecules unique to NOD mice was sufficient to trigger mTEC apoptosis via non-B cell-mediated mechanisms. Thymic tissue sections from NOD mice had clear evidence of apoptosis, and such apoptotic cells were almost exclusively cytokeratin V⁺ mTECs. Apoptosis of cytokeratin V⁺ mTECs

was also evident in NOD- μ MT^{-/-} mice, although the proportion of apoptotic cells seemed lower than that for B cell sufficient NOD mice. In contrast to the NOD strains, we could not see any apoptotic cells in the B6 control mouse thymic tissue section. We quantified the frequency of apoptotic cells in the thymic sections of the respective strains of mice (Figure 6). We counted a total of 4×10^4 DAPI⁺ cells/mm² per section, ascertaining similar frequencies of cytokeratin V⁺ mTECs for each tissue sections examined. As shown in Figure 6b, ~6% of DAPI⁺ cells were apoptotic in NOD mice. This frequency of apoptosis was two-fold higher than seen for NOD- μ MT^{-/-} mice (~3%). In contrast to the NOD strains, <1% of cells were apoptotic in control B6 mice. We were curious to determine if apoptotic cytokeratin V⁺ mTECs in NOD mice expressed insulin. Thymic tissue sections from 11 week old female NOD mice were incubated with anti-cytokeratin V⁺ and anti-insulin antibodies and apoptosis determined by TUNEL staining as before. As a control, we similarly analysed thymic tissue sections from age-matched, female B6 mice. Interestingly, within the apoptotic cytokeratin V⁺ mTEC pool in NOD mice resided cytokeratin V⁺ mTECs that expressed insulin, although it is important to note that some insulin⁺ cytokeratin V⁺ mTECs were not apoptotic suggesting there is not a complete loss of insulin⁺ cytokeratin V⁺ mTECs but a reduction in their numbers. Similarly, some apoptotic mTECs did not express insulin (Supplementary Figure 7).

Taken together, these data suggest that B cell-mediated autoimmune targeting of cytokeratin V⁺ mTECs results in the loss of a distinct population of cytokeratin V⁺ mTECs, some of which express insulin, and this key feature occurs before sustained autoimmune attack in the pancreas.

Thymic B cells enhance premature egress of T cells from the thymus.

The evidence that thymic stroma had bound autoantibodies and the presence of these autoantibodies correlated with increased apoptosis of thymic stroma, including some insulin⁺ mTECs, we investigated the impact this may have on thymocytes capable of responding to islet antigen, particularly insulin. We isolated the thymocytes from NOD mice thymi and cultured the cells in the presence of bone-marrow derived dendritic cells and either whole insulin or proinsulin peptide15:23 [38]. The proliferative response to the CD4SP and CD8SP thymocytes to the respective stimulants was assessed by flow cytometric analysis of Ki67 (Figure 7, Supplementary Figure 8a). As controls we included B6 mice stimulated with whole insulin, and B cell-deficient NOD- μ MT^{-/-} mice stimulated with whole insulin or proinsulin peptide 15:23. For CD4SP cells only those isolated from B cell sufficient NOD mice responded to both whole insulin, although the response was not significant in comparison to control mice. (Figure 7a). The responses to whole insulin for thymocytes from B6 and NOD- μ MT^{-/-} mice being close to baseline. In contrast, CD4SP thymocytes from NOD mice exhibited a significantly increased response to proinsulin P15:23 with respect to NOD- μ MT^{-/-} mice. The responses of CD8SP thymocytes was slightly different; whereas thymocytes isolated from NOD mice responded to the whole insulin molecule, the responses for individual mice was quite diverse- some responded well, others' response close to baseline levels for B6 control mice (Figure 7b). Similarly, CD8SP thymocytes from NOD- μ MT^{-/-} mice had some diversity in responsiveness to whole insulin, although it was noted that even the best responders still responded weaker than that seen for NOD mice. In contrast, CD8SP thymocytes from NOD mice responded far better to proinsulin P15:23, than those from NOD- μ MT^{-/-} mice, although the response was not significantly enhanced. In these same mice the responses to the proinsulin peptide were less diverse and above baseline levels.

456 We initially wondered whether this increased response for insulin and proinsulin peptide by
457 NOD thymocytes was representative of increased survival of autoreactive T cells, and thus a
458 breakdown in negative selection. In particular, we queried whether thymocytes that had
459 very recently rearranged their TcR escaped from the thymus before completing negative
460 selection. If this held true, we expected an increase in RAG-GFP^{hi} T cells in the blood; RAG-
461 GFP levels normally fall during negative selection due to the time to complete the process
462 and as such, peripheral T cells are usually RAG-GFP^{int} [30]. To test this hypothesis we
463 performed flow cytometry analysis of total GFP levels of T cells in the peripheral blood of
464 NOD-RAG-GFP mice in comparison to control FVB-RAG-GFP mice and B cell deficient
465 NOD- μ MT^{-/-}-RAG-GFP mice. Representative flow cytometry plots showing the gating
466 strategy is shown in Supplementary Figure 8b. As shown in Figure 7c, in the NOD murine
467 strains the frequency of total GFP⁺ T cells in peripheral blood was greater than seen for the
468 control FVB strain, for NOD mice this increase being significant. Furthermore, this increased
469 frequency of GFP⁺ T cells in the peripheral blood of the NOD strains was almost entirely due
470 to GFP^{hi} cells, as GFP^{int} cells were only slightly increased in frequency in comparison to
471 control FVB-RAG-GFP mice, again NOD mice showing a significant increase. Importantly,
472 although not significant, it was clear that the frequency of RAG-GFP^{hi} cells in B cell
473 sufficient NOD-RAG-GFP mice was higher than in B cell deficient NOD- μ MT^{-/-}-RAG-GFP
474 highlighting the importance of B cells in the early release of T cells from the thymus prior to
475 negative selection.

Discussion

Ablation of efficient purging of autoreactive T cells in the thymus and the role of B cells in T1D seem two distinct entities in understanding how immunological tolerance is broken in this chronic autoimmune condition. Here, we establish that inappropriate accumulation of B cells in the NOD mouse thymus is a unique feature of the disease process, and these thymic B cells may play a role in the egress of pre-negatively selected T cells.

T1D progression in both man and NOD mice occurs over time. The initial stages of T1D, where priming of the immune response to islet antigen occurs but not overt β cell destruction, is characterised by autoantibodies to β antigens [39]. It is accepted that following priming of the autoreactive T cell repertoire to β cell antigens, the activity of the autoreactive T cells is kept in check by regulatory mechanisms. Ultimately, such regulation fails, and leading to β cell destruction. Little is known as to why regulation of autoreactive T cells fails over time, although paucity of, or dysfunction of, T regulatory cells is speculated to contribute to the phenomenon [40; 41; 42]. Our data adds a new dimension to our understanding of the immunological changes that occur at the late insulitic- pre-diabetic phase that may tip the autoreactive T cell response in favour of β cell destruction; targeted thymic B cell autoimmune attack of thymic stroma expressing β cell antigens.

B cells are present in the thymus of mammals from fetal age to adulthood, their numbers remaining relatively static in ontogeny and equating to those of thymic dendritic cells [5; 11; 13]. Previous studies in NOD mouse strains documented B cell accumulation in the thymus of aged mice [43; 44]. Here we extended on these early studies showing that in NOD mice, thymic B cell numbers are not static, their numbers significantly increase at the late insulitic- pre-diabetic phase suggesting the restricted B cell niche normally present expands. This change in B cell numbers occurs at the same time as increased numbers of RAG⁺ B cells are

detected in the thymus, but decreased homeostatic proliferation. Together these findings suggest that permissiveness of B cell development that can normally occur within the thymus [33; 45; 46; 47] is enhanced in NOD mice as they age, and the increase in B cell numbers potentially reflects this increased rate of development rather than *in situ* proliferation. Although we cannot exclusively discount that peripheral B cells migrating to the thymus contribute to the thymic B cell population, we, like others, have found peripheral B cells have little propensity to traffic to the thymus (data not shown, [47]). Future studies in how the NOD mouse thymic environment potentially nurtures B cell development and retention will be informative.

The phenotype of thymic B cells in NOD mice resembles that of thymic B cells in non-autoimmune strains of mice; they predominantly express B2 follicular cell markers, and have a predominantly activated phenotype with high MHC and costimulatory molecule expression (not shown, [45; 48]). The location of thymic B cells in NOD mice is also reminiscent of reports in other murine strains- positioned predominantly at the cortico-medullary junction- but in contrast to non-autoimmune prone mice, large B cell follicles form and this is age-dependent. Furthermore, the hallmarks of germinal centres are readily detectable in the NOD thymus; IL-21 and T follicular helper (T_{fh}) cells. Abnormalities in levels of IL-21 and T_{fh} cells in peripheral tissues, and blood, have been strongly associated with T1D [37; 49]. Here, we show similar abnormalities exist in the thymus occurring specifically at the late insulitic-pre-diabetic phase of the T1D condition. In addition the thymus of NOD mice has enhanced levels of AID mRNA transcripts, suggesting increased *in situ* somatic hypermutation and class switching of the B cell repertoire activity. Plasma cells are also increased in the thymus of NOD mice with respect to control animals which taken all this information together implies ectopic germinal centres are a feature of the NOD thymus, not just their pancreas [50]. Our evidence that the NOD mouse thymus is populated with significantly increased

525 numbers of B cells with IgG, IgA and IgE receptors with respect to non-autoimmune prone
526 mice, as well as enhanced levels of soluble IgG1 and IgA antibodies supports our rationale of
527 ectopic germinal centre formation in this primary lymphoid tissue.

528 The significance of these unique changes in the NOD mouse thymus as mice progress along
529 the T1D pathway, we believe, is that they have the potential to impact on the capacity of
530 negative selection of autoreactive T cells to occur effectively. The importance of mTEC
531 expression of TSAs for efficient deletion of developing T cells bearing autoreactive T cell
532 receptors is well established [51]. Our evidence that a selective population of mTECs have
533 autoantibodies bound *in situ*, and in the presence of thymic B cells a proportion of mTECs
534 undergo apoptosis, a number of which express insulin, is likely to have implications on
535 negative selection of islet-reactive T cells. The antigenic specificity of the intrathymic
536 autoantibodies target is unknown, and we do not believe that they must recognise insulin to
537 impact of T1D progression. It is possible that the intrathymic autoantibodies recognise and
538 promote apoptosis of particular mTECs that express certain TSAs that are associated with
539 other autoimmune conditions NOD mice develop [52; 53; 54]. It follows that reduction in
540 insulin expressing mTECs may happen inadvertently. Alternatively, or in addition, it is
541 possible that thymic cognate B-T cell interactions promote survival of developing
542 autoreactive T cells as opposed to their deletion [55].

543 Our data is supportive of the rationale that pre-negatively selected T cells are potentially
544 released from the thymus prematurely. Two photon microscopy has documented that
545 developing T cells reside in medulla for 3-5 days to complete negative selection [56]. In
546 RAG2p-GFP reporter mice, this duration in the medulla equates to decreased GFP intensity
547 due to the 54 hour half-life of the molecule [30]. Our evidence that in NOD mice CD3⁺GFP^{hi}
548 cells are significantly enhanced in peripheral blood with respect to non-autoimmune prone
549 mice suggests an aborted time, or failed entry into, the medulla of GFP^{hi} T cells, and as a

550 consequence failed negative selection. It follows that the increased export of non-negatively
551 selected T cells could overpower waning regulatory mechanisms in the islets leading to the
552 final sustained attack of the β cells.

553 The fledgling field of thymic B cell research is starting to unravel the importance of this
554 unique population of cells in the immune system. Our data highlight a new relationship
555 between thymic B cells and type 1 diabetes development. Future studies that define the *in situ*
556 developmental pathway and receptor specificity of thymic B cells will be important for
557 identifying key therapeutic strategies for type 1 diabetes and other autoimmune conditions in
558 which thymic B cells make a contribution.

559 **Acknowledgements**

560 We would like to thank staff of the Biology Technical Facility and Biology Support Services
561 for their technical support. We also thank Dr. Lena Petersone for advice on IL-21 detection by
562 flow cytometry approaches. This work was supported by the Diabetes UK and the Jean Shanks
563 Foundation.

564 **Authors contributions:**

565 AP, JS, MK, KH and EAG performed the experiments and analysed data; AP and EAG wrote
566 the paper.

567 **Conflict of Interest Statement**

568 All authors declare there was no conflict of interest.

References

- [1] J.W. Kappler, N. Roehm, and P. Marrack, T cell tolerance by clonal elimination in the thymus. *Cell* 49 (1987) 273-80.
- [2] A.J. Bonito, C. Aloman, M.I. Fiel, N.M. Danzl, S. Cha, E.G. Weinstein, S. Jeong, Y. Choi, M.C. Walsh, and K. Alexandropoulos, Medullary thymic epithelial cell depletion leads to autoimmune hepatitis. *Journal of Clinical Investigation* 123 (2013) 3510-3524.
- [3] B. Kyewski, J. Derbinski, J. Gotter, and L. Klein, Promiscuous gene expression and central T-cell tolerance: more than meets the eye. *Trends in Immunology* 23 (2002) 364-371.
- [4] M.S. Anderson, E.S. Venanzi, L. Klein, Z. Chen, S.P. Berzins, S.J. Turley, H. von Boehmer, R. Bronson, A. Dierich, C. Benoist, and D. Mathis, Projection of an immunological self shadow within the thymus by the aire protein. *Science* 298 (2002) 1395-401.
- [5] L. Klein, B. Kyewski, P.M. Allen, and K.A. Hogquist, Positive and negative selection of the T cell repertoire: what thymocytes see (and don't see). *Nat Rev Immunol* 14 (2014) 377-91.
- [6] J. Oh, and J.S. Shin, The Role of Dendritic Cells in Central Tolerance. *Immune Netw* 15 (2015) 111-20.
- [7] F. Frommer, and A. Waisman, B Cells Participate in Thymic Negative Selection of Murine Auto-reactive CD4(+) T Cells. *Plos One* 5 (2010).
- [8] T. Yamano, J. Nadjic, M. Hinterberger, M. Steinert, S. Koser, S. Pinto, N. Gerdes, E. Lutgens, N. Ishimaru, M. Busslinger, B. Brors, B. Kyewski, and L. Klein, Thymic B Cells Are Licensed to Present Self Antigens for Central T Cell Tolerance Induction. *Immunity* 42 (2015) 1048-1061.
- [9] P. Kleindienst, I. Chretien, T. Winkler, and T. Brocker, Functional comparison of thymic B cells and dendritic cells in vivo. *Blood* 95 (2000) 2610-6.
- [10] K. Nango, M. Inaba, K. Inaba, Y. Adachi, S. Than, T. Ishida, T. Kumamoto, M. Uyama, and S. Ikehara, ONTOGENY OF THYMIC B-CELLS IN NORMAL MICE. *Cellular Immunology* 133 (1991) 109-115.
- [11] P.G. Isaacson, A.J. Norton, and B.J. Addis, The human thymus contains a novel population of B lymphocytes. *Lancet* 2 (1987) 1488-91.
- [12] J. Spencer, M. Choy, T. Hussell, L. Papadaki, J.P. Kington, and P.G. Isaacson, PROPERTIES OF HUMAN THYMIC B-CELLS. *Immunology* 75 (1992) 596-600.

- [13] J.L. Andreu-Sanchez, J. Faro, J.M. Alonso, C.J. Paige, C. Martinez, and M.A. Marcos, Ontogenic characterization of thymic B lymphocytes. Analysis in different mouse strains. *Eur J Immunol* 20 (1990) 1767-73.
- [14] B. Christensson, P. Biberfeld, and G. Matell, B-cell compartment in the thymus of patients with myasthenia gravis and control subjects. *Ann N Y Acad Sci* 540 (1988) 293-7.
- [15] I.R. Mackay, M. Masel, and F.M. Burnet, Thymic Abnormality in Systemic Lupus Erythematosus. *Australas Ann Med* 13 (1964) 5-14.
- [16] T. Sato, S. Ishikawa, K. Akadegawa, T. Ito, H. Yurino, M. Kitabatake, H. Yoneyama, and K. Matsushima, Aberrant B1 cell migration into the thymus results in activation of CD4 T cells through its potent antigen-presenting activity in the development of murine lupus. *Eur J Immunol* 34 (2004) 3346-58.
- [17] D.L. Kaufman, M. Clare-Salzler, J. Tian, T. Forsthuber, G.S. Ting, P. Robinson, M.A. Atkinson, E.E. Sercarz, A.J. Tobin, and P.V. Lehmann, Spontaneous loss of T-cell tolerance to glutamic acid decarboxylase in murine insulin-dependent diabetes. *Nature* 366 (1993) 69-72.
- [18] R. Tisch, X.D. Yang, S.M. Singer, R.S. Liblau, L. Fugger, and H.O. McDevitt, Immune response to glutamic acid decarboxylase correlates with insulitis in non-obese diabetic mice. *Nature* 366 (1993) 72-5.
- [19] S.M. Lieberman, A.M. Evans, B. Han, T. Takaki, Y. Vinnitskaya, J.A. Caldwell, D.V. Serreze, J. Shabanowitz, D.F. Hunt, S.G. Nathenson, P. Santamaria, and T.P. DiLorenzo, Identification of the beta cell antigen targeted by a prevalent population of pathogenic CD8+ T cells in autoimmune diabetes. *Proc Natl Acad Sci U S A* 100 (2003) 8384-8.
- [20] Q. He, Y.M. Morillon, 2nd, N.A. Spidale, C.J. Kroger, B. Liu, R.B. Sartor, B. Wang, and R. Tisch, Thymic development of autoreactive T cells in NOD mice is regulated in an age-dependent manner. *J Immunol* 191 (2013) 5858-66.
- [21] K. Kachapati, D. Adams, K. Bednar, and W.M. Ridgway, The non-obese diabetic (NOD) mouse as a model of human type 1 diabetes. *Methods Mol Biol* 933 (2012) 3-16.
- [22] P. Leete, A. Willcox, L. Krogvold, K. Dahl-Jorgensen, A.K. Foulis, S.J. Richardson, and N.G. Morgan, Differential Insulitic Profiles Determine the Extent of beta-Cell Destruction and the Age at Onset of Type 1 Diabetes. *Diabetes* 65 (2016) 1362-9.
- [23] V. Parikka, K. Nanto-Salonen, M. Saarinen, T. Simell, J. Ilonen, H. Hyoty, R. Veijola, M. Knip, and O. Simell, Early seroconversion and rapidly increasing autoantibody

- concentrations predict prepubertal manifestation of type 1 diabetes in children at genetic risk. *Diabetologia* 55 (2012) 1926-36.
- [24] H. Noorchashm, Y.K. Lieu, N. Noorchashm, S.Y. Rostami, S.A. Greeley, A. Schlachterman, H.K. Song, L.E. Noto, A.M. Jevnikar, C.F. Barker, and A. Naji, I-Ag7-mediated antigen presentation by B lymphocytes is critical in overcoming a checkpoint in T cell tolerance to islet beta cells of nonobese diabetic mice. *J Immunol* 163 (1999) 743-50.
- [25] D.V. Serreze, H.D. Chapman, D.S. Varnum, M.S. Hanson, P.C. Reifsnyder, S.D. Richard, S.A. Fleming, E.H. Leiter, and L.D. Shultz, B lymphocytes are essential for the initiation of T cell-mediated autoimmune diabetes: analysis of a new "speed congenic" stock of NOD.Ig mu null mice. *J Exp Med* 184 (1996) 2049-53.
- [26] C.Y. Hu, D. Rodriguez-Pinto, W. Du, A. Ahuja, O. Henegariu, F.S. Wong, M.J. Shlomchik, and L. Wen, Treatment with CD20-specific antibody prevents and reverses autoimmune diabetes in mice. *J Clin Invest* 117 (2007) 3857-67.
- [27] M.D. Pescovitz, C.J. Greenbaum, H. Krause-Steinrauf, D.J. Becker, S.E. Gitelman, R. Goland, P.A. Gottlieb, J.B. Marks, P.F. McGee, A.M. Moran, P. Raskin, H. Rodriguez, D.A. Schatz, D. Wherrett, D.M. Wilson, J.M. Lachin, J.S. Skyler, and C.D.S.G. Type 1 Diabetes TrialNet Anti, Rituximab, B-lymphocyte depletion, and preservation of beta-cell function. *N Engl J Med* 361 (2009) 2143-52.
- [28] D.V. Serreze, S.A. Fleming, H.D. Chapman, S.D. Richard, E.H. Leiter, and R.M. Tisch, B lymphocytes are critical antigen-presenting cells for the initiation of T cell-mediated autoimmune diabetes in nonobese diabetic mice. *Journal of Immunology* 161 (1998) 3912-3918.
- [29] G.M. Brodie, M. Wallberg, P. Santamaria, F.S. Wong, and E.A. Green, B-cells promote intra-islet CD8+ cytotoxic T-cell survival to enhance type 1 diabetes. *Diabetes* 57 (2008) 909-17.
- [30] T.M. McCaughy, M.S. Wilken, and K.A. Hogquist, Thymic emigration revisited. *Journal of Experimental Medicine* 204 (2007) 2513-2520.
- [31] M.J. Smith, T.A. Packard, S.K. O'Neill, C.J.H. Dunand, M. Huang, L. Fitzgerald-Miller, D. Stowell, R.M. Hinman, P.C. Wilson, P.A. Gottlieb, and J.C. Cambier, Loss of Anergic B Cells in Prediabetic and New-Onset Type 1 Diabetic Patients. *Diabetes* 64 (2015) 1703-1712.
- [32] J.H. Li, E. Lu, T.S. Yi, and J.G. Cyster, EBI2 augments Tfh cell fate by promoting interaction with IL-2-quenching dendritic cells. *Nature* 533 (2016) 110-+.

- [33] K. Akashi, L.I. Richie, T. Miyamoto, W.H. Carr, and I.L. Weissman, B lymphopoiesis in the thymus. *J Immunol* 164 (2000) 5221-6.
- [34] D. Zotos, J.M. Coquet, Y. Zhang, A. Light, K. D'Costa, A. Kallies, L.M. Corcoran, D.I. Godfrey, K.M. Toellner, M.J. Smyth, S.L. Nutt, and D.M. Tarlinton, IL-21 regulates germinal center B cell differentiation and proliferation through a B cell-intrinsic mechanism. *J Exp Med* 207 (2010) 365-78.
- [35] H.M. McGuire, A. Vogelzang, N. Hill, M. Flodstrom-Tullberg, J. Sprent, and C. King, Loss of parity between IL-2 and IL-21 in the NOD Idd3 locus. *Proc Natl Acad Sci U S A* 106 (2009) 19438-43.
- [36] A.P. Sutherland, T. Van Belle, A.L. Wurster, A. Suto, M. Michaud, D. Zhang, M.J. Grusby, and M. von Herrath, Interleukin-21 is required for the development of type 1 diabetes in NOD mice. *Diabetes* 58 (2009) 1144-55.
- [37] R.C. Ferreira, H.Z. Simons, W.S. Thompson, A.J. Cutler, X.C. Dopico, D.J. Smyth, M. Mashar, H. Schuilenburg, N.M. Walker, D.B. Dunger, C. Wallace, J.A. Todd, L.S. Wicker, and M.L. Pekalski, IL-21 production by CD4(+) effector T cells and frequency of circulating follicular helper T cells are increased in type 1 diabetes patients. *Diabetologia* 58 (2015) 781-790.
- [38] F.S. Wong, and L. Wen, Innate and adaptive immune responses are highly interconnected at many levels. *Curr Mol Med* 9 (2009) 1-3.
- [39] S.E. Regnell, and A. Lernmark, Early prediction of autoimmune (type 1) diabetes. *Diabetologia* 60 (2017) 1370-1381.
- [40] C. Ferreira, Y. Singh, A.L. Furmanski, F.S. Wong, O.A. Garden, and J. Dyson, Non-obese diabetic mice select a low-diversity repertoire of natural regulatory T cells. *Proc Natl Acad Sci U S A* 106 (2009) 8320-5.
- [41] Y. Zhang, E. Bandala-Sanchez, and L.C. Harrison, Revisiting regulatory T cells in type 1 diabetes. *Curr Opin Endocrinol Diabetes Obes* 19 (2012) 271-8.
- [42] A. Kukreja, G. Cost, J. Marker, C. Zhang, Z. Sun, K. Lin-Su, S. Ten, M. Sanz, M. Exley, B. Wilson, S. Porcelli, and N. Maclaren, Multiple immuno-regulatory defects in type-1 diabetes. *J Clin Invest* 109 (2002) 131-40.
- [43] L.A. O'Reilly, D. Healey, E. Simpson, P. Chandler, T. Lund, M.A. Ritter, and A. Cooke, Studies on the thymus of non-obese diabetic (NOD) mice: effect of transgene expression. *Immunology* 82 (1994) 275-86.
- [44] N.M. Parish, P. Chandler, R. Quartey-Papafio, E. Simpson, and A. Cooke, The effect of bone marrow and thymus chimerism between non-obese diabetic (NOD) and NOD-E

- transgenic mice, on the expression and prevention of diabetes. *Eur J Immunol* 23 (1993) 2667-75.
- [45] I. Ferrero, F. Anjuere, P. Martin, G. Martinez del Hoyo, M.L. Fraga, N. Wright, R. Varona, G. Marquez, and C. Ardavin, Functional and phenotypic analysis of thymic B cells: role in the induction of T cell negative selection. *Eur J Immunol* 29 (1999) 1598-609.
- [46] S. Mori, M. Inaba, A. Sugihara, S. Taketani, H. Doi, Y. Fukuba, Y. Yamamoto, Y. Adachi, K. Inaba, S. Fukuhara, and S. Ikehara, Presence of B cell progenitors in the thymus. *J Immunol* 158 (1997) 4193-9.
- [47] J. Perera, and H. Huang, The development and function of thymic B cells. *Cell Mol Life Sci* 72 (2015) 2657-63.
- [48] M. Inaba, K. Inaba, Y. Adachi, K. Nango, H. Ogata, S. Muramatsu, and S. Ikehara, Functional analyses of thymic CD5+ B cells. Responsiveness to major histocompatibility complex class II-restricted T blasts but not to lipopolysaccharide or anti-IgM plus interleukin 4. *J Exp Med* 171 (1990) 321-6.
- [49] T. Viisanen, E.L. Ihantola, K. Nanto-Salonen, H. Hyoty, N. Nurminen, J. Selvenius, A. Juutilainen, L. Moilanen, J. Pihlajamaki, R. Veijola, J. Toppari, M. Knip, J. Ilonen, and T. Kinnunen, Circulating CXCR5+PD-1+ICOS+ Follicular T Helper Cells Are Increased Close to the Diagnosis of Type 1 Diabetes in Children With Multiple Autoantibodies. *Diabetes* 66 (2017) 437-447.
- [50] E. Astorri, M. Bombardieri, S. Gabba, M. Peakman, P. Pozzilli, and C. Pitzalis, Evolution of ectopic lymphoid neogenesis and in situ autoantibody production in autoimmune nonobese diabetic mice: cellular and molecular characterization of tertiary lymphoid structures in pancreatic islets. *J Immunol* 185 (2010) 3359-68.
- [51] Y. Takahama, I. Ohigashi, S. Baik, and G. Anderson, Generation of diversity in thymic epithelial cells. *Nat Rev Immunol* 17 (2017) 295-305.
- [52] N.F. Bernard, F. Ertug, and H. Margolese, High incidence of thyroiditis and anti-thyroid autoantibodies in NOD mice. *Diabetes* 41 (1992) 40-6.
- [53] C. Thompson, H. Jacobsen, D. Pomeranz Krummel, K. Nagai, and A. Cooke, Non-depleting anti-CD4 antibody not only prevents onset but resolves sialadenitis in NOD mice. *Autoimmunity* 37 (2004) 549-54.
- [54] M. Nakahara, Y. Nagayama, T. Ichikawa, L. Yu, G.S. Eisenbarth, and N. Abiru, The effect of regulatory T-cell depletion on the spectrum of organ-specific autoimmune diseases in nonobese diabetic mice at different ages. *Autoimmunity* 44 (2011) 504-10.

734 [55] F. Frommer, and A. Waisman, B cells participate in thymic negative selection of murine
735 auto-reactive CD4+ T cells. PLoS One 5 (2010) e15372.
736 [56] M. Le Borgne, E. Ladi, I. Dzhagalov, P. Herzmark, Y.F. Liao, A.K. Chakraborty, and E.A.
737 Robey, The impact of negative selection on thymocyte migration in the medulla. Nat
738 Immunol 10 (2009) 823-30.
739

Figure Legends

Figure 1. Intrathymic B cell accumulation precedes β cell destruction. Single cell suspensions were prepared from the respective thymi and all data analyzed on a single cell, live gate. (a) Representative dot plots of thymic CD19⁺ cells: (I) Isotype control for CD19 antibody; (II) 12 week old female B6 mouse; and (III) 12 week old female NOD mouse. (b) Number of B cells in the thymus of 4-6 week old female B6 mice (n= 11), 12-14 week old female B6 mice (n=25), 4-6 week old female NOD mice (n= 13) and 12-14 week old female NOD mice (n= 25). (c) Number of RAG-GFP^{hi} B cells in the thymus of 4-6 week old female FVB-RAG-GFP mice (n=3), 12-14 week old female FVB-RAG-GFP mice (n=8), 4-6 week old female NOD-RAG-GFP mice (n= 4) and 12-14 week old female NOD (n= 8). (d) Frequency of Ki67⁺ B cells in the thymus of 4-6 week old female B6 mice (n= 10), 12-14 week old female B6 mice (n=10), 4-6 week old female NOD (n= 9) and 12-14 week old female NOD mice (n= 8). Data presented as scatter plot, each dot equating to a mouse, the bar representing the mean value. Statistical significance determined using the non-parametric Mann-Whitney U-test, significant P values are shown, ns= not significant.

Figure 2. B cells form follicle-like structures at the cortical-medullary junction in NOD thymi. Flow cytometric analysis of the (a) frequency of B cells displaying a Follicular (FO) or Marginal-zone (MZ) phenotype in the thymus of B6 (n=10) or NOD mice (n=13) and (b) absolute number of B cells displaying FO or MZ phenotypes in the thymus of B6 (n=5) or NOD mice (n=6). Comparisons made between aged-matched, female 11-14 week old mice in a single cell, live gate. Data acquired from at least two independent experiments and is presented as scatter plot; P values were calculated using the Mann-Whitney U test analysis, ns= not significant. (c) Representative confocal immunofluorescence microscopy images of thymi

sections examined for B220 (yellow), cytokeratin V (green) expression and the DNA-intercalating dye DAPI identified nuclei (blue) from 11 week old female NOD or B6 mice. A total of 14 sections from eight NOD mice and a total of six section from three B6 mice were analyzed, and there was consistency in the data obtained from the appropriate strains of mice. Confocal fluorescent images were obtained with a Plan-Apochromat 20x objective.

Figure 3. The NOD thymus has the hallmarks of ectopic GC development. (a) Evaluation of IL-2/IL-21 ratio in cell-free supernatants from spleen and thymic tissue from 11-15 week old B6 (n=5) or NOD (n=4) mice. The data shown is representative of two individual experiments showing similar results. (b) Frequency of CD4⁺ T follicular helper cells (TfH) in the thymus of B6 (n=12) and NOD mice (n=17). (c) Number of CD4⁺ T follicular helper cells in the thymus of B6 (n=10) and NOD mice (n=11). (d) Representative histogram of BCL-6 expression in CD8⁻ CD4⁺ PD1^{hi} ICOS⁺ cells in thymus of NOD (n=5) and B6 (n=5) mice. (e) Number of plasma cells in the thymus of B6 (n=15) and NOD mice (n=15). For (b-e) comparisons made between female, age-matched 10-14 week old B6 and NOD mice. The analysis was performed on a single cell, live gate, and the data is presented as a scatterplot, each dot equating to a mouse, the bar represents mean value. P values were calculated using the Mann-Whitney U test analysis and are shown in the figure, ns= not significant. The data is pooled from at least two independent experiments giving similar results. (f) Quantitative PCR (qPCR) analysis of AID mRNA (*aicda* expression) levels in the whole thymus. Data was normalized to HPRT mRNA as described in methods, and fold change in NOD mice AID mRNA when compared to normalised AID mRNA levels for B6 mice. All mice were 11-14 weeks of age, a total of 5 female B6 mice were compared to 5 female NOD mice. One thymic sample from a female B cell-deficient NOD μ MT^{-/-} mouse was used as a negative control. The data is pooled from two independent experiments and is presented mean \pm standard error mean

(SEM). P values were calculated using the Mann-Whitney U-test and are shown in the figure, ns= not significant.

Figure 4. The NOD thymus harbours a unique pattern of immunoglobulin isotypes (a) Number of IgM⁻ IgD⁻ IgA⁺, IgM⁻ IgD⁻ IgE⁺, IgM⁻ IgD⁻ IgG⁺ B cells in the thymus of 11-14 week old female B6 (n=10) or female NOD mice (n=10). (b) Number of IgM⁻ IgD⁺ IgA⁺, IgM⁻ IgD⁺ IgE⁺, IgM⁻ IgD⁺ IgG⁺ B cells in the thymus of 11-14 week old female B6 (n=10) and female NOD mice (n=10). (c-f) Optical density (OD) values of the respective immunoglobulins in cell free tissue supernatants (c,e) or serum (d,f). A total of 6 female B6 and 6 female NOD mice were assessed in two independent experiments. Data is presented as scatter plot, each dot equating to one mouse and bar representing the mean. P values were calculated using the Mann-Whitney U-test analysis and are shown in the figure, ns= not significant.

Figure 5. IgGs bind to thymic stromal components in NOD mice. (a and b) Representative confocal immunofluorescence microscopy images of thymi sections of NOD (aI-III) and B6 mice (bI-II) examined for cytokeratin V (red), murine IgG (green), and the DNA-intercalating dye DAPI (white). A total of six 11 week old NOD mice and five 11 week old B6 mice, two sections per mouse were examined. (a)I,II and II are derived from different NOD mice. The confocal fluorescent image in AI was obtained with a Plan-Apochromat 20x objective to give a broader view of the extent of immunoglobulin bound to thymic stroma, arrows indicating some of the cells co-positive for cytokeratin V and mouse IgG. The confocal fluorescent images in AII and AIII were obtained with a Plan-Apochromat 63x objective. For (b) the confocal fluorescent image was obtained using a Plan-Apochromat 20x objective. (c) Quantification of murine Ig-bound to stromal cells of age-matched 11 week old, female NOD or B6 mice.

Confocal immunofluorescence microscopy images were subjected to StrataQuest V64 analysis, a total of $2-3 \times 10^4$ DAPI⁺ cells/mm² were counted and the mean fluorescence intensity of DAPI⁺ cells versus mean fluorescence intensity of anti-Ig is presented as a scattergram. The data shown, is representative of two independent mice examined giving similar results.

Figure 6. Increase in thymic B cells was associated to increased apoptosis of stromal cells.

(a) Representative confocal immunofluorescence microscopy images of thymi sections from 9-14 week old female NOD, and NOD- μ MT^{-/-} and B6 mice examined for cytokeratin V (yellow), apoptosis (red), and the DNA-intercalating dye DAPI (white) expression. The data is representative of similar data acquired from 6 female NOD, 6 female NOD- μ MT^{-/-} and 4 female B6 mice, three sections per mouse were examined. In all cases, the confocal fluorescent images were obtained with a Plan-Apochromat 63x objective. Bar represents 20 μ m. (b) Quantification of TUNEL⁺ stromal cells of age-matched 11 week old, female NOD, NOD- μ MT^{-/-} or B6 mice. Confocal immunofluorescence microscopy images were subjected to StrataQuest V64 analysis, a total of 4×10^4 DAPI⁺ cells/mm² were counted and the mean fluorescence intensity of DAPI⁺ cells versus mean fluorescence intensity of TUNEL is presented as a scattergram.

Figure 7. B cells promote premature thymic-release of T cells prior to negative selection.

(a-b) Thymocytes from 11-12 week old B6, NOD and NOD NOD- μ MTKO mice were stimulated with insulin or B15:23 peptide (NOD and NOD- μ MTKO mice, only) for 72 hours and Ki67 expression in CD4SP (a) or CD8SP (b) cells as a measure of proliferation was determined by flow cytometry. The frequency of Ki67⁺ cells for stimulated samples was normalized against the frequency of Ki67⁺ cells in unstimulated samples. (c) Frequency of

837 RTEs (RAG-GFP^{hi}) in peripheral blood of 11-12 week old FVB-GFP (n=8), NOD-GFP (n=8)
838 or NOD- μ MT^{-/-}-GFP (n=5) mice. Data is pooled from two independent experiments and cells
839 were analysed on a live, single, CD3⁺ T cell gate. The data is presented as scatter plot, the bar
840 representing the mean value; P values were calculated using the two-way Anova followed by
841 Tukey multi comparison test and are shown the figure, ns= not significant.