



# ANNALES DE L'INSTITUT FOURIER

Rudolf TANGE

**On the first restricted cohomology of a reductive Lie algebra  
and its Borel subalgebras**


Tome 69, n° 3 (2019), p. 1295-1308.

[http://aif.centre-mersenne.org/item/AIF\\_2019\\_\\_69\\_3\\_1295\\_0](http://aif.centre-mersenne.org/item/AIF_2019__69_3_1295_0)

---

© Association des Annales de l'institut Fourier, 2019,

*Certains droits réservés.*



Cet article est mis à disposition selon les termes de la licence

CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE.

<http://creativecommons.org/licenses/by-nd/3.0/fr/>



# ON THE FIRST RESTRICTED COHOMOLOGY OF A REDUCTIVE LIE ALGEBRA AND ITS BOREL SUBALGEBRAS

by Rudolf TANGE (\*)

---

ABSTRACT. — Let  $k$  be an algebraically closed field of characteristic  $p > 0$  and let  $G$  be a connected reductive group over  $k$ . Let  $B$  be a Borel subgroup of  $G$  and let  $\mathfrak{g}$  and  $\mathfrak{b}$  be the Lie algebras of  $G$  and  $B$ . Denote the first Frobenius kernels of  $G$  and  $B$  by  $G_1$  and  $B_1$ . Furthermore, denote the algebras of regular functions on  $G$  and  $\mathfrak{g}$  by  $k[G]$  and  $k[\mathfrak{g}]$ , and similarly for  $B$  and  $\mathfrak{b}$ . The group  $G$  acts on  $k[G]$  via the conjugation action and on  $k[\mathfrak{g}]$  via the adjoint action. Similarly,  $B$  acts on  $k[B]$  via the conjugation action and on  $k[\mathfrak{b}]$  via the adjoint action. We show that, under certain mild assumptions, the cohomology groups  $H^1(G_1, k[\mathfrak{g}])$ ,  $H^1(B_1, k[\mathfrak{b}])$ ,  $H^1(G_1, k[G])$  and  $H^1(B_1, k[B])$  are zero. We also extend all our results to the cohomology for the higher Frobenius kernels.

RÉSUMÉ. — Soit  $k$  un corps algébriquement clos de caractéristique  $p > 0$  and soit  $G$  un groupe réductif connexe sur  $k$ . Soit  $B$  un sous-groupe de Borel de  $G$  et soit  $\mathfrak{g}$  et  $\mathfrak{b}$  les algèbres de Lie de  $G$  et  $B$ . Notons les premiers noyaux de Frobenius de  $G$  et  $B$  par  $G_1$  et  $B_1$ . De plus, notons les algèbres des fonctions régulières sur  $G$  et  $\mathfrak{g}$  par  $k[G]$  et  $k[\mathfrak{g}]$ , et de même pour  $B$  et  $\mathfrak{b}$ . Le groupe  $G$  agit sur  $k[G]$  par conjugaison et sur  $k[\mathfrak{g}]$  par l'action adjointe. De même,  $B$  agit sur  $k[B]$  par l'action de conjugaison et sur  $k[\mathfrak{b}]$  par l'action adjointe. Nous montrons que, sous certaines hypothèses, les groupes de cohomologie  $H^1(G_1, k[\mathfrak{g}])$ ,  $H^1(B_1, k[\mathfrak{b}])$ ,  $H^1(G_1, k[G])$  et  $H^1(B_1, k[B])$  sont nuls. Nous étendons aussi nos résultats à la cohomologie pour les noyaux de Frobenius supérieurs.

## Introduction

Let  $k$  be an algebraically closed field of characteristic  $p > 0$ , let  $G$  be a connected reductive group over  $k$ , and let  $\mathfrak{g}$  be the Lie algebra of  $G$ . Recall that  $\mathfrak{g}$  is a restricted Lie algebra: it has a  $p$ -th power map  $x \mapsto x^{[p]} : \mathfrak{g} \rightarrow \mathfrak{g}$ , see [3, I.3.1]. In the case of  $G = \mathrm{GL}_n$  this is just the  $p$ -th matrix power. A

---

*Keywords:* Cohomology, Frobenius kernel, reductive group.

*2010 Mathematics Subject Classification:* 20G05, 20G10.

(\*) I would like to thank H. H. Andersen and J. C. Jantzen for helpful email discussions.

$\mathfrak{g}$ -module  $M$  is called *restricted* if  $(x^{[p]})_M = (x_M)^p$  for all  $x \in M$ . Here  $x_M$  is the endomorphism of  $M$  representing  $x$ .

Recall that an element  $v$  of a  $\mathfrak{g}$ -module  $M$  is called a  $\mathfrak{g}$ -invariant if  $x \cdot v = 0$  for all  $x \in \mathfrak{g}$ . We denote the space of  $\mathfrak{g}$ -invariants in  $M$  by  $M^{\mathfrak{g}}$ . The right derived functors of the left exact functor  $M \mapsto M^{\mathfrak{g}}$  from the category of restricted  $\mathfrak{g}$ -modules to the category of vector spaces over  $k$  are denoted by  $H^i(G_1, \cdot)$ .

Let  $k[\mathfrak{g}]$  be the algebra of polynomial functions on  $\mathfrak{g}$ . If one is interested in describing the algebra of invariants  $(k[\mathfrak{g}]/I)^{\mathfrak{g}}$  for some  $\mathfrak{g}$ -stable ideal  $I$  of  $k[\mathfrak{g}]$ , then it is of interest to know if  $H^1(G_1, k[\mathfrak{g}]) = 0$ , because then we have an exact sequence

$$k[\mathfrak{g}]^{\mathfrak{g}} \rightarrow (k[\mathfrak{g}]/I)^{\mathfrak{g}} \rightarrow H^1(G_1, I) \rightarrow 0$$

by the long exact cohomology sequence. So, in this case,  $(k[\mathfrak{g}]/I)^{\mathfrak{g}}$  is built up from the image of  $k[\mathfrak{g}]^{\mathfrak{g}}$  in  $k[\mathfrak{g}]/I$ , and  $H^1(G_1, I)$ .

The paper is organised as follows. In Section 1 we state some results from the literature that we need to prove our main result. This includes a description of the algebra of invariants  $k[\mathfrak{g}]^G$ , the normality of the nilpotent cone  $\mathcal{N}$ , and some lemmas on graded modules over graded rings. In Section 2 we prove Theorems 2.1 and 2.2 which state that, under certain mild assumptions on  $p$ ,  $H^1(G_1, k[\mathfrak{g}])$  and  $H^1(B_1, k[\mathfrak{b}])$  are zero. In Section 3 we prove Theorems 3.1 and 3.2 which state that, under certain mild assumptions on  $p$ ,  $H^1(G_1, k[G])$  and  $H^1(B_1, k[B])$  are zero. In Section 4 we extend these four theorems to the cohomology for the higher Frobenius kernels  $G_r$  and  $B_r$ ,  $r \geq 2$ .

We briefly indicate some background to our results. For convenience we only discuss the  $G$ -module  $k[\mathfrak{g}]$ . As is well-known, under certain mild assumptions  $k[\mathfrak{g}]$  has a good filtration, see [6] or [11, II.4.22]. So a natural first approach to prove that  $H^1(G_1, k[\mathfrak{g}]) = 0$  would be that to show that  $H^1(G_1, \nabla(\lambda)) = 0$  for all induced modules  $\nabla(\lambda)$  that show up in a good filtration of  $k[\mathfrak{g}]$ . However, this isn't true: even for  $p > h$ ,  $h$  the Coxeter number, one can easily deduce counterexamples from [1, Cor. 5.5] (or [11, II.12.15]).<sup>(1)</sup> It is also easy to see that we cannot have  $H^i(G_1, k[\mathfrak{g}]) = 0$  for all  $i > 0$ : the trivial module  $k$  is direct summand of  $k[\mathfrak{g}]$ , and for  $p > h$  we have  $H^\bullet(G_1, k) \cong k[\mathcal{N}]$  where the degrees of  $k[\mathcal{N}]$  are doubled, see [11, II.12.14].

The idea of our proof that  $H^1(G_1, k[\mathfrak{g}]) = 0$  is as follows. Noting that  $H^1(G_1, k[\mathfrak{g}])$  is a  $k[\mathfrak{g}]^G$ -module, we interpret a certain localisation of

---

<sup>(1)</sup>This approach *does* work when proving (the well-known fact) that  $H^1(G, k[\mathfrak{g}]) = 0$ .

$H^1(G_1, k[\mathfrak{g}])$  as the cohomology group of the coordinate ring of the generic fiber of the adjoint quotient map  $\mathfrak{g} \rightarrow \mathfrak{g} // G$ . It is easy to see that this cohomology group has to be zero, so we are left with showing that  $H^1(G_1, k[\mathfrak{g}])$  is torsion-free over the invariants  $k[\mathfrak{g}]^G$ . To prove the latter we use Hochschild's characterisation of the first restricted cohomology group and a "Nakayama Lemma type result". The ideas of the proofs of the other main results are completely analogous.

## 1. Preliminaries

Throughout this paper  $k$  is an algebraically closed field of characteristic  $p > 0$ . For the basics of representations of algebraic groups we refer to [11].

### 1.1. Restricted representations and restricted cohomology

Let  $G$  be a linear algebraic group over  $k$  with Lie algebra  $\mathfrak{g}$ . Let  $G_1$  be the first Frobenius kernel of  $G$  (see [11, Ch. I.9]). It is an infinitesimal group scheme with  $\dim k[G_1] = p^{\dim(\mathfrak{g})}$ . Its category of representations is equivalent to the category of restricted representations of  $\mathfrak{g}$ , see the introduction.

Let  $M$  be an  $G_1$ -module. By [7] (see also [11, I.9.19]) we have

$$(1.1) \quad H^1(G_1, M) = \{\text{restricted derivations : } \mathfrak{g} \rightarrow M\} / \{\text{inner derivations of } M\}.$$

Here a *derivation* from  $\mathfrak{g}$  to  $M$  is a linear map  $D : \mathfrak{g} \rightarrow M$  satisfying

$$D([x, y]) = x \cdot D(y) - y \cdot D(x)$$

for all  $x, y \in \mathfrak{g}$ . Such a derivation is called *restricted* if

$$D(x^{[p]}) = (x_M)^{p-1}(D(x))$$

for all  $x \in \mathfrak{g}$ , where  $x_M$  is the vector space endomorphism of  $M$  given by the action of  $x$ , and  $-^{[p]}$  denotes the  $p$ -th power map of  $\mathfrak{g}$ . An *inner derivation* of  $M$  is a map  $x \mapsto x \cdot u : \mathfrak{g} \rightarrow M$  for some  $u \in M$ . If  $M$  is restricted, then every inner derivation is restricted. Clearly  $H^1(G_1, M)$  is an  $G$ -module with trivial  $\mathfrak{g}$ -action: If  $D$  is a derivation and  $y \in \mathfrak{g}$ , then  $[y, D]$  is the inner derivation given by  $D(y)$ . Note also that  $H^1(G_1, k[\mathfrak{g}])$  is a  $k[\mathfrak{g}]^{\mathfrak{g}}$ -module, since the restricted derivations  $\mathfrak{g} \rightarrow k[\mathfrak{g}]$  form a  $k[\mathfrak{g}]^{\mathfrak{g}}$ -module and the map  $f \mapsto (x \mapsto x \cdot f)$  from  $k[\mathfrak{g}]$  to the restricted derivations  $\mathfrak{g} \rightarrow k[\mathfrak{g}]$  is  $k[\mathfrak{g}]^{\mathfrak{g}}$ -linear.

### 1.2. Actions of restricted Lie algebras

Let  $\mathfrak{g}$  be a restricted Lie algebra over  $k$ . Following [17] we define an action of  $\mathfrak{g}$  on an affine variety  $X$  over  $k$  to be a homomorphism  $\mathfrak{g} \rightarrow \text{Der}_k(k[X])$  of restricted Lie algebras, where  $\text{Der}_k(k[X])$  is the (restricted) Lie algebra of  $k$ -linear derivations of  $k[X]$ . It is easy to see that this includes the case that  $X$  is a restricted  $\mathfrak{g}$ -module. If  $\mathfrak{g}$  acts on  $X$  and  $x \in X$ , then we define  $\mathfrak{g}_x$  to be the stabiliser in  $\mathfrak{g}$  of the maximal ideal  $\mathfrak{m}_x$  of  $k[X]$  corresponding to  $x$ . In case  $X$  is a closed subvariety of a restricted  $\mathfrak{g}$ -module, then we have  $\mathfrak{g}_x = \{y \in \mathfrak{g} \mid y \cdot x = 0\}$ .

LEMMA 1.1. — *Let  $\mathfrak{g}$  be a restricted Lie algebra over  $k$  acting on a normal affine variety  $X$  over  $k$ . If  $\max_{x \in X} \text{codim}_{\mathfrak{g}} \mathfrak{g}_x = \dim X$ , then  $k[X]^{\mathfrak{g}} = k[X]^p$ .*

*Proof.* — By [17, Thm. 5.2(5)] we have  $[k(X) : k(X)^{\mathfrak{g}}] = p^{\dim(X)}$ . By [4, Cor. 3 to Thm. V.16.6.4] we have  $[k(X) : k(X)^p] = p^{\dim(X)}$ . So  $k(X)^{\mathfrak{g}} = k(X)^p$ , since we always have  $\supseteq$ . Clearly,  $k(X)^p = \text{Frac}(k[X]^p)$ ,  $k(X)^{\mathfrak{g}} = \text{Frac}(k[X]^{\mathfrak{g}})$  and  $k[X]^{\mathfrak{g}}$  is integral over  $k[X]^p$ . Since  $X$  is normal variety,  $k[X]^p \cong k[X]$  is a normal ring. It follows that  $k[X]^{\mathfrak{g}} = k[X]^p$ .  $\square$

### 1.3. Two lemmas on graded rings and modules

We recall a version of the graded Nakayama lemma which follows from [14, Ch. 13, Lem. 4, Ex. 3, Lem. 3].

LEMMA 1.2 ([14, Ch. 13]). — *Let  $R = \bigoplus_{i \geq 0} R^i$  be a positively graded ring with  $R^0$  a field, let  $M$  be a positively graded  $R$ -module and let  $(x_i)_{i \in I}$  be a family of homogeneous elements of  $M$ . Put  $R^+ = \bigoplus_{i > 0} R^i$ .*

- (1) *If the images of the  $x_i$  in  $M/R^+M$  span the vector space  $M/R^+M$  over  $R^0$ , then the  $x_i$  generate  $M$ .*
- (2) *If  $M$  is projective and the images of the  $x_i$  in  $M/R^+M$  form an  $R^0$ -basis of  $M/R^+M$ , then  $(x_i)_{i \in I}$  is an  $R$ -basis of  $M$ .*

LEMMA 1.3. — *Let  $R$  be a positively graded ring with  $R^0$  a field and let  $N$  be a positively graded  $R$ -module which is projective.*

- (1) *Let  $M$  be a submodule of  $N$  with  $(R^+N) \cap M \subseteq R^+M$ . Then  $M$  is free and a direct summand of  $N$ .*
- (2) *Let  $M$  be a positively graded  $R$ -module, let  $\varphi : M \rightarrow N$  be a graded  $R$ -linear map and let  $\bar{\varphi} : M/R^+M \rightarrow N/R^+N$  be the induced  $R^0$ -linear map. Assume the canonical map  $M \rightarrow M/R^+M$  maps  $\text{Ker}(\varphi)$  onto  $\text{Ker}(\bar{\varphi})$ . Then  $\text{Im}(\varphi)$  is free and a direct summand of  $N$ .*

*Proof.*

(1). — From the assumption it is immediate that the natural map  $M/R^+M \rightarrow N/R^+N$  is injective. Now choose an  $R^0$ -basis  $(\bar{x}_i)_{i \in I}$  of  $M/R^+M$  and extend it to a basis  $(\bar{x}_i)_{i \in I \cup J}$  of  $N/R^+N$ . Let  $(x_i)_{i \in I \cup J}$  be a homogeneous lift of this basis to  $N$ . Then this is a basis of  $N$  by Lemma 1.2(2). Furthermore,  $(x_i)_{i \in I}$  must span  $M$  by Lemma 1.2(2). So  $M$  is (graded-) free and has the  $R$ -span of  $(x_i)_{i \in J}$  as a direct complement.

(2). — By (1) it suffices to show that  $(R^+N) \cap \text{Im}(\varphi) \subseteq R^+ \text{Im}(\varphi)$ . Let  $x \in M$  and assume that  $\varphi(x) \in R^+N$ . Then  $\bar{x} := x + R^+M \in \text{Ker}(\bar{\varphi})$ . By assumption there exists  $x_1 \in \text{Ker}(\varphi)$  such that  $\bar{x} = \bar{x}_1$ . Then  $x - x_1 \in R^+M$  and  $\varphi(x) = \varphi(x - x_1) \in R^+ \text{Im}(\varphi)$ .  $\square$

There is also an obvious version of Lemma 1.3 (and of course of Lemma 1.2) for a local ring  $R$ : simply assume  $R$  local, omit the gradings everywhere and replace  $R^+$  by the maximal ideal of  $R$ .

#### 1.4. The standard hypotheses and consequences

In the remainder of this paper  $G$  is a connected reductive group over  $k$  and  $\mathfrak{g}$  is its Lie algebra. Recall that  $\mathfrak{g}$  is a restricted Lie algebra, see [3, I.3.1], we denote its  $p$ -th power map by  $x \mapsto x^{[p]}$ . The group  $G$  acts on  $\mathfrak{g}$  and the nilpotent cone  $\mathcal{N}$  via the adjoint action and on  $G$  via conjugation, and therefore it also acts on their algebras of regular functions:  $k[\mathfrak{g}]$ ,  $k[\mathcal{N}]$  and  $k[G]$ . Fix a maximal torus  $T$  of  $G$  and let  $\mathfrak{t}$  be its Lie algebra. We fix an  $\mathbb{F}_p$ -structure on  $G$  for which  $T$  is defined and split over  $\mathbb{F}_p$ . Then  $\mathfrak{g}$  has an  $\mathbb{F}_p$ -structure and  $\mathfrak{t}$  is  $\mathbb{F}_p$ -defined. Denote the  $\mathbb{F}_p$ -defined regular functions on  $\mathfrak{g}$  and  $\mathfrak{t}$  by  $\mathbb{F}_p[\mathfrak{g}]$  and  $\mathbb{F}_p[\mathfrak{t}]$ . We will need the following standard hypotheses, see [10, 6.3, 6.4] or [12, 2.6, 2.9]:

- (H1) The derived group  $DG$  of  $G$  is simply connected,
- (H2)  $p$  is good for  $G$ ,
- (H3) There exists a  $G$ -invariant non-degenerate bilinear form on  $\mathfrak{g}$ .

Put  $G_x = \{g \in G \mid \text{Ad}(g)(x) = x\}$  and  $\mathfrak{g}_x = \{y \in \mathfrak{g} \mid [y, x] = 0\}$ . Assuming (H1)–(H3) we have by [12, 2.9] that  $\text{Lie}(G_x) = \mathfrak{g}_x$  for all  $x \in \mathfrak{g}$ . See also [15, Sect. 2.1]. Put  $n = \dim(T)$ . We call  $x \in \mathfrak{g}$  *regular* if  $\dim G_x$  (or  $\dim \mathfrak{g}_x$ ) equals  $n$ , the minimal value. Under assumptions (H1) and (H3) we have that  $d\alpha \neq 0$  for all roots  $\alpha$ , so restriction of functions defines an isomorphism  $k[\mathfrak{g}]^G \xrightarrow{\sim} k[\mathfrak{h}]^W$ , see [12, Prop. 7.12]. The set of regular semisimple elements in  $\mathfrak{g}$  is the nonzero locus of the regular function  $f_{\text{rs}}$  on

$\mathfrak{g}$  which corresponds under the above isomorphism to the product of the differentials of the roots. Note that  $f_{rs} \in \mathbb{F}_p[\mathfrak{g}]$ :  $f_{rs}$  is defined over  $\mathbb{F}_p$ .

Under assumptions (H1)–(H3) it follows from work of Demazure [5] that  $k[t]^W$  is a polynomial algebra in  $n$  homogeneous elements defined over  $\mathbb{F}_p$ , see [10, 9.6 end of proof]. We denote the corresponding elements of  $\mathbb{F}_p[\mathfrak{g}]$  by  $s_1, \dots, s_n$ . Assuming (H1)–(H3) the vanishing ideal of  $\mathcal{N}$  in  $k[\mathfrak{g}]$  is generated by the  $s_i$ , see [12, 7.14], and all regular orbit closures are normal, in particular  $\mathcal{N}$  is normal, see [12, 8.5].

We call  $g \in G$  regular if  $G_g := \{h \in G \mid hgh^{-1} = g\}$  has dimension  $n$ , the minimal value. Restriction of functions defines an isomorphism  $k[G]^G \xrightarrow{\sim} k[T]^W$ , see [19, 6.4]. The set of regular semisimple elements in  $G$  is the nonzero locus of the regular function  $f'_{rs}$  on  $G$  which corresponds under the above isomorphism to  $\prod_{\alpha \text{ a root}} (\alpha - 1)$ . If  $G$  is semisimple, simply connected, then  $k[G]^G$  is a polynomial algebra in the characters  $\chi_1, \dots, \chi_n$  of the irreducible  $G$ -modules whose highest weights are the fundamental dominant weights. Furthermore, the schematic fibers of the adjoint quotient  $G \rightarrow G//G$  are reduced and normal and they are regular orbit closures. See [19] and [8, 4.24]. One can also deduce from (H1)–(H3) that  $\text{Lie}(G_g) = \mathfrak{g}_g := \{x \in \mathfrak{g} \mid \text{Ad}(g)(x) = x\}$ .

## 2. The cohomology groups $H^1(G_1, k[\mathfrak{g}])$ and $H^1(B_1, k[\mathfrak{b}])$

Throughout this section we assume that hypotheses (H1)–(H3) from Section 1.4 hold.

**THEOREM 2.1.** —  $H^1(G_1, k[\mathfrak{g}]) = 0$ .

*Proof.* — Let  $K$  be an algebraic closure of the field of fractions of  $R := k[\mathfrak{g}]^G$ . Since the action of  $\mathfrak{g}$  on  $k[\mathfrak{g}]$  is  $R$ -linear we have  $H^1(G_1, k[\mathfrak{g}]) = H^1((G_1)_R, k[\mathfrak{g}])$ , where  $-_R$  denotes base change from  $k$  to  $R$ , see [11, I.1.10]. So, by the Universal Coefficient Theorem [11, Prop. I.4.18], we have

$$K \otimes_R H^1(G_1, k[\mathfrak{g}]) = H^1((G_1)_K, K \otimes_R k[\mathfrak{g}]) = H^1((G_K)_1, K \otimes_R k[\mathfrak{g}]).$$

For  $i \in \{1, \dots, n\}$  denote the regular function on  $\mathfrak{g}_K$  corresponding to  $s_i \in k[\mathfrak{g}]$  by  $\tilde{s}_i$ . Then  $K \otimes_R k[\mathfrak{g}] = K[\mathfrak{g}_K]/(\tilde{s}_1 - s_1, \dots, \tilde{s}_n - s_n) = K[F]$ , where  $F \subseteq \mathfrak{g}_K$  is the fiber of the morphism

$$x \mapsto (\tilde{s}_1(x), \dots, \tilde{s}_n(x)) : \mathfrak{g}_K \rightarrow \mathbb{A}_K^n$$

over the point  $(s_1, \dots, s_n) \in \mathbb{A}_K^n$ . Let  $f_{rs} \in \mathbb{F}_p[\mathfrak{g}] \cap k[\mathfrak{g}]^G$  be the polynomial function from Section 1.4 with nonzero locus the set of regular semisimple elements in  $\mathfrak{g}$ , and let  $\tilde{f}_{rs}$  be the corresponding polynomial function on

$\mathfrak{g}_K$ . Then we have for all  $x \in F$  that  $\tilde{f}_{rs}(x) = f_{rs} \neq 0$ . So  $F$  consists of regular semisimple elements. By [20, Lem. 3.7, Thm. 3.14] this means that  $F = G_K/S$  for some maximal torus  $S$  of  $G_K$ . In particular,  $K[F]$  is an injective  $G_K$ -module. But then it is also injective as a  $(G_K)_1$ -module, see [11, Rem. I.4.12, Cor. I.5.13b)]. So  $K \otimes_R H^1(G_1, k[\mathfrak{g}]) = H^i((G_K)_1, K[F]) = 0$  for all  $i > 0$ .

So it now suffices to show that  $H^1(G_1, k[\mathfrak{g}])$  has no  $R$ -torsion. We are going to apply Lemma 1.3(2) to the  $R$ -linear map

$$\varphi = f \mapsto (x \mapsto x \cdot f) : k[\mathfrak{g}] \rightarrow \text{Hom}_k(\mathfrak{g}, k[\mathfrak{g}]).$$

Here the grading of  $\text{Hom}_k(\mathfrak{g}, k[\mathfrak{g}])$  is given by

$$\text{Hom}_k(\mathfrak{g}, k[\mathfrak{g}])^i = \text{Hom}_k(\mathfrak{g}, k[\mathfrak{g}]^i).$$

As explained in [12, 7.13, 7.14] the conditions of [16, Prop. 10.1] are satisfied under the assumptions (H1)–(H3), so  $k[\mathfrak{g}]$  is a free  $R$ -module. So  $\text{Hom}_k(\mathfrak{g}, k[\mathfrak{g}])$  is also a free  $R$ -module. We have  $k[\mathfrak{g}]/R^+k[\mathfrak{g}] = k[\mathcal{N}]$ , and

$$\bar{\varphi} = f \mapsto (x \mapsto x \cdot f) : k[\mathcal{N}] \rightarrow \text{Hom}_k(\mathfrak{g}, k[\mathcal{N}]).$$

By [12, 6.3,6.4], we have  $\min_{x \in \mathcal{N}} \dim \mathfrak{g}_x = n$  and  $\dim \mathcal{N} = \dim \mathfrak{g} - n$ . So from Lemma 1.1 it is clear that the restriction map  $k[\mathfrak{g}] \rightarrow k[\mathcal{N}]$  maps the  $\mathfrak{g}$ -invariants of  $k[\mathfrak{g}]$  onto those of  $k[\mathcal{N}]$ . But  $k[\mathfrak{g}]^{\mathfrak{g}} = \text{Ker}(\varphi)$  and  $k[\mathcal{N}]^{\mathfrak{g}} = \text{Ker}(\bar{\varphi})$ . So, by Lemma 1.3(2),  $\text{Im}(\varphi)$  is a direct  $R$ -module summand of  $\text{Hom}_k(\mathfrak{g}, k[\mathfrak{g}])$ . In particular,  $\text{Hom}_k(\mathfrak{g}, k[\mathfrak{g}])/\text{Im}(\varphi)$  is isomorphic to an  $R$ -submodule of  $\text{Hom}_k(\mathfrak{g}, k[\mathfrak{g}])$  and therefore  $R$ -torsion-free. From (1.1) in Section 1.1 it is clear that  $H^1(G_1, k[\mathfrak{g}])$  is isomorphic to an  $R$ -submodule of  $\text{Hom}_k(\mathfrak{g}, k[\mathfrak{g}])/\text{Im}(\varphi)$ , so it is also  $R$ -torsion-free.  $\square$

Let  $B$  be a Borel subgroup of  $G$  containing  $T$ , let  $\mathfrak{b}$  be its Lie algebra and let  $\mathfrak{u}$  be the Lie algebra of the unipotent radical  $U$  of  $B$ .

**THEOREM 2.2.** —  $H^1(B_1, k[\mathfrak{b}]) = 0$ .

*Proof.* — Consider the restriction map  $k[\mathfrak{b}]^B \rightarrow k[\mathfrak{t}]$ . Under the assumptions (H1)–(H3)  $\mathfrak{t}$  contains elements which are regular in  $\mathfrak{g}$ . Furthermore, the set of regular semisimple elements in  $\mathfrak{g}$  is open in  $\mathfrak{g}$ . So the regular semisimple elements of  $\mathfrak{g}$  in  $\mathfrak{b}$  are dense in  $\mathfrak{b}$ . Since the union of the  $B$ -conjugates of  $\mathfrak{t}$  is the set of all semisimple elements in  $\mathfrak{b}$ , by [3, Prop. 11.8], it is also dense in  $\mathfrak{b}$ . This shows that the map  $k[\mathfrak{b}]^B \rightarrow k[\mathfrak{t}]$  is injective. Furthermore,  $\text{Ad}(g)(x) - x \in \mathfrak{u}$  for all  $g \in B$  and  $x \in \mathfrak{b}$  by [3, Prop. 3.17], since  $DB \subseteq U$ . So if we extend  $f \in k[\mathfrak{t}]$  to a regular function  $f$  on  $\mathfrak{b}$  by  $f(x + y) = f(x)$  for all  $x \in \mathfrak{t}$  and  $y \in \mathfrak{u}$ , then  $f \in k[\mathfrak{b}]^B$ . So the map



$k[\mathfrak{b}]^B \rightarrow k[\mathfrak{t}]$  is surjective, that is, restriction of functions defines an isomorphism

$$k[\mathfrak{b}]^B \xrightarrow{\sim} k[\mathfrak{t}].$$

Extend a basis of  $\mathfrak{t}^*$  to (linear) functions  $\xi_1, \dots, \xi_n$  on  $\mathfrak{b}$  in the manner indicated above. Then these functions are algebraically independent generators of  $k[\mathfrak{b}]^B$ , and  $k[\mathfrak{b}]$  is a free  $k[\mathfrak{b}]^B$ -module. Clearly, the vanishing ideal of  $\mathfrak{u}$  in  $k[\mathfrak{b}]$  is generated by the  $\xi_i$ . Furthermore,  $\min_{x \in \mathfrak{u}} \dim \mathfrak{b}_x = n$ , see [12, 6.8]. We can now follow the same arguments as in the proof of Theorem 2.1. Just replace  $G, \mathfrak{g}, \mathcal{N}, k[\mathfrak{g}]^G$  and the  $s_i$  by  $B, \mathfrak{b}, \mathfrak{u}, k[\mathfrak{b}]^B$  and the  $\xi_i$ , and replace  $f_{rs}$  by its restriction to  $\mathfrak{b}$ .  $\square$

*Remark 2.3.* — We have  $k[\mathcal{N}] = \text{ind}_B^G k[\mathfrak{u}]$ . Using [11, Lem. II.12.12a)] and the arguments from [11, II.12.2] it follows that  $H^1(G_1, k[\mathcal{N}]) = \text{ind}_B^G H^1(B_1, k[\mathfrak{u}])$ . From this one can easily deduce examples with  $H^1(G_1, k[\mathcal{N}]) \neq 0$ .

### 3. The cohomology groups $H^1(G_1, k[G])$ and $H^1(B_1, k[B])$

Assume first that  $G = \text{GL}_n$ . Put  $R = k[\mathfrak{g}]^G$  and  $R_1 = R[\det^{-1}]$ . Then, using the fact that the  $\mathfrak{g}$ -action on  $k[G]$  is  $R_1$ -linear, the Universal Coefficient Theorem and Theorem 2.1, we obtain

$$\begin{aligned} H^1(G_1, k[G]) &= H^1((G_1)_{R_1}, k[G]) = R_1 \otimes_R H^1((G_1)_R, k[\mathfrak{g}]) \\ &= R_1 \otimes_R H^1(G_1, k[\mathfrak{g}]) = 0. \end{aligned}$$

Similarly, we obtain  $H^1(B_1, k[B]) = 0$ .

To prove our result for the case of arbitrary reductive  $G$  we assume in this section the following:

*There exists a central (see [3, 22.3]) surjective morphism  $\psi : \tilde{G} \rightarrow G$  where  $\tilde{G}$  is a direct product of groups of the following types:*

- (1) a simply connected simple algebraic group of type  $\neq A$  for which  $p$  is good,
- (2)  $\text{SL}_m$  for  $p \nmid m$ ,
- (3)  $\text{GL}_m$ ,
- (4) a torus.

**THEOREM 3.1.** —  $H^1(G_1, k[G]) = 0$ .

*Proof.* — First we reduce to the case that  $G$  is of one of the above four types. Let  $\psi : \tilde{G} \rightarrow G$  be as above. Then  $G$  is the quotient of  $\tilde{G}$  by a (schematic) central diagonalisable closed subgroup scheme  $\tilde{Z}$ , see [11,

II.1.18]. Let  $N$  be the image of  $\tilde{G}_1$  in  $G_1$ . Then  $N$  is normal in  $G_1$  and  $G_1/N$  is diagonalisable. So  $H^i(G_1, k[G]) = H^i(N, k[G])^{G_1/N}$ , by [11, I.6.9(3)]. Furthermore,  $H^i(N, k[G]) = H^i(\tilde{G}_1, k[G])$ , by [11, I.6.8(3)], since the kernel of  $\tilde{G}_1 \rightarrow N$  is central.

The group scheme  $\tilde{Z}$  also acts via the right multiplication action on  $k[\tilde{G}]$  and this action commutes with the conjugation action of  $\tilde{G}$ . So  $k[G] = k[\tilde{G}]^{\tilde{Z}}$  is a direct  $\tilde{G}$ -module summand of  $k[\tilde{G}]$ . So it suffices to show that  $H^1(\tilde{G}_1, k[\tilde{G}]) = 0$ . By the Künneth Theorem we may now assume that  $G$  is of one of the above four types.

For  $G$  a torus the assertion is obvious, and for  $G = \text{GL}_n$  we have already proved the assertion. Now assume that  $G$  is of type (1) or (2). Then  $G$  satisfies (H1)–(H3) and  $G$  is simply connected simple. By [20, 2.15] the centraliser of a semisimple group element is connected, so when the element is also regular, its centraliser is a maximal torus. As in the proof of Theorem 2.1 we are now reduced to showing that  $H^1(G_1, k[G])$  has no torsion over  $R := k[G]^G$ .

For this it is enough that  $R_{\mathfrak{m}} \otimes_R H^1(G_1, k[G]) = H^1(G_1, R_{\mathfrak{m}} \otimes_R k[G])$  has no torsion over  $R_{\mathfrak{m}}$  for all maximal ideals  $\mathfrak{m}$  of  $R$ . By [19, 6.11, 7.16, 8.1] the conditions of [16, Prop. 10.1] are satisfied, so  $k[G]$  is a free  $R$ -module and  $k[G]_{\mathfrak{m}} = R_{\mathfrak{m}} \otimes_R k[G]$  is a free  $R_{\mathfrak{m}}$ -module for all maximal ideals  $\mathfrak{m}$  of  $R$ . Furthermore,  $k[G]_{\mathfrak{m}}/\mathfrak{m}_{\mathfrak{m}}k[G]_{\mathfrak{m}} = k[G]/\mathfrak{m}k[G]$  is the coordinate ring of a fiber  $F$  of the adjoint quotient map. We know  $F$  is normal of codimension  $n$ , and a regular orbit closure, so  $k[F]^{\mathfrak{g}} = k[F]^p$  by Lemma 1.1. By the local version of Lemma 1.3 the  $R_{\mathfrak{m}}$ -linear map  $\varphi = f \mapsto (x \mapsto x \cdot f) : k[G]_{\mathfrak{m}} \rightarrow \text{Hom}_k(\mathfrak{g}, k[G]_{\mathfrak{m}})$  we now get that  $H^1(G_1, R_{\mathfrak{m}} \otimes_R k[G])$  has no  $R_{\mathfrak{m}}$ -torsion. □

Let  $B$  be a Borel subgroup of  $G$ .

**THEOREM 3.2.** —  $H^1(B_1, k[B]) = 0$ .

*Proof.* — This follows by modifying the proof of Theorem 3.1 in the same way as the proof of Theorem 2.1 was modified to obtain the proof of Theorem 2.2. □

*Remark 3.3.* — One can also prove Theorem 3.2 assuming (H1)–(H3). The point is that it is obvious that restriction of functions always defines an isomorphism  $k[B]^B \xrightarrow{\sim} k[T]$ .

*Remark 3.4.* — We briefly discuss the  $B$ -cohomology of  $k[B]$  and  $k[\mathfrak{b}]$ . From [13, Thm. 1.13, Thm. 1.7(a)(ii)] it is immediate that  $H^i(B, k[B]) = 0$  for all  $i > 0$ . Now assume that there exists a central surjective morphism

$\psi : G \rightarrow G$  where  $\tilde{G}$  is a direct product of groups of the types (1)–(4) mentioned before, except that for type (2) we drop the condition on  $p$ . Then we deduce from the arguments from the proof of [1, Prop. 4.4] that  $H^i(B, k[\mathfrak{b}]) = 0$  for all  $i > 0$  as follows. First we reduce as in the proof of Theorem 3.1 to the case that  $G$  is simple of type (1) or (2) and then we deal with type (2) as in [1]. Now assume  $G$  is of type (1) and let  $I$  be the vanishing ideal of  $B$  in  $k[G]$ . As in [1] write

$$(3.1) \quad \mathfrak{m} = M \oplus \mathfrak{m}^2$$

where  $\mathfrak{m}$  is the vanishing ideal in  $k[G]$  of the unit element of  $G$  and  $M \cong \mathfrak{g}^*$  as  $G$ -modules. It suffices to show that  $I = I \cap M + I \cap \mathfrak{m}^2$ , since then we get a decomposition analogous to (3.1) for  $k[B]$  and we can finish as in [1]. Let  $f \in I$ . Then the  $M$ -component of  $f$  correspond to  $df \in \mathfrak{g}^*$  which vanishes on  $\mathfrak{b}$ . This means it corresponds under the trace form of the chosen representation  $\rho : G \rightarrow V$  (the adjoint representation for exceptional types) to an element  $x \in \mathfrak{u}$ . So the  $M$ -component of  $f$  is  $g \mapsto \text{tr}(\rho(g)d\rho(x))$  which vanishes on  $B$ . But then the  $\mathfrak{m}^2$ -component of  $f$  must also vanish on  $B$ .

#### 4. The cohomology groups for the higher Frobenius kernels

In this section we will generalise the results from the previous two sections to all Frobenius kernels  $G_r, r \geq 1$ .

LEMMA 4.1. — *Let  $G$  be a linear algebraic group over  $k$  acting on a normal affine variety  $X$  over  $k$ . If  $\max_{x \in X} \text{codim}_{\mathfrak{g}} \mathfrak{g}_x = \dim X$ , then  $k[X]^{G_r} = k[X]^{p^r}$  for all integers  $r \geq 1$ .*

*Proof.* — Since  $\text{codim}_{\mathfrak{g}} \mathfrak{g}_x \leq \text{codim}_G G_x \leq \dim(X)$  and  $\max_{x \in X} \text{codim}_{\mathfrak{g}} \mathfrak{g}_x = \dim X$  we must have that for  $x \in X$  with  $\text{codim}_{\mathfrak{g}} \mathfrak{g}_x = \dim X$  the schematic centraliser of  $x$  in  $G$  is reduced. So  $(G_r)_x = (G_x)_r$  and

$$\begin{aligned} (G_r : (G_r)_x) &:= \dim(k[G_r]) / \dim(k[(G_r)_x]) \\ &= p^{r \dim(G)} / p^{r \dim(G_x)} = p^{r \dim(X)}. \end{aligned}$$

By [17, Thm. 2.1 (5)] we get  $[k(X) : k(X)^{G_r}] = p^{r \dim(X)}$ . By [4, Cor. 3 to Thm. V.16.6.4] and the tower law we have  $[k(X) : k(X)^{p^r}] = p^{r \dim(X)}$ . So  $k(X)^{G_r} = k(X)^{p^r}$ , since we always have  $\supseteq$ . Clearly,  $k(X)^{p^r} = \text{Frac}(k[X]^{p^r})$ ,  $k(X)^{G_r} = \text{Frac}(k[X]^{G_r})$  and  $k[X]^{G_r}$  is integral over  $k[X]^{p^r}$ . Since  $X$  is normal variety,  $k[X]^{p^r} \cong k[X]$  is a normal ring. It follows that  $k[X]^{G_r} = k[X]^{p^r}$ . □

THEOREM 4.2. — *Let  $r$  be an integer  $\geq 1$ .*

(1) *Under the assumptions of Section 2 we have*

$$H^1(G_r, k[\mathfrak{g}]) = 0 \text{ and } H^1(B_r, k[\mathfrak{b}]) = 0.$$

(2) *Under the assumptions of Section 3 we have*

$$H^1(G_r, k[G]) = 0 \text{ and } H^1(B_r, k[B]) = 0.$$

*Proof.*

(1). — Let  $(H, M)$  be the group and module in question, i.e.  $(G, k[\mathfrak{g}])$  or  $(B, k[\mathfrak{b}])$ . Put  $R = k[\mathfrak{h}]^H$ . Let  $\varphi$  be the first map in the Hochschild complex of the  $H_r$ -module  $M$ , see [11, I.4.14]:

$$\varphi = f \mapsto (\Delta_M(f) - 1 \otimes f) : M \rightarrow k[H_r] \otimes M.$$

Then the induced map  $\bar{\varphi} : M/R^+M \rightarrow k[H_r] \otimes (M/R^+M)$  is the first map in the Hochschild complex of the  $H_r$ -module  $M/R^+M$  which is  $k[\mathcal{N}]$  or  $k[\mathfrak{u}]$ . So  $\text{Ker}(\varphi) = M^{H_r}$  and  $\text{Ker}(\bar{\varphi}) = (M/R^+M)^{H_r}$ . Now the proof is the same as that of the corresponding result in Section 2, except that we work with the above map  $\varphi$  and instead of Lemma 1.1 we apply Lemma 4.1.

(2). — Let  $(H, M)$  be the group and module in question, i.e.  $(G, k[G])$  or  $(B, k[B])$ . As in the proof of the corresponding result in Section 3 we reduce to the case that  $G$  is simple of type (1) or (2). Put  $R = k[H]^H$ . Fix a maximal ideal  $\mathfrak{m}$  of  $R$ . Let  $\varphi$  be the first map in the Hochschild complex of the  $H_r$ -module  $M_{\mathfrak{m}}$ . Then the induced map  $\bar{\varphi} : M_{\mathfrak{m}}/\mathfrak{m}_{\mathfrak{m}}M_{\mathfrak{m}} \rightarrow k[H_r] \otimes M_{\mathfrak{m}}/\mathfrak{m}_{\mathfrak{m}}M_{\mathfrak{m}}$  is the first map in the Hochschild complex of the  $H_r$ -module  $M_{\mathfrak{m}}/\mathfrak{m}_{\mathfrak{m}}M_{\mathfrak{m}} = M/\mathfrak{m}M$  which is the coordinate ring of the fiber of  $H \rightarrow H//H$  over the point  $\mathfrak{m}$ . So  $\text{Ker}(\varphi) = (M_{\mathfrak{m}})^{H_r}$  and  $\text{Ker}(\bar{\varphi}) = (M/\mathfrak{m}M)^{H_r}$ . Now the proof is the same as that of the corresponding result in Section 3, except that we work with the above map  $\varphi$  and instead of Lemma 1.1 we apply Lemma 4.1.  $\square$

*Remark 4.3.* — For  $G$  classical with natural module  $V = k^n$  we consider the cohomology groups  $H^1(G_r, S^i V)$  and  $H^1(G_r, S^i(V^*))$ .

Results about these modules can mostly easily be deduced from results on induced modules in the literature. For induced modules one can reduce to  $B_r$ -cohomology using the following result of Andersen–Jantzen for general  $G$ . Let  $B$  be a Borel subgroup of  $G$  with unipotent radical  $U$  and let  $T$  be a maximal torus of  $B$ . For  $\lambda \in X(T)$ , the character group of  $T$ , we denote by  $\nabla(\lambda)$ , the  $G$ -module induced from the 1-dimensional  $B$ -module given by  $\lambda$ . We call the roots of  $T$  in the opposite Borel subgroup  $B^+$  positive. By [11, II.12.2] we have for  $\lambda$  dominant

$$(4.1) \quad H^1(G_r, \nabla(\lambda))^{[-r]} \cong \text{ind}_B^G(H^1(B_r, \lambda)^{[-r]}).$$

Below we will always take  $\lambda = \varpi_1$  the first non-constant diagonal matrix coordinate. First take  $G = \text{GL}_n$ . Let  $B$  and  $T$  be the lower triangular matrices and the diagonal matrices. Then the character group  $X(T)$  of  $T$  identifies with  $\mathbb{Z}^n$ . Let  $\varepsilon_1$  be the first standard basis element of  $X(T)$ , i.e. the character  $D \mapsto D_{ii}$ . Then  $S^i V = \nabla(i\varepsilon_1)$  and  $S^i(V^*) = \nabla(-i\varepsilon_n)$ . Replacing  $\mathbf{u}^{*[s]}$  by  $\lambda \otimes \mathbf{u}^{*[s]}$  for  $\lambda = i\varepsilon_1$  or  $\lambda = -i\varepsilon_n$  in the proof of [11, Lem. II.12.1] and using (4.1) we obtain  $H^1(G_r, S^i V) = H^1(G_r, S^i(V^*)) = 0$ .

Now take  $G = \text{SL}_n$ . Then  $S^i V = \nabla(i\varpi_1)$  and  $S^i(V^*) = \nabla(i\varpi_{n-1})$ , where  $\varpi_j$  denotes the  $j$ -th fundamental dominant weight. From [2, Cor. 3.2(a)] we easily deduce that  $H^1(G_r, S^i V) \neq 0$  if and only if  $H^1(G_r, S^i(V^*)) \neq 0$  if and only if  $n = 2$  and  $p^r \mid i + 2p^s$  for some  $s \in \{0, \dots, r - 1\}$ , or  $n = 3$ ,  $p = 2$  and  $2^r \mid i - 2^{r-1}$ .

For  $G = \text{Sp}_n$ ,  $n \geq 4$  even, we deduce using  $S^i(V) = \nabla(i\varpi_1)$  and [2, Cor. 3.2(a)] that  $H^1(G_r, S^i V) \neq 0$  if and only if  $p = 2$  and  $i$  is odd.

Now let  $G$  be the special orthogonal group  $\text{SO}_n$ ,  $n \geq 4$ , as defined in [18, Ex. 7.4.7(3), (4), (6), (7)] (when  $p = 2$  this is an abuse of notation). Note that  $V \cong V^*$  unless  $n$  is odd and  $p = 2$ . Although the simply connected cover  $\tilde{G} \rightarrow G$  need not be separable, it still follows from [11, I.6.8(3), I.6.9(3)] that  $H^1(G_r, M) = H^1(\tilde{G}_r, M)^{T^r}$  for any  $G$ -module  $M$ , and  $H^1(B_r, M) = H^1(\tilde{B}_r, M)^{T^r}$  for any  $B$ -module  $M$ . So one has to pick out the weight spaces of the weights in  $p^r X(T) \subseteq p^r X(\tilde{T})$ . For  $n \geq 8$  it follows from [2, Cor. 3.2(a)] that  $H^1(\tilde{G}_r, \nabla(i\varpi_1)) = 0$  for all  $i \geq 0$ . For general  $n \geq 4$  we proceed as follows. From [2, Sect. 2.5–2.7] we deduce that all weights of  $H^1(B_r, i\varpi_1)$  are of the form  $i\varpi_1 + p^s \alpha$  for some  $s \in \{0, \dots, r - 1\}$  and some  $\alpha$  simple or “long” (i.e. there is a shorter root). Since such weights don’t occur in  $p^r X(T)$  for  $\text{SO}_n$ ,  $n \geq 4$ , we get that  $H^1(B_r, i\varpi_1) = 0$ , and therefore by (4.1)  $H^1(G_r, \nabla(i\varpi_1)) = 0$  for all  $i \geq 0$ . By [11, II.2.17,18]  $S^i(V^*)$  has a filtration with sections  $\nabla(i\varpi_1), \nabla((i - 2)\varpi_1), \dots$ . So  $H^1(G_r, S^i(V^*)) = 0$  for all  $i \geq 0$ .

The fact that the weights of  $H^1(B_r, i\varpi_1)$  have the form stated above can be seen more directly as follows. First one observes that 1-cocycles in the Hochschild complex of a  $U_r$ -module  $M$  can be seen as the linear maps  $D : \text{Dist}^+(U_r) \rightarrow M$  with  $D(ab) = aD(b)$  for all  $a \in \text{Dist}(U_r)$  and  $b \in \text{Dist}^+(U_r)$ . Here  $\text{Dist}^+(U_r)$  denotes the distributions without constant term, i.e. the distributions  $a$  with  $a(1) = 0$ . Then one shows that, outside type  $G_2$ ,  $\text{Dist}(U_r)$  is generated by the  $\text{Dist}(U_{-\alpha, r})$  with  $\alpha$  simple or long.<sup>(2)</sup>

<sup>(2)</sup> If  $p$  is not special in the sense of [9], then (also in type  $G_2$ )  $\text{Dist}(U_r)$  is generated by the  $\text{Dist}(U_{-\alpha, r})$  with  $\alpha$  simple.

It follows that  $H^1(U_r, M)$  is a subquotient of  $M \otimes \bigoplus_{\alpha, 0 \leq s < r} \mathfrak{u}_{-\alpha}^{*[s]}$ , the  $\alpha$  simple or long. Now use that, for  $M$  a  $B_r$ -module,  $H^1(B_r, M) = H^1(U_r, M)^{T_r}$ .

## BIBLIOGRAPHY

- [1] H. H. ANDERSEN & J. C. JANTZEN, “Cohomology of induced representations for algebraic groups”, *Math. Ann.* **269** (1984), no. 4, p. 487-525.
- [2] C. P. BENDEL, D. K. NAKANO & C. PILLEN, “Extensions for Frobenius kernels”, *J. Algebra* **272** (2004), no. 2, p. 476-511.
- [3] A. BOREL, *Linear algebraic groups*, second ed., Graduate Texts in Mathematics, vol. 126, Springer, 1991, xii+288 pages.
- [4] N. BOURBAKI, *Elements of Mathematics: Algebra II. Chapters 4–7*, Springer, 1990, Translated from the French by P. M. Cohn and J. Howie, vii+461 pages.
- [5] M. DEMAZURE, “Invariants symétriques entiers des groupes de Weyl et torsion”, *Invent. Math.* **21** (1973), p. 287-301.
- [6] S. DONKIN, “On conjugating representations and adjoint representations of semisimple groups”, *Invent. Math.* **91** (1988), no. 1, p. 137-145.
- [7] G. HOCHSCHILD, “Cohomology of restricted Lie algebras”, *Am. J. Math.* **76** (1954), p. 555-580.
- [8] J. E. HUMPHREYS, *Conjugacy classes in semisimple algebraic groups*, Mathematical Surveys and Monographs, vol. 43, American Mathematical Society, 1995, xviii+196 pages.
- [9] J. C. JANTZEN, “First cohomology groups for classical Lie algebras”, in *Representation theory of finite groups and finite-dimensional algebras (Bielefeld, 1991)*, Progress in Mathematics, vol. 95, Birkhäuser, 1991, p. 289-315.
- [10] ———, “Representations of Lie algebras in prime characteristic”, in *Representation theories and algebraic geometry (Montreal, PQ, 1997)*, NATO ASI Series. Series C. Mathematical and Physical Sciences, vol. 514, Kluwer Academic Publishers, 1998, Notes by Iain Gordon, p. 185-235.
- [11] ———, *Representations of algebraic groups*, second ed., Mathematical Surveys and Monographs, vol. 107, American Mathematical Society, 2003, xiv+576 pages.
- [12] ———, “Nilpotent orbits in representation theory”, in *Lie theory*, Progress in Mathematics, vol. 228, Birkhäuser, 2004, p. 1-211.
- [13] W. VAN DER KALLEN, “Longest weight vectors and excellent filtrations”, *Math. Z.* **201** (1989), no. 1, p. 19-31.
- [14] D. S. PASSMAN, *A course in ring theory*, The Wadsworth & Brooks/Cole Mathematics Series, Wadsworth & Brooks/Cole Advanced Books & Software, 1991, x+306 pages.
- [15] A. PREMÉT & D. I. STEWART, “Rigid orbits and sheets in reductive Lie algebras over fields of prime characteristic”, *J. Inst. Math. Jussieu* **17** (2018), no. 3, p. 583-613.
- [16] R. W. RICHARDSON, “The conjugating representation of a semisimple group”, *Invent. Math.* **54** (1979), no. 3, p. 229-245.
- [17] S. SKRYABIN, “Invariants of finite group schemes”, *J. Lond. Math. Soc.* **65** (2002), no. 2, p. 339-360.
- [18] T. A. SPRINGER, *Linear algebraic groups*, second ed., Progress in Mathematics, vol. 9, Birkhäuser, 1998, xiv+334 pages.
- [19] R. STEINBERG, “Regular elements of semisimple algebraic groups”, *Publ. Math., Inst. Hautes Étud. Sci.* (1965), no. 25, p. 49-80.
- [20] ———, “Torsion in reductive groups”, *Adv. Math.* **15** (1975), p. 63-92.

Manuscrit reçu le 19 février 2018,  
révisé le 28 avril 2018,  
accepté le 12 juin 2018.

Rudolf TANGE  
University of Leeds  
School of Mathematics  
LS2 9JT, Leeds (UK)  
R.H.Tange@leeds.ac.uk