

This is a repository copy of *Scenarios of land use and land cover change and their multiple impacts on natural capital in Tanzania*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/133225/>

Version: Accepted Version

Article:

Capitani, Claudia, van Soesbergen, Arnout, Mukama, Kusaga et al. (17 more authors) (2019) Scenarios of land use and land cover change and their multiple impacts on natural capital in Tanzania. *Environmental conservation*. pp. 17-24. ISSN: 0376-8929

<https://doi.org/10.1017/S0376892918000255>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Scenarios of land use and land cover change and their multiple impacts on natural capital in Tanzania

Journal:	<i>Environmental Conservation</i>
Manuscript ID	EC-17-06-141.R2
Manuscript Type:	Research Paper
Date Submitted by the Author:	05-Feb-2018
Complete List of Authors:	<p>Capitani, Claudia; University of York, Environment Department van Soesbergen, Arnout; United Nations Environment Programme World Conservation Monitoring Centre, ; King's College London, Mukama, Kusaga; WWF Tanzania Country, Forest Program; Tanzania Forest Service, Tanga City District Malugu, Isaac; WWF Tanzania Country, Forest Program Mbilinyi, Boniface; Sokoine University of Agriculture, Department of Agricultural Engineering and Land Planning Chamuya, Nurdin ; Ministry of Natural Resources and Tourism, Forest and Beekeeping Division Kempen, Bas; ISRIC World Soil Information Mant, Rebecca; United Nations Environment Programme World Conservation Monitoring Centre Malimbwi, Rogers; Sokoine University of Agriculture, Forest mensuration and Management Department Munishi, Pantaleo; Sokoine University, Forest Biology Njana, Marco; Sokoine University of Agriculture, National Carbon Monitoring Centre (NMC) Ortmann, Antonia; Food and Agriculture Organization of the United Nations Uganda Office Platts, Philip; University of York, Environment Runsten, Lisen; United Nations Environment Programme World Conservation Monitoring Centre Sassen, Marieke; United Nations Environment Programme World Conservation Monitoring Centre Sayo, Philippina; WWF Tanzania Country, Forest Program Shirima, Deo; Sokoine University of Agriculture, Department of Ecosystems and Conservation Zahabu, Eliakimu; Sokoine University of Agriculture Faculty of Forestry and Nature Conservation Burgess, Neil; United Nations Environment Programme World Conservation Monitoring Centre; University of Copenhagen, Center for Macroecology, Evolution and Climate, The Natural History Museum Marchant, Robert; York Institute for Tropical Ecosystem Dynamics, Environment Department</p>
Keywords:	REDD+, ecosystem services trade-offs, decision-making, participatory gis, carbon, biodiversity, water

1 Scenarios of land use and land cover change and their multiple impacts on natural
2 capital in Tanzania

3 CLAUDIA CAPITANI¹, ARNOOUT VAN SOESBERGEN^{2,3}, KUSAGA MUKAMA⁴,
4 ISAAC MALUGU⁴, BONIFACE MBILINYI⁵, NURDIN CHAMUYA⁶, BAS KEMPEN⁷,
5 ROGERS MALIMBWI⁸, REBECCA MANT², PANTELEO MUNISHI⁹, MARCO
6 ANDREW NJANA¹⁰, ANTONIA ORTMANN¹¹, PHIL PLATTS¹², LISEN
7 RUNSTEN², MARIEKE SASSEN², PHILIPPINA SAYO⁴, DEO SHIRIMA¹³,
8 ELIKAMU ZAHABU¹⁴, NEIL D. BURGESS^{2,15}, ROB MARCHANT¹

9

10 ¹ York Institute for Tropical Ecosystems, University of York, Environment,
11 Heslington, York, North Yorkshire, UK. Corresponding author contact:
12 claudia.capitani@york.ac.uk

13 ² UN Environment World Conservation Monitoring Centre, Cambridge,
14 Cambridgeshire, UK

15 ³ King's College London, London, London, UK

16 ⁴ WWF Tanzania Country, Forest Program, Dar es Salaam, TZ

17 ⁵ Sokoine University of Agriculture, Department of Agricultural Engineering and Land
18 Planning, Morogoro, TZ

19 ⁶ Ministry of Natural Resources and Tourism, Forest and Beekeeping Division, Dar
20 es Salaam, Dar es Salaam, TZ

21 ⁷ ISRIC World Soil Information, Wageningen, Gelderland, NL

22 ⁸ Sokoine University of Agriculture, Department of Forest mensuration and
23 Management, Morogoro, Morogoro, TZ

24 ⁹ Sokoine University of Agriculture, Forest Biology, Morogoro, Morogoro, TZ

25 ¹⁰ Sokoine University of Agriculture, National Carbon Monitoring Centre (NCCM),

26 Morogoro, Morogoro, TZ
27 11 Food and Agriculture Organization of the United Nations Uganda Office
28 Kampala, Kampala, UG
29 12 University of York, Department of Biology, Heslington, York, UK
30 13 Sokoine University of Agriculture, Department of Ecosystems and Conservation,
31 Morogoro, Morogoro, TZ
32 14 Sokoine University of Agriculture, Faculty of Forestry and Nature Conservation
33 Morogoro, Morogoro, TZ
34 15 University of Copenhagen, Center for Macroecology, Evolution and Climate, The
35 Natural History Museum, Copenhagen, DK
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 SUMMARY

52 REDD+ (reducing emissions from deforestation, and forest degradation, plus the
53 conservation of forest carbon stocks, sustainable management of forests, and
54 enhancement of forest carbon stocks, in developing countries) requires information
55 on land use and land cover changes (LULCC) and carbon emissions trends from the
56 past to the present and into the future. Here we use the results of participatory
57 scenario development in Tanzania, to assess the potential interacting impacts on
58 carbon stock, biodiversity and water yield of alternative scenarios where REDD+ is
59 effectively implemented or not by 2025, the green economy (GE) and the business
60 as usual (BAU) respectively. Under the BAU scenario, land use and land cover
61 changes causes 296 MtC national stock loss by 2025, reduces the extent of suitable
62 habitats for endemic and rare species, mainly in encroached protected mountain
63 forests, and produce changes of water yields. In the GE scenario, national stock loss
64 decreases to 133 MtC. In this scenario, consistent LULCC impacts occur within small
65 forest patches with high carbon density, water catchment capacity and biodiversity
66 richness. Opportunities for maximising carbon emissions reductions nationally are
67 largely related to sustainable woodland management but also contain trade-offs with
68 biodiversity conservation and changes in water availability.

69
70
71
72
73
74
75

76 INTRODUCTION

77 Many countries across the tropics face major challenges around meeting the needs
78 of rapid developing and growing populations, maintaining viable ecosystem services
79 while tackling the impacts of climate change through mitigation and adaptation
80 strategies. The REDD+ mechanism has been proposed as a climate change
81 mitigation framework with the potential for reducing greenhouse gas emissions while
82 addressing rural poverty and conserving forest biodiversity and ecosystem services
83 in the 2010 16th Conference of the Parties (COP 16) of the United Nations
84 Framework Convention on Climate Change (UNFCCC). The international
85 discussions on REDD+ evolved and diversified over time (Angelsen *et al.* 2012;
86 Pistorius 2012; Lund *et al.* 2016), delivering hope, discouragement, support and
87 criticism on its feasibility and capacity to provide win-win solutions to climate change
88 mitigation - while also contributing to livelihoods, sustainable development,
89 enhanced governance, and biodiversity conservation (Sunderlin *et al.* 2014;
90 Pasgaard *et al.* 2016; Turnhout *et al.* 2016; Loft *et al.* 2017).

91 Tanzania started its REDD+ readiness process in 2008 (Burgess *et al.* 2010; URT
92 2010). The readiness process set the foundations and tested the carbon emissions
93 monitoring, reporting and evaluation system (MNRT 2015). Tanzania also recently
94 submitted its Intended Nationally Determined Contributions to UNFCCC (URT 2015);
95 these give REDD+ related actions a central national role in both mitigation and
96 adaptation contributions to climate change and development of a low emission
97 growth pathway. More recently the country has submitted its Forest Reference
98 Emission Level (FREL) to UNFCCC, currently undergoing technical assessment,
99 which estimates annual deforestation rate at 580,000 ha year⁻¹ over the 2002-2013
100 period (URT 2017). Several factors drive deforestation either directly (e.g. demand

101 for farmland and biomass energy) or indirectly (e.g. high population growth rate,
102 governance weakness and unsecure land tenure (Burgess et al. 2010, Kweka et al.
103 2015).

104 The Norwegian government funded a series of REDD+ pilot projects in Tanzania,
105 which mainly focused on the local implementation of REDD+, in isolation from other
106 policy mechanisms (Blomley *et al.* 2015). Although useful, these local insights are of
107 limited use for scaling to the national context, or for creating long-term future
108 sustainable development strategies (Abidoye *et al.* 2015). A key part of the REDD+
109 mechanism in Tanzania is to estimate trade-offs between carbon emission reduction
110 and multiple co-benefits potentially achievable under REDD+, such as food and
111 energy provisions, water availability and biodiversity conservation in relation to
112 national development strategies (e.g. Tanzania Development Vision 2025, URT
113 2005). An initial assessment of potential REDD+ co-benefits in Tanzania (Miles *et al.*
114 2009; Runsten *et al.* 2013) has been followed by efforts to produce increasingly
115 specific and nation-based datasets, analyses (Augustino *et al.* 2014), scenarios
116 method (Capitani *et al.* 2016) and REDD+ Social and Environmental Safeguard
117 Standards (VPO 2013a). In this study, we present a quantitative evaluation of the
118 potential interacting impacts of two alternative socio-economic and land use and land
119 cover changes scenarios (LULCC) on carbon stock and two non-carbon forest
120 ecosystem services, biodiversity and water regulation. We analyse the spatial
121 distribution of potential win-win or conflicting outcomes from the two scenarios. Then,
122 we discuss the potential contribution of scenario analysis to the Forest Reference
123 Emission level reporting, and for identifying potential synergies or conversely
124 preventing unintended impacts, within the framework of the Tanzania national
125 climate change and development strategies and International pledges.

126

127 METHODS

128 Our study focused on the mainland of the United Republic of Tanzania, the largest
129 country in East Africa with a population of 44.9 million people (NBS & OGCS 2013).
130 Forests cover ca. 48.1 million hectares (Mha), corresponding to 55% of Tanzania
131 mainland (National Forest Resources Monitoring and Assessment, NAFORMA,
132 MNRT 2015). This figure is higher than estimates obtained from satellite data (38.3%
133 in 2010, MNRT 2013). In Tanzania forests are managed either in protected areas-
134 various designations comprising about half of the woody volume where forest
135 management ranges from total protection (e.g. nature reserves) to regulated
136 harvesting (e.g. forest reserves), or in 'village' and 'general land' (15.4 Mha, MNRT
137 2015). An estimated 4 Mha falls under community forest management regimes under
138 Participatory Forest Management (PFM, MNRT 2008).

139

140 Scenarios development

141 We developed land use and land cover changes scenarios for Tanzania to 2025
142 following four-steps within a mixed participatory and modelling scenario framework
143 (Supplementary 1.1) that engaged 240 stakeholders from civil society and authorities
144 at local, regional and national level (WWF 2015, Capitani *et al.* 2016). First we
145 broadly defined two alternative scenarios: the business as usual (BAU)- policies
146 framework, demand for commodities, and implementation of REDD+ follow the
147 current development trajectory, and the green economy (GE)- a shift toward
148 sustainable practices is envisaged for agriculture, forestry and energy sectors
149 supported by governance enforcement, effective REDD+ implementation, and
150 enhanced productivity. Then, regional stakeholders developed locally tailored,

qualitative and semi-quantitative scenarios trajectories, associated with specific spatial patterns and likelihood of LULCC. Next, LULCC scenarios were modelled by allocating demand for cultivated land and wood biomass according to LULCC likelihood spatial layers (Table S1), as expected by stakeholders and validated with secondary data. By using the national land use and land cover map for 2010 (MNRT 2013, Fig. S1a) as baseline and the World Database on Protected Areas (IUCN & UNEP-WCMC 01/2015), changes were modelled from specific land use and land cover classes to arable land (cultivation expansion), to mixed cultivated-wooded land (shifting cultivation), and to classes having lower tree cover and biomass without cultivation replacement (degradation, e.g. from closed woodland to bushland). Preliminary results were validated in a national level workshop in 2015 and refined thereafter to create the results presented here. The spatial resolution of scenario outputs was ca. 100 m. To maintain the local representativeness of change pressures in the national scale impacts assessment on carbon and non-carbon benefits, we applied a double resampling process that has reduced the accuracy of our analysis (see Discussion and Supplementary 1.2).

Carbon stock

Biomass carbon stock was estimated for the Tanzania mainland using a national dataset for above ground biomass (AGB, Ortmann 2014) based on NAFORMA forest inventory data, and from land-cover-specific ratios for below ground biomass (MNRT 2015), litter and deadwood biomass (Willcock *et al.* 2012). The wood dry matter biomass was converted to carbon by applying a 0.47 conversion factor, following the national protocol (URT 2017). Top soil organic carbon content for the 0-30cm layer was estimated by multiplying carbon concentration data from a national map

(Kempen *et al.* 2014) by the corresponding volume and bulk density obtained from the Soil and Terrain Database (SOTER) of Southern Africa (Dijkshoorn 2003). Both scenarios and the associated LULCC change imply C stock losses by 2025, though lower in the GE than in the BAU scenario (Capitani *et al.* 2016), reflecting the need of ensuring food and energy security, while allowing infrastructure development. For LULCC driven carbon stock changes estimate, the baseline (Fig. S2a) was created from biomass and top soil carbon datasets resampled from the original ca. 250-m resolution to ca. 100-m resolution by using the nearest neighbour method. We assumed that cultivation expansion depletes the five carbon pools, while shifting cultivation and degradation deplete the above ground and dead wood biomass only. For newly created cultivated land or shifting cultivation, carbon stocks in the scenarios were estimated as the average stock of the respective classes for the baseline. Carbon stock for degraded areas in the scenarios was estimated by decreasing the baseline biomass proportionally to the average biomass loss for the specific LULCC types expected in each pixel. Carbon stock changes were calculated as the pixel base difference between the baseline and the scenarios. The final results were then aggregated at 1-km resolution.

Biodiversity

We assessed the potential impacts of LULCC on biodiversity under the two scenarios focusing on terrestrial vertebrate species as derived from the IUCN Red List database (mammals, birds, amphibians and reptiles, IUCN, 2016 and BirdLife International & NatureServe 2015). Species sensitive to the modelled LULCC (hence LULCC-sensitive species) were selected following the IUCN classification of threats from cultivation expansion (threat class 2.1, 2.2.1), livestock rearing (class 2.3),

201 wood harvesting for energy and timber (class 5.3), fire (class 7.1), and urbanization
 202 (class 1) (see Salafsky *et al.* 2008). For every species, extent of occurrences (EOO)
 203 layers in Tanzania were clipped to the occupied habitats by matching the associated
 204 IUCN habitat classes with global cover land use types (Foden *et al.* 2013) and then
 205 with our reference land use and land cover classes to generate Extent of Suitable
 206 Habitat (ESH) polygons. We collected spatial distribution data and generated ESHs
 207 for 164 amphibians, 311 mammals, 58 reptiles, 1002 birds species on the Tanzanian
 208 mainland. Out of these 1535 terrestrial vertebrates, 177 are either classified by IUCN
 209 (2016) as endemic (127) or included in the IUCN categories 'Critically Endangered,
 210 Endangered and Vulnerable' (hence threatened species, 140) or both (90). We
 211 calculated ESH reduction in the two scenarios for LULCC-sensitive species, focusing
 212 on endemic species and threatened species with at least 1% of their range included
 213 on the Tanzania mainland. We calculated a spatially explicit biodiversity index
 214 prioritising species richness and rarity (BRRI, modified from van Soesbergen *et al.*
 215 2016, Fig. S2b,) across Tanzania at 1-km resolution, by summing over all occurring
 216 species in each grid-cell (richness) the ESH weighted by the species distribution
 217 range size in Tanzania and over the globe (rarity, see Supplementary 1.3 for
 218 equations).

219

220 Water yield

221 To assess the impacts of LULCC under the two scenarios on water yields we used
 222 the WaterWorld V2 (Mulligan 2013) model at a resolution of 1 km. WaterWorld is a
 223 fully distributed, process-based hydrological model that utilises remotely sensed and
 224 globally available datasets. Baseline climate data is based on a long term
 225 climatology from WorldClim (Hijmans *et al.* 2005). Land use and land cover in the

model is represented by fractional values for three functional vegetation types (tree, herb and bare). We calculated these fractional values for each land use class in the baseline and scenarios using the nearest mean fractional value for a group of cells of that class for MODIS VCF data for the year 2010 (DiMicelli *et al.* 2011) thus retaining variability within land use classes as well as within country. Calculations were made at the ca. 100-m scenario resolution by resampling the MODIS VCF data. Final baseline and scenario fractional vegetation maps were then aggregated to 1-km resolution and used to run the model. Changes in water yields under each scenario were analysed as changes in pixel based water balance in mm year⁻¹ between the baseline (Fig. S2c) and the scenarios.

Multi-dimensions scenarios assessment

We assessed spatial patterns of synergies and trade-offs between carbon stock, biodiversity and water yield changes in the two scenarios. We focused on LULCC subjected areas, though we acknowledge that impacts could also be reflected outside, particularly for water. Changes in the three dimensions compared to the baseline were standardised, based on the scenarios and baseline statistical distribution of each dimension, and merged into a composite Red-Green-Blue (RGB) plot. We defined as increasing impacts between the scenarios and the baseline the decline of C stock, of BRRI index, and either positive or negative changes in water yield diverging from 0. Here, we report and discuss trade-offs across scenarios by comparing high to low impacts on the three dimensions.

RESULTS

In the BAU scenario, cultivated land is expected to expand by 5.4 Mha (0.36 Mha

year⁻¹) by 2025 (Fig. S1b). In addition, shifting cultivation expands over 3.5 Mha (0.23 Mha year⁻¹) and degradation over 3.4 Mha (0.22 Mha year⁻¹) by 2025. In the BAU scenario, 11% of LULCC occur within protected areas, mainly in state managed forest reserves. In the GE scenario (Fig. S1c), cultivation expansion is reduced to 4.5 Mha (0.3 Mha year⁻¹) and degradation occurs over 3.6 Mha (0.23 Mha year⁻¹).

Carbon

In the BAU scenario, the envisaged land cover changes are estimated to result in ca. 296 million tonnes of carbon (MtC) national stock loss by 2025 compared to 2010. The countrywide estimated carbon stock loss in the GE scenario is ca. 133 MtC by 2025 (Fig. 1). In the GE scenario, 37 MtC avoided emissions within protected areas accounts for 23% of the emissions difference compared to the BAU scenarios. Countrywide the C stock changes mostly occur within open woodland in both scenarios, ranging between 58% (GE) and 65% (BAU) of total change (Table 1). Under the GE scenario, following forest protection and sustainable management enforcement LULCC are partially displaced to habitats with lower management safeguards, such as bushland, grassland and mangrove forests.

Biodiversity

In the BAU scenario 326 LULCC-sensitive species are impacted by habitat conversion; this includes 100 Tanzania endemic and 120 threatened species. In the BAU scenario the extent of suitable habitat (ESH) reduction averages 20% for the endemic species and 6.5 % for the 37 non-endemic threatened species. Under BAU six species (*Arthroleptis kutogundua*, *Afrixalus morerei*, *Churamiti maridadi*, *Galagoides rondoensis*, *Nectophrynoides laticeps* and *Nectophrynoides paulae*) lose

50% or more of their ESH. In the GE scenario, 317 LULCC-sensitive species are impacted by LULCC. The mean ESH reduction decreases to 4% for the 91 impacted endemic species and to less than 1% for the 36 non-endemic threatened species. The biodiversity richness and rarity index (BRRI) is highly variable across Tanzania, with the highest values mainly concentrated within the Eastern Arc Mountains (EAM) biodiversity hotspot (Meng *et al.* 2016, Fig. S2b). In both scenarios (Fig. 2), the highest potential impact in high BRRI areas occurs in mountain forest patches. Compared to the Ge scenario, in the BAU scenario BRRI losses were locally higher, due to larger habitat losses of LULCC-sensitive species, but the BRRI gains were slightly wider, due to generalist species expansion in habitats with reduced canopy compared to the baseline. In the GE scenario, BRRI losses extended in species-rich regions not exposed to LULCC in the BAU scenario.

Water yield

Changes in water yields, expressed as changes in water balance, are greater under the BAU scenario than the GE scenario, with a mean increase in water balance of 3.9 mm year⁻¹ (+2%) versus 1.9 mm year⁻¹ (+1%), across BAU and GE scenario respectively (Fig. 3). Under the BAU scenario, nearly 10% of the country sees a change in water balance of more than 50%, while under the GE scenario this is 6.2%. In both scenarios mountain and lowland forest and closed woodland face the most intensive changes in water balance (per hectare), but woodland and wetlands contribute the largest observed absolute change at national scale because they cover a much bigger area than forests. Increases in water yield are generally the result of land degradation, reducing the amount of water use by vegetation and thus increasing available water for runoff, more closely following the rainfall pattern. In

addition to water use by vegetation, trees can also play an important role in 'capturing' occult precipitation within cloud forests (Bruijnzeel *et al.* 2011) and favouring precipitation infiltration within miombo (Kashaigili & Majaliwa 2013). In the baseline this contributes up to 17% of the water balance in montane forested areas of the Eastern Arc, the northern volcanoes, and in the west near lake Tanganyika. Forest degradation in those areas therefore is more likely to result in a reduction in available water.

Multi-dimensions scenarios assessment

The simultaneous assessments of impacts of LULCC on carbon, biodiversity and water yield gives a complex pattern for both scenarios. Few land use patches show matching degrees of impact (e.g. either low or high impact in every variable); while in most areas LULCC generate different combinations of impact intensity (Fig. 4). In the BAU scenario, simultaneous high impacts in every dimension are mainly focused in protected forests and woodlands across EAM and south-western Tanzania (Fig. 4). In the GE scenario, 40% of LULCC are avoided, and simultaneous high impacts on carbon, biodiversity and water yield decrease. Increased impact on carbon, biodiversity and water yield is more frequent outside managed areas. In the GE scenario, about 19% of LULCC occur in different areas than in the BAU scenario (potential displacement). In about one-third of displaced LULCC areas, low impact on carbon is associated with high impact on either biodiversity or water yield.

DISCUSSION

Studies that assess potential future trade-offs and interactions between carbon and non-carbon benefits of natural habitat conservation are rare for East Africa (e.g. van

Soesbergen *et al.* 2016). Synergies and trade-offs between ecosystem services, as their provision and demand change (Locatelli *et al.* 2013), with simultaneous assessment of carbon and non-carbon benefits at large scale being highly challenging (Busch & Grantham 2013).

In the highly diverse landscape of Tanzania, under land change scenarios spatial patterns of impacts on carbon storage, biodiversity and water yield are not homogeneous. Consistent patterns are identifiable to some extent in relation to the different habitats and forest management regimes. In montane and lowland forests LULCC driven impacts are usually consistent and result in high carbon stock loss, biodiversity loss and water yield change. This increased water availability could benefit farmers locally, but could cause severe impacts downstream (e.g. Enfors & Gordon 2007; Kashaigili & Majaliwa 2013). In species-rich dry woodlands of north-eastern Tanzania LULCC impact is higher on biodiversity than on carbon stock. In addition, cultivated land expansion result in relatively low rates of carbon stock loss per unit area but are locally associated with cumulated water deficit, and increased irrigation demand. Site-specific trade-offs between carbon and non-carbon benefits impacts require joined up action by decision-makers, for example management interventions that link water provision with carbon storage.

Lessons for REDD+ implementation

The Tanzania National REDD+ Strategy (VPO 2013b) identifies three broad categories of REDD+ implementation actions: improved management and restoration of protection and production forest reserves, community-based forest management (including non-reserved areas), and plantation forestry. Our findings suggest that strictly protected forests conserve carbon, preserve biodiversity and

351 maintain the water catchment, albeit over relatively small areas. Sustainable
352 management of productive forests can support carbon emission reduction in the GE
353 scenario, but with trade-offs for biodiversity and water yield. Maximising the potential
354 benefits depends on the simultaneous enforcement of management and adequate
355 resolution of conflicts, while ensuring current and future human communities' needs
356 are met (Persha & Meshack 2015). Critical to REDD+ implementation is the risk of
357 avoiding deforestation leakage (Pfeifer *et al.* 2012). In the GE scenario, LULCC
358 impacts on biodiversity shift from rare forest species to species-rich communities in
359 semi-open habitats that have lower carbon value and hence of slightly lower priority
360 in Tanzania REDD+ framework. This suggests that ambitious REDD+ targets are
361 needed for carbon emission and habitat conversion reduction to meet biodiversity
362 conservation objectives in Tanzania.

363 Protected areas and community-based forest management areas alone are not
364 sufficient to achieve emission reductions required to fulfil the Tanzanian national
365 commitment (URT 2015), meanwhile ensuring food, water and energy security to the
366 increasing population. At the national scale in both scenarios most carbon stock
367 changes, as well as water yield and biodiversity disturbance, are anticipated in
368 general land, particularly focused along the commercial development corridors (e.g.
369 SAGCOT and Tanga). Addressing land and natural resource degradation outside
370 managed areas requires better integration of a landscape-centred REDD+ (Turnhout
371 *et al.* 2016), development (e.g. poverty reduction, food security and education) and
372 conservation policies based on broader consensus and engagement by a wide range
373 of actors that have political will and support from Government ministries, NGOs and
374 community based organisations.

375 The Forest Reference Emission Level (FREL) assessment for Tanzania estimated

ca. 58 MtCO₂ year⁻¹ emitted due to deforestation (URT 2017), comparable to ca. 61 MtCO₂ year⁻¹ estimated in the BAU scenario using the same deforestation definition, though a different methodology. Our demand driven LULCC scenarios provide a useful estimate on the magnitude of deforestation fraction not detectable from satellite images (Hojas-Gascon *et al.* 2015). The multi-dimensional quantitative assessment can contribute to ongoing national and international debates surrounding expectations for carbon and co-benefits values; these can be used to chart the triple wins or compounded losses of potential futures. The scenarios, and importantly the wider information behind these, can be used to support current negotiations of desirable or undesirable impacts across diverse beneficiaries of forest services, in relation to REDD+, the Intergovernmental Platform on Biodiversity and Ecosystem Services and the Sustainable Development Goals.

Caveats and limitations

As with all results from scenario analysis, our findings have inherent uncertainty. The presented results are not predictions but depict potential impacts within the range of our scenario trajectories. To maximise relevance and legitimacy, to represent multiple scale perspectives, interaction of key components of water, carbon and biodiversity, and to overcome consistent challenges of time series data quality and scarcity for Tanzania, we put great efforts in model and datasets customization.

However, the uncertainties generated by this approach should be considered when drawing conclusions from the presented results.

Dataset resampling has affected the accuracy of impacts spatial patterns and of the multi-dimension assessment at pixel level. The choice of indices also influenced the presented findings. For example, the adopted biodiversity index has the advantage

of being sensitive to LULCC. However, it does not consider other essential aspects of biodiversity (Supplementary 1.3) or interactions with other sources of disturbance (e.g. climate change disturbance (Foden et al. 2013). Prioritization of biodiversity and ecosystem services conservation should account for internal feedbacks characterised by connectivity and complementarity (Kukkala and Moilanen 2017), which are not captured by pixel-based analysis.

The selected thematic and temporal scopes influence our findings. Considering additional dimensions (e.g. social) and different impacts thresholds (e.g. negotiated amongst stakeholders) could change the outcomes of multiple co-benefits assessment. The limited scenarios temporal horizon was set to comply with tangible objectives such as the Tanzania Development Vision 2025 (URT 2005) and the REDD+ roadmap, but this could limit the scope for green development assessment. In respect to the relevance for supporting decision making, we successfully engaged with a broad range of stakeholders from across the country to co-produce scenarios, build local assessment capacity and consensus around the scenarios outputs. Such approaches need integrating into institutional frameworks to effectively influence policy formulation and implementation to mainstream biodiversity conservation and ecosystem services provision in future land use planning.

Supplementary material

For supplementary material accompanying this paper, visit www.cambridge.org/core/journals/environmental-conservation

Acknowledgements

We thank the 180 stakeholders engaged in the scenarios development across Tanzania regions and 60 nationally focused decision-makers and delegates of the

NGO community in Tanzania who provided validation and feedback at the national synthesis workshop. We especially thank staff within the WWF Tanzania office.

Financial support

This work was funded by the Norwegian Government International Climate and Forest Initiative through their Royal Embassy, and the Ministry for Foreign Affairs of Finland, the University of York and the UNEP-WCMC.

REFERENCES

- Abidoye B, Mungatana E, Mahlalela L, Sacolo T, Babalola F. (2015) Forest ecosystems in the transition to a green economy and the role of REDD + in the United Republic of Tanzania. UNEP 2015. eds. I Mulder, B Mbote, Nairobi, Kenya.
- Angelsen A, Brockhaus M, Sunderlin WD, Verchot LV (2012) Analysing REDD +. Challenges and choices. eds. A Angelsen, M Brockhaus, WD Sunderlin, LV Verchot. Bogor, Indonesia.
- Augustino S, Bowles P, Carr J, Cox N, Hicks C, Mant R, Mbilinyi B, Meng H, Ravilious C, Runsten L, Salvaterra T, Dos Santos S, Tognelli MF, Zahabu E (2014). Environmental and social spatial data to support REDD+ safeguards and planning in Tanzania. SUA, Morogoro, URT; UNEP-WCMC, Cambridge, UK; IUCN Global Species Programme, Cambridge UK and Washington DC, USA.
- BirdLife International, NatureServe (2015) Bird species distribution maps of the world. BirdLife International, Cambridge, UK and NatureServe, Arlington, USA.
- Blomley T, Edwards K, Kingazi S, Lukumbuzya K, Vesa L (2015) REDD + hits the ground REDD + hits the ground. ed. J Mayers. London, UK.
- Bruijnzeel LA, Mulligan M, Scatena FN (2011) Hydrometeorology of tropical montane cloud forests: emerging patterns. *Hydrological Processes* **25**: 465-498.

- 451 Burgess ND, Bahane B, Clairs T, Danielsen F, Dalsgaard S, Funder M, Hagelberg N,
 452 Harrison P, Haule C, Kabalimu K, Kilahama F, Kilawe E, Lewis SL, Lovett JC, Lyatuu
 453 G, Marshall AR, Meshack C, Miles L, Milledge SAH, Munishi PKT, *et al.* (2010)
 454 Getting ready for REDD+ in Tanzania: a case study of progress and challenges.
 455 *Oryx* **44**: 339–351.
- 456 Busch J, Grantham HS (2013) Parks versus payments: reconciling divergent policy
 457 responses to biodiversity loss and climate change from tropical deforestation.
 458 *Environmental Research Letters* **8**: 34028.
- 459 Capitani C, Mukama K, Mbilinyi B, Malugu IO, Munishi PKT, Burgess ND, Platts PJ,
 460 Sallu SM, Marchant R (2016) From Local Scenarios to National Maps: A
 461 Participatory Framework for Envisioning the Future of Tanzania. *Ecology and*
 462 *Society* **21**: 4.
- 463 Dijkshoorn JA (2003) SOTER database for Southern Africa. ISRIC – World Soil
 464 Information, Wageningen, NL.
- 465 DiMicelli CM, Carroll ML, Sohlberg RA, Huang C, Hansen MC, Townshend JRG
 466 (2011) Annual global automated MODIS vegetation continuous fields (mod44b) at
 467 250 m spatial resolution for data years beginning day 65, 2000 - 2010, collection 5
 468 percent tree cover. University of Maryland, College Park, MD, USA.
- 469 Enfors E I, Gordon LJ (2007) Analysing resilience in dryland agro-ecosystems: a
 470 case study of the Makanya catchment in Tanzania over the past 50 years. *Land*
 471 *degradation and development* **18**: 680-696.
- 472 Foden WB, Butchart SHM, Stuart SN, Vié J-C, Akçakaya HR, Angulo A, DeVantier L
 473 M, Gutsche A, Turak E, Cao L, Donner SD, Katariya V, Bernard R, Holland RA,
 474 Hughes AF, O'Hanlon SE, Garnett ST, Şekercioğlu ÇH, Mace GM, Hoffmann M, *et*
 475 *al.* (2013) Identifying the World's Most Climate Change Vulnerable Species: A

- 476 Systematic Trait-Based Assessment of all Birds, Amphibians and Corals. *PLoS ONE*
 477 **8**: e65427 URL
- 478 Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A (2005) Very high resolution
 479 interpolated climate surfaces for global land areas. *International Journal of*
 480 *Climatology* **25**: 1965–1978.
- 481 Hojas-Gascon L, Cerutti PO, Eva H, Nasi R, Martius C (2015) Monitoring
 482 deforestation and forest degradation in the context of REDD + Lessons from
 483 Tanzania. Bogor, Indonesia: CIFOR. DOI: 10.17528/cifor/005642
- 484 IUCN, UNEP-WCMC. The World Database on Protected Areas (WDPA). Cambridge,
 485 2015. Downloaded 01/2015. [www dataset]. URL <http://www.protectedplanet.net>.
- 486 IUCN 2016. The IUCN Red List of Threatened Species. Version 2016-3.
 487 Downloaded 05/2016. [www dataset]. URL <http://www.iucnredlist.org>.
- 488 Kashaigili JJ, Majaliwa AM (2013) Implications of land use and land cover changes
 489 on hydrological regimes of the Malagarasi river, Tanzania. *Journal of agricultural*
 490 *sciences and applications* **2**: 45-50
- 491 Kempen B, Kaaya A, Mhaiki CN, Kiluvia S, Ruiperez-Gonzalez M (2014) Modelling
 492 and mapping the topsoil organic carbon content for Tanzania. *Geophysical Research*
 493 *Abstracts* **16**: 10413.
- 494 Kukkala SA, Moilanen A (2017) Ecosystem services and connectivity in spatial
 495 conservation prioritization. *Landscape Ecology* **32**:5–14.
- 496 Locatelli B, Imbach P, Wunder S (2013) Synergies and trade-offs between
 497 ecosystem services in Costa Rica. *Environmental Conservation* **41**: 27–36.
- 498 Loft L, Pham TT, Wong GY, Brockhaus M, Le DN, Tjajadi JS, Luttrell C (2017) Risks
 499 to REDD+: potential pitfalls for policy design and implementation. *Environmental*
 500 *Conservation* **44**:44–55.

- 501 Lund JF, Sungusia E, Mabele MB, Scheba A (2016) Promising Change, Delivering
 502 Continuity: REDD+ as Conservation Fad. *World Development* **89**:124-139.
- 503 Meng H, Carr J, Beraducci J, Bowles P, Branch W, Capitani C, Chenga J, Cox N,
 504 Howell K, Malonza P, Marchant R, Mbilinyi B, Mukama K, Msuya C, Platts P, Safari
 505 I, Spawls S, Shennan-Farpon Y, Wagner P, Burgess N (2016) Tanzania's reptile
 506 biodiversity: Distribution, threats and climate change vulnerability. *Biological
 507 Conservation* **4**:72-82.
- 508 Miles L, Kabalimu K, Bahane B, Ravilious C, Dunning E, Bertzky M, Kapos V,
 509 Dickson B (2009) Carbon, Biodiversity and Ecosystem Services: Exploring Co-
 510 Benefits. Tanzania. Prepared by UNEP-WCMC, Cambridge, UK, Forestry and
 511 Beekeeping Division, Ministry of Natural Resources and Tourism, Dar es Salaam,
 512 URT. UN-REDD Programme, URT. [www document]. URL [http://www.unep-](http://www.unep-wcmc.org/climate/publications.aspx)
 513 [wcmc.org/climate/publications.aspx](http://www.unep-wcmc.org/climate/publications.aspx).
- 514 Ministry of Natural Resources and Tourism (MNRT) (2008) Participatory Forest
 515 Management in Tanzania: Facts and Figures. Dar es Salaam, URT.
- 516 Ministry of Natural Resources and Tourism (MNRT) (2013) Tanzania Mainland Land
 517 Use - Land Cover. Tanzania Forest Services Agency, Dar es Salaam, URT.
- 518 Ministry of Natural Resources and Tourism (MNRT) (2015) National Forest
 519 Resources Monitoring and Assessment of Tanzania Mainland. Dar es Salaam, URT.
- 520 Mulligan M (2013). WaterWorld: a self parameterising, physically based model for
 521 application in data-poor but problem-rich environments. *Hydrology Research* **44**:
 522 748–769.
- 523 National Bureau of Statistics (NBS), Office of Chief Government Statistician (OCGS),
 524 Zanzibar (2013) The 2012 Population and Housing Census: Population distribution
 525 by administrative area. Dar es Salaam, URT.

- Ortmann A. (2014) Estimating land cover changes and current carbon stock in Tanzania in the context of REDD+. Master thesis, Global Change Ecology Graduate Program (M.Sc.) at the University of Bayreuth within the Elite Network of Bavaria, DE.
- Pasgaard M, Sun Z, Müller D, Mertz O (2016) Challenges and opportunities for REDD+: A reality check from perspectives of effectiveness, efficiency and equity. *Environmental Science and Policy* **63**: 161–169.
- Persha L, Meshack C (2015) Is Tanzania's Joint Forest Management Program a Triple Win? Understanding Causal Pathways for Livelihoods, Governance and Forest Condition Impacts. 3ie Grantee Final Report. New Delhi: International Initiative for Impact Evaluation (3ie).
- Pfeifer M, Burgess ND, Swetnam RD, Platts PJ, Willcock S, Marchant R (2012) Protected Areas: Mixed Success in Conserving East Africa's Evergreen Forests. *PLoS ONE* **7**: e39337.
- Pistorius T (2012) From RED to REDD+: The Evolution of a Forest-Based Mitigation Approach for Developing Countries. *Current Opinion in Environmental Sustainability* **4**: 6 638–45.
- Runsten L, Ravilious C, Kashindye A, Giliba R, Hailakwahi V, Kashaga LRA, Khalid S, Mbilinyi B, Mwampashi Y, Pastory M, Mant R, Osti M, Crête P, Pekkarinen A, Leppanen M, Miles L (2013) Using Spatial Information to Support Decisions on Safeguards and Multiple Benefits for REDD + in Tanzania. UNEP-WCMC, Cambridge, UK. Ministry of Natural Resources and Tourism, Dar es Salaam, URT. [www document]. URL <http://www.un-redd.org/tabid/5954/Default.aspx>
- Salafsky N, Salzer D, Stattersfield AJ, Hilton-Taylor C, Neugarten R, Butchart SHM, Collen B, Cox N, Master LL, O'Connor S, Wilkie D (2008) A Standard Lexicon for

- 551 Biodiversity Conservation: Unified Classifications of Threats and Actions.
 552 *Conservation Biology* **22**: 897–911.
- 553 Sunderlin WD, Pratama CD, Bos AB, Avitabile V, Sills EO, Sassi C de, Joseph S,
 554 Agustavia M, Pribadi UA, Anandadas A (2014) REDD+ on the ground: The need for
 555 scientific evidence In *REDD+ on the ground: A case book of subnational initiatives*
 556 *across the globe*. eds. EO Sills, SS Atmadja, C de Sassi, AE Duchelle, DL Kweka,
 557 IAP Resosudarmo, WD Sunderlin, pp. 2-32. Bogor, Indonesia: CIFOR.
- 558 Turnhout E, Gupta A, Weatherley-Singh J, Vijge MJ, de Koning J, Visseren-
 559 Hamakers IJ, Herold M, Lederer M (2016) Envisioning REDD+ in a post-Paris era:
 560 between evolving expectations and current practice. *Wiley Interdisciplinary Reviews*
 561 *Climate Change* 8:e425.
- 562 United Republic of Tanzania (URT) (2005) Tanzania Development Vision 2025.
 563 Planning Commission, Dar es Salaam, URT. [WWW document]. URL
 564 <http://www.tzonline.org/pdf/theTanzaniadevelopmentvision.pdf>
- 565 United Republic of Tanzania (URT) (2010) Readiness Preparation Proposal (R-PP).
 566 Dar es Salaam, Tanzania. [www document]. URL
 567 http://www.forestcarbonpartnership.org/sites/forestcarbonpartnership.org/files/Documents/PDF/Jun2010/Tanzania-Revised_R-PP_main_document_V9-10.06.2010.pdf
- 568
 569 United Republic of Tanzania (URT) (2015) Intended Nationally Determined
 570 Contributions to the UNFCCC. Dar es Salaam, URT.
- 571 United Republic of Tanzania (URT) (2017) Tanzania's forest reference emission
 572 level submission to the UNFCCC. Dar es Salaam, URT.
- 573 van Soesbergen A, Arnell AP, Sassen M, Stuch B, Schaldach R, Göpel J, Vervoort
 574 J, Mason-D'Croz D, Islam S, Palazzo A (2016) Exploring future agricultural
 575 development and biodiversity in Uganda, Rwanda and Burundi: a spatially explicit

scenario-based assessment. *Regional Environmental Change*. doi:10.1007/s10113-016-0983-6

Vice President's Office (VPO) (2013a) The Tanzania REDD+ social and environmental safeguard standards, Final Draft Dar es Salaam, URT.

Vice President's Office (VPO) (2013b) National strategy for reduced emissions from deforestation and forest degradation (REDD+). Dar es Salaam, URT.

Willcock S, Phillips OL, Platts PJ, Balmford A, Burgess ND, Lovett JC, Ahrends A, Bayliss J, Doggart N, Doody K, Fanning E, Green J, Hall J, Howell KL, Marchant R, Marshall AR, Mbilinyi B, Munishi PKT, Owen N, Swetnam RD, *et al.* (2012) Towards Regional, Error-Bounded Landscape Carbon Storage Estimates for Data-Deficient Areas of the World. *PloS One* 7: e44795.

World Wildlife Fund (WWF) Tanzania Country Office (2015) WWF REDD+ pilot project. Final project report. WWF, Dar es Salaam, URT. [www document]. URL http://d2ouvy59p0dg6k.cloudfront.net/downloads/wwf__redd__final_project_report__10th_april_2015_1.pdf.

601 FIGURE LEGENDS

602 Figure 1. Changes in total carbon stock (carbon tonnes per hectare, $C\ t\ ha^{-1}$) in the
603 business as usual (BAU) and green economy (GE) scenarios across Tanzania by
604 2025.

605

606 Figure 2. Changes in biodiversity richness and rarity index (BRRI) for terrestrial
607 vertebrates in the business as usual (BAU) and green economy (GE) scenarios
608 across Tanzania by 2025. Negative and positive changes relates to prevalent losses
609 and gains of species suitable habitats, respectively.

610

611 Figure 3. Changes in water yields per year ($mm\ year^{-1}$) in the business as usual
612 (BAU) and green economy (GE) scenarios across Tanzania by 2025. In both
613 scenarios yield increment (blue shades) compared to the baseline is more frequent
614 than yield decrease (red shades).

615

616 Figure 4: Red-Green-Blue (RGB) plot of combined impacts on carbon stocks (black
617 to green), biodiversity (BRRI, black to red) and water yield (black to blue) under the
618 business as usual (BAU) and green economy (GE) scenarios across Tanzania by
619 2025. Areas mapped in black indicate low impact values and light colours high
620 impact values for all three dimensions. The three-dimensional legend is represented
621 in two visions at the bottom left of the figure. The upper vision shows, for each cube
622 face, the colour combinations of the three dimensions when one is at its maximum
623 value and the other two are varying. The lower vision shows, for each cube face, the
624 colour combination of the three dimensions when one is at its lowest value and the
625 other two are varying.

626 Table 1. Share of carbon (C) stock changes (million tonnes, Mt, and percentage of
 627 the total land cover class C stock, %) by different land cover classes in the business
 628 as usual (BAU) and green economy (GE) scenarios. Classes are grouped according
 629 to the national definition of forests, other wooded land and other land, URT 2017.
 630 Carbon stock losses in the BAU scenario within protected areas (PAs) are reported
 631 separately.

	BAU	BAU: losses within PAs	GE
	C Mt - (%)	C Mt - (%)	C Mt - (%)
Forests			
Mountain & lowland forest	6 (5)	4 (3)	<1 (<1)
Closed woodland	42 (7)	5 (1)	5 (1)
Open woodland	192 (20)	4 (2)	79 (8)
Mangrove forest	<1 (3)	<1 (<1)	<1 (5)
Thicket	3 (21)	<1 (<1)	0 (0)
Other wooded land			
Bushland	33 (9)	2 (<1)	39 (10)
Grassland	5 (3)	<1 (<1)	8 (6)
Other land			
Wetlands	15 (19)	2 (2)	5 (6)




Figure 1. Changes in total carbon stock (carbon tonnes per hectare, C t ha-1) in the business as usual (BAU) and green economy (GE) scenarios across Tanzania by 2025.

92x103mm (300 x 300 DPI)




Figure 2. Changes in biodiversity richness and rarity index (BRR) for terrestrial vertebrates in the business as usual (BAU) and green economy (GE) scenarios across Tanzania by 2025. Negative and positive changes relates to prevalent losses and gains of species suitable habitats, respectively.

93x105mm (600 x 600 DPI)




Figure 3. Changes in water yields per year (mm year-1) in the business as usual (BAU) and green economy (GE) scenarios across Tanzania by 2025. In both scenarios yield increment (blue shades) compared to the baseline is more frequent than yield decrease (red shades).

93x105mm (300 x 300 DPI)




Figure 4: Red-Green-Blue (RGB) plot of combined impacts on carbon stocks (black to green), biodiversity (BRRRI, black to red) and water yield (black to blue) under the business as usual (BAU) and green economy (GE) scenarios across Tanzania by 2025. Areas mapped in black indicate low impact values and light colours high impact values for all three dimensions. The three-dimensional legend is represented in two visions at the bottom left of the figure. The upper vision shows, for each cube face, the colour combinations of the three dimensions when one is at its maximum value and the other two are varying. The lower vision shows, for each cube face, the colour combination of the three dimensions when one is at its lowest value and the other two are varying.

99x119mm (600 x 600 DPI)

Supplementary material – (.pdf 722 Kb)

This Supplementary provides details on the scenario development framework; the baseline maps for carbon and non-carbon benefits assessment, along with caveats and potential sources of error in datasets manipulation; and, the biodiversity richness and rarity index.

1.1 Scenarios development framework

Our scenario development framework aimed to tackle the challenges of translating qualitative narratives into quantitative scenarios incorporating indigenous and local knowledge. Following a mixed participatory and modelling framework (Table S1), our approach allows translation of stakeholders’ developed qualitative and semi-quantitative scenarios trajectories and land use and land cover change patterns into quantitative and spatially explicit information.

Table S1. Steps of the participatory scenario development framework

Step 1 Scenarios definitions	Business as usual: policy framework, demand for commodities, and implementation of REDD+ follow the current development trajectory. Green economy: shift toward sustainable practices for agriculture, forestry and energy sectors supported by governance enforcement, effective REDD+ implementation, and enhanced productivity.
Step 2 Scenarios developm ent by stakeholde rs	a) Development of qualitative and semi-quantitative socio-economic and environmental trajectories of change and relative drivers by main livelihood sectors identified at regional level by multiple stakeholders. b) Identification of specific spatial patterns of land use and land cover changes (LULCC) related to expected trajectories and drivers of change (e.g. “ <i>high likelihood of conversion from closed woodland to</i>

	<i>grassland due to charcoal production near roads and in districts where governance is weak in region X").</i>
Step 3 Modeling	<p>a) Quantification of demand for cultivated land and wood biomass according to secondary data¹ and expected trajectories. In this study, the business as usual scenario refers to the BAU2 quantitative scenario detailed in Capitani et al. (2016; Appendix 2).</p> <p>Business as usual: 30% expansion for both cultivated and mixed cultivated-wooded land; pro-capita annual wood volume demand = 0.87 m³.</p> <p>Green economy: 10% increase in crop productivity no expansion of shifting cultivation; 50% reduction of wood biomass harvesting exceeding available sustainable cut.</p> <p>b) Spatial allocation of LULCC based on scalar composite indicators of likelihood of change calculated for different types of LULCC following the stakeholders' assessment and calculated from global and national reference datasets (corrected through locally obtained information when necessary)¹ according to the formula:</p> $SI_{lulcc} = (sp_1 + sp_2 + sp_3) \times m \times pas$ <p>SI_{lulcc}, composite indicators of likelihood of each specific LULCC; reclassified and standardized spatial datasets affecting LULCC likelihood (sp_n); $m = 0/1$ masking factor derived from crop suitability and slope to mask out unsuitable areas for cultivation expansion; pas, protected areas mask used to limit LULCC likelihood according to the rules: likelihood of LULCC occurring within protected areas decreasing</p>

	<p>with the distance from protected areas border in the BAU scenario (<i>pas</i> decreasing from 1 to 0); LULCC not occurring within protected areas in the GE scenario (<i>pas</i> = 0).</p> <p>Demand for land and for biomass is allocated through specific LULCC from the pixels with the highest likelihood of change until demand is fulfilled.</p>
Step 4 Iteration	Validation of preliminary results, feedback and synthesis workshop with regional and national stakeholders; model and outputs refinement.

¹ See Appendix 2 Capitani et al. 2016.

1.2 Scenarios and baseline maps

The scenario outputs (Fig. S1) were generated with a spatial resolution of ca. 100 m, in agreement with the population density dataset (WorldPop, Tatem 2017ⁱ), representing one of the major driving forces of land changes in our scenarios. Impacts from land use and land cover change scenarios in Tanzania on carbon, biodiversity and water yield were calculated using datasets derived from different inputs, at different resolution and with different methods (Fig. S2).




Figure S1. Land use and land cover reference map for 2010 (a, MNRT 2013) and for b) the business as usual and c) the green economy scenarios. Scenario output maps can be obtained upon request from the authors.




Figure S2 - Baseline maps for total carbon stock (a, ton ha^{-1}), biodiversity richness and rarity index of terrestrial vertebrates (b, range between 0 and 0.89) and water yield (c, mm year^{-1}) in Tanzania mainland. In b) the Eastern Arc Mountains biodiversity hotspot boundaries are represented by the purple line.

The high resolution adopted for the scenario analysis was helpful in incorporating local knowledge collected during the regional workshops, e.g. for simulating local patterns of small forest patches encroachment. To transfer the local representativeness of change pressures into the national scale impacts assessment

on carbon and non-carbon benefits, we altered the spatial resolution of the layers used to calculate carbon stock, biodiversity and water yield change, in order to match the ca. 100-m scenario resolution. Then we generalised the results at 1-km resolution. This double resampling process has determined a loss of accuracy in the analysis.

For biodiversity and water yield, the downscaling of the original input datasets at the scenario resolution was applied to match the reference habitat types and land cover classes with those used for the scenario analysis. Then the biodiversity and the water yield indices and their changes were calculated at 1-km resolution.

For carbon stock, the biomass and soil carbon stock layers were downscaled from ca. 250 to ca. 100 m resolution, to apply the change pressure on biomass and land determined by the specific land change expected in the scenarios (e.g. from forest to cultivated land, from closed woodland to bushland). Then changes were aggregate at 1-km resolution. The total amount of carbon biomass removed is upper limited by land and biomass demand set for the scenarios. However, the pixel-base allocation for the carbon stock change is influenced by the pixel-base carbon density, particularly for soil stock, and therefore is affected by the resampling process.

1.3 Biodiversity richness and rarity index

The Biodiversity richness and rarity index in the baseline $BRRI_{gt_0}$ was calculated for each grid-cell (g) by the formula:

$$BRRI_{gt_0} = \sum_1^i \left(\frac{ESH_{igt_0}}{ESH_{it_0}} \times R_i \right)$$

with ESH_{igt_0} the extent of suitable habitat of the i species in each pixel g , ESH_{it_0} the total extent of suitable habitat of the i species in Tanzania and R_i the ratio of the distribution range of the i species in Tanzania over the globe, at the time t_0 .

Changes between the scenarios and the baseline were calculated for each pixel (g)

$$BRRI_g = \sum_1^i \left(\frac{ESH_{igt_1} - ESH_{igt_0}}{ESH_{it_0}} \times R_i \right)$$

with ESH_{igt} the extent of suitable habitat of the i species in each pixel g in the scenario (t_1) or in the baseline (t_0), ESH_{it_0} the total extent of suitable habitat of the i species in Tanzania in the baseline and R_i the ratio of the distribution range of the i species.

When calculating the BRRI changes in the future scenarios we assumed that:

- LULCC-sensitive species abandon habitats converted to cultivated land or degraded;
- non-LULCC-sensitive species lose habitat due to conversion to cultivated land (e.g. species mainly associated with forest or closed canopy woodland or generalist species reported not to be tolerant to agriculture activities);
- non-LULCC-sensitive species mainly found in grassland can gain habitat following degradation of woodland and bushland, when degradation is above $15\text{m}^3 \text{ha}^{-1}$ wood biomass loss.

These rules are based on the reported habitat preference for the speciesⁱⁱ, on the

reference land use and land cover classes, and on the biomass changes calculated for the scenarios; gains are considered only within the extent of occurrence of each species. We did not consider other factors than habitat that could affect species capacity of moving or adapting to changes.

The adopted biodiversity richness and rarity index (BRRI) has the advantages of being calculated from data relatively easy to obtain on a large scale, and of being directly sensitive to LULCC, compared to other quantitative indices (e.g. species abundance, richness, diversity). However, it doesn't consider multiple aspect of biodiversity complexity, e.g. functional or taxonomic diversity, connectivity, complementarity, species adaptation capacity. In Tanzania the BRRI represents well the highly endemic montane forests and species-rich woodlands, and particularly emphasized the impacts of habitat changes on rare species. Using other indices, or other prioritisation approaches, different spatial pattern would emerge, e.g. weighting all species equally as in the species richness index.

ⁱ Tatem, A. J. (2017) WorldPop, open data for spatial demography. Sci. Data 4:170004 doi: 10.1038/sdata.2017.4

ⁱⁱ IUCN 2016. The IUCN Red List of Threatened Species. Version 2016-3. Downloaded 05/2016. [www dataset]. URL <http://www.iucnredlist.org>.