

This is a repository copy of *"Back-to-Front" Indole Synthesis using Silver(I) Catalysis: Unexpected C-3 Pyrrole Activation Mode Supported by DFT.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/133416/>

Version: Published Version

Article:

Clarke, Aimee K., Lynam, Jason M. orcid.org/0000-0003-0103-9479, Taylor, Richard John Kenneth orcid.org/0000-0002-5880-2490 et al. (1 more author) (2018) "Back-to-Front" Indole Synthesis using Silver(I) Catalysis: Unexpected C-3 Pyrrole Activation Mode Supported by DFT. *ACS Catalysis*. pp. 6844-6850. ISSN: 2155-5435

<https://doi.org/10.1021/acscatal.8b00745>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

“Back-to-Front” Indole Synthesis Using Silver(I) Catalysis: Unexpected C-3 Pyrrole Activation Mode Supported by DFT


Aimee K. Clarke, Jason M. Lynam,* Richard J. K. Taylor,* and William P. Unsworth*

University of York, Heslington, York, YO10 5DD, United Kingdom

Supporting Information

ABSTRACT: An efficient silver(I)-catalyzed method is reported for the synthesis of substituted indoles, most notably 5-hydroxy-derivatives, via π -acidic alkyne activation. Most methods for the preparation of indoles involve annulation of a benzene precursor, but the method reported herein is unusual in that pyrrole precursors are used. Density Functional Theory (DFT) studies suggest that these reactions proceed via initial activation of the pyrrole C-3 position before undergoing subsequent rearrangement, contradicting the conventional wisdom that pyrroles are more nucleophilic through C-2.

KEYWORDS: indoles, alkynes, silver(I), cyclization, spirocycles



INTRODUCTION

Indole derivatives are crucial in human biology (Figure 1); for example, tryptophan 1 is an essential amino acid, and serotonin




Figure 1. Biologically important indole derivatives.

2 and melatonin are neurotransmitters.^{1,2} Indoles are also ubiquitous in the plant world, and they are important components of agrochemicals, functional polymers/sensors, fragrances, and dyes. However, the significance of indole derivatives (especially 5-hydroxy indoles) is best highlighted by their biomedical applications (Figure 1).

As well as well-known natural products such as the anticancer vinca alkaloids (vincristine/vinblastine) and tran-

quilizing reserpine alkaloids, major indole-based drugs include pindolol (3, antihypertensive), ondansetron (4, antiemetic), Sumatriptan (5, migraine), indomethacin (6, arthritis), oxypertine (7, schizophrenia), and Arbidol (8, influenza).^{1,2} Furthermore, many more indoles with high pharmaceutical potential are in the pipeline, for example, evodiamine 9³ and dragsmacidin D.⁴

The range of applications of indole derivatives has stimulated much research into the search for efficient methods for their synthesis.^{1,5,6} As can be seen from Figure 1, versatile synthetic approaches are needed in terms of substituent variation. Such a synthetic challenge, with such valuable targets, ensures that novel approaches to indole synthesis are continually being reported.^{1,5–7} Of course, the Fischer indole synthesis,⁶ in which a phenylhydrazine is treated with an aldehyde or ketone, is still enormously important both in small scale and in industrial procedures. The vast majority of subsequently developed procedures also employ arylhydrazines, anilines, nitrobenzenes, or related benzene derivatives, and we have termed these the “standard” (or “front-to-back”) approaches, with many being named reactions (Scheme 1a).

In contrast, there are far fewer indole syntheses that commence from pyrrole or substituted pyrroles (Scheme 1b), and those that are known often afford little scope for substituent variation.^{8,9} The Muratake–Natsume indole synthesis (a Friedel–Crafts-type approach) is perhaps the only established named reaction of this type,^{8a} although a few other more recent procedures have also been reported.⁹


The research described herein concerns the development of new silver(I)-catalyzed, pyrrole-based routes to indoles, of the

Received: February 22, 2018

Revised: May 29, 2018


Published: June 13, 2018

Scheme 1. Standard (a) and “Back-to-Front” (b) Approaches to Substituted Indoles^{6–9}



rare “back-to-front” type, as outlined in Scheme 1b. We have recently developed a number of efficient metal-catalyzed ynone and ynol cyclization procedures for the synthesis of various cyclic scaffolds, including spirocyclic indolenines,^{10,11} other spirocycles,^{10a,d,12} carbazoles,^{10b,c} quinolines,^{10b,e} and quinolizidines.¹³ In this Article, two new silver(I)-catalyzed procedures for the conversion of ynols **11** into indoles **12**, and ynones **13** into 5-hydroxy-indoles **14**, are reported (Scheme 2). Both approaches can be used to prepare indoles

Scheme 2. “Back-to-Front” Ynol/Ynone Cyclization Approaches to Substituted Indoles




with a range of substitution patterns from closely related starting materials. Furthermore, they are simple to perform and use mild reaction conditions, and hence represent a practical and versatile new approach to these important scaffolds. The reactions were also studied using Density Functional Theory (DFT), and, intriguingly, these studies suggest that the reactions proceed via initial cyclization at the pyrrole C-3 position before undergoing subsequent rearrangement, contradicting the conventional wisdom that pyrroles are more nucleophilic through C-2.

RESULTS AND DISCUSSION


We started by examining the cyclizations of ynol derivatives **11a–f** (Scheme 3). The requisite starting materials (as well as ynones **13a–o** discussed later) were all prepared by standard literature procedures or minor variations thereof,¹⁴ with full experimental and characterization details included in the Supporting Information. First, it was found that pyrrole **11a** ($R^1 = R^2 = H$) could be converted into unsubstituted indole **12a** in reasonable yield upon stirring with 10 mol % $AgNO_3$ and 5 mol % Ag_2O for 24 h at rt in CH_2Cl_2 ; these conditions were chosen as they had been shown to be effective for the cyclization of indole-tethered ynols in our earlier work.^{10b} Monosubstituted indoles **12b** and **12c** were also formed in high yield using the same $AgNO_3/Ag_2O$ conditions. Dis-

Scheme 3. Ynol Cyclization to Indoles **12a–f**



substituted indole **12d** was also formed in high yield in the same way, but these conditions were less successful for the formation of disubstituted indoles **12e** and **12f** (just 28% and 19% conversion, respectively, under the same conditions). However, using a silica-supported $AgNO_3$ catalyst ($AgNO_3 \cdot SiO_2$), which was developed as part of our earlier work on alkyne activation,^{10d} the desired products were isolated in 43% and 99% yield, respectively. The role of Ag_2O in these reactions is not entirely clear; Ag_2O has been shown to improve reaction turnover when included as an additive in related $Ag(I)$ -mediated cyclization reactions^{10b} (as observed in this study), but it is not catalytically active in its own right; for example, ynol **11c** did not react when treated with 15 mol % Ag_2O under the standard reaction conditions. Thus, the role of this basic additive may instead be to modulate the acidity of the reaction (e.g., by quenching any adventitious protic acids present in the reaction), which may help prevent unwanted side reactions and/or catalyst degradation.^{10b} Alternatively, its basicity may help to drive the dehydration step.

Next, attention turned to the reactions of ynone derivatives of the form **13** (Scheme 4). These cyclization reactions are arguably of greater importance due to the fact that a 5-hydroxy substituent is produced in the product **14**; 5-hydroxylated indoles are ubiquitous in nature (e.g., serotonin **2**/indomethacin **6**), and this method could be used to generate analogues of these important compounds.¹⁵ The hydroxy group may also serve as a reactive handle to enable further chemical modifications, for example, alkylation, cross-coupling, etc. First, simple 5-hydroxy indole **14a** was formed in 64% yield using 10 mol % $AgNO_3$ as catalyst, again upon stirring for 1.5 h at rt in CH_2Cl_2 , with the Ag_2O additive not being necessary in this case. The Ag_2O additive was also not needed during the formation of disubstituted indoles **14b–i**, and the catalyst loading of $AgNO_3$ could also be lowered to 5 mol %. The yields were consistently high across this series, with aliphatic and electron-rich/electron-poor aromatic groups being well tolerated, as were tethered alkyl chloride and protected amine substituents. In the case of indole **14h**, its structural assignment is supported by X-ray crystallographic data.¹⁶ Finally, four trisubstituted indoles **14j–m** were also


Scheme 4. Ynone Cyclization to Indoles 14a–m^a

^aDouble the standard catalyst loading was required to achieve full conversion in this case.

generated; these substrates generally reacted more slowly than their disubstituted analogues, and some required the addition of Ag₂O to proceed to completion, but nonetheless each was formed in high yield under these modified conditions.

In terms of the mechanism for these cyclization reactions, two main pathways were considered, illustrated in Scheme 5

Scheme 5. Mechanistic Possibilities




for the ynone series. First, coordination of the alkyne to the active π -acidic silver species^{17,18} increases its electrophilicity and activates it toward attack from the electron-rich pyrrole ring, either via the pyrrole C-2 position (15 \rightarrow 16) or via its C-3 position (15 \rightarrow 17). Cyclisation via C-2 (route A) was considered to be the most likely at the outset of this study, in view of the well-accepted wisdom that pyrroles are more nucleophilic through C-2 than C-3, and the fact that this was judged to be the less sterically encumbered of the two

alternative pathways. Attack via C-2 is also the most direct route to the product, and thus from intermediate 16, protodemetalation and tautomerization completes the reaction to form indole 14. A second possibility, which was considered less likely in view of the above, but could not be ruled out, was that cyclization proceeds via C-3 (route B) to form a spirocyclic intermediate of the form 17, before undergoing 1,2-migration (17 \rightarrow 16), followed by protodemetalation and tautomerisation as in route A. For the ynol series, the same mechanism is likely to operate, except that the tautomerization step is replaced by the elimination of water (not shown).

However, subsequent examples led us to question our notion that C-2 cyclization is the more favorable pathway. Thus, when each of ynones 13n and 13o were reacted with 10 mol % AgNO₃ and 5 mol % Ag₂O at rt in CH₂Cl₂, none of the expected indole products 14n and 14o were formed, and instead spirocycles 18n and 18o were isolated (Scheme 6).

Scheme 6. Dearomatizing Spirocyclization Reactions of Ynones 13m and 13n




These results were surprising, not only because dearomatizing spirocyclization reactions through pyrrole C-3 positions are rare,^{19,20} but also because spirocyclization had taken place in preference to C-2 annulation. Resubjecting the spirocyclic products 18n and 18o to the reaction conditions led to no discernible reaction (when 1,2-migration to convert them into 14n and 14o, respectively, might have been expected), even when heated to reflux, attesting to their surprising stability.

To gain a better understanding of the potential mechanistic pathways underpinning the silver-catalyzed reactions, a number of possible pathways for C–C bond formation were probed with Density Functional Theory (DFT, see the Supporting Information for details of the computational methodology employed).²¹ Thus, one example from the “normal” indole-forming series (13e to 14e) was examined and compared to another (13n) in which the spirocycle 18n was formed unexpectedly. The calculations indicated that the Gibbs energies for the transformation of 13e to 14e and (putative) 18e (Scheme 7a) were both exergonic, but that indole 14e is the thermodynamic product of the reaction ($\Delta G_{298} = -223$ kJ mol⁻¹) and spirocycle 18e would be a kinetic product ($\Delta G_{298} = -58$ kJ mol⁻¹). Although employing the substituted pyrrole 13n changed the outcome of the experimental reaction, it did not significantly alter this picture with 14n being more stable than 18n, indicating that this reaction is under kinetic control.

Silver(I) complexes are kinetically labile, and in the case of simple salts such as AgNO₃ the precise nature of the active species is unclear. However, the reaction of AgOTf with 1 equiv of PPh₃ results in the formation of known complex Ag(OTf)(PPh₃).²² It was anticipated that the ynone substrate would displace the weakly bound OTf ligand in this complex

Scheme 7. DFT-Calculated Energies for (a) the Formation of Compounds 14e/n and 18e/n from 13e/n and (b) Silver-Catalyzed C–C Bond Formation from Alkyne Complex A^a



^aEnergies are Gibbs energies at 298 K at the D3-PBE0/def2-TZVPP//BP86/SVP(P) level with COSMO solvent correction in CH_2Cl_2 in kJ mol^{-1} .

to give a cationic silver(I) phosphine species that (1) would promote C–C bond formation and (2) could be effectively modeled by DFT. To demonstrate the feasibility of this approach, AgOTf (10 mol %) was mixed with PPh_3 (10 mol %) in CH_2Cl_2 solution to generate $\text{Ag}(\text{OTf})(\text{PPh}_3)$. Subsequent reaction with 13e for 3 h resulted in the formation of 14e in 77% isolated yield, demonstrating that this was indeed a viable catalyst system.

Therefore, the cyclization of ynones 13e and 13n was modeled by DFT using $[\text{Ag}(\text{PPh}_3)]^+$ as the catalyst system. The silver complex A, in which the alkyne is bound through the ynone oxygen atom, was taken as the reference state for the calculations (Scheme 7b). In the case of the cyclization reaction from 13e, the $\eta^2\text{-(CC)}$ bound alkyne complex Be lies at +2 kJ mol^{-1} , but C–C bond formation takes place from the slipped alkyne complex Ce, which may be viewed as being a formal carbocation. Two transition states for C–C bond formation were located. Transition state ts_{CD} is best viewed as nucleophilic attack by the C-3 atom of the pyrrole at the carbocation of C and leads to the silver-coordinated spirocycle De. A second transition state, ts_{CE} , was also located, which corresponds to C–C bond formation between the C-2 position of the pyrrole with the carbocation leading to Ee, and subsequent keto–enol tautomerization would then give Fe.


On the basis of these calculations, the potential energy surface is almost flat, with ts_{CD} and ts_{CE} at +18 and +28 kJ mol^{-1} , respectively. Although the formation of De will occur more rapidly than Ee based on these data, the equilibrium position between Ae, Be, Ce, and De is expected to favor Ae. Therefore, as Ee is of significantly lower energy (–32 kJ mol^{-1}), this would be the expected product from the reaction, as long as protodemetalation is slow. All attempts to find a transition state connecting De and Ee (i.e., a formal 1,2-migration) resulted in ts_{CD} .²³

In the case of 13n, the situation is somewhat different, as the corresponding complex Dn is now lower in energy than An, Bn,²⁴ and Cn. This may provide an explanation for the different reaction outcome in this case, as, in contrast to 13e, the complex Dn will dominate the equilibrium between the states A–D (i.e., it will be the major resting state prior to the slower formation of En) and hence should be more prone to protodemetalation to give the spirocyclic product 18n. This is consistent with 18n being a kinetic product formed from the lower lying transition state (ts_{CD}).

Examination of the predicted structure of De revealed that there are no elemental intramolecular steps that could lead to protodemetalation. Therefore, this step must involve an intermolecular deprotonation/protonation pathway; however,

modeling such steps is fraught with difficulty as the precise nature of the base is unclear. To obtain at least an indicative picture of this process, triflate was used as a model base. The neutral metalated spirocycle **Ge** and triflic acid were used as the reference point for the calculation (Scheme 8).

Scheme 8. DFT-Calculated Energies for Proton Transfer Reactions for Metalated Spirocycle^a



^aEnergies are Gibbs energies at 298 K at the D3-PBE0/def2-TZVPP//BP86/SVP(P) level with COSMO solvent correction in CH₂Cl₂ in kJ mol⁻¹.


Protodemetalation proceeds via an initial encounter complex **He** in which the HOTf has a weak interaction with the M–C bond. Proton transfer through TS_{HeIe} is essentially barrierless,²⁵ implying that protonation of the Ag–C bond by HOTf is rapid. Protonation ultimately leads to **Ie**, which has a T-shaped geometry with the silver coordinated to PPh₃, a triflate anion, and an O-bound spirocycle.

A further state, **Ke**, was also located, which corresponds to **De** with a hydrogen bonded triflate anion. This is significantly lower in energy than both the reference state and those involved in proton transfer. These data imply that the rate of the proton transfer reaction would be controlled by the deprotonation of **De** rather than the protonation of the M–C bond, at least in the case of HOTf.²⁶

Interestingly, the DFT results suggest that dearomatization through the pyrrole C-3 position (C to D) is lower in energy than dearomatization through the pyrrole C-2 position (C to E) in both ynone systems **13e** and **13n** (and, indeed, this is likely to be the case for most/all of the related ynone **13** in this study), but spirocyclic products are generally not isolated due to slow protodemetalation of the vinyl silver intermediate **D**. On the basis of this hypothesis, we reasoned that if the rate of protodemetalation of **D** could be increased relative to isomerization to **E**, this might enable C-3 spirocyclic products to be prepared by trapping intermediate **D**. To test this, we first attempted the reaction of ynone **13e** at lower temperature (0 °C), but this resulted in no reaction. Instead, the same reaction was then attempted using 5 mol % of the silica supported catalyst (AgNO₃·SiO₂) in place of AgNO₃ at rt; AgNO₃·SiO₂ was found to accelerate protodemetalation in our

earlier work on related indole spirocyclization reactions, and hence we predicted that the use of this catalyst might enable spirocyclic vinyl silver intermediate **De** to be trapped out prior to rearrangement.^{10d} Pleasingly, this experiment worked well; although the use of AgNO₃·SiO₂ did not lead to a complete switch in chemoselectivity, spirocycle **18e** was indeed formed, and isolated in 28% yield, alongside indole **14e**, which was formed in 38% yield (Scheme 9).²⁷ These results add strong

Scheme 9. Synthesis of Spirocycle **18e**



support to those obtained in the DFT study that spirocyclic vinyl silver species **De** is an intermediate in the synthesis of indole **14e**, and also show that under appropriate conditions, this intermediate can be intercepted, enabling synthetically challenging C-3 spirocyclic pyrrole derivatives to be prepared.

In summary, a new “back-to-front” method to generate substituted indoles from pyrrole ynone and ynone precursors has been developed. The synthetic procedures are promoted by relatively mild silver(I) catalysts, proceed at room temperature, and are insensitive to air and moisture. DFT studies conducted on the pyrrole-tethered ynone series (**13**) suggest that these reactions proceed via nucleophilic attack onto the activated alkyne via the pyrrole C-3 position, going against the generally accepted view that pyrroles are most nucleophilic through C-2. Formal C-2 annulation products are still formed in most cases (likely via ring opening and reclosing), but this discovery could be highly significant in the design of new dearomatizing spirocyclization methods for pyrrole precursors; methods to prepare C-3 spirocycles from pyrroles are very rare, but the knowledge that they are transiently formed en route to C-2 annulated products should help to inform new strategies by which the spirocycles can be intercepted prior to rearrangement.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acscatal.8b00745.

Full characterization data, images of NMR spectra and experimental and computational details for all of the materials produced in this study, as well as a link to the raw data (PDF)

■ AUTHOR INFORMATION

Corresponding Authors

*E-mail: jason.lynam@york.ac.uk.

*E-mail: richard.taylor@york.ac.uk.

*E-mail: william.unsworth@york.ac.uk.

ORCID

William P. Unsworth: 0000-0002-9169-5156

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We would like to thank the EPSRC (A.K.C. EP/R013748/1), the University of York (A.K.C., W.P.U.), and the Leverhulme Trust (for an Early Career Fellowship, ECF-2015-013, W.P.U.) for financial support. We also thank the EPSRC (grants EP/H011455/1 and EP/K031589/1) for funding the computational resources used in this study.

REFERENCES

- (1) For a comprehensive overview, see: *Comprehensive Heterocyclic Chemistry III*; Katritzky, A. R.; Ramsden, C. A.; Scriven, E. F. V.; Taylor, R. J. K., Eds.; Elsevier: Oxford, 2008, and references therein (especially Vol. 3; Ramsden, C. A.; Jones, G., Eds.).
- (2) For a review of the biomedical applications of indole derivatives, see: Kaushik, N. K.; Kaushik, N.; Attri, P.; Kumar, N.; Kim, C. H.; Verma, A. K.; Choi, E. H. Biomedical Importance of Indoles. *Molecules* **2013**, *18*, 6620–6662.
- (3) (a) Unsworth, W. P.; Kitsiou, C.; Taylor, R. J. K. Direct Imine Acylation: Rapid Access to Diverse Heterocyclic Scaffolds. *Org. Lett.* **2013**, *15*, 258–262. (b) Kitsiou, C.; Unsworth, W. P.; Coulthard, G.; Taylor, R. J. K. Substrate Scope in the Direct Imine Acylation of Ortho-Substituted Benzoic Acid Derivatives: the Total Synthesis (\pm)-Cavidine. *Tetrahedron* **2014**, *70*, 7172–7180 and references therein..
- (4) Zhang, F.; Wang, B.; Prasad, P.; Capon, R. J.; Jia, Y. Asymmetric Total Synthesis of (+)-Dragmacidin D Reveals Unexpected Stereo-complexity. *Org. Lett.* **2015**, *17*, 1529–1532 and references therein..
- (5) (a) Gribble, G. W. *Indole Ring Synthesis: From Natural Products to Drug Discovery*; Wiley: New York, 2016. (b) Gribble, G. W. J. Recent developments in indole ring synthesis—methodology and applications. *Chem. Soc., Perkin Trans. 1* **2000**, 1045–1075. (c) Inman, M.; Moody, C. J. Indole Synthesis – Something Old, Something New. *Chem. Sci.* **2013**, *4*, 29–41.
- (6) Robinson, B. *The Fischer Indole Synthesis*; Wiley: New York, 1982.
- (7) For recent approaches, see: (a) Zhang, Z.-Z.; Liu, B.; Xu, J.-W.; Yan, S.-Y.; Shi, B.-F. Indole Synthesis via Cobalt(III)-Catalyzed Oxidative Coupling of *N*-Arylureas and Internal Alkynes. *Org. Lett.* **2016**, *18*, 1776–1779. (b) Wu, C.-J.; Meng, Q.-Y.; Lei, T.; Zhong, J.-J.; Liu, W.-Q.; Zhao, L.-M.; Li, Z.-J.; Chen, B.; Tung, C.-H.; Wu, L.-Z. An Oxidant-Free Strategy for Indole Synthesis via Intramolecular C–C Bond Construction under Visible Light Irradiation: Cross-Coupling Hydrogen Evolution Reaction. *ACS Catal.* **2016**, *6*, 4635–4639. (c) Hermann, G. N.; Jungab, C. L.; Bolm, C. Mechanochemical Indole Synthesis by Rhodium-Catalysed Oxidative Coupling of Acetanilides and Alkynes under Solventless Conditions in a Ball Mill. *Green Chem.* **2017**, *19*, 2520–2523. (d) Tussing, S.; Ohland, M.; Wicker, G.; Flörke, G.; Paradies, J. Borane-Catalyzed Indole Synthesis through Intramolecular Hydroamination. *Dalton Trans.* **2017**, 46, 1539–1545. (e) Ilies, L.; Isomura, M.; Yamauchi, S.-I.; Nakamura, T.; Nakamura, E. Indole Synthesis via Cyclative Formation of 2,3-Dizincindoles and Regioselective Electrophilic Trapping. *J. Am. Chem. Soc.* **2017**, *139*, 23–26. (f) Agasti, S.; Deya, A.; Maiti, D. Palladium-Catalyzed Benzofuran and Indole Synthesis by Multiple C–H Functionalizations. *Chem. Commun.* **2017**, 53, 6544–6556.
- (8) (a) Muratake, H.; Mikawa, A.; Natsume, M. Total synthesis of (+)-Herbindole A, (+)-Herbindole B, and (+)-Herbindole C. Determination of the Absolute Configuration of the Natural Herbindoles. *Tetrahedron Lett.* **1992**, *33*, 4595–4598. (b) Kozikowski, A. P.; Sato, K.; Basu, A.; Lazo, J. S. Synthesis and Biological Studies of Simplified Analogs of Lyngbyatoxin A; use of an Isoxazoline-based Indole Synthesis. Quest for Protein Kinase C modulators. *J. Am. Chem. Soc.* **1989**, *111*, 6228–6234 and references therein. See also ref 5a, section 10..
- (9) (a) Jackson, P. M.; Moody, C. J. Diels-Alder Reactions of 2,3-Dimethylenepyrrole Analogues; a New Synthesis of Indoles. *J. Chem. Soc., Perkin Trans. 1* **1990**, 2156–2158. (b) Jackson, P. M.; Moody, C. J. Diels-Alder Reactions of 1,5-Dihydropyrano[3,4-b]pyrrol-5(1H)-ones, Pyrrole-2,3-quinodimethane Analogues; a New Synthesis of Indoles. *Tetrahedron* **1992**, *48*, 7447–7466. (c) Andrews, J. F. P.; Jackson, P. M.; Moody, C. J. Pyrrole-2,3-quinodimethane Analogues in the Synthesis of Indoles. Part 2.1 Synthesis and Diels-Alder Reactions of 1,6-Dihydropyrano[4,3-b]pyrrol-6(1H)-ones. *Tetrahedron* **1993**, *49*, 7353–7372. (d) Hashmi, A. S. K.; Yang, W.; Rominger, F. Gold(I)-Catalyzed Rearrangement of 3-Silyloxy-1,5-enynes: An Efficient Synthesis of Benzo[b]thiophenes, Dibenzothio-phenes, Dibenzofurans, and Indole Derivatives. *Chem. - Eur. J.* **2012**, *18*, 6576–6580. (e) Martins, G. M.; Zeni, G.; Back, D. F.; Kaufman, T. S.; Silveira, C. C. Expedient Iodocyclization Approach Toward Polysubstituted 3H-Benzo[e]indoles. *Adv. Synth. Catal.* **2015**, *357*, 3255–3261. (f) Li, Y.; Waser, J. Platinum-Catalyzed Domino Reaction with Benziodoxole Reagents for Accessing Benzene-Alkynylated Indoles. *Angew. Chem., Int. Ed.* **2015**, *54*, 5438–5442. (g) Kanchupalli; Joseph, D.; Katukojvala, S. Pyridazine N-Oxides as Precursors of Metallocarbenes: Rhodium-Catalyzed Transannulation with Pyrroles. *Org. Lett.* **2015**, *17*, 5878–5881.
- (10) (a) James, M. J.; Cuthbertson, J. D.; O'Brien, P.; Taylor, R. J. K.; Unsworth, W. P. Silver(I)- or Copper(II)-Mediated Dearomatization of Aromatic Ynones: Direct Access to Spirocyclic Scaffolds. *Angew. Chem., Int. Ed.* **2015**, *54*, 7640–7643. (b) James, M. J.; Clubley, R. E.; Palate, K. Y.; Procter, T. J.; Wyton, A. C.; O'Brien, P.; Taylor, R. J. K.; Unsworth, W. P. Silver(I)-Catalyzed Dearomatization of Alkyne-Tethered Indoles: Divergent Synthesis of Spirocyclic Indolenines and Carbazoles. *Org. Lett.* **2015**, *17*, 4372–4375. (c) Liddon, J. T. R.; James, M. J.; Clarke, A. K.; O'Brien, P.; Taylor, R. J. K.; Unsworth, W. P. Catalyst-Driven Scaffold Diversity: Selective Synthesis of Spirocycles, Carbazoles and Quinolines from Indolyl Ynones. *Chem. - Eur. J.* **2016**, *22*, 8777–8780. (d) Clarke, A. K.; James, M. J.; O'Brien, P.; Taylor, R. J. K.; Unsworth, W. P. Silica-Supported Silver Nitrate as a Highly Active Dearomatising Spirocyclisation Catalyst: Synergistic Alkyne Activation by Silver Nanoparticles and Silica. *Angew. Chem., Int. Ed.* **2016**, *55*, 13798–13802. (e) Liddon, J. T. R.; Clarke, A. K.; Taylor, R. J. K.; Unsworth, W. P. Preparation and Reactions of Indoleninyl Halides: Scaffolds for the Synthesis of Spirocyclic Indole Derivatives. *Org. Lett.* **2016**, *18*, 6328–6331.
- (11) For a review of spirocyclic indolenine synthesis, see: James, M. J.; O'Brien, P.; Taylor, R. J. K.; Unsworth, W. P. Synthesis of Spirocyclic Indolenines. *Chem. - Eur. J.* **2016**, *22*, 2856–2881.
- (12) (a) Unsworth, W. P.; Cuthbertson, J. D.; Taylor, R. J. K. Total Synthesis of Spirobacillene A. *Org. Lett.* **2013**, *15*, 3306–3309. (b) Clarke, A. K.; Liddon, J. T. R.; Cuthbertson, J. D.; Taylor, R. J. K.; Unsworth, W. P. Dearomatisation Approaches to Spirocyclic Dienones via the Electrophilic Activation of Alkynes. *Org. Biomol. Chem.* **2017**, *15*, 233–245.
- (13) (a) James, M. J.; Grant, N. D.; O'Brien, P.; Taylor, R. J. K.; Unsworth, W. P. Catalytic Dearomatization Approach to Quinolizidine Alkaloids: Five Step Total Synthesis of (\pm)-Lasubine II. *Org. Lett.* **2016**, *18*, 6256–6259. (b) Ho, H. E.; James, M. J.; O'Brien, P.; Taylor, R. J. K.; Unsworth, W. P. Ag(I)-Catalyzed Synthesis of Azabicyclic Alkaloid Frameworks from Ketimine-Tethered Ynones: Total Synthesis of Indolizidine 209D. *Org. Lett.* **2018**, *20*, 1439–1443.
- (14) (a) Trost, B. M.; Lumb, J.-P.; Azzarelli, J. M. An Atom-Economic Synthesis of Nitrogen Heterocycles from Alkynes. *J. Am. Chem. Soc.* **2011**, *133*, 740–743. (b) Yasuhara, A.; Takeda, Y.; Suzuki, N.; Sakamoto, T. Synthesis of 2,3-Disubstituted Indole Using Palladium(II)-Catalyzed Cyclization with Alkenylation Reaction. *Chem. Pharm. Bull.* **2002**, *50*, 235–238. (c) Ravinder, K.; Reddy, A. V.; Mahesh, K. C.; Narasimulu, M.; Venkateswarlu, Y. Simple and Selective Removal of the *t*-Butyloxycarbonyl (Boc) Protecting Group on Indoles, Pyrroles, Indazoles, and Carbolines. *Synth. Commun.* **2007**, *37*, 281–287. (d) Yadav, J. S.; Reddy, B. V. S.; Satheesh, G. InBr₃/Cu(OTf)₂-Catalyzed C-Alkylation of Pyrroles and Indoles with α -Diazocarbonyl Compounds. *Tetrahedron Lett.* **2003**, *44*, 8331–8334. (e) Bray, B. L.; Mathies, P. H.; Naef, R.; Solas, D. R.; Tidwell, T. T.; Artis, D. R.; Muchowski, J. M. *N*-(Triisopropylsilyl)pyrrole. A

Progenitor "Par Excellence" of 3-Substituted Pyrroles. *J. Org. Chem.* **1990**, *55*, 6317–6328.

(15) LaPorte, M.; Hong, K. B.; Xu, J.; Wipf, P. 5-Hydroxyindoles by Intramolecular Alkynol–Furan Diels–Alder Cycloaddition. *J. Org. Chem.* **2013**, *78*, 167–174 and references therein..

(16) CCDC 1554901 contains the crystallographic data for compound **14h**, see: www.ccdc.cam.ac.uk/data_request/cif (data of access: 9th June 2017).

(17) For silver(I)-mediated alkyne activation, see refs **10–13** and citations therein, as well as: Álvarez-Corral, M.; Muñoz-Dorado, M.; Rodríguez-García, I. Silver-Mediated Synthesis of Heterocycles. *Chem. Rev.* **2008**, *108*, 3174–3198.

(18) For related processes involving alkyne activation with other π -acidic catalysts, see: (a) Loh, C. C. J.; Badorrek, J.; Raabe, G.; Enders, D. Merging Organocatalysis and Gold Catalysis: Enantioselective Synthesis of Tetracyclic Indole Derivatives through a Sequential Double Friedel–Crafts Type Reaction. *Chem. - Eur. J.* **2011**, *17*, 13409–13414. (b) Peshkov, V. A.; Pereshivko, O. P.; Van der Eycken, E. V. Synthesis of Azocino[5,4-b]indoles via Gold-Catalyzed Intramolecular Alkyne Hydroarylation. *Adv. Synth. Catal.* **2012**, *354*, 2841–2848. (c) Heffernan, S. J.; Tellam, J. P.; Queru, M. E.; Silvanus, A. C.; Benito, D.; Mahon, M. F.; Hennessy, A. J.; Andrews, B. I.; Carbery, D. R. Double Gold-Catalysed Annulation of Indoles by Enynes. *Adv. Synth. Catal.* **2013**, *355*, 1149–1159. (d) Corkey, B. K.; Heller, S. T.; Wang, Y.-M.; Toste, F. D. Enantioselective Cyclization of Enamide-yne and Application to the Synthesis of the Kopsifoline Core. *Tetrahedron* **2013**, *69*, 5640–5646. (e) Gronnier, C.; Faudot dit Bel, P.; Henrion, G.; Kramer, S.; Gagosz, F. Divergent Product Selectivity in Gold- versus Silver-Catalyzed Reaction of 2-Propynyloxy-6-fluoropyridines with Arylamines. *Org. Lett.* **2014**, *16*, 2092–2095. (f) Xu, W.; Wang, W.; Wang, X. Gold-Catalyzed Cyclization Leads to a Bridged Tetracyclic Indolenine that Represses β -Lactam Resistance. *Angew. Chem., Int. Ed.* **2015**, *54*, 9546–9549. (g) Schroder, F.; Ojeda, M.; Erdmann, N.; Jacobs, J.; Luque, R.; Noel, T.; Van Meervelt, L.; Van der Eycken, J.; Van der Eycken, E. Supported Gold Nanoparticles as Efficient and Reusable Heterogeneous Catalyst for Cycloisomerization Reactions. *Green Chem.* **2015**, *17*, 3314–3318.

(19) For general reviews on dearomatization reactions, see ref **11**, as well as: (a) Zheng, C.; You, S. L. Catalytic Asymmetric Dearomatization by Transition-Metal Catalysis: A Method for Transformations of Aromatic Compounds. *Chem.* **2016**, *1*, 830–857. (b) Roche, S. P.; Youte Tendoung, J. J.; Tréguier, B. Advances in Dearomatization Strategies of Indoles. *Tetrahedron* **2015**, *71*, 3549–3591. (c) Zhuo, C. X.; Zheng, C.; You, S. L. Transition-Metal-Catalyzed Asymmetric Allylic Dearomatization Reactions. *Acc. Chem. Res.* **2014**, *47*, 2558–2573. (d) Zhuo, C.-X.; Zhang, W.; You, S. L. Catalytic Asymmetric Dearomatization Reactions. *Angew. Chem., Int. Ed.* **2012**, *51*, 12662–12686.

(20) For rare exceptions of C-3 pyrrole dearomatizing spirocyclization processes, see ref **10d**, as well as: (a) Shabalina, D. A.; Dvorkoa, M. Y.; Schmidt, E. Y.; Ushakov, I. A.; Protsuk, N. I.; Kobychiev, V. B.; Soshnikov, D. Y.; Trofimov, A. B.; Vitkovskaya, N. M.; Mikhaleva, A. I.; Trofimov, B. A. 3H-Pyrroles from Ketoximes and Acetylene: Synthesis, Stability and Quantum-Chemical Insight. *Tetrahedron* **2015**, *71*, 3273–3281. (b) Chambers, S. J.; Coulthard, G.; Unsworth, W. P.; O'Brien, P.; Taylor, R. J. K. From Heteroaromatic Acids and Imines to Azaspirocycles: Stereoselective Synthesis and 3D Shape Analysis. *Chem. - Eur. J.* **2016**, *22*, 6496–6500.

(21) For related DFT studies, see: Alcaide, B.; Almerndros, P.; Alonso, J. M.; Busto, E.; Fernández, I.; Ruiz, M. P.; Xiaokaiti, G. Versatile Synthesis of Polyfunctionalized Carbazoles from (3-Iodoindol-2-yl)butynols via a Gold-Catalyzed Intramolecular Iodine-Transfer Reaction. *ACS Catal.* **2015**, *5*, 3417–3421 and reference cited therein..

(22) (a) Bardaji, M.; Crespo, O.; Laguna, A.; Fische, A. K. Structural Characterization of Silver(I) Complexes $[\text{Ag}(\text{O}_3\text{SCF}_3)(\text{L})]$ ($\text{L} = \text{PPh}_3$, PPh_2Me , SC_4H_8) and $[\text{AgLn}](\text{CF}_3\text{SO}_3)$ ($n = 2-4$), ($\text{L} = \text{PPh}_3$, PPh_2Me). *Inorg. Chim. Acta* **2000**, *304*, 7. (b) Lettko, L.; Wood, J. S.;

Rausch, M. D. Synthesis of (Phosphine)silver(I) Trifluoromethanesulfonate Complexes and the Molecular Structure of Di- μ -trifluoromethylsulfonate-(tetrakis-triphenylphosphine)disilver(I). *Inorg. Chim. Acta* **2000**, *308*, 37.

(23) A similar stepwise rearrangement was described (with DFT support) for the overall 1,2-migration of aza-spiroindolenines, see: Wu, Q.-F.; Zheng, C.; Zhuo, C.-X.; You, S.-L. Highly Efficient Synthesis and Stereoselective Migration Reactions of Chiral Five-membered aza-Spiroindolenines: Scope and Mechanistic Understanding. *Chem. Sci.* **2016**, *7*, 4453–4459.

(24) (a) Ciano, L.; Fey, N.; Halliday, C. J. V.; Lynam, J. M.; Milner, L. M.; Mistry, N.; Pridmore, N. E.; Townsend, N. S.; Whitwood, A. C. Dispersion, Solvent and Metal Effects in the Binding of Gold Cations to Alkynyl Ligands: Implications for Au(I) Catalysis. *Chem. Commun.* **2015**, *51*, 9702–9705. (b) Rocchigiani, L.; Fernandez-Cestau, J.; Agonigi, G.; Chambrier, I.; Budzelaar, P. H. M.; Bochmann, M. Gold(III) Alkyne Complexes: Bonding and Reaction Pathways. *Angew. Chem., Int. Ed.* **2017**, *56*, 13861–13865.

(25) The difference in free energies of states **He** and **TS_{Hie}** was $<10 \text{ kJ mol}^{-1}$ at all levels of theory employed. Therefore, the apparent artifact in **Scheme 8** that **TS_{Hie}** is lower in energy than **He** may simply reflect the precision of the calculations.

(26) A transition state connecting **Ke** and **De** + HOTf could not be located.

(27) As was found for spirocycles **18n** and **18o**, spirocycle **18e** does not rearrange when resubjected to the reaction conditions (with both AgNO_3 and $\text{AgNO}_3 \cdot \text{SiO}_2$) or upon reflux in CH_2Cl_2 .