

This is a repository copy of *Investigating the function of prehistoric stone bowls and griddle stones in the Aleutian Islands by lipid residue analysis*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/133427/>

Version: Accepted Version

Article:

Admiraal, Marjolein orcid.org/0000-0002-9830-8032, Lucquin, Alexandre Jules Andre orcid.org/0000-0003-4892-6323, Von Tersch, Matthew et al. (2 more authors) (2018) Investigating the function of prehistoric stone bowls and griddle stones in the Aleutian Islands by lipid residue analysis. QUATERNARY RESEARCH. ISSN: 0033-5894

<https://doi.org/10.1017/qua.2018.31>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

1 **Title:**

2 Investigating the function of prehistoric stone bowls and griddle stones in the Aleutian Islands by lipid
3 residue analysis

4

5 **Authors:**

6 Marjolein Admiraal^a

7 Alexandre Lucquin^b

8 Matthew von Tersch^b

9 Peter D. Jordan^a

10 Oliver E. Craig^b

11

12 ^a Arctic Centre, University of Groningen, Aweg 30, 9718 CW Groningen, the Netherlands.

13 ^b BioArCh, University of York, Wentworth Way, YO10 5DD, York, United Kingdom.

ABSTRACT

The earliest durable cooking technologies found in Alaska are stone bowls and griddle stones recovered from the Aleutian Islands. This paper aims to identify the function of these artefacts. Molecular and chemical analysis of carbonised residues found on their surfaces confirm that these artefacts were used to process marine resources. Both artefacts have high lipid content and C:N ratios, suggesting they were used to process oily substances. Stable isotope results of individual lipids suggest that they were used to process different sets of resources within the aquatic spectrum as griddle stones have slightly more ^{13}C depleted lipids than stone bowls, possibly indicating more variable use. Integration of these results with archaeological and ethnographic data lead us to infer that griddle stones were used for the cooking of a diversity of aquatic resources, possibly with the addition of plant foods, whereas stone bowls were specifically used to render marine mammal fats. We further hypothesize that a sudden peak in stone bowl frequencies at 4000-3000 cal yr BP was connected to a Neoglacial cold spell bringing sea-ice conditions to the Aleutian Islands. This may have led to new subsistence strategies in which the rendering of marine mammal fats played a central role.

Keywords: durable container technologies, Aleutian Islands, stone bowls, griddle stones, oil rendering, cooking, organic residue analysis, lipids, compound specific isotopes, maritime adaptation, Neoglacial

INTRODUCTION

The use of durable container technologies in prehistory has often been connected to increasingly sedentary lifestyles, generally linked to agriculture. But it was not only the introduction of farming that led people to stay in one place. Seasonal abundance of aquatic resources in specific parts of the landscape can also facilitate increasing sedentism (Jordan and Zvelebil, 2009). The sub-Arctic Aleutian Islands are an ecological hotspot where early sedentism occurred based on the year-round abundance of marine resources. Through exploiting marine mammals and fish the Aleutian tradition grew out to become "...one of the world's most highly specialized and successful maritime hunter-gatherer adaptations" (Corbett and

Yarborough, 2016: p.607). Terrestrial resources were scarce with only a limited range of plant and terrestrial animals available. However, birds such as ptarmigan and waterfowl were abundant. Nevertheless, people focused their main efforts on the sea.

Heavy stone vessels such as bowls and flat cooking stones known as griddle stones were a technology central to this subsistence economy. The procurement, manufacture and maintenance of these tools required investment of time and effort (Jeanotte et al., 2012). But despite their apparent importance, the function of these artefacts remains unclear. Carbonised deposits on griddle stone surfaces, bowl rims and exteriors hint at the use of these artefacts as food processing tools using direct heating methods. Knecht et al. (2001: p.49) and Knecht and Davis (2008: p.73) suggested that stone bowls were used for the hot rendering of sea mammal oil, one of the most important commodities in the life of Northern peoples. However, this has never been tested. Little is known regarding the use of griddle stones although it has been suggested that they were used for cooking sea food (Jeanotte et al., 2012).

In this paper we aim to identify the function of stone bowls and griddle stones. The organic residues preserved on these artefacts offer the opportunity to identify different cooking and storage practices. Through organic residue analysis we test the hypotheses that 1) these artefacts were used for the processing of aquatic resources, and 2) stone bowls and griddle stones may have been used for different purposes. Building on our finds, our second aim is to explore why stone bowl frequencies peak so suddenly and to explore the role of climate change in the emergence and abundance of this artefact type.

Culture historical phasing

Humans first arrived in the Aleutian Islands around 9000 cal yr BP. Their subsistence practice is considered to have been focused on maritime resources despite the terrestrial character of their toolkit. Possibly these people were late Paleoarctic terrestrial game hunters that came to the Aleutian Islands using a route across landfast sea ice (Davis et al., 2016: p.293). Knecht and Davis (2001) divided the Anangula tradition in an early (9000-7000 cal yr BP) and a late stage (7000-4000 cal yr BP). It has been argued that an influx of Ocean Bay I people from Kodiak Island (Fig.1) around 7000 cal yr BP further added to the foundation of the specialized maritime adaptation that the Unanga people (better known as the Aleut) are known for (Dumond, 1977; Dumond and Bland, 1995). People were attracted to the region because of an abundance of fish (cod and halibut) and sea mammals such as: harbor seal, whales, porpoise, sea lion and sea otter but also a variety of bird species. The earliest stone bowls (n=2) and griddle stones (n=1) are found in low numbers at a few sites dating to the Early Anangula phase. They become more numerous during later phases.

The Margaret Bay phase (4000-3000 cal yr BP) is a period of both climatic and cultural change. Based on the faunal assemblages of the Margaret Bay (3800-3000 cal yr BP) and Amaknak Bridge (3500-2500 cal yr BP) sites, Davis (2001) and Crockford and Frederick (2007) argue for the presence of sea-ice in the region generated by the onset of the colder sea-surface temperatures of the Neoglacial. This induced marine productivity and new species appeared in the region such as walrus, ringed seal and polar bear. The new situation presented challenges and opportunities for the Unanga. Subsistence practices were adapted to the new circumstances and focused more on ice-bound marine mammal hunting (Knecht and Davis, 2008). Stone bowls peak during this phase (Table 1) with high occurrences at the Margaret Bay (n=434), and Amaknak Bridge (n=71) sites, which suggests the substantial importance of these artefacts in Aleutian daily life at these sites.

The Amaknak phase of 3000-1000 cal yr BP can be considered the start of the florescence of the Aleutian tradition (Davis et al., 2016: p.286) with a complex and varied toolkit representing the continuous further development of the long established maritime adaptation. Stone bowls seem to go out of use during this period (Davis, et al., 2016: p. 286), while the occurrence of griddle stones increases from this time onwards (Knecht and Davis, 2003; Jeanotte et al., 2012). These two technologies are hardly ever found together. Temperatures fluctuated and possibly influenced the human populations in the area. Colder temperatures led to increased marine productivity, which could have induced cultural expansion as suggested by Maschner (2016: p.340). During the Late Aleutian phase of 1000 to 2000 cal yr BP tensions rose along the Pacific coast of SW Alaska. Fortified sea stacks and refuge sites indicate warfare, possibly with the newly established Koniag tradition of Kodiak Island, but also among neighbouring Unanga groups (Davis et al., 2016: p.286).

MATERIALS AND METHODS

Stone Bowls

These heavy, non-portable artefacts are made of ground volcanic tuff and come in different textures and colours. Although no complete specimens have been recovered to date, (partial) reconstructions show that shapes varied from oval to rectangular and sizes range from 12 to 45 cm in diameter, and 3 to 12 cm in depth (Figs. 2 and 3). Bowls are distinguished from lamps mainly by their relative depth and base thickness. Where lamps are often shallow with a thick base, bowls have higher walls with a base that is always thinner than the walls and which allows for cooking using a direct heating source. Another distinction is the absence of a wick in bowls whereas some lamps have a raised platform for the wick. Stone bowls occur in large numbers during the colder Margaret Bay phase (4000-3000 cal yr BP). At the Margaret Bay site a total of 434 fragments were recovered, 75% of which dated to around 3300-3100 cal yr BP, the final phase of occupation. At the Amaknak Bridge site 71 fragments were found dating towards the very end of the phase around 2780 cal yr BP. Six fragments were reported from the base of the Chaluka mound dated 3700 cal yr BP (Denniston, 1966: p.108). A few fragments (n=6) were found at the

lower levels of the Tanaxtaxak site, also ascribed to the Margaret Bay phase based on artefact assemblage (Knecht and Davis, 2003: p.45). Stone bowls are scarce outside this period though a few older fragments were found at the earlier levels of Margaret Bay (Knecht et al., 2001) and at the Anangula Blade site (n=1) (McCartney and Veltre, 1996) and the Oiled Blade site (n=1) on Hog Island at 9000 cal yr BP (Knecht and Davis, 2001: p.273). With the abandonment of the Margaret Bay and Amaknak Bridge sites stone bowls also seem to disappear from the Aleutian Islands archaeological record (Knecht et al., 2001: p.49).

Griddle Stones

Referred to as “stone frying-pans” by (Jochelson, 1925: p.109), the presence of these grease-covered stone slabs goes back 9000 years in the Aleutians (Fig. 4b). No complete specimens of griddle stones are known. Like the stone bowls, they are all fragmented, perhaps fractured during use, or purposefully broken after their use-life was completed. Jeanotte et al. (2012) showed that at the ADK-011 site on Adak Island the majority of griddle stone raw material was carefully selected from a source some 5 km away from the site, while a lesser quality source was also available much closer to site. This indicates that these artefacts were not just flat stones selected randomly. Acquiring them would have been costly both in time and effort.

Despite the importance of these food processing techniques in the Aleutian subsistence economy, the subject has received little attention in current archaeological literature. Jeanotte et al. (2012) were the first to perform analysis on the residues associated with the griddle stones by using bulk carbon isotope analysis and visible/near infrared spectrometry but were not able to offer any specific identifications. Here we aim to investigate the function of stone bowls and griddle stones through the structural and isotopic analysis of lipids that are preserved in the greasy crusts on the artefact surfaces. This approach has been shown to be highly effective distinguishing marine and terrestrial products formed during the use of archaeological artefacts (Colonese et al., 2017; Craig et al., 2013; Farrell et al., 2014; Shoda et al., 2017).

Lipid Extraction of archaeological food crusts

Twenty charred surface residue samples of approximately 100mg were collected of stone bowls from the Margaret Bay (n=11), Amaknak Bridge (n=8), and Tanaxtaxak (n=1) sites. Where available multiple samples were taken to compare interior with exterior residues or base with rim residues. Most of the bowls however only had encrustations on the exterior. Charred surface deposits were also collected from eight griddle stones (~100mg). One sample dates to the Early Anangula phase (9000-8000 cal yr BP) Oiled Blade site, while the other sampled griddle stones were much younger with two specimens from the Tanaxtaxak site on Unalaska (around 500 cal yr BP) and five samples from the Ulyagan site on Carlisle Island, part of the Islands of the Four Mountains (around 400 cal yr BP). The Tanaxtaxak griddle stones were sampled on both sides for comparative reasons. Samples were acquired by scraping off surface residues using a sterile scalpel, and homogenized by grinding the samples to a fine powder using a mortar and pestle.

Approximately 20 mg of the sample was weighed out for lipid extraction using acidified methanol and following established protocols (Colonese et al., 2017; Papakosta et al., 2015). This approach has been extremely efficient in extracting lipids from carbonised deposits, especially where intact and partially degraded acyl lipids are unlikely to survive (Craig et al., 2007; Lucquin et al., 2016b).

One mL of methanol was added to the sample which was subsequently ultrasonicated for 15 min. Then 200 µL sulphuric acid (H₂SO₄) was added after which the samples were heated for 4 hours at 70°C. The samples were then centrifuged at 3000 rpm for 5 minutes. The supernatant was transferred to a sterile vial and then extracted three times by adding 2 mL of hexane, mixing, separating and removing the supernatant. The sample was neutralized by passing through a pipette with glass wool and potassium carbonate (K₂CO₃). Eventually the extracts were dried under a gentle stream of nitrogen (N₂) and an internal standard (10 µL C36 alkane) was added to all samples (lipid quantities ranging from 40 to 8600 µg/g) before further analysis by GC (gas chromatography), GC-MS (GC-mass spectrometry) and GC-c-IRMS (GC-combustion-isotope ratio MS). The majority of acid extracts was also silylated after acid

extraction by adding 100 µL of BSTFA (N,O-bis(trimethylsilyl)trifluoroacetamide) and heating the sample at 70°C for 60 min in order to determine the presence of dihydroxy acids (Hansel and Evershed, 2009).

Collagen extraction of archaeological bones

A selection of archaeological bone material from the Tanaxtaxak, Margaret Bay and Summer Bay sites as well as the Brooks River area on the Alaska Peninsula was collected to serve as collagen reference material for bulk isotope analysis. Species were: Fin whale (n=2), Porpoise (n=2), Right whale (n=2), Narwhal/Beluga whale (n=3) (all determined using ZooMS, courtesy of the University of York), and sea lion (n=2), seal (n=4), sea otter (n=2), eagle (n=1), bear (n=2), caribou (n=5), anadromous fish (n=2) and marine fish (n=5). Sampling was done by removing a small section of mechanically cleaned bone using a sterile Dremel saw.

Collagen of 32 bone samples was extracted using a modified Longin method (Brown et al., 1988). Samples (200-300 mg) were demineralized using 0.6 M hydrochloric acid (HCl) at 4°C for several days depending on the sample. Samples were rinsed with distilled water after demineralization. Then they were gelatinised with 0.001 M HCl at 80°C for 48 hours after which the samples were first filtered using Poly-ethylene Ezee filters (Elkay Laboratories Ltd., 9 mL, pore size 60-90 µm). Subsequently the samples were ultrafiltered (30 kDa, Amicon® Ultra-4 centrifugal filter units, Millipore, MA, USA). Finally the samples were frozen and lyophilised.

GC-MS

The equipment used for GC-MS analysis was an Agilent 7890A series chromatograph attached to an Agilent 5975C Inert XL mass-selective detector with a quadrupole mass analyser (Agilent technologies, Cheshire, UK). A splitless injector was used and kept at 300 °C. The GC column was inserted into the ion source of the mass spectrometer directly. The carrier gas used was Helium with a constant flow rate of 3 mL min⁻¹. The ionisation energy of the MS was 70eV and spectra were obtained by

scanning between m/z 50 and 800. A DB-5ms (5%-phenyl)-methylpolysiloxane column (30 m x 0.250 mm x 0.25 mm; J&W Scientific, Folsom, CA, USA) was used for scanning. The temperature was set at 50 °C for 2 min, then raised by 10 °C min⁻¹ until it reached 325 °C where it was held for 15 min.

All extracts were also analysed on a DB-23 (50%-Cyanopropyl)- methylpolysiloxane column (60 m x 0.250 mm x 0.25 mm; J & Scientific, Folsom, CA, USA) in SIM mode to identify isoprenoid fatty acids and ω - (o-alkylphenyl) alkanolic acids as aquatic biomarkers (Cramp and Evershed, 2014) and to resolve the mixture of phytanic acid diastereomers (Lucquin et al., 2016a). The temperature was set at 50 °C for 2 min, then raised by 10 °C min⁻¹ until it reached 100 °C, then raised by 4 °C min⁻¹ to 140 °C, then by 0.5 °C min⁻¹ to 160 °C, then by 20 °C min⁻¹ to 250 °C where it was maintained for 10 min. The first group of ions (m/z 74, 87, 213, 270) corresponding 4,8,12- trimethyltridecanoic acid (TMTD) fragmentation, the second group of ions (m/z 74, 88, 101, 312) corresponding to pristanic acid, the third group of ions (m/z 74, 101, 171, 326) corresponding to phytanic acid and the fourth group of ions (m/z 74, 105, 262, 290, 318, 346) corresponding to ω -(o-alkylphenyl) alkanolic acids of carbon length C16 to C22 were monitored, respectively. Helium was used as the carrier gas with a flow rate of 2.4 mL min⁻¹. The relative abundance of two diastereomers of phytanic acids was obtained by the integration of the ion m/z 101.

Bulk isotope analysis: carbon/nitrogen

Thirty-one surface residue samples of which 21 stone bowls, seven griddle stones and three lamps as well as 32 bone collagen samples were analysed by elemental analysis - isotope ratio mass spectrometry (EA-IRMS). The residue samples were ground into a homogenised powder. The residue and collagen samples were weighed out in duplicate into tin capsules (~0.9 mg). The bulk stable nitrogen ($\delta^{15}\text{N}$) and carbon ($\delta^{13}\text{C}$) isotope values were measured based on previously described methods (Craig et al., 2007). Precision of instrument on repeated measurement was $\pm 0.2\text{‰}$ (s.e.m.), $\delta^{13}\text{C}$, $\delta^{15}\text{N} = [(\text{R}_{\text{sample}}/\text{R}_{\text{standard}} - 1)] \times 1000$, where $\text{R} = {}^{13}\text{C}/{}^{12}\text{C}$ and ${}^{15}\text{N}/{}^{14}\text{N}$. Accuracy was determined by measurements of international standard reference materials within each analytical run. These were IAEA 600 $\delta^{13}\text{C}_{\text{raw}} = -27.69 \pm 0.02$, $\delta^{13}\text{C}_{\text{true}} = -27.77 \pm 0.04$, $\delta^{15}\text{N}_{\text{raw}} = 1.49 \pm 0.38$, $\delta^{15}\text{N}_{\text{true}} = 1.0 \pm 0.2$; IAEA N2 $\delta^{15}\text{N}_{\text{raw}} = 20.9$

± 0.33 , $\delta^{15}\text{N}_{\text{true}} = 20.3 \pm 0.2$; IA Cane, $\delta^{13}\text{C}_{\text{raw}} -11.76 \pm 0.10$; $\delta^{13}\text{C}_{\text{true}} = -11.64 \pm 0.03$. Data were normalised to these international standards. All samples with %N values below 1% and %C below 10% were excluded.

Gas chromatography combustion isotope ratio mass spectrometry

Eleven stone bowl and seven griddle stone samples were measured in duplicate for stable carbon isotope values of methyl palmitate ($\text{C}_{16:0}$) and methyl stearate ($\text{C}_{18:0}$) derived from precursor fatty acids by GC-c-IRMS, following existing procedure (Craig et al., 2012). The instrument used for the analysis was a Delta V Advantage isotope ratio mass spectrometer (Thermo Fisher, Bremen, Germany) linked to a Trace Ultra gas chromatograph (Thermo Fisher) with a GC Isolink II interface (Cu/Ni combustion reactor held at $1000\text{ }^{\circ}\text{C}$; Thermo Fisher) to oxidise all the carbon species to CO_2 . The carrier gas used was ultra-high purity grade helium with a flow rate of 2 mL min^{-1} and parallel acquisition of the molecular data was realised by deriving a small part of the flow to an ISQ mass spectrometer (Thermo Fisher). Samples were diluted in hexane and $1\text{ }\mu\text{L}$ of each sample was injected into DB-5MS ultra-inert fused-silica column ($60\text{ m} \times 0.25\text{ mm} \times 0.25\text{ }\mu\text{m}$; J&W Scientific). The temperature was set at $50\text{ }^{\circ}\text{C}$ for 0.5 min and raised by $25\text{ }^{\circ}\text{C min}^{-1}$ to $175\text{ }^{\circ}\text{C}$, then raised by $8\text{ }^{\circ}\text{C min}^{-1}$ to $325\text{ }^{\circ}\text{C}$ where it was held for 20 min. A clear resolution and a baseline separation of the analysed peaks were achieved. Eluted products were ionized in the mass spectrometer by electron impact and ion intensities of m/z 44, 45 and 46 were recorded for automatic computing of the $^{13}\text{C}/^{12}\text{C}$ ratio of each peak in the extracts. Computation was made with Isodat software (version 3.0; Thermo Fisher) and was based on comparisons with standard reference gas (CO_2) of known isotopic composition that was repeatedly measured. The results of the analysis were expressed in per mill (‰) relative to an international standard, VPDB.

The accuracy of the instrument was determined on *n*-alkanoic acid ester standards of known isotopic composition (Indiana standard F8-3). The mean \pm S.D. values of these were $-29.60 \pm 0.21\text{‰}$ and $-23.02 \pm 0.29\text{‰}$ for the methyl ester of $\text{C}_{16:0}$ (reported mean value vs. VPDB $-29.90 \pm 0.03\text{‰}$) and $\text{C}_{18:0}$ (reported mean value vs. VPDB $-23.24 \pm 0.01\text{‰}$) respectively. Precision was determined on a laboratory standard

mixture injected regularly between samples (28 measurements). The mean \pm S.D. value of *n*-alkanoic acid esters were $-31.65 \pm 0.27\text{‰}$ for the methyl ester of C_{16:0} and $-26.01 \pm 0.26\text{‰}$ for the methyl ester of C_{18:0}. Each sample was measured in replicate (average s.d. 0.07‰ for C_{16:0} and 0.13‰ for C_{18:0}). Values were also corrected subsequent to analysis to account for the methylation of the carboxyl group that occurs during acid extraction. Corrections were based on comparisons with a standard mixture of C_{16:0} and C_{18:0} fatty acids of known isotopic composition processed in each batch under identical conditions.

RESULTS

Lipid preservation

The preservation of organic residues on the artefacts is very good in general. Both griddle stones and stone bowls provided high quantities of lipids per sample. Thirty of 33 samples ranged from 400 to 8600 $\mu\text{g g}^{-1}$, indicative of exceptional preservation. The griddle stones ($n=8$, mean = 4200 $\mu\text{g/g}$), were richer in lipids than stone bowls ($n=22$, mean = 2477 $\mu\text{g g}^{-1}$). However, the majority of stone bowls sampled here are about 1500 years older than the griddle stones so this may be the result of degradation. Both of these artefact types contained a much greater amount of lipids than commonly found on charred deposits associated with ceramic cooking pots. For example, the mean lipid concentration from 14 charred deposits on pottery from the sub-Arctic Sakhalin Islands extracted under identical conditions was 298 $\mu\text{g g}^{-1}$ (Gibbs et al. 2017). Only two stone bowl samples showed lower lipid preservation with lipid quantities ranging from 40 to 130 $\mu\text{g g}^{-1}$ (Supplementary Table 1). The oldest stone bowl sample in the Aleutian Islands from the Anangula Blade site (Quimby, 1945; McCartney and Veltre, 1996) yielded no lipid biomarker results and based on the associated %N value of 0.71 we discarded this sample.

GC-MS analysis

Thirty of 33 samples of both the stone bowls (20 of 23) and the griddle stones (10 of 10) contained isoprenoid acids: TMTD (4,8,12-trimethyltridecanoic acid), pristanic acid (2,6,10,14-

tetramethylpentadecanoic acid), and phytanic acid (3,7,11,15-tetramethylhexadecanoic acid) (Fig. 5), as well as ω -(*o*-alkylphenyl) alkanoic acids (APAAs) of carbon length 16 to 22 (Fig. 7). These meet the established criteria for the identification of aquatic resources in archaeology (Hansel et al., 2004; Evershed et al., 2008; Hansel and Evershed, 2009; Lucquin et al., 2016a). Interestingly, APAAs are only formed during the prolonged heating of tri-unsaturated fatty acids at a temperature of at least 270°C and therefore the aquatic oils must have been heated on these artefacts presumably during their processing. These data rule out the contamination of degraded aquatic oils that may be present in the soils as these are unlikely to have been heated. Additionally, the presence of APAAs on these stone artefacts suggests that formation of these compounds is not necessarily dependent on the presence of a ceramic matrix as stated by Evershed, et al. (2008: p.111). To date no evidence of pottery has been found in the Aleutian Islands. Isoprenoid acids are degradation products of phytol, a constituent of chlorophyll, and occur widely in marine organisms. Phytanic acid also occurs in the tissues of ruminant animals. The contribution SSR:SRR diastereomers of phytanic acid (SSR%) provides a means to discriminate these sources (Lucquin et al., 2016a). As expected, because of the lack of ruminants in the area, the data obtained confirms the aquatic origin of phytanic acid in all the Aleutian samples as compared to modern references (Fig. 6).

Saturated fatty acids range from C₈ to C₃₂ and unsaturated fatty acids, even numbered from C_{16:1} to C_{24:1}, with some extending up to C_{26:1}. All samples contain longer chain FA and UFAs with the exception of the three badly preserved samples. Dicarboxylic acids (or diacids), most likely degradation products of unsaturated fatty acids, are also widely present in all samples ranging mostly from 7 to 15 carbon length (Fig. 5). Experimental work by Evershed et al. (2008) showed that diacids of carbon length 8 to 11 form during the heating of aquatic oils. Following derivatization of the acid extract with BSTFA, trace amounts of long chain *n*-alkanols (C₂₂ - C₃₂) were also present in many of the samples with an even number of carbon atoms. As these were found in trace amounts there is a possibility they are derived from the burial environment, as they are a common lipid component of soils (van Bergen et al., 1998), derived from wax compounds in higher plants. However, these compounds were much more abundant in the charred deposits from several of the griddle stones from the Ulyagan site, along with the matching distribution of

long chain fatty acids (C_{22} - C_{32}). In this case, it is conceivable that plant products were directly processed on these artefacts.

Stable isotope analysis of individual fatty acids

Based on the lipid residue analysis results, integrated with contextual archaeological information of the materials, it seems very likely that the tested artefacts were used for the processing of aquatic resources. But what kind of aquatic resources were processed? Were the different artefacts used for different purposes? To further differentiate within the aquatic spectrum we analysed stable isotopes of individual fatty acids C_{16} and C_{18} using GC-c-IRMS. This approach serves as a means to discriminate aquatic animals based on their habitat (marine, anadromous, and freshwater), with the marine species relatively enriched in ^{13}C compared to the others (Fig 8).

The carbon isotope values show some difference between the two technological groups (Fig. 8). In general griddle stones are more depleted in both $\delta^{13}C_{16}$ and $\delta^{13}C_{18}$ than stone bowls. Two stone bowls are more depleted than the others. Of the seven tested griddle stones, five are separated from the stone bowls and two have similar fatty acid isotope values to the stone bowls. One of these is the only exceptionally old specimen from the Early Anangula phase (9000-8000 cal yr BP) site of Oiled Blade (UNL-318). It is unlikely that the isotopic approach deployed here can be used to distinguish between marine mammal and marine fish oil in this case. Although no authentic lipid carbon isotopes values have been measured from the Aleutian Islands, the $\delta^{13}C$ values of collagen extracted from 17 marine mammals (mean $\delta^{13}C_{coll} = -14.47 \pm 0.04$) and five marine fish (mean $\delta^{13}C_{coll} = -11.96 \pm 0.03$) from SW Alaska (Supplementary Table 2) are similar. More depleted lipid sources could include salmon ($\delta^{13}C_{coll} = -15.44 \pm 0.05$) or potentially terrestrial resources including plants. The latter would be consistent with degraded wax esters found on the griddle stones. However, we stress that overall both the stone bowls and griddle stones have strong marine isotope signatures and aquatic lipid profiles, so any other products are only a minor component in these residues.

Bulk $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ isotopes

The $\delta^{15}\text{N}$ ratios of the carbonised deposits associated with the stone bowls and griddle stones are generally within the range expected for marine tissues (i.e. $>10\text{‰}$; Fig 9). These values can be tentatively compared to $\delta^{15}\text{N}$ values of collagen from associated fauna (fish, marine mammals, terrestrial fauna) by assuming that any of the $\delta^{15}\text{N}$ in the surface residues are derived from animal tissue protein and the $\Delta^{15}\text{N}_{\text{tissue-collagen}} = \text{ca. } +2\text{‰}$ (Fernandes et al., 2015). Interestingly, the range of $\delta^{15}\text{N}$ values observed in the carbonised deposits is at the lower end of the marine mammal and fish range (Supplementary Table 1).

Compared to pottery vessels from the Sakhalin Islands, which are assumed to have been used for cooking a range of marine tissues (Gibbs et al. 2017), all the Aleutian artefacts have higher C:N ratios, indicative of a relatively higher lipid content. These data are more comparable to the so called ‘blubber lamps’ of the European Mesolithic where it is thought that marine mammal oil was burned for illumination (Heron et al., 2013). One caveat to this interpretation is that extensive microbial degradation or percolation with groundwater might lead to a preferential loss of protein, effectively increasing the C:N ratio (Heron and Craig, 2015) although this would seem less likely given the environment is so conducive to molecular preservation. The $\delta^{13}\text{C}$ values in all but one case are less than -25‰ (Supplementary Table 1) which are consistent with values reported from pottery from coastal sites with clear marine lipid signatures (Craig et al., 2011, 2013).

DISCUSSION

Our main question centred on the specific function of griddle stones and stone bowls within the Aleutian subsistence economy. We hypothesized that these artefacts were used to process aquatic resources but were used in different ways. The results of this research cautiously support our hypotheses. All artefacts with sufficient preservation ($n=30$) show strong evidence for the processing of aquatic resources. Minor isotopic differences between stone bowls and griddle stones may indicate a more variable use of the

latter. However, the lipid concentrations and C:N ratios from both artefacts are consistent with their use for processing aquatic oils and fats. By integrating the organic residue results with information of archaeological contexts, ethnography and climate, we discuss the possible function and role of these artefacts placed in a framework of the wider subsistence strategies of the ancient Unangaʼ.

Griddle stone function

Griddle stone charred residues show a clear aquatic signal. This is visible in the presence of all aquatic biomarkers as well as in compound specific-, and bulk isotope results. We hypothesized that griddle stones had a more general use for cooking foodstuffs. The residue results are cautiously supportive of this notion. Interpretation is based on comparisons with stone bowls that we assume were used for a very specific purpose, namely the rendering of marine oil. The residues on griddle stones seem to be derived from a bigger diversity of resources. Despite this, aquatic oils still make up the majority of the sample. Depleted $\delta^{13}\text{C}$ values may indicate the contribution of salmon and plant products to the sample. Furthermore the presence of *n*-alkanols on the Ulyagan griddle stones supports the possibility that plant products contributed to the otherwise predominantly aquatic sample. The lower C:N ratio values attest to a higher presence of proteins possibly caused by the cooking of flesh as opposed to fats.

Not all griddle stones show the same consistent residue results. We analysed the earliest griddle stone in the Aleutian Islands, a partial specimen from the Oiled Blade (UNL-318) site, dating to the Early Anangula phase at 9000-8000 cal yr BP. The compound specific $\delta^{13}\text{C}$ isotope data and C:N ratio value of this particular griddle stone are closer to stone bowl values. This possibly indicates a less diverse use on this ancient griddle stone as opposed to griddle stones from later periods.

Ethnographic resources are of great value when considering function because griddle stones were still widely in use by the Unangaʼ during early contact times. A report by C.I. Shade describes the traditional Unangan way to prepare cod soup using a griddle stone: *“The traditional method of making soup was to dig a fire pit and place over it a stone, flush with the ground. Then a very thin beach stone was placed on*

the fire stone and clay walls built upon this base. The liquid was cooked in this. A bluish clay called qudii u was used for the walls of this vessel which turned white when heated. This kind of fire pit was called unaalu. The same vessel was used more than once. One way of preparing the cod soup was with seaweed and seal oil" (in: Johnson, 2004: p.52).

This is an interesting notion suggesting griddle stones were actually also used as containers. It also attests to the use of marine mammal oil in cooking practices, agreeable with our findings. No evidence for the use of clay has ever been detected in the archaeological record of the Aleutians. However, some griddle stones are clean in the centre and have a thick edge of greasy and carbonized material around this clean area (Fig. 4a). This use-wear pattern could represent the process described above. Not all griddle stones show this residue distribution though, some show residues in the centre (Fig. 4b). This may suggest different methods of cooking with the use of griddle stones.

Use patterns possibly changed through time with the earlier griddle stone used for the processing of a single commodity, while griddle stones of the Late Aleutian phase were probably used to cook dishes of a more diverse character, although still predominantly aquatic.

Stone bowl function

The charred residues distributed mainly along the rims on both interior and exterior, but also on the bases of the bowls, seem to be solely aquatic in origin. The stone bowls have comparable lipid distribution, C:N ratios and bulk isotope characteristics to prehistoric oil lamps from Europe that were unequivocally used to burn aquatic oils for fuel (Heron et al., 2013; Heron and Craig, 2015; Piezonka et al., 2016; Oras et al., 2017).

Aquatic oil played an important role in the lives of prehistoric and historic peoples of the (sub)Arctic. Not only was the substance used as a fuel to burn in oil lamps, it is known to be an important part of the Unanga diet (Unger, 2014), and critical for the storage of various foodstuffs (Frink and Giordano, 2015).

Knecht (2001, 2008) has suggested multiple times that stone bowls were used for the purpose of rendering aquatic oils. And despite the absence of stone bowls in archaeological sites dating to later phases, one ethnographic source refers to an artefact used during contact times which description sounds remarkably like that of a stone bowl: *"The stone of which these lamps are made is very soft, and may be hollowed out with others of greater hardness, not merely for this purpose, but also for deep pots, in which they boil their fish. They use them however, but seldom, preferring mostly the iron and copper kettles, which they procure from the Russians"* (Sarychev, 1806: p.73).

Another argument supporting the use of stone bowls for the rendering of aquatic oils is the residue distribution on the rims but not on the bottom (Fig. 2). Oil may have been rendered by placing cuts of fat in boiling water. The rendered oil could be scooped off the surface leaving the bottom of the bowl clean but the rims stained. It seems probable that the oil came from marine mammals because they yield much more fat than fish do and were readily available as is evident from archaeological faunal material.

Explaining the peak in stone bowl frequencies

The high stone bowl occurrence during the Margaret Bay phase at the Margaret Bay, and Amaknak Bridge sites is remarkable. What were the driving forces behind the sudden spike in the occurrence of such a specialist artefact type? It is possible that this change in stone bowl frequency is the product of a sampling error. After all, the Margaret Bay (13,500 artefacts, 434 stone bowls) and Amaknak Bridge (3000 artefacts, 71 stone bowls) sites were extensively excavated in comparison to sites of earlier phases (e.g. Oiled Blade: 800 artefacts, 1 stone bowl). However, other sites have also seen extensive investigation and yielded no evidence of stone bowls. For example the Summer Bay site where 564 m² was excavated, yielding 3300 artefacts but no stone bowls, the same goes for the upper levels of Tanaxtaxak, 3500 artefacts total but no stone bowls after the Margaret Bay phase (Knecht and Davis, 2001: p.270, Table 1). The only exception of stone bowl occurrence after the Margaret Bay phase is a surface find that may or may not belong to the Late Aleutian Eider Point site (Fig. 3).

454 Here we explore the notion that stone bowl frequencies spike during the Margaret Bay phase and go out
455 of use after this period ends. Why did frequencies peak at this specific time? What changed?
456 Furthermore, if aquatic oil was rendered using stone bowls then why does this artefact only occur at this
457 frequency during the Margaret Bay phase (4000-3000 cal yr BP)? Assuming oil rendering was important
458 throughout the entire prehistoric and historic sequence of the Aleutian Islands, one would expect to see
459 high frequencies of stone bowls throughout the whole sequence. Were they replaced by another tool
460 type, was the method for rendering oil changed?

461
462 At around 3500 cal yr BP a cold spell ascribed to the Neoglacial brought colder temperatures to the
463 Aleutian Islands (Fig. 10). Marine mammals became more abundant as lower sea surface temperatures
464 increased marine productivity. This induced cultural expansion and increased marine mammal hunting
465 practices. Archaeological bone material is preserved for the first time in the Aleutian sequence and it
466 shows the presence of sea-ice dependent species such as polar bear, ringed seal, and walrus that are
467 not present in later phases when there is no sea-ice (Crockford and Frederick, 2007; Davis, 2001; Knecht
468 and Davis, 2008).

469
470 The increased presence of marine mammals rich in fats during this period of high marine productivity may
471 have increased the rendering of marine oil. On the other hand the unpredictability of climate change could
472 have posed problems for the rendering of oil using a cold method where pieces of fat were stored in a
473 cleaned seal skin, referred to as a *seal poke*, and left to slowly self-render into oil (Frink and Giordano,
474 2015). Temperature was of the utmost importance to this process. Under no circumstances was the
475 substance allowed to freeze, nor should it become too warm. Therefore it was stored in a cool and dark
476 place, often a submerged pit, to prevent the oil from becoming rancid (Frink and Giordano, 2015). Semi-
477 subterranean houses in the Aleutians dating to before 3000 cal yr BP often had subfloor storage pits lined
478 with stone slabs (Knecht and Davis, 2008). It is possible that these pits were used for this purpose.

479
480 We contend that decreasing temperatures in the Aleutians as demonstrated by faunal remains of the
481 Margaret Bay and Amaknak Bridge sites (Davis, 2001; Crockford and Frederick, 2007), may have

induced a change in the method for rendering oil. The hot rendering of oil is not only more controlled but also quicker. Although stone containers could be used for cold rendering as well, for hot rendering the use of a durable container such as a stone bowl was a necessity. Fat was cut up and boiled in water using a container, either by means of stone boiling, or by heating the container directly over a fire. The latter seems to have been the case for the examined specimens as evidenced by thick carbonated encrustations stuck to the bases of the bowls.

The adoption of durable cooking technologies in the Circumpolar North

Aleutian stone bowls and griddle stones are among the earliest durable, non-portable cooking technologies in the circumpolar North. We argue here that stone bowls were used for the rendering of marine fats while griddle stones were probably used for cooking food with high contributions of marine oils. But why were they adopted in the first place? Were stone bowls unique in light of their function? Were they replaceable? How does the adoption of stone bowls relate to the wider debate of early durable container technologies in the circumpolar North?

Stone bowls were a means to an end, to render marine oil. Alternative methods could be used to reach that same end. The seal poke system allowed for the rendering of fat into oil using a cold method that could have been employed before and after the period when bowls became abundant. Climate change may have made this system to render oil more prone to failure and induced the introduction of the stone bowl. However, sedentism also plays an important role here. It was sedentism that allowed for the manufacture and maintenance of this expensive container technology. The fact that a lot of early pottery is associated with the processing of aquatic resources (Jordan and Zvelebil, 2009; Craig et al., 2013) may be closely linked to the notion that a stable abundance of aquatic resources induced sedentism, instead of the idea that aquatic resource processing demanded the use of durable containers such as stone bowls and pottery.

That said, climatic circumstances could very well have encouraged the local invention of the stone bowl in the Aleutian Islands. The hot rendering of aquatic oil would have been difficult, or even impossible with

different, less durable technologies such as basketry or seal pokes. Therefore we ascribe the sudden peak in stone bowl frequencies to a change in oil rendering methods brought on by the Neoglacial cold spell. The introduction and use of griddle stones as a cooking technology is more gradual and consistent. The demand for a container or grill plate to cook on would have easily led to the utilization of these flat stones that are abundant in the Aleutian Islands.

CONCLUSIONS

In this paper our aim was to identify the function of stone bowls and griddle stones. We argue that stone bowls were used for the rendering of marine oil using direct heating. We ascribe the sudden peak in stone bowl occurrence to a shift in temperature caused by the Neoglacial cold spell that subsequently induced a change in oil rendering methods and brought an abundance of sea-ice dependent species, rich in fats, to the area. Our results also suggest that griddle stones were used for the processing of a slightly more diverse set of resources although still predominantly marine. We ascribe this diversity to cooking practices as opposed to oil rendering in stone bowls.

This is the first systematic research into the function of Aleutian cooking technologies employing molecular and chemical analysis of carbonised residues. Future work could test the hypotheses raised in this paper by analysing a larger set of samples, especially regarding griddle stones to more firmly establish trends. The differences between the two technological groups are minor but apparent. Differentiation within the aquatic spectrum is still poorly understood and advances in experimental techniques involving APAA isomer ratios and phytanic acid ratios (Fig. 6), but also an expansion of archaeological reference data from, for example, bone lipids, are necessary to further understand the origin of varying signals.

ACKNOWLEDGEMENTS

This research is a part of MA's PhD research project, hosted by the University of Groningen Arctic Centre as well as the BioArCh laboratory of the University of York and co-supervised by co-authors PJ and OC.

538 The PhD is funded by the University of Groningen, Faculty of Arts. Additional support for residue analysis
539 came from the Arts and Humanities Research Council (AH/L0069X/1). We would like to thank the
540 Museum of the Aleutians and the Ounalashka Corporation for supporting this research and granting
541 access to the artefacts. We also thank the Oregon Museum of Natural and Cultural History and the
542 Katmai National Park Service for allowing the sampling of archaeological bone material. Thank you to
543 Virginia Hatfield for providing the AMK3 griddle stone samples, to Frits Steenhuisen for his help with
544 Figure 1, and to Krista McGrath and Dr. Camilla Speller for the identification of archaeological bone using
545 ZooMS. MA would like to thank Dr. Shinya Shoda and Dr Laura Llorente-Rodriguez for their valuable
546 contribution to her lab training. Special thanks go out to the guest editors Bre MacInnes, Ben Fitzhugh,
547 Kirsten Nicolaysen and Virginia Hatfield for their efforts in bringing this special issue of Quaternary
548 Research together.

549 **REFERENCES**

- 550 Bell, J.G., Preston, T., Henderson, R.J., Strachan, F., Bron, J.E., Cooper, K., Morrison, D.J., 2007.
551 Discrimination of wild and cultured European sea bass (*Dicentrarchus labrax*) using chemical and isotopic
552 analyses. *Journal of Agricultural and Food Chemistry* 55, 5934–5941.
- 553 Brown, T.A., Nelson, D.E., Vogel, J.S., Southon, J.R., 1988. Improved collagen extraction by modified
554 Longin method. *Radiocarbon* 30, 171–177.
- 555 Choy, K., Potter, B.A., McKinney, H.J., Reuther, J.D., Wang, S.W., Wooller, M.J., 2016. Chemical profiling
556 of ancient hearths reveals recurrent salmon use in Ice Age Beringia. *Proceedings of the National*
557 *Academy of Sciences* 113, 9757-9762.
- 558 Colonese, A.C., Farrell, T., Lucquin, A., Firth, D., Charlton, S., Robson, H.K., Alexander, M., Craig, O.E.,
559 2015. Archaeological bone lipids as palaeodietary markers. *Rapid Communications in Mass Spectrometry*
560 29, 611–618.
- 561 Colonese, A.C., Lucquin, A., Guedes, E.P., Thomas, R., Best, J., Fothergill, B.T., Sykes, N., Foster, A.,
562 Miller, H., Poole, K., Maltby, M., Von Tersch, M., Craig, O.E., 2017. The identification of poultry
563 processing in archaeological ceramic vessels using in-situ isotope references for organic residue
564 analysis. *Journal of Archaeological Science* 78, 179–192.
- 565 Corbett, D., Yarborough, M., 2016. The Aleutian tradition: the last 4,000 years. In: Friesen, M. and
566 Mason, O. (Eds.), *The Oxford Handbook of the Prehistoric Arctic*. Oxford University Press, New York,
567 pp.607-630.
- 568 Craig, O.E., Allen, R.B., Thompson, A., Stevens, R.E., Steele, V.J., Heron, C., 2012. Distinguishing wild
569 ruminant lipids by gas chromatography/combustion/isotope ratio mass spectrometry. *Rapid*
570 *Communications in Mass Spectrometry* 26, 2359–2364.
- 571 Craig, O.E., Forster, M., Andersen, S.H., Koch, E., Crombé, P., Milner, N.J., Stern, B., Bailey, G.N.,
572 Heron, C.P., 2007. Molecular and isotopic demonstration of the processing of aquatic products in
573 northern European prehistoric pottery. *Archaeometry* 49, 135–152.

574 Craig, O.E., Saul, H., Lucquin, A., Nishida, Y., Tache, K., Clarke, L., Thompson, A., Altoft, D.T.,
575 Uchiyama, J., Ajimoto, M., Gibbs, K., Isaksson, S., Heron, C.P., Jordan, P., 2013. Earliest evidence for
576 the use of pottery. *Nature* 496, 351–354.

577 Craig, O.E., Steele, V.J., Fischer, A., Hartz, S., Andersen, S.H., Donohoe, P., Glykou, A., Saul, H., Jones,
578 D.M., Koch, E., Heron, C.P., 2011. Ancient lipids reveal continuity in culinary practices across the
579 transition to agriculture in Northern Europe. *Proceedings of the National Academy of Sciences* 108,
580 17910–17915.

581 Cramp, L., Evershed, R.P., 2014. Reconstructing aquatic resource exploitation in human prehistory using
582 lipid biomarkers and stable isotopes. In: Holland, H.D. and Turekian, K.K. (Eds.) *Treatise on*
583 *geochemistry: archaeology and anthropology*. Elsevier, Oxford, 14, pp. 319–339.

584 Cramp, L.J.E., Jones, J., Sheridan, A., Smyth, J., Whelton, H., Mulville, J., Sharples, N., Evershed, R.P.,
585 2014. Immediate replacement of fishing with dairying by the earliest farmers of the Northeast Atlantic
586 archipelagos. *Proceedings of the Royal Society B – Biological Sciences* 281, 20132372.

587 Crockford, S.J., Frederick, S.G., 2007. Sea ice expansion in the Bering Sea during the Neoglacial:
588 evidence from archaeozoology. *Holocene* SAGE Publications 17, 699–706.

589 Davis, R., 2001. Sea mammal hunting and the Neoglacial: environmental change and subsistence
590 technology at Margaret Bay. In: Dumond, D.E. (Ed.), *Archaeology in the Aleut Zone of Alaska*. University
591 of Oregon, Eugene, pp. 71-86.

592 Davis, R., Knecht, R., Rogers, J., 2016. First maritime cultures of the Aleutians. In: Friesen, M. and
593 Mason, O. (Eds.), *The Oxford Handbook of the Prehistoric Arctic*. Oxford University Press, New York, pp.
594 279-302.

595 Debono Spiteri, C., 2012. Pottery use at the transition to agriculture in the western Mediterranean:
596 evidence from biomolecular and isotopic characterisation of organic residues in Impressed/Cardial Ware
597 vessels. University of York, <http://etheses.whiterose.ac.uk/3391/>.

598 Denniston, G.B., 1966. Cultural change at Chaluka, Umnak Island: stone artifacts and features. *Arctic*

599 Anthropology 3, 84-124.

600 Dumond, D.E., 1977. The Eskimos and Aleuts. Thames and Hudson, London.

601 Dumond, D.E., Bland, R.L., 1995. Holocene prehistory of the northernmost North Pacific. *Journal of World*
602 *Prehistory* 9, 401–451.

603 Evershed, R.P., Copley, M.S., Dickson, L., Hansel, F.A., 2008. Experimental evidence for the processing
604 of marine animal products and other commodities containing polyunsaturated fatty acids in pottery
605 vessels. *Archaeometry* 50, 101–113.

606 Farrell, T.F.G., Jordan, P., Taché, K., Lucquin, A., Gibbs, K., Jorge, A., Britton, K., Craig, O.E., Knecht,
607 R., 2014. Specialized processing of aquatic resources in prehistoric Alaskan pottery? A lipid-residue
608 analysis of ceramic sherds from the Thule-period site of Nunalleq, Alaska. *Arctic Anthropology* 51, 86–
609 100.

610 Fernandes, R., Grootes, P., Nadeau, M.-J., Nehlich, O., 2015. Quantitative diet reconstruction of a
611 Neolithic population using a Bayesian mixing model (FRUITS): The case study of Ostorf (Germany).
612 *American Journal of Physical Anthropology* 158, 325-340.

613 Frink, L., Giordano, C., 2015. Women and subsistence food technology: the Arctic seal poke storage
614 system. *Food and Foodways* 23, 251–272.

615 Gibbs, K., Isaksson, S., Craig, O.E., Lucquin, A., Grishchenko, V.A., Farrell, T., Thompson, A., Kato, H.,
616 Vasilevski, A.A., Jordan, P.D., 2017. Exploring the emergence of an 'Aquatic' Neolithic in the Russian Far
617 East: organic residue analysis of early hunter-gatherer pottery from Sakhalin Island. *Antiquity* 91 (360),
618 1484-1500.

619 Hansel, F.A., Copley, M.S., Madureira, L.A.S., Evershed, R.P., 2004. Thermally produced ω -(o-
620 alkylphenyl)alkanoic acids provide evidence for the processing of marine products in archaeological
621 pottery vessels. *Tetrahedron Letters* 45, 2999–3002.

622 Hansel, F.A., Evershed, R.P., 2009. Formation of dihydroxy acids from Z-monounsaturated alkenoic acids

623 and their use as biomarkers for the processing of marine commodities in archaeological pottery vessels.
 624 Tetrahedron Letters 50, 5562–5564.

625 Heron, C., Andersen, S., Fischer, A., Glykou, A., Hartz, S., Saul, H., Steele, V., Craig, O., 2013.
 626 Illuminating the Late Mesolithic: residue analysis of “blubber” lamps from Northern Europe. *Antiquity* 87,
 627 178–188.

628 Heron, C., Craig, O.E., 2015. Aquatic resources in foodcrusts: identification and implication. *Radiocarbon*
 629 57, 707-719.

630 Horiuchi, A., Miyata, Y., Kamijo, N., Cramp, L., Evershed, R.P., 2015. A dietary study of the Kamegaoka
 631 culture population during the final Jomon Period, Japan, using stable isotope and lipid analyses of
 632 ceramic residues. *Radiocarbon* 57, 721–736.

633 Jeanotte, R., Nicolaysen, K., Dowell, F.E., Johnson, T., West, D., 2012. Griddle stones from Adak Island,
 634 Alaska: their provenance and the biological origins of organic residues from cooking. In: West, D.,
 635 Hatfield, V.L., Wilmerding, E.G., Lefèvre, C., Gualtieri, L. (Eds.), *The People Before. The geology,*
 636 *paleoecology and archaeology of Adak Island, Alaska.* Archaeopress, Oxford, pp. 269–287.

637 Jochelson, W., 1925. *Archaeological investigations in the Aleutian Islands.* Carnegie Institution,
 638 Washington D.C.

639 Johnson, L.L., 2004. Prehistoric fishing technologies and species targeted in the Aleutian Islands:
 640 archaeological and ethnohistoric evidence. *New Zealand Journal of Archaeology* 24, 45–59.

641 Jordan, P., Zvelebil, M., 2009. *Ceramics before farming. The dispersal of pottery among prehistoric*
 642 *Eurasian hunter-gatherers.* Left Coast Press Inc., Walnut Creek.

643 Knecht, R., Davis, R.S., 2001. A prehistoric sequence for the Eastern Aleutians. In: Dumond, D.E. (Ed.),
 644 *Archaeology in the Aleut Zone of Alaska.* University of Oregon, Eugene.

645 Knecht, R., Davis, R.S., 2003. Archaeological evaluation of Tanaxtaxak, the Amaknak Spit site (UNL-
 646 00055) final report. The Museum of the Aleutians, Unalaska.

647 Knecht, R., Davis, R.S., 2008. The Amaknak Bridge site: cultural change and the Neoglacial in the
648 Eastern Aleutians. *Arctic Anthropology* 45, 61–78.

649 Knecht, R., Davis, R.S., Carver, G.A., 2001. The Margaret Bay site and eastern Aleutian prehistory. In:
650 Dumond, D.E. (Ed.), *Archaeology in the Aleut Zone of Alaska*. University of Oregon, Eugene.

651 Lucquin, A., Colonese, A.C., Farrell, T.F.G., Craig, O.E., 2016a. Utilising phytanic acid diastereomers for
652 the characterisation of archaeological lipid residues in pottery samples. *Tetrahedron Letters* 57, 703–707.

653 Lucquin, A., Gibbs, K., Uchiyama, J., Saul, H., Ajimoto, M., Eley, Y., Radini, A., Heron, C.P., Shoda, S.,
654 Nishida, Y., Lundy, J., Jordan, P., Isaksson, S., Craig, O.E., 2016b. Ancient lipids document continuity in
655 the use of early hunter–gatherer pottery through 9,000 years of Japanese prehistory. *Proceedings of the*
656 *National Academy of Sciences* 113(15), 3991–3996.

657 Maschner, H., 2016. Archaeology of the Eastern Aleut Region. In: Friesen, M. and Mason, O. (Eds.), *The*
658 *Oxford Handbook of the Prehistoric Arctic*. Oxford University Press, New York, pp. 323–348.

659 McCartney, A.P., Veltre, D.W., 1996. Anangula Core and Blade site. In: West, F.H. (Ed.), *American*
660 *beginnings: the prehistory and palaeoecology of Beringia*. The University of Chicago Press, Ltd., London,
661 pp. 443–450.

662 Oras, E., Higham, T.F.G., Cramp, L.J.E., Bull, I.D., 2017. Archaeological science and object biography: a
663 Roman bronze lamp from Kavastu bog (Estonia). *Antiquity* 91, 124–138.

664 Outram, A.K., Stear, N.A., Bendrey, R., Olsen, S., Kasparov, A., Zaibert, V., Thorpe, N., Evershed, R.P.,
665 2009. The earliest horse harnessing and milking. *Science* 323, 1332–1335.

666 Papakosta, V., Smittenberg, R.H., Gibbs, K., Jordan, P., Isaksson, S., 2015. Extraction and derivatization
667 of absorbed lipid residues from very small and very old samples of ceramic potsherds for molecular
668 analysis by gas chromatography–mass spectrometry (GC-MS) and single compound stable carbon
669 isotope analysis by gas chromatography–combustion–isotope ratio mass spectrometry (GC-C-IRMS).
670 *Microchemical Journal* 123, 196–200.

671 Piezonka, H., Meadows, J., Hartz, S., Kostyleva, E., Nedomolkina, N., Ivanishcheva, M., Kosorukova, N.,
672 Terberger, T., 2016. Stone Age pottery chronology in the Northeast European forest zone: new AMS and
673 EA-IRMS results on foodcrusts. *Radiocarbon* 58, 267-289.

674 Quimby, G.I., 1945. Pottery from the Aleutian Islands. *Fieldiana Anthropology* 36, 1–13.

675 Sarychev, G., 1806. Account of a voyage of discovery to the 1807 north-east of Siberia, the frozen ocean
676 and the north-east sea. Translated from the Russian and embellished with engravings. Vol.1, 1806; Vol.2,
677 1807, J.G. Bernard, London.

678 Shoda, S., Lucquin, A., Ahn, J.-H., Hwang, C.-J., Craig, O.E., 2017. Pottery use by early Holocene
679 hunter-gatherers of the Korean peninsula closely linked with the exploitation of marine resources.
680 *Quaternary Science Reviews* 170, 164–173.

681 Taché, K., Craig, O.E., 2015. Cooperative harvesting of aquatic resources and the beginning of pottery
682 production in north-eastern North America. *Antiquity* 89, 177–190.

683 Unger, S., 2014. Qaqamiigux: traditional foods and recipes from the Aleutian and Pribilof Islands:
684 nourishing our mind, body and spirit for generations. Aleutian Pribilof Islands Association, Incorporated,
685 Anchorage.

686 Van Bergen, P.F., Nott, C.J., Bull, I.D., Poulton, P.R., Evershed, R.P., 1998. Organic geochemical studies
687 of soils from the Rothamsted Classical Experiments-IV. Preliminary results from a study of the effect of
688 soil pH on organic matter decay. *Organic Geochemistry* 29, 1779–1795.

689 **List of Tables + captions:**

690 Table 1: Prehistory of the Eastern Aleutian Islands: dates, site characteristics, subsistence trends
691 and climatic influences. Based on: 1: Davis et al, (2016), 2: Hatfield (2010), 3: Maschner (2016), 4: Mason
692 (2001), 5: Magny and Haas (2004), 6: Knecht and Davis, 2008, 7: Corbett and Yarborough (2016)

693 Supplementary Table 1: Summary of organic residue analysis results on stone bowls and griddle
694 stones from the Aleutian Islands. Blank cells = no analysis, - = compound not present

695 Supplementary Table 2: Bone collagen data from archaeological contexts of the Aleutian Islands
696 and the Alaska Peninsula. Species were identified by ZooMS at the University of York for the following
697 samples: 201, 204-5, 209-10, 310-12, 316, and 319.

698 **List of Figures + captions**

699 Figure 1: map of the Aleutian Islands, Alaska Peninsula and Kodiak Island with emphasis on site
700 locations mentioned in the text at Unalaska Island (Amaknak Bridge, Margaret Bay, Oiled Blade,
701 Tanaxtaxak), Umnak Island (Anangula Blade), and Carlisle Island (Ulyagan).

702 Figure 2: Example of two different types of stone bowls fashioned out of differing textured volcanic
703 tuff and varying in size and shape: a) UNL48-57: red, more crude textured tuff, thick rim, thin base; b)
704 UNL50-51: sand colored fine tuff, finely ground both inside and out c) UNL50-51: base with carbonized
705 encrustations (Photographs by M. Admiraal, courtesy of the Museum of the Aleutians).

706 Figure 3: Stone bowl with encrustation on the interior. Surface find from Eider Point site (UNL-19)
707 probably dating to the Late Aleutian phase (1,000-2,000 cal BP) (Photograph by M. Admiraal, courtesy of
708 the Museum of the Aleutians).

709 Figure 4: A. UNL55-39 griddle stone with clean center, Late Aleutian phase; B. UNL318-47 griddle
710 stone with encrustations in the center, Early Anangula phase (Photographs by M. Admiraal, courtesy of
711 the Museum of the Aleutians).

712 Figure 5: a typical total ion current (TIC) of an acid/methanol extract of a stone bowl from the

713 Margaret Bay site (UNL48-61b) showing saturated fatty acids, diacids (DC), branched (br), isoprenoid
714 acids (4,8,12-TMTD, pristanic, and phytanic acid) and long-chain unsaturated fatty acids.

715 Figure 6: percentage of SSR (SSR%) diastereomer in total phytanic acid in Aleutian artefacts
716 compared to modern ruminant and aquatic resources (Lucquin et al., 2016a, 2016b).

717 Figure 7: Partial summed mass chromatogram (m/z 105) showing APAA distribution in griddle
718 stone sample AMK3-1030 run on DB23 using AQUASIM method. *: C₁₆, +: C₁₈, #: C₂₀, °: C₂₂

719 Figure 8: a. Modern reference samples of anadromous fish (pink triangles), freshwater fish (green
720 diamonds), marine fish (blue circles), marine mammals (blue squares) and aquatic birds (yellow
721 downward triangles) (Bell et al., 2007; Outram et al., 2009; Craig et al., 2011, 2013; Debono Spiteri, 2012;
722 Cramp et al., 2014; Colonese et al., 2015; Horiuchi et al., 2015; Taché and Craig, 2015; Choy et al.,
723 2016). b. GC-c-IRMS results showing isotopic values of C_{16:0} and C_{18:0} fatty acids of stone bowls (green
724 circles), griddle stones (yellow squares) and one lamp (blue triangle).

725 Figure 9: Bulk isotope results of stone bowls (green circles), griddle stones (yellow squares) and
726 lamps (blue triangles) compared to Sakhalin pottery (open diamonds) (Gibbs et al., 2017) and European
727 oil lamps (open triangles) (Heron et al., 2013; Heron and Craig, 2015; Piezonka et al., 2016; Oras et al.,
728 2017) against archaeological bone collagen data from the Aleutian Islands and the Alaska Peninsula. The
729 collagen $\delta^{15}\text{N}$ values were adjusted by +2‰ to correct for the collagen to tissue offset in order to make
730 these values more comparable with the food crusts (Fernandes et al., 2015).

731 Figure 10: Schematic diagram of stone bowl and griddle stone relative abundance against the
732 Margaret Bay phase sea-ice presence in the Aleutian Islands as inferred by (Crockford and Frederick,
733 2007) on the basis of archaeological faunal assemblages.