

This is a repository copy of *Ehresmann monoids: Adequacy and expansions*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/134067/>

Version: Accepted Version

Article:

Branco, Mario J. J., Gomes, Gracinda M. S., Gould, Victoria Ann Rosalind orcid.org/0000-0001-8753-693X et al. (1 more author) (2018) *Ehresmann monoids: Adequacy and expansions*. *Journal of Algebra*. pp. 344-367. ISSN: 1090-266X

<https://doi.org/10.1016/j.jalgebra.2018.06.036>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

EHRESMANN MONOIDS: ADEQUACY AND EXPANSIONS

MÁRIO J.J. BRANCO, GRACINDA M.S. GOMES, VICTORIA GOULD,
AND YANHUI WANG

ABSTRACT. It is known that an Ehresmann monoid $\mathcal{P}(T, Y)$ may be constructed from a monoid T acting via order-preserving maps on both sides of a semilattice Y with identity, such that the actions satisfy an appropriate compatibility criterion. Our main result shows that if T is cancellative and equidivisible (as is the case for the free monoid X^*), the monoid $\mathcal{P}(T, Y)$ not only is Ehresmann but also satisfies the stronger property of being adequate.

Fixing T , Y and the actions, we characterise $\mathcal{P}(T, Y)$ as being unique in the sense that it is the initial object in a suitable category of Ehresmann monoids. We also prove that the operator \mathcal{P} defines an expansion of Ehresmann monoids.

INTRODUCTION

Ehresmann monoids have their roots in the work of Ehresmann on local structures in differential geometry [5], and were formally introduced in the literature by Lawson [17]. They may be defined in various ways but here we take the modern approach and consider them as monoids equipped with two basic unary operations (usually denoted $+$ and $*$), that is, as *bi-unary monoids*. As such, Ehresmann monoids form a variety Ehr and so the free Ehresmann monoid on any set exists. The variety Ehr contains the quasi-variety Adq of adequate monoids. A rich theory has developed surrounding adequate monoids since their introduction by Fountain [6]; they are precisely those monoids with commuting idempotents such that every principal one-sided ideal is projective. From [14], and also [3] and our results here, the free Ehresmann monoid coincides with the free adequate monoid on any set, so that Ehr is exactly the variety generated by the quasi-variety Adq . Any inverse monoid is Ehresmann with $a^+ = aa^{-1}$ and $a^* = a^{-1}a$, but Ehresmann monoids are in general very far from being regular.

The importance of understanding algebras by means of their actions is central in mathematics. Examples include the deep theories of R -modules over a ring R and S -acts over a monoid S , Bass-Serre theory of groups acting on trees, and (in a context relatively close to that of this article), McAlister's characterisation of proper inverse monoids by means of groups acting on partially ordered sets. Any inverse monoid has a proper cover [19, 20], and any proper inverse monoid

1991 *Mathematics Subject Classification.* Primary 20M10; Secondary 20M99.

Key words and phrases. Ehresmann monoid, adequate monoid, expansion.

embeds into a semidirect product of a semilattice by a group [23]. Note that the free inverse monoid on any set is proper [22, 24].

The crucial results of McAlister in the inverse case have been extended to various classes of unary and bi-unary monoids, requiring a sequence of new ideas as one moves further away from regularity. Many difficulties arise, and the theory that emerges splits into one and two-sided cases. We refer the reader to the work of Fountain, Gomes, Gould, Jones, Lawson, Kambites and Kudryavtseva [6, 7, 8, 9, 12, 14, 15, 16, 17] to see the development of the theory.

In [2], Branco, Gomes and Gould initiated a new approach to the study of left Ehresmann and left adequate monoids, where certainly the techniques involving semidirect products fail. In a (left/right) Ehresmann monoid M , the image of the unary operation(s) forms a semilattice, say Y , named the *semilattice of projections*; given a submonoid T , the monoid M is said to be *T -generated* if it is generated as a semigroup¹ by $T \cup Y$. They introduced for a T -generated left Ehresmann monoid the concept of *T -proper* (the analogue of the aforementioned concept of proper is not useful here); proved that any left Ehresmann monoid has an X^* -proper cover for some set X , where X^* is the free monoid on X , and deduced that the free left Ehresmann monoid on any set X is X^* -proper. In a subsequent paper [10], Gomes and Gould took a monoid T acting via order-preserving maps on the left of a semilattice Y with identity, and constructed a T -generated T -proper left Ehresmann monoid with semilattice of projections Y , which we denote here $\mathcal{P}_\ell(T, Y)$. They proved that if T is right cancellative with trivial group of units, then $\mathcal{P}_\ell(T, Y)$ is left adequate; and that the free left Ehresmann monoid on X is of the form $\mathcal{P}_\ell(X^*, Y)$, coinciding therefore with the free left adequate monoid on X . We remark that the monoid $\mathcal{P}_\ell(T, Y)$ may be characterised as being the unique T -proper T -generated left Ehresmann monoid having *uniqueness of T -normal forms*. Here, we do not concern ourselves with uniqueness of T -normal forms [2, 10], but point out that their presence implies the property of being T -proper.

Naturally, one would like to have an analogous theory for the two-sided case. This paper is the second of a pair (the first being [3]) initiating, developing and implementing a theory for *two-sided* Ehresmann and adequate monoids, corresponding to that in the one-sided case. It is very far from true that combining the left and the right cases is sufficient in itself to make progress: new methods are required.

In [3] the first three authors introduced the notions of *T -proper* and *strongly T -proper* for T -generated Ehresmann monoids. The main thrust was to construct a strongly T -proper Ehresmann monoid $\mathcal{P}(T, Y)$ from a semilattice Y with identity acted upon *on both sides* by a monoid T via order-preserving maps satisfying the so called *compatibility conditions* for the actions. Let us encode the monoid T ,

¹The signature can also be that of a monoid or (bi-)unary monoid, without the concept being affected.

the semilattice Y and the actions via a \mathcal{P} -quadruple $\mathcal{T} = (T, Y, \cdot, \circ)$. It is proved in [3] that any Ehresmann monoid admits a strongly X^* -proper Ehresmann cover and that the free Ehresmann monoid is of the form $\mathcal{P}(X^*, Y)$. Observe that the existence of covers for Ehresmann monoids is also discussed in [13].

A matter missing from [3] concerns the claim of adequacy for $\mathcal{P}(T, Y)$ in the case where T is cancellative. This the major question that gave rise to this paper.

For ease of reference we recall in Section 1 some basic facts concerning adequate and Ehresmann monoids, and the construction of $\mathcal{P}(T, Y)$. In Section 2, we tackle the above question and answer it in full when T is equidivisible.

Theorem 2.10 *Let T be an equidivisible cancellative monoid acting on both sides upon a semilattice Y with identity, satisfying the compatibility conditions. Then $\mathcal{P}(T, Y)$ is adequate.*

As a consequence of this theorem, $\mathcal{P}(X^*, Y)$ is adequate, thus confirming, as mentioned before, that the free Ehresmann monoid on X is in fact the free adequate monoid, a result essentially shown in [14].

In Section 3, we return to the consideration of arbitrary monoids T . First we show that fixing T , Y and the actions of T on Y , the operator \mathcal{P} determines an expansion of Ehresmann monoids (Theorem 3.4). Next, we aim to find an abstract characterisation of the Ehresmann monoids of the form $\mathcal{P}(T, Y)$, built from a \mathcal{P} -quadruple \mathcal{T} . Notice that, unlike the one-sided case, $\mathcal{P}(T, Y)$ does not have uniqueness of T -normal forms [3], and so it cannot be distinguished by such a property.

Given a \mathcal{P} -quadruple $\mathcal{T} = (T, Y, \cdot, \circ)$, we define a category $\mathcal{C}(\mathcal{T})$ of Ehresmann monoids, that are T -generated with semilattice of projections Y . Our second main result can be stated as follows.

Theorem 3.6 *Let $\mathcal{T} = (T, Y, \cdot, \circ)$ be a \mathcal{P} -quadruple. Then $\mathcal{P}(T, Y)$ is the initial object in the category $\mathcal{C}(\mathcal{T})$.*

Section 4 ends the article by posing some open questions.

1. PRELIMINARIES

Before we recall some basic definitions and results concerning adequate and Ehresmann monoids we describe the route that has led us to viewing the monoid $\mathcal{P}(T, Y)$ as a generalisation of a semidirect product. For further details, of both background and technicalities, we refer the reader to [3] and [11].

As remarked in the Introduction, the crucial properties and resulting structure of proper inverse semigroups and monoids have inspired the search for analogous results for other classes. We focus here for convenience on the case for monoids, although many of the following statements do not require the presence of an identity. Let S be an inverse monoid, so that S is Ehresmann with semilattice of projections $E = E(S)$. Any product of elements of S of the form $w = a_0 e_1 a_1 e_1 \dots e_n a_n$ where $a_i \in S, e_j \in E$, for $0 \leq i \leq n, 1 \leq j \leq n$, can be manipulated into an expression $w = e a_0 a_1 \dots a_n$ where $e \in E$. This is due to the

fact that if a is an element of an inverse monoid and e is idempotent, then there is an idempotent f with $ea = af$, and vice versa. In fact, inverse monoids satisfy the *ample* identities $xy^+ = (xy)^+x$ and $y^*x = x(yx)^*$. This is a key factor in co-ordinatising elements of a proper inverse monoid S by *two* co-ordinates - an idempotent and an element of the group S/σ - and the embeddability of proper inverse monoids into semidirect products. A number of authors have considered generalisations of inverse monoids satisfying one or both of the ample identities, as appropriate, where again semidirect products feature (see, for example, [4, 8, 9]). Many of these are unary or bi-unary monoids having natural representations as mappings. One such case is that of left restriction monoids, which are unary monoids represented by monoids of partial mappings where the unary operation takes an element to the identity in its domain. Once one moves from monoids of mappings to more general monoids of relations, the ample identity may be lost. Indeed, a motivating example is that of \mathcal{B}_X , the bi-unary monoid of relations on a set X , with unary operations of domain and range. The monoid \mathcal{B}_X is Ehresmann but does not satisfy the ample identities. It follows that in attempting to understand an Ehresmann monoid M in terms of the projections E and the *reduced* Ehresmann monoid M/σ , we must consider sequences of the form $a_0e_1a_1e_1 \dots e_na_n$ without recourse to any quick simplification. This leads us naturally to replacing semidirect products by quotients of free products. The congruences are determined by action(s) that persist from earlier cases without recourse to the ample identities.

Let M be a monoid with set of idempotents $E(M)$. For any $a, b \in M$,

$$a \mathcal{R}^* b \Leftrightarrow \forall x, y \in M (xa = ya \Leftrightarrow xb = yb)$$

and

$$a \mathcal{L}^* b \Leftrightarrow \forall x, y \in M (ax = ay \Leftrightarrow bx = by).$$

Clearly, \mathcal{R}^* is a left congruence and \mathcal{L}^* a right congruence.

Recall that a monoid M is *adequate* if every \mathcal{R}^* -class and every \mathcal{L}^* -class contains an idempotent and $E(M)$ forms a semilattice. From the commutativity of idempotents it follows that each \mathcal{R}^* -class and \mathcal{L}^* -class of an element a contains a unique idempotent, denoted by a^+ and a^* , respectively. Thus we have two unary operations on M given by $a \mapsto a^+$ and $a \mapsto a^*$, whence M becomes an algebra with signature $(2, 1, 1, 0)$; as such we refer to it as a *bi-unary monoid*. The class of adequate monoids forms a quasi-variety of algebras in this signature. The defining quasi-identities are those for monoids together with

$$x^+x = x, \quad (x^+y^+)^+ = x^+y^+ = y^+x^+ \quad \text{and} \quad (xy)^+ = (xy^+)^+,$$

$$x^2 = x \rightarrow x = x^+ \quad \text{and} \quad xy = zy \rightarrow xy^+ = zy^+,$$

and their left-right duals.

An *Ehresmann monoid* is a bi-unary monoid M , in which we again denote the unary operations by $^+$ and * , satisfying the identities for monoids together with

$$x^+x = x, \quad (x^+y^+)^+ = x^+y^+ = y^+x^+ \quad \text{and} \quad (xy)^+ = (xy^+)^+,$$

their left-right duals, and

$$(x^*)^+ = x^* \quad \text{and} \quad (x^+)^* = x^+.$$

The identities $x^+ = x^+x^+$ and $(x^+)^+ = x^+$, and their left-right duals, are a consequence of those above. Putting $E = \{s^+ : s \in M\} = \{s^* : s \in M\}$ we have that E is a semilattice, the *semilattice of projections*. We have already remarked that the variety generated by the quasi-variety of adequate monoids is the variety of Ehresmann monoids. In particular, an adequate monoid M is Ehresmann, with $E(M) = E$.

Another approach to Ehresmann monoids is based on relations $\tilde{\mathcal{R}}_E$ and $\tilde{\mathcal{L}}_E$, which themselves contain \mathcal{R}^* and \mathcal{L}^* , respectively. We do not pursue this route here, the interested reader may consult [3].

Let M be an Ehresmann monoid with semilattice of projections E and submonoid T . Recall from the Introduction that M is *T -generated* if M is generated (as a semigroup) by $T \cup E$: we denote this by $M = \langle T \cup E \rangle_{(2)}$.

Lemma 1.1. [3, Lemmas 2.2 and 2.3] *Let M be a T -generated Ehresmann monoid with semilattice of projections E . Then T acts on the left and on the right of E by order-preserving maps defining, for $t \in T$ and $e \in E$,*

$$t \cdot e = (te)^+ \quad \text{and} \quad e \circ t = (et)^*.$$

On the other hand, for any $a \in M$ and $e, f \in E$,

$$(eaf)^+ = e(a(eaf)^*)^+ \quad \text{and} \quad (eaf)^* = ((eaf)^+a)^*f.$$

On an Ehresmann monoid M with semilattice of projections E , the relation σ (σ_E for emphasis) is the *semigroup congruence on S generated by $E \times E$* . It is clear that σ is also the bi-unary monoid congruence on the same generators.

A T -generated Ehresmann monoid M with semilattice of projections E is said to be *T -proper* if for any $s, t \in T$ and $e \in E$,

$$[(se)^+ = (te)^+ \text{ and } se \sigma te] \Rightarrow se = te$$

and, dually,

$$[(es)^* = (et)^* \text{ and } es \sigma et] \Rightarrow es = et.$$

Further, M is said to be *strongly T -proper* if, for any $s, t \in T$

$$s \sigma t \Rightarrow s = t.$$

Note that strongly T -proper implies T -proper; the exact relationship between the two conditions is still under investigation.

We recall from [3] the recipe of the first three authors for constructing T -proper Ehresmann monoids from monoids acting on semilattices.

Let T be a monoid with identity 1_T and let Y be a semilattice with identity 1_Y . To avoid any ambiguity we assume that $T \cap Y = \emptyset$. Let $T * Y$ be the free semigroup product of T and Y . We say that $x \in T * Y$ has a T -beginning if x begins with a $t \in T$, that is, $x = tz$ for some $z \in T * Y$. Dually, x has a T -end if $x = zt$ for some $t \in T$ and $z \in T * Y$. Correspondingly, we say that x has a Y -beginning (Y -end) if $x = ez$ ($x = ze$) for some $e \in Y$ and $z \in T * Y$.

If, for example, x has a T -beginning and Y -end, we write x as

$$x = t_0 e_1 t_1 e_2 \dots t_{n-1} e_n$$

where $t_i \in T$ and $e_j \in Y$, $0 \leq i \leq n-1$ and $1 \leq j \leq n$.

Suppose that T acts on the left of Y via order-preserving maps. We denote the action of $t \in T$ on $y \in Y$ by $t \cdot y$. It follows that there exists a monoid morphism

$$\phi_\ell : T \rightarrow \mathcal{O}_Y^*, (t\phi_\ell)(y) = t \cdot y,$$

where \mathcal{O}_Y^* is the monoid of order-preserving maps of Y with maps composed from right to left. Now, Y acts on the left of itself by order-preserving maps via multiplication, hence there is a monoid morphism, also denoted ϕ_ℓ , given by

$$\phi_\ell : Y \rightarrow \mathcal{O}_Y^*, (z\phi_\ell)(y) = zy.$$

By the universal property of free products, we obtain a semigroup morphism

$$\phi_\ell : T * Y \rightarrow \mathcal{O}_Y^*$$

defined by

$$(s_1 \dots s_n)\phi_\ell = (s_1\phi_\ell) \dots (s_n\phi_\ell),$$

where each $s_i \in T \cup Y$. We thus have a semigroup left action of $T * Y$ on Y , which we may without ambiguity denote by \cdot , given by

$$s_1 \dots s_n \cdot y = s_1 \cdot (s_2 \cdot (\dots (s_n \cdot y) \dots)).$$

We now define u^+ , for $u \in T * Y$, to be

$$u^+ = u \cdot 1_Y.$$

Therefore $e^+ = e$ for all $e \in Y$. We remark that for any $u \in T * Y$, if v is obtained from u via insertion or deletion of elements 1_Y or 1_T , then $u^+ = v^+$. Notice also that $1_T^+ = 1_Y$. The free product $T * Y$ is now a unary semigroup.

Lemma 1.2. [3, Lemma 4.1] *If $u, v \in T * Y$ and $e \in Y$, then $(uv)^+ = u \cdot v^+ = (uv^+)^+$, $(eu)^+ = eu^+$, $(uv)^+ \leq u^+$ and $(uev)^+ \leq (uv)^+$.*

We also suppose we have a right action of T on Y via order-preserving maps. Denoting the right action of $t \in T$ on $y \in Y$ by $y \circ t$, there exists a monoid morphism

$$\phi_r : T \rightarrow \mathcal{O}_Y, (y)(t\phi_r) = y \circ t,$$

where \mathcal{O}_Y denotes the dual monoid of \mathcal{O}_Y^* (where maps are composed from left to right). Again Y acts on the right on itself by order-preserving maps via

multiplication, and we may consider the monoid morphism, also denoted ϕ_r , given by

$$\phi_r : Y \rightarrow \mathcal{O}_Y, (y)(z\phi_r) = yz.$$

As before, by the universal property of free products, we have a semigroup morphism

$$\phi_r : T * Y \rightarrow \mathcal{O}_Y$$

defined by

$$(s_1 \dots s_n)\phi_r = (s_1\phi_r) \dots (s_n\phi_r),$$

where each $s_i \in T \cup Y$. We thus have a semigroup action of $T * Y$ on Y , which we may without ambiguity denote by \circ , where

$$y \circ s_1 \dots s_n = ((\dots (y \circ s_1) \dots) \circ s_{n-1}) \circ s_n.$$

We now define u^* (for $u \in T * Y$) to be

$$u^* = 1_Y \circ u,$$

so that $e^* = e$ for all $e \in Y$. As before, we remark that for any $u \in T * Y$, if v is obtained from u via insertion or deletion of elements 1_Y or 1_T , then $u^* = v^*$. Notice also that $1_T^* = 1_Y$ and we have that the free product $T * Y$ has become a bi-ary semigroup.

Lemma 1.3. [3, Lemma 4.2] *If $u, v \in T * Y$ and $e \in Y$, then $(uv)^* = u^* \circ v = (u^*v)^*$, $(ue)^* = u^*e$, $(uv)^* \leq v^*$ and $(uev)^* \leq (uv)^*$.*

From Lemmas 1.2 and 1.3, we observe that for any $u \in T * Y$ and $e \in Y$, we get

$$u \cdot e = (ue)^+ \text{ and } e \circ u = (eu)^*.$$

However, $T * Y$ is not Ehresmann, since for example it does not satisfy the identity $x^+x = x$.

To proceed, we require the actions to satisfy compatibility conditions that we now define.

Definition 1.4. Let T be a monoid acting on both sides upon a semilattice Y . We say that the *compatibility conditions* are satisfied if, for any $t \in T$ and $e, f \in Y$:

$$(CC1) \quad e(t \cdot f) = e(t \cdot ((e \circ t)f))$$

and

$$(CC2) \quad (e \circ t)f = ((e(t \cdot f)) \circ t)f.$$

Definition 1.5. A \mathcal{P} -quadruple is a quadruple $\mathcal{T} = (T, Y, \cdot, \circ)$ where T is a monoid acting by \cdot on the left and \circ on the right of a semilattice Y with identity via order preserving maps satisfying the compatibility conditions.

Observe that by Lemma 1.1, given a T -generated Ehresmann monoid M , the monoid T acts on both sides upon E satisfying the compatibility conditions and with these actions $\mathcal{T} = (T, E, \cdot, \circ)$ is a \mathcal{P} -quadruple.

Aiming at constructing the Ehresmann monoid $\mathcal{P}(T, Y)$, now let

$$H_\ell = \{(u^+u, u) : u \in T * Y\} \cup \{(1_T, 1_Y)\}$$

and

$$H_r = \{(uu^*, u) : u \in T * Y\} \cup \{(1_T, 1_Y)\}.$$

We use \sim to denote the semigroup congruence on $T * Y$ generated by $H_\ell \cup H_r$. Thus for any $u, v \in T * Y$, we have that $u \sim v$ if and only if $u = v$ or there is a sequence

$$u = z_0, z_1, \dots, z_n = v$$

where $n \in \mathbb{N}$ and for $0 \leq i \leq n-1$ we have

$$z_i = c_i \alpha_i d_i, z_{i+1} = c_i \beta_i d_i$$

for some $c_i, d_i \in (T * Y)^1$ and $(\alpha_i, \beta_i) \in (H_\ell \cup H_r) \cup (H_\ell \cup H_r)^{-1}$.

If $n = 1$ and $c_1, d_1 \in T * Y$, we say that $u \sim v$ *via a basic step*. The relation \sim is not just a congruence on $T * Y$ but it is also a bi-unary congruence [3, Lemma 4.9].

An element $z \in T * Y$ can take one of four forms, depending on whether z has T - or Y -beginning and T - or Y -end. For convenience, we introduce a new symbol \square which we regard as an adjoined identity to the monoid T . By writing an element $z \in T * Y$ as $\square e_1 z_1 \dots e_n z_n$, where $e_1, \dots, e_n \in Y$ and $z_1, \dots, z_n \in T$, we are indicating that $z = e_1 z_1 \dots e_n z_n$ has Y -beginning, with similar conventions for Y -ends. The \square symbol serves as a marker to help us control places in products of elements in $T * Y$.

Lemma 1.6. [3, Lemma 4.12] *The map $\tau : T * Y \rightarrow T$ given by: $\tau(t) = t$ if $t \in T$, $\tau(y) = 1_T$ if $y \in Y$, and*

$$\tau(u) = t_0 t_1 \dots t_n,$$

for $u = t_0 e_1 t_1 \dots e_n t_n$ with $t_0, t_n \in T \cup \{\square\}$, $t_1, \dots, t_{n-1} \in T$ and $e_1, e_2, \dots, e_n \in Y$, is a well-defined monoid morphism with $\sim \subseteq \ker \tau$.

Theorem 1.7. [3, Theorem 4.18] *Let $\mathcal{T} = (T, Y, \cdot, \circ)$ be a \mathcal{P} -quadruple. The quotient $\mathcal{P}(T, Y) := (T * Y) / \sim$ is an Ehresmann monoid with semilattice of projections*

$$Y' = \{[e] : e \in Y\}$$

where $[u]^+ = [u^+]$ and $[u]^ = [u^*]$ for any $u \in T * Y$, and $1_{\mathcal{P}(T, Y)} = [1_T] = [1_Y]$.*

*Further, Y' is isomorphic to Y and the submonoid $T' = \{[t] : t \in T\}$ of $\mathcal{P}(T, Y)$ is isomorphic to T under the natural morphism $\nu_{\mathcal{T}} : T * Y \rightarrow \mathcal{P}(T, Y)$.*

The monoid $\mathcal{P}(T, Y)$ is T' -generated, with $\mathcal{P}(T, Y)/\sigma_{Y'} \simeq T'$ and so strongly T' -proper, hence T' -proper.

We end this section by remarking that \sim is generated by H_ℓ together with H_r , and it is precisely the interaction between the two kinds of generators that causes us difficulties at many stages in proof when comparing to left Ehresmann and right Ehresmann monoids. This is highlighted in, for example, Lemma 2.8 and Theorem 2.10.

2. SUFFICIENT CONDITIONS FOR $\mathcal{P}(T, Y)$ TO BE ADEQUATE

The aim of this section is to show that $\mathcal{P}(T, Y)$ is an adequate monoid when T is an equidivisible cancellative monoid.

Recall that a monoid T is *equidivisible* if for any $a, b, c, d \in T$, if $ab = cd$ then for some $u \in T$, $a = cu$ and $ub = d$, or $au = c$ and $b = ud$. Groups and free monoids are clearly the first examples. The direct product of a free monoid and a group is equidivisible and cancellative, but excluding degenerate cases in general is neither a group, nor a free monoid, nor has trivial group of units. The non-negative real numbers under addition provide another example of an equidivisible cancellative monoid. A classical example of an equidivisible cancellative monoid with trivial group of units that is not free can be found in [21, Example 6.24]. In fact as proved in [18], a monoid is free if and only if it is graded and equidivisible.

Throughout this section, we assume that T is a cancellative monoid acting on both sides upon a semilattice Y with identity by order-preserving maps, satisfying the compatibility conditions. We will denote the group of units of T by $U(T)$ and the (group) inverse of an element $t \in U(T)$ by t^{-1} .

Lemma 2.1. *Let $u = t_0 e_1 t_1 \dots e_n t_n \in T * Y$ be such that $\tau(u) = 1_T$, where $t_i \in T$ ($0 \leq i \leq n$) and $e_j \in Y$ ($1 \leq j \leq n$). Then*

$$(t_i e_{i+1} t_{i+1} \dots e_n t_n u^+)^+ = (t_i e_{i+1} t_{i+1} \dots e_n t_n u)^+ \leq (t_i t_{i+1} \dots t_n u)^+ \leq e_i$$

for any i with $1 \leq i \leq n$. In particular, $(t_n u^+) = (t_n u)^+ \leq e_n$.

Proof. By Lemma 1.2, we get the equalities as well as

$$(t_i e_{i+1} t_{i+1} \dots e_n t_n u)^+ \leq (t_i t_{i+1} \dots t_n u)^+,$$

for any i with $1 \leq i \leq n$. Since $\tau(u) = 1_T$, we have that $t_0 \dots t_n = 1_T$. Then, as T is cancellative, $t_0, \dots, t_n \in U(T)$ and for any $i \in \{1, \dots, n\}$, we obtain $(t_0 t_1 \dots t_{i-1})^{-1} = t_i t_{i+1} \dots t_n = t_{i-1}^{-1} \dots t_1^{-1} t_0^{-1}$. Thus

$$\begin{aligned}
(t_i t_{i+1} \dots t_n u)^+ &= (t_{i-1}^{-1} \dots t_1^{-1} e_1 t_1 \dots e_n t_n)^+ \\
&\leq (t_{i-1}^{-1} \dots t_1^{-1} t_1 e_2 t_2 \dots e_n t_n)^+ \quad (\text{by Lemma 1.2}) \\
&= (t_{i-1}^{-1} \dots t_2^{-1} e_2 t_2 \dots e_n t_n)^+ \\
&\vdots \\
&= (e_i t_i \dots e_n t_n)^+ \\
&\leq e_i \quad (\text{by Lemma 1.2}),
\end{aligned}$$

as required. \square

Lemma 2.2. *If $u \in T * Y$ is such that $\tau(u) = 1_T$, then $u \sim u^+ \sim u^*$.*

Proof. We begin by supposing that u has a T -beginning and a T -end. Then

$$u = t_0 e_1 t_1 e_1 \dots t_{n-1} e_n t_n$$

where $t_i \in T$ and $e_j \in Y$, $0 \leq i \leq n$ and $1 \leq j \leq n$. Since $\tau(u) = 1_T$, we have that $u^+ = 1_Y u^+ \sim 1_T u^+ = t_0 t_1 \dots t_n u^+$, and so

$$\begin{aligned}
u^+ &\sim t_0 t_1 \dots t_n u^+ \\
&\sim t_0 t_1 \dots t_{n-1} (t_n u^+)^+ t_n u^+ \\
&= t_0 t_1 \dots t_{n-1} e_n (t_n u^+)^+ t_n u^+ \quad (\text{by Lemma 2.1}) \\
&\sim t_0 t_1 \dots t_{n-1} e_n t_n u^+ \\
&\vdots \\
&\sim t_0 t_1 \dots t_i e_{i+1} t_{i+1} \dots e_n t_n u^+ \\
&\sim (t_0 t_1 \dots t_{i-1}) (t_i e_{i+1} t_{i+1} \dots e_n t_n u^+)^+ t_i e_{i+1} t_{i+1} \dots e_n t_n u^+ \\
&= (t_0 t_1 \dots t_{i-1}) e_i (t_i e_{i+1} t_{i+1} \dots e_n t_n u^+)^+ t_i e_{i+1} t_{i+1} \dots e_n t_n u^+ \quad (\text{by Lemma 2.1}) \\
&\sim (t_0 t_1 \dots t_{i-1}) e_i t_i e_{i+1} t_{i+1} \dots e_n t_n u^+ \\
&\vdots \\
&\sim t_0 e_1 t_1 \dots e_n t_n u^+ \\
&= u u^+.
\end{aligned}$$

Dually, we can show that $u^* \sim u^* u$. As Y is a semilattice, it follows that

$$u^+ u^* \sim u u^+ u^* = u u^* u^+ \sim u u^+ \sim u^+,$$

and similarly, $u^+ u^* \sim u^*$. We now deduce that $u^* \sim u^+$ and finally

$$u^+ \sim u u^+ \sim u u^* \sim u.$$

Now, suppose that u has a Y -beginning. Then $u = 1_Y u$, so that $u \sim 1_T u$ and $1_T u$ has a T -beginning. Similarly, if u has a Y -end, we get $u \sim u 1_T$ and $u 1_T$ has a T -end. Notice that $\tau(1_T u) = \tau(u 1_T) = \tau(1_T u 1_T)$. Thus if u has either a

Y -beginning or a Y -end, $u \sim v$ for some v with T -beginning and T -end such that $\tau(v) = \tau(u)$. So $v \sim v^+ \sim v^*$ by the previous case. From $u \sim v$, we get $u^+ \sim v^+$ and $u^* \sim v^*$, since \sim is a bi-unary congruence, and so

$$u \sim v \sim v^+ \sim u^+$$

and

$$u \sim v \sim v^* \sim u^*$$

as required. \square

We now locate the full set of idempotents of $\mathcal{P}(T, Y)$ in the current case.

Proposition 2.3. *We have $E(\mathcal{P}(T, Y)) = Y'$.*

Proof. We only need to show that $E(\mathcal{P}(T, Y)) \subseteq Y'$. If $[x]^2 = [x]$, then $x^2 \sim x$. It follows from Lemma 1.6 that $\tau(x)^2 = \tau(x)$, which implies that $\tau(x) = 1_T$ since T is cancellative. So by Lemma 2.2 we obtain that $x \sim x^+$. Thus $[x] = [x^+] = [x]^+ \in Y'$. \square

Lemma 2.4. *Let $h \in U(T)$. Then for any $u \in T * Y$,*

$$hu^+h^{-1} \sim (hu)^+.$$

If in addition $\tau(u) = h^{-1}$, then

$$hu^+h^{-1} \sim hu.$$

Proof. Clearly $\tau(hu^+h^{-1}) = 1_T$. Applying Lemmas 1.2 and 2.2 we get
 $hu^+h^{-1} \sim (hu^+)^+hu^+h^{-1} = (hu)^+hu^+h^{-1} \sim (hu)^+(hu^+h^{-1})^+ = (hu^+h^{-1})^+(hu)^+$
 $= ((hu^+h^{-1})^+hu)^+ \sim (hu^+h^{-1}hu)^+ = (hu^+1_Tu)^+ \sim (hu^+u)^+ \sim (hu)^+.$
 If $\tau(u) = h^{-1}$, then $\tau(hu) = 1_T$ and $(hu)^+ \sim hu$ by Lemma 2.2. \square

To proceed to our main results, we need to consider factorisations in $T * Y$.

The next lemma, whose proof is clear, tells us that a factorisation wea , we or ea , where $w = w_0h_1w_1 \dots h_pw_p$ has T -end and fixed “length” p (meaning that consecutive symbols in the expression $w = w_0h_1w_1 \dots h_pw_p$ do not belong both to T or to Y), a has T -beginning and $e \in Y$, is unique; and dually.

Lemma 2.5. *Let $w, v, a, b \in T * Y$ be such that*

$$w = w_0h_1w_1 \dots h_pw_p, \quad v = v_0g_1v_1 \dots g_pv_p,$$

and

$$a = a_0e_1a_1 \dots e_na_n, \quad b = b_0f_1b_1 \dots f_nb_n,$$

where $w_0, v_0 \in T \cup \{\square\}$, $w_i, v_i \in T$ and $h_i, g_i \in Y$ for $1 \leq i \leq p$, and $a_{k-1}, b_{k-1} \in T$ and $e_k, f_k \in Y$ for $1 \leq k \leq n$, and $a_n, b_n \in T \cup \{\square\}$. Then for any $e, f \in Y$, if $wea = vfb$, then $w = v, e = f$ and $a = b$. Similar claims follow for equalities of the form $we = vf$ and $ea = fb$.

The analogue of the above is true if w, v have Y -ends, a, b have Y -beginnings and $wsa = vtb$, $ws = vt$ or $sa = tb$, for some $s, t \in T$.

We require a series of technical results on factorisations of the elements of $T * Y$ in case T is equidivisible, starting with a folklore result on the general case.

Lemma 2.6. *Let U and S be semigroups. If $w, a, v, b \in (U * S)^1$ with $wa = vb \neq 1$, then one of the following cases holds:*

- (I) $w = v$ and $a = b$;
- (II) there exists $u \in U * S$ such that $w = vu$ and $ua = b$;
- (III) there exists $u \in U * S$ such that $v = wu$ and $ub = a$;
- (IV) there exist $w', a' \in (U * S)^1$ and $e, f, g, h \in U$ or $e, f, g, h \in S$ such that $w = w'e$, $fa' = a$, $v = w'g$, $ha' = b$ and $ef = gh$.

Case (IV) may be refined to our case for T and Y .

Lemma 2.7. *Let T be equidivisible. If $w, a, v, b \in T * Y$ with $wa = vb$, then one of the following cases holds:*

- (I) $w = v$ and $a = b$;
- (II) there exists $u \in T * Y$ such that $w = vu$ and $ua = b$;
- (III) there exists $u \in T * Y$ such that $v = wu$ and $ub = a$;
- (IV) there exist $w', a' \in T * Y$ and $e, f, g, h \in Y$ such that $w = w'e$, $fa' = a$, $v = w'g$, $ha' = b$ and $ef = gh$.

Proof. In view of the previous lemma, we only need to analyse the case when $w', a' \in (T * Y)^1$ and e, f, g, h lie in T or in Y such that $w = w'e$, $fa' = a$, $v = w'g$, $ha' = b$ and $ef = gh$. Since T and Y are monoids we can assume $w', a' \in T * Y$. If $e, f, g, h \in Y$ we have the new case (IV). If $e, f, g, h \in T$ then as T is equidivisible we have three possibilities to discuss. If $e = g$, then $f = h$ and then we are in case (I). If $e = gu$ and $uf = h$ for some $u \in T$, then $w = w'gu = vu$ and $b = ufa' = ua$ and we are in case (II). Finally, if $g = eu$ and $uh = f$ for some $u \in T$, then $v = w'eu = wu$ and $a = uha' = ub$, the case (III). \square

In the proof of the next lemma we frequently call upon Lemmas 1.2 and 1.3, as well as the definition of \sim , without specific mention. The proof also requires Lemma 2.7 at various instances. We recall that, as stated in Theorem 2.10 in the Introduction, we need to show that $\mathcal{P}(T, Y)$ is adequate under the given conditions, in particular, we have to prove that $[a] \mathcal{R}^* [a]^+$ in $\mathcal{P}(T, Y)$. In our journey to establish this fact we must consider both kinds of generators of \sim at every stage: in this context there is no symmetry involved between H_ℓ and H_r .

Lemma 2.8. *Let T be equidivisible and let $x, a, z \in T * Y$ such that $xa \sim z$ via a basic step. Then there exist $h \in U(T)$ and $y, b \in T * Y$ such that*

$$z = yb, \quad \tau(x) = \tau(y)h, \quad h^{-1}\tau(b) = \tau(a) \quad \text{and} \quad xa^+ \sim yb^+h.$$

Proof. Since $xa \sim z$ via a basic step, there exist $c, d \in T * Y$ and $(\alpha, \beta) \in (H_\ell \cup H_r) \cup (H_\ell \cup H_r)^{-1}$ such that

$$xa = cad \quad \text{and} \quad c\beta d = z.$$

Thus $\alpha \sim \beta$ and, by Lemma 1.6, we have $\tau(\alpha) = \tau(\beta)$.

According to Lemma 2.7, the equality $xa = (c\alpha)d$ results in one of the factorisations (I), (II), (III), (IV). We discuss each in turn. Note that we only explicitly mention h in one sub-case, in the others $h = 1_T$.

(I) $x = c\alpha$ and $a = d$. Let $y = c\beta$ and $b = a$. Then $z = yb$. Now, $\tau(\alpha) = \tau(\beta)$, $x = c\alpha$ and $y = c\beta$ together imply $\tau(x) = \tau(y)$. Clearly $\tau(a) = \tau(b)$. Notice that $x = c\alpha \sim c\beta = y$ and so $xa^+ \sim ya^+ = yb^+$.

(II) $x = (c\alpha)s$ and $sa = d$, for some $s \in T * Y$. Let $y = c\beta s$ and $b = a$. Then $\tau(x) = \tau(y)$, $\tau(a) = \tau(b)$ and also $z = c\beta d = c\beta sa = yb$. As $x = c\alpha s \sim c\beta s = y$, we obtain $xa^+ \sim yb^+$.

(III) $c\alpha = xs$ and $sd = a$, for some $s \in T * Y$. We call upon Lemma 2.7 again to discuss the four possibilities for factorising $c\alpha = xs$.

(III.1) $c = x$ and $\alpha = s$. Put $y = c$ and $b = \beta d$. Then $z = yb$ and $\tau(x) = \tau(y)$. From $a = sd = \alpha d \sim \beta d = b$, we get $\tau(a) = \tau(b)$ and $a^+ \sim b^+$. We then obtain that $xa^+ \sim yb^+$.

(III.2) $c = xl$ and $l\alpha = s$ with $l \in T * Y$. Put $y = x$, $b = l\beta d$. Then $z = c\beta d = xl\beta d = yb$ and $\tau(x) = \tau(y)$. Also $a = sd = l\alpha d \sim l\beta d = b$, whence $\tau(a) = \tau(b)$ and $a^+ \sim b^+$. Thus $xa^+ \sim yb^+$.

(III.3) $x = ct$ and $ts = \alpha$, for some $t \in T * Y$. The fact that $(\alpha, \beta) \in (H_\ell \cup H_r) \cup (H_\ell \cup H_r)^{-1}$ leads the discussion to the following six cases:

(i) $\beta = \alpha^+ \alpha$. Put $y = c\alpha^+ t$ and $b = a$. Note that $\alpha d = tsd = ta$. Then $z = c\beta d = c\alpha^+ \alpha d = c\alpha^+ ta = yb$, $\tau(a) = \tau(b)$ and $\tau(y) = \tau(c\alpha^+ t) = \tau(ct) = \tau(x)$. Moreover

$$\begin{aligned} yb^+ &= c\alpha^+ tb^+ = c\alpha^+ ta^+ \sim c\alpha^+ (ta)^+ ta^+ \\ &= c\alpha^+ (\alpha d)^+ ta^+ = c(\alpha d)^+ ta^+ = c(ta)^+ ta^+ \sim cta^+ = xa^+. \end{aligned}$$

(ii) $\beta = \alpha \alpha^*$. Put $y = x$ and $b = s\alpha^* d$. Then $z = c\beta d = c\alpha \alpha^* d = cts\alpha^* d = xs\alpha^* d = yb$, $\tau(x) = \tau(y)$ and $\tau(b) = \tau(s\alpha^* d) = \tau(sd) = \tau(a)$. In addition,

$$\begin{aligned} yb^+ &= x(s\alpha^* d)^+ = ct(s\alpha^* d)^+ \sim ctt^*(s\alpha^* d)^+ = ct(t^* s\alpha^* d)^+ = ct(t^* s(ts)^* d)^+ \\ &= ct(t^* s(t^* s)^* d)^+ \sim ct(t^* sd)^+ = ctt^*(sd)^+ \sim ct(sd)^+ = xa^+. \end{aligned}$$

(iii) $\alpha = \beta^+ \beta$. Then $ts = \alpha = \beta^+ \beta$. In the following, we further use Lemma 2.7 to discuss $ts = \beta^+ \beta$ in four cases.

(iii.1) $t = \beta^+$ and $s = \beta$. Put $y = c$ and $b = a$. Then $x = c\beta^+$ and $\beta d = a$. We have $z = c\beta d = yb$, $\tau(y) = \tau(c) = \tau(c\beta^+) = \tau(x)$ and $\tau(a) = \tau(b)$. In addition, we have $yb^+ = ya^+ = c(\beta d)^+ \sim c(\beta^+ \beta d)^+ = c\beta^+ (\beta d)^+ = xa^+$.

(iii.2) $\beta^+ = tu$ and $u\beta = s$, for some $u \in T * Y$. In this case, we then must have $t, u \in Y$. Put $y = c$ and $b = \beta d$. Then $z = c\beta d = yb$, $\tau(y) = \tau(c) = \tau(ct) = \tau(x)$ and $\tau(b) = \tau(\beta d) = \tau(u\beta d) = \tau(sd) = \tau(a)$. Now

$$yb^+ = c(\beta d)^+ \sim c\beta^+ (\beta d)^+ = ctu(\beta d)^+ = ct(u\beta d)^+ = ct(sd)^+ = cta^+ = xa^+.$$

(iii.3) $t = \beta^+ u$ and $us = \beta$, for some $u \in T * Y$. Put $y = cu$ and $b = a$. Then $z = c\beta d = cusd = cua = yb$, $\tau(y) = \tau(cu) = \tau(c\beta^+ u) = \tau(ct) = \tau(x)$ and

$\tau(a) = \tau(b)$. Applying Lemma 1.2 again, we obtain

$$\begin{aligned} yb^+ &= cua^+ = cu(sd)^+ = cus^+(sd)^+ \sim c(us^+)^+us^+(sd)^+ \\ &= c(us)^+u(sd)^+ = c\beta^+u(sd)^+ = cta^+ = xa^+. \end{aligned}$$

(iii.4) $t = lg$, $pr = s$, $\beta^+ = le$ and $fr = \beta$, where $l, r \in T * Y$, $e, f, g, p \in Y$, and $gp = ef$. Note that $l \in Y$, and thus $t \in Y$ too. Put $y = c$ and $b = \beta d$. Then $z = c\beta d = yb$, $\tau(y) = \tau(c) = \tau(ct) = \tau(x)$ and $\tau(b) = \tau(\beta d) = \tau(frd) = \tau(rd) = \tau(prd) = \tau(sd) = \tau(a)$. We also have that

$$yb^+ = c(\beta d)^+ \sim c(\beta^+ \beta d)^+ = c(tsd)^+ = ct(sd)^+ = xa^+.$$

(iv) $\alpha = \beta\beta^*$. Recall that we have $x = ct$, $ts = \alpha = \beta\beta^*$ and $sd = a$.

In the following, once again we use Lemma 2.7 this time to discuss $ts = \beta\beta^*$ in four cases.

(iv.1) $t = \beta$ and $s = \beta^*$. Put $y = c\beta$ and $b = d$. Then $z = c\beta d = yb$. As $y = c\beta = ct = x$, we have $\tau(y) = \tau(x)$. We also have that $\tau(b) = \tau(d) = \tau(\beta^*d) = \tau(sd) = \tau(a)$. In addition, $yb^+ = c\beta d^+ \sim c\beta\beta^*d^+ = c\beta(\beta^*d)^+ = ct(sd)^+ = xa^+$.

(iv.2) $\beta = tu$ and $u\beta^* = s$, for some $u \in T * Y$. Put $y = x$ and $b = ud$. Then $z = c\beta d = ctud = xb = yb$, $\tau(a) = \tau(sd) = \tau(u\beta^*d) = \tau(ud) = \tau(b)$ and clearly $\tau(x) = \tau(y)$. Now

$$\begin{aligned} yb^+ &= x(ud)^+ = ct(ud)^+ \sim ctt^*(ud)^+ \sim ct(t^*ud)^+ \sim ct(t^*u(tu)^*d)^+ \\ &= ct(t^*u\beta^*d)^+ = ctt^*(u\beta^*d)^+ \sim ct(sd)^+ = xa^+. \end{aligned}$$

(iv.3) $t = \beta u$ and $us = \beta^*$, for some $u \in T * Y$. In this case we necessarily have $u, s \in Y$. Put $y = c\beta$ and $b = d$. Thus $z = c\beta d = yb$. Also, we have $\tau(y) = \tau(c\beta) = \tau(c\beta u) = \tau(ct) = \tau(x)$ and $\tau(b) = \tau(d) = \tau(sd) = \tau(a)$. Now

$$yb^+ = c\beta d^+ \sim c\beta\beta^*d^+ = c\beta usd^+ = ct(sd)^+ = xa^+.$$

(iv.4) $t = lg$, $pr = s$, $\beta = le$ and $fr = \beta^*$, where $l, r \in T * Y$ and $e, f, g, p \in Y$ with $gp = ef$. We then must have $r \in Y$ and thus $s \in Y$ too. Put $y = c\beta$ and $b = d$. Then $z = c\beta d = yb$. Also, $\tau(y) = \tau(c\beta) = \tau(cle) = \tau(cl) = \tau(clg) = \tau(ct) = \tau(x)$ and $\tau(b) = \tau(d) = \tau(sd) = \tau(a)$. In addition, we have

$$\begin{aligned} yb^+ &= c\beta d^+ \sim c\beta\beta^*d^+ = clefrd^+ = clgprd^+ = clgsd^+ \\ &= clgsd^+ = ctsd^+ = ct(sd)^+ = xa^+. \end{aligned}$$

(v) $(\alpha, \beta) = (1_Y, 1_T)$. Then $t = s = 1_Y$ as $ts = \alpha$. Put $y = c1_T$ and $b = d$. Then $z = c\beta d = c1_T d = yb$, $\tau(y) = \tau(c1_T) = \tau(c) = \tau(c1_Y) = \tau(ct) = \tau(x)$ and $\tau(a) = \tau(sd) = \tau(1_Y d) = \tau(d) = \tau(b)$. Also,

$$yb^+ = c1_T d^+ \sim c1_Y d^+ = c1_Y 1_Y d^+ = c1_Y(1_Y d)^+ = ct(sd)^+ = xa^+.$$

(vi) $(\alpha, \beta) = (1_T, 1_Y)$. Notice that this is the only situation where we may have $h \neq 1_T$. From $ts = \alpha = 1_T$ we must have that t, s in T and are mutually inverse. Put $y = c1_Y$ and $b = d$. Then $z = c\beta d = c1_Y d = yb$. Also, $\tau(x) = \tau(ct) =$

$\tau(c)\tau(t) = \tau(c1_Y)\tau(t) = \tau(y)t$ and $\tau(a) = \tau(sd) = s\tau(d) = s\tau(b) = t^{-1}\tau(b)$, where $ts = 1_T$. In addition,

$$\begin{aligned} xa^+ &= ct(sd)^+ \sim ct(sd)^+1_T = ct(sd)^+t^t \sim c(tsd)^+t \quad (\text{by Lemma 2.4}) \\ &= c(1_Td)^+t \sim c(1_Yd)^+t \sim c1_Yd^+t = yb^+t. \end{aligned}$$

(III.4) $c = x'g$, $ps' = \alpha$, $x = x'e$ and $fs' = s$, where $x', s' \in T * Y$, $e, f, g, p \in Y$ and $gp = ef$. Put $y = c$ and $b = \beta d$. Then $z = c\beta d = yb$, $\tau(y) = \tau(c) = \tau(x'g) = \tau(x') = \tau(x'e) = \tau(x)$ and $\tau(b) = \tau(\beta d) = \tau(\alpha d) = \tau(ps'd) = \tau(s'd) = \tau(fs'd) = \tau(sd) = \tau(a)$. In addition,

$$\begin{aligned} yb^+ &= c(\beta d)^+ \sim c(\alpha d)^+ = x'g(ps'd)^+ = x'gp(s'd)^+ \\ &= x'ef(s'd)^+ = x'e(fs'd)^+ = x(sd)^+ = xa^+. \end{aligned}$$

(IV) $x = x'e$, $fa' = a$, $c\alpha = x'g$, $\ell a' = d$ for some $x', a' \in T * Y$ and $e, f, g, \ell \in Y$ with $ef = g\ell$. Let $y = c\beta$ and $b = d$. Then $z = c\beta d = yb$, and also we have $\tau(x) = \tau(x'e) = \tau(x'g) = \tau(c\alpha) = \tau(c\beta) = \tau(y)$ and $\tau(b) = \tau(d) = \tau(\ell a') = \tau(fa') = \tau(a)$. Using Lemma 1.2, we see that $yb^+ = c\beta d^+ \sim c\alpha d^+ = x'g(\ell a')^+ = x'g\ell(a')^+ = x'ef(a')^+ = x'e(fa')^+ = xa^+$. \square

Lemma 2.9. *Let T be equidivisible. Let $h, k \in T$ and $x, a, y, b \in T * Y$ with $xa = yb$, $\tau(x) = \tau(y)h$ and $k\tau(b) = \tau(a)$. Then $hk = 1_T$ and $xa^+ \sim yb^+h$.*

Proof. For the first claim, notice that $\tau(y)\tau(b) = \tau(yb) = \tau(xa) = \tau(x)\tau(a) = \tau(y)hk\tau(b)$, so that as T is cancellative $hk = 1_T$.

According to Lemma 2.7 it is necessary to discuss $xa = yb$ in four cases.

(I) When $x = y$ and $a = b$, then $h = k = 1_T$ and the result is obvious.

(II) Assume that $x = yu$ and $ua = b$, for some $u \in T * Y$. Then $\tau(x) = \tau(yu) = \tau(y)\tau(u)$ together with $\tau(x) = \tau(y)h$ give $\tau(u) = h = k^{-1}$, as T is cancellative. Hence $\tau(ku) = \tau(k)\tau(u) = 1_T$. Now we deduce that

$$\begin{aligned} yb^+h &= y(ua)^+h \sim y1_T(ua)^+h = yhk(ua)^+h \\ &\sim yh(kua)^+ \quad (\text{by Lemma 2.4}) \\ &\sim yh((ku)^+a)^+ \quad (\text{by Lemma 2.2, as } \tau(ku) = 1_T) \\ &= yh(ku)^+a^+ \quad (\text{by Lemma 1.2}) \\ &\sim yhkua^+ \quad (\text{by Lemma 2.2 as } \tau(hu) = 1_T) \\ &\sim yua^+ \\ &= xa^+. \end{aligned}$$

(III) This case is dual to (II) and follows in a similar way.

(IV) To conclude, suppose that $x = w'e$, $fs' = a$, $y = w'g$, $ps' = b$, where $w', s' \in T * Y$, $e, f, g, p \in Y$ and $ef = gp$. Then $\tau(x) = \tau(w'e) = \tau(w') =$

$\tau(w'g) = \tau(y)$ and as $\tau(x) = \tau(y)h$ we have $h = 1_T$. Now,

$$\begin{aligned} xa^+ &= w'ea^+ = w'e(fs')^+ \\ &= w'ef(s')^+ \quad (\text{by Lemma 1.2}) \\ &= w'gp(s')^+ \\ &= y(ps')^+ \quad (\text{by Lemma 1.2}) \\ &= yb^+. \end{aligned}$$

The result follows. \square

We now present the main result of this article.

Theorem 2.10. *Let T be an equidivisible cancellative monoid acting on both sides upon a semilattice Y with identity, satisfying the compatibility conditions. Then $\mathcal{P}(T, Y)$ is adequate.*

Proof. The monoid $\mathcal{P}(T, Y)$ is Ehresmann with semilattice of projections Y' and, from Proposition 2.3, we know that $E(\mathcal{P}(T, Y)) = Y'$.

To show that $\mathcal{P}(T, Y)$ is adequate it remains to prove for any $[x], [y], [a] \in \mathcal{P}(T, Y)$, $[x][a] = [y][a]$ implies that $[x][a]^+ = [y][a]^+$, so that $[a] \mathcal{R}^* [a]^+$. If so, the dual argument will give that $[a] \mathcal{L}^* [a]^*$.

Suppose now that $[x], [y], [a] \in \mathcal{P}(T, Y)$ and $[x][a] = [y][a]$, whence $xa \sim ya$. Thus $xa = ya$ or there exists a sequence

$$xa = z'_0 \sim z'_1 \sim \dots \sim z'_n = ya$$

where $(z'_i, z'_{i+1}) = (c'_i \alpha_i d'_i, c'_i \beta_i d'_i)$ for some $(\alpha_i, \beta_i) \in (H_\ell \cup H_r) \cup (H_\ell \cup H_r)^{-1}$ and $c'_i, d'_i \in (T * Y)^1$.

If $xa = ya$ then as T is cancellative, certainly $\tau(x) = \tau(y)$, and so by Lemma 2.9 (with $k = h = 1_T$), we have that $xa^+ \sim ya^+ 1_T \sim ya^+$, giving $[x][a]^+ = [y][a]^+$.

Next, assume that we have a sequence as given above. For convenience, let S and W be either T or Y . We multiply each term of the sequence on the left by 1_S , where x has an S -beginning and on the right by 1_W , where a has a W -end. Let $z_i = 1_S z'_i 1_W$, $c_i = 1_S c'_i$ and $d_i = d'_i 1_W$ for $1 \leq i \leq n$. Now, $c_i, d_i \in T * Y$. We thus have a sequence

$$xa = z_0 \sim z_1 \sim \dots \sim z_n = (1_S y) a,$$

every step of which is basic. It follows from Lemma 1.6 that $\tau(x)\tau(a) = \tau(y)\tau(a)$ which implies that $\tau(x) = \tau(y)$. For convenience, put $x = y_0$ and $a = b_0$. By Lemma 2.8, for $1 \leq i \leq n$ there exist $y_i, b_i \in T * Y$ and $h_i \in U(T)$ such that $z_i = y_i b_i$, $\tau(y_{i-1}) = \tau(y_i) h_i$, $h_i^{-1} \tau(b_i) = \tau(b_{i-1})$ and $y_{i-1} b_{i-1}^+ \sim y_i b_i^+ h_i$. Hence

$$xa^+ = y_0 b_0^+ \sim y_1 b_1^+ h_1 \sim y_2 b_2^+ h_2 h_1 \sim \dots \sim y_n b_n^+ h_n h_{n-1} \dots h_1$$

and so by Lemma 1.6, we obtain that $\tau(x) = \tau(xa^+) = \tau(y_n)h$ where $h = h_n h_{n-1} \dots h_1 \in U(T)$.

From $\tau(x) = \tau(1_S y)$, we get $\tau(1_S y) = \tau(y_n)h$. As we have $(1_S y)a = z_n = y_n b_n$ it then follows that $h^{-1}\tau(b_n) = \tau(a)$, and now Lemma 2.9 yields $(1_S y)a^+ \sim y_n b_n^+ h$. Hence $xa^+ \sim y_n b_n^+ h \sim ya^+$ and so $[x][a]^+ = [y][a]^+$ as required. \square

From [3, Theorem 6.1] and the remarks preceding it, which together tell us that the free Ehresmann monoid on a set X is of the form $\mathcal{P}(X^*, Y)$, we immediately deduce the following corollary. Note that this result is also mentioned in the Section 6 (Remarks) of [14].

Corollary 2.11. *The free Ehresmann monoid on a set X is adequate, and hence coincides with the free adequate monoid on X .*

Corollary 2.12. *The quasi-variety of adequate monoids generates the variety of Ehresmann monoids.*

3. CHARACTERISATION OF $\mathcal{P}(T, Y)$

In this section we return to the consideration of \mathcal{P} -quadruples $\mathcal{T} = (T, Y, \cdot, \circ)$ for an *arbitrary* monoid T . We show that the Ehresmann monoid $\mathcal{P}(\mathcal{T}) = \mathcal{P}(T, Y)$ is unique, in the sense that it is exactly the initial object in a particular category, and do so after proving that the operator \mathcal{P} defines an expansion of a suitable category of Ehresmann monoids.

We start by recalling the definition of an expansion, here in the case of bi-unary monoids.

Definition 3.1. (*cf.* [1]) An *expansion* of a category \mathcal{C} of bi-unary monoids with $(2, 1, 1, 0)$ -morphisms is a “functorial cover”, i.e. a functor \mathcal{E} from \mathcal{C} to itself along with a natural transformation π from \mathcal{E} to the identity functor of \mathcal{C} such that, for each object M of \mathcal{C} , the morphism π_M is onto. Thus, for any objects M_1 and M_2 of \mathcal{C} , the following diagram commutes:

$$\begin{array}{ccc} \mathcal{E}(M_1) & \xrightarrow{\mathcal{E}(\psi)} & \mathcal{E}(M_2) \\ \pi_{M_1} \downarrow & \circlearrowleft & \downarrow \pi_{M_2} \\ M_1 & \xrightarrow[\psi]{} & M_2 \end{array}$$

Let $\mathcal{T} = (T, Y, \cdot, \circ)$ be a \mathcal{P} -quadruple. By Theorem 1.7, we may construct the Ehresmann monoid $\mathcal{P}(\mathcal{T}) := \mathcal{P}(T, Y)$ associated with \mathcal{T} . We have that $\mathcal{P}(\mathcal{T})$ is T' -generated with semilattice of projections Y' , and $\nu_{\mathcal{T}, T} = \nu_{\mathcal{T}}|_T : T \rightarrow T'$ and $\nu_{\mathcal{T}, Y} = \nu_{\mathcal{T}}|_Y : Y \rightarrow Y'$ are isomorphisms, where $T' = \{[t] : t \in T\}$ and $Y' = \{[y] : y \in Y\}$. Further, Lemmas 1.2 and 1.3 give that for any $t \in T$ and $y \in Y$,

$$(t \cdot y)\nu_{\mathcal{T}, Y} = t\nu_{\mathcal{T}, T} \cdot y\nu_{\mathcal{T}, Y} \quad \text{and} \quad (y \circ t)\nu_{\mathcal{T}, Y} = y\nu_{\mathcal{T}, Y} \circ t\nu_{\mathcal{T}, T}.$$

Lemma 3.2. *Let $\mathcal{T}_1 = (T_1, Y_1, \cdot, \circ)$ and $\mathcal{T}_2 = (T_2, Y_2, \cdot, \circ)$ be \mathcal{P} -quadruples, and suppose that $\psi_T : T_1 \rightarrow T_2$ and $\psi_Y : Y_1 \rightarrow Y_2$ are monoid morphisms such that for all $t \in T_1$ and $y \in Y_1$ we have*

$$(t \cdot y)\psi_Y = t\psi_T \cdot y\psi_Y \quad \text{and} \quad (y \circ t)\psi_Y = y\psi_Y \circ t\psi_T.$$

*Let $\psi : T_1 * Y_1 \rightarrow T_2 * Y_2$ be the (semigroup) morphism that extends ψ_T and ψ_Y . Then ψ is a $(2, 1, 1)$ -morphism, and*

$$\mathcal{P}_\psi : \mathcal{P}(\mathcal{T}_1) \rightarrow \mathcal{P}(\mathcal{T}_2)$$

given by

$$[u]\mathcal{P}_\psi = [u\psi]$$

is a $(2, 1, 1, 0)$ -morphism. If ψ_T and ψ_Y are both onto, then so is \mathcal{P}_ψ .

Proof. To show that \mathcal{P}_ψ is well-defined we are required to prove that the congruence \sim giving $\mathcal{P}(\mathcal{T}_1)$ is such that $\sim \subseteq \ker(\psi\nu_{\mathcal{T}_2})$. To do so it is sufficient to see that the generating set $H_\ell \cup H_r$ of \sim lies in $\ker(\psi\nu_{\mathcal{T}_2})$.

First we prove that, for all $u \in T_1 * Y_1$ and $y \in Y_1$,

$$(3.1) \quad (u \cdot y)\psi_Y = u\psi \cdot y\psi_Y \quad \text{and} \quad (y \circ u)\psi_Y = y\psi_Y \circ u\psi.$$

By hypothesis this holds for $u \in T_1$ and if $u \in Y_1$, we have

$$(u \cdot y)\psi_Y = (uy)\psi_Y = (u\psi_Y)(y\psi_Y) = u\psi_Y \cdot y\psi_Y = u\psi \cdot y\psi_Y.$$

Then, by induction on the minimal number of generators from $T_1 \cup Y_1$ of $u \in T_1 * Y_1$, we obtain one half of (3.1) for all $u \in T_1 * Y_1$ and $y \in Y_1$; the other equality may be shown similarly.

Now, for $u \in T_1 * Y_1$, we get

$$u^+\psi = (u \cdot 1_{Y_1})\psi = (u \cdot 1_{Y_1})\psi_Y = u\psi \cdot 1_{Y_1}\psi_Y = u\psi \cdot 1_{Y_2} = (u\psi)^+.$$

Similarly, $u^*\psi = (u\psi)^*$, thus verifying that ψ is a $(2, 1, 1)$ -morphism.

It follows that

$$(u^+u)\psi = (u^+\psi)(u\psi) = (u\psi)^+(u\psi) \sim u\psi.$$

Analogously, $(uu^*)\psi \sim u\psi$. Also $1_{T_1}\psi = 1_{T_1}\psi_T = 1_{T_2} \sim 1_{Y_2} = 1_{Y_1}\psi_Y = 1_{Y_1}\psi$.

Therefore \sim is contained in $\ker(\psi\nu_{\mathcal{T}_2})$, and so \mathcal{P}_ψ is indeed well-defined and clearly is then a semigroup morphism. Given $u \in T_1 * Y_1$,

$$[u]^+\mathcal{P}_\psi = [u^+]\mathcal{P}_\psi = [u^+\psi] = [(u\psi)^+] = [u\psi]^+ = ([u]\mathcal{P}_\psi)^+.$$

In a similar way, \mathcal{P}_ψ respects $*$. As $[1_{T_1}]\mathcal{P}_\psi = [1_{T_1}\psi] = [1_{T_2}]$, we have that \mathcal{P}_ψ is a $(2, 1, 1, 0)$ -morphism.

If both ψ_T and ψ_Y are onto, then so is ψ and hence also \mathcal{P}_ψ . \square

Now let \mathcal{C} be the category whose objects are triples (M, T, Y) , where M is a T -generated Ehresmann monoid with semilattice of projections Y (the objects are over-defined as given M we know Y , however it is useful to mention Y explicitly); by a morphism $\varphi : (M_1, T_1, Y_1) \rightarrow (M_2, T_2, Y_2)$ of \mathcal{C} we mean that $\varphi : M_1 \rightarrow M_2$ is

a $(2, 1, 1, 0)$ -morphism such that $T_1\varphi = T_2$ and $Y_1\varphi = Y_2$. The composition in \mathcal{C} is the usual composition of maps.

Definition 3.3. The category \mathcal{C} defined above is called the category of *marked* Ehresmann monoids.

Let (M, T, Y) be an object of \mathcal{C} . As seen in Lemma 1.1, we then have a \mathcal{P} -quadruple $\mathcal{T} = (T, Y, \cdot, \circ)$ where the actions are the standard ones induced by the action of T on Y in M . By Theorem 1.7, $\mathcal{P}(M, T, Y) := (\mathcal{P}(\mathcal{T}), T\nu_{\mathcal{T}, T}, Y\nu_{\mathcal{T}, Y})$ is an object of \mathcal{C} . Now let (M_1, T_1, Y_1) and (M_2, T_2, Y_2) be objects of \mathcal{C} , and let \mathcal{T}_1 and \mathcal{T}_2 be the corresponding \mathcal{P} -quadruples. Suppose that $\psi : (M_1, T_1, Y_1) \rightarrow (M_2, T_2, Y_2)$ is a morphism of \mathcal{C} . Define $\psi_T : T_1 \rightarrow T_2$ and $\psi_Y : Y_1 \rightarrow Y_2$ as the morphisms induced by ψ . Since $\psi : M_1 \rightarrow M_2$ is a $(2, 1, 1, 0)$ -morphism, it is clear that, for any $t \in T_1$ and $y \in Y_1$,

$$(t \cdot y)\psi_Y = t\psi_T \cdot y\psi_Y \quad \text{and} \quad (y \circ t)\psi_Y = y\psi_Y \circ t\psi_T.$$

With some abuse of notation, letting ψ also denote the semigroup morphism extension of ψ_T and ψ_Y to $T_1 * Y_1$, Lemma 3.2 gives that

$$\mathcal{P}(\psi) := \mathcal{P}_\psi : \mathcal{P}(\mathcal{T}_1) \rightarrow \mathcal{P}(\mathcal{T}_2)$$

is a morphism of Ehresmann monoids. Clearly $(T_1\nu_{\mathcal{T}_1, T_1})\mathcal{P}_\psi = T_2\nu_{\mathcal{T}_2, T_2}$ and $(Y_1\nu_{\mathcal{T}_1, Y_1})\mathcal{P}_\psi = Y_2\nu_{\mathcal{T}_2, Y_2}$. Thus \mathcal{P}_ψ is a morphism in \mathcal{C} .

It follows easily that \mathcal{P} is a functor from \mathcal{C} to \mathcal{C} .

Theorem 3.4. *The functor \mathcal{P} determines an expansion of the category of marked Ehresmann monoids.*

Proof. Let $\mathcal{M} = (M, T, Y)$ be an object of \mathcal{C} . From [3, Theorem 5.2], the morphism $\iota : T * Y \rightarrow M$ extending the inclusion maps $\iota_T : T \rightarrow M$ and $\iota_Y : Y \rightarrow M$ factors through $\mathcal{P}(M, T, Y)$ to produce an onto morphism

$$\pi_{\mathcal{M}} : \mathcal{P}(M, T, Y) \rightarrow (M, T, Y)$$

given by

$$[u_1 \dots u_n]\pi_{\mathcal{M}} = u_1 \dots u_n,$$

where $u_1, \dots, u_n \in T \cup Y$ for $1 \leq i \leq n$, the product on the left hand side is in $T * Y$, and that on the right hand side is taken in M .

It is easy to check that if $\mathcal{M}_1 = (M_1, T_1, Y_1)$ and $\mathcal{M}_2 = (M_2, T_2, Y_2)$ are objects of \mathcal{C} and $\psi : \mathcal{M}_1 \rightarrow \mathcal{M}_2$ is a morphism of \mathcal{C} , then $\pi_{\mathcal{M}_1}\psi = \mathcal{P}_\psi\pi_{\mathcal{M}_2}$, so that the diagram

$$\begin{array}{ccc} \mathcal{P}(\mathcal{M}_1) & \xrightarrow{\mathcal{P}_\psi} & \mathcal{P}(\mathcal{M}_2) \\ \pi_{\mathcal{M}_1} \downarrow & \circlearrowleft & \downarrow \pi_{\mathcal{M}_2} \\ \mathcal{M}_1 & \xrightarrow[\psi]{} & \mathcal{M}_2 \end{array}$$

commutes as required. □

We point out that the expansion determined by \mathcal{P} has associated natural transformation π , where, for each object \mathcal{M} of \mathcal{C} , the morphism $\pi_{\mathcal{M}} : \mathcal{P}(\mathcal{M}) \rightarrow \mathcal{M}$ is defined in the proof of Theorem 3.4.

We now fix a \mathcal{P} -quadruple $\mathcal{T} = (T, Y, \cdot, \circ)$, and define a category $\mathcal{C}(\mathcal{T})$ as follows: the objects are quintuples $(M, S, E, \theta_S, \theta_E)$ where (M, S, E) is an object of \mathcal{C} , $\theta_S : T \rightarrow S$ and $\theta_E : Y \rightarrow E$ are onto monoid morphisms such that for all $t \in T$ and $y \in Y$,

$$(3.2) \quad (t \cdot y)\theta_E = t\theta_S \cdot y\theta_E \quad \text{and} \quad (y \circ t)\theta_E = y\theta_E \circ t\theta_S.$$

A morphism $\psi : (M_1, S_1, E_1, \theta_{S_1}, \theta_{E_1}) \rightarrow (M_2, S_2, E_2, \theta_{S_2}, \theta_{E_2})$ of $\mathcal{C}(\mathcal{T})$ is simply a morphism $\psi : (M_1, S_1, E_1) \rightarrow (M_2, S_2, E_2)$ in \mathcal{C} such that $\theta_{S_1}\psi_S = \theta_{S_2}$ and $\theta_{E_1}\psi_E = \theta_{E_2}$. This can be represented in terms of commutativity of diagrams as follows:

$$\begin{array}{ccc} & T & \\ \theta_{S_1} \swarrow & \circlearrowright & \searrow \theta_{S_2} \\ S_1 & \xrightarrow{\psi_S} & S_2 \end{array} \quad \begin{array}{ccc} & Y & \\ \theta_{E_1} \swarrow & \circlearrowright & \searrow \theta_{E_2} \\ E_1 & \xrightarrow{\psi_E} & E_2 \end{array}$$

Definition 3.5. The category $\mathcal{C}(\mathcal{T})$ is called the category of \mathcal{T} -marked Ehresmann monoids.

Theorem 3.6. Let $\mathcal{T} = (T, Y, \cdot, \circ)$ be a \mathcal{P} -quadruple. Then

$$\mathcal{Q}(\mathcal{T}) = (\mathcal{P}(T, Y), T\nu_{\mathcal{T}, T}, Y\nu_{\mathcal{T}, Y}, \nu_{\mathcal{T}, T}, \nu_{\mathcal{T}, Y})$$

is the initial object in the category $\mathcal{C}(\mathcal{T})$.

Proof. Clearly $\mathcal{Q}(\mathcal{T})$ is an object in $\mathcal{C}(\mathcal{T})$. Let $(M, S, E, \theta_S, \theta_E)$ be an object in $\mathcal{C}(\mathcal{T})$. We must show that there is a unique morphism in $\mathcal{C}(\mathcal{T})$ from $\mathcal{Q}(\mathcal{T})$ to $(M, S, E, \theta_S, \theta_E)$.

Let $\mathcal{U} = (S, E, \cdot, \circ)$ be the \mathcal{P} -quadruple determined by M and notice that by the very definition of $\mathcal{C}(\mathcal{T})$ we have onto monoid morphisms $\theta_S : T \rightarrow S$ and $\theta_E : Y \rightarrow E$ satisfying (3.2) for all $t \in T$ and $y \in Y$.

Let $\theta : T * Y \rightarrow S * E$ be the natural extension of θ_S, θ_E . By Lemma 3.2 we have an onto $(2, 1, 1, 0)$ -morphism

$$\mathcal{P}_\theta : \mathcal{P}(\mathcal{T}) \rightarrow \mathcal{P}(\mathcal{U})$$

in \mathcal{C} given by

$$[u]\mathcal{P}_\theta = [u\theta].$$

By Theorem 3.4, there is an onto $(2, 1, 1, 0)$ -morphism $\pi : \mathcal{P}(\mathcal{U}) \rightarrow M$ that lies in \mathcal{C} and hence $\mathcal{P}_\theta\pi$ lies in \mathcal{C} . For any $t \in T$ we have

$$t\nu_{\mathcal{T}, T}\mathcal{P}_\theta\pi = [t]\mathcal{P}_\theta\pi = [t\theta]\pi = t\theta = t\theta_S,$$

and similarly $y\nu_{\mathcal{T}, Y}\mathcal{P}_\theta\pi = y\theta_E$. Hence $\mathcal{P}_\theta\pi$ lies in $\mathcal{C}(\mathcal{T})$.

Now let $\psi : \mathcal{P}(\mathcal{T}) \rightarrow M$ be any morphism in $\mathcal{C}(\mathcal{T})$. By definition, we must have that for any $t \in T$ and $y \in Y$,

$$t\nu_{\mathcal{T},T}\psi = t\theta_S \quad \text{and} \quad y\nu_{\mathcal{T},Y}\psi = y\theta_E.$$

Then $\mathcal{P}_\theta\pi$ and ψ agree on a set of generators of $\mathcal{P}(\mathcal{T})$, so they must be equal, establishing the uniqueness of $\mathcal{P}_\theta\pi$. The result follows. \square

It is worth remarking that we have shown that if $(M, S, E, \theta_S, \theta_E)$ is an object in $\mathcal{C}(\mathcal{T})$ for some \mathcal{P} -quadruple \mathcal{T} , then the expansion $\mathcal{P}(M, S, E)$ of (M, S, E) also lies in $\mathcal{C}(\mathcal{T})$.

4. OPEN QUESTIONS

We point to some natural questions that arise from our work.

Let $\mathcal{T} = (T, Y, \cdot, \circ)$ be a \mathcal{P} -quadruple.

Open Question 4.1. We have proved that if T is an equidivisible cancellative monoid then the monoid $\mathcal{P}(T, Y)$ is adequate. *Is $\mathcal{P}(T, Y)$ adequate when T is an arbitrary cancellative monoid?*

Open Question 4.2. We know that, unlike the one-sided case, $\mathcal{P}(T, Y)$ does not have uniqueness of T -normal forms. *Is there a uniqueness of T -normal forms of minimal length in $\mathcal{P}(T, Y)$, at least in the cancellative and equidivisible case?*

Open Question 4.3. In the two-sided case, the monoid $\mathcal{P}(T, Y)$ may be thought of as the analogue for Ehresmann monoids of the semidirect product construction known for inverse and for restriction monoids. Similarly for the one-sided case by considering $\mathcal{P}_\ell(T, Y)$, left Ehresmann monoids, and left restriction monoids. *What might be the analogue of the McAlister P -semigroup construction for $\mathcal{P}(T, Y)$ and $\mathcal{P}_\ell(T, Y)$?* Observe that in the two-sided case this may well involve the study of partial actions, in view of the results in [4] for restriction monoids.

Open Question 4.4. That every strongly T -proper Ehresmann monoid M is T -proper is known. *What is the precise connection in the one- and the two-sided cases between the concepts of being strongly T -proper, T -proper, having uniqueness of T -normal forms (the latter in the one-sided case) and indeed other, natural, concepts of T -properness?*

ACKNOWLEDGEMENTS

The first three authors acknowledge financial support of FCT, through project UID/MULTI/04621/2013 of CEMAT-Ciências. The third author thanks the Faculdade de Ciências da Universidade de Lisboa for its hospitality during her visit in June 2016. The fourth author acknowledges NSFC (Grant No:11501331) and Shandong Province Natural Science Foundation (Grant No: BS2015SF002).

REFERENCES

- [1] J.-C. Birget and J. Rhodes, ‘Almost finite expansions of arbitrary semigroups’, *J. Pure Appl. Algebra* **32** (1984), 239-287.
- [2] M.J.J. Branco, G.M.S. Gomes and V. Gould, ‘Left adequate and left Ehresmann monoids’, *Internat. J. Algebra Comput.* **21** (2011), 1259-1284.
- [3] M.J.J. Branco, G.M.S. Gomes and V. Gould, ‘Ehresmann monoids’, *J. Algebra* **443** (2015), 1-34.
- [4] C. Cornock and V. Gould, ‘Proper two-sided restriction semigroups and partial actions’, *J. Pure and Applied Algebra* **216** (2012), 935-949.
- [5] C. Ehresmann, “Oeuvres complètes et commentées” (A. C. Ehresmann, Ed.), Suppl. Cahiers Top Géom. Diff., Amiens, 1980-1984.
- [6] J. Fountain, ‘Adequate semigroups’, *Proc. Edinb. Math. Soc.* **2** (1979), 113-125.
- [7] J. Fountain, ‘A class of right PP monoids’, *Quart. J. Math. Oxford* **28** (1977), 285-300.
- [8] J. Fountain, ‘Free right type A semigroups’, *Glasgow Math. J.* **33** (1991), 135-148.
- [9] J. Fountain, G.M.S. Gomes and V. Gould, ‘The free ample monoid’, *Internat. J. Algebra Comput.* **19** (2009), 527-554.
- [10] G.M.S. Gomes and V. Gould, ‘Left adequate and left Ehresmann monoids II’, *J. Algebra* **348** (2011), 171-195.
- [11] J.M. Howie, ‘Fundamentals of semigroup theory’, *Oxford University Press* 1995.
- [12] P. Jones, ‘Almost perfect restriction semigroups’, *J. Algebra* **445** (2016), 193-220.
- [13] P. Jones, ‘An elementary proof of the existence of proper covers for Ehresmann monoids’, *Private communication* (2014).
- [14] M. Kambites, ‘Free adequate semigroups’, *J. Aust. Math. Soc.* **191** (2011), 365-390.
- [15] M. Kambites, ‘Retracts of trees and free left adequate semigroups’, *Proc. Edinb. Math. Soc.* **54** (2011), 731-747.
- [16] G. Kudryavtseva, ‘Partial monoid actions and a class of restriction semigroups’, *J. Algebra* **429** (2015), 342-370.
- [17] M.V. Lawson, ‘Semigroups and ordered categories I: the reduced case’, *J. Algebra* **141** (1991), 422-462.
- [18] F.W. Levi, ‘On semigroups’, *Bull. Calcutta Math. Soc.* **36** (1944), 141-146.
- [19] D.B. McAlister, ‘Groups, semilattices and inverse semigroups’, *Trans. Amer. Math. Soc.* **192** (1974), 227-244.
- [20] D.B. McAlister, ‘Groups, semilattices and inverse semigroups II’, *Trans. Amer. Math. Soc.* **192** (1974), 351-370.
- [21] J.N. Mordeson and P.S. Nair, ‘Fuzzy Mathematics: an introduction for engineers and scientists’, *Springer Science and Business Media* (2001).
- [22] W.D. Munn, ‘Free inverse semigroups’, *Proc. London Math. Soc.* **29** (1974), 385-404.
- [23] L. O’Carroll, ‘Embedding theorems for proper inverse semigroups’, *J. Algebra* **42** (1976), 26-40.
- [24] H.E. Scheiblich, ‘Free inverse semigroups’, *Semigroup Forum* **4** (1972), 351-358.

DEPARTAMENTO DE MATEMÁTICA AND CEMAT-CIÊNCIAS, FACULDADE DE CIÊNCIAS,
UNIVERSIDADE DE LISBOA, 1749-016, LISBOA, PORTUGAL

E-mail address: `mjbranco@fc.ul.pt`, `gmcunha@fc.ul.pt`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, HESLINGTON, YORK YO10
5DD, UK

E-mail address: `victoria.gould@york.ac.uk`

COLLEGE OF MATHEMATICS AND SYSTEMS SCIENCE, SHANGDONG UNIVERSITY OF SCI-
ENCE AND TECHNOLOGY, QINGDAO, 266590, P. R. CHINA

E-mail address: `yanhuiwang@sdust.edu.cn`