

This is a repository copy of *Tolerance and metabolic responses of Cyanidiophytina (Rhodophyta) towards exposition to Cl4K2Pd and AuCl4K*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/135760/>

Version: Accepted Version

Article:

Sirakov, Maria, Toscano, Elena, Iovinella, Manuela et al. (3 more authors) (2018)
Tolerance and metabolic responses of Cyanidiophytina (Rhodophyta) towards exposition to Cl4K2Pd and AuCl4K. *Phycology International*.

<https://doi.org/10.4081/phycol.2018.55>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

**Tolerance and metabolic responses of Cyanidiophytina (Rhodophyta) towards
exposition to $\text{Cl}_4\text{K}_2\text{Pd}$ and AuCl_4K**

Maria Sirakov,¹ Elena Toscano,¹ Manuela Iovinella,² Seth J. Davis,² Milena Petriccione,³
Claudia Ciniglia¹

¹DISTABIF, “L. Vanvitelli” University of Caserta, Caserta, Italy; ²Department of Biology,
University of York, York, UK; ³Department of Fruit Tree Research Unit, Council for
Agricultural Research and Agricultural Economy Analysis, Caserta, Italy

Correspondence: Claudia Ciniglia, DISTABIF, “L. Vanvitelli” University of Caserta,
Caserta, Italy.

Tel.: +39.0823274582 – Fax: + 39.0823274571

E-mail: claudia.ciniglia@unicampania.it

Key words: Metal tolerance; Red algae; Rare Earth elements; Palladium; Gold.

Contributions: MS, ET: conduction of experiments, analysis of results, contribution to draft
writings; MP: experiments on oxidative stress, analysis of results; MI, SJD: original concept,
provision of resources; CC: original concept, provision of resources, draft editing.

Conflict of interest: the authors declare no potential conflict of interest.

*This article has been accepted for publication and undergone full peer review but has not been through the
copyediting, typesetting, pagination and proofreading process, which may lead to differences between this ver-
sion and the final one.*

Please cite this article as doi: 10.4081/phycol.2018.55

ABSTRACT

Polyextremophilic algae, such as unicellular red algae known as *Cyanidiophyceae*, have the intrinsic capacity to selectively mobilize and adsorb metals, since they are adapted to live in geothermal and volcanic sites characterized by elevated concentration of heavy and rare metals. In this work we evaluated the ability of 3 strains of the genus *Galdieria* (*G. maxima*, *G. sulphuraria*, *G. phlegrea*) along with one strain of *Cyanidium caldarium* to tolerate different concentrations of rare metal as $\text{Cl}_4\text{K}_2\text{Pd}$ and AuCl_4K by monitoring changes in algal growth in culture exposed to different concentration of each metal and investigating algae metabolic response and possible oxidative stress induced by these metals.

INTRODUCTION

In the last decade there is a remarkable and growing demand for Rare Earth Elements (REE) due to their large use in the superconductors, catalysts and electronic industry. On the other side, the issue of their discharge in the environment and the suitability of recycling REE from the electronic waste (e-waste) is of interest of all the population because of their hazard for environment and health, besides their economic value. These issues become evident to the government of the different States and electronic industries prone to develop new methods of removal from the environment, to recycle and re-input of REE in the productive cycle of a “closed loop economy”¹⁻⁴. Recently, biological methods have been developed to ensure the recovery of small quantities of these mineral materials and wastewater systems, using mainly bacteria⁵ or plants known for their ability to immobilize heavy metals in the cell wall and compartmentalization in vacuoles. Interestingly, polyextremophilic algae have the intrinsic properties that make them capable of selective removal and concentration of metals, thanks to their adaptation to live in geothermal and volcanic sites⁶⁻⁸. Geothermal fluids leach out of the hot volcanic rocks and are enriched by enormous amounts of minerals and metals, including lithium, sulfur, boric acid and precious metals such as gold, platinum, palladium and silver⁹. *Cyanidiophyceae*, unicellular red algae, survive in extreme conditions, very low pH (0.0-3.0) and high temperatures (37-55 °C), and colonize acid and hydrothermal sites, but also rocks and muddy soil around hot ponds¹⁰. They belong to 3 genera: *Cyanidioschyzon*, *Cyanidium* and *Galdieria*, which differ in size, cellular shape and growth conditions. *C. merolae* is the only species belonging to *Cyanidioschyzon* genus and differs from the other two groups being lacking of cell wall and dividing by binary fission¹¹. Both *Cyanidium* and *Galdieria* are able to grow both on ammonia and nitrate, the former is an obligatory autotroph whereas the species belonging to *Galdieria* tolerate high concentrations of salts¹² and can grow

Accepted paper

autotrophically and heterotrophically, thus making *Galdieria* the best candidates for biotechnological application into the recovery of REE as it has been already proposed in previous studies from other groups^{13,17}. These algae are one of the few eukaryotes capable of adapting to a very acidic environment. Because of the high temperature and acidic conditions, the environment that these algae live in is usually rich in metals; *Galdieria sulphuraria* is the most suitable alga of all of the Cyanidiophitina for biotechnology experiments and applications because it is the only one of these algae that can grow autotrophically as well as heterotrophically, using over 27 different kinds of sugar and polyols to produce a huge biomass and beneficial compounds¹³. In this report, we have evaluated the ability of *G. maxima*, *G. sulphuraria*, *G. phlegrea* and *C. caldarium* to tolerate different concentrations of REE (such as palladium-Cl₄K₂Pd and Gold-AuCl₄K) by analyzing Maximum Growth Rate (MGR) and the inverse of Generation Time (1/GT)¹⁴. We also investigated metabolic response and possible oxidative stress induced by the metals, by monitoring superoxide dismutase (SOD), catalase (CAT), ascorbate peroxidase (APX) activities^{15,16}.

MATERIALS AND METHODS

Algal strains and culture conditions

The algal strains used in this study belong to the algal collection of the University of Naples (www.acuf.net), namely ACUF 3.4.5 (*G. maxima*), ACUF 7.6.21 (*G. phlegrea*), ACUF9.2.11 (*G. sulphuraria*) and ACUF 626 (*C. caldarium*). All strains were maintained in liquid culture in Allen medium¹⁸, pH 1.5, at 37°C on a plexiglass shaking apparatus under a photon irradiance of 150 μmol photons m⁻² s⁻¹ with continuous light provided by cool-light fluorescent lamps (Philips TLD30w/55).

Experimental set up

3ml of each culture at exponential phase were transferred to a 24well-plate and the medium was supplemented with different metals concentrations (0,1gg/L-1g/L). Cell growth was assessed by recording the optical density (OD) at 750 nm (Bausch & Lomb Spectronic 20), at day 0 and day 4; MGR and 1/GT were calculated according to the formula $[(\text{Log}_{10} \text{OD}_{\text{day4}} - \text{Log}_{10} \text{OD}_{\text{day0}})/4]$ and $1/[\text{Ln}(2)/\text{MGR}]$, respectively. Three replicates were carried out for each test.

Enzyme activity

For measurement of enzyme activity algal cultures were transferred in falcon containing the metal at 1g/L in a final volume of 30 ml and pellet was harvested by centrifugation at 14000 rpm for 10min after 96h of exposure. Pellet was washed using KH_2PO_4 (0.1M pH7.8) followed by centrifugation at 12000 rpm for 4 min at 4°C, twice. Proteins were extracted homogenizing the sample with liquid nitrogen using mortar and pestel, resuspending the powder obtained in 3ml of Lysis Buffer (KH_2PO_4 0.5M pH7.8, DTT 2mM, EDTA 1mM, PMSF 1mM, PEG 1,25mM), samples were centrifuged at 14000rpm for 20 min at 4°C and the supernatant was used for measurement after Bradford quantification.

SOD measurement was determined adding 100 μl of protein extract to the buffer containing KH_2PO_4 50mM pH7.8, Na-EDTA 0.1mM pH7.0, Metionin 13mM, Nitro Blue Tetrazolium (NBT) 75 μM , Riboflavin 2 μM), after 15min exposure to the light samples were read at 560nm using a Spectrophotometer. Enzymatic activity was expressed as units/g of dry weight. CAT measurement was performed adding 200 μl of protein extract to the buffer containing KH_2PO_4 50mM pH7.8, H_2O_2 20mM and H_2O up to 1ml. Absorbance was recorded at 240nm for 100 sec. Enzymatic activity was expressed as nmol H_2O_2 /g of fresh weight. APX measurement was obtained adding 100 μl of protein extract to the buffer containing KH_2PO_4 100mM pH7, Na-EDTA 0,66mM, Ascorbic Acid 0.33mM, H_2O_2 0.35mM. Absorbance at 290nm for 100sec was recorded. Enzymatic activity was expressed as μmol ascorbate/g of fresh weight.

Each condition for each experimental approach was tested 3 times independently.

RESULTS AND DISCUSSION

Cyanidiophyceae are polyextremophilic micro-algae with an intrinsic ability to uptake metals, involving both active and passive mechanisms. Heavy, rare or precious metals can influence algae physiology in various ways, likely inhibiting different physiological processes. In order to evaluate the suitability of *Cyanidiophyceae* for biotechnological application to effectively recover REE, we tested their tolerance to $\text{Cl}_4\text{K}_2\text{Pd}$ and AuCl_4K monitoring the growth and metabolic response of 4 different strains, exposed to each of these metals ranging from 0.1g/L up to 10g / L. The growth was evaluated after 4 days since the single metal exposure and the results are expressed in the form of maximum growth rate (MGR) and the inverse of the Generation Time ($1/\text{GT}$). *G. maxima* was more sensitive to AuCl_4K showing a continuous decline in growth rate (Fig. 1A), while the growth of *G. maxima*

declined at $\text{Cl}_4\text{K}_2\text{Pd}$ 0,1g/L then sharply increased at higher concentrations (Fig.1A). both gold and palladium induced a significant decrease in growth rate of *G. phlegrea* from 0.1g/L to 10g/L (Fig.1B). As shown in Fig.1C, the MGR and 1/GT in *G. sulphuraria* dropped at 0,1g /L in both metals; the algal growth was steady until 1g/L AuCl_4K , then sharply dropped at higher concentration; viceversa, the growth rate significantly and progressively increased at palladium concentrations from 1 to 10 g/L. *C. caldarium* showed a high tolerance to $\text{Cl}_4\text{K}_2\text{Pd}$ whereas AuCl_4K inhibited algal growth with the increase of metal concentration (Fig.1D).

ROS scavenging activities of SOD, CAT and APX were assessed in all algae under $\text{Cl}_4\text{K}_2\text{Pd}$ and AuCl_4K at a concentration of 1g/L, after 24 hours. The antioxidant activity can be considered a measure of the effectiveness of the cell to respond to the impact of the metal, increasing its tolerance as protective mechanisms necessary to remove ROS before they can damage sensitive parts of the cellular machinery¹⁶. SOD catalyses the dismutation of O_2 (singlet oxygen) to O_2 and H_2O_2 , representing the first line of cell defense against ROS production; CAT catalyses the production of H_2O from the degradation of H_2O_2 and ROOH; APX reduces H_2O_2 to H_2O using the ascorbate as an electron donor. The strain/metal specific metabolic responses were quite different as shown in Fig.2. Indeed, a slight increase in enzymatic activity was recorded in *G. maxima* APX and SOD as a response to $\text{Cl}_4\text{K}_2\text{Pd}$; CAT activity significantly decreased both under $\text{Cl}_4\text{K}_2\text{Pd}$ and AuCl_4K (Fig.2A). SOD, CAT and APX activities decreased in presence of $\text{Cl}_4\text{K}_2\text{Pd}$ while increased in presence of AuCl_4K in *G. phlegrea* (Fig.2B); a decrease in all enzymatic activities were recorded in *G. sulphuraria* (Fig.2C); a significant increase in all the enzymatic activities were recorded in *C. caldarium* only under $\text{Cl}_4\text{K}_2\text{Pd}$ (Fig.2D).

A significant increase of the enzymatic activity compared to the control suggests a high scavenging activity of the singlet oxygen in peroxide of hydrogen, which can be expressed as an obvious tolerance of these algae to the metal under examination. An increase in the activity of both antioxidant enzymes is necessary to reduce the concentrations of both singlet oxygen and hydrogen peroxide, minimizing the risks. In general, the modulation of antioxidant enzymes is an important adaptive response to counteract adverse conditions; in fact, the maintenance of a high antioxidant capacity in the cells can be correlated with an increased tolerance against different types of environmental stress^{15, 16, 19}.

CONCLUSIONS

Our results showed a higher tolerance to $\text{Cl}_4\text{K}_2\text{Pd}$ vs AuCl_4K in the cyanidiophyceae strains used in the present work. The growth and the metabolism of *G. phlegrea* was more affected by the presence of both metals. The contribute to the oxidative equilibrium of these polyextremophilic microalgae and the induction of antioxidant enzymes could result from the adaptation of the cell to the development of intracellular ROS; however there is not a clear correlation between any of the enzymatic activity and the better performing growth of the other 3 strains tested. The study from Ju and coworkers¹⁷ showed the ability of *G. sulphuraria* in efficiently recovering both $\text{Cl}_4\text{K}_2\text{Pd}$ and AuCl_4K , without assessing the influence of both metals on growth and physiology. We considered that tolerance is an essential parameter to take into account for biotechnological application, such as REE recovering. Our observations strongly suggest that other strains than *G. sulphuraria* can be used to recover REE, due to their high tolerance to precious and heavy metals, moreover further studies will be necessary to clarify the biological mechanisms underlying the tolerance capacity of *Cyanidiophyceae* and their strategies to respond to the metal toxicity in order to considerer by biotechnological future application.

REFERENCES

1. Jiang M, Ohnuki T, Kozai NM, Tanaka K, Suzuki Y, Sakamoto F, Kamiishi E, Utsumiya S. The Biological nano-mineralization of Ce phosphate by *Saccharomyces cerevisiae*. Chem Geol 2010; 277:61-9
2. Huroda K, Ueda M. Engineering of microorganisms towards recovery of rare metal ions. Appl Microbiol Biotechnol 2010; 87:53-60
3. Güzel Y, Rainer M, Mirza MR, Messner CB, Bonn GK. Highly selective recovery of phosphopeptides using trypsin-assisted digestion of precipitated lanthanide-phosphoprotein complexes. Analyst 2013; 138:2897-2905
4. Homosomi Y, Baba Y, Kubota F, Kamiya N, Goto M. Biosorption of rare earth elements by *Escherichia coli*. J Chem Eng Jpn 2013; 46:450-454
5. Wang M, Tan Q, Chiang F, et al. Recovery of rare and precious metals from urban mines—A review. Front Environ Sci Eng 2017; 11(5): 1
6. Matsunaga T, Takeyama H, Nakao T, Yamazawa A. Screening of marine microalgae for bioremediation of cadmium-polluted seawater. J Biotechnol 1999; 70:33-38
7. Mehta SK, Gaur JP. Use of algae for removing heavy metal ions from wastewater: progress and prospects. Crit Rev Biotechnol 2005; 25:113-152
8. Wang J, Chen C. Biosorbents for heavy metal removal and their future. Biotechnol Adv 2009; 27:195-226
9. Bourcier WL, Lin M, Nix G. Recovery of Minerals and Metals from Geothermal Fluids. In: SME Annual Meeting Cincinnati, OH, United States UCRL-CONF-2005; 215135.
10. Doemel W, & Brock ML. The physiological ecology of *Cyanidium caldarium*. J Gen Microbiol 1971; 67:17-32
11. De Luca P, Taddei R & Varano L. *Cyanidioschyzon merolae*: a new alga of thermal acidic environments. Webbia 1978; 33(1): 37-44
12. Albertano P, Ciniglia C, Pinto G, Pollio A. The taxonomic position of *Cyanidium*, *Cyanidioschyzon* and *Galdieria*: an update. Hydrobiologia 2000; 433:137-143.
13. Minoda A, Sawada H, Suzuki S, Miyashita S, Inagaki K, Yamamoto T, Tsuzuki M. Recovery of rare earth elements from the sulfothermophilic red alga *Galdieria sulphuraria* using aqueous acid. Appl Microbiol Biotechnol 2015; 99(3):1513-9.
14. Dao LHT, Beardall J. Effects of lead on growth, photosynthetic characteristics and production of reactive oxygen species of two freshwater green algae. Chemosphere 2016; 147. 420-429.

15. Foyer CH, Noctor G Redox Homeostasis and Antioxidant Signaling: A Metabolic Interface between Stress Perception and Physiological Responses. *The Plant Cell* 2005; 17: 1866–1875
16. Gill SS, Tuteja N Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiol Biochem* 2010 Dec;48(12): 909-30.
17. Ju X, Igarashi K, Miyashita S, Mitsuhashi H, Inagaki K, Fujii S, Sawada H, Kuwabara T, Minoda A. Effective and selective recovery of gold and palladium ions from metal wastewater using a sulfothermophilic red alga, *Galdieria sulphuraria*. *Bioresour Technol.* 2016; 211:759-64.
18. Allen MM & Stanier RY. Selective isolation of blue-green algae from water and soil. *J Gen Microbiol* 1968; 51: 203-209
19. Okamoto OK, Pinto EL, Latorre R, Bechara EJH, Colepicolo P. Antioxidant modulation in response to metal-induced oxidative stress in algal chloroplasts. *Arch Environ Contam Toxicol* 2001; 40:18-24.

Figure 1. Evaluation of metal tolerance monitoring MGR and 1/GT at day 4

Maximum growth rate (MGR, left) and inverse of Generation Time (1/GT, right) measured at day 4 in *G.maxima* (A), *G.phlegrea* (B), *G.sulphuraria* (C) and *C.caldarium* (D) grown at different concentrations of palladium and gold. MGR and 1/GT in presence of different concentration of palladium ($\text{Cl}_4\text{K}_2\text{Pd}$, orange line/bar) and gold (AuCl_4K , blue line/bar). Error bars represent standard deviation of three replicate cultures; (*) = p-value $\leq 0,001$ calculated by T-test.

Figure 2. Evaluation of enzymatic activities after metals exposition

Enzymatic activity measured in *G.maxima* (A), *G. phlegrea* (B), *G. sulphuraria* (C) and *C. caldarium* (D) treated by 1g/L of palladium ($\text{Cl}_4\text{K}_2\text{Pd}$, orange bar) and gold (AuCl_4K , blue bar) after 96h, Relative units represent: units/g of dry weight (SOD); nmol H_2O_2 /g of fresh weight (CAT); μmol ascorbate/g of fresh weight (APX) Mean (\pm SD) was calculated from three replicates. (*) = p-value $\leq 0,05$ calculated by T-test.

Fig.1




Fig.2

