



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/135892/>

Version: Published Version

Monograph:

Gutacker, Nils, Siciliani, Luigi, Moscelli, Giuseppe et al. (2015) Do patients choose hospitals that improve their health? Discussion Paper. CHE Research Paper . Centre for Health Economics, University of York , York, UK.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

ESHCRU

Economics of
Social and Health Care
Research Unit



Centre For Health Economics

UNIVERSITY *of York*



Do Patients Choose Hospitals That Improve Their Health?

Nils Gutacker, Luigi Siciliani,
Giuseppe Moscelli, Hugh Gravelle

CHE Research Paper 111

Do patients choose hospitals that improve their health?

¹Nils Gutacker

²Luigi Siciliani

¹Giuseppe Moscelli

¹Hugh Gravelle

¹Centre for Health Economics, University of York, UK

²Department of Economics and Related Studies, University of York, UK

May 2015

Background to series

CHE Discussion Papers (DPs) began publication in 1983 as a means of making current research material more widely available to health economists and other potential users. So as to speed up the dissemination process, papers were originally published by CHE and distributed by post to a worldwide readership.

The CHE Research Paper series takes over that function and provides access to current research output via web-based publication, although hard copy will continue to be available (but subject to charge).

Acknowledgements

We are grateful for comments and suggestions from Alistair McGuire, Andrew Street, Karen Bloor, Richard Cookson and those received during presentations at the Universities of Birmingham, Newcastle, York and Melbourne, the King's Fund, and the Leeds meeting of the Health Economic Study Group. This research was funded by the Department of Health in England under the Policy Research Unit in the Economics of Health and Social Care Systems (Ref 103/0001). The views expressed are those of the authors and not necessarily those of the funders. Hospital Episode Statistics are copyright 2015, re-used with the permission of The Health & Social Care Information Centre. All rights reserved.

Disclaimer

Papers published in the CHE Research Paper (RP) series are intended as a contribution to current research. Work and ideas reported in RPs may not always represent the final position and as such may sometimes need to be treated as work in progress. The material and views expressed in RPs are solely those of the authors and should not be interpreted as representing the collective views of CHE research staff or their research funders.

Further copies

Copies of this paper are freely available to download from the CHE website www.york.ac.uk/che/publications/. Access to downloaded material is provided on the understanding that it is intended for personal use. Copies of downloaded papers may be distributed to third-parties subject to the proviso that the CHE publication source is properly acknowledged and that such distribution is not subject to any payment.

Printed copies are available on request at a charge of £5.00 per copy. Please contact the CHE Publications Office, email che-pub@york.ac.uk, telephone 01904 321405 for further details.

Centre for Health Economics
Alcuin College
University of York
York, UK
www.york.ac.uk/che

© Nils Gutacker, Luigi Siciliani, Giuseppe Moscelli, Hugh Gravelle

Abstract

Many health care systems collect and disseminate information on provider quality in order to facilitate patient choice and induce competitive behaviour amongst providers. The Department of Health in England has recently mandated the collection of patient-reported health outcome measures (PROMs) for the purpose of performance assessment and consumer information. This is the first attempt to routinely measure the gain in health that patients experience as the result of care and thus offer a more comprehensive picture of hospital quality than existing 'failure measures' such as mortality or readmission rates. In this paper we test whether hospital demand responds to hospital quality measures based on health gains in addition to more conventional measures. We estimate hospital choice models for elective hip replacement surgery using rich administrative data for all publicly-funded patients in the English NHS in 2010-2012. Our focus is on two key aspects of hospital choice: 1) the extent to which patients are more likely to choose hospitals which are expected to achieve larger improvements in patients' health and 2) whether patients' response to quality differs with their morbidity, as measured by pre-operative health status, and other characteristics such as age or income deprivation. In order to address potential endogeneity bias we implement an empirical strategy based on lagged explanatory variables, hospital fixed effects and a control group design based on demand for emergency hip replacement. Our results suggest that hospitals can increase demand by 9% if they increase the average health gains that patients experience by one standard deviation. Hospital demand has a higher elasticity with respect to average health gains than emergency readmission or mortality rates. Elective patients are twice as willing as emergency hip replacement patients to travel further for an increase in quality.

Keywords: Patient choice, hospital demand, demand elasticity, quality of care, health outcomes

1. Introduction

Many European healthcare systems have recently extended patients' right to choose their provider of elective hospital care (Vrangbaek et al. 2012). Enhanced choice can accommodate patients' preferences for provider characteristics (e.g. proximity, quality or availability of amenities) and create market conditions that incentivise providers to compete (Besley and Ghatak 2003). Patients in the English National Health Service (NHS) have to be referred to inpatient services by their general practitioner, who acts as a gatekeeper, but are free to choose their preferred provider of care. Prices for hospital care are set nationally and patients do not bear the cost of treatment, so providers are expected to compete for elective patients on the basis of quality. Two prerequisites for such quality competition are that patients and their agents¹ have access to reliable, meaningful and understandable information about the quality of care offered by alternative providers, and that they act upon such information (Besley and Ghatak 2003; Marshall et al. 2004; Faber et al. 2009).

English patients can access comparative information on hospital quality through several channels, including the NHS Choices website, the Health & Social Care Information Centre (HSCIC) website and the Dr Foster Hospital Guide. These present information on risk-adjusted 28-day mortality and emergency readmission rates, calculated from routine hospital discharge data. Such indicators have been criticised as being incomplete and noisy measures of quality, revealing little about the changes in health that the vast majority of patients will experience as the result of treatment (Appleby and Devlin 2004; Lilford and Pronovost 2010). This is especially so for mortality rates for common elective operations such as hip (0.3%) and knee replacement (0.2%), which are generally very low (Berstock et al. 2014; Belmont et al. 2014).

New hospital quality measures that address these concerns are increasingly available. Since April 2009, all providers in the English NHS have been required to collect patient-reported outcome measures (PROMs) for all NHS-funded patients undergoing unilateral hip and knee replacement, varicose vein surgery or groin hernia repair (Department of Health 2008). PROMs are validated questionnaires used to elicit patients' health status and health-related quality of life (HRQoL)². Each eligible patient is invited to complete a PROM questionnaire before and three or six months after their surgery. The changes in scores can be interpreted as the improvement in patients' health and are used for hospital benchmarking (Nuttall et al. 2015; Gutacker, Bojke et al. 2013).

Hospital quality measures derived from PROMs improve over 'failure' measures such as mortality or emergency readmission rates in several ways. First, they capture the entire spectrum of health (Appleby and Devlin 2004; Gutacker, Bojke et al. 2013) and thus allow inferences about improvements in health as a consequence of treatment. Second, because post-operative health status is adjusted for pre-operative status, it can be argued that they adjust better for case-mix. Finally, PROMs reflect the patients' view on their health and health improvement. This, one may argue, makes them especially relevant for prospective patients who are about to choose their provider.

It has been the English Department of Health's expressed ambition to establish patients' self-reported outcomes as an important component of hospital quality assessment. It was also hoped that such

¹ These may include the patient's general practitioner (GP) as well as family, friends and others. Some patients may not be willing or able to make a choice and their referring GP may choose the most appropriate hospital for them, i.e. the GP acts as an agent to the patient. It is generally not possible to distinguish between decision makers using administrative data. For simplicity, we will henceforward denote the decision-maker as the patient.

² PROMs, such as the EuroQol-5D or the Oxford Hip Score, are widely used in the evaluation of health technologies. The National Institute for Health and Care Excellence (NICE) mandates the use of PROMs for outcome measurement.

information would be used “*by patients and GPs exercising choice*” (Department of Health 2008, p.6). Consequently, provider-specific average risk-adjusted changes in health status have been disseminated online on a regular basis since the beginning of the national PROM programme (Health & Social Care Information Centre 2013b). Some patients might access this information directly, whereas others might rely on their general practitioners, who act as their agent, to retrieve, interpret and communicate this information.

In this paper we test whether hospital demand responds to PROM-based measures of hospital quality in addition to more conventional measures such as mortality and readmission rates. We estimate a hospital choice model for elective hip replacement surgery in the English NHS to identify how hospital choice responds to hospital and patient characteristics. Our focus is on two key aspects of hospital choice: 1) whether hospitals with better PROM-derived quality (as measured by the changes in patients’ Oxford Hip Score (OHS)) face higher demand and 2) whether patients’ response to quality differs according to their morbidity, as measured by the pre-operative health status, and other characteristics such as age or income deprivation. To address potential endogeneity we use lagged quality and waiting times. We also undertake robustness checks using hospital fixed effects and by comparing the effects of quality on choices by elective hip replacement patients with those by emergency hip replacement patients who we expect to be less sensitive to quality.

This is the first study which explores whether hospital demand responds to quality as measured by average patient health gains at provider level, which are derived from patient self-reported outcome measures. The existing literature has predominantly focused on failure measures such as mortality rates, either measured at aggregate hospital level or for specific conditions (e.g Sivey 2008; Beckert et al. 2012; Moscone et al. 2012; Gaynor, Propper et al. 2012), readmission rates (Varkevisser et al. 2012; Moscone et al. 2012), as well as hospital reputation and other composite scores (Pope 2009; Varkevisser et al. 2010; Varkevisser et al. 2012; Ruwaard and Douven 2014); see Brekke et al. (2014) for an overview. These studies have typically reported a positive relation between quality and hospital demand. Second, we make novel use of pre-operative individual level PROMs data to explore such questions as whether sicker patients travel farther and choose hospitals with higher quality of care as often assumed in the literature (Gowrisankaran and Town 1999; Geweke et al. 2003). Previous studies have either relied on instrumental variable approaches to approximate the role of (unobserved) pre-operative health status on demand (Gowrisankaran and Town 1999; Geweke et al. 2003) or have used measures of comorbidity burden and past utilisation as proxies for health status. Our data allow us to explore this issue more directly. Third, our paper contributes to the small literature on hospital choice in publicly funded health systems where demand is rationed by waiting time (Sivey 2012; Beckert et al. 2012; Moscone et al. 2012; Gaynor, Propper et al. 2012). Our analysis differs from Beckert et al. (2012), who also study choice of provider for hip replacement surgery in England, in that we use provider quality measures which are procedure-specific and more directly related to the quality of care provided³, explore the role of pre-operative health status, and model the entire relevant market, including private providers of NHS-funded care.

We find that hospitals with higher PROM scores faces larger demand, and that this effect is unlikely to be driven entirely by omitted variables. The elasticity of provider demand to average health gains is 1.3 or

³ Beckert et al. (2012) model hospital quality using hospital-wide mortality rates and MRSA infection rates. Aggregate hospital level quality indicators, such as the summary hospital mortality indicator (SHMI) used in the English NHS, do not correlate well with hip procedure specific outcome measures (Gravelle et al. 2014). In 2010/11, the Pearson correlation coefficients between SHMI and the quality measures used in this study were -0.09 (OHS), -0.05 (emergency readmission rate) and 0.10 (mortality rate), respectively.

about 34 patients for a standard deviation increase in average OHS. A one standard deviation increase in hospital average health gains corresponds to a willingness to travel of just over one additional kilometre. Demand has a higher elasticity with respect to quality measures based on average health gains than procedure-specific mortality or readmission rates.

The remainder of the paper is structured as follows: in the next section we describe the data used in this study in more detail. Section 3 describes our econometric model and sets out our strategy to mitigate potential endogeneity bias. In Section 4 we present the estimated marginal utilities of hospital characteristics and show how these vary with observed patient characteristics. Section 5 presents the estimated effects of changes in providers' quality on their own demand and that of their competitors. Finally, the last section offers a discussion of the results.

2. Data

We use patient-level data from Hospital Episode Statistics (HES) for all elective admissions for patients aged 18 or over who underwent NHS-funded primary (i.e. non-revision) hip replacement surgery⁴ between April 2010 and March 2013 in NHS or private providers. HES contains rich information on patients' demographic and medical characteristics, small area of residence and on the hospital stay. Privately funded patients treated in the private sector are not included in HES and are excluded from our analysis⁵.

We derive four patient variables from HES: patients' age, gender, the number of emergency admissions during the 365 days prior to their hip replacement admission, and the number of Elixhauser comorbid conditions recorded in admissions in the previous year (Elixhauser et al. 1998; Gutacker, Bloor et al. 2015). These are available for all patients. We use the 2004 Index of Multiple Deprivation (Noble et al. 2006) to attribute to each patient the proportion of residents claiming means-tested benefits in their Lower Super Output Area (LSOA)⁶, which we interpret as a measure of income deprivation. We measure a patient's distance from a hospital as the straight-line distance from the centroid of their LSOA⁷.

The PROM survey invites all NHS-funded hip replacement patients to report their health status and HRQoL before and six months after surgery using a paper-based questionnaire. The pre-operative questionnaire is administered by the hospital either as part of the admission process or during the last outpatient appointment preceding admission. The post-operative questionnaire is administered by a central agency and posted to the patient. Participation in the PROM survey is compulsory for providers but optional for patients. Approximately 60% of patients provide complete pre- and postoperative PROM questionnaires that can be linked to their HES record (Hutchings et al. 2014; Gutacker, Street et al. forthcoming).

Each PROM questionnaire contains three instruments: the Oxford Hip Score (OHS), the EuroQoL-5D (EQ-5D) descriptive system, and the EuroQol Visual Analogue Scale (EQ-VAS). The OHS is a condition-specific instrument that consists of 12 questionnaire items regarding hip-related functioning and pain (Dawson et al. 1996). Each item is scored on a five-point scale, with four indicating no problems and zero indicating severe problems. The overall score is calculated as the sum of all items and ranges from zero (worst) to 48 (best). Both EuroQol instruments are generic PROMs, i.e. they can be applied to different health conditions, and are described in detail elsewhere (Brooks 1996). Previous analysis showed substantial correlation between the EQ-5D and OHS (Neuburger et al. 2013). Since the OHS is a condition-specific measure and hence plausibly more likely to affect hospital choice for hip replacements we focus on the OHS throughout this paper.

We use PROMs data in two ways. First, we obtained risk-adjusted hospital-specific PROM change scores for the OHS from the HSCIC website (Health & Social Care Information Centre 2013b). Data are

⁴ See Department of Health (2008) for procedure codes. We exclude patients that underwent revision surgery to ensure a more homogeneous sample and because these are believed to be likely to return to the place of initial surgery, independent of observed hospital attributes

⁵ Approximately 11% of the English population have private (supplementary) insurance and approximately 16% of hip replacement surgeries are funded privately, either out-of-pocket or through private insurance (Hunt et al. 2013; Commission on the Future of Health and Social Care in England 2014)

⁶ HES records patients' locations in terms of the LSOA (2001 census boundaries) in which they reside. Each LSOA contains approximately 1,500 inhabitants and is designed to be homogeneous with respect to tenure and accommodation type.

⁷ We determine a hospital's location on the basis of its headquarter's postcode (for NHS trusts) or the postcode of the individual hospital's site (for ITCs). We do not model NHS hospital sites individually as quality information for these providers is only recorded at trust level. This is likely to induce noise to our distance measure.

reported by financial year, which run from April to March of the next year. The HSCIC excludes from these reports providers with less than 30 valid pre- and post-operative PROM returns due to concerns about statistical validity and patient anonymity. The case-mix adjustment methodology is reported elsewhere (Department of Health 2012).⁸ There is some evidence to suggest that the hospital-specific mean scores are robust to missing data (Gomes et al. forthcoming). Second, in some of our models, we use the information in the individual patients' pre-operative PROMs questionnaires to measure their pre-operative morbidity and investigate whether choice of provider is affected by pre-operative morbidity. Because patients can decline to participate or providers may fail to administer a questionnaire there is scope for missing data and selection bias, and we explore this in the empirical analysis for the subset of models which make use of pre-operative morbidity.

We calculate risk-adjusted hospital-specific 28-day emergency readmission and 28-day mortality rates after hip replacement as additional quality measures. These data are presented on patient information websites (such as NHS Choices). To compute them, we link our HES data to Office of National Statistics death records and apply the HSCIC case-mix adjustment as set out in the readmission outcome indicator specification (Health & Social Care Information Centre 2013a).⁹

We group providers into seven categories used by the National Patient Safety Agency: NHS small / medium / large non-teaching trust, NHS teaching trust, NHS specialised orthopaedic provider, NHS multi-service provider, and NHS Primary Care Trusts (PCTs).¹⁰ We also distinguish NHS hospitals from Independent Sector Treatment Centres (ISTCs) which are private providers treating NHS patients.

Finally, we derive from HES the median time (in months) that patients in each hospital had to wait between the specialist's decision to add the patient to the waiting list and the admission (the inpatient wait). Patients in the English NHS do not pay for their care directly and waiting times thus serve as a rationing mechanism (Iversen and Siciliani 2011). We use the median rather than the mean because it is less affected by a small number of patients with very long wait and thus more representative of the expected waiting time for a prospective patient. We also conduct sensitivity analysis using the proportion of patients in this hospital that waited longer than 120 days.

⁸ The adjustment takes into account a range of patient characteristics including age, sex, pre-operative PROM score, socio-economic status, comorbidity burden, whether the patient lives alone as well as other indicators of disability.

⁹ Both readmission and mortality rates are adjusted for age (in 5-yr bands), sex, socio-economic status, comorbidity burden as captured by the Charlson index and the number of emergency admissions in the last year.

¹⁰ PCTs are responsible for purchasing care for their resident population and, with the exception of the Isle of Wight PCT, do not provide care themselves.

3. Methods

3.1. Model specification

We use a random utility choice model (McFadden 1974). Utility of patient $i = 1, \dots, N$ at provider $j = 1, \dots, J$ at time $t = 1, \dots, T$ is $U_{ijt} = V_{ijt} + \xi_{jt} + \epsilon_{ijt}$, where V_{ijt} depends on observable hospital characteristics and travel distance, ξ_{jt} are unobserved hospital characteristics, and ϵ_{ijt} is unobserved random utility. Patients choose from a set of hospitals $M_{it} \in J$. Assuming ϵ_{ijt} is iid extreme value yields the *multinomial logit* (MNL) model in which the probability that patient i chooses hospital j is

$$P_{ijt} = \exp \frac{V_{ijt} + \xi_{jt}}{\sum_{j \in M_{it}} V_{ij't} + \xi_{j't}} \quad (1)$$

We assume that all patients who require treatment are treated, i.e. there is no outside option.

In our baseline specification, utility is a linear additive function of the distance from the patient's residence to the hospital D_{ij} , distance squared D_{ij}^2 , hospital quality metrics Q_{jt-1} , waiting time W_{jt-1} , and a vector of time-invariant hospital characteristics Z_j , so that

$$U_{ijt} = D'_{ij}\beta_{d,i} + D_{ij}^2\beta_{d^2,i} + Q'_{jt-1}\beta_{q,i} + W'_{jt-1}\beta_{w,i} + Z'_{jt}\beta_{z,i} + \xi_{jt} + \epsilon_{ijt} \quad (2)$$

where ξ_{jt} and ϵ_{ijt} are unobserved. We assume that anticipated utility at a provider is based on its previous period's quality and waiting time because relevant information are available only with a lag (see section 3.2). Varkevisser et al. (2012) make a similar assumption. We also estimate models with contemporaneous waiting time and quality scores in sensitivity analyses.

We allow preferences to vary across patients according to their observed characteristics. Thus the marginal utility of quality for patient i is

$$\beta_{q,i} = \beta_q + X'_i\delta_q \quad (3)$$

and similar for distance, waiting time, and other hospital characteristics. All continuous covariates in X_i are mean centred and base categories for categorical characteristics are set to their mode. Thus, the vectors of coefficients $\beta_d, \beta_{d^2}, \beta_q, \beta_w, \beta_z$ reflect the preferences of an average/modal patient, hereafter referred to as the '*reference patient*'.

We also estimate models which allow for unobserved patient heterogeneity in tastes over quality, with

$$\beta_{q,i} = \beta_q + X'_i\delta_q + \sigma_q\alpha_i \quad (4)$$

where σ_q is the standard deviation of a normal variable with mean zero and α_i is an unobserved patient effect. The latter may capture, for example, differences in the ability to access and interpret quality information. This *random coefficient multinomial logit* (RCMNL) or mixed logit model (Hensher and Greene 2003; Train 2003), unlike the MNL model, allows for unrestricted substitution patterns, thereby relaxing the assumption of independence of irrelevant alternatives (IIA).¹¹ If $\sigma_q = 0$ then the RCMNL model reduces to the MNL model in (2).

¹¹ The IIA states that the probability of choosing one hospital over another depends solely on the characteristics of these two hospitals and not on the characteristics of any other hospital. The standard MNL model imposes the IIA assumption, whereas the RCMNL does not.

While the MNL model has a closed form solution that can be estimated via maximum likelihood, the RCMNL needs to be approximated through simulation. To reduce the computational burden¹² of the RCMNL model we assume uncorrelated normally distributed random coefficients for the quality metrics in Q_{jt-1} and no random coefficients for other covariates. The RCMNL model is estimated with maximum simulated likelihood using 50 Halton draws.

All models are estimated in Stata 13 with *clogit* and the user-written command *mixlogit* (Hole 2007b). Standard errors are clustered at the GP practice level to allow for agent-induced correlation across patients.

3.2. Endogeneity

To interpret β_q as an unbiased estimate of the marginal utility of hospital quality (up to a linear transformation) requires that the unobserved hospital effect ξ_{jt} is uncorrelated with any of the independent variables, i.e. all observed variables are exogenous. This assumption may not hold for four reasons (Varkevisser et al. 2012; Gaynor, Propper et al. 2012; Brekke et al. 2014).

First, hospitals may learn by doing so that higher volume providers have higher quality (Luft et al. 1987; Gaynor, Seider et al. 2005). Thus changes in demand will also affect quality and induce simultaneity bias. Based on the institutional context of this study we argue that this concern can be dismissed. While volume-outcome effects have been reported for elective joint replacement surgery, these scale effects tend to occur only in very low volume hospitals that treat less than 100 patients per year (Judge et al. 2006; Mäkelä et al. 2011). The increasing incidence of hip replacement surgery in England and trends to aggregate services in high-volume hospitals mean that all NHS providers in our sample are comfortably above this threshold and has led commentators to suggest that volume effects are of little relevance in the English NHS (Judge et al. 2006). For private providers we cannot ascertain their true level of activity as treatment of non-NHS patients is not recorded in HES, but we expect those to perform a sufficient number of procedures to operate profitably. The average hospital in our sample treats over 300 patients per year.

Second, because of short run capacity constraints, changes in demand will also affect waiting time in the same period (Gaynor, Propper et al. 2012).¹³ While our primary interest is not in the effect of waiting time on demand, we are concerned that any bias introduced through endogenous variables will filter through to our estimate of β_q (Wooldridge 2002). However, if, as we assume, demand depends on past, rather than current, quality and waiting time, then demand changes in period t cannot affect waiting time at $t - 1$.

Third, sicker patients may choose higher quality hospitals or hospitals may turn away or discourage patients with characteristics that make them less likely to achieve a large improvement in health status. If such systematic selection occurs and is not controlled for in the calculation of hospital quality scores then those scores would in part be determined by patients' choices or provider selection. However, provider quality scores are adjusted for a rich set of demographic, socio-economic, and morbidity patient characteristics, including, in the case of PROMs, the patients' self-reported pre-operative health status.

¹² Even after imposing those constraints the RCMNL model with our baseline specification still took over 5 days to compute on a high-performance computing system.

¹³ It may also be that supply and demand are determined simultaneously, i.e. hospitals react to demand shocks by adjusting their supply, e.g. by performing more surgeries on weekends. We do not consider this in our model explicitly, although the use of lagged waiting time circumvents this problem as well.

Hence, we do not believe that unobserved patient selection is likely to bias the quality scores significantly.

Finally, there may be unobserved hospital characteristics that affect demand and are correlated with observed covariates (Jung et al. 2011). For example, hospitals in areas with better amenities may attract better staff thereby ensuring higher observed clinical quality but also unobserved interpersonal aspects of quality. Our assumption that patients use information on previous period quality and waiting times when choosing hospitals does not remove omitted variable bias operating through unobserved non-transitory hospital characteristics. However, the low correlations between the PROM quality measure and the conventional readmission and mortality measures suggest that omitted variables may not lead to serious bias. We undertake two types of sensitivity analyses to explore the size of the potential omitted variable bias. Our first approach is to estimate the choice model in (2) with alternative-specific time-invariant fixed effects (FEs) (Hodgkin 1996; Monstad et al. 2006; Sivey 2012). These hospital FEs capture the utility of non-transitory unobserved hospital characteristics. The coefficients on observed hospital characteristics are now identified solely through variation within providers over time, thereby removing any endogeneity bias operating through unobserved time-invariant characteristics. However, this approach is quite demanding of the data, and because we only observe providers over three years we expect this approach to result in imprecise estimates of the marginal utility of hospital quality. Also, because our market structure changes over time, due to the opening of new independent sector treatment centres, the FEs do not correspond to observed market shares in each time period. This may bias estimates if incumbent providers differ systematically from new entries. We therefore also estimate a model based on NHS trusts only, whose numbers are relatively stable over time.

Our second approach is to follow Pope (2009) (see also Gaynor, Propper et al. (2012)) and gauge the possible impact of unobserved hospital heterogeneity by using a control group of emergency hip replacement patients whose choice of provider is less responsive to quality and waiting time. The majority of emergency hip replacement patients suffer from a fractured neck of the femur as a result of a fall and official recommendations are that they should be treated within 48 hours (NICE 2011). Further delays are linked to worse outcomes (Moja et al. 2012). We therefore expect provider choice by emergency hip replacement patients to be less affected by publicly reported information on quality and more by distance to providers and time-invariant unobserved factors, such as long-standing reputation or dimensions of accessibility not captured by our distance measure (e.g. parking charges or connection to the public transport system).

If we assume that emergency patients' demand is entirely inelastic to observed quality and they do not wait¹⁴, but value the same unobserved hospital characteristics as elective patients, then their true utility is given by

$$U_{ijt}^{Emer} = D'_{ij}\beta_{d,i}^{Emer} + D'^2_{ij}\beta_{d^2,i}^{Emer} + \xi_{jt} + \epsilon_{ijt} \quad (5)$$

If we estimate the model specified in (2) for emergency patients and find $\hat{\beta}_q^{Emer} \neq 0$, we conclude that $cov(Q_{jt-1}, \xi_{jt}) \neq 0$. Moreover, if we assume that elective and emergency patients have the same preferences for unobserved hospital characteristics, then the effect of quality on elective demand, purged of omitted variable bias, is $\beta_q^\Delta = \beta_q^{Elec} - \beta_q^{Emer}$. Since coefficients in separate MNL models may be scaled differently, we estimate a pooled model for elective and emergency patients by interacting all covariates with an indicator variable for emergency. This forces the scaling to be the same. The coefficients on the

¹⁴ Elective waiting time and associated supply constraints do not apply to emergency patients, i.e. there is always sufficient capacity to treat an emergency patient. Given the urgent nature of the condition, patients will usually be treated within hours of arrival, not weeks or months. Explorations of our data revealed that elective waiting time is only weakly correlated with the volume of emergency patients, suggesting that supply for these distinct groups is separate.

interaction terms are estimates of β_k^Δ for $k \in [d, d^2, q, w, z]$.

If emergency patients are also sensitive to elective quality¹⁵, or emergency quality that correlates with it, or if unobserved hospital characteristics have different effects on choices by emergency and elective patients and are correlated with observed quality, then β_q^Δ can no longer be interpreted as the unbiased effect of quality on elective demand. If unobserved hospital factors are not correlated with quality, then β_k^Δ reflects the differences in preferences in two distinct groups of patients: those that require urgent care and have less time to compare hospitals, and those that have sufficient time to reach an informed decision. In this case, we expect that $\beta_q^\Delta > 0$: elective patients will be more sensitive to quality than emergency patients.

3.3. Elasticities, changes in demand and willingness to travel

The estimated coefficients on quality are estimates of the marginal utility from quality. Since the utility function is unique only up to a linear transformation, the coefficients only convey information about the sign of marginal utility of hospital characteristics and hence about the sign of the effect of quality on demand. The ratio of estimated marginal utilities (the negative of the marginal rate of substitution) is unaffected by linear transformations and so provides quantitative and comparable information about patient preferences. We estimate the reference patient's willingness to travel (WTT) for a one standard deviation (SD) increase in quality as

$$WTT = \frac{\partial D_{ij}}{\partial Q_j} \Big|_{U_{ij}} SD(Q) = -\frac{\partial U_{ij}}{\partial Q_j} / \frac{\partial U_{ij}}{\partial D_{ij}} SD(Q) = \frac{-\beta_q}{\beta_d + 2\beta_{d^2}D} SD(Q) \quad (6)$$

where D is the median distance to hospitals in patients' choice sets. We estimate standard errors by the delta method (Hole 2007a). WTT is the extra distance in kilometres that the reference patient located the median distance away from a provider would be willing to travel to that provider if its quality was increased by $SD(Q)$, where $SD(Q)$ is averaged across hospitals and years.

We are also interested in whether providers could attract more patients by improving their quality. Expected demand at provider j is $Y_{jt} = \sum_{i \in S_{jt}} P_{ijt}$, where S_{jt} is the set of patients whose choice set includes provider j , i.e. for whom $j \in M_{it}$. Following Santos et al. (forthcoming) we calculate the average partial effect of a one SD increase in quality on provider j 's demand, i.e. demand responsiveness to quality, as

$$\frac{\partial Y_{jt}}{\partial Q_{jt-1}} SD(Q) = SD(Q) \sum_{i \in S_{jt}} \frac{\partial P_{ijt}}{\partial Q_{jt-1}} = SD(Q) \sum_{i \in S_{jt}} \beta_q P_{ijt} (1 - P_{ijt}) \quad (7)$$

We report the mean of (7) over all providers and years.

We calculate the elasticity of demand of provider j with respect to own quality as

$$E_{jt}^{Q_{jt-1}} = \sum_{i \in S_{jt}} \frac{\partial Y_{ijt}}{\partial Q_{jt-1}} \frac{Q_{jt-1}}{Y_{ijt}} = \sum_{i \in S_{jt}} \beta_q P_{ijt} (1 - P_{ijt}) \frac{Q_{jt-1}}{\sum_{i \in S_{jt}} P_{ijt}} \quad (8)$$

We report the mean of (8), weighted by providers' predicted demand $\sum_{i \in S_{jt}} P_{ijt}$.

¹⁵ As with elective patients, we do not observe who chooses the hospital for emergency hip replacement. This may be the patient, a family member, GP, or the ambulance crew.

Finally, we compute the cross-elasticity of demand for provider j with respect to the quality of provider j' as

$$E_{jt}^{Q_{j'}} = \sum_{i \in S_{jt} \cap S_{j't}} \frac{\partial P_{ijt}}{\partial Q_{j't-1}} \frac{Q_{j't-1}}{\sum_{i \in S_{jt}} P_{ijt}} = - \sum_{i \in S_{jt} \cap S_{j't}} \beta_q P_{ijt} P_{ij't} \frac{Q_{j't-1}}{\sum_{i \in S_{jt}} P_{ijt}} \quad (9)$$

with $j \neq j'$. Note that for some combinations of j and j' the cross-elasticity is zero because no patients have both providers in their choice sets.

4. Results

4.1. Descriptive statistics

Our main sample is 173,773 elective hip replacement patients treated in 230 providers during the period April 2010 to March 2013.¹⁶ Their average age is 68 years and 40% are male (Table 1). The average pre-operative OHS is 17.5 and 9% of patients have been admitted to hospital as an emergency at least once during the preceding 365 days (average number of admissions = 0.13). Self-reported pre-operative OHS is only weakly correlated with past emergency utilisation ($\rho = -0.10$) and the number of comorbidities ($\rho = -0.14$). This suggests that past emergency utilisation and comorbidity burden are poor proxies for current health status¹⁷ as experienced by the patient.

Table 1: Descriptive statistics - elective sample

Variable	Obs	Mean	SD	ICC
Patient characteristics				
Distance travelled (in km)	173,773	14.7	17.7	
Distance travelled past closest provider (in km)	173,773	5.4	14.8	
Number of providers within 10km radius	173,773	1.6	1.7	
Number of providers within 30km radius	173,773	8.5	7.3	
Age	173,773	68.0	11.5	
Male	173,773	0.40	0.49	
Past utilisation	173,773	0.13	0.49	
Number of Elixhauser conditions	173,773	0.43	0.94	
Income deprivation	173,773	0.12	0.09	
Pre-operative Oxford Hip Score ^a	71,614	17.5	8.2	
Provider characteristics				
Observed volume	571	304.3	209.1	94.7%
Waiting time (in months)	571	2.5	1.1	77.4%
Change in Oxford Hip Score	571	19.8	1.4	57.0%
28-day emergency readmission rate (in %)	571	5.65	2.41	36.8%
28-day mortality rate (in %)	571	0.17	0.36	3.4%

Obs = Observations; SD = Standard deviation; ICC = Intraclass correlation coefficient.

Notes: Patient characteristics for patients choosing provider between April 2010 and March 2013. Provider waiting time, change in Oxford Hip Score, readmission rate, mortality rate are for financial years 2009/10 to 2011/12. Provider characteristics are unweighted.

^a Responders to PROM survey that were treated between April 2010 and March 2012.

On average, within 30km patients have a choice of 8 providers, with over 90% of patients having access to at least two different providers. Even within 10km there are on average 1.6 hospitals and over 20% of patients can choose between two or more providers. To reduce computational burden we restrict patient choice sets to the 50 nearest providers. The 741 patients (or 0.04% of the sample) who chose a provider outside this set were dropped from the analysis.

Patients live on average 14.7 kilometres from their chosen hospital. Figure 1 shows that just over half (53.7%) of patients bypassed the local hospital and nearly a fifth (18.3%) bypassed the nearest three

¹⁶ The number of providers varied slightly over this period because of mergers, changes in coding and market entry, especially with respect to private facilities. There were 157 providers in 2010/11, 202 in 2011/12, and 212 in 2012/13, of which 18 (11.5%) in 2010/11, 62 (30.7%) in 2011/12, and 78 (36.8%) in 2012/13 are privately operated.

¹⁷ We also calculated the correlations between these measures and the EQ-5D utility score, which one may argue is a more holistic measure of health-related quality of life. The correlations are similar: $\rho = -0.10$ for past utilisation, and $\rho = -0.14$ for comorbidity burden.

hospitals. On average, patients travel 5.4 km (SD=14.8) beyond their nearest hospital to be treated.¹⁸




Figure 1: Percentage of elective patients who went to their Nth nearest hospital

The hospital waiting time and quality scores are lagged by one year and are for financial years 2009/10 to 2011/12. The risk-adjusted OHS health again has a mean of 19.8 with a SD of 1.4. There are much larger coefficients of variation for hospital emergency re-admission and mortality rates. The average waiting time at provider level is 2.5 months, which is substantially lower than in previous years (see Appendix Figure A1 and Siciliani et al. 2014). The provider OHS change scores are only weakly correlated with waiting time ($\rho=-0.30$), readmission rates ($\rho=-0.28$) and mortality rates ($\rho=-0.05$). This suggests that choice models that are restricted to mortality and readmission rates may not even indirectly pick up the effect of PROM measures on demand.

The intra-class correlation coefficient (ICC) shows that just over half of the observed variation in OHS is between providers (ICC=57%) rather than over time. Between-provider variation is markedly greater for waiting times (ICC=77%). Most of the variation in readmission rates and mortality is within providers.

4.2. Regression results

4.2.1. Main effects

The results from for the RCMNL model (see Appendix Table A1) suggest no significant variation in the random coefficients on the quality metrics. Hausman tests also did not reject the IIA assumption. We therefore concentrate on the MNL models reported in Tables 2 to 5.

Table 2 is our preferred specification with distance, lagged waiting time, the three lagged quality metrics and indicators for the type of provider as well as interactions with patient age, gender, past utilisation, comorbidity, and local area income deprivation (we explore interactions with pre-operative OHS in

¹⁸ These numbers are somewhat higher than those reported by Beckert et al. (2012), presumably because our data also cover private providers treating NHS-funded patients.

section 4.2.2). This specification does not include hospital FEs. The main effects are the estimated marginal utilities for the reference patient with mean or modal characteristics. The reference patient prefers shorter distances with the marginal disutility from distance declining with distance. She prefers specialised providers to non-specialised providers. She is also more likely to choose a public provider over a private provider after accounting for distance, waiting time and quality¹⁹.

Reference patient demand is increasing with the OHS change score and falling with emergency admission rates²⁰. The estimated WTT for a one SD increase in OHS is 1.3 km or 8.7% of the average distance travelled to the chosen provider. The WTT for a SD decrease in emergency readmission rates is 1.0km. There is no statistically significant effect of procedure-specific mortality rates on demand. Nor does the waiting time affect choice of provider, which may be a result of the historically short waiting time during our study period.

Results are robust to the use of contemporaneous rather than lagged waiting time and quality (Appendix Table A2, model 1). Contemporaneous waiting time has a positive but statistically insignificant coefficient. When we use the proportion of patients waiting longer than 120 days as a waiting time measure the coefficient is negative and statistically significant (Appendix Table A2, model 2). The coefficients on the quality measures are almost unaffected by the use of contemporaneous waiting time and quality.

The HSCIC also produces hospital quality scores based on the case-mix adjusted change in the EQ-5D utility score. This is highly correlated with the OHS change score (Neuburger et al. 2013) and when we estimate the baseline specification with EQ-5D substituted for OHS we find similar WTT (Appendix Table A2, model 3). Results are also robust to exclusion of independent sector treatment centres from patient choice sets (Appendix Table A2, model 4).

4.2.2. Patient heterogeneity

The coefficients on the interaction terms in the lower parts of Table 2 suggest that preferences vary across types of patient. We find, like other studies (Propper et al. 2007; Beckert et al. 2012), that older patients dislike distance more. They care less about waiting time and get greater marginal utility from improvements in the OHS change score, reductions in emergency readmissions and reductions in mortality rates. There is little difference between the preferences of male and female patients except that male patients have a greater dislike for providers with higher mortality. Preferences vary little by morbidity as measured by past emergency admissions. In contrast, patients with more comorbidities have a greater dislike of distance and waiting time, but care less about readmission rates. Finally, patients from neighbourhoods with greater income deprivation care more about distance and less about quality.

The existence of detailed patient reported pre-operative health status measures in our dataset allows us to explore in more detail whether patients in worse health status are more sensitive to quality and more willing to travel, as commonly assumed in the literature on hospital quality (Gowrisankaran and Town 1999; Geweke et al. 2003). The correlations between patients' pre-operative OHS and their routinely available morbidity measures are low, suggesting that they measure different aspects of the patient's

¹⁹ During our study period ISTC were funded through block contracts and paid to provide care to a pre-specified number of NHS patients. However, most ISTCs did not fulfil their quotas although they generally had low waiting times (Naylor and Gregory 2009). Our results are consistent with this observation and suggest a positive preference for public providers by NHS-funded hip replacement patients.

²⁰ We also tested for a non-linear effect of PROM quality on demand by including a squared term for the provider OHS change scores (Ruwaard and Douven 2014). The square term was statistically significant and negative but there was little difference between the linear and non-linear models over the observed range of values.

Table 2: Estimated marginal utilities

Variable	Est	SE
Main effects		
Distance (in km)	-0.184	0.002***
Distance ²	0.000	0.000***
NHS trust - medium	-0.530	0.030***
NHS trust - multi-service	-0.603	0.099***
NHS trust - small	-0.791	0.038***
NHS trust - specialist	1.023	0.072***
NHS trust - teaching	-0.445	0.033***
Independent sector treatment centre	-1.467	0.045***
Primary care trust	-1.159	0.206***
Waiting time (in months)	0.013	0.015
Change in Oxford Hip Score	0.118	0.008***
28-day emergency readmission rate (in %)	-0.052	0.004***
28-day mortality rate (in %)	-0.031	0.026
Interaction with distance		
x Patient age	-0.002	0.000***
x Male	0.002	0.001
x Past utilisation	-0.003	0.002
x Comorbidity count	-0.004	0.001***
x Income deprivation	-0.186	0.017***
Interaction with waiting time		
x Patient age	0.003	0.000***
x Male	-0.009	0.009
x Past utilisation	-0.006	0.012
x Comorbidity count	-0.018	0.007**
x Income deprivation	0.046	0.084
Interaction with change in Oxford Hip Score		
x Patient age	0.001	0.000*
x Male	-0.007	0.005
x Past utilisation	-0.010	0.007
x Comorbidity count	-0.009	0.003**
x Income deprivation	-0.455	0.047***
Interaction with 28-day emergency readmission rate		
x Patient age	-0.0003	0.000*
x Male	0.000	0.003
x Past utilisation	0.011	0.003**
x Comorbidity count	0.001	0.002
x Income deprivation	0.125	0.026***
Interaction with 28-day mortality rate		
x Patient age	-0.001	0.001
x Male	-0.053	0.022*
x Past utilisation	0.046	0.026
x Comorbidity count	-0.009	0.015
x Income deprivation	-0.025	0.168
WTT(OHS change)	1.287	0.085***
WTT(Readmission rate)	-0.981	0.079***
WTT(Mortality rate)	-0.086	0.072
Number of patients	173,032	
Number of providers	230	
BIC	460,994	
Pseudo R ²	0.637	

*** p<0.001; ** p<0.01; * p<0.05

Notes: Conditional logit model of choice of hospital for elective hip replacement patients treated between April 2010 and March 2013. OHS change, waiting time, readmission rate and mortality rate are lagged by one year. Coefficients are marginal utilities. WTT is the ratio of the coefficient on the quality variable to the marginal utility of distance evaluated at the median distance (in km). Interaction terms with distance² and provider type not reported (available on request). Standard errors are clustered at GP practice level.

condition at the time of admission.

The first model in Table 3 is the same as our preferred specification but with additional patient pre-operative OHS interactions. Interaction terms with other patient characteristics are suppressed for brevity. Due to data limitations, we focus on patients treated during April 2010 and March 2012. We find that healthier patients are more willing to travel. Although the marginal utility from higher quality is similar for healthier patients, the reduced distance cost for these patients implies they are more willing to travel for higher quality. Healthier patients are also more likely to choose a private provider, which is consistent with observed differences in intake across provider types (Browne et al. 2008).

The fact that pre-operative OHS data are available for only about 60% of patients raises concerns about response bias if unobserved factors affect propensity to respond and utility from providers²¹. To investigate if responders to the pre-operative PROM questionnaire have different preferences to non-responders we re-estimate the preferred specification of Table 2 for our full sample (responders and non-responders) but interact a dummy variable for responder status with all the main and interacted explanatory variables; pre-operative health status is not modelled.

The pre-operative PROM questionnaire is administered *after* the patient has chosen the provider. Hence, it is unclear whether the response indicator variable reflects patient preferences or whether the choice determines the response indicator. For example, private providers have higher response rates than NHS hospitals (Gomes et al. forthcoming; Gutacker, Street et al. forthcoming) and also tend to have higher observed quality and shorter waiting times. We address this concern by including the observed provider pre-operative response rate as a provider characteristic when modelling the choices of responders and non-responders. This variable is informative about the individual's propensity to fill in a pre-operative PROM questionnaire given the chosen provider²². We find that responders and non-responders have generally very similar revealed preferences, with the exception of preferences for waiting times (non-responders prefer shorter waiting times) and PROM quality (responders derive more utility from health gains and are thus more willing to travel for it). There is no difference with respect to the disutility from travel distance, readmission rates or mortality.

4.3. Omitted variable bias

We also explore the possible impact of omitted hospital characteristics on our estimates of marginal utility for quality and other hospital characteristics. We compare preferences of elective and emergency patients estimated from pooled choice models with a full set of emergency patient dummy variables interacted with all explanatory variables. There are 73,629 emergency patients in our sample. Only 20% of emergency patients bypassed the nearest provider (see Appendix Figure A2). Descriptive statistics for this patient group are reported in Appendix Table A3. Emergency patients' choice sets are the 50 closest providers who carried out hip replacement surgery on at least 30 emergency patients in this year. This rules out private and specialised providers who only treat elective hip replacement patients. 708 (1.0%) emergency patients were dropped because they attended a provider not in their choice set. All main effects still pertain to the elective reference patient.

²¹ We are not concerned about the implications of response rates for the hospital level case-mix adjusted OHS change scores as these have been shown to be robust to variations in response rate (Gomes et al. forthcoming)

²² As a check, we first re-estimate the responder only model with the addition of provider pre-operative response rates. The results are robust to this sensitivity analysis, with the WTT of 1.4km (SE=0.102) for a standard deviation increase in PROM quality being slightly larger than in our preferred specification (full results available on request).

Table 3: Choice models allowing for patient pre-operative Oxford Hip Score

	Patients with pre-op OHS (1)		Patients with pre-op OHS (2)		All patients (3)					
					Responders (3a)		Non-responders (3b)		Difference (3c)	
Variable	Est	SE	Est	SE	Est	SE	Est	SE	Est	SE
Main effects										
Distance (in km)	-0.185	0.002***	-0.185	0.002***	-0.185	0.002***	-0.188	0.007***	0.003	0.007
Distance ²	0.000	0.000***	0.000	0.000***	0.000	0.000***	0.000	0.000**	0.000	0.000
NHS trust - medium	-0.526	0.037***	-0.646	0.037***	-0.641	0.037***	-0.558	0.039***	-0.083	0.034*
NHS trust - multi-service	-0.902	0.133***	-0.973	0.131***	-0.965	0.130***	-0.519	0.104***	-0.446	0.123***
NHS trust - small	-0.820	0.045***	-0.907	0.045***	-0.902	0.044***	-0.857	0.045***	-0.045	0.041
NHS trust - specialist	1.052	0.079***	0.869	0.082***	0.856	0.082***	0.973	0.089***	-0.117	0.070
NHS trust - teaching	-0.445	0.039***	-0.503	0.039***	-0.489	0.039***	-0.608	0.039***	0.119	0.038**
Independent sector treatment centre	-1.373	0.065***	-1.499	0.063***	-1.515	0.063***	-1.670	0.067***	0.155	0.059**
Primary care trust	-0.978	0.223***	-1.301	0.223***	-1.298	0.224***	-1.272	0.235***	-0.026	0.232
Waiting time (in months)	-0.011	0.020	0.035	0.020	0.033	0.020	-0.054	0.023*	0.087	0.018***
Change in Oxford Hip Score	0.161	0.010***	0.139	0.010***	0.137	0.010***	0.104	0.011***	0.033	0.010**
28-day emergency readmission rate (in %)	-0.050	0.006***	-0.046	0.006***	-0.046	0.006***	-0.052	0.006***	0.006	0.006
28-day mortality rate (in %)	-0.135	0.035***	-0.068	0.032*	-0.067	0.032*	-0.014	0.036	-0.053	0.033
Response rate			2.044	0.093***	2.038	0.092***	-2.287	0.080***	4.325	0.086***
Interaction with pre-operative Oxford Hip Score										
x Distance (in km)	0.001	0.000***	0.001	0.000***						
x Distance ²	0.000	0.000***	0.000	0.000***						
x NHS trust - medium	0.000	0.002	-0.001	0.002						
x NHS trust - multi-service	-0.004	0.007	-0.006	0.007						
x NHS trust - small	-0.002	0.002	-0.002	0.002						
x NHS trust - specialist	0.017	0.004***	0.016	0.004***						
x NHS trust - teaching	-0.005	0.002*	-0.008	0.002***						
x Independent sector treatment centre	0.038	0.003***	0.035	0.003***						
x Primary care trust	-0.002	0.008	-0.005	0.008						
x Waiting time (in months)	0.004	0.001***	0.004	0.001***						
x Change in Oxford Hip Score	0.001	0.001*	0.000	0.001						
x 28-day emergency readmission rate (in %)	0.000	0.000	0.000	0.000						
x 28-day mortality rate (in %)	0.002	0.002	0.003	0.002						
x Response rate			0.020	0.005***						
WTT(OHS change)	1.717	0.111***	1.475	0.113***	1.465	0.112***	1.048	0.136***	0.417	0.133**
WTT(Readmission rate)	-0.941	0.107***	-0.867	0.105***	-0.879	0.105***	-0.932	0.132***	0.054	0.127
WTT(Mortality rate)	-0.474	0.122***	-0.238	0.112*	-0.224	0.107*	-0.045	0.113	-0.180	0.107
Number of patients	71,329		71,329		113,751					
Number of providers	206		206		206					
BIC	182,407		179,628		283,989					
Pseudo R ²	0.649		0.654		0.657					

*** p<0.001; ** p<0.01; * p<0.05

Notes: Conditional logit model of choice of hospital for elective hip replacement patients treated between April 2010 and March 2012. OHS change, waiting time, readmission rate and mortality rate are lagged by one year. Coefficients are marginal utilities. WTT is the ratio of the coefficient on the quality variable to the marginal utility of distance evaluated at the median distance (in km). Models in (1) and (2) are for patients reporting a pre-operation OHS. Model in (3) is for all patients and interacts a dummy variable for reporting a pre-operation OHS. Interaction effects are reported in (3c). All models also contain a full set of interactions of age, gender, past utilisation, Elixhauser comorbidities, and deprivation with hospital characteristics and distance (not reported). Standard errors are clustered at GP practice level.

We report results for two different specifications. The first model in Table 4 compares emergency patients with elective patients who choose NHS or independent providers. However, there are some marked differences in observed characteristics between those two groups. For example, emergency patients are on average 12 years older than elective patients and have over twice as many recorded comorbidities. Hence in the second model reported in Table 5 we compare a set of elective and emergency patients matched exactly on age, gender, past emergency admissions, number of comorbidities, income deprivation and year of treatment. Additionally, we restrict the elective patient sample to those who used an NHS provider that treats at least 30 elective and emergency patients in that year; hence the choice sets are identical for elective and emergency conditional on location.

Table 4: Comparison of marginal utilities for elective and emergency patients

Variable	Elective patients		Emergency patients		Difference	
	Est	SE	Est	SE	Est	SE
Distance (in km)	-0.184	0.002***	-0.217	0.004***	-0.033	0.003***
Distance ²	0.000	0.000***	0.001	0.000***	0.000	0.000***
NHS trust - medium	-0.530	0.030***	-0.571	0.045***	-0.041	0.039
NHS trust - multi-service	-0.603	0.099***	-0.935	0.164***	-0.332	0.145*
NHS trust - small	-0.791	0.038***	-0.823	0.050***	-0.032	0.044
NHS trust - specialist	1.023	0.072***	n/a		n/a	
NHS trust - teaching	-0.445	0.033***	-0.609	0.045***	-0.164	0.042***
Independent sector treatment centre	-1.467	0.045***	n/a		n/a	
Primary care trust	-1.159	0.206***	-1.274	0.258***	-0.115	0.176
Waiting time (in months)	0.013	0.015	-0.010	0.022	-0.023	0.021
Change in Oxford Hip Score	0.118	0.008***	0.048	0.013***	-0.070	0.012***
28-day emergency readmission rate (in %)	-0.052	0.004***	-0.046	0.008***	0.006	0.007
28-day mortality rate (in %)	-0.031	0.026	0.056	0.056	0.087	0.057
WTT(OHS change)	1.285	0.085***	0.523	0.143***	-0.763	0.126***
WTT(Readmission rate)	-0.978	0.079***	-0.870	0.150***	0.109	0.134
WTT(Mortality rate)	-0.085	0.072	0.155	0.155	0.241	0.157
Number of patients	173,032		72,921			
Number of providers	230		138			
BIC	570,669					
Pseudo R ²	0.689					

*** p<0.001; ** p<0.01; * p<0.05

Notes: Conditional logit model of choice of hospital for elective and emergency hip replacement patients treated between April 2010 and March 2013. OHS change, waiting time, readmission rate and mortality rate are lagged by one year. Coefficients are marginal utilities for the 'reference patient'. Elective and emergency patients are not matched on observed characteristics but the 'reference patient' in both patient populations is defined according to the average characteristics of the elective patient sample. WTT is the ratio of the coefficient on the quality variable to the marginal utility of distance evaluated at the median distance (in km). Model is estimated with a full set of dummy variables interacted with hospital characteristics and other interaction terms. All models also contain a full set of interactions of age, gender, past utilisation, Elixhauser comorbidities, and deprivation with hospital characteristics and distance (not reported). Standard errors are clustered at GP practice level.

Table 5: Comparison of marginal utilities for elective and emergency patients - matched sample

Variable	Elective patients		Emergency patients		Difference	
	Est	SE	Est	SE	Est	SE
Distance (in km)	-0.220	0.004***	-0.215	0.004***	0.005	0.005
Distance ²	0.001	0.000***	0.000	0.000***	0.000	0.000***
NHS trust - medium	-0.709	0.050***	-0.560	0.046***	0.149	0.053**
NHS trust - multi-service	-0.710	0.175***	-0.921	0.182***	-0.211	0.213
NHS trust - small	-0.880	0.058***	-0.794	0.053***	0.086	0.062
NHS trust - teaching	-0.468	0.053***	-0.598	0.048***	-0.130	0.058*
Primary care trust	-1.133	0.292***	-1.429	0.314***	-0.296	0.282
Waiting time (in months)	-0.087	0.027**	-0.032	0.024	0.055	0.029
Change in Oxford Hip Score	0.092	0.015***	0.034	0.014*	-0.058	0.016***
28-day emergency readmission rate (in %)	-0.061	0.009***	-0.046	0.008***	0.015	0.010
28-day mortality rate (in %)	-0.050	0.061	0.070	0.063	0.120	0.079
WTT(OHS change)	0.796	0.128***	0.354	0.142*	-0.441	0.149**
WTT(Readmission rate)	-0.893	0.129***	-0.814	0.152***	0.079	0.165
WTT(Mortality rate)	-0.084	0.102	0.140	0.127	0.224	0.148
Number of patients	32,274		32,274			
Number of providers	138		138			
BIC	107,831					
Pseudo R ²	0.771					

*** p<0.001; ** p<0.01; * p<0.05

Notes: Conditional logit model of choice of hospital for elective and emergency hip replacement patients treated between April 2010 and March 2013. OHS change, waiting time, readmission rate and mortality rate are lagged by one year. Coefficients are marginal utilities for the 'reference patient'. Elective and emergency patients are matched exactly on observed characteristics (age, gender, past emergency utilisation in last year (none, once, or more), income deprivation of neighbourhood, number of Elixhauser comorbid conditions, year of treatment) and the 'reference patient' in both patient populations is defined according to the average (prior to matching) characteristics of the elective patient sample. Choice sets include only providers that treat at least 30 elective and 30 emergency hip replacement patient in this period. WTT is the ratio of the coefficient on the quality variable to the marginal utility of distance evaluated at the median distance (in km). Model is estimated with a full set of dummy variables interacted with hospital characteristics and other interaction terms. All models also contain a full set of interactions of age, gender, past utilisation, Elixhauser comorbidities, and deprivation with hospital characteristics and distance (not reported). Standard errors are clustered at GP practice level.

Both models suggest that emergency patients care less about provider OHS changes but have similar preferences over the more traditional quality measures using readmission and mortality rates. In the second specification, with closely matched patients, the estimated marginal utility of OHS changes ($\beta_q^{Emer}=0.034$) is just over one third of that for elective patients ($\beta_q^{Elec}=0.092$) and significant at $p<0.05$. If we assume that emergency patients' demand is entirely inelastic to variation in observed elective quality and that the estimated association for emergency patients is a result of omitted variables that affect emergency and elective patients in the same way, then the difference in the marginal utility of OHS changes ($\beta_q^{\Delta}=0.058$) can be interpreted as a lower bound estimate of the true effect of OHS change score on elective patient utility. The WTT for a one SD increase in OHS change scores then is 0.4km (SE=0.149), which is smaller than that reported in Table 2.

Table 6 shows the main effects from our preferred specification estimated with additional hospital FEs. We find that PROM quality still has a statistically significant effect on demand, whereas emergency readmission rates no longer do. The WTT to travel for PROM quality is however 87% lower than that calculated from the results in Table 2 (0.2km vs 1.3km). This is likely to be due to the fixed effect absorbing part of the effect of time-invariant quality on choice. Results are broadly similar when patients' choice sets are restricted to NHS hospitals, although we now find a counter-intuitive positive effect of waiting time on demand.

Table 6: Choice model controlling for unobserved time-invariant hospital effects

	All providers (1)		NHS providers only (2)	
	Est	SE	Est	SE
Distance (in km)	-0.202	0.002***	-0.231	0.003***
Distance ²	0.000	0.000***	0.001	0.000***
Waiting time (in months)	0.021	0.024	0.053	0.012***
Change in Oxford Hip Score	0.017	0.006**	0.016	0.007*
28-day emergency readmission rate (in %)	0.005	0.004	0.000	0.004
28-day mortality rate (in %)	0.038	0.020	0.021	0.026
WTT(OHS change)	0.168	0.060**	0.124	0.055*
WTT(Readmission rate)	0.089	0.066	-0.0001	0.053
WTT(Mortality rate)	0.095	0.051	0.031	0.039
Number of patients	173,032		148,629	
Number of providers	230		144	
BIC	411,541		260,299	
Pseudo R ²	0.678		0.742	

*** p<0.001; ** p<0.01; * p<0.05

Notes: Conditional logit model of choice of hospital for elective hip replacement patients treated between April 2010 and March 2013. OHS change, waiting time, readmission rate and mortality rate are lagged by one year. Coefficients are marginal utilities. WTT is the ratio of the coefficient on the quality variable to the marginal utility of distance evaluated at the median distance (in km). Model in (1) does not impose restrictions on the type of provider in patients' choice sets. Model in (2) is based on a restricted choice set of NHS providers, thereby excluding patients that selected ISTCs. All models include indicator variables for hospitals (not reported). All models also contain a full set of interactions of age, gender, past utilisation, Elixhauser comorbidities, and deprivation with hospital characteristics and distance (not reported). Standard errors are clustered at GP practice level.

5. The economic effects of quality on demand

We use the results from choice models to illustrate the effect of quality differentiation on hospital demand. Column four and five of Table 7 provide the marginal utilities of the different quality measures and the willingness to travel for a one SD increase in these measures. The sixth and seventh columns show the average total and relative change in demand from a one SD increase in quality, and column eight gives the own quality demand elasticities. We base our calculations on the estimates for our preferred specification in Table 2. This should be kept in mind when interpreting the results presented in this section.

Table 7: Effect sizes of hospital quality measures

Quality indicator	Observed		Marginal utility	Effect of SD increase in quality			Elasticity
	Mean	SD		WTT	Demand change	% Demand change	
Change in Oxford Hip Score	19.8	1.4	0.118	1.3	33.9	9.4	1.3
Emergency readmission rate (in %)	-5.6	2.4	-0.052	-1.0	-25.3	-7.0	-0.2
Mortality rate (in %)	-0.2	0.4	-0.031	-0.1	-2.2	-0.6	0.0

Notes: All calculations based on estimated marginal utilities reported in Table 2. WTT is the ratio of the coefficient on the quality variable to the marginal utility of distance evaluated at the median distance (in km). Changes in volume and elasticities are averaged across hospital-year observations and are weighted by predicted demand $\hat{Y}_{ijt} = \sum_{i \in M_{it}} P_{ijt}$.

The expected increase in demand for a SD increase in OHS is approximately 34 patients, or 9.4% of predicted demand at current quality levels. Increases in readmission and mortality rates are associated with decreases in demand, although the association of mortality and demand is not statistically significant. The effect of a one SD increase in OHS is larger than that of a one SD decrease in readmission rate.

There is substantial variation across providers in the effect of OHS change scores on own demand (Figure 2). The estimated elasticities range from 0.2 to 2.4 (mean = 1.3). About 42% of the variation in elasticities is explained by the amount of competition a provider faces, here measured by the Herfindahl-Hirschman Index (HHI)²³. Providers in more competitive areas (low HHI) face larger quality elasticities than those in less competitive areas (high HHI), with elasticities falling by approximately 0.29 per 0.1 increase in HHI (assuming a linear effect; $p < 0.001$) (Figure 3). Markets are more competitive in areas where independent sector treatment centres are active.

We also examine the effect of changes in the quality of other providers on a provider's demand. Higher cross-quality demand elasticities make it more likely that increases in one provider's quality will trigger an increase in the quality of other providers. Figure 4 shows how cross-quality elasticities decline rapidly as the distances between providers increase. Whereas a 1% increase in a competitor's PROM quality is associated with a -0.63% reduction in demand if the competitor is located within 10 km, this reduces to -0.23% when the competitor is 30km away. To put this into context, Santos et al. (forthcoming) report an average cross-quality elasticity of demand for general practitioners in England of -0.04 (using a 10km radius).

²³ The HHI for provider j is calculated as the sum of the squared market shares of all providers $j^* = 1, \dots, J^*$ that service LSOA $a = 1, \dots, A$, here denoted as s_{aj} , weighted by the proportion of the provider's observed total activity originating from this LSOA, s_{ja} , so that

$$HHI_j = \sum_a s_{ja} * \left[\sum_{j^*} (s_{aj^*})^2 \right] \quad (10)$$

Hospital catchment areas are defined as all LSOAs within a radius of 30km around the hospital.




Figure 2: Distribution of changes in hospital demand as a result of a SD increase in Oxford Hip Score change scores and quality elasticity of demand




Figure 3: Differences in quality elasticity of demand between providers in competitive (low HHI) and non-competitive (high HHI) markets




Figure 4: Percentage change in demand as a result of percentage change in competitor's quality

6. Discussion and concluding remarks

The collection of patient-reported outcome measures has been introduced in England with the ambition that these new metrics of hospital quality would influence patient choice of hospital (Department of Health 2008). This paper is the first to test the relationship between observed hospital PROM quality and demand for elective hip replacement surgery. It uses data on observed choices for all NHS-funded patients treated between April 2010 and March 2013 in private and public hospitals in England. In order to address potential endogeneity bias we implement an empirical strategy based on lagged explanatory variables, hospital fixed effects and a control group design based on demand for emergency hip replacement.

Our results suggest that elective hospital demand is statistically significantly associated with observed quality as measured by PROMs and other metrics. While individual patients are not very sensitive to quality differences - the estimated willingness to travel for a standard deviation increase in PROM quality is less than 1.3km - the number of potential patients in a hospital's market implies that hospitals can attract an increase in elective activity of approximately 34 new patients, or 9% of existing activity levels, if they find ways to improve PROM quality by one standard deviation. Hospital demand is more responsive to a one standard deviation of PROM quality than one standard deviation of emergency readmission rates, and there is no statistically significant association with mortality rates after hip replacement surgery.

Our findings that choice responds to quality suggest that providers could compete on quality. However, the change in activity that would arise after a change in quality may be modest. First, a standard deviation increase in OHS (equivalent to 1.4 points) would be a substantial improvement in quality for any provider and difficult to achieve. For comparison, the average year-on-year improvement in hospital PROM scores is 0.196 OHS points, or less than 15% of the observed standard deviation. Second, we show that the effect of quality changes on the providers' ability to attract patients away from local competitors diminishes rapidly as distance increases. This may result in local quasi-monopolies where quality improvements have little effect on demand. Finally, our estimated effect is likely to be an upper bound estimate and our analysis on emergency patients suggests that the coefficient of demand to quality could be up to 30% smaller. Taken together, the incentive effect of patients 'voting with their feet' and demanding higher quality is likely to be limited.

There are several policy levers which may be used to ensure that PROM quality information is used to inform hospital choice (Marshall et al. 2004; Faber et al. 2009). Many patients may still not know about hospital PROM scores and more active dissemination to the general public may be required (e.g. by adding the information to the Choose & Book system). Some patients may find it difficult to access this information, for example if they do not have access to the internet. There is a lack of evidence on the extent to which patients and general practitioners are aware of this information and consider it as part of their decision-making process. Similarly, the information may not be sufficiently meaningful to them in its current format. A recent study by Hildon et al. (2012) showed that a high proportion of patients and doctors do not consider the reported PROMs to have an intuitive metric and thus struggle to interpret provider scores. Finally, some patients may not consider variation between hospitals sufficiently large to be considered important. Some of these points may resolve over time, whereas others require targeted policy intervention to improve the dissemination of quality information.

We also explore whether patient preferences vary according to observed and unobserved patient characteristics. We find that the preference for PROM quality increases with age and decreases with income deprivation, comorbidity burden and past utilisation. Qualitatively similar results are obtained for preferences for quality as approximated by emergency readmission rates. Interestingly, we do not

find evidence that preferences for quality vary with pre-operative health status as reported by the patient herself. But because healthier patients are more willing to travel, they have *ceteris paribus* a higher willingness to travel for quality. Hence, the '*distance bias*' described by Gowrisankaran and Town (1999) is likely to occur not because more morbid patients request higher quality, but because they derive different disutility from travel. This finding may be specific to the condition under study as osteoarthritis and other conditions that require hip replacement reduce patients' mobility, and more severely morbid patients thus may be less able or willing to travel.

There remains scope for further research. For example, we cannot disentangle whether the estimated effect is driven by patients' choices versus general practitioners choices acting on their behalf. We conjecture it is due to both. We also did not test whether the first release of PROM information in 2009/10 constituted news to patients and how this impacted on providers' demand.

In conclusion, the results in this paper provide some first evidence to suggest that hospital demand for hip replacement responds to hospital quality as captured by changes in patient-reported health status.

References

- Appleby, J. and N. Devlin (2004). *Measuring success in the NHS: using patient assessed health outcomes to manage the performance of health care providers*. The King's Fund. London.
- Beckert, W., M. Christensen and K. Collyer (2012). 'Choice of NHS-funded hospital services in England'. *The Economic Journal* 122, 400–417.
- Belmont, P. J., G. P. Goodman, B. R. Waterman, J. O. Bader and A. J. Schoenfeld (2014). 'Thirty-Day Postoperative Complications and Mortality Following Total Knee Arthroplasty - Incidence and Risk Factors Among a National Sample of 15,321 Patients'. *The Journal of Bone & Joint Surgery* 96 (1), 20–26.
- Berstock, J. R., A. D. Beswick, E. Lenguerrand, M. R. Whitehouse and A. W. Blom (2014). 'Mortality after total hip replacement surgery: A systematic review'. *Bone and Joint Research* 3 (6), 175–182.
- Besley, T. and M. Ghatak (2003). 'Incentives, Choice, and Accountability in the Provision of Public Services'. *Oxford Review of Economic Policy* 19 (2), 235–249.
- Brekke, K., H. Gravelle, L. Siciliani and O. Straume (2014). 'Patient Choice, Mobility and Competition Among Health Care Providers'. In: *Health Care Provision and Patient Mobility*. Ed. by R. Levaggi and M. Montefiori. Springer.
- Brooks, R. (1996). 'EuroQol: the current state of play'. *Health Policy* 37, 53–72.
- Browne, J., L. Jamieson, J. Lewsey, J. van der Meulen, L. Copley and N. Black (2008). 'Case-mix & patients' reports of outcome in Independent Sector Treatment Centres: Comparison with NHS providers'. *BMC Health Services Research* 8 (1), 78.
- Commission on the Future of Health and Social Care in England (2014). *The UK private health market*. Appendix to 'A new settlement for Health and Social Care'. London: The King's Fund.
- Dawson, J., R. Fitzpatrick, A. Carr and D. Murray (1996). 'Questionnaire on the perceptions of patients about total hip replacement'. *Journal of Bone & Joint Surgery, British Volume* 78-B, 185–190.
- Department of Health (2008). *Guidance on the routine collection of Patient Reported Outcome Measures (PROMs)*. The Stationary Office, London.
- (2012). *Patient Reported Outcome Measures (PROMs) in England: The case-mix adjustment methodology*. The Stationary Office, London.
- Elixhauser, A., C. Steiner, D. Harris and R. Coffey (1998). 'Comorbidity measures for use with administrative data'. *Medical Care* 36 (1), 8–27.
- Faber, M., M. Bosch, H. Wollersheim, S. Leatherman and R. Grol (2009). 'Public reporting in health care: how do consumers use quality-of-care information?: A systematic review'. *Medical Care* 47 (1), 1–8.
- Gaynor, M., H. Seider and W. Vogt (2005). 'The Volume-Outcome Effect, Scale Economies, and Learning by Doing'. *American Economic Review* 95 (2), 243–247.
- Gaynor, M., C. Propper and S. Seiler (2012). *Free to Choose? Reform and Demand Response in the English National Health Service*. Working Paper 18574. National Bureau of Economic Research.
- Geweke, J., G. Gowrisankaran and R. J. Town (2003). 'Bayesian Inference for Hospital Quality in a Selection Model'. *Econometrica* 71 (4), 1215–1238.
- Gomes, M., N. Gutacker, C. Bojke and A. Street (forthcoming). 'Addressing missing data in patient-reported outcome measures (PROMs): implications for comparing provider performance.' *Health Economics*.
- Gowrisankaran, G. and R. J. Town (1999). 'Estimating the quality of care in hospitals using instrumental variables'. *Journal of Health Economics* 18 (6), 747–767.
- Gravelle, H., R. Santos and L. Siciliani (2014). 'Does a hospital's quality depend on the quality of other hospitals? A spatial econometrics approach'. *Regional Science and Urban Economics* 49, 203–216.

- Gutacker, N., K. Bloor and R. Cookson (2015). 'Comparing the performance of the Charlson/Deyo and Elixhauser comorbidity measures across five European countries and three conditions'. *The European Journal of Public Health* 25 (Suppl 1), 15–20.
- Gutacker, N., C. Bojke, S. Daidone, N. Devlin and A. Street (2013). 'Hospital Variation in Patient-Reported Outcomes at the Level of EQ-5D Dimensions: Evidence from England'. *Medical Decision Making* 33 (6), 804–818.
- Gutacker, N., A. Street, M. Gomes and C. Bojke (forthcoming). 'Should English healthcare providers be penalised for failing to collect patient-reported outcome measures (PROMs)?' *Journal of the Royal Society of Medicine*.
- Health & Social Care Information Centre (2013a). *Emergency readmissions to hospital within 28 days of discharge: primary hip replacement surgery*. Compendium of Population Health Indicators. Accessed on 24/11/2014. URL: https://indicators.ic.nhs.uk/download/NCHOD/Specification/Spec_33D_533ISP4CPP1_12_V1.pdf.
- (2013b). *Finalised Patient Reported Outcome Measures (PROMs) in England: April 2011 to March 2012*. Accessed on 26/03/2015. URL: <http://www.hscic.gov.uk/catalogue/PUB11359/final-proms-eng-apr11-mar12-fin-report-v2.pdf>.
- Hensher, D. A. and W. H. Greene (2003). 'The Mixed Logit model: The state of practice'. *Transportation* 30, 133–176.
- Hildon, Z., D. Allwood and N. Black (2012). 'Making data more meaningful. Patients' views of the format and content of quality indicators comparing health care providers'. *Patient Education and Counseling* 88, 298–304.
- Hodgkin, D. (1996). 'Specialized service offerings and patients' choice of hospital: the case of cardiac catheterization'. *Journal of Health Economics* 15 (3), 305–332.
- Hole, A. R. (2007a). 'A comparison of approaches to estimating confidence intervals for willingness to pay measures'. *Health Economics* 16 (8), 827–840.
- (2007b). 'Fitting mixed logit models by using maximum simulated likelihood'. *The Stata Journal* 7 (3), 388–401.
- Hunt, L. P., Y. Ben-Shlomo, E. M. Clark, P. Dieppe, A. Judge, A. J. MacGregor, J. H. Tobias, K. Vernon and A. W. Blom (2013). '90-day mortality after 409,096 total hip replacements for osteoarthritis, from the National Joint Registry for England and Wales: a retrospective analysis'. *The Lancet* 382 (9898), 1097–1104.
- Hutchings, A., J. Neuburger, J. van der Meulen and N. Black (2014). 'Estimating recruitment rates for routine use of patient reported outcome measures and the impact on provider comparisons.' *BMC Health Services Research* 14 (1), 66.
- Iversen, T. and L. Siciliani (2011). 'Non-price rationing and waiting times'. In: *The Oxford Handbook of Health Economics*. Ed. by S. Glied and P. C. Smith. Oxford: Oxford University Press.
- Judge, A., J. Chard, I. Learmonth and P. Dieppe (2006). 'The effects of surgical volumes and training centre status on outcomes following total joint replacement: analysis of the Hospital Episode Statistics for England'. *Journal of Public Health* 28 (2), 116–124.
- Jung, K., R. Feldman and D. Scanlon (2011). 'Where would you go for your next hospitalization?' *Journal of Health Economics* 30 (4), 832–841.
- Lilford, R. and P. Pronovost (2010). 'Using hospital mortality rates to judge hospital performance: a bad idea that just won't go away'. *BMJ* 340, 955–957.
- Luft, H. S., S. S. Hunt and S. C. Maerki (1987). 'The Volume-Outcome Relationship: Practice-Makes-Perfect or Selective-Referral Patterns?' *Health Services Research* 22 (2), 157–182.
- Mäkelä, K. T., U. Häkkinen, M. Peltola, M. Linna, H. Kröger and V. Remes (2011). 'The effect of hospital volume on length of stay, re-admissions, and complications of total hip arthroplasty'. *Acta Orthopaedica* 82 (1), 20–26.

- Marshall, M. N., P. S. Romano and H. T. O. Davies (2004). 'How do we maximize the impact of the public reporting of quality of care?' *International Journal for Quality in Health Care* 16 (S1), 57–63.
- McFadden, D. (1974). 'Conditional logit analysis of qualitative choice behaviour'. In: *Frontiers in economics*. Ed. by P. Zarembka. Vol. 4. New York: Academic Press, 105–142.
- Moja, L., A. Piatti, V. Pecoraro, C. Ricci, G. Virgili, G. Salanti, L. Germagnoli, A. Liberati and G. Banfi (2012). 'Timing Matters in Hip Fracture Surgery: Patients Operated within 48 Hours Have Better Outcomes. A Meta-Analysis and Meta-Regression of over 190,000 Patients'. *PLoS ONE* 7 (10), e46175.
- Monstad, K., L. Engester and B. Espehaug (2006). *Patient preferences for choice of hospital*. Tech. rep. Health Economics Bergen Working Paper No 05/06.
- Moscone, F., E. Tosetti and G. Vittadini (2012). 'Social interaction in patients' hospital choice: evidence from Italy'. *Journal of the Royal Statistical Society: Series A (Statistics in Society)* 175 (2), 453–472.
- Naylor, C. and S. Gregory (2009). 'Independent sector treatment centres'. The King's Fund Briefing.
- Neuburger, J., A. Hutchings, J. van der Meulen and N. Black (2013). 'Using patient-reported outcomes (PROs) to compare the provider of surgery: does the choice of measure matter?' *Medical Care* 51 (6), 517–523.
- NICE (2011). *Hip fracture: The management of hip fracture in adults*. NICE Guidelines 124. London.
- Noble, M., G. Wright, G. Smith and C. Dibben (2006). 'Measuring multiple deprivation at the small-area level'. *Environment and Planning A* 38 (1), 169–185.
- Nuttall, D., D. Parkin and N. Devlin (2015). 'Inter-provider comparison of patient-reported outcomes: developing an adjustment to account for differences in patient case mix'. *Health Economics* 24 (1), 41–54.
- Pope, D. (2009). 'Reacting to rankings: Evidence from "America's best hospitals"'. *Journal of Health Economics* 28 (6), 1154–1165.
- Propper, C., M. Damiani, G. Leckie and J. Dixon (2007). 'Impact of patients' socioeconomic status on the distance travelled for hospital admission in the English National Health Service'. *Journal of Health Services Research & Policy* 12 (3), 153–159.
- Ruwaard, S. and R. Douven (2014). *Quality and hospital choice for cataract treatments: the winner takes most*. CPB Netherlands Bureau for Economic Policy Analysis.
- Santos, R., H. Gravelle and C. Propper (forthcoming). 'Does quality affect patients' choice of doctor? Evidence from England'. *The Economic Journal*.
- Siciliani, L., V. Moran and M. Borowitz (2014). 'Measuring and comparing health care waiting times in OECD countries'. *Health Policy* 118 (3), 292–303.
- Sivey, P. (2008). 'The effect of hospital quality on choice of hospital for elective heart operations in England'. PhD thesis. University of York.
- (2012). 'The effect of waiting time and distance on hospital choice for English cataract patients'. *Health Economics* 21, 444–456.
- Train, K. E. (2003). *Discrete Choice Methods with Simulation*. 1st ed. Cambridge: Cambridge University Press.
- Varkevisser, M., S. A. Geest and F. T. Schut (2010). 'Assessing hospital competition when prices don't matter to patients: the use of time-elasticities'. *International Journal of Health Care Finance and Economics* 10 (1), 43–60.
- (2012). 'Do patients choose hospitals with high quality ratings? Empirical evidence from the market for angioplasty in the Netherlands'. *Journal of Health Economics* 31 (2), 371–378.
- Vrangbaek, K., R. Robertson, U. Winblad, H. Van de Bovenkamp and A. Dixon (2012). 'Choice policies in Northern European health systems'. *Health Economics, Policy and Law* 7 (S1), 47–71.
- Wooldridge, J. M. (2002). *Econometric Analysis of Cross Section and Panel Data*. Cambridge, Massachusetts: MIT Press.

Appendix

Table A1: Mixed logit choice model

Variable	Mean		Standard deviation	
	Est	SE	Est	SE
Distance (in km)	-0.184	0.002***		
Distance ²	0.000	0.000***		
NHS trust - medium	-0.530	0.030***		
NHS trust - multi-service	-0.603	0.099***		
NHS trust - small	-0.791	0.038***		
NHS trust - specialist	1.023	0.072***		
NHS trust - teaching	-0.445	0.033***		
Independent sector treatment centre	-1.467	0.045***		
Primary care trust	-1.159	0.206***		
Waiting time (in months)	0.013	0.015		
Change in Oxford Hip Score	0.118	0.008***	0.000	0.001
28-day emergency readmission rate (in %)	-0.052	0.004***	0.000	0.001
28-day mortality rate (in %)	-0.031	0.026	-0.002	0.005
WTT(OHS change)	1.287	0.085***		
WTT(Readmission rate)	-0.981	0.079***		
WTT(Mortality rate)	-0.086	0.072		
Number of patients	173,032			
Number of providers	230			
BIC	461,041			

*** p<0.001; ** p<0.01; * p<0.05

Notes: Random coefficient (mixed) multinomial logit model of choice of hospital for elective hip replacement patients treated between April 2010 and March 2013. OHS change, waiting time, readmission rate and mortality rate are lagged by one year. Coefficients are marginal utilities. Random coefficients are specified for OHS change, readmission rate and mortality rate and estimates. WTT is the ratio of the coefficient on the quality variable to the marginal utility of distance evaluated at the median distance (in km). Interaction terms with patient characteristics not reported (available on request). Standard errors are clustered at GP practice level. The mean coefficients do differ from those reported in Table 2 for the conditional logit model if there is less rounding.

Table A2: Sensitivity analyses

Variable	Model (1)		Model (2)		Model (3)		Model (4)	
	Est	SE	Est	SE	Est	SE	Est	SE
Main effects								
Distance (in km)	-0.183	0.002***	-0.184	0.002***	-0.184	0.002***	-0.207	0.003***
Distance ²	0.000	0.000***	0.000	0.000***	0.000	0.000***	0.001	0.000***
NHS trust - medium	-0.508	0.032***	-0.530	0.030***	-0.540	0.031***	-0.643	0.036***
NHS trust - multi-service	-0.656	0.090***	-0.615	0.099***	-0.612	0.097***	-0.604	0.098***
NHS trust - small	-0.740	0.037***	-0.792	0.038***	-0.790	0.038***	-0.909	0.042***
NHS trust - specialist	1.019	0.071***	1.002	0.072***	1.065	0.073***	1.047	0.083***
NHS trust - teaching	-0.502	0.033***	-0.432	0.033***	-0.412	0.032***	-0.513	0.037***
Independent sector treatment centre	-1.472	0.068***	-1.522	0.037***	-1.398	0.044***		
Primary care trust	-1.071	0.200***	-1.153	0.206***	-1.111	0.206***	-1.195	0.213***
Waiting time (in months)	0.033	0.018	-0.193	0.076*	0.007	0.015	-0.029	0.018
Change in PROM score	0.104	0.007***	0.115	0.008***	4.935	0.284***	0.121	0.010***
28-day emergency readmission rate (in %)	-0.054	0.004***	-0.052	0.004***	-0.054	0.004***	-0.054	0.006***
28-day mortality rate (in %)	-0.044	0.024	-0.032	0.026	-0.026	0.026	0.033	0.035
Interaction with distance								
x Patient age	-0.001	0.000***	-0.002	0.000***	-0.002	0.000***	-0.002	0.000***
x Male	0.001	0.001	0.002	0.001	0.002	0.001	0.005	0.002**
x Past utilisation	-0.003	0.002	-0.003	0.002	-0.003	0.002	-0.001	0.002
x Comorbidity count	-0.005	0.001***	-0.004	0.001***	-0.003	0.001***	-0.001	0.001
x Income deprivation	-0.179	0.015***	-0.186	0.017***	-0.189	0.018***	-0.241	0.022***
Interaction with waiting time								
x Patient age	0.003	0.000***	0.008	0.002**	0.003	0.000***	0.003	0.001***
x Male	-0.018	0.010	-0.068	0.051	-0.010	0.009	0.001	0.012
x Past utilisation	-0.005	0.012	0.049	0.059	-0.007	0.012	0.008	0.013
x Comorbidity count	-0.036	0.007***	-0.057	0.033	-0.018	0.007**	-0.016	0.007*
x Income deprivation	0.032	0.086	1.180	0.458*	0.083	0.084	0.007	0.107
Interaction with change in PROM score								
x Patient age	0.001	0.000***	0.001	0.000*	0.047	0.010***	0.001	0.000
x Male	0.004	0.005	-0.007	0.005	-0.300	0.214	-0.002	0.006
x Past utilisation	-0.014	0.006*	-0.008	0.007	-0.327	0.240	-0.012	0.007
x Comorbidity count	-0.009	0.003**	-0.009	0.003**	-0.475	0.128***	-0.004	0.004
x Income deprivation	-0.389	0.046***	-0.448	0.047***	-18.634	1.698***	-0.568	0.056***
Interaction with 28-day emergency readmission rate								
x Patient age	0.000	0.000*	0.000	0.000*	0.000	0.000	-0.001	0.000**
x Male	0.002	0.003	0.000	0.003	0.000	0.003	-0.003	0.004
x Past utilisation	0.012	0.004**	0.012	0.003***	0.012	0.004***	0.010	0.004**
x Comorbidity count	0.003	0.002	0.001	0.002	0.001	0.002	0.003	0.002
x Income deprivation	0.082	0.026**	0.124	0.026***	0.138	0.026***	0.121	0.034***
Interaction with 28-day mortality rate								
x Patient age	-0.002	0.001*	-0.001	0.001	-0.001	0.001	-0.001	0.001
x Male	-0.051	0.021*	-0.053	0.022*	-0.049	0.022*	-0.058	0.031
x Past utilisation	0.039	0.030	0.049	0.026	0.047	0.026	0.060	0.032
x Comorbidity count	-0.001	0.015	-0.010	0.015	-0.009	0.015	-0.004	0.018
x Income deprivation	-0.059	0.172	-0.023	0.168	-0.049	0.167	-0.230	0.210
WTT(PROM change)	1.174	0.086***	1.261	0.086***	1.284	0.076***	1.078	0.089***
WTT(Readmission rate)	-1.042	0.081***	-0.983	0.080***	-1.034	0.079***	-0.824	0.089***
WTT(Mortality rate)	-0.125	0.067	-0.091	0.073	-0.074	0.074	0.055	0.059
Number of patients	176,471		173,032		171,737		148,629	
Number of providers	233		230		225		144	
BIC	473,568		460,956		450,787		299,221	
Pseudo R ²	0.645		0.637		0.639		0.701	

*** p<0.001; ** p<0.01; * p<0.05

Notes: Conditional logit model of choice of hospital for elective hip replacement patients treated between April 2010 and March 2013. PROM change, waiting time, readmission rate and mortality rate are lagged by one year if not otherwise stated. Coefficients are marginal utilities. WTT is the ratio of the coefficient on the quality variable to the marginal utility of distance evaluated at the median distance (in km). Interaction terms with distance² and provider type not reported (available on request). Standard errors are clustered at GP practice level.

Model (1) - PROM change, waiting time, readmission rate and mortality rate are contemporaneous. Based on observed choices for patients treated between April 2009 and March 2012. Since April 2012 PROM scores have been reported separately for primary and revision hip replacement surgeries, so that our measures of PROM quality are no longer comparable.

Model (2) - Proportion of patients waiting longer than 120 days substituted for waiting time (both lagged).

Model (3) - Lagged EQ-5D change scores substituted for lagged OHS change scores.

Model (4) - Patients' choice sets exclude independent sector treatment centres.

Table A3: Descriptive statistics - emergency sample

Variable	Obs	Mean	SD	ICC
Patient characteristics				
Distance travelled (in km)	73,629	14.2	27.1	
Distance travelled past closest provider (in km)	73,629	4.2	25.4	
Number of providers within 10km radius	73,629	1.0	1.4	
Number of providers within 30km radius	73,629	5.4	5.4	
Age	73,629	80.9	9.8	
Male	73,629	0.27	0.44	
Past utilisation	73,629	0.65	1.17	
Number of Elixhauser conditions	73,629	0.99	1.56	
Income deprivation	73,629	0.14	0.10	
Provider characteristics				
Observed volume	394	186.9	87.1	80.7%
Waiting time (in months)	394	3.0	0.7	46.1%
Change in Oxford Hip Score	394	19.4	1.3	49.1%
28-day emergency readmission rate (in %)	394	5.99	2.20	38.2%
28-day mortality rate (in %)	394	0.20	0.25	5.3%

Obs = Observations; SD = Standard deviation; ICC = Intraclass correlation coefficient.

Notes: Patient characteristics for patients choosing provider between April 2010 and March 2013. Provider waiting time, change in Oxford Hip Score, readmission rate, mortality rate are based on elective patients treated by the respective providers and are for financial years 2009/10 to 2011/12.




Figure A1: Waiting time for elective hip replacement surgery in England - 2000 to 2012




Figure A2: Percentage of emergency patients who went to their Nth nearest hospital