



UNIVERSITY OF LEEDS

This is a repository copy of *Left atrial size and function in a South Asian population and their potential influence on the risk of atrial fibrillation*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/137957/>

Version: Accepted Version

---

**Article:**

O'Neill, J, Swoboda, PP [orcid.org/0000-0001-7162-7079](https://orcid.org/0000-0001-7162-7079), Plein, S et al. (1 more author) (2018) Left atrial size and function in a South Asian population and their potential influence on the risk of atrial fibrillation. *Clinical cardiology*, 41 (10). pp. 1379-1385. ISSN 0160-9289

<https://doi.org/10.1002/clc.23064>

---

© 2018 Wiley Periodicals, Inc. This is the peer reviewed version of the following article: O'Neill, J, Swoboda, PP, Plein, S et al. (1 more author) (2018) Left atrial size and function in a South Asian population and their potential influence on the risk of atrial fibrillation. *Clinical cardiology*, 41 (10). pp. 1379-1385, which has been published in final form at <https://doi.org/10.1002/clc.23064>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

**Supplemental Table 1.** Univariable and multivariable linear regression analysis for LAV<sub>min</sub>/BSA, LAV<sub>max</sub>/BSA

<b>LAV<sub>min</sub>/BSA</b>						
	Univariable model			Multivariable model		
	Beta coefficient	Standard error	p value	Beta coefficient	Standard error	p value
Age	0.457	1.327	<b>&lt;0.001</b>	0.383	0.034	<b>&lt;0.001</b>
Male gender	0.153	1.029	0.094	-0.012	0.987	0.894
South Asian	-0.274	1.001	<b>0.001</b>	-0.201	0.837	<b>0.008</b>
Body mass index	0.259	0.098	<b>0.004</b>	0.104	0.086	0.183
LV ejection fraction	0.108	0.043	0.239			
LV mass/BSA	0.403	0.046	<b>&lt;0.001</b>	0.357	0.048	<b>&lt;0.001</b>
Hypertension	0.229	1.756	<b>0.012</b>	0.083	1.518	0.286
Diabetes mellitus	0.014	2.388	0.882			
<b>LAV<sub>max</sub>/BSA</b>						
Age	0.356	0.064	<b>&lt;0.001</b>	0.291	0.066	<b>0.001</b>
Male gender	0.112	1.882	0.221			
South Asian	-0.325	1.792	<b>&lt;0.001</b>	-0.267	1.625	<b>0.001</b>
Body mass index	0.206	0.181	<b>0.024</b>	0.083	0.166	0.317
LV ejection fraction	0.093	0.079	0.312			
LV mass/BSA	0.339	0.085	<b>&lt;0.001</b>	0.277	0.078	0.001
Hypertension	0.203	3.214	<b>0.026</b>	0.089	2.970	0.288
Diabetes mellitus	0.000	4.346	0.999			

**Abbreviations:** LAV<sub>min</sub>, minimum left atrial volume; LAV<sub>max</sub>, maximum left atrial volume; LV, left ventricular; BSA, body surface area.