

A multiple testing approach to the regularisation of large sample correlation matrices*

Natalia Bailey
Monash University

M. Hashem Pesaran
University of Southern California and Trinity College, Cambridge

L. Vanessa Smith
University of York

October 14, 2018

Abstract

This paper proposes a regularisation method for the estimation of large covariance matrices that uses insights from the multiple testing (MT) literature. The approach tests the statistical significance of individual pair-wise correlations and sets to zero those elements that are not statistically significant, taking account of the multiple testing nature of the problem. The effective p-values of the tests are set as a decreasing function of N (the cross section dimension), the rate of which is governed by the nature of dependence of the underlying observations, and the relative expansion rates of N and T (the time dimension). In this respect, the method specifies the appropriate thresholding parameter to be used under Gaussian and non-Gaussian settings. The MT estimator of the sample correlation matrix is shown to be consistent in the spectral and Frobenius norms, and in terms of support recovery, so long as the true covariance matrix is sparse. The performance of the proposed MT estimator is compared to a number of other estimators in the literature using Monte Carlo experiments. It is shown that the MT estimator performs well and tends to outperform the other estimators, particularly when N is larger than T .

JEL Classifications: C13, C58

Keywords: High-dimensional data, Multiple testing, Non-Gaussian observations, Sparsity, Thresholding, Shrinkage

*Corresponding author: L. Vanessa Smith, Department of Economics, University of York, Heslington, York, YO10 5DD, UK. Email: vanessa.smith@york.ac.uk.

1 Introduction

Improved estimation of covariance matrices is a problem that features prominently in a number of areas of multivariate statistical analysis. In finance it arises in portfolio selection and optimisation (Ledoit and Wolf (2003)), risk management (Fan *et al.* (2008)) and testing of capital asset pricing models (Sentana (2009)). In global macroeconomic modelling with many domestic and foreign channels of interactions, error covariance matrices must be estimated for impulse response analysis and bootstrapping (Pesaran *et al.* (2004); Dees *et al.* (2007)). In the area of bioinformatics, covariance matrices are required when inferring gene association networks (Carroll (2003); Schäfer and Strimmer (2005)). Such matrices are further encountered in fields including meteorology, climate research, spectroscopy, signal processing and pattern recognition.

Importantly, the issue of consistently estimating the population covariance matrix, $\Sigma = (\sigma_{ij})$, becomes particularly challenging when the number of variables, N , is larger than the number of observations, T . In this case, one way of obtaining a suitable estimator for Σ is to appropriately restrict the off-diagonal elements of its sample estimate denoted by $\hat{\Sigma}$. Numerous methods have been developed to address this challenge, predominantly in the statistics literature. See Pourahmadi (2011) for an extensive review and references therein. Some approaches are regression-based and make use of suitable decompositions of Σ such as the Cholesky decomposition (see Pourahmadi (1999), Pourahmadi (2000), Rothman *et al.* (2010), Abadir *et al.* (2014), among others). Others include banding or tapering methods as proposed, for example, by Bickel and Levina (2004), Bickel and Levina (2008b) and Wu and Pourahmadi (2009), which assume that the variables under consideration follow a natural ordering. Two popular regularisation techniques in the literature that do not make use of any ordering assumptions are those of thresholding and shrinkage.

Thresholding involves setting off-diagonal elements of the sample covariance matrix that are in absolute terms below certain threshold values to zero. This approach includes ‘universal’ thresholding put forward by El Karoui (2008) and Bickel and Levina (2008a), and

‘adaptive’ thresholding proposed by Cai and Liu (2011). Universal thresholding applies the same thresholding parameter to all off-diagonal elements of the unconstrained sample covariance matrix, while adaptive thresholding allows the threshold value to vary across the different off-diagonal elements of the matrix. Furthermore, the selected non-zero elements of $\hat{\Sigma}$ can either be set to their sample estimates or can be adjusted downward. This relates to the concepts of ‘hard’ and ‘soft’ thresholding, respectively. The thresholding approach traditionally assumes that the underlying (population) covariance matrix is *sparse*, where sparsity is loosely defined as the presence of a sufficient number of zeros on each row of Σ such that it is absolute summable row (column)-wise, or more generally in the sense defined by El Karoui (2008). However, Fan *et al.* (2011) and Fan *et al.* (2013) show that such regularisation techniques can be applied even if the underlying population covariance matrix is not sparse, so long as the non-sparsity is characterised by an approximate factor structure. The main challenge in applying this approach lies in the estimation of the thresholding parameter, which is primarily calibrated by cross-validation.

In contrast to thresholding, the shrinkage approach reduces all sample estimates of the covariance matrix towards zero element-wise. More formally, the shrinkage estimator of Σ is defined as a weighted average of the sample covariance matrix and an invertible covariance matrix estimator known as the shrinkage target - see Friedman (1989). A number of shrinkage targets have been considered in the literature that take advantage of *a priori* knowledge of the data characteristics under investigation. Examples of covariance matrix targets can be found in Ledoit and Wolf (2003), Daniels and Kass (1999), Daniels and Kass (2001), Fan *et al.* (2008), and Hoff (2009), among others. Ledoit and Wolf (2004) suggest a modified shrinkage estimator that involves a linear combination of the unrestricted sample covariance matrix with the identity matrix. This is recommended by the authors for more general situations where no natural shrinking target exists. On the whole, shrinkage estimators tend to be stable, but yield inconsistent estimates if the purpose of the analysis is the estimation of the true and false positive rates of the underlying true sparse covariance matrix (the so called ‘support recovery’ problem).

This paper considers an alternative approach using a multiple testing (*MT*) procedure to set the thresholding parameter. A similar idea has been suggested by El Karoui (2008) - p. 2748, who considers testing the $N(N - 1)/2$ null hypotheses that $\sigma_{ij} = 0$, for all $i \neq j$, jointly. But no formal theory has been developed in the literature for this purpose. In our application of this idea we focus on testing the significance of the correlation coefficients, $\rho_{ij} = \sigma_{ij}/\sigma_{ii}^{1/2}\sigma_{jj}^{1/2}$ for all $i \neq j$, which avoids the scaling problem associated with the use of σ_{ij} , and allows us to obtain a universal threshold for all i and j pairs. We use ideas from the multiple testing literature to control the rate at which the spectral and Frobenius norms of the difference between the true correlation matrix $\mathbf{R} = (\rho_{ij})$, and our proposed estimator of it, $\tilde{\mathbf{R}}_{MT} = (\tilde{\rho}_{ij,T})$, tends to zero, and will not be particularly concerned with controlling the overall size of the joint $N(N - 1)/2$ tests of $\rho_{ij} = 0$, for all $i \neq j$.

We establish that $\tilde{\mathbf{R}}_{MT}$ converges to \mathbf{R} in spectral norm at the rate of $O_p\left(\frac{m_N c_p(N)}{\sqrt{T}}\right)$, where m_N is the maximum number of non-zero elements in the off-diagonal rows of \mathbf{R} , $c_p(N) = \Phi^{-1}\left(1 - \frac{p}{2N^\delta}\right)$, $\Phi^{-1}(\cdot)$ is the inverse of the cumulative distribution of a standard normal variate, p is the nominal size of the test, and the choice of $\delta > 0$ is related to the degree of non-Gaussianity of the underlying observations. This is equivalent to the corresponding $O_p\left(m_N \sqrt{\frac{\ln(N)}{T}}\right)$ rate established for the threshold estimator of Σ in the literature, considering that $c_p^2(N)/\ln(N) \rightarrow 2\delta$ as $N \rightarrow \infty$. The main difference between the two approaches is that we use a multiple testing critical value to set the threshold, whilst the literature uses cross validation. It is perhaps also worth noting that our results are established under weaker moment conditions than sub-Gaussianity typically assumed in the literature while comparable to the polynomial-type tail conditions considered in Bickel and Levina (2008a) or Cai and Liu (2011).

In terms of the Frobenius norm, we show that the *MT* estimator converges at the rate of $O_p\left(\sqrt{\frac{m_N N}{T}}\right)$, for suitable choices of the critical value function in our MT procedure. This result holds even if the underlying observations are non-Gaussian. To the best of our knowledge, the only work that addresses the theoretical properties of the thresholding estimator for the Frobenius norm is Bickel and Levina (2008a), who establish the slower

rate of $O_p\left(\sqrt{\frac{m_N N \ln(N)}{T}}\right)$. We also establish conditions under which our proposed estimator consistently recovers the support of the population covariance matrix under Gaussian and non-Gaussian observations, and show that the true positive rate tends to one with probability 1, and the false positive rate and the false discovery rate tend to zero with probability 1, even if N tends to infinity faster than T . We provide conditions under which these results hold.

The performance of the MT estimator is investigated using a Monte Carlo simulation study, and its properties are compared to a number of extant regularised estimators in the literature. The simulation results show that the proposed multiple testing estimator is robust to the typical choices of p used in the literature (10%, 5% and 1%), and performs favourably compared to the other estimators, especially when N is large relative to T . The MT procedure also dominates other regularised estimators when the focus of the analysis is on support recovery.

The rest of the paper is organised as follows: Section 2 outlines some preliminaries, introduces the MT procedure and derives its asymptotic properties. The small sample properties of the MT estimator are investigated in Section 3. Concluding remarks are provided in Section 4. Some of the technical proofs and additional material are provided in an online supplement.

Notations

$O(\cdot)$ and $o(\cdot)$ denote the Big O and Little o notations, respectively. If $\{f_N\}_{N=1}^\infty$ is any real sequence and $\{g_N\}_{N=1}^\infty$ is a sequence of positive real numbers, then $f_N = O(g_N)$ if there exists a positive finite constant K such that $|f_N|/g_N \leq K$ for all N . $f_N = o(g_N)$ if $f_N/g_N \rightarrow 0$ as $N \rightarrow \infty$. $O_p(\cdot)$ and $o_p(\cdot)$ are the equivalent orders in probability. If $\{f_N\}_{N=1}^\infty$ and $\{g_N\}_{N=1}^\infty$ are both positive sequences of real numbers, then $f_N = \Theta(g_N)$ if there exists $N_0 \geq 1$ and positive finite constants K_0 and K_1 , such that $\inf_{N \geq N_0} (f_N/g_N) \geq K_0$, and $\sup_{N \geq N_0} (f_N/g_N) \leq K_1$. The largest and the smallest eigenvalues of the $N \times N$ real symmetric matrix $\mathbf{A} = (a_{ij})$ are denoted by $\lambda_{\max}(\mathbf{A})$ and $\lambda_{\min}(\mathbf{A})$, respectively, its trace by $\text{tr}(\mathbf{A}) =$

$\sum_{i=1}^N a_{ii}$, its maximum absolute column sum norm by $\|\mathbf{A}\|_1 = \max_{1 \leq j \leq N} \left(\sum_{i=1}^N |a_{ij}| \right)$, its maximum absolute row sum norm by $\|\mathbf{A}\|_\infty = \max_{1 \leq i \leq N} \left(\sum_{j=1}^N |a_{ij}| \right)$, its spectral radius by $\rho(\mathbf{A}) = |\lambda_{\max}(\mathbf{A})|$, its spectral (or operator) norm by $\|\mathbf{A}\| = \lambda_{\max}^{1/2}(\mathbf{A}'\mathbf{A})$, its Frobenius norm by $\|\mathbf{A}\|_F = \sqrt{\text{tr}(\mathbf{A}'\mathbf{A})}$. $\xrightarrow{a.s.}$ denotes almost sure convergence, and \xrightarrow{p} convergence in probability. $K, K_0, K_1, C, \varkappa, c_\delta, c_d, \varepsilon_0, \epsilon, \gamma$ and η are finite positive constants, independent of N and T . \sup_{it} will be used to denote $\sup_{1 \leq i \leq N, 1 \leq t \leq T}$. All asymptotics are carried out under N and $T \rightarrow \infty$, *jointly*.

2 Regularising the sample correlation matrix: A multiple testing (MT) approach

Let $\{x_{it}, i \in N, t \in T\}$, $N \subseteq \mathbb{N}$, $T \subseteq \mathbb{Z}$, be a double index process where x_{it} is defined on a suitable probability space (Ω, F, P) , and denote the covariance matrix of $\mathbf{x}_t = (x_{1t}, x_{2t}, \dots, x_{Nt})'$ by

$$\text{Var}(\mathbf{x}_t) = \mathbf{\Sigma} = E[(\mathbf{x}_t - \boldsymbol{\mu})(\mathbf{x}_t - \boldsymbol{\mu})'], \quad (1)$$

where $E(\mathbf{x}_t) = \boldsymbol{\mu} = (\mu_1, \mu_2, \dots, \mu_N)'$, and $\mathbf{\Sigma}$ is an $N \times N$ symmetric, positive definite real matrix with (i, j) element, σ_{ij} . We assume that x_{it} is independent over time, t . We consider the regularisation of the sample covariance matrix estimator of $\mathbf{\Sigma}$, denoted by $\hat{\mathbf{\Sigma}}$, with elements

$$\hat{\sigma}_{ij,T} = T^{-1} \sum_{t=1}^T (x_{it} - \bar{x}_i)(x_{jt} - \bar{x}_j), \text{ for } i, j = 1, 2, \dots, N, \quad (2)$$

where $\bar{x}_i = T^{-1} \sum_{t=1}^T x_{it}$. To this end we assume that $\mathbf{\Sigma}$ is (exactly) sparse defined as follows:

Assumption 1 *The population covariance matrix, $\mathbf{\Sigma} = (\sigma_{ij})$, where $\lambda_{\min}(\mathbf{\Sigma}) \geq \varepsilon_0 > 0$, is sparse in the sense that m_N defined by*

$$m_N = \max_{1 \leq i \leq N} \sum_{j=1}^N I(\sigma_{ij} \neq 0), \quad (3)$$

is $O(N^\vartheta)$ for some $0 \leq \vartheta < 1/2$, where $I(A)$ is an indicator function that takes the value of 1 if A holds and zero otherwise.

A comprehensive discussion of the concept of sparsity applied to Σ and alternative ways of defining it are provided in El Karoui (2008) and Bickel and Levina (2008a).

Remark 1 *The concept of sparsity defined by (3) is particularly suited to economic applications where the focus of the analysis is often on connections in a given network, or support recovery of Σ .¹ But our analysis can be readily extended to allow for approximate sparsity entertained in the literature, with (3) replaced by*

$$m_{q,N} = \max_{1 \leq i \leq N} \sum_{j=1}^N |\sigma_{ij}|^q,$$

for $0 \leq q < 1$, and $m_{q,N} = O(N^\vartheta)$, with $0 \leq \vartheta < 1/2$. To simplify the exposition we focus on the concept of exact sparsity as defined by Assumption 1.

We follow the hard thresholding literature but, as noted above, we employ multiple testing to decide on the threshold value. More specifically, we set to zero those elements of $\mathbf{R} = (\rho_{ij})$ that are statistically insignificant and therefore determine the threshold value as part of a multiple testing strategy. We apply the thresholding procedure explicitly to the correlations rather than the covariances. This has the added advantage that one can use a so-called ‘universal’ threshold rather than making entry-dependent adjustments, which in turn need to be estimated when thresholding is applied to covariances. This feature is in line with the method of Bickel and Levina (2008a) or El Karoui (2008) but shares the properties of the adaptive thresholding estimator developed by Cai and Liu (2011).

Specifically, denote the sample correlation of x_{it} and x_{jt} , computed over $t = 1, 2, \dots, T$, by

$$\hat{\rho}_{ij,T} = \hat{\rho}_{ji,T} = \frac{\hat{\sigma}_{ij,T}}{\sqrt{\hat{\sigma}_{ii,T} \hat{\sigma}_{jj,T}}}, \quad (4)$$

¹A similar argument is also made in Fan *et al.* (2011).

where $\hat{\sigma}_{ij,T}$ is defined by (2). For a given i and j , it is well known that under $H_{0,ij} : \sigma_{ij} = 0$, $\sqrt{T}\hat{\rho}_{ij,T}$ is asymptotically distributed as $N(0, 1)$ for T sufficiently large. This suggests using $T^{-1/2}\Phi^{-1}(1 - \frac{p}{2})$ as the threshold for $|\hat{\rho}_{ij,T}|$, where $\Phi^{-1}(\cdot)$ is the inverse of the cumulative distribution of a standard normal variate, and p is the chosen nominal size of the test, typically taken to be 1% or 5%. However, since there are in fact $N(N - 1)/2$ such tests and N is large, then using the threshold $T^{-1/2}\Phi^{-1}(1 - \frac{p}{2})$ for all $N(N - 1)/2$ pairs of correlation coefficients will yield inconsistent estimates of Σ and fail to recover its support.

A popular approach to the multiple testing problem is to control the overall size of the $n = N(N - 1)/2$ tests jointly (known as family-wise error rate) rather than the size of the individual tests. Let the family of null hypotheses of interest be $H_{01}, H_{02}, \dots, H_{0n}$, and suppose we are provided with the corresponding test statistics, $Z_{1T}, Z_{2T}, \dots, Z_{nT}$, with separate rejection rules given by (using a two-sided alternative)

$$\Pr(|Z_{iT}| > CV_{iT} | H_{0i}) \leq p_{iT},$$

where CV_{iT} is some suitably chosen critical value of the test, and p_{iT} is the observed p -value for H_{0i} . Consider now the family-wise error rate (FWER) defined by

$$FWER_T = \Pr[\cup_{i=1}^n (|Z_{iT}| > CV_{iT} | H_{0i})],$$

and suppose that we wish to control $FWER_T$ to lie below a pre-determined value, p . One could also consider other generalized error rates (see for example Abramovich *et al.* (2006) or Romano *et al.* (2008)). Bonferroni (1935) provides a general solution, which holds for all possible degrees of dependence across the separate tests. Using the union bound, we have

$$\begin{aligned} \Pr[\cup_{i=1}^n (|Z_{iT}| > CV_{iT} | H_{0i})] &\leq \sum_{i=1}^n \Pr(|Z_{iT}| > CV_{iT} | H_{0i}) \\ &\leq \sum_{i=1}^n p_{iT}. \end{aligned}$$

Hence to achieve $FWER_T \leq p$, it is sufficient to set $p_{iT} \leq p/n$. Alternative multiple testing procedures advanced in the literature that are less conservative than the Bonferroni procedure can also be employed. One prominent example is the step-down procedure proposed by Holm (1979) that, similar to the Bonferroni approach, does not impose any further restrictions on the degree to which the underlying tests depend on each other. More recently, Romano and Wolf (2005) proposed step-down methods that reduce the multiple testing procedure to the problem of sequentially constructing critical values for single tests. Such extensions can be readily considered but will not be pursued here.

In our application we scale p by a general function of N , which we denote by $f(N) = c_\delta N^\delta$, where c_δ and δ are finite positive constants, and then derive conditions on δ which ensure consistent support recovery and a suitable convergence rate of the error in estimation of $\mathbf{R} = (\rho_{ij})$. In particular, we show that the choice of δ depends on the nature of dependence of the pairs (y_{it}, y_{jt}) , for all $i \neq j$, and on the relative rate at which N and T rise. As will be shown in Section 2.1, the degree of dependence is defined by $K_v = \sup_{ij} K_v(\boldsymbol{\theta}_{ij})$ where $\boldsymbol{\theta}_{ij}$ is a vector of cumulants of (y_{it}, y_{jt}) . When $\rho_{ij} = 0$ for all i and j , $i \neq j$, this parameter is given by $\varphi_{\max} = \sup_{ij} (\varphi_{ij})$ where $\varphi_{ij} = E(y_{it}^2 y_{jt}^2 | \rho_{ij} = 0) > 0$. In the case where y_{it} and y_{jt} are independent under the null, then $\varphi_{\max} = 1$.

Specifically, the multiple testing (MT) estimator of \mathbf{R} , denoted by $\tilde{\mathbf{R}}_{MT} = (\tilde{\rho}_{ij,T})$, is given by

$$\tilde{\rho}_{ij,T} = \hat{\rho}_{ij,T} I [|\hat{\rho}_{ij,T}| > T^{-1/2} c_p(N)], \quad i = 1, 2, \dots, N-1, \quad j = i+1, \dots, N, \quad (5)$$

where

$$c_p(N) = \Phi^{-1} \left(1 - \frac{p}{2f(N)} \right), \quad (6)$$

with $f(N) = c_\delta N^\delta$, $c_\delta, \delta > 0$. The corresponding MT estimator of $\boldsymbol{\Sigma}$ is given by

$$\tilde{\boldsymbol{\Sigma}}_{MT} = \hat{\mathbf{D}}^{1/2} \tilde{\mathbf{R}}_{MT} \hat{\mathbf{D}}^{1/2}, \quad (7)$$

where $\hat{\mathbf{D}} = \text{diag}(\hat{\sigma}_{11,T}, \hat{\sigma}_{22,T}, \dots, \hat{\sigma}_{NN,T})$. The *MT* procedure can also be applied to de-factored observations following the de-factoring approach of Fan *et al.* (2011) and Fan *et al.* (2013).

2.1 Theoretical properties of the *MT* estimator

To investigate the asymptotic properties of the *MT* estimator defined by (5) we make the following assumption on the bivariate distribution of x_{it} and x_{jt} , for any $i \neq j$, and $t = 1, 2, \dots, T$.

Assumption 2 Let $y_{it} = (x_{it} - \mu_i)/\sqrt{\sigma_{ii}}$ with mean $\mu_i = E(x_{it})$, $|\mu_i| < K$, variance $\sigma_{ii} = \text{Var}(x_{it})$, $0 < \sigma_{ii} < K$, and correlation coefficient $\rho_{ij} = \sigma_{ij}/\sqrt{\sigma_{ii}\sigma_{jj}}$, where $\sigma_{ij} = E(y_{it}y_{jt})$, and $|\rho_{ij}| < 1$. Suppose that $\sup_{i,t} E|y_{it}|^{2s} < K$ for some positive integer $s \geq 3$, and let $\boldsymbol{\xi}_{ij,t} = (y_{it}, y_{jt}, y_{it}^2, y_{jt}^2, y_{it}y_{jt})'$ such that for any $i \neq j$ the time series observations $\boldsymbol{\xi}_{ij,t}$, $t = 1, 2, \dots, T$, are random draws from a common distribution which is absolutely continuous with non-zero density on subsets of \mathbb{R}^5 .²

We begin our theoretical derivations with the following proposition.

Proposition 1 Let $y_{it} = (x_{it} - \mu_i)/\sqrt{\sigma_{ii}}$, and suppose that Assumption 2 holds. Consider the sample correlation coefficient given by (4), and note that

$$\hat{\rho}_{ij,T} = \frac{\sum_{t=1}^T (y_{it} - \bar{y}_i)(y_{jt} - \bar{y}_j)}{\left[\sum_{t=1}^T (y_{it} - \bar{y}_i)^2\right]^{1/2} \left[\sum_{t=1}^T (y_{jt} - \bar{y}_j)^2\right]^{1/2}}. \quad (8)$$

Then

$$\rho_{ij,T} = E(\hat{\rho}_{ij,T}) = \rho_{ij} + \frac{K_m(\boldsymbol{\theta}_{ij})}{T} + O(T^{-2}), \quad (9)$$

$$\omega_{ij,T}^2 = \text{Var}(\hat{\rho}_{ij,T}) = \frac{K_v(\boldsymbol{\theta}_{ij})}{T} + O(T^{-2}), \quad (10)$$

²The restrictions on the common distribution imply that Cramér's condition holds. See p.45 of Hall (1992) for further details.

uniformly in the i and j ($i \neq j$) pairs, where³

$$K_m(\boldsymbol{\theta}_{ij}) = -\frac{1}{2}\rho_{ij}(1 - \rho_{ij}^2) + \frac{3}{8}\rho_{ij} [\kappa_{ij}(4, 0) + \kappa_{ij}(0, 4)] - \frac{1}{2} [\kappa_{ij}(3, 1) + \kappa_{ij}(1, 3)] + \frac{1}{4}\rho_{ij}\kappa_{ij}(2, 2), \quad (11)$$

$$K_v(\boldsymbol{\theta}_{ij}) = (1 - \rho_{ij}^2)^2 + \frac{1}{4}\rho_{ij}^2 [\kappa_{ij}(4, 0) + \kappa_{ij}(0, 4)] - \rho_{ij} [\kappa_{ij}(3, 1) + \kappa_{ij}(1, 3)] + \frac{1}{2}(2 + \rho_{ij}^2)\kappa_{ij}(2, 2), \quad (12)$$

$$\kappa_{ij}(4, 0) = E(y_{it}^4) - 3, \quad \kappa_{ij}(0, 4) = E(y_{jt}^4) - 3,$$

$$\kappa_{ij}(3, 1) = E(y_{it}^3 y_{jt}) - 3\rho_{ij}, \quad \kappa_{ij}(1, 3) = E(y_{it} y_{jt}^3) - 3\rho_{ij},$$

$$\kappa_{ij}(2, 2) = E(y_{it}^2 y_{jt}^2) - 2\rho_{ij}^2 - 1,$$

$\boldsymbol{\theta}_{ij} = (\rho_{ij}, \kappa_{ij}(0, 4) + \kappa_{ij}(4, 0), \kappa_{ij}(3, 1) + \kappa_{ij}(1, 3), \kappa_{ij}(2, 2))'$, $\sup_{ij} |K_m(\boldsymbol{\theta}_{ij})| < K$ and $\sup_{ij} K_v(\boldsymbol{\theta}_{ij}) < K$. Under the additional assumption that y_{it} are Gaussian the above expressions simplify to $K_m(\boldsymbol{\theta}_{ij}) = -\frac{1}{2}\rho_{ij}(1 - \rho_{ij}^2)$ and $K_v(\boldsymbol{\theta}_{ij}) = (1 - \rho_{ij}^2)^2$, and it follows that $\sup_{ij} K_v(\boldsymbol{\theta}_{ij}) = 1$.

All proofs are given in the Appendix with supporting Lemmas and technical details provided in an online supplement.

Remark 2 From Gayen (1951) p.232 (eq (54)bis) it follows that $K_v(\boldsymbol{\theta}_{ij}) > 0$ for all correlation coefficients $\rho_{ij} = \sigma_{ij} / \sqrt{\sigma_{ii}\sigma_{jj}}$, such that $|\rho_{ij}| < 1$. Further, in the case where $\rho_{ij} = 0$, by (12),

$$\varphi_{ij} := K_v(\boldsymbol{\theta}_{ij} | \rho_{ij} = 0) = E(y_{it}^2 y_{jt}^2 | \rho_{ij} = 0) > 0, \quad (13)$$

and by (11),

$$\psi_{ij} := K_m(\boldsymbol{\theta}_{ij} | \rho_{ij} = 0) = -0.5 [E(y_{it}^3 y_{jt} | \rho_{ij} = 0) + E(y_{it} y_{jt}^3 | \rho_{ij} = 0)]. \quad (14)$$

Note in addition that when y_{it} and y_{jt} are independently distributed, then $\varphi_{ij} = E(y_{it}^2) E(y_{jt}^2) = 1$, and $\psi_{ij} = 0$. This is also the case when y_{it} are Gaussian.

³See also equations (38) and (39) of Gayen (1951).

The next proposition assists in establishing probability bounds on $\hat{\rho}_{ij,T}$.

Proposition 2 *Consider the standardised correlation coefficient*

$$z_{ij,T} = (\hat{\rho}_{ij,T} - \rho_{ij,T}) / \omega_{ij,T}, \quad (15)$$

where $\hat{\rho}_{ij,T}$ is defined by (4), $\rho_{ij,T}$ and $\omega_{ij,T}^2$ are defined by (9) and (10), respectively, and suppose that Assumptions 1 and 2 hold, and for all i and j ($i \neq j$) $\sup_{ij} E(|z_{ij,T}|^s) < K$, for some finite integer $s \geq 3$. Then the cumulative distribution function of $z_{ij,T}$, denoted by $F_{ij,T}(x) = \Pr(z_{ij,T} \leq x)$, has the following Edgeworth expansion

$$\Pr(z_{ij,T} \leq x) = \Phi(x) + \sum_{r=1}^{s-2} T^{-r/2} g_r(x) \phi(x) + O[T^{-(s-1)/2}] \quad (16)$$

uniformly in $x \in \mathbb{R}$, where $\Phi(x)$ and $\phi(x)$ are the distribution and density functions of the standard Normal $(0, 1)$, respectively, and $g_r(x)$, for $r = 1, 2, \dots, s-2$, are finite-order polynomials in x of degree $3r-1$ whose coefficients do not depend on x . Furthermore, for all finite $s \geq 3$, and $a_T > 0$, we have

$$\Pr(z_{ij,T} \leq -a_T) \leq K e^{-\frac{1}{2}a_T^2} + O\left(T^{-\frac{(s-2)}{2}} a_T^{3(s-2)-1} e^{-\frac{1}{2}a_T^2}\right) + O[T^{-(s-1)/2}], \quad (17)$$

and

$$\Pr(z_{ij,T} > a_T) \leq K e^{-\frac{1}{2}a_T^2} + O\left(T^{-\frac{(s-2)}{2}} a_T^{3(s-2)-1} e^{-\frac{1}{2}a_T^2}\right) + O[T^{-(s-1)/2}]. \quad (18)$$

Remark 3 *This proposition establishes a bound on the probability of $|\hat{\rho}_{ij,T} - \rho_{ij}| > T^{-1/2} c_p(N)$ without requiring sub-Gaussianity, at the expense of the additional order term, $O[T^{-(s-1)/2}]$, which relates the bound to the order of the moments of $z_{ij,T}$.*

Remark 4 *It is also possible to use the Berry-Essen inequality and Cramer-type moderate deviation to obtain the probability bounds in the above proposition which could result in better bounds. On this see, in particular, Delaigle et al. (2011), and the recent contributions by Zhou et al. (2018) and Fan et al. (2018) who make use of a robust covariance estimator*

to tackle heavy-tailed data that yields exponential-type deviation bounds under mild moment conditions. While the focus of these contributions is primarily on large scale dependence-adjusted multiple testing of the mean, application of their approach to our problem could lead to weaker moment conditions.⁴

Using the probability bounds (17) and (18) we first establish the rate of convergence of the MT estimator under the spectral norm which implies convergence in eigenvalues and eigenvectors (see El Karoui (2008), and Bickel and Levina (2008b)).

Theorem 1 (*Convergence under spectral norm*) Consider the sample correlation coefficient of x_{it} and x_{jt} , defined by $\hat{\rho}_{ij,T}$ (see (4)), and denote the associated population correlation matrix by $\mathbf{R} = (\rho_{ij})$. Let $T = c_d N^d$, with $c_d > 0$, and suppose that Assumptions 1 and 2 hold. Further, let

$$c_p(N) = \Phi^{-1} \left(1 - \frac{p}{2f(N)} \right),$$

where $0 < p < 1$, $f(N) = c_\delta N^\delta$, with $c_\delta, \delta > 0$. Suppose also that there exist N_0 such that for all $N > N_0$,

$$1 - \frac{p}{2f(N)} > 0, \tag{19}$$

$$\rho_{\min} > c_p(N)/\sqrt{T}, \tag{20}$$

where $\rho_{\min} = \min_{ij} (|\rho_{ij}|, \rho_{ij} \neq 0)$, and

$$c_p(N)/\sqrt{T} = o(1). \tag{21}$$

Consider values of δ that satisfy condition

$$\delta > \frac{2K_v}{(1-\gamma)^2}, \tag{22}$$

for some small positive constant γ , where $K_v = \sup_{ij} K_v(\boldsymbol{\theta}_{ij})$ and $K_v(\boldsymbol{\theta}_{ij})$ is defined by (12).

⁴We are grateful to Jianqing Fan (Co-Editor) for drawing our attention to this alternative approach which could form the basis of future investigations.

Then for all values of $d > 4/(s - 1)$, where s is defined by Assumption 2, we have

$$\left\| \tilde{\mathbf{R}}_{MT} - \mathbf{R} \right\| = O_p \left(\frac{m_N c_p(N)}{\sqrt{T}} \right), \quad (23)$$

where m_N is defined by (3), and the multiple testing threshold estimator, $\tilde{\mathbf{R}}_{MT}$, is defined by $\tilde{\mathbf{R}}_{MT} = (\tilde{\rho}_{ij,T})$, where $\tilde{\rho}_{ij,T} = \hat{\rho}_{ij,T} I \left[|\hat{\rho}_{ij,T}| > T^{-1/2} c_p(N) \right]$.

Remark 5 The term $c_p(N)$ in (23) directly corresponds to the term $\sqrt{\ln(N)}$ obtained in the literature for the probability order, $O_p \left(\frac{m_N \sqrt{\ln(N)}}{\sqrt{T}} \right)$, of the spectral norm of threshold estimators of \mathbf{R} . This result follows since $\lim_{N \rightarrow \infty} c_p^2(N) / \ln(N) = 2\delta$, for $\delta > 0$.⁵

Remark 6 The parameter d , which controls the rate at which T rises with N , is required to be sufficiently large such that $d > 4/(s - 1)$, and $T^{-1/2} c_p(N) = o(1)$ hold. But from result (a) of Lemma 2 in the online supplement, we have $N^{-d/2} c_p(N) = O \left(\sqrt{N^{-d} \ln(N)} \right)$, and condition $T^{-1/2} c_p(N) = o(1)$ will be met if $N^{-d} \ln(N) = o(1)$. Further, recall that the validity of the Edgeworth expansion that underlies our analysis requires s to be finite, and hence condition $d > 0$ will follow from the moment condition $d > 4/(s - 1)$, for $s \geq 3$ required by Assumption 2.

Remark 7 Condition (19) is met for $\delta > 0$ and N sufficiently large. Condition (20) can be written as

$$\rho_{\min}^2 > \frac{c_p^2(N)}{T} = \frac{c_p^2(N)}{c_d N^d} = c_d^{-1} \left[\frac{c_p^2(N)}{\ln(N)} \right] \left[\frac{\ln(N)}{N^d} \right].$$

Once again since $\lim_{N \rightarrow \infty} c_p^2(N) / \ln(N) = 2\delta$, then condition (20) will be satisfied for any $\delta > 0$, even if ρ_{\min} tends to zero with N , so long as the rate at which ρ_{\min} tends to zero is slower than $\sqrt{\ln(N)/N^d}$.

Remark 8 Note that under Gaussianity where $K_v = \sup_{ij} K_v(\theta_{ij}) = 1$, condition (22) becomes $\delta > 2$. In general, the spectral norm result requires δ to be set above $2 \sup_{ij} K_v(\theta_{ij})$, which turns out to be larger than the value of δ required for the Frobenius norm obtained in the theorem below.

⁵See part (b) of Lemma 2 in the online supplement.

Theorem 2 (Convergence under Frobenius norm) Suppose that conditions of Theorem 1 hold, but (22) is replaced by the weaker condition on δ

$$\delta > (2 - d) \varphi_{\max}, \quad (24)$$

where $\varphi_{\max} = \sup_{ij} E(y_{it}^2 y_{jt}^2 | \rho_{ij} = 0) > 0$, $y_{it} = (x_{it} - \mu_i) / \sqrt{\sigma_{ii}}$ (see Assumption 2), δ and d are the exponents in $f(N) = c_\delta N^\delta$, and $T = c_d N^d$, with $c_\delta, c_d > 0$. Then for all values of

$$d > \max \left(\frac{2 + \vartheta}{s - 1}, \frac{4}{s + 1} \right), \quad (25)$$

where s is defined by Assumption 2, and ϑ ($0 \leq \vartheta < 1/2$) is the degree of sparsity of the correlation matrix, \mathbf{R} , defined by condition (3) in Assumption 1, we have

$$E \left\| \tilde{\mathbf{R}}_{MT} - \mathbf{R} \right\|_F = O \left(\sqrt{\frac{m_N N}{T}} \right), \quad (26)$$

and

$$\left\| \tilde{\mathbf{R}}_{MT} - \mathbf{R} \right\|_F = O_p \left(\sqrt{\frac{m_N N}{T}} \right), \quad (27)$$

where $m_N = O(N^\vartheta)$, and the multiple testing threshold estimator, $\tilde{\mathbf{R}}_{MT}$, is defined by $\tilde{\mathbf{R}}_{MT} = (\tilde{\rho}_{ij,T})$, where $\tilde{\rho}_{ij,T} = \hat{\rho}_{ij,T} I [|\hat{\rho}_{ij,T}| > T^{-1/2} c_p(N)]$.

Remark 9 For the Frobenius norm result to hold condition (24) implies that δ should be set at a sufficiently high level, determined by d (the relative expansion rates of N and T), and φ_{\max} (the maximum degree of dependence between y_{it} and y_{jt} when $\rho_{ij} = 0$). The Frobenius norm result holds even if N rises faster than T , so long as $c_p(N) / \sqrt{T} = o(1)$ and a sufficient number of moments exists such that condition (25) is met. In the case where N and T are of the same order of magnitude (namely, $d = 1$), and where y_{it} and y_{jt} are independently distributed when $\rho_{ij} = 0$ (namely, $\varphi_{\max} = 1$), then the Frobenius norm results, (26) and (27), require $\delta > 1$.

Remark 10 *The number of moments, s , of y_{it} required for the convergence results (23), (26) or (27) to hold is related to the relative rate of expansion of N and T , d . For $d = 1$, $s = 5$ moments of $\hat{\rho}_{ij}$ (which requires x_{it} to have 10 moments) are sufficient to achieve the spectral or Frobenius norm results. Additional moments are required if N is to rise faster than T .*

Remark 11 *The convergence rate of $O_p\left(\sqrt{\frac{m_N N}{T}}\right)$ obtained for the MT estimator under the Frobenius norm compares favourably to the corresponding rate of $O_p\left(\sqrt{\frac{m_N N \ln(N)}{T}}\right)$ obtained for the threshold estimator. See, for example, Theorem 2 of Bickel and Levina (2008a), BL. The slower rate of convergence achieved by BL under the Frobenius norm arises from the fact that their result is derived by explicitly using their spectral norm convergence rate. On the other hand, we consider the derivation of the Frobenius norm convergence rate directly and independently of our spectral norm results. Furthermore, the sparsity condition of Assumption 1 sets an upper bound, m_N , on the number of non-zero units in the rows (columns) of the population covariance matrix Σ , but it is silent as to the number of rows (columns) of Σ with m_N non-zero elements. Whilst this ambiguity does not impact the convergence rate obtained for the spectral norm, it does affect the Frobenius norm. In many economic applications it might be known that only a finite number of rows of Σ , say k , have at most m_N non-zero elements with the rest of the rows only containing a fixed number of non-zero elements, say m_0 , which is bounded in N . Under this notion of sparsity the convergence rate of the Frobenius norm will be given by*

$$\left\| \tilde{\mathbf{R}}_{MT} - \mathbf{R} \right\|_F = O_p\left(\sqrt{\frac{k m_N}{T}}\right) + O_p\left(\sqrt{\frac{(N - k) m_0}{T}}\right),$$

which has a more favourable convergence rate as compared to (27).

Remark 12 *It is interesting to note that application of the Bonferroni procedure to the problem of testing $\rho_{ij} = 0$ for all $i \neq j$, is equivalent to setting $f(N) = N(N - 1)/2$. Our theoretical results suggest that this can be too conservative if $\rho_{ij} = 0$ implies y_{it} and*

y_{jt} are independent, but could be appropriate otherwise depending on the relative rates at which N and T rise. In our Monte Carlo study we consider $\delta = \{1, 2\}$, that corresponds to $\varphi_{\max} = \{1, 1.5\}$.

Consider now the issue of consistent support recovery of \mathbf{R} (or $\mathbf{\Sigma}$) for $T = T(N) = c_d N^d$ and $N \rightarrow \infty$, which is defined in terms of the true positive rate (TPR_N), false positive rate (FPR_N), and false discovery rate (FDR_N) statistics. Consistent support recovery requires $TPR_N \rightarrow 1$, $FPR_N \rightarrow 0$ and $FDR_N \rightarrow 0$, with probability 1 (almost surely) as $N \rightarrow \infty$, and does not follow immediately from the results obtained above on the convergence rates of different estimators of \mathbf{R} . This is addressed in the following theorem.

Theorem 3 (*Support Recovery*) *Suppose that Assumptions 1 and 2 hold, and let*

$$c_p(N) = \Phi^{-1} \left(1 - \frac{p}{2f(N)} \right),$$

where $0 < p < 1$, $f(N) = c_\delta N^\delta$, with $c_\delta, \delta > 0$, and $T = c_d N^d$, with $c_d > 0$. Further, suppose that there exist N_0 such that for all $N > N_0$,

$$1 - \frac{p}{2f(N)} > 0,$$

$$\rho_{\min} > c_p(N)/\sqrt{T}, \tag{28}$$

where $\rho_{\min} = \min_{ij} (|\rho_{ij}|, \rho_{ij} \neq 0)$, and

$$c_p(N)/\sqrt{T} = o(1). \tag{29}$$

Consider the true positive rate (TPR_N), the false positive rate (FPR_N), and the false dis-

covery rate (FDR_N) statistics defined by

$$TPR_N = \frac{\sum_{i \neq j} \sum I(\tilde{\rho}_{ij,T} \neq 0, \text{ and } \rho_{ij} \neq 0)}{\sum_{i \neq j} \sum I(\rho_{ij} \neq 0)} \quad (30)$$

$$FPR_N = \frac{\sum_{i \neq j} \sum I(\tilde{\rho}_{ij,T} \neq 0, \text{ and } \rho_{ij} = 0)}{\sum_{i \neq j} \sum I(\rho_{ij} = 0)}, \quad (31)$$

$$FDR_N = \frac{\sum_{i \neq j} \sum I(\tilde{\rho}_{ij,T} \neq 0, \text{ and } \rho_{ij} = 0)}{\sum_{i \neq j} \sum I(\rho_{ij} \neq 0)}, \quad (32)$$

computed using the multiple testing estimator

$$\tilde{\rho}_{ij,T} = \hat{\rho}_{ij,T} I \left[\left| \hat{\rho}_{ij,T} \right| > T^{-1/2} c_p(N) \right],$$

where $\hat{\rho}_{ij,T}$ is the pair-wise correlation coefficient defined by (4). Then as $N \rightarrow \infty$ we have:

$$TPR_N \xrightarrow{a.s.} 1, \text{ for } \delta > 0, \text{ and } d > 2/(s-1)$$

$$FPR_N \xrightarrow{a.s.} 0, \text{ for } \delta > \varphi_{\max}, \text{ and } d > 2/(s-1)$$

$$FDR_N \xrightarrow{a.s.} 0, \text{ for } \delta > (2 - \vartheta)\varphi_{\max}, \text{ and } d > 2(2 - \vartheta)/(s-1)$$

where ϑ ($0 \leq \vartheta < 1/2$) is the degree of sparsity of the correlation matrix, \mathbf{R} , defined by condition (3) in Assumption 1, $\varphi_{\max} = \sup_{ij} E(y_{it}^2 y_{jt}^2 | \rho_{ij} = 0) > 0$, with $y_{it} = (x_{it} - \mu_i) / \sqrt{\sigma_{ii}}$ (see Assumption 2). Further, as $N \rightarrow \infty$, $TPR_N \rightarrow 1$ and $FPR_N \rightarrow 0$ in probability for any $\delta > 0$ and $d > 2/(s-1)$, and $FDR_N \rightarrow 0$ in probability if $\delta > (1 - \vartheta)\varphi_{\max}$, and $d > 2(1 - \vartheta)/(s-1)$.

Remark 13 We note that

$$\frac{c_p^2(N)}{T} \leq \frac{2[\ln(N) - \ln(p)]}{c_d N^d},$$

and hence condition $T^{-1/2} c_p(N) = o(1)$ will be met if $N^{-d} \ln(N) = o(1)$. Also, since under Assumption 2 $s \geq 3$, it follows from the moment conditions on d that $d > 0$. For a discussion

of the remaining conditions on δ , d , and $\rho_{\min} > T^{-1/2}c_p(N) > 0$, see the above Remarks. In general, the conditions needed for the support recovery results to hold when N is much larger than T are less restrictive as compared to the conditions needed for the validity of the results on the spectral and Frobenius norms.

3 Monte Carlo simulations

We investigate the numerical properties of the proposed multiple testing (*MT*) estimator using Monte Carlo simulations. We compare our estimator with a number of thresholding and shrinkage estimators proposed in the literature, namely the thresholding estimators of Bickel and Levina (2008a) - BL - and Cai and Liu (2011) - CL, and the shrinkage estimator of LW. The thresholding methods of BL and CL require the computation of a theoretical constant, C , that arises in the rate of their convergence. For this purpose, cross-validation is typically employed which we use when implementing these estimators. For the CL approach we also consider the theoretical value of $C = 2$ derived by the authors in the case of Gaussianity. A review of these estimators along with details of the associated cross-validation procedure can be found in the Supplementary Appendix B.

We begin by generating the standardised variates, y_{it} , as

$$\mathbf{y}_t = \mathbf{P}\mathbf{u}_t, t = 1, 2, \dots, T,$$

where $\mathbf{y}_t = (y_{1t}, y_{2t}, \dots, y_{Nt})'$, $\mathbf{u}_t = (u_{1t}, u_{2t}, \dots, u_{Nt})'$, and \mathbf{P} is the Cholesky factor associated with the choice of the correlation matrix $\mathbf{R} = \mathbf{P}\mathbf{P}'$. We consider two alternatives for the errors, u_{it} : (i) the benchmark Gaussian case where $u_{it} \sim IIDN(0, 1)$ for all i and t , and (ii) the case where u_{it} follows a multivariate t-distribution with v degrees of freedom generated as

$$u_{it} = \left(\frac{v-2}{\chi_{v,t}^2} \right)^{1/2} \varepsilon_{it}, \text{ for } i = 1, 2, \dots, N,$$

where $\varepsilon_{it} \sim IIDN(0, 1)$, and $\chi_{v,t}^2$ is a chi-squared random variate with $v > 4$ degrees of

freedom, distributed independently of ε_{it} for all i and t . In order to investigate the robustness of our results to the moment conditions, we experiment with a relatively low degrees of freedom for the t-distribution and set $v=8$, which ensures that $E(y_{it}^6)$ exists and $\varphi_{\max} \leq 2$. Note that under $\rho_{ij} = 0$, $\varphi_{ij} = E(y_{it}^2 y_{jt}^2 | \rho_{ij} = 0) = (v-2)/(v-4)$, and with $v = 8$ we have $\varphi_{ij} = \varphi_{\max} = 1.5$. Given our theoretical findings, it is most likely that we obtain better results if we experiment with higher degrees of freedom. One could further allow for fat-tailed ε_{it} shocks, say, though fat-tail shocks alone (e.g. generating u_{it} as such) do not necessarily result in $\varphi_{ij} > 1$ as shown in Lemma 6 in the online supplementary Appendix A. The same is true for normal shocks under case (i) where $E(y_{it}^2 y_{jt}^2) = 1$ whether $\mathbf{P} = \mathbf{I}_N$ or not. In such cases setting $\delta = 1$ is likely to be sufficient for the Frobenius norms given the (N, T) combinations considered. But for the spectral norm a larger value of δ might be necessary. In order to verify and calibrate the values of δ corresponding to the alternative processes generating y_{it} , we also consider an estimated version of δ . For this purpose we use a cross-validation procedure that corresponds to those used for the BL and CL methods respectively. Details can be found in Section 3.7.

Next, the non-standardised variates $\mathbf{x}_t = (x_{1t}, x_{2t}, \dots, x_{Nt})'$ are generated as

$$\mathbf{x}_t = \mathbf{a} + \gamma f_t + \mathbf{D}^{1/2} \mathbf{y}_t, \quad (33)$$

where $\mathbf{D} = \text{diag}(\sigma_{11}, \sigma_{22}, \dots, \sigma_{NN})$, $\mathbf{a} = (a_1, a_2, \dots, a_N)'$ and $\boldsymbol{\gamma} = (\gamma_1, \gamma_2, \dots, \gamma_N)'$.

We report results for $N = \{30, 100, 200\}$ and $T = 100$, for the baseline case where $\gamma = 0$ and $a = 0$ in (33). The properties of the MT procedure when factors are included in the data generating process are also investigated by drawing γ_i and a_i as $IIDN(1, 1)$ for $i = 1, 2, \dots, N$, and generating f_t , the common factor, as a stationary AR(1) process, but to save space these results are made available upon request. Under both settings we focus on the residuals from an OLS regression of \mathbf{x}_t on an intercept and a factor (if needed).

Given our interest in both the problems of regularisation of $\hat{\boldsymbol{\Sigma}}$ and support recovery of $\boldsymbol{\Sigma}$, we consider two exactly sparse covariance (correlation) matrices:

Monte Carlo design A: Following Cai and Liu (2011) we consider the banded matrix

$$\boldsymbol{\Sigma} = (\sigma_{ij}) = \text{diag}(\mathbf{A}_1, \mathbf{A}_2),$$

where $\mathbf{A}_1 = \mathbf{A} + \epsilon \mathbf{I}_{N/2}$, $\mathbf{A} = (a_{ij})_{1 \leq i, j \leq N/2}$, $a_{ij} = (1 - \frac{|i-j|}{10})_+$ with $\epsilon = \max(-\lambda_{\min}(\mathbf{A}), 0) + 0.01$ to ensure that \mathbf{A} is positive definite, and $\mathbf{A}_2 = 4\mathbf{I}_{N/2}$. $\boldsymbol{\Sigma}$ is a two-block diagonal matrix, \mathbf{A}_1 is a banded and sparse covariance matrix, and \mathbf{A}_2 is a diagonal matrix with 4 along the diagonal. Matrix \mathbf{P} is obtained numerically by applying the Cholesky decomposition to the correlation matrix, $\mathbf{R} = \mathbf{D}^{-1/2} \boldsymbol{\Sigma} \mathbf{D}^{-1/2} = \mathbf{P} \mathbf{P}'$, where the diagonal elements of \mathbf{D} are given by $\sigma_{ii} = 1 + \epsilon$, for $i = 1, 2, \dots, N/2$ and $\sigma_{ii} = 4$, for $i = N/2 + 1, N/2 + 1, \dots, N$.

Monte Carlo design B: We consider a covariance structure that explicitly controls for the number of non-zero elements of the population correlation matrix. First we draw the $N \times 1$ vector $\mathbf{b} = (b_1, b_2, \dots, b_N)'$ with elements generated as *Uniform*(0.7, 0.9) for the first and last N_b ($< N$) elements of \mathbf{b} , where $N_b = \lceil N^\beta \rceil$, and set the remaining middle elements of \mathbf{b} to zero. The resulting population correlation matrix \mathbf{R} is defined by

$$\mathbf{R} = \mathbf{I}_N + \mathbf{b} \mathbf{b}' - \text{diag}(\mathbf{b} \mathbf{b}'), \quad (34)$$

for which $\sqrt{T} \rho_{\min} - c_p(N) > 0$ and $\rho_{\min} = \min_{ij} (|\rho_{ij}|, \rho_{ij} \neq 0) > 0$, in line with Theorem 3. The degree of sparseness of \mathbf{R} is determined by the value of the parameter β . We are interested in weak cross-sectional dependence, so we focus on the case where $\beta < 1/2$ following Pesaran (2015), and set $\beta = 0.25$. Matrix \mathbf{P} is then obtained by applying the Cholesky decomposition to \mathbf{R} defined by (34). Further, we set $\boldsymbol{\Sigma} = \mathbf{D}^{1/2} \mathbf{R} \mathbf{D}^{1/2}$, where the diagonal elements of \mathbf{D} are given by $\sigma_{ii} \sim \text{IID}(1/2 + \chi^2(2)/4)$, $i = 1, 2, \dots, N$.

3.1 Finite sample positive definiteness

As with other thresholding approaches, multiple testing preserves the symmetry of $\hat{\mathbf{R}}$ and is invariant to the ordering of the variables but it does not ensure positive definiteness of the

estimated covariance matrix when $N > T$.

A number of methods have been developed in the literature that produce sparse inverse covariance matrix estimates which make use of a penalised likelihood (D’Aspremont *et al.* (2008), Rothman *et al.* (2008), Rothman *et al.* (2009), Yuan and Lin (2007), and Peng *et al.* (2009)) or convex optimisation techniques that apply suitable penalties such as a logarithmic barrier term (Rothman (2012)), a positive definiteness constraint (Xue *et al.* (2012)), an eigenvalue condition (Liu *et al.* (2014), Fryzlewicz (2013), Fan *et al.* (2013) - FLM). Most of these approaches are rather complex and computationally extensive.

A simpler alternative, which conceptually relates to soft thresholding (such as the smoothly clipped absolute deviation by Fan and Li (2001) and the adaptive lasso by Zou (2006)), is to consider a convex linear combination of $\tilde{\mathbf{R}}_{MT}$ and a well-defined target matrix which is known to result in a positive definite matrix. In what follows, we opt to set as benchmark target the $N \times N$ identity matrix, \mathbf{I}_N , in line with one of the methods suggested by El Karoui (2008). The advantage of doing so lies in the fact that the same support recovery achieved by $\tilde{\mathbf{R}}_{MT}$ is maintained and the diagonal elements of the resulting correlation matrix do not deviate from unity. Given the similarity of this adjustment to the shrinking method, we dub this step shrinkage on our multiple testing estimator (S - MT),

$$\tilde{\mathbf{R}}_{S-MT}(\xi) = \xi \mathbf{I}_N + (1 - \xi) \tilde{\mathbf{R}}_{MT}, \quad (35)$$

with shrinkage parameter $\xi \in (\xi_0, 1]$, and ξ_0 being the minimum value of ξ that produces a non-singular $\tilde{\mathbf{R}}_{S-MT}(\xi_0)$ matrix. Alternative ways of computing the optimal weights on the two matrices can be entertained. We choose to calibrate, ξ , since opting to use ξ_0 in (35), as suggested in El Karoui (2008), does not necessarily provide a well-conditioned estimate of $\tilde{\mathbf{R}}_{S-MT}$. Accordingly, we set ξ by solving the following optimisation problem

$$\xi^* = \arg \min_{\xi_0 + \epsilon \leq \xi \leq 1} \left\| \mathbf{R}_0^{-1} - \tilde{\mathbf{R}}_{S-MT}^{-1}(\xi) \right\|_F^2, \quad (36)$$

where ϵ is a small positive constant, and \mathbf{R}_0 is a reference invertible correlation matrix. Finally, we construct the corresponding covariance matrix as

$$\tilde{\Sigma}_{S-MT}(\xi^*) = \hat{\mathbf{D}}^{1/2} \tilde{\mathbf{R}}_{S-MT}(\xi^*) \hat{\mathbf{D}}^{1/2}.$$

Further details on the S - MT procedure, the optimisation of (36) and choice of reference matrix \mathbf{R}_0 are available in the Supplementary Appendix C.

3.2 Alternative estimators and evaluation metrics

Using the earlier set up and the relevant adjustments to achieve positive definiteness of the estimators of Σ where required, we obtain the following estimates of Σ :

MT_1 : thresholding based on the MT approach applied to the sample correlation matrix ($\tilde{\Sigma}_{MT}$) using $\delta = 1$ ($\tilde{\Sigma}_{MT,1}$)

MT_2 : thresholding based on the MT approach applied to the sample correlation matrix ($\tilde{\Sigma}_{MT}$) using $\delta = 2$ ($\tilde{\Sigma}_{MT,2}$)

$MT_{\hat{\delta}}$: thresholding based on the MT approach applied to the sample correlation matrix ($\tilde{\Sigma}_{MT}$) using cross-validated δ ($\tilde{\Sigma}_{MT,\hat{\delta}}$)

$BL_{\hat{C}}$: BL thresholding on the sample covariance matrix using cross-validated C ($\tilde{\Sigma}_{BL,\hat{C}}$)

CL_2 : CL thresholding on the sample covariance matrix using the theoretical value of $C = 2$ ($\tilde{\Sigma}_{CL,2}$)

$CL_{\hat{C}}$: CL thresholding on the sample covariance matrix using cross-validated C ($\tilde{\Sigma}_{CL,\hat{C}}$)

S - MT_1 : supplementary shrinkage applied to MT_1 ($\tilde{\Sigma}_{S-MT,1}$)

S - MT_2 : supplementary shrinkage applied to MT_2 ($\tilde{\Sigma}_{S-MT,2}$)

S - $MT_{\hat{\delta}}$: supplementary shrinkage applied to $MT_{\hat{\delta}}$ ($\tilde{\Sigma}_{S-MT,\hat{\delta}}$)

$BL_{\hat{C}^*}$: BL thresholding using the Fan *et al.* (2013) - FLM - cross-validation adjustment procedure for estimating C to ensure positive definiteness ($\tilde{\Sigma}_{BL,\hat{C}^*}$)

$CL_{\hat{C}^*}$: CL thresholding using the FLM cross-validation adjustment procedure for estimating C to ensure positive definiteness ($\tilde{\Sigma}_{CL,\hat{C}^*}$)

$LW_{\hat{\Sigma}}$: LW shrinkage on the sample covariance matrix ($\hat{\Sigma}_{LW_{\hat{\Sigma}}}$).

In accordance with the theoretical results and in view of Remark 12, we consider three versions of the MT estimator depending on the choice of $\delta = \{1, 2, \hat{\delta}\}$. The $BL_{\hat{C}}$, CL_2 and $CL_{\hat{C}}$ estimators apply the thresholding procedure without ensuring that the resultant covariance estimators are invertible. The next six estimators yield invertible covariance estimators. The S - MT estimators are obtained using the supplementary shrinkage approach described in Section 3.1. $BL_{\hat{C}^*}$ and $CL_{\hat{C}^*}$ estimators are obtained by applying the additional FLM adjustments. The shrinkage estimator, $LW_{\hat{\Sigma}}$, is invertible by construction. In the case of the MT estimators where regularisation is performed on the correlation matrix, the associated covariance matrix is estimated as $\hat{\mathbf{D}}^{1/2}\tilde{\mathbf{R}}_{MT}\hat{\mathbf{D}}^{1/2}$.

For both Monte Carlo designs A and B, we compute the spectral and Frobenius norms of the deviations of each of the regularised covariance matrices from their respective population Σ :

$$\left\| \Sigma - \hat{\Sigma} \right\| \quad \text{and} \quad \left\| \Sigma - \hat{\Sigma} \right\|_F, \quad (37)$$

where $\hat{\Sigma}$ is set to one of the following estimators $\{\tilde{\Sigma}_{MT,1}, \tilde{\Sigma}_{MT,2}, \tilde{\Sigma}_{MT,\hat{\delta}}, \tilde{\Sigma}_{BL,\hat{C}}, \tilde{\Sigma}_{CL,2}, \tilde{\Sigma}_{CL,\hat{C}}, \tilde{\Sigma}_{S-MT,1}, \tilde{\Sigma}_{S-MT,2}, \tilde{\Sigma}_{S-MT,\hat{\delta}}, \tilde{\Sigma}_{BL,\hat{C}^*}, \tilde{\Sigma}_{CL,\hat{C}^*}, \hat{\Sigma}_{LW_{\hat{\Sigma}}}\}$. The threshold values, $\hat{\delta}$, \hat{C} and \hat{C}^* , are obtained by cross-validation (see Section 3.7 and supplementary appendix B.3 for details). Both norms are also computed for the difference between Σ^{-1} , the population inverse of Σ , and the estimators $\{\tilde{\Sigma}_{S-MT,1}^{-1}, \tilde{\Sigma}_{S-MT,2}^{-1}, \tilde{\Sigma}_{S-MT,\hat{\delta}}^{-1}, \tilde{\Sigma}_{BL,\hat{C}^*}^{-1}, \tilde{\Sigma}_{CL,\hat{C}^*}^{-1}, \hat{\Sigma}_{LW_{\hat{\Sigma}}}^{-1}\}$. Further, we investigate the ability of the thresholding estimators to recover the support of the true covariance matrix via the true positive rate (TPR) and false positive rate (FPR), as defined by (30) and (31), respectively. The statistics TPR and FPR are not relevant to the shrinkage estimator $LW_{\hat{\Sigma}}$ and will not be reported for this estimator.

3.3 Robustness of MT to the choice of p-values

We begin by investigating the sensitivity of the MT estimator to the choice of the p-value, p , and the scaling factor determined by δ used in the formulation of $c_p(N)$ defined by (6).

For this purpose we consider the typical significance levels used in the literature, namely $p = \{0.01, 0.05, 0.10\}$, $\delta = \{1, 2\}$, and a cross-validated version of δ , denoted by $\hat{\delta}$. Tables 1a and 1b summarise the spectral and Frobenius norm losses (averaged over 2000 replications) for Monte Carlo designs A and B respectively, and for both distributional error assumptions (Gaussian and multivariate t). First, we note that neither of the norms is much affected by the choice of the p values when setting $\delta = 1$ or 2 in the scaling factor, irrespective of whether the observations are drawn from a Gaussian or a multivariate t distribution. Similar results are also obtained using the cross validated version of δ . Perhaps this is to be expected since for N sufficiently large the effective p-value which is given by $2p/N^\delta$ is very small and the test outcomes are more likely to be robust to the choice of p values as compared to the choice of δ . The results in Tables 1a and 1b also show that in the case of Gaussian observations, where $\varphi_{\max} = 1$, the scaling factor using $\delta = 1$ is likely to perform better as compared to $\delta = 2$, but the reverse is true if the observations are multivariate t distributed under which the scaling factor using $\delta = 2$ is to be preferred.

It is also interesting that the performance of the MT procedure when using $\hat{\delta}$ is in line with our theoretical findings. The estimates of δ are closer to unity in the case of experiments with $\varphi_{\max} = 1$, and are closer to $\delta = 2$ in the case of experiments with $\varphi_{\max} = 1.5$. The average estimates of $\hat{\delta}$ shown in Tables 1a and 1b are also indicative that a higher value of δ is required when observations are multivariate t distributed. Finally, we note that the norm losses rise with N given that T is kept at 100 almost across the board in all the experiments. Overall, the simulation results support using a sufficiently high value of δ (say around 2) or its estimate, $\hat{\delta}$, obtained by cross validation.

3.4 Norm comparisons of MT , BL , CL , and LW estimators

In comparing our proposed estimators with those in the literature we consider a fewer number of Monte Carlo replications and report the results with norm losses averaged over 100 replications, given the use of the cross-validation procedure in the implementation of MT ,

BL and CL thresholding. This Monte Carlo specification is in line with the simulation set up of BL and CL. Our reported results are also in agreement with their findings.

Tables 2 and 3 summarise the results for the Monte Carlo designs A and B, respectively. Based on the results of Section 3.3, we provide norm comparisons for the MT estimator using the scaling factor where $\delta = 2$ and $\hat{\delta}$, and the conventional significance level of $p = 0.05$. Initially, we consider the threshold estimators, the two versions of MT (MT_2 and $MT_{\hat{\delta}}$) and CL (CL_2 and $CL_{\hat{c}}$) estimators, and BL without further adjustments to ensure invertibility. First, we note that the MT and CL estimators (both versions for each case) dominate the BL estimator in every case, and for both designs. MT performs better than CL , when comparing the versions of the two estimators using their respective theoretical thresholding values and their estimated equivalents. The outperformance of MT is more evident as N increases and when non-Gaussian observations are considered. The same is also true if we compare MT and CL estimators to the LW shrinkage estimator, although it could be argued that it is more relevant to compare the invertible versions of the MT and CL estimators (namely $\tilde{\Sigma}_{CL, \hat{c}^*}$, $\tilde{\Sigma}_{S-MT, 2}$ and $\tilde{\Sigma}_{S-MT, \hat{\delta}}$) with $\hat{\Sigma}_{LW_{\hat{\Sigma}}}$. In such comparisons $\hat{\Sigma}_{LW_{\hat{\Sigma}}}$ performs relatively better, nevertheless, $\hat{\Sigma}_{LW_{\hat{\Sigma}}}$ is still dominated by $\tilde{\Sigma}_{S-MT, 2}$ and $\tilde{\Sigma}_{S-MT, \hat{\delta}}$, with a few exceptions in the case of design A and primarily when $N = 30$. However, no clear ordering emerges when we compare $\hat{\Sigma}_{LW_{\hat{\Sigma}}}$ with $\tilde{\Sigma}_{CL, \hat{c}^*}$.

3.5 Norm comparisons of inverse estimators

Although the theoretical focus of this paper has been on estimation of Σ rather than its inverse, it is still of interest to see how well $\tilde{\Sigma}_{S-MT, 2}^{-1}$, $\tilde{\Sigma}_{S-MT, \hat{\delta}}^{-1}$, $\tilde{\Sigma}_{BL, \hat{c}^*}^{-1}$, $\tilde{\Sigma}_{CL, \hat{c}^*}^{-1}$, and $\hat{\Sigma}_{LW_{\hat{\Sigma}}}^{-1}$ estimate Σ^{-1} , assuming that Σ^{-1} is well defined. Table 4 provides average norm losses for Monte Carlo design B for which Σ is positive definite. Σ for design A is ill-conditioned and will not be considered any further here. As can be seen from the results in Table 4, both $\tilde{\Sigma}_{S-MT, 2}^{-1}$ and $\tilde{\Sigma}_{S-MT, \hat{\delta}}^{-1}$ perform much better than $\tilde{\Sigma}_{BL, \hat{c}^*}^{-1}$ and $\tilde{\Sigma}_{CL, \hat{c}^*}^{-1}$ for Gaussian and multivariate t -distributed observations. In fact, the average spectral norms for $\tilde{\Sigma}_{BL, \hat{c}^*}^{-1}$ and

$\tilde{\Sigma}_{CL,\hat{C}^*}^{-1}$ include some sizeable outliers, especially for $N \leq 100$. However, the ranking of the different estimators remains the same if we use the Frobenius norm which appears to be less sensitive to the outliers. It is also worth noting that $\tilde{\Sigma}_{S-MT,2}^{-1}$ and $\tilde{\Sigma}_{S-MT,\hat{\delta}}^{-1}$ perform better than $LW_{\hat{\Sigma}}$, for all sample sizes and irrespective of whether the observations are drawn as Gaussian or multivariate t . Finally, using $\hat{\delta}$ rather than $\delta = 2$ when implementing the MT method improves the precision of the estimated inverse covariance matrix across all experiments.

3.6 Support recovery statistics

Table 5 reports the true positive and false positive rates (TPR and FPR) for the support recovery of Σ using the multiple testing and thresholding estimators. In the comparison set we include three versions of the MT estimator ($\tilde{\Sigma}_{MT,1}$, $\tilde{\Sigma}_{MT,2}$ and $\tilde{\Sigma}_{MT,\hat{\delta}}$), $\tilde{\Sigma}_{BL,\hat{C}}$, $\tilde{\Sigma}_{CL,2}$, and $\tilde{\Sigma}_{CL,\hat{C}}$. Again we use 100 replications due to the use of cross-validation in the implementation of MT , BL and CL thresholding. We include the MT estimators for choices of the scaling factor where $\delta = 1$ and $\delta = 2$, computed at $p = 0.05$, to see if our theoretical result, namely that for consistent support recovery only the linear scaling factor, where $\delta = 1$, is needed, is borne out by the simulations. Further, we implement MT using $\hat{\delta}$ to verify that the support recovery results under $MT_{\hat{\delta}}$ correspond more closely to those under MT_1 , in line with the findings of Theorem 3. For consistent support recovery we would like to see FPR values near zero and TPR values near unity. As can be seen from Table 5, the FPR values of all estimators are very close to zero, so any comparisons of different estimators must be based on the TPR values. Comparing the results for $\tilde{\Sigma}_{MT,1}$ and $\tilde{\Sigma}_{MT,2}$ we find that as predicted by the theory (Theorem 3 and Remark 13), TPR values of $\tilde{\Sigma}_{MT,1}$ are closer to unity as compared to the TPR values of $\tilde{\Sigma}_{MT,2}$. This is supported by the TPR values of $\tilde{\Sigma}_{MT,\hat{\delta}}$ as well. Similar results are obtained for the MT estimators for different choices of the p values. Table 6 provides results for $p = \{0.01, 0.05, 0.10\}$, and for $\delta = \{1, 2, \hat{\delta}\}$ using 2,000 replications. In this table it is further evident that, in line with the conclusions of Section 3.3, both the TPR and the FPR statistics are relatively robust to the choice of the p values

irrespective of the scaling factor, or whether the observations are drawn from a Gaussian or a multivariate t distribution. This is especially true under design B, since for this specification we explicitly control for the number of non-zero elements in Σ , that ensures the conditions of Theorem 3 are met.

Turning to a comparison with other estimators in Table 5, we find that the MT and CL estimators perform substantially better than the BL estimator. Further, allowing for dependence in the errors causes the support recovery performance of $BL_{\hat{C}}$, CL_2 and $CL_{\hat{C}}$ to deteriorate noticeably while MT_1 , MT_2 and $MT_{\hat{\delta}}$ remain remarkably stable. Finally, again note that TPR values are higher for design B. Overall, the estimators $\tilde{\Sigma}_{MT,1}$ or $\tilde{\Sigma}_{MT,\hat{\delta}}$ do best in recovering the support of Σ as compared to other estimators, although the results of CL and MT for support recovery can be very close, which is in line with the comparative analysis carried out in terms of the relative norm losses of these estimators.

3.7 Cross-validation of δ

We calibrate δ , the parameter of the critical value function, $c_p(N)$, in the MT approach, by following closely the cross-validation procedure implemented in BL and CL. Importantly, Bickel and Levina (2008a) show theoretically the validity of this approach for the ‘sample splitting’, ‘2-fold cross-validation’ and more general ‘V-fold cross-validation’ procedures.

More precisely, we perform a grid search for the choice of δ over the range: $\delta = \{c : \delta_{\min} \leq c \leq \delta_{\max}\}$. We set $\delta_{\min} = 1.0$ and $\delta_{\max} = 2.5$ and impose either fixed increments of 0.1 or N -dependent increments of $1/N$.⁶ At each point of the range, c , we generate x_{it} , $i = 1, 2, \dots, N$, $t = 1, 2, \dots, T$ and select the $N \times 1$ column vectors $\mathbf{x}_t = (x_{1t}, x_{2t}, \dots, x_{Nt})'$, $t = 1, 2, \dots, T$ which we randomly reshuffle over the t -dimension. This yields a new set of $N \times 1$ column vectors $\mathbf{x}_t^{(s)} = (x_{1t}^{(s)}, x_{2t}^{(s)}, \dots, x_{Nt}^{(s)})'$ for the first shuffle $s = 1$. We repeat this reshuffling S times in total where we set $S = 50$. We consider this to be sufficiently large (FLM suggested $S = 20$ while BL recommended $S = 100$ - see also Fang *et al.* (2016)).

⁶The sample size dependent alternative provides slight improvement in estimation precision for δ , but is computationally more expensive as N rises.

For each shuffle $s = 1, 2, \dots, S$, we divide $\mathbf{x}^{(s)} = (\mathbf{x}_1^{(s)}, \mathbf{x}_2^{(s)}, \dots, \mathbf{x}_T^{(s)})$ into two subsamples of size $N \times T_1$ and $N \times T_2$, where $T_2 = T - T_1$. The theoretically ‘justified’ split suggested in BL is given by $T_1 = T \left(1 - \frac{1}{\ln(T)}\right)$ and $T_2 = \frac{T}{\ln(T)}$. In our simulation study we set $T_1 = \frac{2T}{3}$ and $T_2 = \frac{T}{3}$. Let $\hat{\Sigma}_1^{(s)} = (\hat{\sigma}_{1,ij}^{(s)})$ with elements $\hat{\sigma}_{1,ij}^{(s)} = T_1^{-1} \sum_{t=1}^{T_1} x_{it}^{(s)} x_{jt}^{(s)}$ and $\hat{\Sigma}_2^{(s)} = (\hat{\sigma}_{2,ij}^{(s)})$ with elements $\hat{\sigma}_{2,ij}^{(s)} = T_2^{-1} \sum_{t=T_1+1}^T x_{it}^{(s)} x_{jt}^{(s)}$, $i, j = 1, 2, \dots, N$ denote the sample covariance matrices generated using T_1 and T_2 respectively, for each shuffle s . The corresponding sample correlation matrices are given by $\hat{\mathbf{R}}_1^{(s)} = [\hat{\mathbf{D}}_1^{(s)}]^{-1/2} \hat{\Sigma}_1^{(s)} [\hat{\mathbf{D}}_1^{(s)}]^{-1/2}$ and $\hat{\mathbf{R}}_2^{(s)} = [\hat{\mathbf{D}}_2^{(s)}]^{-1/2} \hat{\Sigma}_2^{(s)} [\hat{\mathbf{D}}_2^{(s)}]^{-1/2}$ respectively, where $\hat{\mathbf{D}}_i^{(s)} = \text{diag}(\hat{\sigma}_{i,11}^{(s)}, \hat{\sigma}_{i,22}^{(s)}, \dots, \hat{\sigma}_{i,NN}^{(s)})$, $i = 1, 2$. We regularise $\hat{\mathbf{R}}_1^{(s)}$ using the MT method in (5) and compute the following expression,

$$\hat{J}(c) = \frac{1}{S} \sum_{s=1}^S \left\| \tilde{\mathbf{R}}_1^{(s)}(c) - \hat{\mathbf{R}}_2^{(s)} \right\|_F^2, \quad (38)$$

for each c and

$$\hat{\delta} = \arg \inf_{\delta_{\min} \leq c \leq \delta_{\max}} \hat{J}(c). \quad (39)$$

The final estimator of the correlation matrix is then given by $\tilde{\mathbf{R}}_{\hat{\delta}}$ and the associated covariance matrix estimator, $\tilde{\Sigma}_{\hat{\delta}}$, is computed as in (7).

3.8 Computational demands of the different thresholding methods

Table 7 reports the relative execution times of the different thresholding methods studied. All times are relative to the time it takes to carry out the computations for the MT_2 estimator. The computational times shown for the methods that use a calibrated threshold parameter (i.e. $MT_{\hat{\delta}}$, $BL_{\hat{c}}$ and $CL_{\hat{c}}$) assume a sample-dependent grid in their respective CV procedures. It took 0.010, 0.013, and 0.014 seconds to apply the MT method in Matlab to a sample of $N = \{30, 100, 200\}$, respectively, and $T = 100$ observations using a desktop PC. The execution times of MT_1 and MT_2 are very similar and differ only slightly across the experiments with different p-values. In contrast, the $BL_{\hat{c}}$ and $CL_{\hat{c}}$ thresholding approaches are computationally much more demanding. Their computations took between about 12 and

485257 times (depending on N) longer than the MT_2 approach, for the same sample sizes and computer hardware. The $BL_{\hat{C}}$ method was less demanding than the $CL_{\hat{C}}$ method - it took between about 12 and 584 times longer than the MT_2 approach. Even CL_2 , which does not require estimation of the threshold parameter, took up to 19 times longer than the MT_2 approach. Thus, compared with other thresholding methods, MT_1 and MT_2 procedures have a clear computational advantage over the CL and BL procedures. This is not a surprising outcome, considering that MT_1 and MT_2 do not involve cross validation. But we find similar computational advantages for the MT procedure when we compare its cross-validated version, $MT_{\hat{\delta}}$, with $CL_{\hat{C}}$. The execution times of $MT_{\hat{\delta}}$ were between 1278 and 482038 faster than $CL_{\hat{C}}$. But when compared to $BL_{\hat{C}}$, we find that $BL_{\hat{C}}$ is somewhere between 24 and 2634 faster to compute than $MT_{\hat{\delta}}$. However, when using a fixed point increment in the implementation of the $MT_{\hat{\delta}}$ procedure, the computational advantage of $BL_{\hat{C}}$ over $MT_{\hat{\delta}}$ disappears.

4 Concluding Remarks

This paper considers regularisation of large covariance matrices particularly when the cross section dimension N of the data under consideration exceeds the time dimension T . In this case the sample covariance matrix, $\hat{\Sigma}$, becomes ill-conditioned and is not a satisfactory estimator of the population covariance.

A regularisation estimator is proposed which makes use of insights from the multiple testing literature to obtain threshold values for sample correlation coefficients. The proposed MT estimator of the correlation coefficient (ρ_{ij}) is set to zero when the sample correlation, in absolute value, is below the threshold, otherwise the MT estimator is set to the sample correlation coefficient. It is shown that the resultant estimator has a convergence rate of the order of $m_N c_p(N)/\sqrt{T}$ under the spectral norm, and $\sqrt{m_N N/T}$ under the Frobenius norm, where N is the number of units each observed T times, m_N measures the degree of sparsity of the population correlation matrix, and $c_p(N) = \Phi^{-1}\left(1 - \frac{p}{2N^\delta}\right)$, where $\Phi^{-1}(\cdot)$ is

the inverse of the cumulative distribution of a standard normal variate, and p is the nominal size of the test. $c_p(N)$ directly corresponds to $\sqrt{\ln(N)}$. Our analysis allows for non-Gaussian observations and provides guidance as to the choice of critical value function for thresholding in terms of the degree to which underlying observations are dependent even if $\rho_{ij} = 0$. The choice of δ depends on the degree of non-Gaussianity of the underlying observations and yields spectral norm results that are similar to the rates obtained in the literature. But for the Frobenius norm we obtain better rates than those established in the literature.

The numerical properties of the proposed estimator are investigated using Monte Carlo simulations. It is shown that the MT estimator performs well, and generally better than the other estimators proposed in the literature. The simulations also show that in terms of spectral and Frobenius norm losses, the MT estimator is reasonably robust to the choice of p in the threshold criterion, $|\hat{\rho}_{ij}| > T^{-1/2}\Phi^{-1}\left(1 - \frac{p}{2f(N)}\right)$, where $f(N) = c_\delta N^\delta$, with c_δ and δ being finite positive constants, particularly when setting $\delta = 2$. For support recovery, better results are obtained if $\delta = 1$.

Acknowledgements

The authors are grateful to Jianqing Fan (Co-Editor), an Associate Editor and three anonymous reviewers for valuable comments and suggestions, as well as Elizaveta Levina and Martina Mincheva for helpful email correspondence with regard to implementation of their approaches. They would also like to thank Alex Chudik, George Kapetanios, Yuan Liao, Ron Smith and Michael Wolf for useful comments. Financial support under ESRC Grant ES/I031626/1 is gratefully acknowledged by the authors.

Table 1a: Spectral and Frobenius norm losses for the MT estimator using significance levels $p = \{0.01, 0.05, 0.10\}$ and scaling factors with $\delta = \{1, 2, \hat{\delta}\}$, for $T = 100$

		Monte Carlo design A								
		$\delta = 1$			$\delta = 2$			$\hat{\delta}$		
$N \setminus p$		0.01	0.05	0.10	0.01	0.05	0.10	0.01	0.05	0.10
$\mathbf{u}_{it} \sim \text{Gaussian}$										
<i>Spectral norm</i>										
30		1.70(0.49)	1.68(0.49)	1.71(0.49)	1.89(0.51)	1.79(0.50)	1.75(0.50)	1.71(0.49)	1.68(0.49)	1.69(0.49)
100		2.61(0.50)	2.51(0.50)	2.50(0.50)	3.11(0.50)	2.91(0.50)	2.84(0.50)	2.62(0.50)	2.52(0.50)	2.51(0.50)
200		3.04(0.48)	2.92(0.49)	2.89(0.49)	3.67(0.47)	3.46(0.47)	3.37(0.47)	3.05(0.48)	2.93(0.49)	2.90(0.49)
<i>Frobenius norm</i>										
30		3.17(0.45)	3.14(0.50)	3.20(0.53)	3.49(0.42)	3.32(0.43)	3.26(0.43)	3.19(0.44)	3.13(0.48)	3.16(0.52)
100		6.67(0.45)	6.51(0.51)	6.60(0.55)	7.75(0.40)	7.34(0.41)	7.17(0.42)	6.70(0.45)	6.52(0.50)	6.57(0.54)
200		9.87(0.46)	9.60(0.53)	9.73(0.58)	11.76(0.40)	11.15(0.41)	10.89(0.42)	9.91(0.46)	9.62(0.52)	9.69(0.57)
$\mathbf{u}_{it} \sim \text{multivariate } t\text{-distributed with 8 degrees of freedom}$										
<i>Spectral norm</i>										
30		2.26(1.08)	2.42(1.20)	2.55(1.26)	2.29(0.90)	2.24(0.99)	2.24(1.03)	2.23(0.95)	2.32(1.04)	2.39(1.08)
100		3.85(4.84)	4.20(5.28)	4.46(5.48)	3.78(3.78)	3.71(4.12)	3.71(4.27)	3.67(3.81)	3.83(4.11)	3.93(4.21)
200		4.49(3.46)	5.04(4.34)	5.44(4.77)	4.26(1.80)	4.20(2.21)	4.19(2.37)	4.20(2.43)	4.45(2.78)	4.57(2.94)
<i>Frobenius norm</i>										
30		4.06(1.14)	4.35(1.32)	4.60(1.40)	4.12(0.90)	4.04(1.00)	4.03(1.06)	4.03(1.00)	4.19(0.13)	4.32(1.19)
100		8.88(5.17)	9.75(5.67)	10.49(5.87)	9.04(4.04)	8.80(4.40)	8.74(4.57)	8.65(4.16)	9.09(4.48)	9.41(4.59)
200		12.96(4.23)	14.50(5.41)	15.81(5.95)	13.25(2.10)	12.85(2.54)	12.71(2.76)	12.57(2.97)	13.25(3.48)	13.73(3.67)

$N \setminus p$	Cross validated values of δ		
	0.01	0.05	0.10
$\mathbf{u}_{it} \sim \text{Gaussian}$			
30	1.08(0.11)	1.10(0.12)	1.12(0.13)
100	1.04(0.06)	1.05(0.07)	1.06(0.08)
200	1.03(0.05)	1.03(0.06)	1.04(0.06)
$\mathbf{u}_{it} \sim \text{multivariate } t\text{-distr. with 8 dof}$			
30	1.13(0.18)	1.19(0.22)	1.25(0.25)
100	1.12(0.18)	1.18(0.22)	1.23(0.25)
200	1.15(0.20)	1.20(0.23)	1.24(0.25)

Note: The MT approach is implemented using $\delta = 1$, $\delta = 2$, and $\hat{\delta}$, computed using cross-validation. Norm losses and estimates of δ , $\hat{\delta}$, are averages over 2,000 replications. Simulation standard deviations are given in parentheses.

Table 1b: Spectral and Frobenius norm losses for the MT estimator using significance levels $p = \{0.01, 0.05, 0.10\}$ and scaling factors with $\delta = \{1, 2, \hat{\delta}\}$, for $T = 100$

		Monte Carlo design B								
		$\delta = 1$			$\delta = 2$			$\hat{\delta}$		
$N \setminus p$		0.01	0.05	0.10	0.01	0.05	0.10	0.01	0.05	0.10
$\mathbf{u}_{it} \sim \text{Gaussian}$										
<i>Spectral norm</i>										
30		0.48(0.16)	0.50(0.16)	0.53(0.16)	0.50(0.20)	0.49(0.18)	0.48(0.17)	0.48(0.17)	0.49(0.16)	0.49(0.16)
100		0.75(0.34)	0.76(0.32)	0.78(0.31)	0.89(0.43)	0.81(0.39)	0.79(0.37)	0.76(0.35)	0.76(0.34)	0.76(0.34)
200		0.71(0.22)	0.74(0.20)	0.77(0.20)	0.85(0.33)	0.78(0.28)	0.75(0.26)	0.72(0.24)	0.72(0.22)	0.72(0.22)
<i>Frobenius norm</i>										
30		0.87(0.17)	0.91(0.18)	0.97(0.19)	0.89(0.20)	0.87(0.17)	0.86(0.17)	0.86(0.17)	0.88(0.17)	0.88(0.17)
100		1.56(0.24)	1.66(0.24)	1.77(0.24)	1.67(0.34)	1.60(0.29)	1.58(0.27)	1.56(0.25)	1.58(0.24)	1.58(0.25)
200		2.16(0.18)	2.32(0.20)	2.50(0.21)	2.25(0.24)	2.19(0.21)	2.16(0.20)	2.15(0.18)	2.18(0.19)	2.18(0.20)
$\mathbf{u}_{it} \sim \text{multivariate } t\text{-distributed with 8 degrees of freedom}$										
<i>Spectral norm</i>										
30		0.70(0.39)	0.78(0.43)	0.84(0.45)	0.67(0.33)	0.67(0.35)	0.67(0.37)	0.67(0.33)	0.68(0.35)	0.68(0.36)
100		1.16(0.97)	1.32(1.10)	1.42(1.18)	1.15(0.75)	1.11(0.80)	1.10(0.83)	1.10(0.72)	1.10(0.77)	1.11(0.80)
200		1.36(1.73)	1.65(2.05)	1.83(2.20)	1.14(1.03)	1.13(1.21)	1.14(1.28)	1.16(1.06)	1.19(1.20)	1.20(1.27)
<i>Frobenius norm</i>										
30		1.23(0.42)	1.40(0.48)	1.53(0.51)	1.15(0.35)	1.16(0.38)	1.17(0.39)	1.17(0.36)	1.19(0.38)	1.20(0.39)
100		2.39(1.12)	2.90(1.31)	3.25(1.40)	2.17(0.77)	2.15(0.86)	2.16(0.90)	2.17(0.76)	2.22(0.85)	2.24(0.89)
200		3.57(2.13)	4.52(2.54)	5.18(2.72)	2.97(1.21)	2.98(1.43)	3.01(1.53)	3.06(1.27)	3.17(1.48)	3.21(1.57)

$N \setminus p$	Cross validated values of δ		
	0.01	0.05	0.10
$\mathbf{u}_{it} \sim \text{Gaussian}$			
30	1.27(0.27)	1.46(0.35)	1.61(0.36)
100	1.25(0.24)	1.43(0.31)	1.56(0.32)
200	1.23(0.22)	1.36(0.26)	1.49(0.27)
$\mathbf{u}_{it} \sim \text{multivariate } t\text{-distr. with 8 dof}$			
30	1.45(0.38)	1.72(0.39)	1.87(0.35)
100	1.59(0.41)	1.76(0.40)	1.85(0.37)
200	1.68(0.44)	1.78(0.41)	1.85(0.39)

The MT approach is implemented using $\delta = 1$, $\delta = 2$, and $\hat{\delta}$, computed using cross-validation. Norm losses and estimates of δ , $\hat{\delta}$, are averages over 2,000 replications. Simulation standard deviations are given in parentheses.

Table 2: Spectral and Frobenius norm losses for different regularised covariance matrix estimators ($T = 100$) - Monte Carlo design A

	$N = 30$		$N = 100$		$N = 200$	
	Norms		Norms		Norms	
	Spectral	Frobenius	Spectral	Frobenius	Spectral	Frobenius
$\mathbf{u}_{it} \sim \text{Gaussian}$						
<i>Error matrices ($\Sigma - \tilde{\Sigma}$)</i>						
MT_2	1.85(0.53)	3.38(0.40)	2.83(0.50)	7.29(0.42)	3.45(0.43)	11.17(0.38)
$MT_{\hat{\delta}}$	1.75(0.55)	3.21(0.49)	2.44(0.50)	6.48(0.50)	2.95(0.45)	9.65(0.48)
$BL_{\hat{C}}$	5.30(2.16)	7.61(1.23)	8.74(0.06)	16.90(0.10)	8.94(0.04)	24.26(0.13)
CL_2	1.87(0.55)	3.39(0.44)	2.99(0.49)	7.57(0.44)	3.79(0.47)	11.88(0.42)
$CL_{\hat{C}}$	1.82(0.58)	3.33(0.56)	2.54(0.50)	6.82(0.51)	3.02(0.46)	10.22(0.59)
$S-MT_2$	3.36(0.78)	4.45(0.63)	5.83(0.34)	10.95(0.47)	6.47(0.21)	16.64(0.35)
$S-MT_{\hat{\delta}}$	2.67(0.81)	3.85(0.65)	5.08(0.40)	9.70(0.51)	5.79(0.27)	14.91(0.46)
$BL_{\hat{C}^*}$	7.09(0.10)	8.62(0.09)	8.74(0.06)	16.90(0.10)	8.94(0.04)	24.25(0.10)
$CL_{\hat{C}^*}$	7.05(0.16)	8.58(0.12)	8.71(0.07)	16.85(0.11)	8.94(0.04)	24.23(0.09)
$LW_{\hat{\Sigma}}$	2.99(0.47)	6.49(0.29)	5.20(0.34)	16.70(0.19)	6.28(0.20)	26.84(0.14)
$\mathbf{u}_{it} \sim \text{multivariate } t\text{-distributed with 8 degrees of freedom}$						
<i>Error matrices ($\Sigma - \tilde{\Sigma}$)</i>						
MT_2	2.17(0.72)	4.02(0.88)	3.44(0.98)	8.52(1.17)	4.00(0.83)	12.79(1.66)
$MT_{\hat{\delta}}$	2.27(0.88)	4.20(1.11)	3.59(1.39)	8.76(1.65)	4.32(1.53)	13.28(2.83)
$BL_{\hat{C}}$	6.90(0.82)	8.75(0.55)	8.74(0.10)	17.26(0.30)	9.00(0.42)	24.93(1.02)
CL_2	2.55(0.93)	4.53(1.00)	4.63(1.11)	10.35(1.48)	5.92(0.81)	16.43(1.74)
$CL_{\hat{C}}$	2.27(0.76)	4.24(0.94)	3.85(1.51)	9.44(2.33)	5.04(2.04)	15.65(4.71)
$S-MT_2$	3.28(0.80)	4.76(0.77)	5.84(0.45)	11.47(0.62)	6.48(0.32)	17.27(0.71)
$S-MT_{\hat{\delta}}$	2.86(0.92)	4.51(0.97)	5.30(0.52)	10.76(0.77)	6.00(0.39)	16.36(1.04)
$BL_{\hat{C}^*}$	7.06(0.13)	8.84(0.30)	8.74(0.10)	17.25(0.31)	8.95(0.08)	24.84(0.55)
$CL_{\hat{C}^*}$	7.01(0.16)	8.77(0.30)	8.73(0.11)	17.23(0.29)	8.94(0.08)	24.77(0.53)
$LW_{\hat{\Sigma}}$	3.35(0.51)	7.35(0.50)	5.67(0.46)	18.04(0.45)	6.60(0.43)	28.18(0.53)

Note: Norm losses are averages over 100 replications. Simulation standard deviations are given in parentheses. $\tilde{\Sigma} = \{\tilde{\Sigma}_{MT,2}, \tilde{\Sigma}_{MT,\hat{\delta}}, \tilde{\Sigma}_{BL,\hat{C}}, \tilde{\Sigma}_{CL,2}, \tilde{\Sigma}_{CL,\hat{C}}, \tilde{\Sigma}_{S-MT,2}, \tilde{\Sigma}_{S-MT,\hat{\delta}}, \tilde{\Sigma}_{BL,\hat{C}^*}, \tilde{\Sigma}_{CL,\hat{C}^*}, \tilde{\Sigma}_{LW_{\hat{\Sigma}}}\}$. $\tilde{\Sigma}_{MT,2}, \tilde{\Sigma}_{MT,\hat{\delta}}, \tilde{\Sigma}_{S-MT,2}$ and $\tilde{\Sigma}_{S-MT,\hat{\delta}}$ are computed using $p = 0.05$. $(MT_2, S-MT_2)$ and $(MT_{\hat{\delta}}, S-MT_{\hat{\delta}})$ are thresholding based on multiple testing with critical value $\Phi^{-1}\left(1 - \frac{p}{2f(N)}\right)$, where $f(N) = N^2$ and $f(N) = N^{\hat{\delta}}$, respectively, with $\hat{\delta}$ being estimated by cross-validation. BL is Bickel and Levina universal thresholding, CL is Cai and Liu adaptive thresholding, $\tilde{\Sigma}_{MT,2}$ and $\tilde{\Sigma}_{MT,\hat{\delta}}$ are based on MT_2 and $MT_{\hat{\delta}}$. $\tilde{\Sigma}_{S-MT,2}$ and $\tilde{\Sigma}_{S-MT,\hat{\delta}}$ apply supplementary shrinkage to $\tilde{\Sigma}_{MT,2}$ and $\tilde{\Sigma}_{MT,\hat{\delta}}$, $\tilde{\Sigma}_{BL,\hat{C}}$ and $\tilde{\Sigma}_{CL,\hat{C}}$ are based on \hat{C} which is obtained by cross-validation, $\tilde{\Sigma}_{BL,\hat{C}^*}$ and $\tilde{\Sigma}_{CL,\hat{C}^*}$ employ the further adjustment to the cross-validation coefficient, \hat{C}^* , proposed by Fan *et al.* (2013), $\tilde{\Sigma}_{CL,2}$ is CL's estimator with $C = 2$ (the theoretical value of C). $\tilde{\Sigma}_{LW_{\hat{\Sigma}}}$ is Ledoit and Wolf's shrinkage estimator applied to the sample covariance matrix.

Table 3: Spectral and Frobenius norm losses for different regularised covariance matrix estimators ($T = 100$) - Monte Carlo design B

	$N = 30$		$N = 100$		$N = 200$	
	Norms		Norms		Norms	
	Spectral	Frobenius	Spectral	Frobenius	Spectral	Frobenius
$\mathbf{u}_{it} \sim \text{Gaussian}$						
<i>Error matrices ($\Sigma - \hat{\Sigma}$)</i>						
MT_2	0.49(0.18)	0.89(0.19)	0.87(0.37)	1.63(0.28)	0.73(0.24)	2.15(0.19)
$MT_{\hat{\delta}}$	0.48(0.14)	0.89(0.16)	0.79(0.31)	1.57(0.23)	0.67(0.18)	2.15(0.17)
$BL_{\hat{C}}$	0.91(0.50)	1.35(0.43)	1.40(0.95)	2.25(0.78)	2.53(0.55)	3.49(0.32)
CL_2	0.49(0.17)	0.90(0.18)	1.00(0.48)	1.77(0.44)	0.90(0.37)	2.30(0.30)
$CL_{\hat{C}}$	0.49(0.15)	0.92(0.17)	0.83(0.31)	1.71(0.28)	1.14(0.83)	2.54(0.58)
$S-MT_2$	0.68(0.27)	1.08(0.21)	1.53(0.53)	2.16(0.38)	1.23(0.41)	2.44(0.26)
$S-MT_{\hat{\delta}}$	0.66(0.23)	1.07(0.18)	1.45(0.44)	2.08(0.29)	1.12(0.30)	2.38(0.19)
$BL_{\hat{C}^*}$	1.19(0.46)	1.63(0.40)	3.32(0.20)	3.90(0.14)	2.73(0.11)	3.61(0.08)
$CL_{\hat{C}^*}$	1.08(0.46)	1.53(0.46)	3.34(0.15)	3.92(0.06)	2.73(0.10)	3.61(0.08)
$LW_{\hat{\Sigma}}$	1.05(0.13)	2.07(0.10)	2.95(0.26)	4.47(0.09)	2.46(0.06)	6.01(0.03)
$\mathbf{u}_{it} \sim \text{multivariate } t\text{-distributed with 8 degrees of freedom}$						
<i>Error matrices ($\Sigma - \hat{\Sigma}$)</i>						
MT_2	0.64(0.24)	1.12(0.24)	1.05(0.45)	2.13(0.49)	1.29(2.32)	3.15(2.66)
$MT_{\hat{\delta}}$	0.66(0.25)	1.15(0.26)	1.03(0.42)	2.17(0.53)	1.30(1.90)	3.29(2.22)
$BL_{\hat{C}}$	1.36(0.40)	1.84(0.35)	2.70(0.94)	3.58(0.74)	2.70(0.29)	4.08(0.67)
CL_2	0.71(0.29)	1.21(0.30)	1.69(0.70)	2.73(0.70)	1.62(0.57)	3.31(0.65)
$CL_{\hat{C}}$	0.80(0.39)	1.33(0.39)	2.03(1.08)	3.07(0.90)	2.19(0.78)	3.72(0.62)
$S-MT_2$	0.69(0.26)	1.18(0.23)	1.41(0.57)	2.36(0.47)	1.32(0.79)	3.02(0.87)
$S-MT_{\hat{\delta}}$	0.69(0.25)	1.19(0.22)	1.36(0.49)	2.34(0.42)	1.30(0.78)	3.10(0.87)
$BL_{\hat{C}^*}$	1.49(0.26)	1.98(0.21)	3.33(0.24)	4.07(0.18)	2.77(0.37)	4.04(0.56)
$CL_{\hat{C}^*}$	1.26(0.40)	1.79(0.40)	3.35(0.17)	4.08(0.14)	2.73(0.14)	4.01(0.42)
$LW_{\hat{\Sigma}}$	1.13(0.15)	2.25(0.11)	3.14(0.21)	4.68(0.11)	2.52(0.08)	6.18(0.13)

See the note to Table 2.

Table 4: Spectral and Frobenius norm losses for the inverses of different regularised covariance matrix estimators for Monte Carlo design B - $T = 100$

	$N = 30$		$N = 100$		$N = 200$	
	Norms		Norms		Norms	
	Spectral	Frobenius	Spectral	Frobenius	Spectral	Frobenius
<i>Error matrices</i> ($\Sigma^{-1} - \hat{\Sigma}^{-1}$)						
$\mathbf{u}_{it} \sim \text{Gaussian}$						
$S\text{-}MT_2$	4.44(1.23)	2.66(0.32)	15.81(2.63)	5.90(0.45)	14.24(2.37)	5.50(0.38)
$S\text{-}MT_{\hat{\delta}}$	4.36(1.22)	2.64(0.31)	15.25(2.78)	5.80(0.48)	13.36(2.47)	5.39(0.37)
$BL_{\hat{C}^*}$	$3.8 \times 10^3 (2.4 \times 10^4)$	19.56(58.88)	$1.2 \times 10^3 (1.1 \times 10^4)$	12.16(33.25)	41.07(143.74)	7.66(3.17)
$CL_{\hat{C}^*}$	$1.9 \times 10^3 (1.7 \times 10^4)$	10.92(42.39)	51.99(241.39)	8.16(4.23)	28.45(24.37)	7.35(1.11)
$LW_{\hat{\Sigma}}$	11.03(0.58)	4.26(0.09)	31.04(0.64)	8.62(0.06)	31.81(0.21)	9.40(0.05)
$\mathbf{u}_{it} \sim \text{multivariate } t\text{-distributed with 8 degrees of freedom}$						
$S\text{-}MT_2$	3.45(1.61)	2.44(0.39)	12.78(3.13)	5.55(0.55)	11.57(4.17)	5.58(0.66)
$S\text{-}MT_{\hat{\delta}}$	3.43(1.63)	2.45(0.40)	12.37(3.27)	5.51(0.59)	11.28(3.97)	5.65(0.67)
$BL_{\hat{C}^*}$	$157.26 (1.0 \times 10^3)$	6.11(11.28)	$349.35 (3.1 \times 10^3)$	9.80(17.03)	28.58(22.06)	7.77(1.04)
$CL_{\hat{C}^*}$	85.82(546.85)	5.53(7.84)	$517.27 (4.8 \times 10^3)$	10.07(21.25)	25.61(3.55)	7.54(0.50)
$LW_{\hat{\Sigma}}$	12.08(1.19)	4.48(0.20)	31.78(1.32)	8.74(0.23)	32.06(1.00)	9.50(0.33)

Note: $\hat{\Sigma}^{-1} = \{\tilde{\Sigma}_{S\text{-}MT,2}^{-1}, \tilde{\Sigma}_{S\text{-}MT,\hat{\delta}}^{-1}, \tilde{\Sigma}_{BL,\hat{C}^*}^{-1}, \tilde{\Sigma}_{CL,\hat{C}^*}^{-1}, \hat{\Sigma}_{LW_{\hat{\Sigma}}}^{-1}\}$. See also the note to Table 2.

Table 5: Support recovery statistics for different multiple testing and thresholding estimators - $T = 100$

N	Monte Carlo design A						N	Monte Carlo design B					
	MT_1	MT_2	$MT_{\hat{\delta}}$	$BL_{\hat{C}}$	CL_2	$CL_{\hat{C}}$		MT_1	MT_2	$MT_{\hat{\delta}}$	$BL_{\hat{C}}$	CL_2	$CL_{\hat{C}}$
$\mathbf{u}_{it} \sim \text{Gaussian}$													
30	TPR 0.80	0.71	0.79	0.29	0.72	0.78	30	TPR 1.00	0.98	1.00	0.64	0.98	1.00
	FPR 0.00	0.00	0.00	0.04	0.00	0.00		FPR 0.00	0.00	0.00	0.00	0.00	0.00
100	TPR 0.69	0.57	0.69	0.00	0.56	0.68	100	TPR 1.00	0.98	1.00	0.80	0.94	0.99
	FPR 0.00	0.00	0.00	0.00	0.00	0.00		FPR 0.00	0.00	0.00	0.00	0.00	0.00
200	TPR 0.66	0.53	0.66	0.00	0.50	0.65	200	TPR 1.00	0.96	0.99	0.11	0.88	0.78
	FPR 0.00	0.00	0.00	0.00	0.00	0.00		FPR 0.00	0.00	0.00	0.00	0.00	0.00
$\mathbf{u}_{it} \sim \text{multivariate } t\text{-distributed with 8 degrees of freedom}$													
30	TPR 0.80	0.72	0.79	0.03	0.62	0.74	30	TPR 1.00	0.98	0.99	0.26	0.89	0.82
	FPR 0.01	0.00	0.00	0.00	0.00	0.00		FPR 0.01	0.00	0.00	0.00	0.00	0.00
100	TPR 0.69	0.58	0.67	0.00	0.43	0.57	100	TPR 1.00	0.97	0.98	0.27	0.70	0.57
	FPR 0.00	0.00	0.00	0.00	0.00	0.00		FPR 0.00	0.00	0.00	0.00	0.00	0.00
200	TPR 0.66	0.53	0.64	0.00	0.35	0.47	200	TPR 0.99	0.93	0.95	0.05	0.57	0.30
	FPR 0.00	0.00	0.00	0.00	0.00	0.00		FPR 0.00	0.00	0.00	0.00	0.00	0.00

Note: TPR is the true positive rate and FPR is the false positive rate defined by (30) and (31), respectively. MT estimators are computed with $p = 0.05$. For a description of other estimators see the note to Table 2. The TPR and FPR numbers are averages over 100 replications

Table 6: Support recovery statistics for the multiple testing estimator computed with $p = \{0.01, 0.05, 0.10\}$ - $T = 100$

N	Monte Carlo design A										Monte Carlo design B																			
	$p = 0.01$					$p = 0.05$					$p = 0.10$					$p = 0.01$					$p = 0.05$					$p = 0.10$				
	MT_1	MT_2	MT_δ	MT_1	MT_2	MT_δ	MT_1	MT_2	MT_δ	N	MT_1	MT_2	MT_δ	MT_1	MT_2	MT_δ	MT_1	MT_2	MT_δ	MT_1	MT_2	MT_δ	MT_1	MT_2	MT_δ					
$\mathbf{u}_{it} \sim \text{Gaussian}$																														
30	TPR 0.75	0.67	0.75	0.80	0.71	0.79	0.81	0.73	0.80	30	TPR 1.00	0.97	1.00	1.00	0.99	1.00	1.00	0.99	1.00	1.00	0.99	1.00	1.00	0.99	1.00	1.00				
	FPR 0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		FPR 0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00				
100	TPR 0.65	0.54	0.64	0.69	0.57	0.69	0.71	0.59	0.70	100	TPR 1.00	0.97	0.99	1.00	0.98	1.00	1.00	0.98	1.00	1.00	0.99	1.00	1.00	0.99	1.00	1.00				
	FPR 0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		FPR 0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00				
200	TPR 0.62	0.49	0.61	0.66	0.53	0.66	0.68	0.54	0.67	200	TPR 0.99	0.92	0.99	1.00	0.96	0.99	1.00	0.96	0.99	1.00	0.97	0.99	1.00	0.97	0.99	0.99				
	FPR 0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		FPR 0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00				
$\mathbf{u}_{it} \sim \text{multivariate } t\text{-distributed with 8 degrees of freedom}$																														
30	TPR 0.76	0.68	0.74	0.80	0.81	0.78	0.81	0.73	0.79	30	TPR 0.99	0.96	0.98	1.00	0.98	1.00	0.98	0.99	1.00	0.99	1.00	0.99	1.00	0.99	1.00	0.99				
	FPR 0.00	0.00	0.00	0.01	0.01	0.00	0.01	0.00	0.01		FPR 0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.00	0.00				
100	TPR 0.65	0.54	0.64	0.71	0.71	0.67	0.71	0.59	0.68	100	TPR 0.99	0.96	0.97	1.00	0.97	1.00	0.97	0.98	1.00	0.97	0.98	1.00	0.98	1.00	0.98	0.98				
	FPR 0.00	0.00	0.00	0.01	0.01	0.00	0.01	0.00	0.00		FPR 0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.00				
200	TPR 0.62	0.50	0.60	0.68	0.68	0.63	0.68	0.55	0.65	200	TPR 0.99	0.91	0.93	0.99	0.94	0.99	0.94	0.95	1.00	0.94	0.95	1.00	0.96	0.96	0.96	0.96				
	FPR 0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		FPR 0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00				

Note: TPR is the true positive rate and FPR is the false positive rate defined by (30) and (31), respectively. MT estimators are computed with $p = 0.05$. For a description of other estimators see the note to Table 2. The TPR and FPR numbers are averages over 2000 replications

Table 7: Relative execution times of different thresholding methods

	$T = 100$		
	$N = 30$	$N = 100$	$N = 200$
MT_2	1.000	1.000	1.000
MT_1	0.996	0.971	1.017
$MT_{\hat{\delta}}$	35.84	497.4	3219
$BL_{\hat{C}}$	11.53	106.3	584.8
CL_2	1.924	5.629	19.12
$CL_{\hat{C}}$	1314	63481	485257

Note: All times are relative to the MT_2 estimator.
 See Table 2 for a note on the thresholding methods.

Appendix: Mathematical proofs of theorems for the MT estimator

The lemmas referred to in this Appendix are stated and proved in a supplement which will be available online.

Proof of Proposition 1. The results for $E(\hat{\rho}_{ij,T})$ and $Var(\hat{\rho}_{ij,T})$ are established in Gayen (1951) using a bivariate Edgeworth expansion approach. This confirms earlier findings obtained by Tschuprow (1925) (English Translation, 1939) who shows that results (9) and (10) hold for any law of dependence between x_{it} and x_{jt} . See, in particular, p. 228 and equations (53) and (54) in Gayen (1951). To see that these results hold uniformly in the i and j ($i \neq j$) pairs, first note we have that $\sup_{ij} |\rho_{ij}| < 1$. Using (10) and (12) $\lim_{T \rightarrow \infty} [TVar(\hat{\rho}_{ij,T})] = K_v(\theta_{ij})$. The uniform boundedness of $|K_m(\theta_{ij})|$ and $K_v(\theta_{ij})$ follows directly from Assumption 2 that the sixth-order moment of y_{it} is uniformly bounded and application of Holder's and the Cauchy-Schwarz inequalities. Application of these inequalities establishes the uniform boundedness of the moments $E(y_{it}^3 y_{jt})$ and $E(y_{it}^2 y_{jt}^2)$ as given below:

$$\begin{aligned} \sup_{ij,t} |E(y_{it}^2 y_{jt}^2)| &\leq \sup_{ij,t} E(|y_{it}^2 y_{jt}^2|) \leq \sup_{ij,t} \left\{ [E(|y_{it}|^4)]^{1/2} [E(|y_{jt}|^4)]^{1/2} \right\} \\ &\leq \sup_{i,t} [E(|y_{it}|^4)]^{1/2} \sup_{j,t} [E(|y_{jt}|^4)]^{1/2} < K \end{aligned}$$

and

$$\begin{aligned} \sup_{ij,t} |E(y_{it} y_{jt}^3)| &\leq \sup_{ij,t} E(|y_{it} y_{jt}^3|) \leq \sup_{ij,t} \left\{ [E(|y_{it}|^4)]^{1/4} [E(|y_{jt}|^{4/3})]^{3/4} \right\} \\ &= \sup_{ij,t} \left\{ [E(|y_{it}|^4)]^{1/4} [E(|y_{jt}|^4)]^{3/4} \right\} = \sup_{i,t} E(|y_{it}|^4) < K. \end{aligned}$$

The remaining terms included in $O(T^{-2})$ in (9) and (10) as can be seen from Gayen (1951) are also a function of ρ_{ij} and $\kappa_{ij}(\cdot, \cdot)$ up to order $\kappa_{ij}(4, 0)$. Hence, the results of Gayen

(1951) hold uniformly across all i and j pair of correlations.

Consider now the case where y_{it} for all i are Gaussian. Then $E(y_{it}^4) = 3$, and for all $i \neq j$ we have

$$y_{it} = \rho_{ij}y_{jt} + \eta_{jt},$$

where $E(\eta_{jt}) = 0$, $Var(\eta_{jt}) = 1 - \rho_{ij}^2$, and η_{jt} and y_{jt} are independently distributed. Hence,

$$E(y_{jt}^3 y_{it}) = E[y_{jt}^3 (\rho_{ij} y_{jt} + \eta_{jt})] = \rho_{ij} E(y_{jt}^4) = 3\rho_{ij}.$$

$$E(y_{it}^2 y_{jt}^2) = E[y_{jt}^2 (\rho_{ij}^2 y_{jt}^2 + \eta_{jt}^2 + 2\rho_{ij} y_{jt} \eta_{jt})] = 3\rho_{ij}^2 + (1 - \rho_{ij}^2) = 1 + 2\rho_{ij}^2.$$

Using the above results it now follows that

$$\kappa_{ij}(4, 0) = \kappa_{ij}(0, 4) = 0, \quad \kappa_{ij}(3, 1) = \kappa_{ij}(1, 3) = 0,$$

which in turn establishes, when used in (11) and (12), that $K_m(\boldsymbol{\theta}_{ij}) = -\frac{1}{2}\rho_{ij}(1 - \rho_{ij}^2)$ and $K_v(\boldsymbol{\theta}_{ij}) = (1 - \rho_{ij}^2)^2$. ■

Proof of Proposition 2. Under Assumption 2, for a given i and j , set $\boldsymbol{\xi}_t = (y_{it}, y_{jt}, y_{it}^2, y_{jt}^2, y_{it}y_{jt})' = (\xi_{1t}, \xi_{2t}, \dots, \xi_{5t})'$, where $y_{it} = (x_{it} - \mu_i) / \sqrt{\sigma_{ii}}$. To simplify the notation we drop the subscripts i, j . Define

$$\bar{\boldsymbol{\xi}}_T = T^{-1} \sum_{t=1}^T \boldsymbol{\xi}_t = (\bar{\xi}_{1T}, \bar{\xi}_{2T}, \dots, \bar{\xi}_{5T})',$$

and note that by Assumption 2, $\boldsymbol{\xi}_t$, for $t = 1, 2, \dots, T$, are random draws from a common distribution with non-zero density, the elements of $\boldsymbol{\xi}_t$ are continuously differentiable functions of $\mathbf{y}_t = (y_{it}, y_{jt})'$. $\hat{\rho}_{ij,T}$, the sample correlation coefficient of x_{it} and x_{jt} , can be written as

$$\hat{\rho}_{ij,T} = H(\bar{\boldsymbol{\xi}}_T) = \frac{\bar{\xi}_{5T} - \bar{\xi}_{1T}\bar{\xi}_{2T}}{\left(\bar{\xi}_{3T} - \bar{\xi}_{1T}^2\right)^{1/2} \left(\bar{\xi}_{4T} - \bar{\xi}_{2T}^2\right)^{1/2}},$$

where $\bar{\xi}_{3T} > \bar{\xi}_{1T}^2$, and $\bar{\xi}_{4T} > \bar{\xi}_{2T}^2$. See also Bhattacharya and Ghosh (1978) - p. 434.

It is also easily seen that $\boldsymbol{\mu}_\xi = E(\bar{\boldsymbol{\xi}}_T) = (0, 0, 1, 1, \rho_{ij})'$, and $H(\boldsymbol{\mu}_\xi) = \rho_{ij}$, and hence

$\sqrt{T} [H(\bar{\boldsymbol{\xi}}_T) - H(\boldsymbol{\mu}_\xi)] = \sqrt{T} (\hat{\rho}_{ij,T} - \rho_{ij})$, where $H(\boldsymbol{\xi})$ is continuous and differentiable in $\boldsymbol{\xi}$, and all derivatives of $H(\boldsymbol{\xi})$ are continuous in a neighbourhood of $\boldsymbol{\mu}_\xi$; $1, \xi_{1t}, \xi_{2t}, \dots, \xi_{5t}$ are linearly independent, and $E|\xi_{kt}|^s < \infty$, for $k = 1, 2, \dots, 5$, for some positive integer $s \geq 3$. Hence, Theorem 2 of Bhattacharya and Ghosh (1978) can be applied to $\hat{\rho}_{ij,T}$, which establishes the validity of the Edgeworth expansion, (16). To prove (17) using (16) we first note that (for some $a_T > 0$)

$$\begin{aligned} \Pr(z_{ij,T} > a_T) &= 1 - \Pr(z_{ij,T} \leq a_T) \\ &= 1 - \Phi(a_T) - \sum_{r=1}^{s-2} T^{-r/2} g_r(a_T) \phi(a_T) + O[T^{-(s-1)/2}] \\ &= \Phi(-a_T) - (2\pi)^{-1/2} \exp\left(-\frac{a_T^2}{2}\right) \sum_{r=1}^{s-2} T^{-r/2} g_r(a_T) + O[T^{-(s-1)/2}], \end{aligned}$$

and by the inequality (A.1) (in the online supplement), we have

$$\Pr(z_{ij,T} > a_T) \leq \frac{1}{2} \exp\left(-\frac{a_T^2}{2}\right) + (2\pi)^{-1/2} \exp\left(-\frac{a_T^2}{2}\right) \sum_{r=1}^{s-2} T^{-r/2} |g_r(a_T)| + O[T^{-(s-1)/2}]. \quad (40)$$

But $g_r(a_T)$ is a polynomial of degree $3r - 1$ in a_T , which is odd for even r , and even for odd r . For $r = 1$ and $r = 2$ we have

$$|g_1(x)| \leq |g_{11}| + |g_{12}| |x|^2, \text{ and } |g_2(x)| \leq |g_{21}| |x| + |g_{22}| |x|^3 + |g_{23}| |x|^5,$$

where g_{ij} are fixed coefficients that depend on the cumulants of $\boldsymbol{\xi}$. Result (17) now follows from (40) by separating the constant terms of $g_r(a_T)$ from the powers of a_T . Similarly, using (16) for $a_T > 0$ we have

$$\Pr(z_{ij,T} \leq -a_T) = \Phi(-a_T) - \sum_{r=1}^{s-2} T^{-r/2} g_r(-a_T) \phi(a_T) + O[T^{-(s-1)/2}],$$

which upon using (A.1) yields

$$\Pr(z_{ij,T} \leq -a_T) \leq \frac{1}{2} \exp\left(-\frac{a_T^2}{2}\right) + (2\pi)^{-1/2} \exp\left(-\frac{a_T^2}{2}\right) \sum_{r=1}^{s-2} T^{-r/2} |g_r(-a_T)| + O[T^{-(s-1)/2}],$$

and result (18) follows. ■

Proof of Theorem 1. First we note that (see Horn and Johnson (1985) - p.297)

$$\|\tilde{\mathbf{R}} - \mathbf{R}\| \leq \|\tilde{\mathbf{R}} - \mathbf{R}\|_\infty = \max_{1 \leq i \leq N} \sum_j |\tilde{\rho}_{ij,T} - \rho_{ij}|,$$

where

$$\tilde{\rho}_{ij,T} = \hat{\rho}_{ij,T} I[|\hat{\rho}_{ij,T}| > \theta(N, T)], \quad i = 1, 2, \dots, N-1, \quad j = i+1, \dots, N,$$

$\theta(N, T) = T^{-1/2} c_p(N)$, and $c_p(N) = \Phi^{-1}\left(1 - \frac{p}{2f(N)}\right) > 0$. Note that $\theta(N, T) > 0$, and $\theta(N, T) = o(1)$ by assumption. Let $\rho_{ij}^* = \rho_{ij} I[|\rho_{ij}| > \theta(N, T)]$, and further note that

$$\max_i \sum_j |\tilde{\rho}_{ij,T} - \rho_{ij}| \leq \max_i \sum_j |\tilde{\rho}_{ij,T} - \rho_{ij}^*| + \max_i \sum_j |\rho_{ij}^* - \rho_{ij}|, \quad (41)$$

where to simplify the notation we will be using \max_i for $\max_{1 \leq i \leq N}$. We begin with the second term of (41) and write

$$\begin{aligned} \max_i \sum_j |\rho_{ij}^* - \rho_{ij}| &= \max_i \sum_j |\rho_{ij} I[|\rho_{ij}| > \theta(N, T)] - \rho_{ij}| = \\ &\leq \max_i \sum_j |\rho_{ij}| I[|\rho_{ij}| \leq \theta(N, T)]. \end{aligned}$$

But $|\rho_{ij}| I[|\rho_{ij}| \leq \theta(N, T)] \leq \theta(N, T)$, and hence, in view of (3) we have

$$\max_i \sum_j |\rho_{ij}^* - \rho_{ij}| < K\theta(N, T) \max_i \left(\sum_{j, \rho_{ij} \neq 0} 1 \right) = O[\theta(N, T) m_N]. \quad (42)$$

Consider now the first term of (41), and following Bickel and Levina (2008a) note that

$$\begin{aligned}
\max_i \sum_j |\tilde{\rho}_{ij,T} - \rho_{ij}^*| &\leq \max_i \sum_j |\hat{\rho}_{ij,T}| I [|\hat{\rho}_{ij,T}| > \theta(N, T), |\rho_{ij}| < \theta(N, T)] \\
&\quad + \max_i \sum_j |\rho_{ij}| I [|\hat{\rho}_{ij,T}| < \theta(N, T), |\rho_{ij}| > \theta(N, T)] \\
&\quad + \max_i \sum_j |\hat{\rho}_{ij,T} - \rho_{ij}| I [|\hat{\rho}_{ij,T}| > \theta(N, T), |\rho_{ij}| > \theta(N, T)] \\
&= \mathcal{A} + \mathcal{B} + \mathcal{C}.
\end{aligned} \tag{43}$$

Starting with \mathcal{C} we have

$$\mathcal{C} = \max_i \sum_j |\hat{\rho}_{ij,T} - \rho_{ij}| I [|\hat{\rho}_{ij,T}| > \theta(N, T), |\rho_{ij}| > \theta(N, T)].$$

But $I [|\hat{\rho}_{ij,T}| > \theta(N, T), |\rho_{ij}| > \theta(N, T)] \leq I [|\rho_{ij}| > \theta(N, T)]$ and also $I (|\rho_{ij}| > \theta(N, T) | \rho_{ij} = 0) = 0$. Hence

$$\begin{aligned}
\mathcal{C} &\leq \max_{ij} |\hat{\rho}_{ij,T} - \rho_{ij}| \max_i \sum_j I [|\rho_{ij}| > \theta(N, T)] \\
&\leq \max_{ij} |\hat{\rho}_{ij,T} - \rho_{ij}| \max_i \sum_j I [|\rho_{ij}| > \theta(N, T) | \rho_{ij} \neq 0] \\
&\quad + \max_{ij} |\hat{\rho}_{ij,T} - \rho_{ij}| \max_i \sum_j I [|\rho_{ij}| > \theta(N, T) | \rho_{ij} = 0] \\
&= \max_{ij} |\hat{\rho}_{ij,T} - \rho_{ij}| \max_i \sum_j I [|\rho_{ij}| > \theta(N, T) | \rho_{ij} \neq 0] \leq \max_{ij} |\hat{\rho}_{ij,T} - \rho_{ij}| m_N.
\end{aligned} \tag{44}$$

However, using (A.3) of Lemma 3 in the online supplement and noting that $c_p^2(N) = T\theta^2(N, T)$, we have

$$\begin{aligned}
\sup_{ij} \Pr [|\hat{\rho}_{ij,T} - \rho_{ij}| > \theta(N, T)] &\leq K e^{-\frac{1}{2} \frac{c_p^2(N)}{K_v}} + O \left(T^{-\frac{(s-2)}{2}} \left[\frac{c_p^2(N)}{K_v} \right]^{\frac{3(s-2)-1}{2}} e^{-\frac{1}{2} \frac{c_p^2(N)}{K_v}} \right) \\
&\quad + O(T^{-(s-1)/2}),
\end{aligned}$$

where $K_v = \sup_{ij} K_v(\boldsymbol{\theta}_{ij}) < K$, and $K_v(\boldsymbol{\theta}_{ij})$ is defined by (12), with $K_v(\boldsymbol{\theta}_{ij}) > 0$. By the

first-order Bonferroni inequality we have

$$\Pr \left[\max_{ij} |\hat{\rho}_{ij,T} - \rho_{ij}| > \theta(N, T) \right] \leq KN^2 e^{-\frac{1}{2} \frac{c_p^2(N)}{K_v}} + O \left(N^2 T^{-\frac{(s-2)}{2}} \left[\frac{c_p^2(N)}{K_v} \right]^{\frac{3(s-2)-1}{2}} e^{-\frac{1}{2} \frac{c_p^2(N)}{K_v}} \right) + O(N^2 T^{-(s-1)/2}),$$

which can also be written as (noting that the middle term of the above is dominated by the first term)

$$\Pr \left[\max_{ij} |\hat{\rho}_{ij,T} - \rho_{ij}| > \theta(N, T) \right] = O \left(N^2 e^{-\frac{1}{2} \frac{c_p^2(N)}{K_v}} \right) + O(N^2 T^{-(s-1)/2}). \quad (45)$$

Also, using result (b) of Lemma 2 in the online supplement, $e^{-\frac{1}{2} \frac{c_p^2(N)}{K_v}} = O(N^{-\delta/K_v})$, and therefore

$$\Pr \left[\max_{ij} |\hat{\rho}_{ij,T} - \rho_{ij}| > \theta(N, T) \right] = O(N^{2-\delta/K_v}) + O(N^2 T^{-(s-1)/2}). \quad (46)$$

Since $T = O(N^d)$, then it follows that

$$\max_{ij} |\hat{\rho}_{ij,T} - \rho_{ij}| = O_p[\theta(N, T)], \quad (47)$$

so long as $\delta > 2K_v$ and $d > 4/(s-1)$. Using this result in (44) now yields

$$\mathcal{C} = O_p[m_N \theta(N, T)]. \quad (48)$$

For \mathcal{A} we have

$$\begin{aligned} \mathcal{A} &= \max_i \sum_j |\hat{\rho}_{ij,T}| I [|\hat{\rho}_{ij,T}| > \theta(N, T), |\rho_{ij}| < \theta(N, T)] \\ &\leq \max_i \sum_j |\hat{\rho}_{ij,T} - \rho_{ij}| I [|\hat{\rho}_{ij,T}| > \theta(N, T), |\rho_{ij}| < \theta(N, T)] + \max_i \sum_j |\rho_{ij}| I [|\rho_{ij}| < \theta(N, T)] \\ &\leq \mathcal{A}_1 + \mathcal{A}_2, \end{aligned}$$

where

$$\mathcal{A}_2 = \max_i \sum_j |\rho_{ij}| I [|\rho_{ij}| < \theta(N, T)] = O[\theta(N, T) m_N]. \quad (49)$$

Also, for any $\gamma \in (0, 1)$, we have

$$\begin{aligned} \mathcal{A}_1 &= \max_i \sum_j |\hat{\rho}_{ij,T} - \rho_{ij}| I [|\hat{\rho}_{ij,T}| > \theta(N, T), |\rho_{ij}| < \theta(N, T)] \\ &\leq \max_i \sum_j |\hat{\rho}_{ij,T} - \rho_{ij}| I [|\hat{\rho}_{ij,T}| > \theta(N, T), |\rho_{ij}| \leq \gamma\theta(N, T)] \\ &\quad + \max_i \sum_j |\hat{\rho}_{ij,T} - \rho_{ij}| I [|\hat{\rho}_{ij,T}| > \theta(N, T), \gamma\theta(N, T) < |\rho_{ij}| < \theta(N, T)] \\ &\leq \mathcal{A}_{11} + \mathcal{A}_{12}, \end{aligned}$$

where

$$\mathcal{A}_{11} = \max_{ij} |\hat{\rho}_{ij,T} - \rho_{ij}| \max_i \sum_j I [|\hat{\rho}_{ij,T} - \rho_{ij}| > (1 - \gamma)\theta(N, T)], \quad (50)$$

$$\mathcal{A}_{12} = \max_{ij} |\hat{\rho}_{ij,T} - \rho_{ij}| \max_i \sum_j I [|\hat{\rho}_{ij,T}| > \theta(N, T), \gamma\theta(N, T) < |\rho_{ij}| < \theta(N, T)]. \quad (51)$$

But

$$\Pr \left[\max_i \sum_j I [|\hat{\rho}_{ij,T} - \rho_{ij}| > (1 - \gamma)\theta(N, T)] > 0 \right] = \Pr \left[\max_{ij} |\hat{\rho}_{ij,T} - \rho_{ij}| > (1 - \gamma)\theta(N, T) \right], \quad (52)$$

and by (45) we have,

$$\Pr \left[\max_{ij} |\hat{\rho}_{ij,T} - \rho_{ij}| > (1 - \gamma)\theta(N, T) \right] = O \left(N^2 e^{-\frac{1}{2} \frac{(1-\gamma)^2 c_p^2(N)}{K_v}} \right) + O(N^2 T^{-(s-1)/2}).$$

Using a similar line of reasoning as above there exist $\delta > 2K_v/(1 - \gamma)^2$ and $d > 4/(s - 1)$

such that

$$\Pr \left[\max_{ij} |\hat{\rho}_{ij,T} - \rho_{ij}| > (1 - \gamma)\theta(N, T) \right] \rightarrow 0.$$

Using this result in conjunction with (52) and (47) in (50) it follows that $\mathcal{A}_{11} = O_p[m_N \theta(N, T)]$.

For \mathcal{A}_{12} , we first note that

$$\begin{aligned}
& \max_i \sum_j I [|\hat{\rho}_{ij,T}| > \theta(N, T), \gamma\theta(N, T) < |\rho_{ij}| < \theta(N, T)] \\
& \leq \max_i \sum_j I [\gamma\theta(N, T) < |\rho_{ij}| < \theta(N, T)] \\
& = \max_i \sum_j I \{[\gamma\theta(N, T) < |\rho_{ij}| < \theta(N, T)] \mid \rho_{ij} \neq 0\} \leq m_N,
\end{aligned}$$

which gives $\mathcal{A}_{12} = O_p[m_N\theta(N, T)]$, and together with the result for \mathcal{A}_{11} we have $\mathcal{A}_1 = O_p[\theta(N, T)m_N]$. Overall using (49) we obtain

$$\mathcal{A} = O_p[m_N\theta(N, T)]. \quad (53)$$

Finally, for \mathcal{B} we have

$$\begin{aligned}
\mathcal{B} & = \max_i \sum_j |\rho_{ij}| I [|\hat{\rho}_{ij,T}| < \theta(N, T), |\rho_{ij}| > \theta(N, T)] \\
& \leq \max_i \sum_j |\hat{\rho}_{ij,T} - \rho_{ij}| I [|\hat{\rho}_{ij,T}| < \theta(N, T), |\rho_{ij}| > \theta(N, T)] \\
& \quad + \max_i \sum_j |\hat{\rho}_{ij,T}| I [|\hat{\rho}_{ij,T}| < \theta(N, T), |\rho_{ij}| > \theta(N, T)] \\
& = \mathcal{B}_1 + \mathcal{B}_2.
\end{aligned}$$

But as before

$$\begin{aligned}
\mathcal{B}_1 & = \max_i \sum_j |\hat{\rho}_{ij,T} - \rho_{ij}| I [|\hat{\rho}_{ij,T}| < \theta(N, T), |\rho_{ij}| > \theta(N, T)] \\
& \leq \max_{ij} |\hat{\rho}_{ij,T} - \rho_{ij}| \max_i \sum_j I [|\hat{\rho}_{ij,T}| < \theta(N, T), |\rho_{ij}| > \theta(N, T)] \\
& = \max_{ij} |\hat{\rho}_{ij,T} - \rho_{ij}| \max_i \sum_j I \{[|\hat{\rho}_{ij,T}| < \theta(N, T), |\rho_{ij}| > \theta(N, T)] \mid \rho_{ij} \neq 0\} \\
& \leq m_N \max_{ij} |\hat{\rho}_{ij,T} - \rho_{ij}| = O_p[m_N\theta(N, T)], \quad (54)
\end{aligned}$$

$$\begin{aligned}
\mathcal{B}_2 &= \max_i \sum_j |\hat{\rho}_{ij,T}| I \left[|\hat{\rho}_{ij,T}| < \theta(N, T), \quad |\rho_{ij}| > \theta(N, T) \right] \\
&\leq \max_i \sum_{j, \rho_{ij} \neq 0} |\hat{\rho}_{ij,T}| I \left[|\hat{\rho}_{ij,T}| < \theta(N, T) \right] \\
&\leq \theta(N, T) m_N.
\end{aligned}$$

Hence $\mathcal{B}_2 = O_p[\theta(N, T) m_N]$, which in conjunction with (54) yields,

$$\mathcal{B} = O_p[\theta(N, T) m_N]. \quad (55)$$

Substituting results from (53), (55) and (48) in (43), and using the outcome with (42) in (41) we obtain

$$\left\| \tilde{\mathbf{R}} - \mathbf{R} \right\| = O_p[\theta(N, T) m_N] = O_p\left(\frac{m_N c_p(N)}{\sqrt{T}}\right),$$

as required. ■

Proof of Theorem 2. Consider the squared Frobenius norm,

$$\left\| \tilde{\mathbf{R}} - \mathbf{R} \right\|_F^2 = \sum_{i \neq j} \sum (\tilde{\rho}_{ij,T} - \rho_{ij})^2,$$

and recall that

$$\tilde{\rho}_{ij,T} - \rho_{ij} = (\hat{\rho}_{ij,T} - \rho_{ij}) I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \right) - \rho_{ij} \left[1 - I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \right) \right].$$

Hence

$$\begin{aligned}
(\tilde{\rho}_{ij,T} - \rho_{ij})^2 &= (\hat{\rho}_{ij,T} - \rho_{ij})^2 I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \right) + \rho_{ij}^2 \left[1 - I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \right) \right]^2 \\
&\quad - 2\rho_{ij} (\hat{\rho}_{ij,T} - \rho_{ij}) I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \right) \left[1 - I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \right) \right].
\end{aligned}$$

However,

$$I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \right) \left[1 - I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \right) \right] = 0,$$

and

$$\left[1 - I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \right) \right]^2 = 1 - I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \right).$$

Therefore, we have

$$\begin{aligned} \sum_{i \neq j} \sum (\tilde{\rho}_{ij,T} - \rho_{ij})^2 &= \sum_{i \neq j} \sum (\hat{\rho}_{ij,T} - \rho_{ij})^2 I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \right) \\ &\quad + \sum_{i \neq j} \sum \rho_{ij}^2 \left[1 - I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \right) \right] \\ &= \sum_{i \neq j} \sum (\hat{\rho}_{ij,T} - \rho_{ij})^2 I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \right) \\ &\quad + \sum_{i \neq j} \sum \rho_{ij}^2 I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| \leq c_p(N) \right). \end{aligned}$$

Taking expectations we have the following decomposition

$$E \left(\left\| \tilde{\mathbf{R}} - \mathbf{R} \right\|_F^2 \right) = \sum_{i \neq j} \sum E (\tilde{\rho}_{ij,T} - \rho_{ij})^2 = \mathcal{D} + \mathcal{E} + \mathcal{F}, \quad (56)$$

where

$$\begin{aligned} \mathcal{D} &= \sum_{i \neq j, \rho_{ij} \neq 0} \sum \rho_{ij}^2 E \left[I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| \leq c_p(N) \mid \rho_{ij} \neq 0 \right) \right], \\ \mathcal{E} &= \sum_{i \neq j, \rho_{ij} \neq 0} \sum E \left[(\hat{\rho}_{ij,T} - \rho_{ij})^2 I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} \neq 0 \right) \right], \\ \mathcal{F} &= \sum_{i \neq j, \rho_{ij} = 0} \sum E \left[\hat{\rho}_{ij,T}^2 I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} = 0 \right) \right]. \end{aligned}$$

Consider now the orders of the above three terms in turn, starting with \mathcal{D} . We have $\rho_{\min} = \min_{ij} (|\rho_{ij}|, \rho_{ij} \neq 0)$ and $\rho_{\max} = \max_{ij} (|\rho_{ij}|, \rho_{ij} \neq 0)$ such that $\rho_{\max} < 1$. Then

$$\begin{aligned} \mathcal{D} &\leq \rho_{\max}^2 N m_N \sup_{ij} E \left[I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| \leq c_p(N) \mid \rho_{ij} \neq 0 \right) \right] \\ &= \rho_{\max}^2 N m_N \sup_{ij} \Pr \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| \leq c_p(N) \mid \rho_{ij} \neq 0 \right), \end{aligned}$$

and using (A.6) of Lemma 3 in the online supplement for $|\rho_{ij}| > c_p(N)/\sqrt{T}$, we have

$$\mathcal{D} \leq K \rho_{\max}^2 N m_N K e^{-\frac{T(\rho_{\min} - T^{-1/2} c_p(N))^2}{2K_v}} \left[1 + O\left(T^{\frac{2(s-2)-1}{2}}\right) \right] + O(N m_N T^{-(s-1)/2}),$$

where $K_v = \sup_{ij} |K_v(\boldsymbol{\theta}_{ij})| < K$. By assumption $T^{-1/2} c_p(N) = o(1)$, and since $\rho_{\min} > 0$, then the first term of the above will tend to zero with N and $T \rightarrow \infty$. Therefore, \mathcal{D} is of order $O(N m_N N^{-d(s-1)/2}) = O(N^{1+\vartheta-d(s-1)/2})$, and \mathcal{D} tends to zero as $N \rightarrow \infty$, for values of $d > 2(1+\vartheta)/(s-1)$ and under (21), where by assumption $0 \leq \vartheta < 1/2$.

Consider now \mathcal{E} . Recalling that $\hat{\rho}_{ij,T} = \omega_{ij,T} z_{ij,T} + \rho_{ij,T}$ we have the following decomposition of \mathcal{E} , $\mathcal{E} = \mathcal{E}_1 + \mathcal{E}_2 + 2\mathcal{E}_3$, where

$$\begin{aligned} \mathcal{E}_1 &= \sum_{i \neq j, \rho_{ij} \neq 0} \sum \omega_{ij,T}^2 E \left[z_{ij,T}^2 I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} \neq 0 \right) \right], \\ \mathcal{E}_2 &= \sum_{i \neq j, \rho_{ij} \neq 0} \sum (\rho_{ij,T} - \rho_{ij})^2 E \left[I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} \neq 0 \right) \right], \\ \mathcal{E}_3 &= \sum_{i \neq j, \rho_{ij} \neq 0} \sum (\rho_{ij,T} - \rho_{ij}) \omega_{ij,T} E \left[z_{ij,T} I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} \neq 0 \right) \right]. \end{aligned}$$

Again, using (9) and (10),

$$\omega_{ij,T}^2 = \frac{K_v(\boldsymbol{\theta}_{ij})}{T} + O(T^{-2}), \quad (57)$$

$$(\rho_{ij,T} - \rho_{ij})^2 = \frac{K_m^2(\boldsymbol{\theta}_{ij})}{T^2} + O(T^{-3}), \quad (58)$$

$$(\rho_{ij,T} - \rho_{ij}) \omega_{ij,T} = \frac{K_v^{1/2}(\boldsymbol{\theta}_{ij}) K_m(\boldsymbol{\theta}_{ij})}{T^{3/2}} + O(T^{-5/2}). \quad (59)$$

$$\begin{aligned} \mathcal{E}_1 &= \sum_{i \neq j, \rho_{ij} \neq 0} \sum \omega_{ij,T}^2 E \left[z_{ij,T}^2 I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} \neq 0 \right) \right] \\ &\leq \frac{N m_N}{T} \left[\sup_{ij} K_v(\boldsymbol{\theta}_{ij}) + O(T^{-1}) \right] \sup_{ij} E \left[z_{ij,T}^2 I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} \neq 0 \right) \right]. \end{aligned}$$

Using (A.7) in Lemma 4 in the online supplement with $r = 2$, and noting that $\sup_{ij} K_v(\boldsymbol{\theta}_{ij}) <$

K , then $\mathcal{E}_1 = O(m_N N/T)$. Similarly, since $E \left[I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} \neq 0 \right) \right] \leq 1$, we have

$$\begin{aligned} \mathcal{E}_2 &= \sum_{i \neq j, \rho_{ij} \neq 0} \sum (\rho_{ij,T} - \rho_{ij})^2 E \left[I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} \neq 0 \right) \right] \\ &\leq N m_N \left[\frac{K_m^2(\boldsymbol{\theta}_{ij})}{T^2} + O(T^{-3}) \right] = O\left(\frac{N m_N}{T^2}\right). \end{aligned}$$

Also, using (A.7) in Lemma 4 in the online supplement with $r = 1$, and using (59) we have

$$\begin{aligned} \mathcal{E}_3 &= \sum_{i \neq j, \rho_{ij} \neq 0} \sum (\rho_{ij,T} - \rho_{ij}) \omega_{ij,T} E \left[z_{ij,T} I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} \neq 0 \right) \right] \\ &\leq K N m_N \left[\frac{\sup_{ij} K_v^{1/2}(\boldsymbol{\theta}_{ij}) K_m(\boldsymbol{\theta}_{ij})}{T^{3/2}} + O(T^{-5/2}) \right] = O\left(\frac{N m_N}{T^{3/2}}\right). \end{aligned}$$

Therefore, overall $\mathcal{E} = O\left(\frac{m_N N}{T}\right)$. Consider now the following decomposition of \mathcal{F} , in (56):

$$\begin{aligned} \mathcal{F} &= \sum_{i \neq j, \rho_{ij} = 0} \sum E \left[\hat{\rho}_{ij,T}^2 I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} = 0 \right) \right] \\ &= \sum_{i \neq j, \rho_{ij} = 0} \sum \omega_{ij,T}^2 E \left[z_{ij,T}^2 I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} = 0 \right) \right] \\ &\quad + \sum_{i \neq j, \rho_{ij} = 0} \sum \rho_{ij,T}^2 E \left[I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} = 0 \right) \right] \\ &\quad + 2 \sum_{i \neq j, \rho_{ij} = 0} \sum \rho_{ij,T} \omega_{ij,T} E \left[z_{ij,T} I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} = 0 \right) \right] \\ &= \mathcal{F}_1 + \mathcal{F}_2 + \mathcal{F}_3. \end{aligned}$$

Consider \mathcal{F}_1 and using (57) note that

$$\mathcal{F}_1 \leq \frac{N(N - m_N - 1)}{T} \left[\sup_{ij} K_v(\boldsymbol{\theta}_{ij}) + O(T^{-1}) \right] \sup_{ij} E \left[z_{ij,T}^2 I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} = 0 \right) \right].$$

Then using (A.8) in Lemma 4 of the online supplement with $r = 2$, we have

$$\mathcal{F}_1 \leq K \frac{N(N - m_N - 1)}{T} \left[\sup_{ij} K_v(\boldsymbol{\theta}_{ij}) + O(T^{-1}) \right] \times \left[K e^{-\frac{1}{2} \frac{c_p^2(N)}{\varphi_{\max}}} + O \left(T^{-\frac{(s-2)}{2}} \left[\frac{c_p^2(N)}{\varphi_{\max}} \right]^{\frac{3(s-2)-1}{2}} e^{-\frac{1}{2} \frac{c_p^2(N)}{\varphi_{\max}}} \right) + O(T^{-(s-1)/2}) \right],$$

where $\varphi_{\max} = \sup_{ij} \varphi_{ij} > 0$. Now noting that $\sup_{ij} |K_v(\boldsymbol{\theta}_{ij})| = K_v < \infty$, and $m_N/N = o(1)$, then

$$\begin{aligned} \mathcal{F}_1 &= O \left(N^2 T^{-1} e^{-\frac{1}{2} \frac{c_p^2(N)}{\varphi_{\max}}} \right) + O \left(N^2 T^{-1} T^{-(s-1)/2} \right) \\ &= O \left(e^{-\left(\frac{1}{2\varphi_{\max}}\right) \ln N \left[\frac{c_p^2(N)}{\ln N} - 2(2-d)\varphi_{\max} \right]} \right) + O \left(N^{2-d(s-1)/2-d} \right). \end{aligned}$$

Therefore, since $\lim_{N \rightarrow \infty} c_p^2(N)/\ln(N) = 2\delta$ (see result (b) of Lemma 2 in the online supplement), then $\mathcal{F}_1 \rightarrow 0$, as $N \rightarrow \infty$, if $\delta > (2-d)\varphi_{\max}$, and $d > 4/(s+1)$. Similarly, using (59), we have

$$\begin{aligned} \mathcal{F}_2 &= \sum_{i \neq j} \sum_{\rho_{ij}=0} \rho_{ij,T}^2 E \left[I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} = 0 \right) \right] \\ &\leq N(N - m_N - 1) \left\{ \frac{\sup_{ij} [K_m^2(\boldsymbol{\theta}_{ij}) \mid \rho_{ij} = 0]}{T^2} + O(T^{-3}) \right\} \sup_{ij} E \left[I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} = 0 \right) \right]. \end{aligned}$$

Now using (A.4) of Lemma 3 in the online supplement we have

$$\mathcal{F}_2 \leq K \frac{N(N - m_N - 1) [\psi_{\max}^2 + O(T^{-1})]}{T^2} \times \left[K e^{-\frac{1}{2} \frac{c_p^2(N)}{\varphi_{\max}}} + O \left(T^{-\frac{(s-2)}{2}} \left[\frac{c_p^2(N)}{\varphi_{\max}} \right]^{\frac{3(s-2)-1}{2}} e^{-\frac{1}{2} \frac{c_p^2(N)}{\varphi_{\max}}} \right) + O(T^{-(s-1)/2}) \right],$$

where $\psi_{\max} = \sup_{ij} \psi_{ij} < K$, and ψ_{ij} is defined by (14). Once again, since $m_N/N = o(1)$, then

$$\mathcal{F}_2 = O \left(e^{2(1-d) \ln N - \frac{1}{2} \frac{c_p^2(N)}{\varphi_{\max}}} \right) + O \left(N^2 T^{-2} T^{-(s-1)/2} \right),$$

and following similar arguments as above, it follows that $\mathcal{F}_2 \rightarrow 0$ as $N \rightarrow \infty$, if $\delta > 2(1-d)\varphi_{\max}$, and $d > 4/(s+3)$. Both of these conditions are met if $\delta > (2-d)\varphi_{\max}$ and $d > 4/(s+1)$, since $(2-d)\varphi_{\max} > 2(1-d)\varphi_{\max}$, and $s > 0$. Consider now \mathcal{F}_3 and, using (9) and (10) evaluated at $\rho_{ij} = 0$, note that

$$\begin{aligned} \mathcal{F}_3 &= \sum_{i \neq j} \sum_{\rho_{ij}=0} \rho_{ij,T} \omega_{ij,T} E \left[z_{ij,T} I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} = 0 \right) \right] \\ &\leq \sum_{i \neq j} \sum_{\rho_{ij}=0} |\rho_{ij,T}| |\omega_{ij,T}| E \left[|z_{ij,T}| I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} = 0 \right) \right] \\ &\leq N(N - m_N - 1) \left[\frac{\sup_{ij} |\psi_{ij}|}{T} + O(T^{-2}) \right] \left[\frac{\sup_{ij} \sqrt{\varphi_{ij}}}{\sqrt{T}} + O(T^{-3/2}) \right] \\ &\quad \times \sup_{ij} E \left[|z_{ij,T}| I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| \leq c_p(N) \mid \rho_{ij} = 0 \right) \right]. \end{aligned}$$

Further using (A.8) in Lemma 4 of the online supplement with $r = 1$, we obtain (recall that $\sup_{ij} |\psi_{ij}| < K$ and $\sup_{ij} \sqrt{\varphi_{ij}} < K$)

$$\begin{aligned} \mathcal{F}_3 &\leq K \frac{N(1 - m_N/N - 1/N)}{T^{3/2}} \times \\ &\quad \left[K e^{-\frac{1}{2} \frac{c_p^2(N)}{\varphi_{\max}}} + O \left(T^{-\frac{(s-2)}{2}} \left[\frac{c_p^2(N)}{\varphi_{\max}} \right]^{\frac{3(s-2)-1}{2}} e^{-\frac{1}{2} \frac{c_p^2(N)}{\varphi_{\max}}} \right) + O(T^{-(s-1)/2}) \right], \end{aligned}$$

which establishes that $\mathcal{F}_3 \rightarrow 0$, as $N \rightarrow \infty$, if $\delta > (2-d)\varphi_{\max}$ and $d > 4/(s+1)$ (using the same type of derivations as above). Therefore, overall $\mathcal{F} \rightarrow 0$, under the same conditions.

Using this result, together with the results obtained for \mathcal{D} and \mathcal{E} above, in (56) we obtain $E \left\| \tilde{\mathbf{R}} - \mathbf{R} \right\|_F^2 = O\left(\frac{m_N N}{T}\right)$, and (26) follows as required. Also by the Markov inequality

$$\Pr \left(\sqrt{\frac{T}{m_N N}} \left\| \tilde{\mathbf{R}} - \mathbf{R} \right\|_F \geq \varepsilon \right) = \Pr \left(\frac{T}{m_N N} \left\| \tilde{\mathbf{R}} - \mathbf{R} \right\|_F^2 \geq \varepsilon^2 \right) \leq \frac{\frac{T}{m_N N} E \left\| \tilde{\mathbf{R}} - \mathbf{R} \right\|_F^2}{\varepsilon^2} \leq \frac{K}{\varepsilon^2},$$

for some small $\varepsilon > 0$. Hence,

$$\sqrt{\frac{T}{m_N N}} \left\| \tilde{\mathbf{R}} - \mathbf{R} \right\|_F = O_p(1),$$

and result (27) follows. ■

Proof of Theorem 3. Recall that $T = c_d N^d$, $c_d > 0$, and consider first the FPR_N statistic given by (31) which can be written equivalently as

$$FPR_N = |FPR_N| = \frac{\sum_{i \neq j} \sum I(|\hat{\rho}_{ij,T}| > T^{-1/2} c_p(N) | \rho_{ij} = 0)}{N(N - m_N - 1)}. \quad (60)$$

Note that the elements of FPR_N are either 0 or 1 and so $|FPR_N| = FPR_N$. Taking the expectation of (60) we have

$$E |FPR_N| = \frac{\sum_{i \neq j} \sum \Pr(|\hat{\rho}_{ij,T}| > T^{-1/2} c_p(N) | \rho_{ij} = 0)}{N(N - m_N - 1)} \leq \sup_{ij} \Pr(|\hat{\rho}_{ij,T}| > T^{-1/2} c_p(N) | \rho_{ij} = 0).$$

Hence, using (A.4) in Lemma 3 of the online supplement we have

$$E |FPR_N| = O\left(e^{-\frac{1}{2} \frac{c_p^2(N)}{\varphi_{\max}}}\right) + O(T^{-(s-1)/2}),$$

where $\varphi_{\max} = \sup_{ij} \varphi_{ij} < K$ by Assumption 2 (see also Proposition 1). Hence, as $N \rightarrow \infty$ for any $d > 0$ (recalling that $T = O(N^d)$), $E |FPR_N| \rightarrow 0$, noting that $c_p^2(N) \rightarrow \infty$, and $\varphi_{\max} > 0$. Further, by the Markov inequality applied to $|FPR_N|$ we have, for some $\eta > 0$,

$$\Pr(|FPR_N| > \eta) \leq \frac{E(|FPR_N|)}{\eta} = O\left(e^{-\frac{1}{2} \frac{c_p^2(N)}{\varphi_{\max}}}\right) + O(T^{-(s-1)/2}). \quad (61)$$

It therefore follows that $\lim_{N \rightarrow \infty} \Pr(|FPR_N| > \eta) = 0$, and $FPR_N \xrightarrow{p} 0$ as $N \rightarrow \infty$, for any $d > 0$. For almost sure convergence by the Borel Cantelli lemma it suffices to show that

$$\sum_{N=1}^{\infty} \Pr(|FPR_N| > \eta) < \infty. \quad (62)$$

From result (b) of Lemma 2 we have

$$e^{-\frac{1}{2} \frac{c_p^2(N)}{\varphi_{\max}}} = O\left(N^{-\frac{\delta}{\varphi_{\max}}}\right), \quad (63)$$

and from (61) it follows that (for $\eta > 0$)

$$\Pr(|FPR_N| > \eta) = O\left(N^{-\frac{\delta}{\varphi_{\max}}}\right) + O\left(N^{-d(s-1)/2}\right). \quad (64)$$

Hence (62) holds if $\sum_{N=1}^{\infty} N^{-\frac{\delta}{\varphi_{\max}}}$ and $\sum_{N=1}^{\infty} N^{-d(s-1)/2}$ converge, which is ensured if $\delta > \varphi_{\max}$, and $d > 2/(s-1)$, which establishes that $FPR_N \xrightarrow{a.s.} 0$, as $N \rightarrow \infty$.

Consider now the TPR_N statistic given by (30) and note that

$$TPR_N = \frac{\sum_{i \neq j} \sum I(\tilde{\rho}_{ij,T} \neq 0, \text{ and } \rho_{ij} \neq 0)}{\sum_{i \neq j} \sum I(\rho_{ij} \neq 0)}.$$

Hence

$$X_N = 1 - TPR_N = \frac{\sum_{i \neq j} \sum I(\tilde{\rho}_{ij,T} = 0, \text{ and } \rho_{ij} \neq 0)}{Nm_N}.$$

Since $|X_N| = X_N$, then

$$E|X_N| = E(X_N) = \frac{\sum_{i \neq j} \sum \Pr\left(|\hat{\rho}_{ij,T}| < T^{-1/2}c_p(N) | \rho_{ij} \neq 0\right)}{Nm_N} \leq \sup_{ij} \Pr\left(|\hat{\rho}_{ij,T}| < T^{-1/2}c_p(N) | \rho_{ij} \neq 0\right).$$

From (A.6) of Lemma 3 of the online supplement we further have that

$$\begin{aligned} E|X_N| &\leq \sup_{ij} \Pr\left(|\hat{\rho}_{ij,T}| < T^{-1/2}c_p(N) | \rho_{ij} \neq 0\right) \\ &\leq K e^{-\frac{T(\rho_{\min} - T^{-1/2}c_p(N))^2}{2K_v}} \left[1 + O\left(T^{\frac{2(s-2)-1}{2}}\right)\right] + O\left(T^{-(s-1)/2}\right), \end{aligned}$$

where $\rho_{\min} = \min_{ij} (|\rho_{ij}|, \rho_{ij} \neq 0) > 0$, and $K_v = \sup_{ij} K_v(\boldsymbol{\theta}_{ij}) < K$. Hence, since by assumption $T^{-1/2}c_p(N) = o(1)$, and $T = c_d N^d$, with $c_d, d > 0$, it follows that $\lim_{N \rightarrow \infty} E|X_N| =$

0, as $N \rightarrow \infty$. Further, by the Markov inequality, $\Pr(|X_N| > \eta) \leq \frac{E|X_N|}{\eta}$ for some $\eta > 0$, and it follows that

$$\Pr(|TPR_N - 1| > \eta) \leq \frac{E(|TPR_N - 1|)}{\eta} = O\left(e^{-\frac{T(\rho_{\min} - T^{-1/2}c_p(N))^2}{2K_v}}\right) + O(N^{-d(s-1)/2}). \quad (65)$$

Once again since by assumption $T^{-1/2}c_p(N) = o(1)$, $d > 0$, and $\rho_{\min} > 0$, then for any $\eta > 0$, $\lim_{N \rightarrow \infty} \Pr(|TPR_N - 1| > \eta) = 0$, and $TPR_N \xrightarrow{p} 1$, as $N \rightarrow \infty$. For almost sure convergence it is further required that

$$\sum_{N=1}^{\infty} \Pr(|TPR_N - 1| > \eta) < \infty. \quad (66)$$

From (65) we have that

$$\sum_{N=1}^{\infty} \Pr(|TPR_N - 1| > \eta) = O\left(\sum_{N=1}^{\infty} a_N\right) + O\left(\sum_{N=1}^{\infty} b_N\right),$$

where (setting $c_d = 1$ to simplify the notations)

$$a_N = e^{-\frac{N^d}{2K_v}(\rho_{\min} - N^{-d/2}c_p(N))^2}, \text{ and } b_N = N^{-d(s-1)/2}.$$

Hence, for (66) to hold, the series $\sum_{N=1}^{\infty} a_N$ and $\sum_{N=1}^{\infty} b_N$ must converge. Using the direct comparison test for convergence of infinite series, this will be the case if

$$N^{-d(s-1)/2} \leq N^{-1-\epsilon}, \quad (67)$$

and

$$e^{-\frac{N^d}{2K_v}(\rho_{\min} - T^{-1/2}c_p(N))^2} \leq N^{-1-\epsilon} \quad (68)$$

for all $N \geq N_0$, where N_0 is some finite positive integer, since for $\epsilon > 0$, we have $\sum_{N=1}^{\infty} N^{-1+\epsilon} < \infty$

K . Condition (68) can be written equivalently as

$$\frac{1}{2K_v} (\rho_{\min} - T^{-1/2} c_p(N))^2 > (1 + \epsilon) T^{-1} \ln(N),$$

which is satisfied since by assumption $N^{-d/2} c_p(N) = T^{-1/2} c_p(N) = o(1)$, $d > 0$, $\rho_{\min} > 0$, and K_v is a bounded positive constant. Hence, under the conditions of the theorem it follows that $TPR_N \xrightarrow{a.s.} 1$ as $N \rightarrow \infty$, if $d > 2/(s-1)$.

Finally, consider the FDR statistic defined by (32), and note that

$$FDR_N = \left(\frac{N - m_N - 1}{m_N} \right) FPR_N.$$

Now noting that $\frac{(N - m_N - 1)}{m_N} = \Theta(N^{1-\vartheta})$, and using (64) we have

$$E|FDR_N| = O\left(N^{1-\vartheta} N^{-\frac{\delta}{\varphi_{\max}}}\right) + O\left(N^{1-\vartheta} N^{-d(s-1)/2}\right).$$

Hence, $\lim_{N \rightarrow \infty} E|FDR_N| = 0$, as $N \rightarrow \infty$, if $\delta > (1 - \vartheta)\varphi_{\max}$ and $d > 2(1 - \vartheta)/(s - 1)$.

Also, applying Markov inequality to $|FDR_N|$, for some $\eta > 0$ we have

$$\Pr(|FDR_N| > \eta) \leq \frac{E(|FDR_N|)}{\eta} = O\left(N^{1-\vartheta} N^{-\frac{\delta}{\varphi_{\max}}}\right) + O\left(N^{1-\vartheta} N^{-d(s-1)/2}\right). \quad (69)$$

Almost sure convergence requires

$$\sum_{N=1}^{\infty} \Pr(|FDR_N| > \eta) < \infty, \quad (70)$$

and using (69) this follows if $\delta > (2 - \vartheta)\varphi_{\max}$ and $d > 2(2 - \vartheta)/(s - 1)$, then $FDR_N \xrightarrow{a.s.} 1$ as $N \rightarrow \infty$. ■

References

- K. M. Abadir, W. Distaso, and F. Zikes. Design-free estimation of large variance matrices. *Journal of Econometrics*, 181:165–180, 2014.
- F. Abramovich, Y. Benjamini, D. Donoho, and I. Johnstone. Adapting to unknown sparsity by controlling the false discovery rate. *Annals of Statistics*, 34:584–653, 2006.
- R. N. Bhattacharya and J. K. Ghosh. On the validity of the formal Edgeworth expansion. *The Annals of Statistics*, 6:434–451, 1978.
- P. J. Bickel and E. Levina. Some theory for Fisher’s linear discriminant function, ‘naive Bayes’, and some alternatives when there are many more variables than observations. *Bernoulli*, 10:989–1010, 2004.
- P. J. Bickel and E. Levina. Covariance regularisation by thresholding. *The Annals of Statistics*, 36:2577–2604, 2008.
- P. J. Bickel and E. Levina. Regularised estimation of large covariance matrices. *The Annals of Statistics*, 36:199–227, 2008.
- C. Bonferroni. Il calcolo delle assicurazioni su gruppi di teste. *Studi in Onore del Professore Salvatore Ortu Carboni, Rome, Italy*, 1935.
- T. Cai and W. Liu. Adaptive thresholding for sparse covariance matrix estimation. *Journal of American Statistical Association*, 106:672–684, 2011.
- R. J. Carroll. Variances are not always nuisance parameters. *Biometrics*, 59:211–220, 2003.
- M. J. Daniels and R. Kass. Nonconjugate Bayesian estimation of covariance matrices in hierarchical models. *Journal of American Statistical Association*, 94:1254–1263, 1999.
- M. J. Daniels and R. Kass. Shrinkage estimators for covariance matrices. *Biometrics*, 57:1173–1184, 2001.

- A. D'Aspremont, O. Banerjee, and L. Elghaoui. First-order methods for sparse covariance selection. *SIAM Journal on Matrix Analysis and Applications*, 30:56–66, 2008.
- S. Dees, F. di Mauro, M. H. Pesaran, and L. V. Smith. Exploring the international linkages of the euro area: A Global VAR analysis. *Journal of Applied Econometrics*, 22:1–38, 2007.
- A. Delaigle, P. Hall, and J. Jin. Robustness and accuracy of methods for high dimensional data analysis based on student's t-statistic. *Journal of Royal Statistical Society, Series B*, 73:283–301, 2011.
- N. El Karoui. Operator norm consistent estimation of large dimensional sparse covariance matrices. *The Annals of Statistics*, 36:2717–2756, 2008.
- J. Fan and R. Li. Variable selection via nonconcave penalized likelihood and its oracle properties. *Journal of American Statistical Association*, 96:1348–1360, 2001.
- J. Fan, Y. Fan, and J. Lv. High dimensional covariance matrix estimation using a factor model. *Journal of Econometrics*, 147:186–197, 2008.
- J. Fan, Y. Liao, and M. Mincheva. High dimensional covariance matrix estimation in approximate factor models. *The Annals of Statistics*, 39:3320–3356, 2011.
- J. Fan, Y. Liao, and M. Mincheva. Large covariance estimation by thresholding principal orthogonal complements. *Journal of Royal Statistical Society, Series B*, 75:1–44, 2013.
- J. Fan, Y. Ke, Q. Sun, and W-X Zhou. Farmtest: Factor-adjusted robust multiple testing with approximate false discovery control. *Journal of the American Statistical Association*, *Forthcoming*, 2018.
- Y. Fang, B. Wang, and Y. Feng. Tuning parameter selection in regularised estimations of large covariance matrices. *Journal of Statistical Computation and Simulation*, 86:494–509, 2016.

- J. H. Friedman. Regularized discriminant analysis. *Journal of the American Statistical Association*, 84:165–175, 1989.
- P. Fryzlewicz. High-dimensional volatility matrix estimation via wavelets and thresholding. *Biometrika*, 100:921–938, 2013.
- A. K. Gayen. The frequency distribution of the product-moment correlation coefficient in random samples of any size drawn from non-normal universes. *Biometrika*, 38:219–247, 1951.
- P. Hall. *The Bootstrap and Edgeworth Expansion*. Springer-Verlag New York, Inc, 1992.
- P. D. Hoff. A hierarchical eigenmodel for pooled covariance estimation. *Journal of Royal Statistical Society, Series B*, 71:971–992, 2009.
- S. Holm. A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics*, 6:65–70, 1979.
- R. A. Horn and C. R. Johnson. *Matrix Analysis*. Cambridge University Press: Cambridge, 1985.
- O. Ledoit and M. Wolf. Improved estimation of the covariance matrix of stock returns with an application to portfolio selection. *Journal of Empirical Finance*, 10:603–621, 2003.
- O. Ledoit and M. Wolf. A well-conditioned estimator for large-dimensional covariance matrices. *Journal of Multivariate Analysis*, 88:365–411, 2004.
- H. Liu, L. Wang, and T. Zhao. Sparse covariance matrix estimation with eigenvalue constraints. *Journal of Computational and Graphical Statistics*, 23:439–459, 2014.
- J. Peng, P. Wang, N. Zhou, and J. Zhu. Partial correlation estimation by joint sparse regression models. *Journal of American Statistical Association*, 104:735–746, 2009.

- M. H. Pesaran, T. Schuermann, and S. M. Weiner. Modeling regional interdependencies using a Global Error Correcting Macroeconometric model. *Journal of Business and Economic Statistics*, 22:129–162, 2004.
- M. H. Pesaran. Testing weak cross-sectional dependence in large panels. *Econometric Reviews*, 34:1089–1117, 2015.
- M. Pourahmadi. Joint mean-covariance models with applications to longitudinal data: Unconstrained parameterisation. *Biometrika*, 86:677–690, 1999.
- M. Pourahmadi. Maximum likelihood estimation of generalised linear models for multivariate normal covariance matrix. *Biometrika*, 87:425–435, 2000.
- M. Pourahmadi. Covariance estimation: GLM and regularisation perspectives. *Statistical Science*, 26:369–387, 2011.
- P. J. Romano and M. Wolf. Exact and approximate stepdown methods for multiple hypothesis testing. *Journal of the American Statistical Association*, 100(469):94–108, 2005.
- P. J. Romano, A. M. Shaikh, and M. Wolf. Formalised data snooping based on generalised error rates. *Econometric Theory*, 24:404–447, 2008.
- A. J. Rothman, P. J. Bickel, E. Levina, and J. Zhu. Sparse permutation invariant covariance estimation. *Electronic Journal of Statistics*, 2:494–515, 2008.
- A. J. Rothman, E. Levina, and J. Zhu. Generalised thresholding of large covariance matrices. *Journal of American Statistical Association*, 104:177–186, 2009.
- A. J. Rothman, P. J. Bickel, E. Levina, and J. Zhu. A new approach to Cholesky-based estimation of high-dimensional covariance matrices. *Biometrika*, 97:539–550, 2010.
- A. J. Rothman. Positive definite estimators of large covariance matrices. *Biometrika*, 99:733–740, 2012.

- J. Schäfer and K. Strimmer. A shrinkage approach to large-scale covariance matrix estimation and implications for functional Genomics. *Statistical Applications in Genetics and Molecular Biology*, 4, Article 32, 2005.
- E. Sentana. The econometrics of mean-variance efficiency tests: A survey. *The Econometrics Journal*, 12:65–101, 2009.
- A. Tschuprow. *Grundbegriffe und Grundprobleme der Korrelationstheori*. Leipzig: B.G. Teubner, 1925.
- W. B. Wu and M. Pourahmadi. Banding sample covariance matrices for stationary processes. *Statistica Sinica*, 19:1755–1768, 2009.
- L. Xue, S. Ma, and H. Zou. Positive-definite ℓ_1 -penalized estimation of large covariance matrices. *Journal of the American Statistical Association*, 107:1480–1491, 2012.
- M. Yuan and Y. Lin. Model selection and estimation in the Gaussian graphical model. *Biometrika*, 94:19–35, 2007.
- W-X Zhou, K. Bose, J. Fan, and H. Liu. A new perspective on robust M-estimation: Finite sample theory and applications to dependence-adjusted multiple testing. *The Annals of Statistics*, 46:1904–1931, 2018.
- H. Zou. The adaptive lasso and its oracle properties. *Journal of American Statistical Association*, 101:1418–1429, 2006.

An Online Supplement for
A multiple testing approach to the regularisation of large sample correlation
matrices

Natalia Bailey
Department of Econometrics and Business Statistics, Monash University

M. Hashem Pesaran
Department of Economics & USC Dornsife INET,
University of Southern California, USA, and Trinity College, Cambridge

L. Vanessa Smith
Department of Economics, University of York

October 14, 2018

Supplementary Appendix A

Technical Lemmas

A.1 Statement of technical lemmas

We begin by stating a number of technical lemmas that are needed for the proofs of the main results.

Lemma 1 Consider the distribution function of a standard normal variate defined by

$$\Phi(x) = (2\pi)^{-1/2} \int_{-\infty}^x e^{-\frac{u^2}{2}} du.$$

Then, for $x > 0$

$$\Phi(-x) = 1 - \Phi(x) \leq \frac{1}{2} \exp\left(-\frac{x^2}{2}\right). \quad (\text{A.1})$$

Lemma 2 Consider the critical value function⁷

$$c_p(N) = \Phi^{-1}\left(1 - \frac{p}{2f(N)}\right),$$

where $\Phi^{-1}(\cdot)$ is the inverse function of the cumulative standard normal distribution, $0 < p < 1$, $f(N) = c_\delta N^\delta$, where c_δ and δ are finite positive constants, and suppose there exists finite N_0 such that for all $N > N_0$

$$1 - \frac{p}{2f(N)} > 0. \quad (\text{A.2})$$

Then for $0 < \varkappa \leq 1$, we have

$$(a) \quad c_p(N) = O\left([\ln(N)]^{1/2}\right);$$

$$(b) \quad \exp\left[-\varkappa c_p^2(N)/2\right] = \Theta\left(N^{-\delta\varkappa}\right), \quad \lim_{N \rightarrow \infty} c_p^2(N)/\ln(N) = 2\delta;$$

$$(c) \quad \text{if } \delta > 1/\varkappa, \text{ then } N \exp\left[-\varkappa c_p^2(N)/2\right] \rightarrow 0, \text{ as } N \rightarrow \infty.$$

Lemma 3 Consider the sample correlation coefficient, $\hat{\rho}_{ij,T}$, defined by (4), and suppose that Assumptions 1 and 2 hold, $T = c_d N^d$, with $c_d > 0$. Then, there exists N_0 such that for all $N \geq N_0$,⁸

$$\begin{aligned} \Pr\left[|\hat{\rho}_{ij,T} - \rho_{ij}|\right] &\leq K e^{-\frac{1}{2} \frac{c_p^2(N)}{K_v(\boldsymbol{\theta}_{ij})}} + O\left(T^{-\frac{(s-2)}{2}} \left[\frac{c_p^2(N)}{K_v(\boldsymbol{\theta}_{ij})}\right]^{\frac{3(s-2)-1}{2}} e^{-\frac{1}{2} \frac{c_p^2(N)}{K_v(\boldsymbol{\theta}_{ij})}}\right) \\ &\quad + O\left(T^{-(s-1)/2}\right), \end{aligned} \quad (\text{A.3})$$

⁷We would like to thank George Kapetanios for his help with the proof of (b) and (c) of this Lemma.

⁸To simplify the notation we have dropped the lower order terms $e^{-\frac{1}{2} \frac{c_p(N)}{\sqrt{K_v(\boldsymbol{\theta}_{ij})}} \frac{K}{\sqrt{T}}}$ and $e^{-\frac{K}{T}}$, as they do not affect the results, and can be absorbed in the remainder order term.

and

$$\Pr \left[|\hat{\rho}_{ij,T}| > T^{-1/2} c_p(N) | \rho_{ij} = 0 \right] \leq K e^{-\frac{1}{2} \frac{c_p^2(N)}{\varphi_{ij}}} + O \left(T^{-\frac{(s-2)}{2}} \left[\frac{c_p^2(N)}{\varphi_{ij}} \right]^{\frac{3(s-2)-1}{2}} e^{-\frac{1}{2} \frac{c_p^2(N)}{\varphi_{ij}}} \right) + O \left(T^{-(s-1)/2} \right), \quad (\text{A.4})$$

where $K_v(\boldsymbol{\theta}_{ij})$ is given by (12), $\varphi_{ij} = E(y_{it}^2 y_{jt}^2 | \rho_{ij} = 0)$, and

$$c_p(N) = \Phi^{-1} \left(1 - \frac{p}{2f(N)} \right) > 0, \quad (\text{A.5})$$

with $0 < p < 1$, and $f(N) = c_\delta N^\delta$ where c_δ and δ are finite positive constants. Further, if $|\rho_{ij}| > c_p(N)/\sqrt{T}$ then we have

$$\Pr \left[|\hat{\rho}_{ij,T}| \leq T^{-1/2} c_p(N) | \rho_{ij} \neq 0 \right] \leq K e^{-\frac{T(|\rho_{ij}| - T^{-1/2} c_p(N))^2}{2K_v(\boldsymbol{\theta}_{ij})}} \left[1 + O \left(T^{\frac{2(s-2)-1}{2}} \right) \right] + O \left(T^{-(s-1)/2} \right). \quad (\text{A.6})$$

Lemma 4 Consider the standardised sample correlation coefficient

$$z_{ij,T} = [\text{Var}(\hat{\rho}_{ij,T})]^{-1/2} [\hat{\rho}_{ij,T} - E(\hat{\rho}_{ij,T})],$$

where $\hat{\rho}_{ij,T}$ is defined by (4) and $E(\hat{\rho}_{ij,T})$ and $\text{Var}(\hat{\rho}_{ij,T}) > 0$ are given by (9) and (10), respectively. Suppose that Assumptions 1 and 2 hold, $c_p(N) = \Phi^{-1} \left(1 - \frac{p}{2f(N)} \right)$, and condition (A.2) holds. Also let $T = c_d N^d$, with $c_d > 0$. Then, there exists N_0 such that for $N \geq N_0$, and for $r \geq 0$,

$$\sup_{ij} E \left[|z_{ij,T}|^r I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) | \rho_{ij} \neq 0 \right) \right] < K, \quad (\text{A.7})$$

and

$$\sup_{ij} E \left[|z_{ij,T}|^r I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) | \rho_{ij} = 0 \right) \right] \leq K e^{-\frac{1}{2} \frac{c_p^2(N)}{\varphi_{\max}}} + O \left(T^{-\frac{(s-2)}{2}} \left[\frac{c_p^2(N)}{\varphi_{\max}} \right]^{\frac{3(s-2)-1}{2}} e^{-\frac{1}{2} \frac{c_p^2(N)}{\varphi_{\max}}} \right) + O \left(T^{-(s-1)/2} \right), \quad (\text{A.8})$$

where $\varphi_{\max} = \sup_{ij} \varphi_{ij}$, φ_{ij} is defined by (13), and $s \geq 3$ is defined by Assumption 2.

Lemma 5 Consider the data generating process

$$\mathbf{y}_t = \mathbf{P} \mathbf{u}_t,$$

where \mathbf{y}_t and \mathbf{u}_t are $N \times 1$ vectors of random variables, and \mathbf{P} is an $N \times N$ matrix of fixed constants, such that $\mathbf{P} \mathbf{P}' = \mathbf{R}$, where \mathbf{R} is a correlation matrix. Suppose that \mathbf{u}_t follows a

multivariate t -distribution with v degrees of freedom generated as

$$\mathbf{u}_t = \left(\frac{v-2}{\chi_{v,t}^2} \right)^{1/2} \boldsymbol{\varepsilon}_t,$$

where $\boldsymbol{\varepsilon}_t = (\varepsilon_{1t}, \varepsilon_{2t}, \dots, \varepsilon_{Nt})' \sim IIDN(\mathbf{0}, \mathbf{I}_N)$, and $\chi_{v,t}^2$ is a chi-squared random variate with $v > 4$ degrees of freedom distributed independently of $\boldsymbol{\varepsilon}_t$. Then we have that

$$E(y_{it}^2 y_{jt}^2) = \frac{(v-2) [(\mathbf{P}'_i \mathbf{P}_i)^2 + (\mathbf{P}'_j \mathbf{P}_j)^2]}{(v-4)},$$

where \mathbf{p}'_i is the i^{th} row of \mathbf{P} . In the case where $\mathbf{P} = \mathbf{I}_N$, $E(y_{it}^2 y_{jt}^2) = (v-2)/(v-4)$ and

$$E(y_{it}^2 y_{jt}) = E(y_{jt}^2 y_{it}) = 0.$$

Lemma 6 *Fat-tailed shocks do not necessarily generate $E(y_{it}^2 y_{jt}^2) > 1$.*

A.2 Proofs of lemmas for the MT estimator

Proof of Lemma 1. Using results in Chiani et al. (2003) - eq. (5), we have

$$\text{erf c}(x) = \frac{2}{\sqrt{\pi}} \int_x^\infty e^{-u^2} du \leq \exp(-x^2), \quad (\text{A.9})$$

where $\text{erf c}(x)$ is the complement of the $\text{erf}(x)$ error function defined by

$$\text{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-u^2} du. \quad (\text{A.10})$$

But

$$1 - \Phi(x) = (2\pi)^{-1/2} \int_x^\infty e^{-\frac{u^2}{2}} du = \frac{1}{2} \text{erf c} \left(\frac{x}{\sqrt{2}} \right),$$

and using (A.9) we have

$$1 - \Phi(x) = \frac{1}{2} \text{erf c} \left(\frac{x}{\sqrt{2}} \right) \leq \frac{1}{2} \exp \left[- \left(\frac{x}{\sqrt{2}} \right)^2 \right] = \frac{1}{2} \exp \left(-\frac{x^2}{2} \right).$$

■

Proof of Lemma 2. First note that

$$\Phi^{-1}(z) = \sqrt{2} \text{erf}^{-1}(2z - 1), \quad z \in (0, 1),$$

where $\Phi(x)$ is cumulative distribution function of a standard normal variate, and $\text{erf}(x)$ function is defined by (A.10). Consider now the inverse complementary error function $\text{erfc}^{-1}(x)$ given by

$$\text{erf c}^{-1}(1 - x) = \text{erf}^{-1}(x).$$

Using results in Chiani et al. (2003) on p.842, we have

$$\operatorname{erf} c^{-1}(x) \leq \sqrt{-\ln(x)}.$$

Applying the above results to $c_p(N)$ we have

$$\begin{aligned} c_p(N) &= \Phi^{-1} \left(1 - \frac{p}{2f(N)} \right) \\ &= \sqrt{2} \operatorname{erf}^{-1} \left\{ 2 \left[1 - \frac{p}{2f(N)} \right] - 1 \right\} \\ &= \sqrt{2} \operatorname{erf}^{-1} \left[1 - \frac{p}{f(N)} \right] = \sqrt{2} \operatorname{erf} c^{-1} \left[\frac{p}{f(N)} \right] \\ &\leq \sqrt{2} \sqrt{-\ln \left[\frac{p}{f(N)} \right]} = \sqrt{2 [\ln f(N) - \ln(p)]}. \end{aligned}$$

Therefore, for $f(N) = c_\delta N^\delta$ we have

$$c_p^2(N) \leq 2 [\delta \ln(N) - \ln(p)] = O[\ln(N)],$$

which establishes part (a).

Further, by Proposition 24 of Dominici (2003) we have that

$$\lim_{N \rightarrow \infty} c_p(N)/LW \left\{ \frac{1}{2\pi \left[\left(1 - \frac{p}{2f(N)} \right) - 1 \right]^2} \right\}^{1/2} = 1,$$

where LW denotes the Lambert W function which satisfies $\lim_{N \rightarrow \infty} LW(N)/\{\ln(N) - \ln[\ln(N)]\} = 1$ as $N \rightarrow \infty$. We note that $\lim_{N \rightarrow \infty} \ln(N)/\{\ln(N) - \ln[\ln(N)]\} = 1$ as $N \rightarrow \infty$. So

$$\lim_{N \rightarrow \infty} \frac{LW \left\{ \frac{1}{2\pi \left[\left(1 - \frac{p}{2f(N)} \right) - 1 \right]^2} \right\}^{1/2}}{\left\{ 2 \ln \left(\frac{\sqrt{2f(N)}}{\sqrt{\pi p}} \right) \right\}^{1/2}} = 1.$$

Hence, for any $0 < \varkappa \leq 1$,

$$\lim_{N \rightarrow \infty} \frac{\exp[-\varkappa c_p^2(N)/2]}{\exp \left[-\frac{\varkappa \left\{ \left[2 \ln \left(\frac{\sqrt{2f(N)}}{\sqrt{\pi p}} \right) \right]^{1/2} \right\}^2}{2} \right]} = \lim_{N \rightarrow \infty} \frac{\exp[-\varkappa c_p^2(N)/2]}{[f(N)]^{-\varkappa} \pi^{\varkappa/2} p^{\varkappa} 2^{-\varkappa/2}} = 1, \text{ as } N \rightarrow \infty,$$

and substituting $c_\delta N^\delta$ for $f(N)$ yields,

$$\lim_{N \rightarrow \infty} \frac{\exp[-\varkappa c_p^2(N)/2]}{c_\delta^{-\varkappa} N^{-\delta \varkappa}} = \frac{2^{\varkappa/2} c_\delta^\varkappa}{\pi^{\varkappa/2} p^\varkappa}. \quad (\text{A.11})$$

It follows from (A.11) that $\exp[-\varkappa c_p^2(N)/2] = \Theta(N^{-\delta\varkappa})$, as required. From this result it also follows that $[-\varkappa c_p^2(N)/2] = \Theta(-\delta\varkappa \ln N)$, which in turn yields $\lim_{N \rightarrow \infty} c_p^2(N)/\ln(N) = 2\delta$. This completes the proof of part (b). Finally, it readily follows from (b) that $N \exp[-\varkappa c_p^2(N)/2] = \Theta(N^{1-\delta\varkappa})$, and therefore $N \exp[-\varkappa c_p^2(N)/2] \rightarrow 0$ when $\delta > 1/\varkappa$, as required for the proof of part (c). ■

Proof of Lemma 3. We first note that

$$\begin{aligned} & \Pr \left[\left| \hat{\rho}_{ij,T} - \rho_{ij} \right| > T^{-1/2} c_p(N) \right] \\ &= \Pr \left[\hat{\rho}_{ij,T} - \rho_{ij} > T^{-1/2} c_p(N) \right] + \Pr \left[-(\hat{\rho}_{ij,T} - \rho_{ij}) > T^{-1/2} c_p(N) \right] \\ &= \Pr \left[\hat{\rho}_{ij,T} - \rho_{ij} > T^{-1/2} c_p(N) \right] + \Pr \left[(\hat{\rho}_{ij,T} - \rho_{ij}) < -T^{-1/2} c_p(N) \right]. \end{aligned} \quad (\text{A.12})$$

and

$$\hat{\rho}_{ij,T} - \rho_{ij} = \omega_{ij,T} z_{ij,T} + (\rho_{ij,T} - \rho_{ij}),$$

where $z_{ij,T}$ is the standardised sample correlation coefficient defined by (15), $\rho_{ij,T} = E(\hat{\rho}_{ij,T})$ and $\omega_{ij,T}^2 = \text{Var}(\hat{\rho}_{ij,T}) > 0$ are given by (9) and (10), respectively. Hence

$$\begin{aligned} \Pr \left[\hat{\rho}_{ij,T} - \rho_{ij} > T^{-1/2} c_p(N) \right] &= \Pr \left[\omega_{ij,T} z_{ij,T} + (\rho_{ij,T} - \rho_{ij}) > T^{-1/2} c_p(N) \right] \\ &= \Pr \left(z_{ij,T} > a_{ij,T} \right), \end{aligned}$$

where

$$a_{ij,T} = \frac{T^{-1/2} c_p(N) - (\rho_{ij,T} - \rho_{ij})}{\omega_{ij,T}}.$$

Similarly,

$$\Pr \left[\hat{\rho}_{ij,T} - \rho_{ij} < -T^{-1/2} c_p(N) \right] = \Pr \left(z_{ij,T} < -b_{ij,T} \right),$$

where

$$b_{ij,T} = \frac{T^{-1/2} c_p(N) + (\rho_{ij,T} - \rho_{ij})}{\omega_{ij,T}}$$

But using (9) and (10) we have (note that by assumption $\sup_{ij} |K_m(\boldsymbol{\theta}_{ij})|$ and $\sup_{ij} K_v^{1/2}(\boldsymbol{\theta}_{ij}) < K$)

$$a_{ij,T} = \frac{T^{-1/2} c_p(N) - \frac{K_m(\boldsymbol{\theta}_{ij})}{T} + O(T^{-2})}{\sqrt{T^{-1} K_v(\boldsymbol{\theta}_{ij}) + O(T^{-2})}} = \frac{c_p(N)}{K_v^{1/2}(\boldsymbol{\theta}_{ij})} + O(T^{-1/2}), \quad (\text{A.13})$$

and

$$b_{ij,T} = \frac{T^{-1/2} c_p(N) + \frac{K_m(\boldsymbol{\theta}_{ij})}{T} + O(T^{-2})}{\sqrt{T^{-1} K_v(\boldsymbol{\theta}_{ij}) + O(T^{-2})}} = \frac{c_p(N)}{K_v^{1/2}(\boldsymbol{\theta}_{ij})} + O(T^{-1/2}). \quad (\text{A.14})$$

Using the above results in (A.12) we now have

$$\begin{aligned} \Pr \left[\left| \hat{\rho}_{ij,T} - \rho_{ij} \right| > \theta(N, T) \right] &= \Pr \left(z_{ij,T} > \frac{c_p(N)}{K_v^{1/2}(\boldsymbol{\theta}_{ij})} + O(T^{-1/2}) \right) \\ &\quad + \Pr \left(z_{ij,T} \leq \frac{-c_p(N)}{K_v^{1/2}(\boldsymbol{\theta}_{ij})} + O(T^{-1/2}) \right), \end{aligned}$$

where $\theta(N, T) = T^{-1/2}c_p(N)$. Result (A.3) now follows using (17) and (18) with a_T replaced by (A.13), and ignoring the higher order terms $e^{-\frac{K}{\sqrt{T}} \frac{c_p(N)}{\sqrt{K_v(\boldsymbol{\theta}_{ij})}}}$ and $e^{-\frac{K}{T}}$ that arise from squaring $a_T = K_v^{-1/2}(\boldsymbol{\theta}_{ij})c_p(N) + O(T^{-1/2})$. Result (A.4) can be obtained as a special case by setting $\rho_{ij} = 0$. Finally, to establish (A.6), using similar line of reasoning as above, we first note that

$$\begin{aligned} \Pr [|\hat{\rho}_{ij,T}| \leq \theta(N, T) | \rho_{ij} \neq 0] &= \Pr \left(z_{ij,T} \leq -\frac{\sqrt{T}\rho_{ij} - c_p(N)}{K_v^{1/2}(\boldsymbol{\theta}_{ij})} + O(T^{-1/2}) \right) \\ &\quad - \Pr \left(z_{ij,T} \leq -\frac{c_p(N) + \sqrt{T}\rho_{ij}}{K_v^{1/2}(\boldsymbol{\theta}_{ij})} + O(T^{-1/2}) \right) \\ &\leq \Pr \left(z_{ij,T} \leq -\frac{\sqrt{T}[\rho_{ij} - T^{-1/2}c_p(N)]}{K_v^{1/2}(\boldsymbol{\theta}_{ij})} + O(T^{-1/2}) \right). \end{aligned} \quad (\text{A.15})$$

Suppose that $\rho_{ij} > T^{-1/2}c_p(N) > 0$, then using (17) we have (again ignoring higher order terms in T^{-1})

$$\Pr [|\hat{\rho}_{ij,T}| \leq T^{-1/2}c_p(N) | \rho_{ij} \neq 0] \leq K e^{-\frac{T(\rho_{ij} - T^{-1/2}c_p(N))^2}{2K_v(\boldsymbol{\theta}_{ij})}} \left[1 + O\left(T^{\frac{2(s-2)-1}{2}}\right) \right] + O[T^{-(s-1)/2}]. \quad (\text{A.16})$$

A similar result follows when $\rho_{ij} < 0$. In this case we consider writing (A.15) equivalently as

$$\begin{aligned} \Pr [|\hat{\rho}_{ij,T}| \leq T^{-1/2}c_p(N) | \rho_{ij} \neq 0] &= 1 - \Pr \left(z_{ij,T} > -\frac{\sqrt{T}\rho_{ij} - c_p(N)}{K_v^{1/2}(\boldsymbol{\theta}_{ij})} + O(T^{-1/2}) \right) \\ &\quad - 1 + \Pr \left(z_{ij,T} > -\frac{c_p(N) + \sqrt{T}\rho_{ij}}{K_v^{1/2}(\boldsymbol{\theta}_{ij})} + O(T^{-1/2}) \right) \\ &\leq \Pr \left(z_{ij,T} > \frac{\sqrt{T}[-\rho_{ij} + T^{-1/2}c_p(N)]}{K_v^{1/2}(\boldsymbol{\theta}_{ij})} + O(T^{-1/2}) \right), \end{aligned}$$

where by assumption $-\rho_{ij} + T^{-1/2}c_p(N) > 0$. Now applying (18) to the right hand side of the above yields the outcome in (A.16) with ρ_{ij} replaced by $-\rho_{ij}$. Thus the desired result (A.6) is established for positive and negative values of ρ_{ij} such that $|\rho_{ij}| - T^{-1/2}c_p(N) > 0$.

Proof of Lemma 4. We first note that since $\inf_{ij} \text{Var}(\hat{\rho}_{ij,T}) > 0$, and $\hat{\rho}_{ij,T}$ is a correlation coefficient, $|\hat{\rho}_{ij,T}| \leq 1$, there exists T_0 such that for all $T > T_0$

$$|z_{ij,T}| \leq \frac{|\hat{\rho}_{ij,T}| + |E(\hat{\rho}_{ij,T})|}{\sqrt{\text{Var}(\hat{\rho}_{ij,T})}} \leq 2 \sup_{i,j} \left(\frac{1}{\sqrt{\text{Var}(\hat{\rho}_{ij,T})}} \right) < K.$$

Hence, $E |z_{ij,T}|^r < K$ for any finite r . Also, by the Cauchy–Schwarz inequality

$$\begin{aligned} E \left[|z_{ij,T}|^r I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} \neq 0 \right) \right] &\leq [E (z_{ij,T}^{2r})]^{1/2} \left\{ E \left[I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} \neq 0 \right) \right] \right\}^{1/2} \\ &= [E (z_{ij,T}^{2r})]^{1/2} \Pr \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} \neq 0 \right) \\ &\leq K. \end{aligned}$$

which establishes (A.7), as required. Similarly,

$$E \left[|z_{ij,T}|^r I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} = 0 \right) \right] \leq K \Pr \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} = 0 \right),$$

and using result (A.4) of Lemma 3, we have

$$\begin{aligned} E \left[|z_{ij,T}|^r I \left(\left| \sqrt{T} \hat{\rho}_{ij,T} \right| > c_p(N) \mid \rho_{ij} = 0 \right) \right] &\leq K e^{-\frac{1}{2} \frac{c_p^2(N)}{\varphi_{ij}}} + O \left(T^{-\frac{(s-2)}{2}} \left[\frac{c_p^2(N)}{\varphi_{ij}} \right]^{\frac{3(s-2)-1}{2}} e^{-\frac{1}{2} \frac{c_p^2(N)}{\varphi_{ij}}} \right) \\ &\quad + O(T^{-(s-1)/2}). \end{aligned}$$

which establishes (A.8). ■

Proof of Lemma 5. First we note that

$$\begin{aligned} E \left(\frac{1}{\chi_{v,t}^2} \right) &= \frac{1}{v-2}, \quad Var \left(\frac{1}{\chi_{v,t}^2} \right) = \frac{2}{(v-2)^2(v-4)} \\ E \left(\frac{1}{\chi_{v,t}^2} \right)^2 &= \frac{2}{(v-2)^2(v-4)} + \left(\frac{1}{v-2} \right)^2 = \frac{v-2}{(v-2)^2(v-4)}. \end{aligned} \quad (\text{A.17})$$

Then

$$E(\mathbf{u}_t \mathbf{u}_t') = E \left[\left(\frac{v-2}{\chi_v^2} \right) \boldsymbol{\varepsilon}_t \boldsymbol{\varepsilon}_t' \right] = E \left(\frac{v-2}{\chi_{v,t}^2} \right) E(\boldsymbol{\varepsilon}_t \boldsymbol{\varepsilon}_t') = \mathbf{I}_N,$$

and

$$E(\mathbf{y}_t) = \mathbf{0}, \quad E(\mathbf{y}_t \mathbf{y}_t') = \mathbf{P} \mathbf{P}' = \mathbf{R}.$$

It is clear that y_{it} has mean zero and a unit variance. Denote the i^{th} row of \mathbf{P} by \mathbf{p}_i' and note that $y_{it} = \mathbf{p}_i' \mathbf{u}_t = \left(\frac{v-2}{\chi_{v,t}^2} \right)^{1/2} \mathbf{p}_i' \boldsymbol{\varepsilon}_t$, and hence

$$\mu_{ij}(2, 2) = E(y_{it}^2 y_{jt}^2) = E \left[\left(\frac{v-2}{\chi_{v,t}^2} \right)^2 (\mathbf{p}_i' \boldsymbol{\varepsilon}_t)^2 (\mathbf{p}_j' \boldsymbol{\varepsilon}_t)^2 \right],$$

and since $\boldsymbol{\varepsilon}_t$ and $\chi_{v,t}^2$ are distributed independently using (A.17) we have

$$E(y_{it}^2 y_{jt}^2) = \frac{(v-2)^3}{(v-2)^2(v-4)} E[(\boldsymbol{\varepsilon}_t' \mathbf{A}_i \boldsymbol{\varepsilon}_t) (\boldsymbol{\varepsilon}_t' \mathbf{A}_j \boldsymbol{\varepsilon}_t)],$$

where $\mathbf{A}_i = \mathbf{p}_i \mathbf{p}_i'$. But since $\boldsymbol{\varepsilon}_t \sim N(\mathbf{0}, \mathbf{I}_N)$, using results in Magnus (1978) we have

$$\begin{aligned} E[(\boldsymbol{\varepsilon}_t' \mathbf{A}_i \boldsymbol{\varepsilon}_t)(\boldsymbol{\varepsilon}_t' \mathbf{A}_j \boldsymbol{\varepsilon}_t)] &= \text{tr}(\mathbf{p}_i \mathbf{p}_i') \text{tr}(\mathbf{p}_j \mathbf{p}_j') + \text{tr}(\mathbf{p}_i \mathbf{p}_i' \mathbf{p}_j \mathbf{p}_j') \\ &= (\mathbf{p}_i' \mathbf{p}_i)^2 + (\mathbf{p}_i' \mathbf{p}_j)^2. \end{aligned}$$

Hence

$$E(y_{it}^2 y_{jt}^2) = \frac{(v-2) [(\mathbf{p}_i' \mathbf{p}_i)^2 + (\mathbf{p}_i' \mathbf{p}_j)^2]}{(v-4)}.$$

When \mathbf{P} is an identity matrix then $\mathbf{p}_i' \mathbf{p}_i = 1$ and $\mathbf{p}_i' \mathbf{p}_j = 0$, and hence $E(y_{it}^2 y_{jt}^2) = (v-2)/(v-4)$. Also

$$E(y_{it}^2 y_{jt}) = E \left[\left(\frac{v-2}{\chi_{v,t}^2} \right)^{3/2} \right] E[(\boldsymbol{\varepsilon}_t' \mathbf{A}_i \boldsymbol{\varepsilon}_t) \mathbf{p}_j' \boldsymbol{\varepsilon}_t] = 0.$$

■

Proof of Lemma 6. Consider the data generating process $\mathbf{y}_t = \mathbf{P} \mathbf{u}_t$ where the elements of $\mathbf{u}_t = (u_{1t}, u_{2t}, \dots, u_{Nt})'$, u_{it} , are generated as a standardized independent chi-squared distribution with v_i degrees of freedom, namely

$$u_{it} = \frac{\chi_{it}^2(v_i) - v_i}{\sqrt{2v_i}}, \text{ for all } i \text{ and } t.$$

Then it is clear that $E(u_{it}) = 0$, $E(u_{it}^2) = 1$, as well as $E(u_{it}^2 u_{jt}^2) = E(u_{it}^2) E(u_{jt}^2) = 1$, and $E(\mathbf{u}_t \mathbf{u}_t') = \mathbf{I}_N$. Let \mathbf{p}_i' be the i^{th} row of \mathbf{P} and note that

$$\begin{aligned} E(y_{it} y_{jt}) &= \mathbf{p}_i' E(\mathbf{u}_t \mathbf{u}_t') \mathbf{p}_j = \mathbf{p}_i' \mathbf{p}_j = \rho_{ij} \\ \mathbf{p}_i' \mathbf{p}_i &= \sum_{r=1}^N p_{ir}^2 = 1. \end{aligned}$$

Also

$$\begin{aligned} E(y_{it}^2 y_{jt}^2) &= E[(\mathbf{p}_i' \mathbf{u}_t \mathbf{u}_t' \mathbf{p}_i)(\mathbf{p}_j' \mathbf{u}_t \mathbf{u}_t' \mathbf{p}_j)] \\ &= \sum_r \sum_{r'} \sum_s \sum_{s'} p_{ir} p_{ir'} p_{js} p_{js'} E(u_{rt} u_{r't} u_{st} u_{s't}). \end{aligned}$$

But

$$\begin{aligned} E(u_{rt} u_{r't} u_{st} u_{s't}) &= 0 \text{ if } r \neq r' \text{ or } s \neq s' \\ &= E(u_{rt}^2 u_{st}^2) = 1 \text{ if } r = r' \text{ and } s = s', \end{aligned}$$

and hence

$$E(y_{it}^2 y_{jt}^2) = \sum_r \sum_s p_{ir}^2 p_{js}^2 = \left(\sum_{r=1}^N p_{ir}^2 \right)^2 = 1.$$

Therefore, fat-tailed shocks do not necessarily generate $\mu_{ij}(2, 2) = E(y_{it}^2 y_{jt}^2) > 1$. ■

Supplementary Appendix B

An overview of key regularisation techniques

Here we provide an overview of three main covariance estimators proposed in the literature which we use in our Monte Carlo experiments for comparative analysis, namely the thresholding methods of Bickel and Levina (2008), and Cai and Liu (2011), and the shrinkage approach of Ledoit and Wolf (2004).

B.1 Bickel-Levina (BL) thresholding

The method developed by Bickel and Levina (2008) - BL - employs ‘universal’ thresholding of the sample covariance matrix $\hat{\Sigma} = (\hat{\sigma}_{ij})$, $i, j = 1, 2, \dots, N$. Under this approach Σ is required to be sparse as they define on p. 2580. The BL thresholding estimator is given by

$$\tilde{\Sigma}_{BL,C} = \left(\hat{\sigma}_{ij} I \left[|\hat{\sigma}_{ij}| \geq C \sqrt{\frac{\ln(N)}{T}} \right] \right), \quad i = 1, 2, \dots, N-1, \quad j = i+1, i+2, \dots, N \quad (\text{B.18})$$

where $I(\cdot)$ is an indicator function and C is a positive constant which is unknown. The choice of thresholding function - $I(\cdot)$ - implies that (B.18) implements ‘hard’ thresholding.

The consistency rate of the BL estimator is $m_N \sqrt{\frac{\ln(N)}{T}}$ under the spectral norm of the error matrix $(\tilde{\Sigma}_{BL,C} - \Sigma)$. The potential computational burden in the implementation of this approach is the estimation of the thresholding parameter, C . This is usually calibrated by a separate cross-validation (CV) procedure. The quality of the performance of the BL estimator is rooted in the specification chosen for the implementation of CV.⁹ Details of the BL cross-validation procedure are given in Section B.3.

As argued by BL, thresholding maintains the symmetry of $\hat{\Sigma}$ but does not ensure positive definiteness of $\tilde{\Sigma}_{BL,\hat{C}}$ in finite samples. BL show that their threshold estimator is positive definite if

$$\left\| \tilde{\Sigma}_{BL,C} - \tilde{\Sigma}_{BL,0} \right\| \leq \epsilon \text{ and } \lambda_{\min}(\Sigma) > \epsilon, \quad (\text{B.19})$$

where $\|\cdot\|$ is the spectral or operator norm and ϵ is a small positive constant. This condition is not met unless T is sufficiently large relative to N . ‘Universal’ thresholding on $\hat{\Sigma}$ performs best when the units x_{it} , $i = 1, 2, \dots, N$, $t = 1, 2, \dots, T$ are assumed homoskedastic (i.e. $\sigma_{11} = \sigma_{22} = \dots = \sigma_{NN}$).

B.2 Cai and Liu (CL) thresholding

Cai and Liu (2011) - CL - proposed an improved version of the BL approach by incorporating the unit specific variances in their ‘adaptive’ thresholding procedure. In this way, unlike ‘universal’ thresholding on $\hat{\Sigma}$, their estimator is robust to heteroscedasticity. Specifically,

⁹Fang et al. (2013) provide useful guidelines regarding the specification of various parameters used in cross-validation through an extensive simulation study.

the thresholding estimator $\tilde{\Sigma}_{CL,C}$ is defined as

$$\tilde{\Sigma}_{CL,C} = (\hat{\sigma}_{ij} s_{\tau_{ij}} [|\hat{\sigma}_{ij}| \geq \tau_{ij}]), \quad i = 1, 2, \dots, N-1, \quad j = i+1, i+2, \dots, N \quad (\text{B.20})$$

where $\tau_{ij} > 0$ is an entry-dependent adaptive threshold such that $\tau_{ij} = \sqrt{\hat{\theta}_{ij} \omega_T}$, with $\hat{\theta}_{ij} = T^{-1} \sum_{i=1}^T (x_{it} x_{jt} - \hat{\sigma}_{ij})^2$ and $\omega_T = C \sqrt{\ln(N)/T}$, for some constant $C > 0$. CL implement their approach using the general thresholding function $s_{\tau}(\cdot)$ rather than $I(\cdot)$, but point out that all their theoretical results continue to hold for the hard thresholding estimator. The consistency rate of the CL estimator is $C_0 m_N \sqrt{\ln(N)/T}$ under the spectral norm of the error matrix $(\tilde{\Sigma}_{CL,C} - \Sigma)$. The parameter C can be fixed to a constant implied by theory ($C = 2$ in CL) or chosen via cross-validation. Details of the CL cross-validation procedure are provided in Section B.3.

As with the BL estimator, thresholding in itself does not ensure positive definiteness of $\tilde{\Sigma}_{CL,\hat{C}}$. In light of condition (B.19), Fan et al. (2013) - FLM - extend the CL approach and propose setting a lower bound on the cross-validation grid when searching for C such that the minimum eigenvalue of their threshold estimator is positive, $\lambda_{\min}(\tilde{\Sigma}_{FLM,\hat{C}}) > 0$. This idea originated from Fryzlewicz (2013). Further details of this procedure can be found in Section B.3. We apply this extension to both BL and CL procedures (see Section B.3 for the relevant expressions).

B.3 Cross-validation

We perform a grid search for the choice of C over a specified range: $C = \{c : C_{\min} \leq c \leq C_{\max}\}$. In the BL procedure, we set $C_{\min} = \left| \min_{ij} \hat{\sigma}_{ij} \right| \sqrt{\frac{T}{\ln N}}$ and $C_{\max} = \left| \max_{ij} \hat{\sigma}_{ij} \right| \sqrt{\frac{T}{\ln N}}$ and impose increments of $\frac{C_{\max} - C_{\min}}{N}$. In CL cross-validation, we set $C_{\min} = 0$ and $C_{\max} = 4$, and impose increments of c/N for $c = 1$. In each point of the respective ranges, c , we use x_{it} , $i = 1, 2, \dots, N$, $t = 1, 2, \dots, T$ and select the $N \times 1$ column vectors $\mathbf{x}_t = (x_{1t}, x_{2t}, \dots, x_{Nt})'$, $t = 1, 2, \dots, T$ which we randomly reshuffle over the t -dimension. This gives rise to a new set of $N \times 1$ column vectors $\mathbf{x}_t^{(s)} = (x_{1t}^{(s)}, x_{2t}^{(s)}, \dots, x_{Nt}^{(s)})'$ for the first shuffle $s = 1$. We repeat this reshuffling S times in total where we set $S = 50$. We consider this to be sufficiently large (FLM suggested $S = 20$ while BL recommended $S = 100$ - see also Fang et al. (2013)). In each shuffle $s = 1, 2, \dots, S$, we divide $\mathbf{x}^{(s)} = (\mathbf{x}_1^{(s)}, \mathbf{x}_2^{(s)}, \dots, \mathbf{x}_T^{(s)})$ into two subsamples of size $N \times T_1$ and $N \times T_2$, where $T_2 = T - T_1$. A theoretically 'justified' split suggested in BL is given by $T_1 = T \left(1 - \frac{1}{\ln(T)}\right)$ and $T_2 = \frac{T}{\ln(T)}$. In our simulation study we set $T_1 = \frac{2T}{3}$ and $T_2 = \frac{T}{3}$. Let $\hat{\Sigma}_1^{(s)} = (\hat{\sigma}_{1,ij}^{(s)})$, with elements $\hat{\sigma}_{1,ij}^{(s)} = T_1^{-1} \sum_{t=1}^{T_1} x_{it}^{(s)} x_{jt}^{(s)}$, and $\hat{\Sigma}_2^{(s)} = (\hat{\sigma}_{2,ij}^{(s)})$ with elements $\hat{\sigma}_{2,ij}^{(s)} = T_2^{-1} \sum_{t=T_1+1}^T x_{it}^{(s)} x_{jt}^{(s)}$, $i, j = 1, 2, \dots, N$, denote the sample covariance matrices generated using T_1 and T_2 respectively, for each shuffle s . We threshold $\hat{\Sigma}_1^{(s)}$ as in (B.18), (B.20) using $I(\cdot)$ as the thresholding function, where for CL both $\hat{\theta}_{ij}$ and ω_T are adjusted to

$$\hat{\theta}_{1,ij}^{(s)} = \frac{1}{T_1} \sum_{t=1}^{T_1} (x_{it}^{(s)} x_{jt}^{(s)} - \hat{\sigma}_{1,ij}^{(s)})^2,$$

and

$$\omega_{T_1}(c) = c \sqrt{\frac{\ln(N)}{T_1}}.$$

Then (B.20) becomes

$$\tilde{\Sigma}_1^{(s)}(c) = \left(\hat{\sigma}_{1,ij}^{(s)} I \left[\left| \hat{\sigma}_{1,ij}^{(s)} \right| \geq \tau_{1,ij}^{(s)}(c) \right] \right),$$

for each c , where

$$\tau_{1,ij}^{(s)}(c) = \sqrt{\hat{\theta}_{1,ij}^{(s)}} \omega_{T_1}(c) > 0,$$

and $\hat{\theta}_{1,ij}^{(s)}$ and $\omega_{T_1}(c)$ are defined above. The following expression is computed for BL and CL,

$$\hat{G}(c) = \frac{1}{S} \sum_{s=1}^S \left\| \tilde{\Sigma}_1^{(s)}(c) - \hat{\Sigma}_2^{(s)} \right\|_F^2, \quad (\text{B.21})$$

for each c and

$$\hat{C} = \arg \min_{C_{\min} \leq c \leq C_{\max}} \hat{G}(c). \quad (\text{B.22})$$

If several values of c attain the minimum of (B.22), then \hat{C} is chosen to be the smallest one. The final estimator of the covariance matrix is then given by $\tilde{\Sigma}_{\hat{C}}$. The thresholding approach does not necessarily ensure that the resultant estimate, $\tilde{\Sigma}_{\hat{C}}$, is positive definite. To ensure that the threshold estimator is positive definite FLM propose setting a lower bound on the cross-validation grid for the search of C such that $\lambda_{\min}(\tilde{\Sigma}_{\hat{C}}) > 0$ - see Fryzlewicz (2013). Therefore, for BL and CL we modify (B.22) so that

$$\hat{C}^* = \arg \min_{C_{pd} + \epsilon \leq c \leq C_{\max}} \hat{G}(c), \quad (\text{B.23})$$

where C_{pd} is the lowest c such that $\lambda_{\min}(\tilde{\Sigma}_{C_{pd}}) > 0$ and ϵ is a small positive constant. We do not conduct thresholding on the diagonal elements of the covariance matrices which remain in tact.

B.4 Ledoit and Wolf (LW) shrinkage

Ledoit and Wolf (2004) - LW - considered a shrinkage estimator for regularisation which is based on a linear combination of the sample covariance matrix, $\hat{\Sigma}$, and an identity matrix \mathbf{I}_N , and provide formulae for the appropriate weights. The LW shrinkage is expressed as

$$\hat{\Sigma}_{LW} = \hat{\rho}_1 \mathbf{I}_N + \hat{\rho}_2 \hat{\Sigma}, \quad (\text{B.24})$$

with the estimated weights given by

$$\hat{\rho}_1 = m_T b_T^2 / d_T^2, \quad \hat{\rho}_2 = a_T^2 / d_T^2$$

where

$$\begin{aligned} m_T &= N^{-1} \operatorname{tr}(\hat{\Sigma}), \quad d_T^2 = N^{-1} \operatorname{tr}(\hat{\Sigma}^2) - m_T^2, \\ a_T^2 &= d_T^2 - b_T^2, \quad b_T^2 = \min(\bar{b}_T^2, d_T^2), \end{aligned}$$

and

$$\bar{b}_T^2 = \frac{1}{NT^2} \sum_{t=1}^T \left\| \dot{\mathbf{x}}_t \dot{\mathbf{x}}_t' - \hat{\Sigma} \right\|_F^2 = \frac{1}{NT^2} \sum_{t=1}^T \operatorname{tr}[(\dot{\mathbf{x}}_t \dot{\mathbf{x}}_t')(\dot{\mathbf{x}}_t \dot{\mathbf{x}}_t')] - \frac{2}{NT^2} \sum_{t=1}^T \operatorname{tr}(\dot{\mathbf{x}}_t' \hat{\Sigma} \dot{\mathbf{x}}_t) + \frac{1}{NT} \operatorname{tr}(\hat{\Sigma}^2),$$

and noting that $\sum_{t=1}^T \operatorname{tr}(\dot{\mathbf{x}}_t' \hat{\Sigma} \dot{\mathbf{x}}_t) = \sum_{t=1}^T \operatorname{tr}(\hat{\Sigma} \sum_{t=1}^T \dot{\mathbf{x}}_t \dot{\mathbf{x}}_t') = T \sum_{t=1}^T \operatorname{tr}(\hat{\Sigma}^2)$, we have

$$\bar{b}_T^2 = \frac{1}{NT^2} \sum_{t=1}^T \left(\sum_{i=1}^N \dot{x}_{it}^2 \right)^2 - \frac{1}{NT} \operatorname{tr}(\hat{\Sigma}^2),$$

with $\dot{\mathbf{x}}_t = (\dot{x}_{1t}, \dot{x}_{2t}, \dots, \dot{x}_{Nt})'$ and $\dot{x}_{it} = (x_{it} - \bar{x}_i)$.¹⁰

$\hat{\Sigma}_{LW}$ is positive definite by construction. Thus, the inverse $\hat{\Sigma}_{LW}^{-1}$ exists and is well conditioned.

Supplementary Appendix C

Shrinkage on MT estimator (S-MT)

Recall the shrinkage on the multiple testing estimator (*S-MT*) expression displayed in Section 3.1,

$$\tilde{\mathbf{R}}_{S-MT}(\xi) = \xi \mathbf{I}_N + (1 - \xi) \tilde{\mathbf{R}}_{MT},$$

where the $N \times N$ identity matrix \mathbf{I}_N is set as benchmark target, the shrinkage parameter is denoted by $\xi \in (\xi_0, 1]$, and ξ_0 is the minimum value of ξ that produces a non-singular $\tilde{\mathbf{R}}_{S-MT}(\xi_0)$ matrix. Note that shrinkage is deliberately implemented on the correlation matrix $\tilde{\mathbf{R}}_{MT}$ rather than on $\tilde{\Sigma}_{MT}$. In this way we ensure that no shrinkage is applied to the variances. Further, shrinkage is applied to the non-zero elements of $\tilde{\mathbf{R}}_{MT}$, and as a result the shrinkage estimator, $\tilde{\mathbf{R}}_{S-MT}$, also consistently recovers the support of \mathbf{R} , since it has the same support recovery property as $\tilde{\mathbf{R}}_{MT}$. With regard to the calibration of the shrinkage parameter, ξ , we solve the following optimisation problem

$$\xi^* = \arg \min_{\xi_0 + \epsilon \leq \xi \leq 1} \left\| \mathbf{R}_0^{-1} - \tilde{\mathbf{R}}_{S-MT}^{-1}(\xi) \right\|_F^2,$$

¹⁰Note that LW scale the Frobenius norm by $1/N$, and use $\|\mathbf{A}\|_F^2 = \operatorname{tr}(\mathbf{A}'\mathbf{A})/N$. See Definition 1 of Ledoit and Wolf (2004, p. 376). Here we use the standard notation for this norm.

where ϵ is a small positive constant, and \mathbf{R}_0 is a reference invertible correlation matrix. Let $\mathbf{A} = \mathbf{R}_0^{-1}$ and $\mathbf{B}(\xi) = \tilde{\mathbf{R}}_{S-MT}^{-1}(\xi)$. Note that since \mathbf{R}_0 and $\tilde{\mathbf{R}}_{S-MT}$ are symmetric

$$\left\| \mathbf{R}_0^{-1} - \tilde{\mathbf{R}}_{S-MT}^{-1}(\xi) \right\|_F^2 = \text{tr}(\mathbf{A}^2) - 2 \text{tr}[\mathbf{A}\mathbf{B}(\xi)] + \text{tr}[\mathbf{B}^2(\xi)].$$

The first order condition for the above optimisation problem is given by

$$\frac{\partial \left\| \mathbf{R}_0^{-1} - \tilde{\mathbf{R}}_{S-MT}^{-1}(\xi) \right\|_F^2}{\partial \xi} = -2 \text{tr} \left(\mathbf{A} \frac{\partial \mathbf{B}(\xi)}{\partial \xi} \right) + 2 \text{tr} \left(\mathbf{B}(\xi) \frac{\partial \mathbf{B}(\xi)}{\partial \xi} \right),$$

where

$$\begin{aligned} \frac{\partial \mathbf{B}(\xi)}{\partial \xi} &= -\tilde{\mathbf{R}}_{S-MT}^{-1}(\xi) \left(\mathbf{I}_N - \tilde{\mathbf{R}}_{MT} \right) \tilde{\mathbf{R}}_{S-MT}^{-1}(\xi) \\ &= -\mathbf{B}(\xi) \left(\mathbf{I}_N - \tilde{\mathbf{R}}_{MT} \right) \mathbf{B}(\xi). \end{aligned}$$

Hence, ξ^* is obtained as the solution of

$$f(\xi) = -\text{tr} \left[(\mathbf{A} - \mathbf{B}(\xi)) \mathbf{B}(\xi) \left(\mathbf{I}_N - \tilde{\mathbf{R}}_{MT} \right) \mathbf{B}(\xi) \right] = 0,$$

where $f(\xi)$ is an analytic differentiable function of ξ for values of ξ close to unity, such that $\mathbf{B}(\xi)$ exists.

The resulting $\tilde{\mathbf{R}}_{S-MT}(\xi^*)$ is guaranteed to be positive definite since

$$\lambda_{\min} \left[\tilde{\mathbf{R}}_{S-MT}(\xi) \right] = \xi \lambda_{\min}(\mathbf{I}_N) + (1 - \xi) \lambda_{\min}(\tilde{\mathbf{R}}_{MT}) > 0,$$

for any $\xi \in [\xi_0, 1]$, where $\xi_0 = \max \left(\frac{\epsilon - \lambda_{\min}(\tilde{\mathbf{R}}_{MT})}{1 - \lambda_{\min}(\tilde{\mathbf{R}}_{MT})}, 0 \right)$.

C.1 Derivation of S-MT shrinkage parameter

We need to solve $f(\xi) = 0$ for ξ^* such that $f(\xi^*) = 0$ for a given choice of \mathbf{R}_0 .¹¹

Abstracting from the subscripts, note that

$$f(1) = -\text{tr} \left[(\mathbf{R}^{-1} - \mathbf{I}_N) \left(\mathbf{I}_N - \tilde{\mathbf{R}} \right) \right],$$

or

$$\begin{aligned} f(1) &= -\text{tr} \left[-\mathbf{R}^{-1} \tilde{\mathbf{R}} + \mathbf{R}^{-1} - \mathbf{I}_N + \tilde{\mathbf{R}} \right] \\ &= \text{tr} \left(\mathbf{R}^{-1} \tilde{\mathbf{R}} \right) - \text{tr} \left(\mathbf{R}^{-1} \right), \end{aligned}$$

which is generally non-zero. Also, $\xi = 0$ is ruled out, since $\tilde{\mathbf{R}}_{S-MT}(0) = \tilde{\mathbf{R}}$ need not be

¹¹The code for computing \mathbf{R}_0 of our choice is available upon request (see Section C.2).

non-singular.

Thus we need to assess whether $f(\xi) = 0$ has a solution in the range $\xi_0 < \xi < 1$, where ξ_0 is the minimum value of ξ such that $\tilde{\mathbf{R}}_{S-MT}(\xi_0)$ is non-singular. First, we can compute ξ_0 by implementing naive shrinkage as an initial estimate:

$$\tilde{\mathbf{R}}_{S-MT}(\xi_0) = \xi_0 \mathbf{I}_N + (1 - \xi_0) \tilde{\mathbf{R}}.$$

The shrinkage parameter $\xi_0 \in [0, 1]$ is given by

$$\xi_0 = \max \left(\frac{\epsilon - \lambda_{\min}(\tilde{\mathbf{R}})}{1 - \lambda_{\min}(\tilde{\mathbf{R}})}, 0 \right),$$

where in our simulation study we set $\epsilon = 0.01$. Here, $\lambda_{\min}(\mathbf{A})$ stands for the minimum eigenvalue of matrix \mathbf{A} . If $\tilde{\mathbf{R}}$ is already positive definite and $\lambda_{\min}(\tilde{\mathbf{R}}) > 0$, then ξ_0 is automatically set to zero. Conversely, if $\lambda_{\min}(\tilde{\mathbf{R}}) \leq 0$, then ξ_0 is set to the smallest possible value that ensures positivity of $\lambda_{\min}(\tilde{\mathbf{R}}_{S-MT}(\xi_0))$.

Second, we implement the optimisation procedure. In our simulation study we employ a grid search for $\xi^* = \{\xi : \xi_0 + \epsilon \leq \xi \leq 1\}$ with increments of 0.005. The final ξ^* is given by

$$\xi^* = \arg \min_{\xi} [f(\xi)]^2.$$

C.2 Specification of reference matrix \mathbf{R}_0

Implementation of the above procedure requires the use of a suitable reference matrix \mathbf{R}_0 . Our experimentations suggested that the shrinkage estimator of Ledoit and Wolf (2004) - LW - applied to the correlation matrix is likely to work well in practice, and is to be recommended. Schäfer and Strimmer (2005) consider LW shrinkage on the correlation matrix. In our application we also take account of the small sample bias of the correlation coefficients in what follows. We set as reference matrix \mathbf{R}_0 the shrinkage estimator of LW applied to the sample correlation matrix:

$$\hat{\mathbf{R}}_0 = \theta \mathbf{I}_N + (1 - \theta) \hat{\mathbf{R}},$$

with shrinkage parameter $\theta \in [0, 1]$, and $\hat{\mathbf{R}} = (\hat{\rho}_{ij})$. The optimal value of the shrinkage parameter that minimizes the expectation of the squared Frobenius norm of the error of estimating \mathbf{R} by $\hat{\mathbf{R}}_0$:

$$E \left\| \hat{\mathbf{R}}_0 - \mathbf{R} \right\|_F^2 = \sum_{i \neq j} \sum_{i \neq j} E (\hat{\rho}_{ij} - \rho_{ij})^2 + \theta^2 \sum_{i \neq j} \sum_{i \neq j} E (\hat{\rho}_{ij}^2) - 2\theta \sum_{i \neq j} \sum_{i \neq j} E [\hat{\rho}_{ij} (\hat{\rho}_{ij} - \rho_{ij})], \quad (\text{C.25})$$

is given by

$$\theta^* = \frac{\sum_{i \neq j} \sum_{i \neq j} E [\hat{\rho}_{ij} (\hat{\rho}_{ij} - \rho_{ij})]}{\sum_{i \neq j} \sum_{i \neq j} E (\hat{\rho}_{ij}^2)} = 1 - \frac{\sum_{i \neq j} \sum_{i \neq j} E (\hat{\rho}_{ij} \rho_{ij})}{\sum_{i \neq j} \sum_{i \neq j} E (\hat{\rho}_{ij}^2)}, \quad (\text{C.26})$$

with

$$\hat{\theta}^* = 1 - \frac{\sum_{i \neq j} \sum \hat{\rho}_{ij} \left[\hat{\rho}_{ij} - \frac{\hat{\rho}_{ij}(1-\hat{\rho}_{ij}^2)}{2T} \right]}{\frac{1}{T} \sum_{i \neq j} \sum (1 - \hat{\rho}_{ij}^2)^2 + \sum_{i \neq j} \sum \left[\hat{\rho}_{ij} - \frac{\hat{\rho}_{ij}(1-\hat{\rho}_{ij}^2)}{2T} \right]^2}.$$

Note that $\lim_{T \rightarrow \infty}(\hat{\theta}^*) = 0$ for any N . However, in small samples values of $\hat{\theta}^*$ can be obtained that fall outside the range $[0, 1]$. To avoid such cases, if $\hat{\theta}^* < 0$ then $\hat{\theta}^*$ is set to 0, and if $\hat{\theta}^* > 1$ it is set to 1, or $\hat{\theta}^{**} = \max(0, \min(1, \hat{\theta}^*))$.

References

- Bickel, P. J. and E. Levina (2008). Covariance regularisation by thresholding. *The Annals of Statistics* 36, 2577-2604.
- Cai, T. and W. Liu (2011). Adaptive thresholding for sparse covariance matrix estimation. *Journal of American Statistical Association* 106, 672-684.
- Chiani, M., D. Dardari, and M. K. Simon (2003). New exponential bounds and approximations for the computation of error probability in fading channels. *IEEE Transactions on Wireless Communications* 2(4), 840-845.
- Dominici, D. E. (2003). The inverse of the cumulative standard normal probability function. *Integral Transforms and Special Functions* 14, 281-292.
- Fan, J., Y. Liao, and M. Mincheva (2013). Large covariance estimation by thresholding principal orthogonal complements. *Journal of Royal Statistical Society, Series B* 75, 1-44.
- Fang, Y., B. Wang, and Y. Feng (2016). Tuning parameter selection in regularised estimations of large covariance matrices. *Journal of Statistical Computation and Simulation* 86, 494-509.
- Fryzlewicz, P. (2013). High-dimensional volatility matrix estimation via wavelets and thresholding. *Biometrika* 100, 921-938.
- Ledoit, O. and M. Wolf (2004). A well-conditioned estimator for large-dimensional covariance matrices. *Journal of Multivariate Analysis* 88, 365-411.
- Magnus, J. R. (1978). The moments of products of quadratic forms in normal variables. *Statistica Neerlandica* 32, 201-210.
- Schäfer, J. and K. Strimmer (2005). A shrinkage approach to large-scale covariance matrix estimation and implications for functional Genomics. *Statistical Applications in Genetics and Molecular Biology* 4, Article 32.