



This is a repository copy of *Acceptability of post-mortem imaging among Muslim and non-Muslim communities*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/138245/>

Version: Accepted Version

---

**Article:**

Ben Taher, M., Pearson, J., Cohen, M. et al. (1 more author) (2018) Acceptability of post-mortem imaging among Muslim and non-Muslim communities. *British Journal of Radiology*, 91 (1091). 20180295. ISSN 0007-1285

<https://doi.org/10.1259/bjr.20180295>

---

© 2018 The Authors. This is an author produced version of a paper subsequently published in *British Journal of Radiology*: Mohamed Ben Taher, Jean Pearson, Marta Cohen, and Amaka C Offiah, Acceptability of post-mortem imaging among Muslim and non-Muslim communities, *The British Journal of Radiology* 2018 91:1091, <https://doi.org/10.1259/bjr.20180295>

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# The British Journal of Radiology

## Acceptability of Post-Mortem Imaging among Muslim and non-Muslim Communities --Manuscript Draft--

<b>Manuscript Number:</b>	
<b>Full Title:</b>	Acceptability of Post-Mortem Imaging among Muslim and non-Muslim Communities
<b>Short Title:</b>	Acceptability of Post-Mortem Imaging
<b>Article Type:</b>	Full Paper
<b>Section/Category:</b>	Diagnostic Radiology
<b>Corresponding Author:</b>	Mohamed Ben Taher Sheffield, UNITED KINGDOM
<b>Corresponding Author Secondary Information:</b>	
<b>Corresponding Author's Institution:</b>	
<b>Corresponding Author's Secondary Institution:</b>	
<b>First Author:</b>	Mohamed Ben Taher
<b>First Author Secondary Information:</b>	
<b>Order of Authors:</b>	Mohamed Ben Taher Jean Pearson Marta Cohen Amaka C Offiah
<b>Order of Authors Secondary Information:</b>	
<b>Abstract:</b>	<p><b>Abstract</b></p> <p><b>Objectives:</b> People's reactions towards autopsy vary according to their cultural and religious beliefs. This paper aims to determine the reaction towards this procedure among Muslims resident in Libya (Group 1) and Muslims (Group 2) and non-Muslims (Group 3) resident in the UK.</p> <p><b>Methods:</b> 400 questionnaires were distributed to the three communities, interrogating belief about post-mortem investigations and after what type of death these were appropriate. Descriptive statistics and non-parametric statistics were used to analyse the data.</p> <p><b>Results:</b> Of the 400 distributed questionnaires, there was a high return rate of 320 (80%). All groups felt that children should be buried sooner than adults (<math>p &lt; 0.001</math>) but 77% of Libyan Muslims thought that children should be buried within 12 hours of death compared to 16% of UK Muslims and only 7% of UK non-Muslims (<math>p &lt; 0.001</math>). More non-Muslims were unconcerned about a negative impact of traditional autopsy on the dignity of the corpse than Muslims (<math>p &lt; 0.001</math>) and more Muslims responded that autopsy has a negative emotional effect on the family (<math>p &lt; 0.001</math>). Type of death altered what sort of investigations were desired. In the case of homicide, Libyan Muslims were less likely to prefer CT (<math>p &lt; 0.001</math>) or MRI (<math>p = 0.001</math>). Sex had no effect on the results of the survey.</p> <p><b>Conclusion:</b> Post-mortem imaging is acceptable to both Muslims and non-Muslims in Libya and the UK but Muslims have a significant preference for post-mortem imaging compared to autopsy, except in homicidal cases.</p> <p><b>Advances in knowledge:</b> *The ability of post-mortem imaging to preserve the dignity of the corpse is independent of religion, however significantly more Muslims feel that autopsy has a negative emotional effect on the family of the deceased</p>

	<p>*A significant majority of Muslims in Libya prefer to bury children within 12 hours of death, while a delay of up to a week is acceptable in the UK</p> <p>*Muslims resident in the UK have an attitude closer to that of the indigenous (non-Muslim) population and therefore educational programmes may be successful in changing attitudes of Muslims in Libya and other predominantly Muslim countries</p>
<p><b>Suggested Reviewers:</b></p>	<p>Non Non image42@gmail.com</p>
	<p>Non Non image44@gmail.com</p>
	<p>Non Non image44@gmail.com</p>
<p><b>Opposed Reviewers:</b></p>	



**Acceptability of Post-Mortem Imaging among Muslim and non-Muslim Communities**

Mohamed Ben Taher<sup>1</sup>, Jean Pearson<sup>2</sup>, Marta Cohen<sup>3</sup>, Amaka C Offiah<sup>1,4</sup>

1 Department of Oncology and Metabolism, University of Sheffield, UK

2 Corporate Information and Computing Services, University of Sheffield

3 Department of Pathology, Sheffield Children's NHS Foundation Trust, Western Bank, Sheffield, UK

4 Department of Radiology, Sheffield Children's NHS Foundation Trust, Western Bank, Sheffield, UK

Correspondence: Mr Mohamed Ben Taher Academic Unit of Child Health, University of Sheffield, Damer Street Building, Sheffield Children's NHS Foundation Trust, Western Bank, Sheffield, S10 2TH

mbentaher1@sheffield.ac.uk

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Introduction

Autopsy is regarded as the gold standard for determining cause of death (1). Clinically important information is discovered in up to 46% of neonatal and infant autopsies and the procedure is important for eliminating abuse as a cause of death (2). In the last three decades, however, autopsy rates across the world have declined by 40-50% (3). This decline has been influenced by religious, cultural and emotional factors (4). Furthermore, fewer parents are willing to consent and fewer clinicians are asking for permission to perform the autopsy (2, 5). Other reasons given for this decline include administrative bottlenecks when requesting an autopsy and delays in providing the autopsy report (6).

Muslims teach that Allah stressed the importance of maintaining the dignity of the body before and after death. Islam, therefore, calls for respect for the body and recommends that the body should not be disfigured, based on the Hadith, “The breaking of the bone of a dead person is like breaking the bone of a live person” (7). Furthermore, Islam requires that the body be buried soon after death, and that there be no cremation (8). An Islamic fatwa (opinion), issued in 1982, however, states that the benefits of autopsy may be greater than its disadvantages, if it serves justice (9).

Over the last three decades, post-mortem imaging has been increasingly used as part of the forensic examination and provides significant information. Post-mortem computed tomography (PMCT) or magnetic resonance imaging (PMMRI) scans can be used to detect some causes of death as an addition to, or instead of, a conventional autopsy (1). Medical imaging is particularly useful when consent for conventional autopsy has been withheld. PMCT offers a rapid method of scanning the whole body (including inside a body bag) (2) and is now widely used in forensic medicine in adults.

Currently, however, neither MRI nor CT are sufficiently accurate to replace an autopsy as the post-mortem (PM) investigation of choice. In recent studies of PMCT, misinterpretation of PM change and/or poor imaging have led to the cause of death being misdiagnosed (10, 11). Caution and the development of expertise in interpretation are therefore required. In their study, Sieswerda-Hoogendoorn and van Rijn (2010) state that CT identifies bone fractures more reliably than autopsy, including sites such as the face, which might be overlooked during an autopsy examination (3). MRI has been shown to be more acceptable to some parents than conventional autopsy for identifying the cause of death of their child (12, 13). Recently, some UK healthcare centres have begun issuing death certificates which include reports of PMMRI findings that are accepted as medicolegal documents (14).

PM imaging has the advantages of being non-invasive and less time consuming than conventional autopsy. Data storage offers the chance to review cases in later years and the ability to highlight areas of interest before (and thereby guiding) the forensic pathology investigation (15). Despite these advantages, conventional autopsy is still the only modality available in Libya for both children and adults. A search of the literature in PubMed, Medline and the Cochrane Systematic Review databases was conducted to identify studies related to the opinion of Muslims on the use of PM imaging to diagnose cause of death; no relevant publications were identified. Prior to the (potentially widespread) introduction of PM imaging to Libya, we aimed to ascertain the opinions and preferences of Muslims and non-Muslims regarding PM imaging (PMCT and PMMRI).

## Methods

### Study Design

This study used a non-validated questionnaire (Figure 1) divided into three sections, the first being related to respondents' demographics. The second section addressed respondents' knowledge of PM examinations (autopsy, CT and MRI). The third and final section explored reasons why PM examinations might be unacceptable to respondents. A total of 400 questionnaires were

1  
2  
3  
4 distributed to adult volunteers as follows: Group 1: 75 Muslim adults attending an out-patient clinic at [REDACTED]  
5 Hospital, which is one of the main hospitals in the central region of Libya and 75 Muslim students and teaching staff at [REDACTED]  
6 University in Libya, Libya; Group 2: 50 Muslim Libyans self-selected from those attending a regular monthly Libyan community  
7 meeting in [REDACTED], UK; Group 3: 200 non-Muslims (110 distributed to members of a community centre in [REDACTED] and 90 to  
8 staff and students at the University of [REDACTED]). The questionnaire was in Arabic for all respondents in Libya and in English for  
9 those respondents resident in the UK.  
10  
11  
12  
13  
14  
15  
16

### 17 Statistical Analysis

18 Descriptive statistics summarise respondents' demographics and non-parametric tests were used to compare between groups and  
19 methods of investigation. Comparison between groups was by the chi-squared test. Monte-Carlo significances were calculated  
20 rather than the traditional asymptotic analysis approach; this removed the need to worry over small sample sizes. For age, a  
21 Kruskal-Wallis test was used and a Mann-Whitney for sex. Cochran-W was used to compare the differences between the  
22 different investigations. Statistical analyses were performed using the Statistical Package for the Social Sciences, version 24  
23 (IBM, Armonk NY). Statistical significance was set at  $p < 0.05$ .

24 Ethical approval was granted through the University of [REDACTED] approval process (Reference Number 007234).  
25  
26

### 27 Results

28 Of the 400 distributed questionnaires, there was a high return rate of 320 (80%) (Table 1, Figure 2). There was a significant  
29 difference in the age of the three groups (KW=  $p < 0.01$ ) with median (90% CI) for Groups 1, 2 and 3 being 30 (27-32), 38 (35-  
30 41) and 27 (24-28) years respectively.  
31  
32

33 In response to whether they had previously heard about post-mortem imaging (PMCT and/or PMMRI), (16% PMCT and 14%  
34 PMMRI) of Group 1, (7% PMCT and 6% PMMRI) of Group 2 and (31% PMCT and 80% PMMRI) of Group 3 answered  
35 positively ( $p < 0.001$ ).  
36  
37

38 For both adult and child burials, differences in opinion concerning an acceptable delay in burial were significant when comparing  
39 the three respondent groups and when comparing all Muslims to non-Muslims ( $p$  ranged from  $< 0.001$  to  $< 0.011$ , Table 2).  
40 Muslims in Libya preferred more rapid burial, particularly for children; 77% of Group 1, 16% of Group 2 and only 7% of Group  
41 3 preferring to bury a child within 12 hours of death ( $p < 0.001$ ).  
42  
43

44 More of Groups 1 and 2 (88% and 91% respectively) than Group 3 (72%) felt that autopsy leads to an unnecessary delay in burial  
45 ( $p < 0.001$ , Table 3). More non-Muslims were unconcerned about the impact of traditional autopsy on the dignity of the corpse  
46 than Muslims (Table 3, Figure 3). The ability of post-mortem imaging to preserve the dignity of the corpse was independent of  
47 religion, however Muslims felt differently about the emotional impact of autopsy, with 93%, 98% and 64% of Groups 1, 2 and 3  
48 respectively, responding that autopsy has a negative emotional effect on the family ( $p < 0.001$ , Figure 4).  
49  
50

51 In terms of identifying the cause of homicidal deaths, 58% of Group 1, 64% of Group 2 and 52% of Group 3 felt that autopsy  
52 should be used ( $p = 0.289$ ). In contrast, 42% of Group 1, 47% of Group 2 and 26% of Group 3 preferred the use of CT to  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 investigate the causes of natural and unexplained death ( $p = 0.004$ ), while no respondent from Group 1, 2% of Group 2 and 16%  
5 of Group 3 preferred conventional autopsy to investigate natural expected deaths ( $p < 0.001$ ). Finally, 31% of Group 1, 51% of  
6 Group 2 and 52% of Group 3 ( $p = 0.001$ ) preferred the use of MRI over autopsy to investigate the causes of suspicious deaths  
7 (Table 4).  
8  
9

10  
11 There was a significant difference in annual income between Muslim and non-Muslim respondents ( $p < 0.001$ ), however, there  
12 was no difference in salary between those who approved of conventional autopsy and those who did not ( $p=0.894$ ).  
13  
14

## 15 **Discussion**

16 The recent Arab uprising has been associated with considerable damage to infrastructure and a significant number of people have  
17 been killed (16); a total of 21,490 persons were killed in Libya between February 2011 and February 2012. Due to the high  
18 number of criminal offences and the limited number of consultant pathologists in Libya, there are difficulties with investigating  
19 and explaining the circumstances of death (17). Post-mortem imaging may play an important role.  
20  
21

22 As far as we are aware, this study is the first to formally evaluate the acceptability of PM CT and MRI of Libyan Muslims and  
23 UK Muslims of Libyan descent with that of non-Muslim UK residents.  
24  
25

26 The fact that the questionnaire had response rates of 92% from Muslims and 69% from non-Muslims shows the importance given  
27 to this subject amongst the respondents. This study shows that both Muslims and non-Muslims perceive conventional autopsy to  
28 have a negative emotional effect on family members due to its invasiveness and (for Muslims) the delay to burial that it causes.  
29  
30

31 Most non-Muslim participants had heard about post-mortem imaging compared to only a small minority of Muslims (mainly  
32 doctors and/or those resident in the UK). By comparison, most Muslim and non-Muslim participants were aware of conventional  
33 autopsy as an investigational procedure. These differences in awareness between Muslim and non-Muslim participants may be  
34 attributed to the fact that several UK healthcare providers now routinely offer post-mortem MRI in children and PMCT in adults  
35 and that some families may have participated in or have been aware of previous/on-going diagnostic accuracy studies of PM  
36 imaging. In Libya on the other hand, conventional autopsy is the only available means of PM investigation. We pre-empted this  
37 lack of awareness when distributing the questionnaires by providing a short background to conventional autopsy, PMCT and  
38 PMMRI as an introduction to the questionnaire.  
39  
40

41 The time to burial of the body is important in the Libyan culture and is an essential issue for the Muslim family of the deceased,  
42 since the culture enforces the religious belief that the body should be buried as soon as possible to reduce the emotional effect on  
43 the family and to respect the deceased. It is clearly demonstrated in this study that Muslim participants in Libya support this view,  
44 with a significant number feeling that burial should occur within 12 hours of death. This contrasts with the views of non-Muslims,  
45 for which no respondent felt that burial was necessary within 24 hours of death and that delays of up to a week were acceptable.  
46  
47 Age appears to be negatively related to time of burial but is best interpreted as an artefact of the data collection, where UK  
48 Muslims tended to be older than Libyan Muslims and UK non-Muslims i.e. there is a lack of mature non-Muslims in our study  
49 population. These views were irrespective of sex. Of interest, more Muslims in the UK had no concerns if burial was delayed for  
50 up to 3 days following death. This might be due to differences in the process of obtaining approval for burial and/or due to their  
51 living in the UK and assimilating the views of that population. Our results support those of Gatrad et al., that Muslims prefer to  
52 bury the body immediately or as soon as possible after death (8, 18). Another study showed that three days is generally considered  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1  
2  
3  
4 the maximum delay before burial of a body in the Muslim world (7), which is in keeping with the attitudes of the Muslims we  
5 surveyed in the UK. Lishimpi et al. (2001), who studied the guardians and parents of deceased children in Zambia, also found  
6 that concerns about time delay before burial had an influence on decisions to refuse an autopsy, although the religion of their  
7 study participants was not provided (19). Only a small number of Muslim participants in our study thought (perhaps incorrectly)  
8 that CT and MRI could lead to unnecessary delay to burial.  
9

10  
11 “Mutilation” of the body is an important reason that might lead a family to refuse a PM examination (9) along with a fear that  
12 organs might be sold for transplantation (19). Less invasive methods such as medical imaging can help maintain the body’s  
13 dignity, which Muslim and non-Muslim participants of this study also believe. On the other hand, non-Muslim participants were  
14 more prepared to accept that a conventional autopsy would not violate the body’s dignity and here, the influence of religion is  
15 clearly seen. This result is supported by Ben-Sasi et al. who pointed out that generally, traditional autopsy was perceived  
16 comparatively favourably (scored 8 out of 10), with certain demographic factors affecting the overall autopsy acceptability,  
17 including ethnicity (more Caucasian and African individuals preferred autopsy compared to Asian or Arabic individuals) and  
18 religion (Christians and those with no religious beliefs found autopsy more acceptable than did those of Muslim or Sikh faiths)  
19 (20). Other studies supporting our results include 1) Lynch, who found that Hindus and those of other religions are considered to  
20 have a less intrinsic objection to autopsy than Muslims (21). 2) Cox et al who reported from Uganda, that 59% of relatives  
21 (Muslim and non-Muslim) were opposed to autopsy for reasons including delayed burial, body mutilation and associated reasons  
22 of a religious nature. Furthermore, the rate of autopsies decreased by approximately 9% due to cultural beliefs and fears that it  
23 might lead to infertility among women (22). 3) Loughrey who showed that relatives and parents might not consider the benefits  
24 of an autopsy and may prefer to “maintain the physical dignity” of their loved one, rather than define the precise cause of death  
25 (23) and 4) Parmar and Rathod, most of whose study participants refused conventional autopsy due to the delay in burial and  
26 concern about the cutting of the body or removal of organs (24). Furthermore, consenting to post-mortem examination, especially  
27 for infants or children, is psychologically distressing for all guardians involved.  
28  
29

30  
31 Despite the overall preference for PM imaging over conventional autopsy, it was interesting to find that Muslim participants  
32 preferred conventional autopsy for the investigation of homicides and PM imaging for the investigation of unsuspecting deaths.  
33 This might imply more belief, by Muslims, in the intrinsic superiority of conventional autopsy to identify the cause of any death.  
34 This might not be a misguided belief, since Hussain et al showed that autopsy explained 78% of cases of homicidal deaths (11).  
35 On the other hand, PMMRI has been shown to be accurate in detecting abnormal pathology in fetuses (sensitivity 77%, specificity  
36 95%), with slightly lower specificity and sensitivity in children (25), while PMCT identified the main pathologic process leading  
37 to death in 39 of 40 adults (26), such that in certain instances, PM imaging is not inferior to conventional autopsy.  
38

39  
40 Our survey included 15 physicians (all Muslim) who did not respond significantly differently to the non-physicians. It has been  
41 shown that some physicians find the request from relatives for PM examination of a loved one, to be one of the most difficult  
42 and unpleasant quarters of paediatric medical practice (22). Interestingly, in their survey of general practitioners and clinicians,  
43 Midelfart and Aase showed that the number of doctors participating in their study who found that the value of autopsy had  
44 decreased due to improvements in CT and MRI techniques was 81% and 71% for each modality respectively (27). We disagree  
45 that the value of autopsy has decreased and do not perceive it as a case of performing one or the other technique, but rather we  
46 believe that PM imaging should be viewed as an adjunct to conventional autopsy and a replacement only when consent for  
47 conventional autopsy is withheld. This is a view we have previously expressed and that is held by others (28, 29, 30).  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 Roberts et al. pointed out that the cost implications of PM imaging may be a concern; MRI in particular is more expensive than  
5 traditional autopsy (31). In Group 1, the average salary was particularly low and in Libya, healthcare is paid for by the individual.  
6 It might be expected that in Libya, caution over the cost of MRI would be a concern, which is congruent with over half of the  
7 people in Group 1 being cautious of using this technique. A similar caution may be applied to CT. Healthcare in the UK is free  
8 at the point of delivery and therefore cost is not necessarily of personal concern to the individual. Sex had no effect on the results  
9 of the survey. As far as we know, there are no published studies that have previously measured the impact of variables such as  
10 sex and income on the acceptance rate of autopsies.  
11  
12  
13  
14  
15

16 In conclusion, religious beliefs and age of the deceased (child versus adult) affect individual preference for PM investigational  
17 methods. The preference amongst Muslims for PM imaging is mainly related to the perception that it leads to less delay to burial  
18 and is less invasive. Interestingly, conventional autopsy is preferred by Muslims when the cause of death is suspicious. Muslims  
19 resident in the UK have an attitude closer to that of the indigenous (non-Muslim) population and therefore in conjunction with  
20 developing expertise in performing and reporting on post-mortem imaging investigations, educational programmes may be  
21 successful in changing attitudes of Muslims in Libya and other predominantly Muslim countries.  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## References

1. Underwood J. Post-mortem imaging and autopsy: rivals or allies? *The Lancet*. 2012;379(9811):100-2.
2. Thayyil S, Chitty L, Robertson N, Taylor A, Sebire N. Minimally invasive fetal postmortem examination using magnetic resonance imaging and computerised tomography: current evidence and practical issues. *Prenat Diagn*. 2010;30(8):713-8.
3. Stawicki S, Aggrawal A, Dean A, Bahner D, Steiberg S, Stehly C. Postmortem use of advanced imaging techniques: Is autopsy going digital. *OPUS*. 2008;12:17-26.
4. Noda Y, Yoshimura K, Tsuji S, Ohashi A, Kawasaki H, Kaneko K, et al. Postmortem Computed Tomography Imaging in the Investigation of Nontraumatic Death in Infants and Children. *Biomed Res Int*. 2013;2013.
5. Sieswerda-Hoogendoorn T, van Rijn RR. Current techniques in postmortem imaging with specific attention to paediatric applications. *Pediatr Radiol*. 2010;40(2):141-52.
6. Oluwasola OA, Fawole OI, Otegbayo AJ, Ogun GO, Adebamowo CA, Bamigboye AE. The autopsy: knowledge, attitude, and perceptions of doctors and relatives of the deceased. *Arch Pathol Lab Med*. 2009;133(1):78-82.
7. Mohammed M, Kharoshah MA. Autopsy in Islam and current practice in Arab Muslim countries. *J Forensic Leg Med*. 2014;23:80-3.
8. Gatrad AR. Muslim customs surrounding death, bereavement, postmortem examinations, and organ transplants. *BMJ*. 1994;309(6953):521-3.
9. Puranik R, Gray B, Lackey H, Yeates L, Parker G, Duflo J, et al. Comparison of conventional autopsy and magnetic resonance imaging in determining the cause of sudden death in the young. *Journal of cardiovascular magnetic resonance : J Cardiovasc Magn Reson*. 2014;16:44.
10. Oyake Y, Aoki T, Shiotani S, Kohno M, Ohashi N, Akutsu H, et al. Postmortem computed tomography for detecting causes of sudden death in infants and children: retrospective review of cases. *Radiat Med*. 2006;24(7):493-502.
11. Hussain Z, Shah MM, Afridi HK, Arif M, Ayub J. Homicidal deaths by firearms in Peshawar: an autopsy study. *J Ayub Med Coll Abbottabad*. 2006;18(1):44-7.
12. Cannie M, Votino C, Moerman P, Vanheste R, Segers V, Van Berkel K, et al. Acceptance, reliability and confidence of diagnosis of fetal and neonatal virtuopsy compared with conventional autopsy: a prospective study. *Ultrasound Obstet Gynecol*. 2012;39(6):659-65.
13. Wichmann D, Obbelode F, Vogel H, Hoepker WW, Nierhaus A, Braune S, et al. Virtual autopsy as an alternative to traditional medical autopsy in the intensive care UnitA prospective cohort study. *Ann Intern Med*. 2012;156(2):123-30.
14. Thayyil S. Less invasive autopsy: an evidenced based approach. *Arch Dis Child*. 2010;archdischild165704.
15. Underwood J. Post-mortem imaging and autopsy: rivals or allies? *The Lancet*. 379(9811):100-2.
16. Daw MA, El-Bouzedi A, Dau AA. Libyan armed conflict 2011: Mortality, injury and population displacement. *Afr J Emerg Med*. 2015;5(3):101-7.
17. news Ao. <http://ara.tv/zcdz5> [
18. Black J. Broaden your mind about death and bereavement in certain ethnic groups in Britain. *Br Med J (Clin Res Ed)*. 1987;295(6597):536.
19. Lishimpi K, Chintu C, Lucas S, Mudenda V, Kaluwaji J, Story A, et al. Necropsies in African children: consent dilemmas for parents and guardians. *Arch Dis Child*. 2001;84(6):463-7.
20. Ben-Sasi K, Chitty L, Franck L, Thayyil S, Judge-Kronis L, Taylor A, et al. Acceptability of a minimally invasive perinatal/paediatric autopsy: healthcare professionals' views and implications for practice. *Prenat Diagn*. 2013;33(4):307-12.
21. Lynch M. Forensic pathology: Redefining medico-legal death investigation. *J Law Med*. 1999;7:67-74.
22. Cox JA, Lukande RL, Kateregga A, Mayanja-Kizza H, Manabe YC, Colebunders R. Autopsy acceptance rate and reasons for decline in Mulago Hospital, Kampala, Uganda. *Trop Med Int Health*. 2011;16(8):1015-8.
23. Loughrey M, McCluggage W, Toner P. The declining autopsy rate and clinicians' attitudes. *Ulster Med J*. 2000;69(2):83.
24. Parmar P, Rathod GB. Study of knowledge, attitude and perception regarding medico-legal autopsy in general population. *Int J Med Pharm Sci*. 2013;3(06):1-6.
25. Arthurs OJ, Owens CM, Olsen OE, Thayyil S, Addison S, Jones R, et al. Diagnostic accuracy of post mortem MRI for abdominal abnormalities in fetuses and children. *Eur J Radiol*. 2015;84(3):474-81.
26. Ross S, Ebner L, Flach P, Brodhage R, Bolliger SA, Christe A, et al. Postmortem whole-body MRI in traumatic causes of death. *AJR Am J Roentgenol*. 2012;199(6):1186-92.
27. MIDELFART J, AASE S. The value of autopsy from a clinical point of view. *Apmis*. 1998;106(7-12):693-8.
28. Lewis C, Hill M, Arthurs OJ, Hutchinson JC, Chitty LS, Sebire N. Health professionals' and coroners' views on less invasive perinatal and paediatric autopsy: a qualitative study. *Arch Dis Child*. 2018;archdischild-2017-314424.
29. Whitby EH, Offiah AC, Cohen MC. Initial experiences of a minimally invasive autopsy service. A report of the techniques and observations in the first 11 cases. *Pediatric and Developmental Pathology*. 2015;18(1):24-9.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

30. Cohen MC, Whitby E, Fink MA, Collett JM, Offiah AC. Running a postmortem service—a business case and clinical experience. *Pediatr Radiol* . 2015;45(4):501-8.

31. Roberts ISD, Benamore RE, Benbow EW, Lee SH, Harris JN, Jackson A, et al. Post-mortem imaging as an alternative to autopsy in the diagnosis of adult deaths: a validation study. *The Lancet*.379(9811):136-42.

32. Smit PR, de Jong RR, Bijleveld CC. Homicide data in Europe: definitions, sources, and statistics. *Handbook of European homicide research*: Springer; 2012. p. 5-23.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Figure Legends:**

Figure 1: Non-validated questionnaire used for this study

\* Homicide is the term used when the cause of death of one person can be attributed to another (32). Natural death occurs in the course of nature with no unusual circumstances. Unexpected deaths are sudden deaths from natural causes. Suspicious deaths may include accidents, murder and suicide.

Figure 2: Distribution of questionnaires/response rate

The figure summarises distribution of the questionnaire and response rate from the various groups of respondents

Figure 3: Post-mortem methods investigations that preserve the dignity

Figure 4: Post-mortem methods lead to a negative emotional effect on the family

**Table Legends:**

Table 1: Respondents' demographics

\* Group 1 = **Muslims in Libya** Group 2 = **Muslims (in UK)** Group 3 = **Non-Muslims (in UK)**

Table 2: Acceptable time to burial of children and adults according to religion and the country of residence of participants

\* Group 1 = **Muslims in Libya** Group 2 = **Muslim (in UK)** Group 3 = **Non-Muslim (in UK)**

Table 3: Respondents' impression of investigations that lead to delay in burial and preserve the dignity of the corpse

Table 4: Respondents' preference for post-mortem investigation depending on nature of death

\* Group 1 = **Muslims in Libya** Group 2 = **Muslims (in UK)** Group 3 = **Non-Muslims (in UK)**

Public Perception Of Using Medical Imaging To Identify Causes Of Death Amongst The Libyan Community In Libya And XXXX.

**Background**

The examination of a body after death to find out why somebody has died is termed post-mortem examination (PM). The traditional and gold standard method for this is autopsy, where body parts are opened up surgically and examined by specialists. An alternative method is conducting this examination using CT (computed tomography) or MRI (magnetic resonance imaging), which do not require the body to be surgically opened.

We are conducting a survey to assess the views and experience of XXXX resident in XXX, and XXXX and non XXX (muslims/non-muslims) resident in XXXX, regarding post-mortem examinations (either autopsy or CT/MRI).

By filling out and returning this questionnaire, you agree to us using your responses for the purposes of our research. We reassure you that it will not be possible for anyone to identify you from the answers that you give.

Thank you for your time in reading the information sheet and in considering whether or not to take part in this study.

**Section 1:**

**About You**

1. **Age (years):** \_\_\_\_\_

2. **Occupation:** \_\_\_\_\_

3. **Sex:**

Male

Female

4. **Religion:**

Muslim

Christian

Hindu

None

Other  Please specify \_\_\_\_\_

5. **Ethnicity:** \_\_\_\_\_

6. **Country of origin:** \_\_\_\_\_

7. **Highest qualification:**

None

University

Secondary School

Primary School

8. **Annual household income:**

< £10600

£10601– £31000

£31001 - £785000

£785001 - £150000

> £150,000

**Section 2:**

**Your Experience of PM (Autopsy, CT or MRI) Examination**

1. Have you come across PM examination (autopsy) before?  
Yes  No
2. Have you heard of PM examination by computed tomography (CT)?  
Yes  No
3. Have you heard of PM examination by magnetic resonance imaging (MRI)?  
Yes  No
4. Do you know anyone (family, relative or friend) who had a PM?  
Yes  No
5. If yes to Question 4, please complete the table below.

**Section 3:**

Cases	Age	Sex	When	Type of Autopsy		
				Conventional autopsy	CT scan	MRI
1.						
2.						
3.						
4.						
5.						

**Your Views on PM Examination (Autopsy)**

1. Performing PM examination leads to a delay in burial. What length of delay do you think is acceptable?

For An Adult:

< 12 hours

12 – 24 hours

1 - 3 days

3 – 7 days

> 1 week

For A Child:

< 12 hours

12 – 24

1- 3 days

3 – 7 days

> 1 week

2. In general, which form of PM would you accept CT/MRI or conventional autopsy in the following situations:

- a. Homicide/suicide cases. CT  MRI  Autopsy  None
- b. Suspicious but natural deaths CT  MRI  Autopsy  None
- c. Natural, but unexplained deaths CT  MRI  Autopsy  None
- d. Natural and expected. CT  MRI  Autopsy  None

3. What concerns you about PM examinations?

- a. Delays in burial.
  - CT
  - MRI
  - Autopsy
  - None
- b. Dignity and sacredness of the body.
  - CT
  - MRI
  - Autopsy
  - None
- c. Emotional burden on the family.
  - CT
  - MRI
  - Autopsy
  - None
- d. Cost effectiveness.
  - CT
  - MRI
  - Autopsy
  - None

Do you have other comments you would like to make or do you have any other anxieties about PM examinations?

.....

.....

Thank you for your time and cooperation. XXXXXX


PhD Student, University of XXXXX. If you have any questions, please do not hesitate to contact me on:  
XXXXXX


Or contact

Supervisor: XXXXX at the XXXXXX Telephone: XXXXX



Figure 2: Distribution of Questionnaires/Response Rate








Table 1: Respondents' Demographics

Analysis Groups *									
Religion		Group 1		Group 2		Group 3		Total	
Age	<= 25	43	31%	0	0%	63	46%	106	33%
	26 - 30	35	25%	6	13%	28	20%	69	22%
	31 - 40	24	17%	21	47%	21	15%	66	21%
	41 - 65	34	25%	18	40%	23	17%	75	23%
	66+	2	1%	0	0%	2	2%	4	1%
Sex	Male	88	64%	31	69%	73	53%	192	60%
	Female	50	36%	14	31%	64	47%	128	40%
Ethnicity	Mixed	138	100%	45	100%	1	1%	184	58%
	White	0	0%	0	0%	86	63%	86	27%
	African	0	0%	0	0%	18	13%	18	6%
	Black	0	0%	0	0%	7	5%	7	2%
	Asian	0	0%	0	0%	24	18%	24	8%
	Other	0	0%	0	0%	1	1%	1	0%
Income	Less than £250	30	2%	4	9%	1	1%	35	11%
	£250 - 500	14	10%	2	4%	0	0%	16	5%
	£500 - 750	30	22%	4	9%	16	12%	50	16%
	£750 - 1000	45	33%	11	24%	37	27%	93	29%
	£1000 - 2000	15	11%	14	31%	16	12%	45	14%
	£2000 - 3000	2	1%	5	11%	48	35%	55	17%
	More than £3000	2	1%	5	11%	19	14%	26	8%

\* Group 1 = Muslims in Libya Group 2 = UK Muslims Group 3 = UK Non-Muslims

Table 2: Acceptable time before burial of children and adults

Religion/ Residence*	Time to Burial (Child)					Time to Burial (Adult)				
	< 12 hrs	12-24 hrs	1-3 days	4 -7days	>7days	< 12 hrs	12-24 hrs	1-3 days	4 -7days	>7days
<b>Group1</b>	101	37	0	0	0	62	54	22	0	0
<b>Group 2</b>	21	13	11	0	0	12	17	16	0	0
<b>Group 3</b>	9	18	43	34	33	4	11	43	29	50

## \* p values:

**Child Burials**

Group 1 Vs Group 2: p &lt; 0.001

Group 2 Vs Group 3: p &lt; 0.001

Muslims (All) Vs Group 3: p &lt; 0.001

**Adult Burials**

Group 1 Vs Group 2: p &lt; 0.011

Group 2 Vs Group 3: p &lt; 0.001

Muslims (All) Vs Group 3 p &lt; 0.001

**Child versus Adult Burials**

Group 1: chi-squared test was statistically significant at p &lt; 0.001

Group 2: chi-squared test was statistically significant at p &lt; 0.001

Group 3: chi-squared test was statistically significant at p &lt; 0.001

\* **Group 1 = Muslims in Libya Group 2 = UK Muslims Group 3 = UK Non-Muslims**

Table 3: Unacceptable delay in burial and preservation of the corpse's dignity

<b>Causes an Unacceptable Delay in Burial n (%)</b>				
<b>Post-mortem method</b>	<b>Muslims in Libya (Group 1)</b>	<b>UK Muslims (Group 2)</b>	<b>UK Non-Muslims (Group 3)</b>	<b>P value</b>
<b>CT</b>	6 (4%)	0 (0%)	27 (20%)	<0.001
<b>MRI</b>	11 (8%)	4 (9%)	32 (23%)	0.001
<b>Autopsy</b>	122 (88%)	41 (91%)	99 (72%)	<0.001
<b>Dignity of the Corpse</b>				
<b>CT</b>	59 (43%)	30 (67%)	70 (51%)	0.019
<b>MRI</b>	85 (62%)	29 (64%)	73 (53%)	0.264
<b>Autopsy</b>	3 (2%)	3 (7%)	22 (16%)	< 0.001

Table 4: PM imaging versus conventional autopsy

<b>Mode of Death</b>				
<b>Homicide/suicide n (%)</b>				
<b>Religion</b>	<b>Group 1</b>	<b>Group 2</b>	<b>Group 3</b>	<b>p value</b>
<b>CT</b>	29 (21%)	15 (33%)	66 (48%)	< 0.001
<b>MRI</b>	30 (22%)	19 (42%)	56 (41%)	0.001
<b>Autopsy</b>	80 (58%)	29 (64%)	71 (52%)	0.289
<b>Natural but unexplained</b>				
<b>Religion</b>	<b>Group 1</b>	<b>Group 2</b>	<b>Group 3</b>	<b>p value</b>
<b>CT</b>	58 (42%)	21 (47%)	35 (26%)	0.004
<b>MRI</b>	55 (40%)	29 (64%)	54 (39%)	0.008
<b>Autopsy</b>	27 (20%)	4 (9%)	73 (53%)	< 0.001
<b>Natural and expected</b>				
<b>Religion</b>	<b>Group 1</b>	<b>Group 2</b>	<b>Group 3</b>	<b>p value</b>
<b>CT</b>	89 (65%)	33 (73%)	21 (15%)	< 0.001
<b>MRI</b>	51 (37%)	17 (38%)	36 (26%)	0.119
<b>Autopsy</b>	0 (0%)	1 (2%)	22 (16%)	< 0.001
<b>Suspicious</b>				
<b>Religion</b>	<b>Group 1</b>	<b>Group 2</b>	<b>Group 3</b>	<b>p value</b>
<b>CT</b>	52 (38%)	18 (40%)	50 (37%)	0.903
<b>MRI</b>	43 (31%)	23 (51%)	71 (52%)	0.001
<b>Autopsy</b>	45 (33%)	16 (36%)	44 (32%)	0.935

\* Group 1 = Muslims in Libya Group 2 = UK Muslims Group 3 = UK Non-Muslims