

This is a repository copy of *Closing the global ozone yield gap: Quantification and co-benefits for multi-stress tolerance*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/138312/>

Version: Accepted Version

Article:

Mills, G, Sharps, Katrina, Simpson, David et al. (8 more authors) (2018) Closing the global ozone yield gap: Quantification and co-benefits for multi-stress tolerance. *Global Change Biology*. pp. 4869-4893. ISSN 1354-1013

<https://doi.org/10.1111/gcb.14381>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Closing the global ozone yield gap: Quantification and co-benefits for multi-stress tolerance

Journal:	<i>Global Change Biology</i>
Manuscript ID	GCB-18-0501.R1
Wiley - Manuscript type:	Primary Research Articles
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Mills, Gina; Centre for Ecology and Hydrology Bangor; University of Gothenburg, Department of Biological and Environmental Sciences, PO Box 461</p> <p>Sharps, Katrina; Centre for Ecology and Hydrology Bangor</p> <p>Simpson, David; Norwegian Meteorological Institute, ; Chalmers University of Technology,</p> <p>Pleijel, Håkan; University of Gothenburg, Department of Biological and Environmental Sciences</p> <p>Frei, Michael; University of Bonn, INRES - Plant Nutrition</p> <p>Burkey, kent; USDA-ARS</p> <p>Emberson, Lisa; Stockholm Environment Institute, Environment</p> <p>Uddling, Johan; University of Gothenburg, Department of Biological and Environmental Sciences</p> <p>Broberg, Malin; University of Gothenburg, Department of Biological and Environmental Sciences</p> <p>Feng, Zhaozhong; Chinese Academy of Sciences, State Key Laboratory of Urban and Regional Ecology</p> <p>Kobayashi, Kazuhiko; The University of Tokyo, Graduate School of Agricultural and Life Sciences</p> <p>Agrawal, Madhoolika; Banaras Hindu University, Department of Botany</p>
Keywords:	Ozone, wheat, soybean, maize, rice, pests and diseases, aridity, heat stress
Abstract:	<p>Increasing both crop productivity and the tolerance of crops to abiotic and biotic stresses are major challenges for global food security in our rapidly changing climate. For the first time, we show how the spatial variation and severity of tropospheric ozone effects on yield compare with effects of other stresses on a global scale, and discuss mitigating actions against the negative effects of ozone. We show that the sensitivity to ozone declines in the order soybean > wheat > maize > rice, with genotypic variation in response being most pronounced for soybean and rice. Based on stomatal uptake, we estimate that ozone (mean of 2010 - 2012) reduces global yield annually by 12.4%, 7.1%, 4.4% and 6.1% for soybean, wheat, rice and maize, respectively (the 'ozone yield gaps'), adding up to 227 Tg of lost yield. Our modelling shows that the highest ozone-induced production losses for soybean are in North and South America whilst for wheat they are in India and China, for rice in parts of India, Bangladesh, China and</p>

	<p>Indonesia, and for maize in China and the USA. Crucially, we also show that the same areas are often also at risk of high losses from pests and diseases, heat stress and to a lesser extent aridity and nutrient stress. In a solution-focussed analysis of these results, we provide a crop ideotype with tolerance of multiple stresses (including ozone) and describe how ozone effects could be included in crop breeding programmes. We also discuss altered crop management approaches that could be applied to reduce ozone impacts in the shorter-term. Given the severity of ozone effects on staple food crops in areas of the world that are also challenged by other stresses, we recommend increased attention to the benefits that could be gained from addressing the ozone yield gap.</p>

SCHOLARONE™
Manuscripts

For Review Only

Closing the global ozone yield gap: Quantification and co-benefits for multi-stress tolerance

Authors: Gina Mills^{1,2*}, Katrina Sharps¹, David Simpson^{3,4}, Håkan Pleijel², Michael Frei⁵,
Kent Burkey⁶, Lisa Emberson⁷, Johan Uddling², Malin Broberg², Zhaozhong Feng⁸,
Kazuhiko Kobayashi⁹ and Madhoolika Agrawal¹⁰

Affiliations

¹Centre for Ecology and Hydrology, Bangor, UK.

²Department of Biological and Environmental Sciences, University of Gothenburg, Sweden.

³EMEP MSC-W, Norwegian Meteorological Institute, Norway.

⁴Department of Space, Earth & Environment, Chalmers University of Technology, Sweden.

⁵Institute of Crop Science and Resource Conservation, University of Bonn, Germany

⁶USDA-ARS, Raleigh, North Carolina, USA

⁷Stockholm Environment Institute, University of York, UK.

⁸State Key Laboratory of Urban and Regional Ecology, Research Center for Eco-

Environmental Sciences, Chinese Academy of Sciences, China.

⁹Department of Global Agricultural Sciences, The University of Tokyo, Japan.

¹⁰Department of Botany, Institute of Science, Banaras Hindu University, India.

***Correspondence** to: Prof. Gina Mills, Centre for Ecology and Hydrology, Environment
Centre Wales, Deiniol Road, Bangor, UK. LL57 2UW. Email: gmi@ceh.ac.uk, Tel. +44
(0)1248 374500.

1 **Running head:** Global benefits of closing the O₃ yield gap

2 **Key words:** Ozone, wheat, soybean, maize, rice, pests and diseases, aridity, nutrient stress,
3 heat stress, stress tolerant ideotype.

4
5 **Paper type:** Primary research

6
7 **Abstract**

8 Increasing both crop productivity and the tolerance of crops to abiotic and biotic stresses are
9 major challenges for global food security in our rapidly changing climate. For the first time,
10 we show how the spatial variation and severity of tropospheric ozone effects on yield
11 compare with effects of other stresses on a global scale, and discuss mitigating actions
12 against the negative effects of ozone. We show that the sensitivity to ozone declines in the
13 order soybean > wheat > maize > rice, with genotypic variation in response being most
14 pronounced for soybean and rice. Based on stomatal uptake, we estimate that ozone (mean of
15 2010 - 2012) reduces global yield annually by 12.4%, 7.1%, 4.4% and 6.1% for soybean,
16 wheat, rice and maize, respectively (the ‘ozone yield gaps’), adding up to 227 Tg of lost
17 yield. Our modelling shows that the highest ozone-induced production losses for soybean are
18 in North and South America whilst for wheat they are in India and China, for rice in parts of
19 India, Bangladesh, China and Indonesia, and for maize in China and the USA. Crucially, we
20 also show that the same areas are often also at risk of high losses from pests and diseases,
21 heat stress and to a lesser extent aridity and nutrient stress. In a solution-focussed analysis of
22 these results, we provide a crop ideotype with tolerance of multiple stresses (including ozone)
23 and describe how ozone effects could be included in crop breeding programmes. We also

1 discuss altered crop management approaches that could be applied to reduce ozone impacts in
2 the shorter-term. Given the severity of ozone effects on staple food crops in areas of the
3 world that are also challenged by other stresses, we recommend increased attention to the
4 benefits that could be gained from addressing the ozone yield gap.

6 **1. Introduction**

7 To feed the rapidly growing global population, we need to develop a new generation of crop
8 cultivars or varieties that will have both high productivity in future climates and high
9 tolerance of the biotic and abiotic stresses that are likely to become more prevalent in the
10 future (Gilliham *et al.*, 2016). Candidate characteristics or traits are currently being tested in
11 ideotype modelling (Semenov and Stratonovitch, 2013) and include improved light
12 conversion efficiency, a longer duration of green leaf area for grain fill, a higher harvest
13 index and optimal phenology. For example, varieties that use less water per unit of carbon
14 fixed will have higher yield under drought conditions (Rebetzke *et al.*, 2002) as will those
15 with ‘stay-green’ characteristics during water stress (Jordan *et al.*, 2012). Whilst it is widely
16 recognised that rapid breeding programmes will have a vital role to play in adaptations of
17 crops to climate change (Atlin *et al.*, 2017), selection of traits for tolerance of one abiotic
18 stress, tropospheric (ground level) ozone pollution, is currently omitted from such breeding
19 programmes (Ainsworth, 2016; Frei *et al.*, 2015). This is happening even though field
20 experiments from nine countries representing three continents have shown that reducing
21 ozone concentrations back to pre-industrial levels would give an average wheat yield benefit
22 of 8.4% globally (Pleijel *et al.*, 2018), a figure that is matched by modelling based on the
23 stomatal uptake of the pollutant (Mills *et al.*, 2018a). Furthermore, an earlier meta-analysis of
24 crop responses to ozone suggested that current ozone levels in the range 31 - 50 ppb (nmol

1 mol^{-1} , v/v) are reducing the yield of major food crops by 5.3 to 19% (Feng & Kobayashi,
2 2009). We undertook this new study to build a case for improving crop yields in our
3 changing climate by closing the ozone-induced yield gap via the inclusion of ozone tolerance
4 in crop breeding programmes, altered crop management and more stringent ozone precursor
5 emission controls.

6 Tropospheric ozone pollution is formed from photochemical reactions involving
7 anthropogenic and biogenic emissions and is involved in a complex web of interactions with
8 ecosystems (Simpson *et al.*, 2014). Whilst concentrations have been beginning to decrease in
9 eastern USA and parts of Europe (2000 – 2014) due to precursor emission controls, they have
10 been increasing rapidly in south (S) and east (E) Asia (Chang *et al.*, 2017). Ozone is a
11 powerful oxidant that is absorbed into leaves via open stomatal pores. Once inside the
12 leaves, ozone reacts with biomolecules to form reactive oxygen species, triggering defence
13 mechanisms that if overwhelmed lead to programmed cell death and a reduced extent and
14 duration of functional green leaf area producing less photosynthate for seed fill (e.g.
15 Ainsworth, 2016). Since pests and diseases (e.g. Oerke, 2006; Huysmans *et al.*, 2017), heat
16 stress (e.g. Driedonks, *et al.*, 2016), drought (e.g. Farooq *et al.*, 2017) or reduced nutrient
17 availability (e.g. Gastal & Lemaire, 2002) usually also reduce the extent and duration of the
18 functional green leaf area, then in simple terms, each of these biotic and abiotic stresses result
19 in the same endpoints – reduced yield quantity that is often associated with reduced quality.

20 So far, most crop breeding programmes have been targeted at increasing or maintaining the
21 yield rather than increasing stability of yield under stress (Gilliham *et al.*, 2017). Because
22 ozone concentrations tend to be very heterogeneous across natural and agricultural regions
23 (Klingberg *et al.*, 2012) as well as over seasons and years, it is not likely that traditional
24 selection would unintentionally favour ozone tolerant crop genotypes. The reverse seems to
25 be the case. For example, an analysis of ozone-exposure yield data for 49 soybean varieties

from 28 field exposure studies showed that ozone sensitivity has increased by an average of 33% between 1960 and 2000 (Osborne *et al.*, 2016). Similarly, modern wheat varieties are more sensitive than older varieties (Biswas *et al.*, 2008; Pleijel *et al.*, 2006). Potentially, this increased sensitivity to ozone over recent decades is related to selective breeding for higher stomatal conductance (Roche, 2015) that inadvertently has increased the ingress of ozone into crops (Biswas *et al.*, 2008; Osborne *et al.*, 2016); further study is required to fully understand the mechanistic basis of this increasing sensitivity with time.

As with many abiotic and biotic stresses, genetic variation in plant response to ozone has been found for every species that has been tested. For the major grain crops, genetic variation in ozone response has been reported for wheat (Zhu *et al.*, 2011), rice (Frei *et al.*, 2008; Shi *et al.*, 2009), soybean (Mulchi *et al.*, 1988; Burkey & Carter, 2009; Jiang *et al.*, 2018), and maize (Yendrek *et al.*, 2017). Variation has also been reported for other crops including snap bean (Burkey *et al.*, 2005; Yuan *et al.*, 2015) and tobacco (Heggestad, 1991). These assessments are based on different criteria including foliar injury and impacts on growth and yield parameters. Taken together, the evidence suggests that sufficient natural genetic variation exists to support improvement in crop stress tolerance either as sources of ozone tolerance genes or providing contrasting genotypes for mechanism studies to identify targets for molecular manipulation. Potential targets for breeding of ozone tolerance that have the greatest likelihood of success include reducing the stomatal uptake of ozone into the leaf and increasing its detoxification once inside the leaf (Feng *et al.*, 2016; Frei *et al.*, 2015).

To target the regions of the world where ozone tolerant crop varieties are most required, we need to understand which crops are most at risk and where they are growing in relation to current high-risk areas for ozone. We know from a recent analysis of ozone concentrations at over 3000 rural sites that the highest ozone values are in many of the world's important crop growing regions, including parts of the USA, Europe, India and China (Mills *et al.*, 2018b).

1 Overall, the latter study showed that the global mean cumulative ozone exposure is double
2 the critical level set by the United Nations as a target for ozone pollution control, above
3 which direct adverse effects on sensitive vegetation may occur according to present
4 knowledge (CLRTAP, 2017). Several studies have modelled ozone concentrations and
5 predicted yield effects using concentration-based yield response functions applied at a range
6 of scales from local (e.g. for India, Lal *et al.*, 2017) to global (e.g. Avnery *et al.*, 2011a,b;
7 Van Dingenen *et al.*, 2009). Whilst these studies indicate effects in the highest ozone areas,
8 they do not take into account the constantly varying effects of soil moisture, air temperature,
9 light and humidity on the uptake of the pollutant via the stomata. In Europe, field evidence
10 for effects of ozone on crops and other types of vegetation shows that risk assessments based
11 on modelled stomatal uptake or flux (Emberson *et al.*, 2000, Simpson *et al.*, 2007) provide a
12 stronger indication of ozone effects than those based on concentration (Mills *et al.*, 2011).
13 Furthermore, dose-response functions for crops that are based on stomatal uptake are better
14 correlated with yield effects than those based on concentration (Pleijel *et al.*, 2000, 2007),
15 providing additional support for their use.

16 With ozone concentrations increasing in rapidly developing regions and predicted to continue
17 to increase in coming decades (Wild *et al.*, 2012), it is timely to consider the options for
18 increasing the tolerance of crops to this abiotic stress. In this study, our analysis included a
19 two-step approach to addressing the ozone problem in crops: (i) a quantitative spatial analysis
20 of the impacts of ozone on crop yield relative to impacts of other abiotic and biotic stresses
21 and (ii) a qualitative analysis of crop traits, including defining an ideotype with multiple
22 stress tolerance. As an initial step, we compiled dose-response data from experiments
23 conducted around the world to determine the scope for breeding ozone tolerant varieties by
24 showing the genotypic range in sensitivity for four staple crops: soybean, wheat, rice and
25 maize. We then used the response functions to model the current impacts of ozone on each

1 crop, showing the regions where the greatest production losses are likely to be occurring.
2 Whilst we wait for ozone effects to be included in predictive crop yield modelling (as
3 suggested by, for example, Challinor *et al.*, 2009; Emberson *et al.*, 2018; Lobell & Asseng,
4 2017), we sought to compare on a global scale the impacts of ozone on yield with the
5 influence of other biotic and abiotic stress. Those selected were: pests and diseases (Oerke *et al.*,
6 2006); aridity (Trabucco & Zomer, 2009); heat stress (developed from Deryng *et al.*
7 (2014) and Teixeira *et al.* (2013)); and soil nutrient stress (GAEZ). The effects of all five
8 stresses were considered in more detail for India where there are major challenges for crop
9 production and food security (Jaswal, 2014) and where global assessments consistently
10 predict high risk from elevated ozone (e.g. Avnery *et al.*, 2011a,b; Van Dingenen *et al.*,
11 2009). In the second part of the study, we conducted an analysis of the plant traits associated
12 with multiple stress tolerance, and considered the trade-offs and benefits of introducing ozone
13 tolerance in crops for cross-tolerance of other biotic and abiotic stresses. This part of the
14 study culminated in the design of an ideotype for an ozone tolerant crop that would also
15 provide tolerance of co-occurring stresses. In an extended discussion, we assess the results
16 from the two parts of the study and consider viable options for reducing the negative effects
17 of ozone on yield, including crop management, breeding and global efforts to reduce ozone
18 pollution.

19

20 **2. Materials and Methods**

21 **2.1 Global spatial analysis of crop yield constraints caused by ozone**

22 **2.1.1 Crop production**

23 Global modelled crop production data (year 2000, 0.0833° (5 arc minute) resolution) was
24 downloaded from the GAEZ (Global Agro-Ecological Zones, v. 3) data portal

(<http://www.fao.org/nr/gaez/en/>) for soybean, wheat, rice and maize. Irrigated and rain-fed production data was collected for each crop. Using ArcMap v. 10.3 (Environmental Systems Research Institute, Redlands, CA, USA), a 1° by 1° global grid was created. For each crop, production was summed per grid cell. Each cell was classed as irrigated or non-irrigated based on the percentage of irrigated crop production per cell. To define a threshold for irrigated versus non-irrigated, we first produced frequency distributions of the percentage of irrigated production for each crop (Supporting Information, Fig. S1). These showed that the majority of cells for each crop were either fully irrigated or fully rain-fed. A threshold of 75% irrigated was used to identify those cells where the majority of the production was on irrigated land. Production for the period 2010-12 was estimated per grid cell by applying a conversion factor from FAOSTAT national production data available, averaged for the years 1999 – 2001 (average production for 2010-12/average production for 1999-2001). Only cells with > 500 tonnes (0.0005 Tg) crop production in 2010-12 were included in the analysis.

As discussed in Mills *et al.* (2018a), each 1° by 1° grid cell was assigned to a climatic zone, using the global ‘Climatic Zone’ GIS raster layer produced by the European Soil Data Centre (ESDAC) at JRC (Joint Research Centre). For each climatic zone, a 90 day growing period was derived per crop (Table S1), with climatic zones illustrated in Fig. S2. Data sources for assigning crop timings are provided with Table S1. For ease of comparison of effects between crops, only the main growing season per year was used for each crop.

2.1.2 Intra- and inter-specific sensitivity of crops to ozone

To determine the relative sensitivity of the four crops to ozone together with the between-variety variation in response to the pollutant, it was necessary to update existing response functions based on ozone concentration as stomatal uptake-based functions are currently only available for wheat. We collated dose-response data from the scientific literature using the

1 method developed by Osborne *et al.* (2016) for soybean and the commonly reported ozone
2 metric, M7 (7 hour mean, averaged from 09:00 to 15:59). The soybean dose response
3 relationship from Osborne *et al.* (2016) was included in our analysis, whilst response
4 functions for wheat, rice and maize provided in Mills & Harmens (2011) were updated with
5 more recent published data (Web of Science and Google Scholar searches conducted between
6 April and October 2017 using the search terms "ozone and yield and *crop name*"). Studies
7 were only included if they met a number of selection criteria. The duration of ozone exposure
8 must have spanned at least 60% of the 90 day growing season for each crop and ozone levels
9 during exposure were up to 100 ppb for wheat and 170 ppb for other crops. Experiments were
10 included if carried out in Open Top Chambers (OTCs), ambient air or large closed
11 chambers/greenhouses (with the air stirred by fans, minimum size 2.6 by 2.2m). Data from
12 both container and field-sown experiments were used to ensure a wide variety of points from
13 different varieties were included. If the seasonal M7 was not given in the text, this was
14 calculated either using the conversion equations provided in Osborne *et al.* 2016 (e.g. for 24
15 hour mean to M7) or information contained in the experimental methodology of the study. As
16 there was no new published data available at the time of analysis for maize, the response
17 function from Mills & Harmens (2011) was used. Yield data from different experiments were
18 standardised as first described by Fuhrer (1997) and recently re-described by Osborne *et al.*
19 2016. Thus, for each set of experimental data, linear regression was used to determine the
20 yield at 0 ppb of ozone (the intercept of the line); this value was the reference for calculating
21 the relative yield (i.e. relative yield = actual yield/yield at 0 ppb).

22 Individual variety dose-response functions were derived for wheat and rice for the four
23 varieties with the most data points. Following Osborne *et al.* 2016, yield reduction estimates
24 ($RYL_{c,p}$) were then calculated for varieties showing statistically significant declines in yield

1 with increasing ozone by calculating the difference in percentage yield loss at 55ppb
2 (representing current M7) relative to that at 23 ppb (representing pre-industrial M7).

3 **2.1.3 Yield constraints caused by ozone**

4 The EMEP MSC-W (European Monitoring and Evaluation Programme, Metereological
5 Synthesising Centre-West) chemical transport model (version 4.16, Simpson *et al.*, 2012,
6 2017) was used to derive daily POD_3IAM (Phytotoxic Ozone Dose above $3 \text{ nmol m}^{-2} \text{ s}^{-1}$,
7 parameterised for Integrated Assessment Modelling, CLRTAP, 2017) values for the years
8 2010 to 2012 per 1° by 1° grid cell as described by Mills *et al.* (2018a). POD_3IAM is
9 parameterised for a generic crop represented by wheat (CLRTAP, 2017) and represents the
10 accumulated stomatal uptake of ozone, modelled from the hourly mean values for ozone,
11 temperature, vapour pressure deficit, irradiance and soil moisture (Mills *et al.* 2018a).
12 Evaluation of the EMEP model performance is also presented in Mills *et al.*, 2018a, and is
13 summarised in the Supporting Information for the current paper (T1).
14 For each crop, the accumulated 90 day POD_3IAM was then calculated per cell using
15 appropriate climate-specific 90 day growing periods (Table S1, Fig. S2), and an average
16 calculated for the period 2010-2012. For example, for soybean in warm temperate climates in
17 the Northern Hemisphere, the time interval was day 182 to day 271. The EMEP model
18 generated irrigated (without soil water limitation) and non-irrigated (rain limited) POD_3IAM
19 values. For grid cells classed as irrigated for each crop (See Section 2.1.1), the irrigated
20 POD_3IAM value was used to calculate percentage yield loss, otherwise the non-irrigated
21 POD_3IAM was used. This approach allowed crop-specific irrigation usage to be taken into
22 account, and was different to Mills *et al.* (2018a) where POD_3IAM values were weighted by
23 the proportion of irrigation use within a $1 \times 1^\circ$ cell. The global distribution of POD_3IAM for
24 each crop is provided in Fig. S3.

Yield loss due to ozone was first calculated for wheat using the most recent methodology adopted by CLRTAP, 2017. This method also differed slightly from that used in our earlier study (Mills *et al.*, 2018a) in that a reference POD₃IAM value to represent ozone uptake at pre-industrial or natural ozone levels was subtracted before crop loss was calculated (CLRTAP, 2017). This value (0.1 mmol m⁻², Equ. 1) was the mean POD₃IAM for the experimental conditions included in the dose-response relationship, assuming constant 10 ppb ozone throughout the 90d period. The equation used to determine percentage yield loss was:

$$\% \text{ Yield loss} = (\text{POD}_3\text{IAM} - 0.1) * 0.64 \quad [\text{Equ. 1}]$$

Where 0.64 is the slope of the relationship between POD₃IAM and percentage yield reduction (Mills *et al.*, 2018a) and represents the percentage reduction per mmol m⁻² POD₃IAM.

For soybean, maize and rice, the climate-specific grid square POD₃IAM values were first used to calculate yield loss using the wheat equation (Equ. 1), and the resultant value was then multiplied by the relative sensitivity of the crop compared to wheat, RS_w. The latter was derived by dividing the slope of the M7 response function for the crop (Fig. 1) by that for wheat. Production loss per crop was calculated per grid square using the following equation:


$$\text{Production loss (tonnes)} = \text{Crop production} * (\% \text{ yield loss}/100) \quad [\text{Equ. 2}]$$

2.2 Global spatial analysis of yield constraints caused by other stresses

2.2.1 Yield constraints caused by pests and diseases

Oerke *et al.* (1994; 2006) provide estimates for pre-harvest crop losses due to weeds, animal pests, (arthropods, nematodes, mammals, slugs and snails, birds), pathogens and viruses for several major global crops, using data compiled from the literature. This database provides regional percentage yield loss estimates up to 2004 for 11 crops, including soybean, wheat, rice and maize, and is available from the Centre for Agriculture and Biosciences International

(CABI) Crop Protection Compendium (CABI, 2005). A value for mean percentage yield loss due to pests and diseases for the period 2002-04 was assigned to each 1° by 1° grid cell, based on the country and region of the world the cell was located in. If a cell contained land from more than one country, it was assigned to a country based on where the majority of the crop was growing in the cell. Data was available for 19 global regions (Oerke *et al.*, 2006). In this study, data were used that represented the remaining crop yield losses after crop protection practices had been applied.



8

Fig. 1. Response functions for (a) soybean, (b) wheat, (c) rice and (d) maize derived from published data using the growing season ozone (7h mean, M7 in ppb) in the experiments. Data points are presented per cultivar/variety, with sources of data provided in the Supporting Information (Table S3). The response functions are: Soybean, $RY = -0.0050x + 1.001$ (r^2 (adj) = 0.625, $p < 0.001$); Wheat, $RY = -0.0048x + 0.96$ (r^2 (adj) = 0.547, $p < 0.001$); Rice, $RY = -0.0021x + 0.987$ (r^2 (adj) = 0.347, $p < 0.001$); and Maize, $RY = -0.0031x + 1.03$ (r^2 (adj) = 0.617, $p < 0.001$).

16

2.2.2 Yield constraints caused by soil nutrients

Soil nutrient classifications (nutrient availability and nutrient retention) at 0.083° by 0.083° resolution were downloaded from the GAEZ (v. 3) data portal (<http://www.fao.org/nr/gaez/en/>) in June, 2017. The soil qualities (nutrient availability and retention) in the GAEZ dataset have been derived from combinations of soil attributes, using data in the Harmonized World Soil Database (HWSD, v. 1.1, FAO/IIASA/ISRIC/ISS CAS/JRC 2009). Nutrient availability refers to soil fertility, and classification is based on soil texture, soil organic carbon, soil pH and total exchangeable bases. The nutrient retention capacity of soil is based on the ability of soil to retain added nutrients against losses due to leaching. Classification of nutrient retention has been derived from soil texture, base saturation, cation exchange capacity of the soil and of the clay fraction and soil pH. In the GAEZ dataset, nutrient availability and retention are classed separately for topsoil (0-30cm) and subsoil (30-100cm) and then combined by weighting based on the prevalence of active roots (Fischer *et al.*, 2012). The GAEZ classes for soil nutrient availability and nutrient retention were combined in this study to produce five soil nutrient stress classes (summarised in Table 1, further details provided in Table S2). The soil nutrient class making up the majority of each 1° by 1° grid cell in areas where crops were growing, was used to represent the class for each cell.

2.2.3 Yield constraints caused by heat stress

Following the methods of Challinor *et al.* (2005), subsequently used by a number of other studies (e.g. Deryng *et al.*, 2014; Teixeira *et al.*, 2013), a heat stress index was calculated per grid cell for each crop to determine if the daily temperature within a 30 day thermal sensitive period (TSP) exceeded the tolerance thresholds for each crop. This method assumes that damage to crops occurs when daily temperatures exceed a critical temperature (T_{crit} , °C) and

1 maximum damage occurs when temperatures exceed the limit temperature (T_{lim} °C). Using
 2 information on the reproductive phase for each crop (FAO), the thermal sensitive period was
 3 designated as days 40-70 of the 90 day growing period (which varies with climate zone for
 4 each crop, Table S1). Following Deryng *et al.*, (2014), the daily effective temperature (T_{eff} ,
 5 °C, (daily mean temp + daily max temp)/2), used as a measure of the daily temperature when
 6 photosynthesis is taking place, was calculated per grid cell using global hourly temperature
 7 data for the period 1990-2014 at 0.5° by 0.5° resolution. The temperature data were from the
 8 European Centre for Medium-Range Weather Forecasts Integrated Forecasting System
 9 (ECMWF-IFS, www.emwf.int/research/ifsdocs/), as prepared for use by the EMEP model.
 10 For each crop, a daily heat stress value (f_{HSD}) was then calculated for each day within the
 11 TSP, per grid cell. As we required an index that could be used to detect increasing levels of
 12 stress (i.e. an index scaled from 0 to 1), heat stress was calculated following Teixeira *et al.*
 13 (2013) (Eqn. 3).

$$f_{HSD} = \begin{cases} 0 & \text{for } T_{eff} < T_{crit} \\ \frac{T_{eff} - T_{crit}}{T_{lim} - T_{crit}} & \text{for } T_{crit} \leq T_{eff} < T_{lim} \\ 1 & \text{for } T_{eff} \geq T_{lim} \end{cases} \quad [\text{Equ. 3}]$$

19 An average value was then calculated across the 30 day TSP to give the final heat stress
 20 index value, (f_{HS}) per grid cell (Eqn. 4).

$$f_{HS} = \frac{\sum_{j=1}^{TSP} (f_{HSD})}{TSP} \quad [\text{Equ. 4}]$$

1 Critical and limiting temperatures per crop were taken from Deryng *et al.* (2014) (maize,
2 wheat and soybean) and Teixeira *et al.* (2013) (rice). These were: Soybean (35°C for T_{crit} and
3 40°C for T_{lim}), wheat (25 and 35 °C), rice (35 and 45°C) and maize (32 and 45°C).

4 **2.2.4 Yield constraints caused by aridity**

5 Global Aridity Index data (Trabucco & Zomer 2009) were downloaded from the CGIAR-CSI
6 GeoPortal (<http://www.csi.cgiar.org>). The mean Aridity Index for the period 1950-2000
7 (0.0083° by 0.0083° resolution) was calculated as:

$$8 \text{ Aridity Index (AI)} = \text{MAP} / \text{MAE} \quad [\text{Equ. 5}]$$

9 Where MAP is the Mean Annual Precipitation and MAE is the Mean Annual Potential
10 Evapo-Transpiration.

11 Mean annual precipitation values were obtained from the WorldClim Global Climate Data
12 (Hijmans *et al.* 2004), for years 1950-2000, while mean annual values of Potential Evapo-
13 Transpiration (PET) were calculated using the average monthly PET values from the Global-
14 PET model (Trabucco & Zomer, 2009). The mean Aridity Index per cell was calculated for
15 each 1° by 1° grid cell where there is production for the crop.

17 **2.3 Comparative analysis of effects of five stresses using a Yield Constraint Score (YCS)**

18
19 As percentage yield loss data was only available for ozone and pests and diseases data, a
20 percentage scale could not be used for all stresses. To overcome this problem, a yield
21 constraint score (YCS) on a scale of 1 – 5 was developed for each abiotic and biotic stress to
22 show spatially where each constraint is predicted to be impacting on yield and to provide
23 some indication of the magnitude of the effect (Table 1). Yield loss was split into the same

1 five percentage yield loss classes for ozone and pests and diseases, with the highest class
2 being >40% and expected to be comparable to severe stress for all yield constraints.
3 Soil nutrient retention and availability were combined to give five overall classes (Table S2).
4 The aridity climate classes used were from the Generalized Climate Classification Scheme
5 (UNEP 1997), while the heat stress index was classified following the methods of Teixeira *et*
6 *al.* (2013). To identify those areas of the world with the highest combined stresses, the YCS
7 for all five stresses were summed (YCS_{all}).

8
9 **Table 1: Categories of Yield Constraint Score (YCS) for ozone, pests and diseases,**
10 **soil nutrients, heat and aridity** (See text for explanations and justifications of
11 categories).

Stress	Attribute	Year(s) of data	Yield constraint score (YCS)				
			1	2	3	4	5
Ozone	% Yield Loss	Mean of 2010-2012	0 to 5	5 to 10	10 to 25	25 to 40	> 40
Pests and diseases	% Yield Loss	Mean of 2002 - 2004	0 to 5	5 to 10	10 to 25	25 to 40	> 40
Soil Nutrients	Retention	HWSD data, 2009, downloaded in June, 2017	None or slight	Slight to moderate	Slight to severe	Moderate to severe	Moderate to very severe
	Availability		None	Slight to moderate	Moderate to severe	Severe	Severe to very severe
	Overall		None	Slight	Moderate	Severe	Very severe
Heat	Index	Mean of 1990 - 2014	0	<0.05	0.05 to 0.15	0.15 to 0.3	> 0.3
Aridity	Index	Mean of 1950 - 2000	>0.65	0.5 to 0.65	0.2 to 0.5	0.03 to 0.2	<0.03
	Climate class		Humid	Dry sub-humid	Semi-arid	Arid	Hyper arid

For description of effects of the five stresses, results are described as regional and national averages, with the mean YCS and YCS_{all} rounded to the nearest integer, reflecting their categorical nature. The regional classification of countries used is that adopted by the Task Force on Hemispheric Transport of Air Pollutants (HTAP) of the Convention on Long-range Transboundary Air Pollution (LRTAP, Dentener & Guizzardi, 2013). Region names are provided in full in the text the first time they are used and thereafter are referred to by the HTAP three letter codes. The region names, three letter codes and a map illustrating the countries included per region are provided in Fig. S4.

2.4 Qualitative analysis of plant traits associated with multiple stress tolerance

The scientific literature on crop stress tolerance was reviewed between June and December, 2017, with the aim of developing an ideotype for an ozone- and multi-stress tolerant crop. This analysis identified target traits to induce ozone tolerance, including reducing the effects on panicles, leaves and roots. It also considered the benefits and trade-offs for tolerance of other stresses, of introducing ozone tolerance into crops.

3. Results

3.1 Quantification of the global impacts of ozone and other stresses on crop yield

3.1.1 Intra-specific sensitivity to ozone

A comprehensive collation of published data on the yield responses of soybean, wheat and rice to ozone resulted in a database representing 52, 18 and 44 varieties, respectively (Fig. 1a-c, with data sources in Table S3). Ozone-response data for these three crops provides good representation of the areas where the crops were grown: Soybean (East Asia (EAS), North America (NAM) and South Asia (SAS)); wheat (Europe (EUR), EAS, SAS) and rice (EAS and SAS). In contrast, only three varieties have been tested to date for yield responses in maize (Fig. 1d), with these experiments being conducted in the USA during the 1980s and

1 early 1990s. For each crop, there was a significant negative response to ozone ($p < 0.001$),
 2 with the slope of the negative relationships declining in the order soybean (-0.0050) > wheat
 3 (-0.0048) > maize (-0.0031) > rice (-0.0021). Within each response function, variation in
 4 ozone sensitivity due to variety provided scatter in the range of sensitivity.

5 For each crop, some varieties were more tolerant to ozone than others, indicating that there is
 6 scope for selecting more tolerant varieties for immediate use or as part of a breeding
 7 programme for new varieties. For soybean, $RYL_{c,p}$ ranged from 13.3% to 37.9%, with the
 8 three most sensitive varieties being the Indian varieties 'PK472', 'Pusa 9712' and 'Pusa
 9 9814' (Osborne *et al.* 2016). For wheat, the $RYL_{c,p}$ for the four varieties with the most data
 10 was 16.4% ('Drabant'), 19.3% ('PBW 343'), 26.4% ('Dragon') and 32.5% ('Albis'). While
 11 the 44 rice varieties showed a range of sensitivities to ozone (Fig. 1c), overall, rice was the
 12 least sensitive of the four crops investigated. Of the four rice varieties with the most data,
 13 'Koshihikari' showed a $RYL_{c,p}$ of 4.7%, 'Nipponbare' showed no significant negative
 14 relationship between relative yield and M7 ($p > 0.05$), while 'Kasalath' and 'Kirara 397' had a
 15 $RYL_{c,p}$ of 6% and 11.1 % respectively. The rice variety 'Pathumthani-1' showed a higher
 16 $RYL_{c,p}$ of 18.1%, however only 5 data points were available, therefore further study may be
 17 required to confirm this result. For maize, the $r^2(\text{adj})$ was 0.62 for the response function
 18 ($p < 0.001$, Fig. 1d), with a $RYL_{c,p}$ for all three varieties of 10%. The $RYL_{c,p}$ for 'Pioneer
 19 3780' with 14 data points was 15.7% whilst $RYL_{c,p}$ was not calculated for 'PAG 397' as the
 20 response function for the 5 data points for this variety was not significant ($p = 0.08$).

21 **3.1.2 Spatial analysis of the global impacts of multiple stresses on crop yield**

22 Soybean

23 The highest ozone-associated production losses (in Tg per $1 \times 1^\circ$ grid square) for soybean are
 24 predicted to be in NAM and South America (SAM) (Table 2), particularly in central and E

1 USA, S Brazil and N Argentina (Fig. 2, Table S4). However, the percentage yield losses are
2 predicted to be lower for Brazil (12.5 – 15%) and Argentina (7.5- 10%) than for the USA (>
3 20% in large areas), showing that in high producing areas where ozone concentrations are
4 more moderate (as indicated by the percentage losses), high total production losses can still
5 be expected. In the rest of the world, production losses due to ozone in excess of 0.01 Tg per
6 1 x 1 ° grid square are predicted for parts of China, India and S and E Europe. In each of
7 these areas, the production loss was not as high as expected from percentage yield losses in
8 excess of 20%, because soybean is not widely grown.

9 The YCSs for ozone were mainly score 3 for the highest producing regions, with some areas
10 with a score of 4 in E USA, NE India and China (Fig. 2, Tables 2 and S4). There is overlap
11 between these areas and the areas with the highest YCS_{all}. Other areas with a relatively high
12 YCS_{all} such as parts of Sub-Saharan Africa (SSA) and SEA are not predicted to have high
13 production losses due to ozone because of lower percentage yield losses and/or low
14 production totals per region. For soybean, the YCS for pests and diseases is 3 or more over
15 most of the growing area, and particularly high (score of 5) in parts of SSA, SAS and SEA.
16 The largest YCS values for nutrient availability (scores of 4 and 5) are in areas of SE USA, S
17 Brazil and SEA including Thailand, Malaysia and Indonesia. Whilst heat stress YCSs are
18 lower than those for aridity, the areas affected by both stresses largely coincided in soybean
19 growing areas.

20 Overall, for soybean, the global mean YCS for ozone of 3 is one category below that for pests
21 and diseases, and one higher than that for nutrients and aridity (Table 2). The mean YCSs for
22 ozone were in the range 2 - 3 for the five highest producing regions (SAM, NAM, EAS, SAS,
23 Russia (RBU)), with YCSs being in the range 3 – 5, 1 – 3, 1 - 2 and 1 - 2 for pests and
24 diseases, nutrients, heat and aridity, respectively (Table 2).

1

2 **Table 2:** Production and regional mean Yield Constraint Score (YCS, rounded to
3 nearest integer) for the five highest producing regions for soybean, wheat,
4 rice and maize.

Soybean	Production (Tg)	Ozone	Pests & diseases	Nutrients	Heat	Aridity
Global	253.6	3	4	2	2	2
South America	125.8	2	4	3	2	1
N America	90.3	3	3	2	2	1
East Asia	14.9	3	4	2	1	2
South Asia	13.2	3	5	1	2	2
Russia	3.6	2	4	1	1	2
Wheat	Production (Tg)	Ozone	Pests & diseases	Nutrients	Heat	Aridity
Global	673.3	2	4	2	3	2
Europe	163.8	2	3	2	2	1
East Asia	118.9	2	3	2	2	2
South Asia	118.1	2	4	2	4	2
N America	84.2	2	4	2	2	2
Russia	66.1	2	4	2	2	1
Rice	Production (Tg)	Ozone	Pests & diseases	Nutrients	Heat	Aridity
Global	716.8	1	4	2	1	2
East Asia	221.2	2	4	2	1	2
South Asia	219.2	2	5	2	2	2
South East Asia	204.5	1	4	3	1	1
South America	22.7	1	4	3	1	1
Sub Saharan Africa	21.1	1	5	2	1	2
Maize	Production (Tg)	Ozone	Pests & diseases	Nutrients	Heat	Aridity
Global	869.1	2	4	2	2	2
N America	313.1	2	3	2	2	2
East Asia	194.4	2	4	2	2	2
South America	91.7	2	4	3	2	2
Europe	75.6	2	3	1	2	1
Sub Saharan Africa	60.0	1	5	2	2	2

5

6 Wheat

By far the highest production losses due to ozone per grid square for wheat are predicted for India and China, with large areas in N India and NW China having over 15% yield losses amounting to production losses in excess of 0.1 Tg (Fig. 3, Table S5). Production losses are also predicted to be high in the highest wheat producing areas of Europe (including France and Germany) and central states of the USA. The mean YCS for ozone globally was 2, reflecting the same mean score in the 9 highest wheat producing regions (Tables 2 and S5), and matching that globally for nutrients, and aridity. The highest predicted production losses due to ozone only overlapped with areas of the highest YCS_{all} in NW India, Pakistan and S USA (Fig. 3). Scores of 3 and above coincide for ozone, pests and diseases, heat and nutrients in a wider area including parts of EAS, SAS and NAM, whilst YCSs for aridity are lower in several parts of this region than for ozone. This study also indicated that the highest YCS values for all stresses are for heat (score 5) in areas of Northern Africa (NAF), SAS, SAM, SSA and SAM (particularly Argentina). Scores for pests and diseases are 3 or more across most of the wheat growing areas. YCSs for nutrient availability are generally the lowest of the five stresses, although there are some high risk areas with values of 4 and above in, for example, NW SAS, Central Asia (CAS) and NE EUR (e.g. Finland), E NAM and SAM (e.g. Brazil). The highest YCSs for aridity are in a zone that includes parts of NAF, the Middle East (MDE) and SAS, CAS and EAS. For the five main wheat producing regions, the mean YCS for ozone was 2 representing 5 - 10% yield loss, whilst it was 3 - 4 for pests and diseases and mainly 1-2 for the other three stresses (Table 2).

Rice

Production losses due to ozone are predicted to only be in excess of 0.1 Tg per grid square in parts of India, Bangladesh, China and Indonesia (Fig. 4). In these areas, the percentage yield losses are mainly in the range 7.5 – 12.5%, resulting in grid square ozone YCSs that are usually either 2 or 3. Across all of the rice growing regions, the mean YCS for ozone per

1 region is either 1 or 2, resulting in a global mean score of 1 and a score of 2 in the two
 2 highest producing regions (Tables 2 and S6). YCSs of 3 for ozone occurred in areas of India
 3 and China where the YCS_{all} was usually in the highest range for rice of 13 – 15. Overall, the
 4 highest mean YCS for this crop is for pests and diseases, being score 5 in most of the rice
 5 growing areas of SSA and SAS. Nutrient YCSs are 4 or more in many parts of SSA, SAM
 6 (particularly Brazil), EAS and SAS. Heat stress is predicted to be less of a problem for rice
 7 than for wheat, with few regions having a score of 3 or more. Indeed, the regional mean YCS
 8 for heat stress in rice is mostly either 1 or 2 (Tables 2 and S6). In the three highest producing
 9 rice regions, EAS, SAS and SEA, irrigation usage is 96%, 74% and 28%, respectively (Table
 10 S5). Here, the aridity score is predicted to be 3 or more only in areas of NW China and W
 11 India.

12 Maize


13 China and the USA are the two countries predicted to have the largest areas where production
 14 losses due to ozone for maize that exceed 0.1 Tg per $1^{\circ} \times 1^{\circ}$ grid square (Fig. 5). In these
 15 areas, the percentage yield losses are mainly in the range 7.5 – 15% for the USA and 12.5 –
 16 15 % for China. There are also high-risk areas in S EUR, for example, in parts of S France
 17 and N Italy and in NAF (particularly Egypt). These areas generally have a YCS_{all} for maize
 18 in the range 10 – 15, and are not in the areas with the highest YCS_{all} for maize of >15. The
 19 latter are mainly found in parts of SAS and SSA, with occasional small areas elsewhere. The
 20 mean YCS for ozone for the four highest maize producing regions (NAM, EAS, SAM and
 21 EUR) is 2 (Tables 2 and S7). For the stresses other than ozone, the highest scores for YCS
 22 are for pests and diseases, with scores of 5 predicted in most of SSA, SAS and SEA. YCSs of
 23 4 and above are predicted for nutrients in much of SSA (particularly in western countries),
 24 SEA, large areas of SAM (particularly Brazil), parts of E USA and small areas of Europe.
 25 For maize, the YCSs for aridity are highest in eastern NAM and SAM, parts of NAF

(particularly W Egypt), MDE, SAS (particularly NW India) and EAS (particularly NE China). Heat stress YCSs are lower for maize than for wheat, indicating that the main areas of concern for this crop are in W SSA, W MDE and SAS. Globally, the mean YCS for maize is 4 for pests and diseases and 2 for each of the other four stresses (Table 2).

3.1.3 Case study - India

It is clear from the results presented above that the five environmental stresses included in this study are all predicted to be having relatively high impacts on yield in several states of India. We selected this country for a more in depth analysis. Although the spatial data for India is present on the global maps in Figs 2 - 5, for ease of interpretation, we have produced additional maps for India for wheat and rice, the two most important crops by production in Fig. S5 and S6, respectively. At the national scale, the mean YCSs for the crop with the highest total production in India, wheat, are 3, 4, 2, 4 and 2 for ozone, pests and diseases, nutrients, heat and aridity, respectively (Table S5). For rice, the second most important crop by Tg produced in India, the YCSs for the same five stresses respectively are 2, 5, 1, 2 and 2 (Table S6). As the data for the risk of losses due to pests and diseases was only available at the national scale for India, with YCSs of 4 for wheat and 5 for rice, these effects were not included in this spatial analysis, conducted at the $1 \times 1^\circ$ scale.

For wheat, the highest production is in the adjacent N states of Uttar Pradesh, Madhya Pradesh, Haryana, Rajasthan and Punjab (Fig. 6). Together, these five states account for 85% of Indian wheat production. Predicted percentage yield losses due to ozone are in the range 15 – 20 % (mean of 16.4%) in most of the wheat producing areas of Uttar Pradesh, the state with the highest wheat production, resulting in a mean ozone YCS of 3 (Fig. 6 and S5). The mean YCS for ozone was 3 for Haryana, Rajasthan and Punjab and 2 for Madhya Pradesh where ozone uptake is lower (Fig. S5). Although the highest percentage yield losses due to



1

2 **Fig. 2: The global effects of five biotic and abiotic stresses on soybean.** All data are
3 presented for the 1 x 1° grid squares where the mean production of soybean was > 500 tonnes
4 (0.0005 Tg). (a) Presents the effects of ozone on crop production (thousand tonnes or 0.001
5 Tg per grid square), and (b) the percentage yield loss due to ozone, averaged for the period
6 2010-2012. In (d) to (h), the Yield Constraint Score (YCS) is presented per grid square on a
7 scale of 1 to 5, where 5 is the highest level of stress (see Table 1) for ozone, pests and
8 diseases, nutrients, heat stress and aridity, respectively, whilst (c) is the total YCS (YCS_{all})
9 calculated from the sum of each of these per grid square. The regional impacts are
10 summarised in Table 2 for the five highest producing regions and Table S4 provides all
11 country and regional means.





Fig. 3: The global effects of five biotic and abiotic stresses on wheat. All data are presented for the $1 \times 1^\circ$ grid squares where the mean production of wheat was > 500 tonnes (0.0005 Tg). (a) Presents the effects of ozone on crop production (thousand tonnes or 0.001 Tg per grid square), and (b) the percentage yield loss due to ozone, averaged for the period 2010-2012. In (d) to (h), the Yield Constraint Score (YCS) is presented per grid square on a scale of 1 to 5, where 5 is the highest level of stress (see Table 1) for ozone, pests and diseases, nutrients, heat stress and aridity, respectively, whilst (c) is the total YCS (YCS_{all}) calculated from the sum of each of these per grid square. The regional impacts are summarised in Table 2 for the five highest producing regions and Table S5 provides all country and regional means.



1
2
3
4
5
6
7
8
9
10
11

Fig. 4: The global effects of five biotic and abiotic stresses on rice. All data are presented for the 1 x 1° grid squares where the mean production of rice was > 500 tonnes (0.0005 Tg). (a) Presents the effects of ozone on crop production (thousand tonnes or 0.001 Tg per grid square), and (b) the percentage yield loss due to ozone, averaged for the period 2010-2012. In (d) to (h), the Yield Constraint Score (YCS) is presented per grid square on a scale of 1 to 5, where 5 is the highest level of stress (see Table 1) for ozone, pests and diseases, nutrients, heat stress and aridity, respectively, whilst (c) is the total YCS (YCS_{all}) calculated from the sum of each of these per grid square. The regional impacts are summarised in Table 2 for the five highest producing regions and Table S6 provides all country and regional means.




Fig. 5: The global effects of five biotic and abiotic stresses on maize. All data are presented for the $1 \times 1^\circ$ grid squares where the mean production of maize was > 500 tonnes (0.0005 Tg). (a) Presents the effects of ozone on crop production (thousand tonnes or 0.001 Tg per grid square), and (b) the percentage yield loss due to ozone, averaged for the period 2010-2012. In (d) to (h), the Yield Constraint Score (YCS) is presented per grid square on a scale of 1 to 5, where 5 is the highest level of stress (see Table 1) for ozone, pests and diseases, nutrients, heat stress and aridity, respectively, whilst (c) is the total YCS (YCS_{all}) calculated from the sum of each of these per grid square. The regional impacts are summarised in Table 2 for the five highest producing regions and Table S7 provides all country and regional means.


1
2 ozone were predicted for states in the far NE of India such as Assam and Manipur, total
3 production losses there were predicted to be minimal as this is not an important wheat
4 growing area. The area of highest ozone impacts on wheat production coincided with the
5 area with the highest YCS for heat stress which covered most of the northern half of the
6 country. Aridity and nutrient YCSs were highest to the W of this region, coinciding with
7 percentage yield losses for ozone predicted to be in the range 5-15% (YCS of 2- 3). For the
8 five highest wheat producing states, the mean YCS for heat stress was 5, with scores for
9 aridity being 2 or 3, and nutrients being 1 – 3 (Fig. 6).

10 Rice growth is much more widely distributed in India than wheat growth, with the highest
11 production being in the N, in part coinciding with wheat growing areas in states such as Uttar
12 Pradesh, and also in E and S states such as West Bengal, West Odisha, Andhra Pradesh and
13 Tamil Nadu (Fig. 6). Together these 5 states produce just below half of India's rice
14 production. The highest percentage yield losses for ozone are predicted to be in the range 10
15 – 15% in the N of the country, including in Uttar Pradesh (mean ozone YCS of 3, Fig. 6 and
16 S6). Lower effects were predicted for Odisha and West Bengal (mean YCS of 2) and the
17 least ozone effects were predicted for rice producing areas in the southern states of Tamil
18 Nadu and Andhra Pradesh (mean YCS of 1), where percentage losses were frequently less
19 than 5%. Heat stress is less of a concern for rice, with a mean YCS of 2 predicted for each of
20 the 5 most important rice producing states. Nutrient stress is predicted to only be important in
21 the far NE states and in isolated grid squares in Rajasthan and along the W coast of India.
22 The mean YCSs for nutrients and aridity for the five highest producing states are either 1 or 2
23 (Fig. 6).

24 **3.2. Plant traits associated with tolerance of ozone and associated stresses in crops**

The derivation of dose-response relationships for 52, 18 and 44 genotypes of soybean, wheat and rice respectively (Fig. 1) has shown that there is clearly scope for the breeding of ozone

(a)



(b)




Fig. 6: Yield Constraint Score (YCS) for five constraints on the yield of (a) wheat and (b) rice in the five Indian states with the highest production per crop. The bars represent the mean YCS per 1 x 1° grid square per state on a scale of 1 to 5, where 5 is the highest level of stress (see Table 1), rounded to the nearest integer. Note: The YCS for pests and disease is only available at the National Scale for India (score 4 for wheat and 5 for rice) and is presented here for information.

tolerant varieties, as many varieties had responses that are above the regression line. As part of this study, we identified a number of traits that could contribute to improved ozone

1 tolerance and have summarized these in an ozone-tolerant crop ideotype, including potential
2 trade-offs and synergies for effects of other stresses that can co-occur with ozone (Fig. 7).

3 Leaf traits for ozone tolerance fall into two categories, the first being processes that limit
4 ozone entry. These include stomatal conductance, and the related trait of water use efficiency
5 (WUE), that reduce ozone uptake while maintaining high rates of photosynthesis. These traits
6 are associated with reduced leaf transpiration and whilst they would be beneficial for water
7 conservation under drought conditions, they may reduce yield and could be potentially
8 deleterious under heat stress by limiting evaporative cooling (Reynolds *et al.*, 2007).

9 Similarly, reduced water uptake associated with lower stomatal conductance has the potential
10 to limit uptake of nutrients such as N from the soil (Zhou *et al.*, 2016). While pathogens are
11 known to have negative effects on leaf gas exchange (Debona *et al.*, 2014), the impact of
12 inherently lower stomatal conductance on disease establishment is less clear although it could
13 be expected that ingress of leaf pathogens that access leaves through the stomatal pores
14 would be reduced.

15 A second category of favourable leaf traits includes antioxidant metabolism and pathways
16 involved in programmed cell death (PCD). Ozone is decomposed into reactive oxygen
17 species (ROS) in the plant apoplast, which either cause direct oxidative damage, or induce
18 signalling cascades similar to a pathogen response, ultimately leading to PCD (Kangasjärvi *et*
19 *al.*, 2005). Thus, balancing the interplay of redox homeostasis and PCD pathways is essential
20 for the breeding of ozone tolerant crop plants. As a first line of defence against ozone stress,
21 high levels of apoplastic antioxidants such as ascorbate may mitigate ROS formation, a
22 concept that has been confirmed in crop plants such as wheat (Feng *et al.*, 2010) and legumes
23 (Yendrek *et al.*, 2015). Breeding for high levels of antioxidants is also assumed to cause




Fig. 7: An ideotype for an ozone-tolerant crop. ‘+’ indicates where there would be a benefit for other stresses of improving tolerance to ozone for the trait, whilst ‘-’ indicates a trade-off, and ‘0’ is no effect.

synergies with other types of abiotic stress tolerance, including for drought and heat, both of which are associated with oxidative stress (Gill & Tuteja, 2010). In the case of some nutrient disorders and biotic stresses, functional redox balance rather than high antioxidant levels *per*

1 *se* are considered as important (Munné-Bosch *et al.*, 2013; Suzuki *et al.*, 2012; Wu *et al.*,
2 2017).

3 PCD is an important pathway of pathogen response in plant leaves (Huysmans *et al.*, 2017),
4 which is controlled by the interplay of ROS, signalling cascades and plant hormones
5 (Kangasjärvi *et al.*, 2005). Breeding for ozone tolerance could thus keep plants from inducing
6 PCD despite the presence of apoplastic ROS. This idea is supported by a study in rice, in
7 which the disruption of the pathogen and ozone responsive apoplastic protein *OsORAP1*,
8 which is involved in cell death, led to enhanced ozone tolerance (Ueda *et al.*, 2015). The
9 potential interference of this strategy with pathogen tolerance in crops is obvious, but it is
10 currently unclear whether a synergistic or rather antagonistic relationship would occur with
11 different classes of pathogens i.e. biotrophic versus necrotrophic ones (Huymans *et al.*,
12 2017). Implications of PCD in other stress types such as heat (Locato *et al.*, 2008), drought
13 (van Doorn 2011) and nutrient deficiency (Siyiannis *et al.*, 2012) have also been reported but
14 the implications for ozone tolerance breeding remain unclear.

15 Root traits that support ozone stress tolerance would include the capacity to efficiently
16 acquire water and nutrient resources under stress environments (Resource Acquisition
17 Efficiency in Fig. 7). Ozone is known to have a greater negative impact on roots than shoots,
18 resulting in the decline in the root/shoot ratio commonly observed (Fiscus *et al.*, 2005). There
19 is evidence that ozone may have an even greater impact on fine roots that acquire water and
20 nutrients from the soil (Vollsnes *et al.*, 2010). Fine roots are the new frontier of future root
21 research. The framework for describing fine root architecture is being refined (McCormack *et*
22 *al.*, 2015; Zobel, 2016), and new techniques are now available to assess fine root dynamics
23 (e.g. measurements of root diameter, Zobel *et al.*, 2007). The challenge ahead is to define and
24 measure fine root traits that contribute to ozone tolerance, and then determine how these traits

1 affect plant response to other stress factors. Presumably, traits that contribute to robust root
2 systems will be of benefit across a range of abiotic stresses.

3 Traits associated with reproductive organs such as panicles or pods are of primary importance
4 in breeding, although the effects of ozone on these organs may be rather secondary, i.e.
5 caused by foliar responses that limit assimilate acquisition (described above) or effects on
6 flowering and pollen viability (Black *et al.*, 2000). Yield losses due to ozone have been
7 ascribed to various yield components in different crops, including reductions in individual
8 seed weight, reduced spikelet number, enhanced spikelet fertility, and reduced panicle or pod
9 number (Ainsworth, 2008; Feng *et al.*, 2008; Morgan *et al.*, 2003), with associated reductions
10 in harvest index (e.g. for wheat, Pleijel *et al.*, 2014). Maintaining high values in these harvest
11 fractions despite ozone stress forms an important breeding target, but synergies or trade-offs
12 with other types of stress would be complex and little information is available to date.

13 Maintaining high crop quality despite ozone stress represents another important breeding
14 goal. Ozone can affect multiple quality traits in seed crops, including protein and starch
15 concentration, as well as visual appearance (Broberg *et al.*, 2015; Wang & Frei, 2011). In
16 many cases, increases in seed protein concentration despite losses in protein yield are
17 observed. This apparent beneficial effect is offset by the negative effects of ozone on seed
18 weight (e.g. for wheat, Broberg *et al.*, 2015). Another quality trait that has been affected in
19 rice by ozone is grain chalkiness, i.e. the formation of milky patches on grains due to
20 inhibited starch loading (Jing *et al.*, 2016). Chalkiness was first described as a typical
21 symptom of heat and drought stress (Wassmann *et al.*, 2009), and lowering plant
22 susceptibility to chalkiness *via* breeding may thus have potential co-benefits with regards to
23 these stresses.

1 A further category of traits that could be targeted by breeders are phenological characteristics.
2 Plants that have a shorter maturity period by entering earlier into reproductive phases might
3 be more tolerant, as they would receive a lower cumulative ozone dose, and might avoid high
4 ozone episodes occurring late in the cropping season. This principle was confirmed in a study
5 by Ueda *et al.* (2015), in which more than 300 genotypes of rice were screened for ozone
6 response, and yield losses were positively correlated with the number of days to maturity. In
7 general, breeding fast-maturing crop varieties may produce substantial synergies, reducing
8 the impacts of growing seasons characterized by high incidence of other stresses, such as
9 drought, heat, nutrient, or biotic stresses.

10 **4. Discussion**

11 In bringing together these datasets and modelling methods to derive YCSs for five stresses
12 and four key crops, we have conducted the first global assessment of the magnitude of ozone
13 stress in relation to other stresses for four staple crops. We have also derived an ideotype for
14 an ozone- and multi-stress tolerant crop. We provide an extended discussion here that first
15 considers the results presented and then considers potential solutions for increasing crop
16 tolerance of ozone, including crop management and breeding approaches.

17 **4.1 The global scale of ozone impacts on crops relative to impacts of other stresses**

18 An in depth evaluation of the spatial analysis conducted here is presented in the Supporting
19 Information (T1) and summarised here. The benefits of impacts modelling based on the
20 stomatal uptake of ozone rather than the concentration above the leaf, together with an
21 evaluation of global modelling of POD₃IAM, are discussed by Mills *et al.* (2018a). In the
22 absence of suitable stomatal uptake dose response relationships for soybean, rice and maize,
23 the RS_w method was developed whereby the effects of ozone on these crops was determined
24 from the POD₃IAM response of wheat. We had to assume that the differences in ozone

1 concentration and sensitivity were a greater driver of response than differences in stomatal
2 uptake and are unable to quantify the uncertainty introduced by this assumption. Whilst
3 experimental data in the M7 response functions used in the RS_w method represented the
4 major crop growing regions for soybean, wheat and rice, the function for maize was limited
5 to relatively old data from NAM only. Thus, the data analysis presented here for maize is
6 likely to be the most relevant for effects described in NAM, and is less certain when applied
7 to other maize-growing regions of the world. For each crop, we assumed only one crop
8 growth period per year. Thus, for those crops such as rice where two or three crop growth
9 cycles may occur per year in major growing areas, assessments based on the main growth
10 period will have an added level of uncertainty. The abiotic and biotic stresses included here
11 were selected as examples for comparison with ozone effects, with heat stress being chosen
12 as representative of effects of extreme climatic events associated with climate change. We
13 acknowledge that other stresses such as flooding may also have catastrophic local effects on
14 yield (e.g. in China, Tao *et al.*, 2017), but have focussed on example stresses for which global
15 data is readily available. Furthermore, global warming impacts on yield could be in a similar
16 range to ozone (e.g. Challinor *et al.*, 2009; Lobell & Asseng, 2017) but have not been
17 considered here. Scores for YCS for pests and diseases may have overestimated current
18 losses as advances in pesticide usage since the 2002-04 dataset was compiled may have
19 reduced total impacts. As it was not possible to base all YCSs on percentage yield loss,
20 uncertainty will have been introduced by comparing effects across stresses. We have
21 acknowledged this uncertainty by using the YCSs to indicate the location of the largest
22 effects rather than to quantify the extent of effects. Lastly, YCS_{all} simply summed all YCSs
23 and provides an indication of where multiple stresses co-occur, without taking into account
24 any interactions that may occur that might lessen or increase the combined effects on yield.
25 Taking into account all of these caveats, this study is the first to present ozone impacts on the

1 global scale together with impacts of other biotic and abiotic stresses, and show spatially
2 where such stresses are likely to co-occur for four major staple crops.

3 At the national scale, the countries identified as having the largest potential effects of ozone
4 (e.g. USA, India and China) match those with the highest monitored ozone concentrations
5 (Mills *et al.*, 2018b) as well as those predicted using concentration-based approaches to have
6 the highest potential yield losses (Avnery *et al.*, 2011a,b; Van Dingenen *et al.*, 2009). At the
7 sub-national scale, however, there were some differences in areas predicted to be at risk,
8 where our stomatal uptake modelling method took into account the modifying effects of
9 climate and soil moisture on ozone uptake rather than simply predicting the largest effects in
10 the areas with the highest ozone concentrations. For example, in India, this study predicts the
11 largest effects on wheat and rice in the northern areas, south of the Himalayas where ozone
12 levels, climatic conditions and irrigation usage promote ozone uptake and subsequent effect.
13 In contrast, an earlier concentration-based study provided little spatial differentiation in
14 effects, predicting widespread and similar effects of ozone in the northern half of India for
15 wheat and across most of India for rice (Van Dingenen *et al.*, 2009). On a global scale, we
16 predict that ozone (mean of 2010 - 2012) reduces soybean yield by 12.4%, wheat yield by
17 7.1%, rice yield by 4.4% and maize yield by 6.1%, adding up to a total of 227 Tg of lost
18 yield. These mean percentage losses are different to those predicted by Avnery *et al.* (2011a)
19 and Van Dingenen *et al.* (2009) using concentration based metrics. Their studies predicted
20 higher losses for wheat (15.4% and 12.3%, respectively) and lower losses for soybean (8.5%
21 and 5.4%, respectively) using AOT40 (Accumulated hourly mean ozone above 40 ppb during
22 daylight hours) for the year 2000.

23 Multivariate analysis of trends in soybean and maize yields in the USA that included a
24 concentration-based ozone metric indicated that the ozone effect is dependent upon
25 temperature and water availability (McGrath *et al.*, 2015). This fits with our earlier

1 conclusion that stomatal uptake-based risk assessment provides a better indication of ozone
2 effects on yield than concentration-based assessments (Mills *et al.*, 2018a). The McGrath *et*
3 *al.* (2015) study indicated a greater sensitivity of maize to ozone than soybean, with maize
4 and soybean yield losses due to ozone over a 31-year period averaging 10% and 5%,
5 respectively. It is possible that our analysis under-estimated the effects of ozone on maize as
6 our analysis was based on experimental data from 1981, 1985, 1991 and 1992. Since newer
7 varieties of wheat and soybean are more sensitive to ozone than older varieties (Biswas *et al.*,
8 2009; Osborne *et al.*, 2016), then newer maize varieties may also be more ozone sensitive
9 leading to larger effects. Partial derivative-linear regression analysis of heat and
10 concentration-based ozone stress impacts on yield data in the USA and Europe indicated
11 similar areas at risk from ozone for maize and soybean to our study, but fewer areas at risk
12 for wheat (Tai & Val Martin, 2017). The latter may reflect that their study omitted soil
13 moisture as a confounding factor, which our earlier modelling study indicated is a particularly
14 important factor in modifying ozone uptake (Mills *et al.*, 2018a). These two statistical
15 studies have confirmed that factors other than ozone concentration need to be taken into
16 account in analysing ozone effects on yield and have drawn attention to potential co-
17 occurrence of heat and ozone stress effects on crop yield.

18 It is clear from our analysis that yield effects due to ozone are within the range of concern for
19 other biotic and abiotic stresses. For example, extreme heat was estimated to have reduced
20 national cereal production by 9-10% (1964 - 2007), with later droughts reducing yields by
21 more than earlier droughts (13.7% for 1985 – 2007 compared to 6.7% for 1964 – 1984, Lesk
22 *et al.*, 2016). If we applied the percentage yield loss ranges used for ozone in YCS (Table 1)
23 to these drought-induced yield losses, the YCS would be 2 or 3 depending on the time period
24 used in the analysis. The YCSs for ozone are mainly in the same range as those predicted for
25 wheat, maize and rice for global impacts of heat stress and aridity (scores 2-3, with rice-heat

1 having a YCS of 1). Across all four crops, the areas predicted to be at the greatest risk of
2 ozone effects on yield are predicted to be: SE NAM, S EUR, N SAS and E EAS, with parts of
3 SAM also predicted to be at risk of yield loss for soybean. In some of these areas, ozone
4 effects are predicted in areas also at risk from heat stress and to a lesser extent aridity, whilst
5 co-occurrence with nutrient stress depended on the crop and tended to be most common in
6 parts of EAS and SAM. Potential impacts on yield due to pests and diseases were predicted
7 to be relatively high in many areas of the world, particularly in those at risk from ozone
8 impacts in SAS and EAS.

9 Ozone impacts are predicted in areas where the largest gaps occur between actual and
10 estimated potential yield, such as parts of SAM, SSA, EAS, SEA and SAS (Neumann *et al.*,
11 2010). Here, yield gaps are already known to be widened by limitations in nutrient and/or
12 irrigation availability (Mueller *et al.*, 2012) and may be further widened by negative effects
13 of ozone pollution. Indeed, in the same regions there has been a plateauing or decrease in the
14 rate of yield increase in recent decades (Grassini *et al.*, 2013; Ray *et al.*, 2012). We suggest
15 that ozone pollution could be contributing to this stagnation, and suggest below how crop
16 tolerance of the pollutant could be improved by breeding or management (Section 4.2.3).

17 India was selected as a case study, as our analysis indicated that ozone pollution may be a
18 particular problem in this country, adding to the existing multi-stress constraints on crop yield
19 (Jaswal, 2014). National mean yield losses due to ozone were predicted to be 15.8%
20 (soybean), 12.6% (wheat), 6.2% (rice) and 7.5% (maize) amounting to 12.6 Tg of lost yield.
21 For wheat, our predicted mean yield loss in Uttar Pradesh of 16% was comparable to a mean
22 17% yield benefit from reducing the ambient ozone from 46 to 5 ppb (M7) by air filtration in
23 field studies conducted from 2004 to 2008 at Varanasi in Uttar Pradesh (Rai *et al.*, 2007,
24 Sarkar *et al.*, 2010). Similarly, at a field site in Haryana, reduction in the M7 by filtration

1 from 37 ppb to 6 ppb, resulted in a 16% yield benefit for wheat (Bhatia *et al.*, 2011), which
2 was similar to our state mean of a 15% yield reduction due to ozone.

3 Wheat yield losses were predicted to be highest in this study in the same regions of India as
4 those predicted by Tang *et al.* (2013) in the first stomatal uptake-based risk assessment for
5 the country. Our analysis, taking into account the added effects of soil moisture and irrigation
6 usage, extended the region of highest ozone effects across the Indo-Gangetic Plain and
7 including Uttar Pradesh and Bihar. Together with Haryana and Punjab, these states are
8 considered to have the highest reductions in yield due to the combined effects of climate
9 change and air pollution, with reductions as high as 50% being predicted in one
10 concentration-based study (Burney & Ramanathan, 2013). Our multi-stress analysis
11 confirmed that heat stress is particularly important in this region (Lobell *et al.*, 2012). Site-
12 specific analysis of the effects of future increases in temperature in 2030-2040, indicated that
13 heat stress is likely to continue to reduce yields in the Indo-Gangetic Plain, especially under
14 climate change (Asseng *et al.*, 2017). Given that from a food security perspective, it is
15 crucial to reduce yield gaps in India, reducing ozone pollution and/or its effects could
16 potentially provide beneficial additional yield in future climates.

17 In considering these comparisons, we are aware that in reality ozone will interact with the
18 other stresses considered and integrated responses in growth and yield will occur. These
19 interactions are generally thought to be determined by factors that might affect gas exchange
20 or metabolic responses to stress. For example, limited water stress may reduce ozone uptake
21 but as water stress becomes more severe, any protection afforded by reduced ozone uptake
22 may be outweighed by drought-induced yield reductions. Additionally, these stresses are
23 thought to impart similar defence mechanisms (Huymans *et al.*, 2017; Kangasjärvi *et al.*,
24 2005; Locato *et al.*, 2008). Whether multiple stresses induce additive or synergistic metabolic
25 responses is open to question and we do not yet have the understanding or tools to be able to

quantify these interactions. Nevertheless, through providing a first global assessment of where these stresses co-occur we have identified which stresses are most important across different global regions. This will help other researchers to identify threats and target future research needs to improve our understanding of responses to multiple stress conditions.

4.2 Options for reducing ozone impacts on crops

The analysis presented here has clearly shown that ozone impacts on yield are occurring in many areas of the world for four staple crops, and that in some regions the YCSs for ozone are as high or higher than for other biotic and abiotic stresses. Whilst these results highlight the ozone problem, we offer here some possible options for reducing ozone effects on crops that might help in closing the ozone yield gap.

4.2.1 Global effort to reduce ozone precursor emissions

The most obvious way of closing the ozone yield gap for crops is to substantially lower the anthropogenic emissions that lead to ozone pollution. Since ozone is a transboundary air pollutant – impacts of emissions in one country can impact on crops grown in countries many 100s and even 1000s of km away – efforts to reduce ozone need to be taken at both local and global scales. One study, using ozone concentration-based metrics, indicated that 100% reductions in anthropogenic precursor emissions from NAM would reduce global yield losses due to ozone for the four crops in our study by between ca. 5% (rice) to ca. 80% (soybean), whilst a complete cut in precursor emissions from SEA would reduce global yield losses by between ca. 20% (soybean) and ca. 95% (rice) (Holloway *et al.*, 2012). Whilst such dramatic cuts in ozone precursor emissions are highly unlikely for the foreseeable future, progress has been made in EUR and NAM, with emission cuts of ca. 40% for major ozone precursors such as NO_x, VOC and CO being made between 1990 and 2013 (Maas & Grennfelt, 2016). These cuts have been associated with significant decreasing trends in the concentration-based metric

1 AOT40 at 26% and 11% of monitoring sites in wheat growing areas of NAM and EUR,
2 respectively over the period 1995-2014 (Mills *et al.*, 2018b), although the dominant trend for
3 EUR remains “no change”. Over the same time period, increases in precursor emissions of
4 20-30% in other areas of the world, including by 50% in India and China have led to
5 increases in ozone concentration in these regions (Maas & Grennfelt, 2016). For example,
6 there has been a significant increase in ozone concentration at nearly 50% of wheat growing
7 monitoring sites in EAS, with average annual increases in AOT40 at these sites being in the
8 range 300–700 ppb h y⁻¹ over the period 1995-2014 (Mills *et al.*, 2018b).

9 Our modelling results suggest that even with declining emissions in NAM, current yield
10 losses due to ozone are in the range 5.3% (rice) to 15.5% (soybean), whilst for EAS with
11 rising emissions, current yield losses are in the range 7.9% (rice) to 19.1% (soybean). With
12 ozone concentrations predicted to continue to rise in EAS and SEA for at least the next 2-3
13 decades even with the most optimistic scenarios (Wild *et al.*, 2012), and as these two regions
14 are predicted to produce 80% of all global ozone precursor emissions by 2050 (Maas &
15 Grennfelt, 2016), there would be considerable benefit for crop yield in the implementation of
16 a concerted effort to reduce precursor emissions in these rapidly developing regions. Actions
17 to reduce ozone are already being considered in some countries. For example, in China, three
18 approaches are being introduced to reduce ozone concentrations: enforcing the European
19 standard V for diesel vehicle emissions; encouraging widespread use of electric vehicles; and
20 discouraging private car use by improving public transport (Feng *et al.*, 2015). Continued
21 effort to reduce ozone is also needed in developed regions such as NAM and EUR as models
22 predict that whilst efforts to reduce peak concentrations have been partially successful in
23 reducing ozone concentrations in recent decades, a stabilisation in ozone concentrations in
24 the next decade or two is likely to be followed by further rises in global background ozone

1 concentration by 2050, primarily driven by increasing CH₄ emissions (Maas & Grennfelt,
2 2016).

3 Whilst reducing global ambient ozone concentrations remains a crucial long-term goal for
4 reducing the ozone yield gap, approaches described below based on crop management and
5 breeding are more likely to provide shorter-term solutions, with some having potential for
6 implementation in the near future.

7 4.2.2 Exploiting existing varietal differences in ozone sensitivity

8 Whilst the analysis presented here has confirmed that intraspecific variation in ozone
9 sensitivity is clearly present for wheat, rice and soybean in experiments conducted over the
10 last 30 - 40 years (Fig. 1, Table S3), of larger importance in the context of closing the ozone
11 yield gap is the potential for selecting ozone tolerance amongst currently grown varieties. To
12 assess this, ideally, varieties should be exposed to ozone under the same environmental
13 conditions, allowing for realistic comparisons of effects on yield and assessments of variety
14 by ozone interactions. Unfortunately, relatively few such experiments have been conducted
15 with two or more varieties in the last decade. Those recent studies showing significant
16 variety by ozone interactions, indicating scope for selecting the more ozone tolerant variety,
17 include examples from SAS and EAS for rice (Akthar *et al.*, 2010; Shi *et al.*, 2009) and
18 wheat (Feng *et al.*, 2010, 2016; Singh *et al.*, 2017; Zhu *et al.*, 2011;), NAM for soybean
19 (Betzberger *et al.*, 2010; Jiang *et al.*, 2018) and maize (Yendrek *et al.*, 2017); and EUR for
20 wheat (Harmens *et al.*, 2018). Further support for the potential benefits of selecting tolerant
21 varieties is also provided by comparisons of yield in filtered air versus non-filtered air
22 (Osborne *et al.*, 2016; Pleijel *et al.*, 2018). For example, in recent studies, reductions of
23 ambient ozone concentration by filtration significantly increased the yield of ozone-sensitive
24 soybean cultivars (PUSA 9712, PUSA 9814) by over 40% (Singh & Agrawal, 2011), rice

1 cultivar Kirara 397 by over 20% (Frei *et al.*, 2012) and wheat cv PBW 343 by 18-20%
2 (Tomer *et al.*, 2015).

3 A modelling study has been conducted to highlight the potential for avoiding production loss
4 in global wheat, maize and soybean by selecting crop varieties with lower than average
5 sensitivity to ozone (Avnery *et al.*, 2013). The variation in sensitivity among varieties was
6 based on the experimental evidence from the large-scale US National Crop Loss Assessment
7 Network (NCLAN) field studies conducted mainly during the 1980s (Heagle, 1989; Heck *et*
8 *al.*, 2013; Heck, 1989). Using a concentration-based method, the study showed that choosing
9 crop varieties with ozone tolerance could improve global crop production by over 140 Tg in
10 2030, equivalent to a 12% increase. Although the older North American varieties may not
11 represent current global variation, and some of the 1980s varieties are no longer used, the
12 approach of Avnery *et al.* (2013) could be extended by conducting new screening
13 experiments with a regional focus to inform farmer choice, modelling and breeding
14 programmes and by using a stomatal uptake based modelling approach.

15 4.2.3 Breeding new varieties with multiple stress tolerance, including ozone

16 The heterogeneity in variety response to ozone for soybean, wheat and rice (Fig. 1) has
17 clearly shown the scope for breeding ozone-tolerant varieties, and an ideotype for an ozone
18 tolerant crop has been defined here (Fig. 7). Ideally, the improved ozone response must not
19 compromise the yield potential or other required agronomic characteristics (e.g. resistance to
20 diseases, shattering, and lodging). Since this study has also shown that ozone stress can
21 typically co-occur with stress caused by heat, pests and diseases, and to a lesser extent aridity
22 and nutrients, the breeding for ozone tolerance traits may cause potential synergies or trade-
23 offs that also need to be considered (Fig. 7). Candidate traits for ozone tolerance were
24 described in Section 3.2.

1 Traditional breeding approaches such as pedigree selection require extensive screening of a
2 large number of plants in multiple locations over extended periods of time (Frei, 2015).
3 Whilst feasible, experimentally maintaining designated ozone concentrations on a sufficiently
4 large scale required for breeding (e.g. in large scale FACE (free air concentration exposure)
5 experiments) seems economically unviable. Therefore, molecular breeding approaches such
6 as marker assisted selection (MAS) appear to be more promising. Phenotypic variation in
7 traits associated with ozone tolerance can be evaluated in smaller-scale controlled ozone
8 fumigation experiments and linked to genetic markers using mapping approaches, including
9 bi-parental quantitative trait locus (QTL) mapping (Frei *et al.*, 2008) and Genome-Wide
10 Association Study (GWAS, Ueda *et al.*, 2015). Theoretically, chromosomal fragments
11 associated with ozone tolerance traits can then be introgressed into recipient varieties using
12 marker assisted backcrossing without the need for large-scale fumigation experiments.

13 Although no large-scale marker-assisted breeding programs for ozone tolerance in crops have
14 been conducted to date, proof of concept has been shown for ozone tolerant rice breeding
15 lines carrying QTL for ozone tolerance (Chen *et al.*, 2011; Frei *et al.*, 2008, 2010) that have a
16 superior performance to the recipient varieties in terms of yield components (Wang *et al.*,
17 2014) and grain quality (Jing *et al.*, 2016). This example should encourage further breeding
18 efforts in rice and other crop species, specifically targeting widely grown mega-varieties of
19 crops grown in ozone-affected parts in the world. As an alternative strategy, traits
20 contributing to ozone tolerance could be incorporated into existing crop varieties through
21 genetic engineering. For example, rice and barley varieties engineered to contain enhanced
22 levels of ascorbate have been engineered and showed enhanced tolerance to a variety of
23 environmental stresses (Ali *et al.*, 2018).

Physiological trait modelling could also be used to understand how different traits intended to confer tolerance for ozone, might influence crop physiology, growth and yield response under a range of environmental conditions and stresses.

4.2.4 Reducing ozone uptake by strategic limitation of irrigation application

Ozone impacts on crops could be reduced by partial stomatal closure induced by reduced irrigation, which could also save water use for irrigated crop production. In the rice growing countries, in response to the increasing water demands by other sectors than agriculture, alternate wetting and drying irrigation (AWD) has become popular in an attempt to reduce water usage and methane emissions (Bouman *et al.*, 2007; Carrijo *et al.*, 2017). This approach could also potentially be exploited to reduce ozone impacts on rice or other crops. A comparison of two studies conducted about 30 km apart in the same city of China suggests such a possibility. In Zhang *et al.* (2009), AWD with moderate water stress increased the growth and yield of rice while reducing stomatal conductance compared to continuously flooded crops, mostly resulting from a greater number of rice grains per panicle under AWD. Interestingly, at a nearby site, elevated ozone reduced rice yield arising from a decrease in the number of grains per panicle in two of the four varieties tested (Shi *et al.*, 2009). This suggests that reduced ozone uptake could be an additional and unintended benefit of AWD for farmers. The potential benefits of the AWD approach require further study.

4.2.5 Fertilizer application to compensate for crop yield losses

Crop loss from ozone exposure could potentially be counteracted by increasing the fertilizer application rate (Cardoso-Vilhena & Barnes, 2001; Chen *et al.*, 2011). However, in addition to the cost of fertilizer, recent analysis has indicated that this mitigation approach may be associated with an aggravation of other environmental problems. It has been shown that the nitrogen/protein yield of wheat is reduced by ozone at a certain level of nitrogen application

and this applies also to other nutrients like phosphorus and potassium (Broberg *et al.*, 2015; 2017). This means that the fraction of nitrogen applied which does not end up in the grain could enhance other environmental problems (Di & Cameron 2002; Mosier *et al.* 1998) such as nitrate leaching, conversion of fertilizer to N₂, emissions of N₂O and even NO, which promotes further ozone formation, as shown for pasture (Sánchez-Martín *et al.*, 2017). Adding nitrogen fertiliser to compensate for reductions in yield may also inadvertently increase the stomatal conductance of leaves of crop plants thereby increasing ozone uptake and subsequent damage (Mills *et al.*, 2016).

4.2.6 Chemical protection against ozone damage

There is scope for investigating the benefits of chemical protection against ozone damage. The most successful antiozonant applied so far has been ethylenediurea (N-[2-(2-oxo-1-imidazolidinyl)ethyl]-N'-phenylurea), abbreviated to EDU, first described by Carnahan *et al.* (1978). This chemical is usually applied as a foliar spray or soil drench, and has been used extensively in experiments and biomonitoring programmes to reduce the effects of ozone pollution, including preventing visible ozone injury on the leaves and growth and yield reductions (Agathokleous *et al.*, 2016; Feng *et al.*, 2010; Jiang *et al.*, 2018; Manning *et al.*, 2011; Pandey *et al.*, 2015; Rai *et al.*, 2015). A meta-analysis suggested that the antiozonant activity of EDU is biochemical rather than biophysical (Feng *et al.*, 2010). Recent results showed that EDU has no negative effects on plants at low O₃ concentration, but increases the crop yield at high O₃ concentration (Ashrafuzzaman *et al.*, 2017). Whilst EDU has not yet been evaluated for application at field scale, concerns have been raised about potential toxicity to aquatic plants (Agathokleous *et al.*, 2016) and more research is needed to determine if this chemical could be extensively used.

Other chemical protectants against ozone could be developed from a knowledge of plant hormonal control of stomatal functioning and stress perception (Wilkinson *et al.*, 2011), and

could potentially provide multi-stress tolerance such as combined tolerance of ozone, heat and drought stress. All three of these stresses induce synthesis of the crop stress hormone, ethylene, and chemicals that inhibit ethylene perception such as 1-MCP (1-methylcyclopropene) have the potential to reduce their effects (Wilkinson & Davies, 2010; Wagg, 2012). Anti-transpirants that reduce stomatal aperture could also reduce ozone effects by reducing ozone uptake in some species. However, there is a growing body of knowledge that chronic exposure to ozone reduces the ability of stomata to respond to abscisic acid under drought conditions, potentially leading to more rather than less ozone uptake (Mills *et al.*, 2016; Wilkinson and Davies, 2009, 2010). An alternative chemical protection approach has also been explored experimentally. Di-1-*p*-methene, a natural terpenic polymer derived from the resin of pine trees that mimics isoprene emissions from plants, has been shown to reduce visible injury in Pinto beans after exposure to 150 ppb of ozone for 4h (Francini *et al.*, 2010).

So far, chemical protection has only been explored at the experimental scale. Given the growing evidence presented here and elsewhere of the negative effects of the pollutant at the global scale, there is considerable scope for developing a chemical protectant against ozone damage, especially if it provides cross-tolerance against other co-occurring stresses.

5. Conclusions

This global-scale study shows that ozone is a very important stress, limiting yields of key crops and comparing in importance with other key stresses. For example in India, where food security concerns are particularly pressing, the mean YCS for effects of ozone on wheat of 3 falls in between those for nutrients and aridity (score 2) and for pests and diseases and heat stress (score 4). Globally, we show that the largest effects of ozone are often in areas already challenged by other stresses such as pests and diseases and heat, particularly in EAS, SAS

1 and SEA. The global mean ozone yield gaps of 4.4 – 12.4 % identified here add up to 227 Tg
2 of lost yield for soybean, wheat, rice and maize. We speculate that, ozone could at least
3 partially, account for the unexplained yield gaps and stagnation in yield improvement seen in
4 many areas of the world in recent years. Thus, international effort to reduce ozone pollution
5 on a global scale would bring clear benefits for agriculture as well as for other types of
6 vegetation, health, materials and climate change (Simpson *et al.*, 2014). However, it is likely
7 to take many decades to achieve the required emission reductions, which is the only long-
8 term solution for reducing the problems caused by tropospheric ozone. Meanwhile, the
9 global population is expected to grow significantly, which together with increasing real
10 income levels, will see increasing demands placed on food production (Tilman *et al.*, 2011).
11 Several interim solutions for closing the ozone yield gap have been outlined in this paper.
12 These include: testing of current varieties for ozone sensitivity and selection of the most
13 tolerant; crop breeding for multiple stress tolerance, including ozone; implementation of
14 protective watering regimes such as AWD; and the development of chemical protection
15 against ozone damage. Given the severity of ozone effects on staple food crops in areas of
16 the world that are also challenged by other stresses, we recommend increased attention to the
17 benefits that could be gained from taking mitigating action to reduce the ozone yield gap.

18

19 **Acknowledgements**

20 We thank the Natural Environment Research Council (NERC projects NEC05574 and
21 NEC06476) for financial support and the Adlerbertska Foundation for supporting this study
22 by funding Gina Mills' guest professorship at the University of Gothenburg (NERC project
23 NEC05831). The Adlerbertska Foundation are also thanked for funding a workshop on the
24 theme of this paper, attended by the authors, as part of the Guest Professorship funding for

Gina Mills. This work has been partially funded by EMEP under UNECE, and the EU project ECLAIRE (project no. 282910). Computer time for EMEP model runs was supported by the Research Council of Norway through the NOTUR project EMEP (NN2890K) for CPU. We also wish to thank Edmar Teixeira for providing advice on calculating a heat stress index.

Data Accessibility Statement:

National-scale data are provided in the Supporting Information. Grid square values for ozone metrics for the LRTAP region can be downloaded from http://www.emep.int/mscw/mscw_data.html.

Statement on Competing Interests: none

The authors declare no competing interests.

References

- Ainsworth EA (2008) Rice production in a changing climate: a meta-analysis of responses to elevated carbon dioxide and elevated ozone concentration. *Global Change Biology* 14, 1642-1650.
- Ainsworth, EA (2016) Understanding and improving global crop response to ozone Pollution. *The Plant Journal* 90, 886–897.
- Agathokleous E, Mouzaki-Paxinou A-C, Saitanis C *et al.* (2016) The first toxicological study of the antiozonant and research tool ethylene diurea (EDU) using a *Lemna minor* L. bioassay: Hints to its mode of action. *Environmental Pollution* 213, 996-1006.

- 1 Akhtar N, Yamaguchi M, Inada H *et al.* (2010) Effects of ozone on growth, yield and leaf gas
2 exchange rates of four Bangladeshi cultivars of rice (*Oryza sativa* L.). *Environmental*
3 *Pollution* 158, 2970-2976.
- 4 Ali B, Pantha S, Acharya R *et al.* Abiotic stress tolerance of transgenic cereal crops with
5 enhanced ascorbate biosynthesis. *Plant Biotechnology Journal*, submitted.
- 6 Atlin GN, Cairns JE, Das B (2017) Rapid breeding and varietal replacement are critical to
7 adaptation of cropping systems in the developing world to climate change. *Global*
8 *Food Security* 12, 31-37.
- 9 Ashrafuzzaman M, Farzana AL, Holtkamp F *et al.* (2017) Diagnosing ozone stress and
10 differential tolerance in rice (*Oryza sativa* L.) with ethylenediurea (EDU).
11 *Environmental Pollution* 230, 339-350.
- 12 Asseng S, Cammarano D, Basso B (2017). Hot spots of wheat yield decline with rising
13 temperatures. *Global Change Biology* 23, 2464–2472.
- 14 Avnery S, Mauzerall DL, Liu J. & Horowitz LW (2011a) Global crop yield reductions due to
15 surface ozone exposure: 1. Year 2000 crop production losses and economic damage.
16 *Atmospheric Environment* 45, 2284-2296.
- 17 Avnery S, Mauzerall DL, Liu J. & Horowitz LW (2011b). Global crop yield reductions due
18 to surface ozone exposure: 2. Year 2030 potential crop production losses and
19 economic damage under two scenarios of O₃ pollution. *Atmospheric Environment* 45,
20 2297-2309.
- 21 Avnery S, Mauzerall DL, Fiore AM (2013) Increasing global agricultural production by
22 reducing ozone damages via methane emission controls and ozone-resistant cultivar
23 selection. *Global Change Biology* 19, 1285-1299.

- 1 Betzelberger AM, Gillespie KM, McGrath JM *et al.* (2010) Effects of chronic elevated ozone
2 concentration on antioxidant capacity, photosynthesis and seed yield of 10 soybean
3 cultivars. *Plant Cell and Environment* 33, 1569-1581.
- 4 Bhatia A, Ghosh A, Kumar V *et al.* (2011) Effect of elevated tropospheric ozone on methane
5 and nitrous oxide emission from rice soil in north India. *Agriculture, Ecosystems and*
6 *Environment* 144, 21–28.
- 7 Biswas DK, Xu H, Yang JC *et al.* (2009) Impacts of methods and sites of plant breeding on
8 ozone sensitivity in winter wheat cultivars. *Agriculture, Ecosystems & Environment*
9 134, 168-177.
- 10 Black VJ, Black CR, Roberts JA, Stewart CA (2000) Tansley Review No. 115 Impact of
11 ozone on the reproductive development of plants. *New Phytologist* 147, 421-447.
- 12 Bouman BAM, Lampayan RM, Tuong TP (2007) Water management in irrigated rice.
13 Coping with water scarcity. Manila, Philippines: International Rice Research Institute.
14 p 53
- 15 Broberg MC, Feng Z, Xin Y, Pleijel H (2015) Ozone effects on wheat grain quality - A
16 summary. *Environmental Pollution* 197, 203-213.
- 17 Broberg MC, Uddling J, Mills G, Pleijel H (2017) Fertilizer efficiency in wheat is reduced by
18 ozone pollution. *Science of the Total Environment* 607-608, 876-880.
- 19 Broberg MC, Feng Z, Xin Y, Pleijel H (2015) Ozone effects on wheat grain quality – a
20 summary. *Environmental Pollution* 197, 203-213.
- 21 Burkey KO, Carter TE (2009) Foliar resistance to ozone injury in the genetic base of U.S.
22 and Canadian soybean and prediction of resistance in descendent cultivars using
23 coefficient of parentage. *Field Crops Research* 111, 207-217.

- 1 Burkey KO, Miller JE, Fiscus EL (2005) Assessment of ambient ozone effects on vegetation
2 using snap bean as a bioindicator species. *Journal of Environmental Quality* 34,
3 1081–1086.
- 4 Burney J & Ramanathan V (2014) Recent climate and air pollution impacts on Indian
5 agriculture. *Proceedings of the National Academy of Sciences* 111, 16319–16324.
- 6 CABI (2005) Crop Protection Compendium, 2005 edition. Wallingford, UK.
- 7 Cardoso-Vilhena J, Barnes J (2001) Does nitrogen supply affect the response of wheat
8 (*Triticum aestivum* cv. Hanno) to the combination of elevated CO₂ and O₃? *Journal*
9 *of Experimental Botany* 52, 1901–1911.
- 10 Carnahan JE, Jenner EL, Wat EKW (1978) Prevention of ozone injury to plants by a new
11 protectant chemical. *Phytopathology* 68, 1225–1229.
- 12 Carrijo DR, Lundy ME, Linquist BA (2017) Rice yields and water use under alternate
13 wetting and drying irrigation: A meta-analysis. *Field Crops Research* 203, 173–180.
- 14 Challinor AJ, Ewert F, Arnold S *et al.* (2009) Crops and climate change: progress, trends, and
15 challenges in simulating impacts and informing adaptation, *Journal of Experimental*
16 *Botany*, 60, 2775–2789.
- 17 Challinor AJ, Wheeler TR, Craufurd PQ, Slingo JM (2005) Simulation of the impact of high
18 temperature stress on annual crop yields. *Agricultural and Forest Meteorology*,
19 135(1–4), 180–189.
- 20 Chang KL, Petropavlovskikh I, Cooper OR *et al.* (2017) Regional trend analysis of surface
21 ozone observations from monitoring networks in eastern North America, Europe and
22 East Asia, *Elementa: Science of the Anthropocene*. doi:
23 <http://doi.org/10.1525/elementa.243>

- 1 Chen CP, Frei M, Wissuwa M (2011a) The OzT8 locus in rice protects leaf carbon
2 assimilation rate and photosynthetic capacity under ozone stress. *Plant Cell and*
3 *Environment* 34, 1141-1149.
- 4 Chen J, Zeng Q, Zhu JG *et al.* (2011b) Nitrogen supply mitigates the effects of elevated [O₃]
5 on photosynthesis and yield in wheat. *Chinese Journal of Plant Ecology* 35, 523–530.
- 6 CLRTAP (2017) Chapter 3 "Mapping Critical Levels for Vegetation". LRTAP Convention
7 Modelling and Mapping Manual. <http://icpvegetation.ceh.ac.uk/>
- 8 Debona D, Rodrigues FÁ, Rios JA *et al.* (2014) Limitations to photosynthesis in leaves of
9 wheat plants infected by *Pyricularia oryzae*. *Phytopathology* 104, 34-39.
- 10 Dentener F, Guizzardi D. 2013. TDHTAP Work Package 2.1. Common set of regions.
11 Available at: [http://iek8wikis.iek.fz-](http://iek8wikis.iek.fz-juelich.de/HTAPWiki/WP2.1?action=AttachFile&do=view&target=_region_definition_document_v4.docx)
12 [juelich.de/HTAPWiki/WP2.1?action=AttachFile&do=view&target=_region_definitio](http://iek8wikis.iek.fz-juelich.de/HTAPWiki/WP2.1?action=AttachFile&do=view&target=_region_definition_document_v4.docx)
13 [n_document_v4.docx](http://iek8wikis.iek.fz-juelich.de/HTAPWiki/WP2.1?action=AttachFile&do=view&target=_region_definition_document_v4.docx) Accessed 11 August, 2016.
- 14 Deryng D, Conway D, Ramankutty N *et al.* (2014) Global crop yield response to extreme
15 heat stress under multiple climate change futures. *Environmental Research Letters*
16 9(3), 034011.
- 17 Di HJ, Cameron KC (2002) Nitrate leaching in temperate agroecosystems: sources, factors
18 and mitigating strategies. *Nutrient Cycling in Agroecosystems* 64, 237 – 256.
- 19 Driedonks N, Rieu I, Vriezen, WH (2016) Breeding for plant heat tolerance at vegetative and
20 reproductive stages. *Plant Reproduction* 29, 67 – 79.
- 21 ECMWF. European Centre for Medium-Range Weather Forecasts European Centre for
22 Medium-Range Weather Forecasts. <http://www.ecmwf.int/> (accessed 2017).

- 1 Emberson LD, Ashmore MR, Simpson D *et al.* (2001) Modelling and Mapping Ozone
2 Deposition in Europe. *Water, Air, and Soil Pollution* 130, 577-582.
- 3 Emberson LD, Pleijel H, Ainsworth EA *et al.* (2018) Ozone effects on crops and
4 consideration in crop models. *Accepted by European Journal of Agronomy*.
- 5 FAO <http://www.fao.org/land-water/databases-and-software/crop-information/soybean/en/>
6 (accessed 2017).
- 7 FAOSTAT. Food and agriculture division of the United Nations statistics division.
8 <http://faostat3.fao.org/home/E> (accessed 2017).
- 9 Farooq M, Gogoi N, Barthakur S *et al.* (2017) Drought Stress in Grain Legumes during
10 Reproduction and Grain Filling. *Journal of Agronomy and Crop Science* 203, 81 –
11 102.
- 12 Feng ZZ, Kobayashi K (2009) Assessing the impacts of current and future concentrations of
13 surface ozone on crop yield with meta-analysis. *Atmospheric Environment* 43, 1510-
14 1519.
- 15 Feng ZZ, Kobayashi K, Ainsworth EA (2008) Impact of elevated ozone concentration on
16 growth, physiology, and yield of wheat (*Triticum aestivum* L.): a meta-analysis.
17 *Global Change Biology* 14, 2696-2708.
- 18 Feng Z, Liu X, Zhang F (2015) Air pollution effects on food security in China: Taking ozone
19 as an example. *Frontiers of Agricultural Science and Engineering* 2, 152-158.
- 20 Feng ZZ, Pang J, Nouchi I *et al.* (2010a) Apoplastic ascorbate contributes to the differential
21 ozone sensitivity in two varieties of winter wheat under fully open-air field
22 conditions. *Environmental Pollution* 158, 3539-3545.

- 1 Feng Z, Wang S, Szantoi Z *et al.* (2010b) Protection of plants from ambient ozone by
2 applications of ethylenediurea (EDU): a meta-analytic review. *Environmental*
3 *Pollution* 158, 3236-3242.
- 4 Feng Z, Wang L, Pleijel H *et al.* (2016) Differential effects of ozone on photosynthesis of
5 winter wheat among cultivars depend on antioxidative enzymes rather than stomatal
6 conductance. *Science of the Total Environment* 572, 404–411.
- 7 Fischer G, Nachtergaele FO, Prieler S *et al.* (2012) Global Agro-ecological Zones
8 Assessment for Agriculture (GAEZ v3.0). Model Documentation. IIASA, Laxenburg,
9 Austria and FAO, Rome, Italy.
- 10 Fiscus EL, Booker FL, Burkey KO (2005) Crop responses to ozone: uptake, modes of action,
11 carbon assimilation and partitioning. *Plant Cell & Environment* 28, 997-1011.
- 12 Francini A, Lorenzini G, Nali C (2011) The Antitranspirant Di1- p -menthene, a Potential
13 Chemical Protectant of Ozone Damage to Plants. *Water Air and Soil Pollution* 219,
14 459-472.
- 15 Frei M (2015) Breeding of ozone resistant rice: Relevance, approaches and challenges.
16 *Environmental Pollution* 197, 144-155.
- 17 Frei, M, Kohno, Y, Tietze S *et al.* (2012) The response of rice grain quality to ozone
18 exposure during growth depends on ozone level and genotype. *Environmental*
19 *Pollution* 163, 199-206.
- 20 Frei M, Tanaka JP, Chen CP, Wissuwa M (2010) Mechanisms of ozone tolerance in rice:
21 characterization of two QTLs affecting leaf bronzing by gene expression profiling and
22 biochemical analyses. *Journal of Experimental Botany* 61, 1405-1417.

- 1 Frei M, Tanaka JP, Wissuwa M (2008) Genotypic variation in tolerance to elevated ozone in
2 rice: dissection of distinct genetic factors linked to tolerance mechanisms. *Journal of*
3 *Experimental Botany* 59, 3741-3752.
- 4 Furher J (1994). The critical level for ozone to protect agricultural crops – An assessment of
5 data from European open-top chamber experiments. *Fuhrer J. & Achermann, B.,*
6 *(Eds). Critical Levels for Ozone. UNECE Workshop Report., 42 – 57.*
- 7 GAEZ. Global Agro-ecological Zones v3.0. <http://www.gaez.iiasa.ac.at/> (accessed 2017).
- 8 Gastal F, Lemaire G (2002) N uptake and distribution in crops: an agronomical and
9 ecophysiological perspective. *Journal of Experimental Botany* 53(370), 789-799.
- 10 Gill SS, Tuteja N (2010) Reactive oxygen species and antioxidant machinery in abiotic stress
11 tolerance in crop plants. *Plant Physiology and Biochemistry* 48, 909-930.
- 12 Gilliham M, Able JA, Roy SJ (2017a) Translating knowledge about abiotic stress tolerance to
13 breeding programmes. *The Plant Journal* 90, 898 – 917.
- 14 Gilliham M, Chapman S, Martin L *et al.* (2017b) The case for evidence-based policy to
15 support stress-resilient cropping systems. *Food and Energy Security* 6, 5 – 11.
- 16 Grassini P, Eskridge KM, Cassman KG (2013) Distinguishing between yield advances and
17 yield plateaus in historical crop production trends. *Nature Communications* 4, 2918 –
18 2929.
- 19 Harmens H, Hayes F, Mills G *et al.* (2018) Wheat yield responses to stomatal uptake of
20 ozone: Peak vs rising background ozone conditions. *Atmospheric Environment* 173, 1-
21 5.
- 22 Heagle AS (1989) Ozone and crop yield. *Annual Review of Phytopathology* 27, 397–423.

- 1 Heggstad HE (1991) Origin of Bel-W3, Bel-C and Bel-B tobacco varieties and their use as
2 indicators of ozone. *Environmental Pollution* 74, 264-291.
- 3 Hijmans RJ, Cameron SE, Parra JL *et al.* (2004) The WorldClim interpolated global
4 terrestrial climate surfaces, version 1.3. Available at <http://biogeo.berkeley.edu/>.
- 5 Hollaway MJ, Arnold SR, Challinor AJ, Emberson LD (2012) Intercontinental trans-
6 boundary contributions to ozone-induced crop yield losses in the Northern
7 Hemisphere. *Biogeosciences* 9, 271-292.
- 8 Huysmans M, Lema A S, Coll NS, Nowack MK (2017) Dying two deaths — programmed
9 cell death regulation in development and disease. *Current Opinion in Plant Biology*
10 35, 37-44.
- 11 HWSD (2009) Harmonized World Soil Database (version 1.1).
12 FAO/IIASA/ISRIC/ISSCAS/JRC. FAO, Rome, Italy and IIASA, Laxenburg, Austria.
- 13 Jaswal S (2014) Challenges to Food Security in India. *IOSR Journal of Humanities and*
14 *Social Science* 19, 93-100.
- 15 Jiang LJ, Feng ZZ, Dai LL *et al.* (2018) Large variability in ambient ozone sensitivity across
16 19 ethylenediurea-treated Chinese cultivars of soybean is driven by total ascorbate.
17 *Journal of Environmental Sciences* 64, 10-22.
- 18 Jing LQ, Dombinov V, Shen SB *et al.* (2016) Physiological and genotype-specific factors
19 associated with grain quality changes in rice exposed to high ozone. *Environmental*
20 *Pollution* 210, 397-408.
- 21 Jordan DR, Hunt CH, Cruickshank AW *et al.* (2012) The relationship between the stay-green
22 trait and grain yield in elite sorghum hybrids grown in a range of environments. *Crop*
23 *Science* 52, 1153–1161.

- 1 Kangasjarvi J, Jaspers P, Kollist H (2005) Signalling and cell death in ozone-exposed plants.
2 *Plant Cell and Environment* 28, 1021-1036.
- 3 Klingberg J, Karlsson PE, Pihl Karlsson G (2012) Variation in ozone exposure in the
4 landscape of southern Sweden with consideration of topography and coastal climate.
5 *Atmospheric Environment* 47, 252-260.
- 6 Lal S, Venkataramani S, Naja M *et al.* (2017). Loss of crop yields in India due to surface
7 ozone: an estimation based on a network of observations. *Environmental Science and*
8 *Pollution Research* 24, 20972 – 20981.
- 9 Lesk C, Rowhani P, Ramankutty N (2016) Influence of extreme weather disasters on global
10 crop production. *Nature*, 529, 84-87.
- 11 Lobell BD & Asseng S (2017) Comparing estimates of climate change impacts from process-
12 based and statistical crop models. *Environmental Research Letters*, 015001.
- 13 Lobell BD, Sibley A & Ortiz-Monasterio JI (2012) Extreme heat effects on wheat senescence
14 in India. *Nature Climate Change* 2, 186–189
- 15 Locato V, Gadaleta C, De Gara L, De Pinto MC (2008) Production of reactive species and
16 modulation of antioxidant network in response to heat shock: a critical balance for
17 cell fate. *Plant Cell and Environment* 31, 1606-1619.
- 18 Maas RP, Grennfelt RP (2016) Towards Cleaner Air. Scientific Assessment Report 2016.
19 EMEP Steering Body and Working Group on Effects of the Convention on Long-
20 Range Transboundary Air Pollution.
- 21 Manning WJ, Paoletti E, Sandermann H, Ernst (2011) Ethylenediurea (EDU): A research tool
22 for assessment and verification of the effects of ground level ozone on plants under
23 natural conditions. *Environmental Pollution* 159, 3283 – 3293.

- 1 McCormack ML, Dickie IA, Eissenstat DM et al. (2015) Redefining fine roots improves
2 understanding of below-ground contributions to terrestrial biosphere processes. *New*
3 *Phytologist* 207, 505–518.
- 4 McGrath JM, Betzelberger AM, Wang S *et al* (2015) An analysis of ozone damage to
5 historical maize and soybean yields in the United States. *Proceedings of the National*
6 *Academy of Sciences* 112(46), 14390–14395.
- 7 Mills G, Harmens H (2011) Ozone Pollution: A hidden threat to food security. Programme
8 Coordination Centre for the ICP Vegetation, Centre of Ecology & Hydrology,
9 Bangor, UK. <http://icpvegetation.ceh.ac.uk/>.
- 10 Mills G, Harmens H, Wagg S *et al.* (2016) Ozone impacts on vegetation in a nitrogen
11 enriched and changing climate. *Environmental Pollution* 208, 898 – 908.
- 12 Mills G, Hayes F, Simpson D *et al.* (2011) Evidence of widespread effects of ozone on crops
13 and (semi-)natural vegetation in Europe (1990–2006) in relation to AOT40- and flux-
14 based risk maps. *Global Change Biology* 17, 592 – 613.
- 15 Mills G, Pleijel H, Malley CS *et al.* (2018b) Tropospheric Ozone Assessment Report: Present
16 day tropospheric ozone distribution and trends relevant to vegetation. *Accepted by*
17 *Elementa: Science of the Anthropocene*.
- 18 Mills G, Sharps K, Simpson D *et al.* (2018a) Ozone pollution will compromise efforts to
19 increase global wheat production. *Global Change Biology*.
20 <https://doi.org/10.1111/gcb.14157>.
- 21 Morgan PB, Ainsworth EA, Long SP (2003) How does elevated ozone impact soybean? A
22 meta-analysis of photosynthesis, growth and yield. *Plant Cell and Environment* 26,
23 1317-1328.

- 1 Mosier A, Kroeze C, Nevison C *et al.* (1998) Closing the global N₂O budget: nitrous oxide
2 emissions through the agricultural nitrogen cycle. *Nutrient Cycling in Agroecosystems*
3 52, 225–248.
- 4 Mueller, ND, Gerber JS, Johnston M *et al.* (2012) Closing yield gaps through nutrient and
5 water management. *Nature* 490, 254-257.
- 6 Mulchi CL, Lee E, Tuthill K, Olinick EV (1988) Influence of ozone stress on growth
7 processes, yields, and grain quality characteristics among soybean cultivars.
8 *Environmental Pollution* 53, 151-169.
- 9 Munne-Bosch S, Queval G, Foyer CH (2013) The Impact of Global Change Factors on
10 Redox Signaling Underpinning Stress Tolerance. *Plant Physiology* 161, 5-19.
- 11 Oerke EC, Dehne H-W, Schönbeck F, Weber A (1994) Crop Production and Crop Protection.
12 Estimated losses in major food and cash crops. Elsevier Science, Amsterdam, the
13 Netherlands, 808 pp.
- 14 Oerke EC (2006) Crop losses to pests. *The Journal of Agricultural Science* 144, 31-43.
- 15 Osborne SA, Mills G, Hayes F *et al.* (2016) Has the sensitivity of soybean cultivars to ozone
16 pollution increased with time? An analysis of published dose–response data. *Global*
17 *Change Biology* 22, 3097 – 3111.
- 18 Pleijel H, Berglen Eriksen A, Danielsson H *et al.* (2006) Differential ozone sensitivity in an
19 old and a modern Swedish wheat cultivar – grain yield and quality, leaf chlorophyll
20 and stomatal conductance. *Environmental and Experimental Botany* 56, 63-71.
- 21 Pleijel H, Broberg MC, Uddling J, Mills G (2018). Current surface ozone concentrations
22 significantly decrease wheat growth, yield and quality. *Science of the Total*
23 *Environment* 613–614, 687–692.

- 1 Pleijel H, Danielsson H, Emberson L *et al.* (2007) Ozone risk assessment for agricultural
2 crops in Europe: Further development of stomatal flux and flux-response relationships
3 for European wheat and potato. *Atmospheric Environment* 41, 3022-3040.
- 4 Pleijel H, Danielsson H, Pihl Karlsson G *et al.* (2000). An ozone flux-response relationship
5 for wheat. *Environmental Pollution* 109, 453-462.
- 6 Pleijel H, Danielsson H, Simpson D, Mills G (2014) Have ozone effects on carbon
7 sequestration been overestimated? A new biomass response function for wheat.
8 *Biogeosciences* 11, 4521–4528.
- 9 Rai R, Agrawal M, Agrawal SB (2007) Assessment of yield losses in tropical wheat using
10 open top chambers, agriculture. *Atmospheric Environment* 41, 9543–9554.
- 11 Ray DK, Ramankutty N, Mueller ND *et al.* (2012) Recent patterns of crop yield growth and
12 stagnation. *Nature Communications* 3, 1293 – 1300.
- 13 Rebetzke GJ, Condon AG, Richards RA, Farquhar GD (2002) Selection for reduced carbon
14 isotope discrimination increases aerial biomass and grain yield of rainfed bread wheat.
15 *Crop Science* 42, 739–745.
- 16 Reynolds MP, Pierre CS, Saad ASI, Vargas M, Condon AG (2007) Evaluating potential
17 genetic gains in wheat associated with stress-adaptive trait expression in elite genetic
18 resources under drought and heat stress. *Crop Science* 47(S3), S172–S189.
- 19 Roche D (2015) Stomatal conductance is essential for higher yield of C3 crops. *Critical*
20 *Reviews in Plant Science* 34, 429 – 453.
- 21 Sánchez-Martín L, Bermejo-Bermejo V, García-Torres L *et al.* (2017) Nitrogen soil
22 emissions and belowground plant processes in Mediterranean annual pastures are
23 altered by ozone exposure and N-inputs. *Atmospheric Environment* 165, 12-22.

- 1 Sarkar A, Agrawal SB (2010) Elevated ozone and two modern wheat cultivars: an assessment
2 of dose dependent sensitivity with respect to growth, reproductive and yield
3 parameters. *Environmental and Experimental Botany* 69, 328–337.
- 4 Semenov MA & Stratonovitch P (2013) Designing high-yielding wheat ideotypes for a
5 changing climate. *Food and Energy Security* 2, 2048 – 3694.
- 6 Shi GY, Yang LX, Wang YX *et al.* (2009) Effects of elevated O₃ concentration on winter
7 wheat and rice yields in the Yangtze River Delta, China. *Agriculture, Ecosystems and*
8 *Environment* 131, 178–184.
- 9 Singh S, Agrawal SB (2011) Cultivar-Specific Response of Soybean (*Glycine max* L.) to
10 ambient and elevated concentrations of ozone under open top chambers. *Water, Air*
11 *and Soil Pollution* 217, 283-302.
- 12 Simpson D, Arneth A, Mills G *et al.* (2014) Ozone — the persistent menace: interactions
13 with the N cycle and climate change. *Current Opinion in Environmental*
14 *Sustainability*, 9–10, 9-19
- 15 Simpson D, Benedictow A, Berge H *et al.* (2012) The EMEP MSC-W chemical transport
16 model – technical description. *Atmospheric Chemistry and Physics*, 12, 7825-
17 7865.
- 18 Simpson, D, Bergström R, Imhof H, Wind P (2017) Updates to the EMEP/MSC-W model,
19 2016-2017 Transboundary particulate matter, photo-oxidants, acidifying and
20 eutrophying components. Status Report 1/2017, The Norwegian Meteorological
21 Institute, Oslo, Norway, www.emep.int, 115-122.
- 22 Simpson D, Emberson L, Ashmore M, Tuovinen J (2007) A comparison of two different
23 approaches for mapping potential ozone damage to vegetation. A model study.
24 *Environmental Pollution* 146, 715-725.

- 1 Singh E, Rai R, Pandey B, Agrawal M (2017) Development of Resistance in Two Wheat
2 Cultivars Against Constant Fumigation of Ozone. *Proceedings of the National*
3 *Academy of Sciences, India Section B: Biological Sciences*, 1-14.
- 4 Siyiannis VF, Protonotarios VE, Zechmann B *et al.* (2012) Comparative spatiotemporal
5 analysis of root aerenchyma formation processes in maize due to sulphate, nitrate or
6 phosphate deprivation. *Protoplasma*, 249, 671-686.
- 7 Suzuki N, Koussevitzky S, Mittler R, Miller G (2012) ROS and redox signalling in the
8 response of plants to abiotic stress. *Plant Cell and Environment* 35, 259-270.
- 9 Tai APK & Martin MV (2017) Impacts of ozone air pollution and temperature extremes on
10 crop yields: Spatial variability, adaptation and implications for future food security.
11 *Atmospheric Environment* 169, 11 -21.
- 12 Tao F, Zhang Z, Zhang S. *et al.* (2016) Variability in crop yields associated with climate
13 anomalies in China over the past three decades. *Regional Environmental Change* 16,
14 1715 – 1723.
- 15
- 16 Tang H, Takigawa M, Liu G *et al.* (2013) A projection of ozone-induced wheat production
17 loss in China and India for the years 2000 and 2020 with exposure-based and flux-
18 based approaches. *Global Change Biology* 19, 2739-2752.
- 19 Teixeira EI, Fischer G, van Velthuisen H *et al.* (2013). Global hot-spots of heat stress on
20 agricultural crops due to climate change. *Agricultural and Forest Meteorology* 170,
21 206-215.
- 22 Tilman D, Balzer C, Hill J, Befort BL (2011) Global food demand and the sustainable
23 intensification of agriculture. *Proceedings of the National Academy of Sciences* 108,
24 20260-20264

- 1 Tomer R, Bhatia A, Kumar V *et al.* (2015) Impact of elevated ozone on growth, yield and
2 nutritional quality of two wheat species in Northern India. *Aerosol and Air Quality*
3 *Research* 15, 329-340.
- 4 Trabucco A & Zomer RJ (2009) Global Aridity Index (Global-Aridity) and Global Potential
5 Evapo-Transpiration (Global-PET) Geospatial Database. *CGIAR Consortium for*
6 *Spatial Information*. Published online, available from the CGIAR-CSI GeoPortal at:
7 <http://www.csi.cgiar.org>.
- 8 Ueda Y, Frimpong F, Qi Y *et al.* (2015a) Genetic dissection of ozone tolerance in rice (*Oryza*
9 *sativa* L.) by a genome-wide association study. *Journal of Experimental Botany* 66,
10 293-306.
- 11 Ueda Y, Siddique S, Frei M (2015b) A Novel Gene, OZONE-RESPONSIVE APOPLASTIC
12 PROTEIN1, Enhances Cell Death in Ozone Stress in Rice. *Plant Physiology* 169,
13 873-889.
- 14 UNEP (United Nations Environment Programme), (1997) World atlas of desertification 2ED.
15 UNEP, London.
- 16 Van Dingenen R, Dentener FJ, Raes F *et al.* (2009) The global impact of ozone on
17 agricultural crop yields under current and future air quality legislation. *Atmospheric*
18 *Environment* 43, 604-618.
- 19 Van Doorn WG (2011) Classes of programmed cell death in plants, compared to those in
20 animals. *Journal of Experimental Botany* 62, 4749-4761.
- 21 Vollsnes AV, Kruse OMO, Eriksen AB *et al.* (2010) *In vivo* root growth dynamics of ozone
22 exposed *Trifolium subterraneum*. *Environmental and Experimental Botany* 69, 183-
23 188.

- 1 Wagg SK (2012) A mechanistic study of the implications of ozone and drought effects on
2 vegetation for global warming. PhD Thesis, University of Lancaster.
- 3 Wang Y, Frei M (2011) Stressed food - The impact of abiotic environmental stresses on crop
4 quality. *Agriculture Ecosystems & Environment* 141, 271-286.
- 5 Wang Y, Yang L, Hoeller M *et al.* (2014) Pyramiding of ozone tolerance QTLs OzT8 and
6 OzT9 confers improved tolerance to season-long ozone exposure in rice.
7 *Environmental and Experimental Botany* 104, 26-33.
- 8 Wassmann R, Jagadish SVK, Heuer S *et al.* (2009) Climate change affecting rice production:
9 the physiological and agronomic basis for possible adaptation strategies. *Advances in*
10 *Agronomy* 101, 59-122.
- 11 Wild O, Fiore AM, Shindell DT *et al.* (2012) Modelling future changes in surface ozone: a
12 parameterized approach. *Atmospheric Chemistry and Physics*, 12, 2037-2054.
- 13 Wilkinson S, Davies WJ (2009) Ozone suppresses soil drying- and abscisic acid (ABA)-
14 induced stomatal closure via an ethylene-dependent mechanism. *Plant, Cell and*
15 *Environment* 32, 949-959.
- 16 Wilkinson S, Davies WJ (2010) Drought, ozone, ABA and ethylene: new insights from cell
17 to plant to community. *Plant, Cell and Environment* 33, 510-525.
- 18 Wilkinson S, Mills G, Illidge R *et al.* (2012) How is ozone pollution reducing our food
19 supply? *Journal of Experimental Botany* 63, 527-536.
- 20 Wu L-B, Ueda Y, Lai S-K, Frei M (2017) Shoot tolerance mechanisms to iron toxicity in rice
21 (*Oryza sativa* L.). *Plant Cell and Environment* 40, 570-584.
- 22 Yendrek CR, Koester RP, Ainsworth EA (2015) A comparative analysis of transcriptomic,
23 biochemical, and physiological responses to elevated ozone identifies species-specific

- 1 mechanisms of resilience in legume crops. *Journal of Experimental Botany* 66, 7101-
2 7112.
- 3 Yendrek CR, Erice G, Montes CM *et al.* (2017) Elevated ozone reduces photosynthetic
4 carbon gain by accelerating leaf senescence of inbred and hybrid maize in a
5 genotype-specific manner. *Plant Cell and Environment* 40, 3088-3100.
- 6 Yuan XY, Calatayud V, Jiang LJ *et al.* (2015) Assessing the effects of ambient ozone in
7 China on snap bean genotypes by using ethylenediurea (EDU). *Environmental*
8 *Pollution* 205, 199-208.
- 9 Zhang H, Xue Y, Wang Z *et al.* (2009) An alternate wetting and moderate soil drying regime
10 improves root and shoot growth in rice. *Crop Science* 49, 2246–2260
- 11 Zhou B, Elazab A, Bort J *et al.* (2016) Agronomic and physiological responses of Chinese
12 facultative wheat genotypes to high-yielding Mediterranean conditions. *Journal of*
13 *Agricultural Science* 154, 870–889.
- 14 Zhu X, Feng Z, Sun T *et al.* (2011) Effects of elevated ozone concentration on yield of four
15 Chinese cultivars of winter wheat under fully open-air field conditions. *Global*
16 *Change Biology* 17, 2697–2706.
- 17 Zobel RW (2016) Fine roots – functional definition expanded to crop species? *New*
18 *Phytologist* 212, 310-312.
- 19 Zobel RW, Baligar VC, Kinraide TB (2007) Fine root diameters can change in response to
20 changes in nutrient concentrations. *Plant and Soil* 297, 243–254.
- 21

Figure legends

Fig. 1. Response functions for (a) soybean, (b) wheat, (c) rice and (d) maize derived from published data using the growing season ozone (7h mean, M7 in ppb) in the experiments. Data points are presented per cultivar/variety, with sources of data provided in the Supporting Information (Table S3). The response functions are: Soybean, $RY = -0.0050x + 1.001$ (r^2 (adj) = 0.625, $p < 0.001$); Wheat, $RY = -0.0048x + 0.96$ (r^2 (adj) = 0.547, $p < 0.001$); Rice, $RY = -0.0021x + 0.987$ (r^2 (adj) = 0.347, $p < 0.001$); and Maize, $RY = -0.0031x + 1.03$ (r^2 (adj) = 0.617, $p < 0.001$).

Fig. 2: The global effects of five biotic and abiotic stresses on soybean. All data are presented for the $1 \times 1^\circ$ grid squares where the mean production of soybean was > 500 tonnes (0.0005 Tg). (a) Presents the effects of ozone on crop production (thousand tonnes or 0.001 Tg per grid square), and (b) the percentage yield loss due to ozone, averaged for the period 2010-2012. In (d) to (h), the Yield Constraint Score (YCS) is presented per grid square on a scale of 1 to 5, where 5 is the highest level of stress (see Table 1) for ozone, pests and diseases, nutrients, heat stress and aridity, respectively, whilst (c) is the total YCS (YCS_{all}) calculated from the sum of each of these per grid square. The regional impacts are summarised in Table 2 for the five highest producing regions and Table S4 provides all country and regional means.

Fig. 3: The global effects of five biotic and abiotic stresses on wheat. All data are presented for the $1 \times 1^\circ$ grid squares where the mean production of wheat was > 500 tonnes (0.0005 Tg). (a) Presents the effects of ozone on crop production (thousand tonnes or 0.001 Tg per grid square), and (b) the percentage yield loss due to ozone, averaged for the period 2010-2012. In (d) to (h), the Yield Constraint Score (YCS) is presented per grid square on a scale of 1 to 5, where 5 is the highest level of stress (see Table 1) for ozone, pests and diseases, nutrients, heat stress and aridity, respectively, whilst (c) is the total YCS (YCS_{all}) calculated from the sum of each of these per grid square. The regional impacts are summarised in Table 2 for the five highest producing regions and Table S5 provides all country and regional means.

Fig. 4: The global effects of five biotic and abiotic stresses on rice. All data are presented for the $1 \times 1^\circ$ grid squares where the mean production of rice was > 500 tonnes (0.0005 Tg). (a) Presents the effects of ozone on crop production (thousand tonnes or 0.001 Tg per grid square), and (b) the percentage yield loss due to ozone, averaged for the period 2010-2012. In (d) to (h), the Yield Constraint Score (YCS) is presented per grid square on a scale of 1 to 5, where 5 is the highest level of stress (see Table 1) for ozone, pests and diseases, nutrients, heat stress and aridity, respectively, whilst (c) is the total YCS (YCS_{all}) calculated from the sum of each of these per grid square. The regional impacts are summarised in Table 2 for the five highest producing regions and Table S6 provides all country and regional means.

1 **Fig. 5: The global effects of five biotic and abiotic stresses on maize.** All data are
2 presented for the 1 x 1° grid squares where the mean production of maize was > 500 tonnes
3 (0.0005 Tg). (a) Presents the effects of ozone on crop production (thousand tonnes or 0.001
4 Tg per grid square), and (b) the percentage yield loss due to ozone, averaged for the period
5 2010-2012. In (d) to (h), the Yield Constraint Score (YCS) is presented per grid square on a
6 scale of 1 to 5, where 5 is the highest level of stress (see Table 1) for ozone, pests and
7 diseases, nutrients, heat stress and aridity, respectively, whilst (c) is the total YCS (YCS_{all})
8 calculated from the sum of each of these per grid square. The regional impacts are
9 summarised in Table 2 for the five highest producing regions and Table S7 provides all
10 country and regional means.

11

12 **Fig. 6: Yield Constraint Score (YCS) for five constraints on the yield of (a) wheat and**
13 **(b) rice in the five Indian states with the highest production per crop.** The bars represent
14 the mean YCS per 1 x 1° grid square per state on a scale of 1 to 5, where 5 is the highest level
15 of stress (see Table 1), rounded to the nearest integer. Note: The YCS for pests and disease is
16 only available at the National Scale for India (score 4 for wheat and 5 for rice) and is
17 presented here for information.

18

19 **Fig. 7: An ideotype for an ozone-tolerant crop.** ‘+’ indicates where there would be a
20 benefit for other stresses of improving tolerance to ozone for the trait, whilst ‘-’ indicates a
21 trade-off, and ‘0’ is no effect.

22

23

- 1
- 2
- 3
- 4

For Review Only




Fig. 1. Response functions for (a) soybean, (b) wheat, (c) rice and (d) maize derived from published data using the growing season ozone (7h mean, M7 in ppb) in the experiments. Data points are presented per cultivar/variety, with sources of data provided in the Supporting Information (Table S3). The response functions are: Soybean, $RY = -0.0050x + 1.001$ (r^2 (adj) = 0.625, $p < 0.001$); Wheat, $RY = -0.0048x + 0.96$ (r^2 (adj) = 0.547, $p < 0.001$); Rice, $RY = -0.0021x + 0.987$ (r^2 (adj) = 0.347, $p < 0.001$); and Maize, $RY = -0.0031x + 1.03$ (r^2 (adj) = 0.617, $p < 0.001$).

149x132mm (600 x 600 DPI)




Fig. 2: The global effects of five biotic and abiotic stresses on soybean. All data are presented for the $1 \times 1^\circ$ grid squares where the mean production of soybean was > 500 tonnes (0.0005 Tg). (a) Presents the effects of ozone on crop production (thousand tonnes or 0.001 Tg per grid square), and (b) the percentage yield loss due to ozone, averaged for the period 2010-2012. In (d) to (h), the Yield Constraint Score (YCS) is presented per grid square on a scale of 1 to 5, where 5 is the highest level of stress (see Table 1) for ozone, pests and diseases, nutrients, heat stress and aridity, respectively, whilst (c) is the total YCS (YCSall) calculated from the sum of each of these per grid square. The regional impacts are summarised in Table 2 for the five highest producing regions and Table S4 provides all country and regional means.

249x290mm (600 x 600 DPI)




Fig. 3: The global effects of five biotic and abiotic stresses on wheat. All data are presented for the 1 x 1° grid squares where the mean production of wheat was > 500 tonnes (0.0005 Tg). (a) Presents the effects of ozone on crop production (thousand tonnes or 0.001 Tg per grid square), and (b) the percentage yield loss due to ozone, averaged for the period 2010-2012. In (d) to (h), the Yield Constraint Score (YCS) is presented per grid square on a scale of 1 to 5, where 5 is the highest level of stress (see Table 1) for ozone, pests and diseases, nutrients, heat stress and aridity, respectively, whilst (c) is the total YCS (YCSall) calculated from the sum of each of these per grid square. The regional impacts are summarised in Table 2 for the five highest producing regions and Table S5 provides all country and regional means.

249x290mm (600 x 600 DPI)




Fig. 4: The global effects of five biotic and abiotic stresses on rice. All data are presented for the $1 \times 1^\circ$ grid squares where the mean production of rice was > 500 tonnes (0.0005 Tg). (a) Presents the effects of ozone on crop production (thousand tonnes or 0.001 Tg per grid square), and (b) the percentage yield loss due to ozone, averaged for the period 2010–2012. In (d) to (h), the Yield Constraint Score (YCS) is presented per grid square on a scale of 1 to 5, where 5 is the highest level of stress (see Table 1) for ozone, pests and diseases, nutrients, heat stress and aridity, respectively, whilst (c) is the total YCS (YCSall) calculated from the sum of each of these per grid square. The regional impacts are summarised in Table 2 for the five highest producing regions and Table S6 provides all country and regional means.

249x290mm (600 x 600 DPI)




Fig. 5: The global effects of five biotic and abiotic stresses on maize. All data are presented for the 1 x 1° grid squares where the mean production of maize was > 500 tonnes (0.0005 Tg). (a) Presents the effects of ozone on crop production (thousand tonnes or 0.001 Tg per grid square), and (b) the percentage yield loss due to ozone, averaged for the period 2010-2012. In (d) to (h), the Yield Constraint Score (YCS) is presented per grid square on a scale of 1 to 5, where 5 is the highest level of stress (see Table 1) for ozone, pests and diseases, nutrients, heat stress and aridity, respectively, whilst (c) is the total YCS (YCSall) calculated from the sum of each of these per grid square. The regional impacts are summarised in Table 2 for the five highest producing regions and Table S7 provides all country and regional means.

249x290mm (600 x 600 DPI)




Fig. 6: Yield Constraint Score (YCS) for five constraints on the yield of (a) wheat and (b) rice in the five Indian states with the highest production per crop. The bars represent the mean YCS per $1 \times 1^\circ$ grid square per state on a scale of 1 to 5, where 5 is the highest level of stress (see Table 1), rounded to the nearest integer. Note: The YCS for pests and disease is only available at the National Scale for India (score 4 for wheat and 5 for rice) and is presented here for information.

168x170mm (150 x 150 DPI)




Fig. 7: An ideotype for an ozone-tolerant crop. '+' indicates where there would be a benefit for other stresses of improving tolerance to ozone for the trait, whilst '-' indicates a trade-off, and '0' is no effect.

249x367mm (600 x 600 DPI)

GCB-18-0501: Response to Reviewers**Closing the global ozone yield gap: Quantification and co-benefits for multi-stress tolerance**

Gina Mills, Katrina Sharps, David Simpson, Håkan Pleijel, Michael Frei, Kent Burkey, Lisa Emberson, Johan Uddling, Malin Broberg, Zhaozhong Feng, Kazuhiko Kobayashi and Madhoolika Agrawal

Note: All page and line numbers refer to the manuscript with changes accepted. An additional version of the manuscript showing all changes made is also provided, for information.

Changes made to improve the integration of the difference components of the paper

Having read the editor's and reviewer's comments, we have made some small changes to the text and structure to improve the integration of the different components of the paper for the reader. Excluding the introduction, for each section, we have introduced a higher-level sub-division that distinguishes between text related to quantifying the problem and text related to solutions for the problem. The structural changes made are:

Introduction

To ensure that the flow of the paper is clearer for the reader, we have modified:

- (i) The last two sentences of the first paragraph to refer back to the title of the paper and introduce the idea that the paper both defines the problem and suggests solutions (P4, L 2- 5).
- (ii) The last paragraph to reflect the (revised) structure of the paper, including specifically mentioning that the paper describes results from two analytical approaches (quantitative spatial analysis and a qualitative analysis leading to an ozone tolerant crop ideotype), and also includes an extended discussion considering options for reducing the negative effects of ozone on yield (P6, L16 to P7, L18).

Methods

The different sub-sections have been re-organised as follows:

2.1 Global spatial analysis of crop yield constraints caused by ozone

- 2.1.1 Crop production
- 2.1.2 Intra- and inter-specific sensitivity of crops to ozone
- 2.1.3 Yield constraints caused by ozone

2.2 Global spatial analysis of yield constraints caused by other stresses

- 2.2.1 Yield constraints caused by pests and diseases
- 2.2.2 Yield constraints caused by soil nutrients
- 2.2.3 Yield constraints caused by heat stress
- 2.2.4 Yield constraints caused by aridity

2.3 Comparative analysis of effects of five stresses using a Yield Constraint Score (YCS)

2.4. Qualitative analysis of plant traits associated with multiple stress tolerance

Results

The different sub-sections have been re-organised as follows:

3.1 Quantification of the global impacts of ozone and other stresses on crop yield

3.1.1 Intra-specific sensitivity to ozone

3.1.2 Spatial analysis of the global impacts of multiple stresses on crop yield

3.1.3 Case study – India

3.2. Plant traits associated with tolerance of ozone and associated stresses in crops

Discussion

The different sub-sections have been re-organised as follows:

4.1 The global scale of ozone impacts on crops relative to impacts of other stresses

4.2 Options for reducing ozone impacts on crops

4.2.1 Global effort to reduce ozone precursor emissions [*Note: new section added, see response to Reviewer 1*]

4.2.2 Exploiting existing varietal differences in ozone sensitivity

4.2.3 Breeding new varieties with multiple stress tolerance, including of ozone

4.2.4 Reducing ozone uptake by strategic limitation of irrigation application

4.2.5 Fertilizer application to compensate for crop yield losses

4.2.6 Chemical protection against ozone damage

Conclusions

Structure unchanged, 2 paragraphs

Reviewer 1 (Reviewer's comments in black font, [our response in blue font](#))

This is very important contribution to the Scientific Community in particular on the relationship between ozone pollution and its impacts on crop production. I think the review paper has summarized almost all recently progress in the area.

[We wish to thank this reviewer for their very supportive comments about the importance of our manuscript for the scientific community and for making suggestions for further improvements.](#)

I have only two minor suggestions: (1) Discussion sector: The options (as a separate paragraph) for reducing surface ozone concentration or pollution should be considered (which can be regarded as the most important option) to reduce ozone impacts crops;

Thank you for this suggestion. We have added a new section, 4.2.1 on “Global effort to reduce ozone precursor emissions” encompassing this suggestion (P40, L11 to P42, L6).

(2) One additional reference which has also addressed this issue could be added in the revision: Feng Z, Liu X, Zhang F. (2015) Air pollution effects on food security in China: Taking ozone as an example. *Front. Agr. Sci. Eng.* 2(2): 152-158.

We have included reference to this paper in the new section 4.2.1 (P41, L17 - 20).

Reviewer 2 (Reviewer's comments in black font, our response in blue font)

Comments to the Author

Based on published data in some literatures, the authors made a comprehensive prediction of ozone impacts on major grain crop yield in the world by using models. They also compared the impacts of ozone with those of other stresses (pest, heat, etc.). They found the largest loss in yield was soybean with significant spatial differences. To my knowledge, this is the first comprehensive report about ozone impacts on the four major grain crops globally. They authors also provided some adaptation strategies in crop improvement and agronomic innovations for ozone impact mitigation for food security.

We wish to thank reviewer 2 for also recognising the importance of our study including its novelty, and for making suggestions for further improvements.

However, there are still some limitations:

(1) The available published data is limited, especially about ozone impacts on maize, the average response functions (Fig. 1) might not represent the specific cropping regions or countries. Moreover, the experimental results from a specific location were not only due to ozone elevation, the results might also due to the interaction between ozone and the local temperature and other environmental factors. Thus the experimental locations in the data base should be representative of the major cropping regions of each grain crop (Soybean, wheat, maize, and rice), especially for the prediction of total production loss.

In producing the response functions, we sought to include all available experimental data that met the inclusion criteria listed in Section 2.1.2. For soybean, wheat and rice, the experimental data provided good representation of the major growing areas for these crops. A new sentence has been added to Section 3.1.1 that lists the HTAP regions represented in the response functions for these three crops (P17, L20 to P18, L1) and reference is now made to this point in the discussion (P35, L2 – 7). Unfortunately, for maize we were limited to data for three cultivars from experiments conducted in the USA during the late 1980s/early 1990s. Thus, the analysis conducted for maize has the highest uncertainty. However, as the highest maize production losses were predicted for the USA where the experiments had been conducted, we considered it relevant to include the analysis for maize within the paper to ensure that we covered all of the main grain crops. We have added new text on the uncertainty associated with the maize analysis in the discussion (Section 4.1, P35, L5 - 8), whereas previously this uncertainty was only mentioned in T1 of the Supporting Information.

(2) In this manuscript, the authors only compared ozone with pest and disease, soil nutrient, heat and aridity. In fact, there are other important stresses, such as global warming, extreme climate events, flooded, etc. These stresses might have more serious impacts on crop production.

We can fully understand the reviewer's point, but unfortunately it was not feasible to consider every stress that could have a potentially large effect on yield in the paper. The stresses chosen were for illustration purposes – to place ozone effects within the context of effects of example other common stresses that also reduce yield. A detailed justification for the stresses chosen, including reference to other studies using these datasets or approaches is provided in the Supporting Information, Text 1. We thought that inclusion of this text within the discussion of the main paper would have made the discussion too long both for the reader and the journal. When choosing stresses, we were largely restricted by availability of datasets and methods suitable for quantifying impacts of current levels of stress. Heat stress was included as an example of an effect of extreme climatic events, with the heat stress index being determined for crop-specific thermal sensitive time-periods over the period 1990-2014.

To address the reviewer's concerns, we have added in the word “example” where we first mention comparison with other abiotic and biotic stress in the Introduction (P7, L5) and discussion (P35, L10 - 12) to show that we are not covering all such stresses. We have also added new text to Section 4.1 of the discussion pointing out that other detrimental stresses such as flooding and global warming have been excluded from this study but their effects can be significant (P35, L12 - 15).

(3) For each prediction, the authors need to give the scenario time and the baseline time. For example, the readers cannot get the information about the prediction years for each Fig. of the manuscript.

Thank you for pointing this out. Whilst this information is included in the methods section, we agree that it was not easy to find quickly in the original manuscript. To avoid overly-long repetitive legends for each figure, we have added a new column to Table 1 ‘Year(s) of data’ (see P16) to facilitate comparisons across stresses and also to make this information easy to find.

(4) About the section of crop ideotype, it would be better more it to the section of discussion.

Whilst writing the paper and after receiving this comment, we discussed at length the best location for this part of the paper. The ideotype is a result in that it is based on a synthesis of information from the scientific literature. Conversely, the associated text explaining the ideotype reads more like a discussion. We are concerned that placing both in the discussion might make the discussion overly long. After further consideration, we think it is better to retain this figure and text in the results (Section 3.2), with a description of how the ideotype could be included in crop breeding located in the discussion (Section 4.2.3). As described above, we have added higher level headings throughout the methods, results and discussion sections to make it clearer when we are quantifying the ozone problem in the context of other stresses, and when we are reporting on options for reducing ozone impacts. We believe that this insertion of extra sub-headings will make the paper more streamlined for the reader and easier to follow.

(5) About the differences in ozone response between crop cultivars, can the authors get the corresponding yield of each cultivar? So the authors can compare the differences the ozone response and the crop yield. Did the higher sensitive cultivars have lower yield under an ambient ozone level? If it is true, this really indicates that there is a potential to enhance crop tolerance to ozone with high yield through cultivar improvement.

This is an important point that we have now added to the discussion (P42, L20 to P43, L2). Because of the different growing and environmental conditions in each experiment it is not possible to compare the actual yield directly across cultivars unless the cultivars have been exposed to ozone at the same location and in the same year. For this reason, the Fuhrer method was developed for international risk assessment. This approach allows cultivars to be compared in response functions based on their relative yield, and has been used in this study to prepare Figure 1. In revising the manuscript, we have inserted new text to provide examples from experiments conducted in the last decade where ozone sensitive varieties of rice, wheat and soybean have significantly higher yield in filtered air with reduced ozone compared to ambient air treatments (P42, L22 to P43, L2).

Or Review Only

1 **Closing the global ozone yield gap: Quantification and co-benefits**
2 **for multi-stress tolerance**

3 **Authors:** Gina Mills^{1,2*}, Katrina Sharps¹, David Simpson^{3,4}, Håkan Pleijel², Michael Frei⁵,
4 Kent Burkey⁶, Lisa Emberson⁷, Johan Uddling², Malin Broberg², Zhaozhong Feng⁸,
5 Kazuhiko Kobayashi⁹ and Madhoolika Agrawal¹⁰

6
7 **Affiliations**

- 8 ¹Centre for Ecology and Hydrology, Bangor, UK.
9 ²Department of Biological and Environmental Sciences, University of Gothenburg, Sweden.
10 ³EMEP MSC-W, Norwegian Meteorological Institute, Norway.
11 ⁴Department of Space, Earth & Environment, Chalmers University of Technology, Sweden.
12 ⁵Institute of Crop Science and Resource Conservation, University of Bonn, Germany
13 ⁶USDA-ARS, Raleigh, North Carolina, USA
14 ⁷Stockholm Environment Institute, University of York, UK.
15 ⁸State Key Laboratory of Urban and Regional Ecology, Research Center for Eco-
16 Environmental Sciences, Chinese Academy of Sciences, China.
17 ⁹Department of Global Agricultural Sciences, The University of Tokyo, Japan.
18 ¹⁰Department of Botany, Institute of Science, Banaras Hindu University, India.

19 ***Correspondence** to: Prof. Gina Mills, Centre for Ecology and Hydrology, Environment
20 Centre Wales, Deiniol Road, Bangor, UK. LL57 2UW. Email: gmi@ceh.ac.uk. Tel. +44
21 [\(0\)1248 374500](tel:+441248374500).

22

Running head: Global benefits of closing the O₃ yield gap

Key words: Ozone, wheat, soybean, maize, rice, pests and diseases, aridity, nutrient stress, heat stress, stress tolerant ideotype.

Paper type: Primary research

Abstract

Increasing both crop productivity and the tolerance of crops to abiotic and biotic stresses are major challenges for global food security in our rapidly changing climate. For the first time, we show how the spatial variation and severity of tropospheric ozone effects on yield compare with effects of other stresses on a global scale, and discuss mitigating actions against the negative effects of ozone. We show that the sensitivity to ozone declines in the order soybean > wheat > maize > rice, with genotypic variation in response being most pronounced for soybean and rice. Based on stomatal uptake, we estimate that ozone (mean of 2010 - 2012) reduces global yield annually by 12.4%, 7.1%, 4.4% and 6.1% for soybean, wheat, rice and maize, respectively (the ‘ozone yield gaps’), adding up to 227 Tg of lost yield. Our modelling shows that the highest ozone-induced production losses for soybean are in North and South America whilst for wheat they are in India and China, for rice in parts of India, Bangladesh, China and Indonesia, and for maize in China and the USA. Crucially, we also show that the same areas are often also at risk of high losses from pests and diseases, heat stress and to a lesser extent aridity and nutrient stress. In a solution-focussed analysis of these results, we provide a crop ideotype with tolerance of multiple stresses (including ozone) and describe how ozone effects could be included in crop breeding programmes. We also

1 | discuss altered crop management approaches that could be applied to reduce ozone impacts [in](#)
2 | [the shorter-term](#). Given the severity of ozone effects on staple food crops in areas of the
3 | world that are also challenged by other stresses, we recommend increased attention to the
4 | benefits that could be gained from addressing the ozone yield gap.

5

6 | **1. Introduction**

7 | To feed the rapidly growing global population, we need to develop a new generation of crop
8 | cultivars or varieties that will have both high productivity in future climates and high
9 | tolerance of the biotic and abiotic stresses that are likely to become more prevalent in the
10 | future (Gilliham *et al.*, 2016). Candidate characteristics or traits are currently being tested in
11 | ideotype modelling (Semenov and Stratonovitch, 2013) and include improved light
12 | conversion efficiency, a longer duration of green leaf area for grain fill, a higher harvest
13 | index and optimal phenology. For example, varieties that use less water per unit of carbon
14 | fixed will have higher yield under drought conditions (Rebetzke *et al.*, 2002) as will those
15 | with ‘stay-green’ characteristics during water stress (Jordan *et al.*, 2012). Whilst it is widely
16 | recognised that rapid breeding programmes will have a vital role to play in adaptations of
17 | crops to climate change (Atlin *et al.*, 2017), selection of traits for tolerance of one abiotic
18 | stress, tropospheric (ground level) ozone pollution, is currently omitted from such breeding
19 | programmes (Ainsworth, 2016; Frei *et al.*, 2015). This is happening even though field
20 | experiments from nine countries representing three continents have shown that reducing
21 | ozone concentrations back to pre-industrial levels would give an average wheat yield benefit
22 | of 8.4% globally (Pleijel *et al.*, 2018), a figure that is matched by modelling based on the
23 | stomatal uptake of the pollutant (Mills *et al.*, 2018a).- Furthermore, an earlier meta-analysis
24 | of crop responses to ozone suggested that current ozone levels in the range 31 - 50 ppb (nmol

mol⁻¹, v/v) are reducing the yield of major food crops by 5.3 to 19% (Feng & Kobayashi, 2009). We undertook this new study to build a case for improving crop yields in our changing climate by closing the ozone-induced yield gap via the inclusion of ozone tolerance in crop breeding programmes, altered crop management and more stringent ozone precursor emission controls. ~~We undertook this new study to build a case for including ozone tolerance in crop breeding programmes. We took a multi-faceted approach that includes a spatial analysis of the impacts of ozone on yield relative to impacts of other abiotic and biotic stresses, defining an ideotype for an ozone tolerant crop and considering options for reducing the negative effects of the pollutant by altered crop management.~~

Tropospheric ozone pollution is formed from photochemical reactions involving anthropogenic and biogenic emissions and is involved in a complex web of interactions with ecosystems (Simpson *et al.*, 2014). Whilst concentrations have been beginning to decrease in eastern USA and parts of Europe (2000 – 2014) due to precursor emission controls, they have been increasing rapidly in south (S) and east (E) Asia (Chang *et al.*, 2017). Ozone is a powerful oxidant that is absorbed into leaves via open stomatal pores. Once inside the leaves, ozone reacts with biomolecules to form reactive oxygen species, triggering defence mechanisms that if overwhelmed lead to programmed cell death and a reduced extent and duration of functional green leaf area producing less photosynthate for seed fill (e.g. Ainsworth, 2016). Since pests and diseases (e.g. Oerke, 2006; Huysmans *et al.*, 2017), heat stress (e.g. Driedonks, *et al.*, 2016), drought (e.g. Farooq *et al.*, 2017) or reduced nutrient availability (e.g. Gastal & Lemaire, 2002) usually also reduce the extent and duration of the functional green leaf area, then in simple terms, each of these biotic and abiotic stresses result in the same endpoints – reduced yield quantity that is often associated with reduced quality.

1 So far, most crop breeding programmes have been targeted at increasing or maintaining the
2 yield rather than increasing stability of yield under stress (Gilliham *et al.*, 2017). Because
3 ozone concentrations tend to be very ~~heterogenous~~heterogeneous across natural and
4 agricultural regions (Klingberg *et al.*, 2012) as well as over seasons and years, it is not likely
5 that traditional selection would unintentionally favour ozone tolerant crop genotypes. The
6 reverse seems to be the case. For example, an analysis of ozone-exposure yield data for 49
7 soybean varieties from 28 field exposure studies showed that ozone sensitivity has increased
8 by an average of 33% between 1960 and 2000 (Osborne *et al.*, 2016). Similarly, modern
9 wheat varieties are more sensitive than older varieties (Biswas *et al.*, 2008; Pleijel *et al.*,
10 2006). Potentially, this increased sensitivity to ozone over recent decades is related to
11 selective breeding for higher stomatal conductance (Roche, 2015) that inadvertently has
12 increased the ingress of ozone into crops (Biswas *et al.*, 2008; Osborne *et al.*, 2016); further
13 study is required to fully understand the mechanistic basis of this increasing sensitivity with
14 time.

15 As with many abiotic and biotic stresses, genetic variation in plant response to ozone has
16 been found for every species that has been tested. For the major grain crops, genetic variation
17 in ozone response has been reported for wheat (Zhu *et al.*, 2011), rice (Frei *et al.*, 2008; Shi *et al.*,
18 2009), soybean (Mulchi *et al.*, 1988; Burkey & Carter, 2009; Jiang *et al.*, 2018), and
19 maize (Yendrek *et al.*, 2017). Variation has also been reported for other crops including snap
20 bean (Burkey *et al.*, 2005; Yuan *et al.*, 2015) and tobacco (Heggestad, 1991). These
21 assessments are based on different criteria including foliar injury and impacts on growth and
22 yield parameters. Taken together, the evidence suggests that sufficient natural genetic
23 variation exists to support improvement in crop stress tolerance either as sources of ozone
24 tolerance genes or providing contrasting genotypes for mechanism studies to identify targets
25 for molecular manipulation. Potential targets for breeding of ozone tolerance that have the

1 greatest likelihood of success include reducing the stomatal uptake of ozone into the leaf and
2 increasing its detoxification once inside the leaf (Feng *et al.*, 2016; Frei *et al.*, 2015).
3 To target the regions of the world where ozone tolerant crop varieties are most required, we
4 need to understand which crops are most at risk and where they are growing in relation to
5 current high-risk areas for ozone. We know from a recent analysis of ozone concentrations at
6 over 3000 rural sites that the highest ozone values are in many of the world's important crop
7 growing regions, including parts of the USA, Europe, India and China (Mills *et al.*,
8 [2018bsubmitted](#)). Overall, the latter study showed that the global mean cumulative ozone
9 exposure is double the critical level set by the United Nations as a target for ozone pollution
10 control, above which direct adverse effects on sensitive vegetation may occur according to
11 present knowledge (CLRTAP, 2017). Several studies have modelled ozone concentrations
12 and predicted yield effects using concentration-based yield response functions applied at a
13 range of scales from local (e.g. for India, Lal *et al.*, 2017) to global (e.g. Avnery *et al.*,
14 2011a,b; Van Dingenen *et al.*, 2009). Whilst these studies indicate effects in the highest
15 ozone areas, they do not take into account the constantly varying effects of soil moisture, air
16 temperature, light and humidity on the uptake of the pollutant via the stomata. In Europe,
17 field evidence for effects of ozone on crops and other types of vegetation shows that risk
18 assessments based on modelled stomatal uptake or flux (Emberson *et al.*, 2000, Simpson *et*
19 *al.*, 2007) provide a stronger indication of ozone effects than those based on concentration
20 (Mills *et al.*, 2011). Furthermore, dose-response functions for crops that are based on
21 stomatal uptake are better correlated with yield effects than those based on concentration
22 (Pleijel *et al.*, 2000, 2007), providing additional support for their use.

23 With ozone concentrations increasing in rapidly developing regions and predicted to continue
24 to increase in coming decades (Wild *et al.*, 2012), it is timely to consider the options for
25 increasing the tolerance of crops to this abiotic stress. In this study, [our analysis included a](#)

1 two-step approach to addressing the ozone problem in crops: (i) a quantitative spatial analysis
 2 of the impacts of ozone on crop yield relative to impacts of other abiotic and biotic stresses
 3 and (ii) a qualitative analysis of crop traits, including defining an ideotype with multiple
 4 stress tolerance. As an initial step, we compiled dose-response data from experiments
 5 conducted around the world to determine the scope for breeding ozone tolerant varieties by
 6 showing the genotypic range in sensitivity for four staple crops: soybean, wheat, rice and
 7 maize. We then used the response functions to modelled the current impacts of ozone on
 8 each crop, showing the regions where the greatest production losses are likely to be
 9 occurring. Whilst we wait for ozone effects to be included in predictive crop yield modelling
 10 (as suggested by, for example, Challinor *et al.*, 2009; Emberson *et al.*, 2018; Lobell &
 11 Asseng, 2017), we sought to compare on a global scale the impacts of ozone on yield with the
 12 influence of other example biotic and abiotic stresses. Those selected were: pests and diseases
 13 (Oerke *et al.*, 2006); aridity (Trabucco & Zomer, 2009); heat stress (developed from Deryng
 14 *et al.* (2014) and Teixeira *et al.* (2013)); and soil nutrient status-stress (GAEZ). The effects of
 15 all five stresses were considered this in more detail for India where there are major challenges
 16 for crop production and food security (Jaswal, 2014) and where global assessments
 17 consistently predict high risk from elevated ozone (e.g. Avnery *et al.*, 2011a,b; Van Dingenen
 18 *et al.*, 2009). In the second part of the study, we conducted an
 19 We also analysis of the plant traits associated with multiple stress tolerance, and considered
 20 the trade-offs and benefits of introducing ozone tolerance in crops for cross-tolerance of other
 21 biotic and abiotic stresses. This part of the study culminated in the design of ed-an ideotype
 22 for an ozone tolerant crop that would also provide tolerance of against the co-occurring
 23 stresses. -In an extended discussion, we assess the results from the two parts of the study and
 24 consider address viable approaches to options for reducing reduce the negative effects of

ozone on yield, including ~~options for~~ crop management, breeding and global efforts to
reducing ozone pollution.

2. Materials and Methods

2.1 Global spatial analysis of crop yield constraints caused by ozone

2.1.1 Crop production

Global modelled crop production data (year 2000, 0.0833° (5 arc minute) resolution) was downloaded from the GAEZ (Global Agro-Ecological Zones, v. 3) data portal (<http://www.fao.org/nr/gaez/en/>) for soybean, wheat, rice and maize. Irrigated and rain-fed production data was collected for each crop. Using ArcMap v. 10.3 (Environmental Systems Research Institute, Redlands, CA, USA), a 1° by 1° global grid was created. For each crop, production was summed per grid cell. Each cell was classed as irrigated or non-irrigated based on the percentage of irrigated crop production per cell. To define a threshold for irrigated versus non-irrigated, we first produced frequency distributions of the percentage of irrigated production for each crop (Supporting Information, Fig. S1). These showed that the majority of cells for each crop were either fully irrigated or fully rain-fed. A threshold of 75% irrigated was used to identify those cells where the majority of the production was on irrigated land. Production for the period 2010-12 was estimated per grid cell by applying a conversion factor from FAOSTAT national production data available, averaged for the years 1999 – 2001 (average production for 2010-12/average production for 1999-2001). Only cells with > 500 tonnes (0.0005 Tg) crop production in 2010-12 were included in the analysis.

As discussed in Mills *et al.* (2018^a), each 1° by 1° grid cell was assigned to a climatic zone, using the global ‘Climatic Zone’ GIS raster layer produced by the European Soil Data Centre (ESDAC) at JRC (Joint Research Centre). For each climatic zone, a 90 day growing period

1 was derived per crop (Table S1), with climatic zones illustrated in Fig. S2. Data sources for
2 assigning crop timings are provided with Table S1. For ease of comparison of effects
3 between crops, only the main growing season per year was used for each crop.

4 **2.1.2 Intra- and inter-specific sensitivity of crops to ozone**

5 To determine the relative sensitivity of the four crops to ozone together with the between-
6 variety variation in response to the pollutant, it was necessary to update existing response
7 functions based on ozone concentration as stomatal uptake-based functions are currently only
8 available for wheat. We collated dose-response data from the scientific literature using the
9 method developed by Osborne *et al.* (2016) for soybean and the commonly reported ozone
10 metric, M7 (7 hour mean, averaged from 09:00 to 15:59). The soybean dose response
11 relationship from Osborne *et al.* (2016) was included in our analysis, whilst response
12 functions for wheat, rice and maize provided in Mills & Harmens (2011) were updated with
13 more recent published data (Web of Science and Google Scholar searches conducted between
14 April and October 2017 using the search terms "ozone and yield and *crop name*"). Studies
15 were only included if they met a number of selection criteria. The duration of ozone exposure
16 must have spanned at least 60% of the 90 day growing season for each crop and ozone levels
17 during exposure were up to 100 ppb for wheat and 170 ppb for other crops. Experiments were
18 included if carried out in Open Top Chambers (OTCs), ambient air or large closed
19 chambers/greenhouses (with the air stirred by fans, minimum size 2.6 by 2.2m). Data from
20 both container and field-sown experiments were used to ensure a wide variety of points from
21 different varieties were included. If the seasonal M7 was not given in the text, this was
22 calculated either using the conversion equations provided in Osborne *et al.* 2016 (e.g. for 24
23 hour mean to M7) or information contained in the experimental methodology of the study. As
24 there was no new published data available at the time of analysis for maize, the response
25 function from Mills & Harmens (2011) was used. Yield data from different experiments were

standardised as first described by Fuhrer (1997) and recently re-described by Osborne *et al.* 2016. Thus, for each set of experimental data, linear regression was used to determine the yield at 0 ppb of ozone (the intercept of the line); this value was the reference for calculating the relative yield (i.e. relative yield = actual yield/yield at 0 ppb).

Individual variety dose-response functions were derived for wheat and rice for the four varieties with the most data points. Following Osborne *et al.* 2016, yield reduction estimates (RYL_{c,p}) were then calculated for varieties showing statistically significant declines in yield with increasing ozone by calculating the difference in percentage yield loss at 55ppb (representing current M7) relative to that at 23 ppb (representing pre-industrial M7).

2.1.3 Yield constraints caused by ozone

The EMEP MSC-W (European Monitoring and Evaluation Programme, Meteorological Synthesising Centre-West) chemical transport model (version 4.16, Simpson *et al.*, 2012, 2017) was used to derive daily POD₃IAM (Phytotoxic Ozone Dose above 3 nmol m⁻² s⁻¹, parameterised for Integrated Assessment Modelling, CLRTAP, 2017) values for the years 2010 to 2012 per 1° by 1° grid cell as described by Mills *et al.* (2018a). POD₃IAM is parameterised for a generic crop represented by wheat (CLRTAP, 2017) and represents the accumulated stomatal uptake of ozone, modelled from the hourly mean values for ozone, temperature, vapour pressure deficit, irradiance and soil moisture (Mills *et al.* 2018a). Evaluation of the EMEP model performance is also presented in Mills *et al.*, 2018a, and is summarised in the Supporting Information for the current paper (T1).

For each crop, the accumulated 90 day POD₃IAM was then calculated per cell using appropriate climate-specific 90 day growing periods (Table S1, Fig. S2), and an average calculated for the period 2010-2012. For example, for soybean in warm temperate climates in the Northern Hemisphere, the time interval was day 182 to day 271. The EMEP model

1 generated irrigated (without soil water limitation) and non-irrigated (rain limited) POD₃IAM
2 values. For grid cells classed as irrigated for each crop (See Section 2.1.1), the irrigated
3 POD₃IAM value was used to calculate percentage yield loss, otherwise the non-irrigated
4 POD₃IAM was used. This approach allowed crop-specific irrigation usage to be taken into
5 account, and was different to Mills *et al.* (2018a) where POD₃IAM values were weighted by
6 the proportion of irrigation use within a 1 x 1° cell. The global distribution of POD₃IAM for
7 each crop is provided in Fig. S3.

8 Yield loss due to ozone was first calculated for wheat using the most recent methodology
9 adopted by CLRTAP, 2017. This method also differed slightly from that used in our earlier
10 study (Mills *et al.*, 2018a) in that a reference POD₃IAM value to represent ozone uptake at
11 pre-industrial or natural ozone levels was subtracted before crop loss was calculated
12 (CLRTAP, 2017). This value (0.1 mmol m⁻², Equ. 1) was the mean POD₃IAM for the
13 experimental conditions included in the dose-response relationship, assuming constant 10 ppb
14 ozone throughout the 90d period. The equation used to determine percentage yield loss was:

15 % Yield loss = (POD₃IAM – 0.1)* 0.64 [Equ. 1]

16 Where 0.64 is the slope of the relationship between POD₃IAM and percentage yield reduction
17 (Mills *et al.*, 2018a) and represents the percentage reduction per mmol m⁻² POD₃IAM.


18 For soybean, maize and rice, the climate-specific grid square POD₃IAM values were first
19 used to calculate yield loss using the wheat equation (Equ. 1), and the resultant value was
20 then multiplied by the relative sensitivity of the crop compared to wheat, RS_w. The latter was
21 derived by dividing the slope of the M7 response function for the crop (Fig. 1) by that for
22 wheat. Production loss per crop was calculated per grid square using the following equation:

23 Production loss (tonnes) = Crop production * (% yield loss/100) [Equ. 2]

24 **2.2 Global spatial analysis of yield constraints caused by other stresses**

2.2.12.4-Yield constraints caused by pests and diseases

Oerke *et al.* (1994; 2006) provide estimates for pre-harvest crop losses due to weeds, animal pests, (arthropods, nematodes, mammals, slugs and snails, birds), pathogens and viruses for several major global crops, using data compiled from the literature. This database provides regional percentage yield loss estimates up to 2004 for 11 crops, including soybean, wheat, rice and maize, and is available from the Centre for Agriculture and Biosciences International (CABI) Crop Protection Compendium (CABI, 2005). A value for mean percentage yield loss due to pests and diseases for the period 2002-04 was assigned to each 1° by 1° grid cell, based on the country and region of the world the cell was located in. If a cell contained land from more than one country, it was assigned to a country based on where the majority of the crop was growing in the cell. Data was available for 19 global regions (Oerke *et al.*, 2006). In this study, data were used that represented the remaining crop yield losses after crop protection practices had been applied.



Formatted: Font: (Default) Times New Roman, 12 pt, Bold

Fig. 1. Response functions for (a) soybean, (b) wheat, (c) rice and (d) maize derived from published data using the growing season ozone (7h mean, M7 in ppb) in the experiments. Data points are presented per cultivar/variety, with sources of data provided in the Supporting Information (Table S3). The response functions are: Soybean, $RY = -0.0050x + 1.001$ (r^2 (adj) = 0.625, $p < 0.001$); Wheat, $RY = -0.0048x + 0.96$ (r^2 (adj) = 0.547, $p < 0.001$); Rice, $RY = -0.0021x + 0.987$ (r^2 (adj) = 0.347, $p < 0.001$); and Maize, $RY = -0.0031x + 1.03$ (r^2 (adj) = 0.617, $p < 0.001$).

Comment [MGE1]: Higher resolution figure inserted

2.2.25 Yield constraints caused by soil nutrients

Soil nutrient classifications (nutrient availability and nutrient retention) at 0.083° by 0.083° resolution were downloaded from the GAEZ (v. 3) data portal (<http://www.fao.org/nr/gaez/en/>) in June, 2017. The soil qualities (nutrient availability and retention) in the GAEZ dataset have been derived from combinations of soil attributes, using

1 data in the Harmonized World Soil Database (HWSD, v. 1.1, FAO/IIASA/ISRIC/ISS
 2 CAS/JRC 2009). Nutrient availability refers to soil fertility, and classification is based on soil
 3 texture, soil organic carbon, soil pH and total exchangeable bases. The nutrient retention
 4 capacity of soil is based on the ability of soil to retain added nutrients against losses due to
 5 leaching. Classification of nutrient retention has been derived from soil texture, base
 6 saturation, cation exchange capacity of the soil and of the clay fraction and soil pH. In the
 7 GAEZ dataset, nutrient availability and retention are classed separately for topsoil (0-30cm)
 8 and subsoil (30-100cm) and then combined by weighting based on the prevalence of active
 9 roots (Fischer *et al.*, 2012). The GAEZ classes for soil nutrient availability and nutrient
 10 retention were combined in this study to produce five soil nutrient stress classes (summarised
 11 in Table 1, further details provided in Table S2). The soil nutrient class making up the
 12 majority of each 1° by 1° grid cell in areas where crops were growing, was used to represent
 13 the class for each cell.

14 **2.2.36 Yield constraints caused by heat stress**

15 Following the methods of Challinor *et al.* (2005), subsequently used by a number of other
 16 studies (e.g. Deryng *et al.*, 2014; Teixeira *et al.*, 2013), a heat stress index was calculated per
 17 grid cell for each crop to determine if the daily temperature within a 30 day thermal sensitive
 18 period (TSP) exceeded the tolerance thresholds for each crop. This method assumes that
 19 damage to crops occurs when daily temperatures exceed a critical temperature (T_{crit} , °C) and
 20 maximum damage occurs when temperatures exceed the limit temperature (T_{lim} °C). Using
 21 information on the reproductive phase for each crop (FAO), the thermal sensitive period was
 22 designated as days 40-70 of the 90 day growing period (which varies with climate zone for
 23 each crop, Table S1). Following Deryng *et al.*, (2014), the daily effective temperature (T_{eff} ,
 24 °C, (daily mean temp + daily max temp)/2), used as a measure of the daily temperature when
 25 photosynthesis is taking place, was calculated per grid cell using global hourly temperature

1 data for the period 1990-2014 at 0.5° by 0.5° resolution. The temperature data were from the
2 European Centre for Medium-Range Weather Forecasts Integrated Forecasting System
3 (ECMWF-IFS, www.emwf.int/research/ifsdocs/), as prepared for use by the EMEP model.
4 For each crop, a daily heat stress value (f_{HSd}) was then calculated for each day within the
5 TSP, per grid cell. As we required an index that could be used to detect increasing levels of
6 stress (i.e. an index scaled from 0 to 1), heat stress was calculated following Teixeira *et al.*
7 (2013) (Eqn. 3).

8

9

10

11

$$f_{\text{HSd}} = \begin{cases} 0 & \text{for } T_{\text{eff}} < T_{\text{crit}} \\ \frac{T_{\text{eff}} - T_{\text{crit}}}{T_{\text{lim}} - T_{\text{crit}}} & \text{for } T_{\text{crit}} \leq T_{\text{eff}} < T_{\text{lim}} \\ 1 & \text{for } T_{\text{eff}} \geq T_{\text{lim}} \end{cases} \quad [\text{Equ. 3}]$$

12

13 An average value was then calculated across the 30 day TSP to give the final heat stress
14 index value, (f_{HS}) per grid cell (Eqn. 4).

15

16

$$f_{\text{HS}} = \frac{\sum_{j=1}^{\text{TSP}} (f_{\text{HSd}})}{\text{TSP}} \quad [\text{Equ. 4}]$$

17 Critical and limiting temperatures per crop were taken from Deryng *et al.* (2014) (maize,
18 wheat and soybean) and Teixeira *et al.* (2013) (rice). These were: Soybean (35°C for T_{crit} and
19 40°C for T_{lim}), wheat (25 and 35 °C), rice (35 and 45°C) and maize (32 and 45°C).

20 | **2.2.47 Yield constraints caused by aridity**

Global Aridity Index data (Trabucco & Zomer 2009) were downloaded from the CGIAR-CSI GeoPortal (<http://www.csi.cgiar.org>). The mean Aridity Index for the period 1950-2000 (0.0083° by 0.0083° resolution) was calculated as:

$$\text{Aridity Index (AI)} = \text{MAP} / \text{MAE} \quad [\text{Equ. 5}]$$

Where MAP is the Mean Annual Precipitation and MAE is the Mean Annual Potential Evapo-Transpiration.

Mean annual precipitation values were obtained from the WorldClim Global Climate Data (Hijmans *et al.* 2004), for years 1950-2000, while mean annual values of Potential Evapo-Transpiration (PET) were calculated using the average monthly PET values from the Global-PET model (Trabucco & Zomer, 2009). The mean Aridity Index per cell was calculated for each 1° by 1° grid cell where there is production for the crop.

2.3 Comparative analysis of effects of five stresses using a Yield Constraint Score (YCS)

As percentage yield loss data was only available for ozone and pests and diseases data, a percentage scale could not be used for all stresses. To overcome this problem, a yield constraint score (YCS) on a scale of 1 – 5 was developed for each abiotic and biotic stress to show spatially where each constraint is predicted to be impacting on yield and to provide some indication of the magnitude of the effect (Table 1). Yield loss was split into the same five percentage yield loss classes for ozone and pests and diseases, with the highest class being >40% and expected to be comparable to severe stress for all yield constraints. Soil nutrient retention and availability were combined to give five overall classes (Table S2). The aridity climate classes used were from the Generalized Climate Classification Scheme

(UNEP 1997), while the heat stress index was classified following the methods of Teixeira *et al.* (2013). To identify those areas of the world with the highest combined stresses, the YCS for all five stresses were summed (YCS_{all}).

Table 1: Categories of Yield Constraint Score (YCS) for ozone, pests and diseases, soil nutrients, heat and aridity (See text for explanations and justifications of categories).

Comment [MGE2]: Table replaced with editable version. New column 'year(s) of data' added.

Formatted: Font: (Default) Times New Roman

Yield constraint score (YCS)		1	2	3	4	5
Ozone	% Yield Loss	0 to 5	5 to 10	10 to 25	25 to 40	> 40
Pests and diseases	% Yield Loss	0 to 5	5 to 10	10 to 25	25 to 40	> 40
Nutrients	Retention	None or slight	Slight to moderate	Slight to severe	Moderate to severe	Moderate to very severe
	Availability	None	Slight to moderate	Moderate to severe	Severe	Severe to very severe
	Overall	None	Slight	Moderate	Severe	Very severe
Heat	Index	0	<0.05	0.05 to 0.15	0.15 to 0.3	> 0.3
Aridity	Index	>0.65	0.5 to 0.65	0.2 to 0.5	0.03 to 0.2	<0.03
	Climate class	Humid	Dry sub-humid	Semi-arid	Arid	Hyper arid

8

2.9 Regional classification of impacts

9

Stress	Attribute	Year(s) of data	Yield constraint score (YCS)				
			1	2	3	4	5
Ozone	% Yield Loss	Mean of 2010-2012	0 to 5	5 to 10	10 to 25	25 to 40	> 40
Pests and diseases	% Yield Loss	Mean of 2002 - 2004	0 to 5	5 to 10	10 to 25	25 to 40	> 40
Soil Nutrients	Retention	HWSD data, 2009, downloaded	None or slight	Slight to moderate	Slight to severe	Moderate to severe	Moderate to very severe

	<u>Availability</u>	<u>in June, 2017</u>	<u>None</u>	<u>Slight to moderate</u>	<u>Moderate to severe</u>	<u>Severe</u>	<u>Severe to very severe</u>
	<u>Overall</u>		<u>None</u>	<u>Slight</u>	<u>Moderate</u>	<u>Severe</u>	<u>Very severe</u>
<u>Heat</u>	<u>Index</u>	<u>Mean of 1990 - 2014</u>	<u>0</u>	<u><0.05</u>	<u>0.05 to 0.15</u>	<u>0.15 to 0.3</u>	<u>> 0.3</u>
<u>Aridity</u>	<u>Index</u>	<u>Mean of 1950 - 2000</u>	<u>>0.65</u>	<u>0.5 to 0.65</u>	<u>0.2 to 0.5</u>	<u>0.03 to 0.2</u>	<u><0.03</u>
	<u>Climate class</u>		<u>Humid</u>	<u>Dry sub-humid</u>	<u>Semi-arid</u>	<u>Arid</u>	<u>Hyper arid</u>

For description of effects of the five stresses, results are described as regional and national averages, with the mean YCS and YCS_{all} rounded to the nearest integer, reflecting their categorical nature. The regional classification of countries used is that adopted by the Task Force on Hemispheric Transport of Air Pollutants (HTAP) of the Convention on Long-range Transboundary Air Pollution (LRTAP, Dentener & Guizzardi, 2013). Region names are provided in full in the text the first time they are used and thereafter are referred to by the HTAP three letter codes. The region names, three letter codes and a map illustrating the countries included per region are provided in Fig. S4.

2.4 Qualitative analysis of plant traits associated with ~~10 Defining a crop ideotype for ozone and multiple stress tolerance~~

The scientific literature on crop stress tolerance was reviewed between June and December, 2017, with the aim of developing an ideotype for an ozone- and multi-stress tolerant crop. This analysis identified target traits to induce ozone tolerance, including reducing the effects on panicles, leaves and roots. It also considered the benefits and trade-offs for tolerance of other stresses, of introducing ozone tolerance into crops.

3. Results

3.1 Quantification of the global impacts of ozone and other stresses on crop yield

3.1.1 Intra-specific sensitivity to ozone

A comprehensive collation of published data on the yield responses of soybean, wheat and rice to ozone resulted in a database representing 52, 18 and 44 varieties, respectively (Fig. 1a-c, with data sources in Table S3). Ozone-response data for these three crops provides good representation of the areas where the crops were grown: Soybean (East Asia (EAS), North America (NAM) and South Asia (SAS)); wheat (Europe (EUR), EAS, SAS) and rice (EAS and SAS). In contrast, only three varieties have been tested to date for yield responses in maize (Fig. 1d), with these experiments being conducted in the USA during the 1980s and early 1990s. For each crop, there was a significant negative response to ozone ($p < 0.001$), with the slope of the negative relationships declining in the order soybean (-0.0050) > wheat (-0.0048) > maize (-0.0031) > rice (-0.0021). Within each response function, variation in ozone sensitivity due to variety provided scatter in the range of sensitivity.

For each crop, some varieties were more tolerant to ozone than others, indicating that there is scope for selecting more tolerant varieties for immediate use or as part of a breeding programme for new varieties. For soybean, $RYL_{c,p}$ ranged from 13.3% to 37.9%, with the three most sensitive varieties being the Indian varieties ‘PK472’, ‘Pusa 9712’ and ‘Pusa 9814’ (Osborne *et al.* 2016). For wheat, the $RYL_{c,p}$ for the four varieties with the most data was 16.4% (‘Drabant’), 19.3% (‘PBW 343’), 26.4% (‘Dragon’) and 32.5% (‘Albis’). While the 44 rice varieties showed a range of sensitivities to ozone (Fig. 1c), overall, indicating rice ~~it~~ was the least sensitive of the four crops investigated. Of the four rice varieties with the most data, ‘Koshihikari’ showed a $RYL_{c,p}$ of 4.7%, ‘Nipponbare’ showed no significant negative relationship between relative yield and M7 ($p > 0.05$), while ‘Kasalath’ and ‘Kirara 397’ had a $RYL_{c,p}$ of 6% and 11.1 % respectively. The rice variety ‘Pathumthani-1’ showed a higher $RYL_{c,p}$ of 18.1%, however only 5 data points were available, therefore further study may be required to confirm this result. For maize, the $r^2(\text{adj})$ was 0.62 for the response

function ($p < 0.001$, Fig. 1d), with a $RYL_{c,p}$ for all three varieties of 10%. The $RYL_{c,p}$ for ‘Pioneer 3780’ with 14 data points was 15.7% whilst $RYL_{c,p}$ was not calculated for ‘PAG 397’ as the response function for the 5 data points for this variety was not significant ($p = 0.08$).

3.1.22 Spatial analysis of the global impacts of multiple stresses on crop yield

3.2.1 Soybean

The highest ozone-associated production losses (in Tg per $1 \times 1^\circ$ grid square) for soybean are predicted to be in ~~North America (NAM)~~ and South America (SAM) (Table 2), particularly in central and E USA, S Brazil and N Argentina (Fig. 2, Table S4). However, the percentage yield losses are predicted to be lower for Brazil (12.5 – 15%) and Argentina (7.5- 10%) than for the USA (> 20% in large areas), showing that in high producing areas where ozone concentrations are more moderate (as indicated by the percentage losses), high total production losses can still be expected. In the rest of the world, production losses due to ozone in excess of 0.01 Tg per $1 \times 1^\circ$ grid square are predicted for parts of China, India and S and E Europe. In each of these areas, the production loss was not as high as expected from percentage yield losses in excess of 20%, because soybean is not widely grown.

The YCSs for ozone were mainly score 3 for the highest producing regions, with some areas with a score of 4 in E USA, NE India and China (Fig. 2, Tables 2 and S4). There is overlap between these areas and the areas with the highest YCS_{all} . Other areas with a relatively high YCS_{all} such as parts of Sub-Saharan Africa (SSA) and ~~South-East Asia (SEA)~~ are not predicted to have high production losses due to ozone because of lower percentage yield losses and/or low production totals per region. For soybean, the YCS for pests and diseases is 3 or more over most of the growing area, and particularly high (score of 5) in parts of SSA, ~~South Asia (SAS)~~ and SEA. The largest YCS values for nutrient availability (scores of 4 and

5) are in areas of SE USA, S Brazil and SEA including Thailand, Malaysia and Indonesia. Whilst heat stress YCSs are lower than those for aridity, the areas affected by both stresses largely coincided in soybean growing areas. Overall, for soybean, the global mean YCS for ozone of 3 is one category below that for pests and diseases, and one higher than that for nutrients and aridity (Table 2). The mean YCSs for ozone were in the range 2 - 3 for the five highest producing regions (SAM, NAM, EAS, SAS, Russia (RBU)), with YCSs being in the range 3 – 5, 1 – 3, 1 - 2 and 1 - 2 for pests and diseases, nutrients, heat and aridity, respectively (Table 2).

9

Table 2: Production and regional mean Yield Constraint Score (YCS, rounded to nearest integer) for the five highest producing regions for soybean, wheat, rice and maize.

Soybean	Production (Tg)	Ozone	Pests & diseases	Nutrients	Heat	Aridity
Global	253.6	3	4	2	2	2
South America	125.8	2	4	3	2	1
N America	90.3	3	3	2	2	1
East Asia	14.9	3	4	2	1	2
South Asia	13.2	3	5	1	2	2
Russia	3.6	2	4	1	1	2
Wheat	Production (Tg)	Ozone	Pests & diseases	Nutrients	Heat	Aridity
Global	673.3	2	4	2	3	2
Europe	163.8	2	3	2	2	1
East Asia	118.9	2	3	2	2	2
South Asia	118.1	2	4	2	4	2
N America	84.2	2	4	2	2	2
Russia	66.1	2	4	2	2	1
Rice	Production (Tg)	Ozone	Pests & diseases	Nutrients	Heat	Aridity
Global	716.8	1	4	2	1	2
East Asia	221.2	2	4	2	1	2
South Asia	219.2	2	5	2	2	2
South East Asia	204.5	1	4	3	1	1

South America	22.7	1	4	3	1	1
Sub Saharan Africa	21.1	1	5	2	1	2
Maize	Production (Tg)	Ozone	Pests & diseases	Nutrients	Heat	Aridity
Global	869.1	2	4	2	2	2
N America	313.1	2	3	2	2	2
East Asia	194.4	2	4	2	2	2
South America	91.7	2	4	3	2	2
Europe	75.6	2	3	1	2	1
Sub Saharan Africa	60.0	1	5	2	2	2

3.2.2 Wheat

By far the highest production losses due to ozone per grid square for wheat are predicted for India and China, with large areas in N India and NW China having over 15% yield losses amounting to production losses in excess of 0.1 Tg (Fig. 3, Table S5). Production losses are also predicted to be high in the highest wheat producing areas of Europe (including France and Germany) and central states of the USA. The mean YCS for ozone globally was 2, reflecting the same mean score in the 9 highest wheat producing regions (Tables 2 and S5), and matching that globally for nutrients, and aridity. The highest predicted production losses due to ozone only overlapped with areas of the highest YCS_{all} in NW India, Pakistan and S USA (Fig. 3). Scores of 3 and above coincide for ozone, pests and diseases, heat and nutrients in a wider area including parts of EAS, SAS and NAM, whilst YCSs for aridity are lower in several parts of this region than for ozone. This study also indicated that the highest YCS values for all stresses are for heat (score 5) in areas of Northern Africa (NAF), SAS, SAM, SSA and SAM (particularly Argentina). Scores for pests and diseases are 3 or more across most of the wheat growing areas. YCSs for nutrient availability are generally the lowest of the five stresses, although there are some high risk areas with values of 4 and above in, for example, NW SAS, Central Asia (CAS) and NE EUR (e.g. Finland), E NAM and SAM (e.g. Brazil). The highest YCSs for aridity are in a zone that includes parts of NAF, the

1 Middle East (MDE) and SAS, CAS and EAS. For the five main wheat producing regions, the
2 mean YCS for ozone was 2 representing 5 - 10% yield loss, whilst it was 3 - 4 for pests and
3 diseases and mainly 1-2 for the other three stresses (Table 2).

4 3.2.3 Rice

5 Production losses due to ozone are predicted to only be in excess of 0.1 Tg per grid square in
6 parts of India, Bangladesh, China and Indonesia (Fig. 4). In these areas, the percentage yield
7 losses are mainly in the range 7.5 – 12.5%, resulting in grid square ozone YCSs that are
8 usually either 2 or 3. Across all of the rice growing regions, the mean YCS for ozone per
9 region is either 1 or 2, resulting in a global mean score of 1 and a score of 2 in the two
10 highest producing regions (Tables 2 and S6). YCSs of 3 for ozone occurred in areas of India
11 and China where the YCS_{all} was usually in the highest range for rice of 13 – 15. Overall, the
12 highest mean YCS for this crop is for pests and diseases, being score 5 in most of the rice
13 growing areas of SSA and SAS. Nutrient YCSs are 4 or more in many parts of SSA, SAM
14 (particularly Brazil), EAS and SAS. Heat stress is predicted to be less of a problem for rice
15 than for wheat, with few regions having a score of 3 or more. Indeed, the regional mean YCS
16 for heat stress in rice is mostly either 1 or 2 (Tables 2 and S6). In the three highest producing
17 rice regions, EAS, SAS and SEA, irrigation usage is 96%, 74% and 28%, respectively (Table
18 S5). Here, the aridity score is predicted to ~~only~~ be 3 or more only in areas of NW China and
19 W India.

20 3.2.4 Maize


21 China and the USA are the two countries predicted to have the largest areas where production
22 losses due to ozone for maize that exceed 0.1 Tg per 1° x 1° grid square (Fig. 5). In these
23 areas, the percentage yield losses are mainly in the range 7.5 – 15% for the USA and 12.5 –
24 15 % for China. There are also high-risk areas in S EUR, for example, in parts of S France

1 and N Italy and in NAF (particularly Egypt). These areas generally have a YCS_{all} for maize
 2 in the range 10 – 15, and are not in the areas with the highest YCS_{all} for maize of >15. The
 3 latter are mainly found in parts of SAS and SSA, with occasional small areas elsewhere. The
 4 mean YCS for ozone for the four highest maize producing regions (NAM, EAS, SAM and
 5 EUR) is 2 (Tables 2 and S7). For the stresses other than ozone, the highest scores for YCS
 6 are for pests and diseases, with scores of 5 predicted in most of SSA, SAS and SEA. YCSs of
 7 4 and above are predicted for nutrients in much of SSA (particularly in western countries),
 8 SEA, large areas of SAM (particularly Brazil), parts of E USA and small areas of Europe.
 9 For maize, the YCSs for aridity are highest in eastern NAM and SAM, parts of NAF
 10 (particularly W Egypt), MDE, SAS (particularly NW India) and EAS (particularly NE
 11 China). Heat stress YCSs are lower for maize than for wheat, indicating that the main areas
 12 of concern for this crop are in W SSA, W MDE and SAS. Globally, the mean YCS for maize
 13 is 4 for pests and diseases and 2 for each of the other four stresses (Table 2).

14 **3.1.3 Case study - India**

15 It is clear from the results presented above that the five environmental stresses included in
 16 this study are all predicted to be having relatively high impacts on yield in several states of
 17 India. We selected this country for a more in depth analysis. Although the spatial data for
 18 India is present on the global maps in Figs 2 - 5, for ease of interpretation, we have produced
 19 additional maps for India for wheat and rice, the two most important crops by production in
 20 Fig. S5 and S6, respectively. At the national scale, the mean YCSs for the crop with the
 21 highest total production in India, wheat, are 3, 4, 2, 4 and 2 for ozone, pests and diseases,


22



Formatted: Font: (Default) Times New Roman, 12 pt

Comment [MGE3]: Journal-ready higher resolution figure inserted


Fig. 2: The global effects of five biotic and abiotic stresses on soybean. All data are presented for the 1 x 1° grid squares where the mean production of soybean was > 500 tonnes (0.0005 Tg). (a) Presents the effects of ozone on crop production (thousand tonnes or 0.001 Tg per grid square), and (b) the percentage yield loss due to ozone, averaged for the period 2010-2012. In (d) to (h), the Yield Constraint Score (YCS) is presented per grid square on a scale of 1 to 5, where 5 is the highest level of stress (see Table 1) for ozone, pests and diseases, nutrients, heat stress and aridity, respectively, whilst (c) is the total YCS (YCS_{all}) calculated from the sum of each of these per grid square. The regional impacts are summarised in Table 2 for the five highest producing regions and Table S4 provides all country and regional means.



Formatted: Font: (Default) Times New Roman, 12 pt

Fig. 3: The global effects of five biotic and abiotic stresses on wheat. All data are presented for the $1 \times 1^\circ$ grid squares where the mean production of wheat was > 500 tonnes (0.0005 Tg). (a) Presents the effects of ozone on crop production (thousand tonnes or 0.001 Tg per grid square), and (b) the percentage yield loss due to ozone, averaged for the period 2010-2012. In (d) to (h), the Yield Constraint Score (YCS) is presented per grid square on a scale of 1 to 5, where 5 is the highest level of stress (see Table 1) for ozone, pests and diseases, nutrients, heat stress and aridity, respectively, whilst (c) is the total YCS (YCS_{all}) calculated from the sum of each of these per grid square. The regional impacts are summarised in Table 2 for the five highest producing regions and Table S5 provides all country and regional means.


Comment [MGE4]: Journal-ready higher resolution figure inserted



Formatted: Font: (Default) Times New Roman, 12 pt

Fig. 4: The global effects of five biotic and abiotic stresses on rice. All data are presented for the 1 x 1° grid squares where the mean production of rice was > 500 tonnes (0.0005 Tg). (a) Presents the effects of ozone on crop production (thousand tonnes or 0.001 Tg per grid square), and (b) the percentage yield loss due to ozone, averaged for the period 2010-2012. In (d) to (h), the Yield Constraint Score (YCS) is presented per grid square on a scale of 1 to 5, where 5 is the highest level of stress (see Table 1) for ozone, pests and diseases, nutrients, heat stress and aridity, respectively, whilst (c) is the total YCS (YCS_{all}) calculated from the sum of each of these per grid square. The regional impacts are summarised in Table 2 for the five highest producing regions and Table S6 provides all country and regional means.

Comment [MGE5]: Journal-ready higher resolution figure inserted



Formatted: Font: (Default) Times New Roman, 12 pt, Bold


Fig. 5: The global effects of five biotic and abiotic stresses on maize. All data are presented for the 1 x 1° grid squares where the mean production of maize was > 500 tonnes (0.0005 Tg). (a) Presents the effects of ozone on crop production (thousand tonnes or 0.001 Tg per grid square), and (b) the percentage yield loss due to ozone, averaged for the period 2010-2012. In (d) to (h), the Yield Constraint Score (YCS) is presented per grid square on a scale of 1 to 5, where 5 is the highest level of stress (see Table 1) for ozone, pests and diseases, nutrients, heat stress and aridity, respectively, whilst (c) is the total YCS (YCS_{all}) calculated from the sum of each of these per grid square. The regional impacts are summarised in Table 2 for the five highest producing regions and Table S7 provides all country and regional means.

Comment [MGE6]: Journal-ready higher resolution figure inserted

1
2 nutrients, heat and aridity, respectively (Table S5). For rice, the second most important crop
3 by Tg produced in India, the YCSs for the same five stresses respectively are 2, 5, 1, 2 and 2
4 (Table S6). As the data for the risk of losses due to pests and diseases was only available at
5 the national scale for India, with YCSs of 4 for wheat and 5 for rice, these effects were not
6 included in this spatial analysis, conducted at the 1 x 1 ° scale.

7 For wheat, the highest production is in the adjacent N states of Uttar Pradesh, Madhya
8 Pradesh, Haryana, Rajasthan and Punjab (Fig. 6). Together, these five states account for 85%
9 of Indian wheat production. Predicted percentage yield losses due to ozone are in the range
10 15 – 20 % (mean of 16.4%) in most of the wheat producing areas of Uttar Pradesh, the state
11 with the highest wheat production, resulting in a mean ozone YCS of 3 (Fig. 6 and S5). The
12 mean YCS for ozone was 3 for Haryana, Rajasthan and Punjab and 2 for Madhya Pradesh
13 where ozone uptake is lower (Fig. S5). Although the highest percentage yield losses due to
14 ozone were predicted for states in the far NE of India such as Assam and Manipur, total
15 production losses there were predicted to be minimal as this is not an important wheat
16 growing area. The area of highest ozone impacts on wheat production coincided with the
17 area with the highest YCS for heat stress which covered most of the northern half of the
18 country. Aridity and nutrient YCSs were highest to the W of this region, coinciding with
19 percentage yield losses for ozone predicted to be in the range 5-15% (YCS of 2- 3). For the
20 five highest wheat producing states, the mean YCS for heat stress was 5, with scores for
21 aridity being 2 or 3, and nutrients being 1 – 3 (Fig. 6).

22 Rice growth is much more widely distributed in India than wheat growth, with the highest
23 production being in the N, in part coinciding with wheat growing areas in states such as Uttar
24 Pradesh, and also in E and S states such as West Bengal, West Odisha, Andhra Pradesh and
25 Tamil Nadu (Fig. 6). Together these 5 states produce just below half of India's rice



Formatted: Font: (Default) Times New Roman, 12 pt, Bold

Fig. 6: Yield Constraint Score (YCS) for five constraints on the yield of (a) wheat and (b) rice in the five Indian states with the highest production per crop. The bars represent the mean YCS per $1 \times 1^\circ$ grid square per state on a scale of 1 to 5, where 5 is the highest level of stress (see Table 1), rounded to the nearest integer. Note: The YCS for pests and disease is only available at the National Scale for India (score 4 for wheat and 5 for rice) and is presented here for information.

Comment [MGE7]: Colour figure replaced with black and white figure

production. The highest percentage yield losses for ozone are predicted to be in the range 10 – 15% in the N of the country, including in Uttar Pradesh (mean ozone YCS of 3, Fig. 6 and S6). Lower effects were predicted for Odisha and West Bengal (mean YCS of 2) and the least ozone effects were predicted for rice producing areas in the southern states of Tamil Nadu and Andhra Pradesh (mean YCS of 1), where percentage losses were frequently less than 5%. Heat stress is less of a concern for rice, with a mean YCS of 2 predicted for each of the 5 most important rice producing states. Nutrient stress is predicted to only be important in the far NE states and in isolated grid squares in Rajasthan and along the W coast of India. The mean YCSs for nutrients and aridity for the five highest producing states are either 1 or 2 (Fig. 6).

3.2. Plant traits associated with tolerance of ozone and associated stresses in crops


4 Crop ideotype for ozone and multiple stress tolerance

The derivation of dose-response relationships for 52, 18 and 44 genotypes of soybean, wheat and rice respectively (Fig. 1) has shown that there is clearly scope for the breeding of ozone tolerant varieties, as many varieties had responses that are above the regression line. As part of this study, we identified a number of traits that could contribute to improved ozone tolerance and have summarized these in an ozone-tolerant crop ideotype, including potential trade-offs and synergies for effects of other stresses that can co-occur with ozone (Fig. 7).

Leaf traits for ozone tolerance fall into two categories, the first being processes that limit ozone entry. These include stomatal conductance, and the related trait of water use efficiency (WUE), that reduce ozone uptake while maintaining high rates of photosynthesis. These traits are associated with reduced leaf transpiration and whilst they would be beneficial for water conservation under drought conditions, they may reduce yield and could be potentially deleterious under heat stress by limiting evaporative cooling (Reynolds *et al.*, 2007).

Formatted: Font: Not Bold, Underline

1 Similarly, reduced water uptake associated with lower stomatal conductance has the potential
2 to limit uptake of nutrients such as N from the soil (Zhou *et al.*, 2016). While pathogens are
3 |



4
5 **Fig. 7: An ideotype for an ozone-tolerant resilient crop.** ‘+’ indicates where there would be
6 a benefit for other stresses of improving tolerance to ozone for the trait, whilst ‘-’ indicates a
7 trade-off, and ‘0’ is no effect.

Comment [MGE8]: Colour figure replaced with black and white figure

1 known to have negative effects on leaf gas exchange (Debona *et al.*, 2014), the impact of
2 inherently lower stomatal conductance on disease establishment is less clear although it could
3 be expected that ingress of leaf pathogens that access leaves through the stomatal pores
4 would be reduced.

5 A second category of favourable leaf traits includes antioxidant metabolism and pathways
6 involved in programmed cell death (PCD). Ozone is decomposed into reactive oxygen
7 species (ROS) in the plant apoplast, which either cause direct oxidative damage, or induce
8 signalling cascades similar to a pathogen response, ultimately leading to PCD (Kangasjärvi *et al.*, 2005). Thus, balancing the interplay of redox homeostasis and PCD pathways is essential
9 for the breeding of ozone tolerant crop plants. As a first line of defence against ozone stress,
10 high levels of apoplastic antioxidants such as ascorbate may mitigate ROS formation, a
11 concept that has been confirmed in crop plants such as wheat (Feng *et al.*, 2010) and legumes
12 (Yendrek *et al.*, 2015). Breeding for high levels of antioxidants is also assumed to cause
13 synergies with other types of abiotic stress tolerance, including for drought and heat, both of
14 which are associated with oxidative stress (Gill & Tuteja, 2010). In the case of some nutrient
15 disorders and biotic stresses, functional redox balance rather than high antioxidant levels *per*
16 *se* are considered as important (Munné-Bosch *et al.*, 2013; Suzuki *et al.*, 2012; Wu *et al.*,
17 2017).

18 PCD is an important pathway of pathogen response in plant leaves (Huysmans *et al.*, 2017),
19 which is controlled by the interplay of ROS, signalling cascades and plant hormones
20 (Kangasjärvi *et al.*, 2005). Breeding for ozone tolerance could thus keep plants from inducing
21 PCD despite the presence of apoplastic ROS. This idea is supported by a study in rice, in
22 which the disruption of the pathogen and ozone responsive apoplastic protein *OsORAP1*,
23 which is involved in cell death, led to enhanced ozone tolerance (Ueda *et al.*, 2015). The
24 potential interference of this strategy with pathogen tolerance in crops is obvious, but it is
25

1 currently unclear whether a synergistic or rather antagonistic relationship would occur with
2 different classes of pathogens i.e. biotrophic versus necrotrophic ones (Huymans *et al.*,
3 2017). Implications of PCD in other stress types such as heat (Locato *et al.*, 2008), drought
4 (van Doorn 2011) and nutrient deficiency (Siyiannis *et al.*, 2012) have also been reported but
5 the implications for ozone tolerance breeding remain unclear.

6 Root traits that support ozone stress tolerance would include the capacity to efficiently
7 acquire water and nutrient resources under stress environments (Resource Acquisition
8 Efficiency in Fig. 7). Ozone is known to have a greater negative impact on roots than shoots,
9 resulting in the decline in the root/shoot ratio commonly observed (Fiscus *et al.*, 2005). There
10 is evidence that ozone may have an even greater impact on fine roots that acquire water and
11 nutrients from the soil (Vollsnes *et al.*, 2010). Fine roots are the new frontier of future root
12 research. The framework for describing fine root architecture is being refined (McCormack *et*
13 *al.*, 2015; Zobel, 2016), and new techniques are now available to assess fine root dynamics
14 (e.g. measurements of root diameter, Zobel *et al.*, 2007). The challenge ahead is to define and
15 measure fine root traits that contribute to ozone tolerance, and then determine how these traits
16 affect plant response to other stress factors. Presumably, traits that contribute to robust root
17 systems will be of benefit across a range of abiotic stresses.

18 Traits associated with reproductive organs such as panicles or pods are of primary importance
19 in breeding, although the effects of ozone on these organs may be rather secondary, i.e.
20 caused by foliar responses that limit assimilate acquisition (described above) or effects on
21 flowering and pollen viability (Black *et al.*, 2000). Yield losses due to ozone have been
22 ascribed to various yield components in different crops, including reductions in individual
23 seed weight, reduced spikelet number, enhanced spikelet fertility, and reduced panicle or pod
24 number (Ainsworth, 2008; Feng *et al.*, 2008; Morgan *et al.*, 2003), with associated reductions
25 in harvest index (e.g. for wheat, Pleijel *et al.*, 2014). Maintaining high values in these harvest

fractions despite ozone stress forms an important breeding target, but synergies or trade-offs with other types of stress would be complex and little information is available to date.

Maintaining high crop quality despite ozone stress represents another important breeding goal. Ozone can affect multiple quality traits in seed crops, including protein and starch concentration, as well as visual appearance (Broberg *et al.*, 2015; Wang & Frei, 2011). In many cases, increases in seed protein concentration despite losses in protein yield are observed. This apparent beneficial effect is offset by the negative effects of ozone on seed weight (e.g. for wheat, Broberg *et al.*, 2015). Another quality trait that has been affected in rice by ozone is grain chalkiness, i.e. the formation of milky patches on grains due to inhibited starch loading (Jing *et al.*, 2016). Chalkiness was first described as a typical symptom of heat and drought stress (Wassmann *et al.*, 2009), and lowering plant susceptibility to chalkiness *via* breeding may thus have potential co-benefits with regards to these stresses.

A further category of traits that could be targeted by breeders are phenological characteristics. Plants that have a shorter maturity period by entering earlier into reproductive phases might be more tolerant, as they would receive a lower cumulative ozone dose, and might avoid high ozone episodes occurring late in the cropping season. This principle was confirmed in a study by Ueda *et al.* (2015), in which more than 300 genotypes of rice were screened for ozone response, and yield losses were positively correlated with the number of days to maturity. In general, breeding fast-maturing crop varieties may produce substantial synergies, reducing the impacts of growing seasons characterized by high incidence of other stresses, such as drought, heat, nutrient, or biotic stresses.

4. Discussion

In bringing together these datasets and modelling methods to derive YCSs for five stresses and four key crops, we have conducted the first global assessment of the magnitude of ozone stress in relation to other stresses for four staple crops. We have also derived an ideotype for an ozone- and multi-stress tolerant crop. We provide an extended discussion here that first considers the results presented and then considers potential solutions for increasing crop tolerance of ozone, including crop management and breeding approaches.

4.21 The global scale of ozone impacts on crops relative to impacts of other stresses

Spatial analysis of the impacts of multiple stresses on crop yield

An in depth evaluation of the spatial analysis conducted here is presented in the Supporting Information (T1) and summarised here. The benefits of impacts modelling based on the stomatal uptake of ozone rather than the concentration above the leaf, together with an evaluation of global modelling of POD₃IAM, are discussed by Mills *et al.* (2018a). In the absence of suitable stomatal uptake dose response relationships for soybean, rice and maize, the RS_w method was developed whereby the effects of ozone on these crops was determined from the POD₃IAM response of wheat. We had to assume that the differences in ozone concentration and sensitivity were a greater driver of response than differences in stomatal uptake and are unable to quantify the uncertainty introduced by this assumption. Whilst experimental data in the M7 response functions used in the RS_w method represented the major crop growing regions for soybean, wheat and rice, the function for maize was limited to relatively old data from NAM only. Thus, the data analysis presented here for maize is likely to be the most relevant for effects described in NAM, and is less certain when applied to other maize-growing regions of the world. For each crop, we assumed only one crop growth period per year. Thus, for those crops such as rice where two or three crop growth cycles may occur per year in major growing areas, assessments based on the main growth period will have an added level of uncertainty, then our estimates of total effects on

Formatted: Subscript

1 ~~production are likely to be underestimated.~~ The abiotic and biotic stresses included here were
 2 ~~selected as examples for comparison with ozone effects, with heat stress being chosen as~~
 3 ~~representative of effects of extreme climatic events associated with climate changewarming~~
 4 ~~and.~~ We acknowledge that other stresses such as flooding may also have catastrophic local
 5 ~~effects on yield (e.g. in China, Tao *et al.*, 2017), but have focussed on example stresses for~~
 6 ~~which global data is readily available. Furthermore, global warming impacts on yield could~~
 7 ~~be in a similar range to ozone (e.g. Challinor *et al.*, 2009; Lobell & Asseng, 2017) but have~~
 8 ~~not been considered here.~~ Scores for YCS for pests and diseases may have overestimated
 9 current losses as advances in pesticide usage since the 2002-04 dataset was compiled may
 10 have reduced total impacts. As it was not possible to base all YCSs on percentage yield loss,
 11 uncertainty will have been introduced by comparing effects across stresses. We have
 12 acknowledged this uncertainty by using the YCSs to indicate the location of the largest
 13 effects rather than to quantify the extent of effects. Lastly, YCS_{all} simply summed all YCSs
 14 and provides an indication of where multiple stresses co-occur, without taking into account
 15 any interactions that may occur that might lessen or increase the combined effects on yield.
 16 Taking into account all of these caveats, this study is the first to present ozone impacts on the
 17 global scale together with impacts of other biotic and abiotic stresses, and show spatially
 18 where such stresses are likely to co-occur for four major staple crops.
 19 At the national scale, the countries identified as having the largest potential effects of ozone
 20 (e.g. USA, India and China) match those with the highest monitored ozone concentrations
 21 (Mills *et al.*, ~~2018bsubmitted~~) as well as those predicted using concentration-based
 22 approaches to have the highest potential yield losses (Avnery *et al.*, 2011-a,b; Van Dingenen
 23 *et al.*, 2009). At the sub-national scale, however, there were some differences in areas
 24 predicted to be at risk, where our stomatal uptake modelling method took into account the
 25 modifying effects of climate and soil moisture on ozone uptake rather than simply predicting

Formatted: Font: Italic

1 the largest effects in the areas with the highest ozone concentrations. For example, in India,
2 this study predicts the largest effects on wheat and rice in the northern areas, south of the
3 Himalayas where ozone levels, climatic conditions and irrigation usage promote ozone
4 uptake and subsequent effect. In contrast, an earlier concentration-based study provided little
5 spatial differentiation in effects, predicting widespread and similar effects of ozone in the
6 northern half of India for wheat and across most of India for rice (Van Dingenen *et al.*, 2009).
7 On a global scale, we predict that ozone (mean of 2010 - 2012) reduces soybean yield by
8 12.4%, wheat yield by 7.1%, rice yield by 4.4% and maize yield by 6.1%, adding up to a total
9 of 227 Tg of lost yield. These mean percentage losses are different to those predicted by
10 Avnery *et al.* (2011a) and Van Dingenen *et al.* (2009) using concentration based metrics.
11 Their studies predicted higher losses for wheat (15.4% and 12.3%, respectively) and lower
12 losses for soybean (8.5% and 5.4%, respectively) using AOT40 (Accumulated hourly mean
13 ozone above 40 ppb during daylight hours) for the year 2000.

14 Multivariate analysis of trends in soybean and maize yields in the USA that included a
15 concentration-based ozone metric indicated that the ozone effect is dependent upon
16 temperature and water availability (McGrath *et al.*, 2015). This fits with our earlier
17 conclusion that stomatal uptake-based risk assessment provides a better indication of ozone
18 effects on yield than concentration-based assessments (Mills *et al.*, 2018a). The McGrath *et*
19 *al.* (2015) study indicated a greater sensitivity of maize to ozone than soybean, with maize
20 and soybean yield losses due to ozone over a 31-year period averaging 10% and 5%,
21 respectively. It is possible that our analysis under-estimated the effects of ozone on maize as
22 our analysis was based on experimental data from 1981, 1985, 1991 and 1992. Since newer
23 varieties of wheat and soybean are more sensitive to ozone than older varieties (Biswas *et al.*,
24 2009; Osborne *et al.*, 2016), then newer maize varieties may also be more ozone sensitive
25 leading to larger effects. Partial derivative-linear regression analysis of heat and

1 concentration-based ozone stress impacts on yield data in the USA and Europe indicated
2 similar areas at risk from ozone for maize and soybean to our study, but fewer areas at risk
3 for wheat (Tai & Val Martin, 2017). The latter may reflect that their study omitted soil
4 moisture as a confounding factor, which our earlier modelling study indicated is a particularly
5 important factor in modifying ozone uptake (Mills *et al.*, 2018a). These two statistical
6 studies have confirmed that factors other than ozone concentration need to be taken into
7 account in analysing ozone effects on yield and have drawn attention to potential co-
8 occurrence of heat and ozone stress effects on crop yield.

9 It is clear from our analysis that yield effects due to ozone are within the range of concern for
10 other biotic and abiotic stresses. For example, extreme heat was estimated to have reduced
11 national cereal production by 9-10% (1964 - 2007), with later droughts reducing yields by
12 more than earlier droughts (13.7% for 1985 – 2007 compared to 6.7% for 1964 – 1984, Lesk
13 *et al.*, 2016). If we applied the percentage yield loss ranges used for ozone in YCS (Table 1)
14 to these drought-induced yield losses, the YCS would be 2 or 3 depending on the time period
15 used in the analysis. The YCSs for ozone are mainly in the same range as those predicted for
16 wheat, maize and rice for global impacts of heat stress and aridity (scores 2-3, with rice-heat
17 having a YCS of 1). Across all four crops, the areas predicted to be at the greatest risk of
18 ozone effects on yield are predicted to be: SE NAM, S EUR, N SAS and E EAS, with parts of
19 SAM also predicted to be at risk of yield loss for soybean. In some of these areas, ozone
20 effects are predicted in areas also at risk from heat stress and to a lesser extent aridity, whilst
21 co-occurrence with nutrient stress depended on the crop and tended to be most common in
22 parts of EAS and SAM. Potential impacts on yield due to pests and diseases were predicted
23 to be relatively high in many areas of the world, particularly in those at risk from ozone
24 impacts in SAS and EAS.

Ozone impacts are predicted in areas where the largest gaps occur between actual and estimated potential yield, such as parts of SAM, SSA, EAS, SEA and SAS (Neumann *et al.*, 2010). Here, yield gaps are already known to be widened by limitations in nutrient and/or irrigation availability (Mueller *et al.*, 2012) and may be further widened by negative effects of ozone pollution. Indeed, in the same regions there has been a plateauing or decrease in the rate of yield increase in recent decades (Grassini *et al.*, 2013; Ray *et al.*, 2012). We suggest that ozone pollution could be contributing to this stagnation, and suggest below how crop tolerance of the pollutant could be improved by breeding or management (Section 4.2.3).

India ~~was~~ ~~was~~ selected as a case study, as our analysis indicated that ozone pollution may be a particular problem in this country, adding to the existing multi-stress constraints on crop yield (Jaswal, 2014). National mean yield losses due to ozone were predicted to be 15.8% (soybean), 12.6% (wheat), 6.2% (rice) and 7.5% (maize) amounting to 12.6 Tg of lost yield. For wheat, our predicted mean yield loss in Uttar Pradesh of 16% was comparable to a mean 17% yield benefit from reducing the ambient ozone from 46 to 5 ppb (M7) by air filtration in field studies conducted from 2004 to 2008 at Varanasi in Uttar Pradesh (Rai *et al.*, 2007, Sarkar *et al.*, 2010). Similarly, at a field site in Haryana, reduction in the M7 by filtration from 37 ppb to 6 ppb, resulted in a 16% yield benefit for wheat (Bhatia *et al.*, 2011), which was similar to our state mean of a 15% yield reduction due to ozone.

Wheat yield losses were predicted to be highest in this study in the same regions of India as those predicted by Tang *et al.* (2013) in the first stomatal uptake-based risk assessment for the country. Our analysis, taking into account the added effects of soil moisture and irrigation usage, extended the region of highest ozone effects across the Indo-Gangetic Plain and including Uttar Pradesh and Bihar. Together with Haryana and Punjab, these states are considered to have the highest reductions in yield due to the combined effects of climate change and air pollution, with reductions as high as 50% being predicted in one

1 concentration-based study (Burney & Ramanathan, 2013). Our multi-stress analysis
2 confirmed that heat stress is ~~potentially~~ particularly important in this region (Lobell *et al.*,
3 2012). Site-specific analysis of the effects of future increases in temperature in 2030-2040,
4 indicated that heat stress is likely to continue to reduce yields in the Indo-Gangetic Plain,
5 especially under climate change (Asseng *et al.*, 2017). Given that from a food security
6 perspective, it is crucial to reduce yield gaps in India, reducing ozone pollution and/or its
7 effects could potentially provide beneficial additional yield in future climates.

8 In considering these comparisons, we are aware that in reality ozone will interact with the
9 other stresses considered and integrated responses in growth and yield will occur. These
10 interactions are generally thought to be determined by factors that might affect gas exchange
11 or metabolic responses to stress. For example, limited water stress may reduce ozone uptake
12 but as water stress becomes more severe, any protection afforded by reduced ozone uptake
13 may be outweighed by drought-induced yield reductions. Additionally, these stresses are
14 thought to impart similar defence mechanisms (Huymans *et al.*, 2017; Kangasjärvi *et al.*,
15 2005; Locato *et al.*, 2008). Whether multiple stresses induce additive or synergistic metabolic
16 responses is open to question and we do not yet have the understanding or tools to be able to
17 quantify these interactions. Nevertheless, through providing a first global assessment of
18 where these stresses co-occur we have identified which stresses are most important across
19 different global regions. This will help other researchers to identify threats and target future
20 research needs to improve our understanding of responses to multiple stress conditions.

21 **4.2 Options for reducing ozone impacts on crops**

22 The analysis presented here has clearly shown that ozone impacts on yield are occurring in
23 many areas of the world for four staple crops, and that in some regions the YCSs for ozone
24 are as high or higher than for other biotic and abiotic stresses. Whilst these results highlight

the ozone problem, we offer here some possible options for reducing ozone effects on crops that might help in closing the ozone yield gap.

4.2.1 Global effort to reduce ozone precursor emissions

Formatted: Underline

The most obvious way of closing the ozone yield gap for crops is to substantially lower the anthropogenic emissions that lead to ozone pollution. Since ozone is a transboundary air pollutant – impacts of emissions in one country can impact on crops grown in countries many 100s and even 1000s of km away – efforts to reduce ozone need to be taken at both local and global scales. One study, ~~using based on ozone concentration-based metrics~~, indicated that 100% reductions in anthropogenic precursor emissions from NAM would reduce global yield losses due to ozone for the four crops in our study by between ca. 5% (rice) to ca. 80% (soybean), whilst a complete cut in precursor emissions from SEA would reduce global yield losses by between ca. 20% (soybean) and ca. 95% (rice) (Holloway *et al.*, 2012). Whilst such dramatic cuts in ozone precursor emissions are highly unlikely for the foreseeable future, progress has been made in EUR and NAM, with emission cuts of ca. 40% for major ozone precursors such as NO_x, VOC and CO being made between 1990 and 2013 (Maas & Grennfelt, 2016). These cuts have been associated with significant decreasing trends in the concentration-based metric AOT40 at 26% and 11% of monitoring sites in wheat growing areas of NAM and EUR, respectively over the period 1995-2014 (Mills *et al.*, 2018b), although the dominant trend for EUR remains “no change”. Over the same time period, increases in precursor emissions of 20-30% in other areas of the world, including by 50% in India and China have led to increases in ozone concentration in these regions (Maas & Grennfelt, 2016). For example, there has been a significant increase in ozone concentration at nearly 50% of wheat growing monitoring sites in EAS, with average annual increases in AOT40 at these sites being in the range 300 – 700 ppb h y⁻¹ over the period 1995-2014 (Mills *et al.*, 2018b).

Formatted: Font: Italic

Formatted: Not Highlight

Formatted: Superscript

Formatted: Font: Italic

Our modelling results suggest predicts that even with declining emissions in NAM, current yield losses due to ozone are in the range 5.3% (rice) to 15.5% (soybean), whilst for EAS with rising emissions, current yield losses are in the range 7.9% (rice) to 19.1% (soybean). With ozone concentrations predicted to continue to rise in EAS and SEA for at least the next 2-3 decades even with the most optimistic scenarios (Wild et al., 2012), and as these two regions are predicted to produce 80% of all global ozone precursor emissions by 2050 (Maas & Grennfelt, 2016), there would be considerable benefit for crop yield in the implementation of a concerted effort to reduce precursor emissions in these rapidly developing regions. Actions to reduce ozone are already being considered in some countries. For example, in China, three approaches are being introduced to reduce ozone concentrations: enforcing European standard V for diesel vehicle emissions; encouraging widespread use of electric vehicles; and discouraging private car use by improving public transport (Feng *et al.*, 2015). Continued effort to reduce ozone is also needed in developed regions such as NAM and EUR as models predict that whilst efforts to reduce peak concentrations have been partially successful in reducing ozone concentrations in recent decades, a stabilisation in ozone concentrations in the next decade or two is likely to be followed by further rises in global background ozone concentration by 2050, primarily driven by increasing CH₄ emissions (Maas & Grennfelt, 2016).

Whilst reducing global ambient ozone concentrations remains a crucial long-term goal for reducing the ozone yield gap, approaches described below based on crop management and breeding are more likely to provide shorter-term solutions, with some having potential for implementation in the near future.

4.2.2 Exploiting existing varietal differences in ozone sensitivity

Formatted: Font: Italic

Formatted: Subscript

1 Whilst the analysis presented here has confirmed that intraspecific variation in ozone
 2 sensitivity is clearly present for wheat, rice and soybean in experiments conducted over the
 3 last 30 - 40 years (Fig. 1, Table S3), of larger importance in the context of closing the ozone
 4 yield gap is the potential for selecting ozone tolerance amongst currently grown varieties. To
 5 assess this, ideally, varieties should be exposed to ozone under the same environmental
 6 conditions, allowing for realistic comparisons of effects on yield and assessments of variety
 7 by ozone interactions. Unfortunately, relatively few such experiments have been conducted
 8 with two or more varieties in the last decade. Those recent studies showing significant
 9 variety by ozone interactions, indicating scope for selecting the more ozone tolerant variety,
 10 include examples from SAS and EAS for rice (Akthar *et al.*, 2010; Shi *et al.*, 2009) and
 11 wheat (Feng *et al.*, 2010, 2016; Singh *et al.*, 2017; Zhu *et al.*, 2011;), NAM for soybean
 12 (Betzberger *et al.*, 2010; Jiang *et al.*, 2018) and maize (Yendrek *et al.*, 2017); and EUR for
 13 wheat (Harmens *et al.*, 2018). Further support for the potential benefits of selecting tolerant
 14 varieties is also provided by comparisons of yield in filtered air versus non-filtered air
 15 (Osborne *et al.*, 2016; Pleijel *et al.*, 2018). For example, in recent studies, reductions of
 16 ambient ozone concentration by filtration significantly increased the yield of ozone-sensitive
 17 soybean cultivars (PUSA 9712, PUSA 9814) by over 40% (Singh & Agrawal, 2011), rice
 18 cultivar Kirara 397 by over 20% (Frei *et al.*, 2012) and wheat cv PBW 343 by 18-20%
 19 (Tomer *et al.*, 2015).

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

21 A modelling study has been conducted to highlight the potential for avoiding production loss
 22 in global wheat, maize and soybean by selecting crop varieties with lower than average
 23 sensitivity to ozone (Avnery *et al.*, 2013). The variation in sensitivity among varieties was
 24 based on the experimental evidence from the large-scale US National Crop Loss Assessment
 25 Network (NCLAN) field studies conducted mainly during the 1980s (Heagle, 1989; Heck *et*

al., 2013; Heck, 1989). Using a concentration-based method, the study showed that choosing crop varieties with ozone tolerance could improve global crop production by over 140 Tg in 2030, equivalent to a 12% increase. Although the older North American varieties may not represent current global variation, and some of the 1980s varieties are no longer used, the approach of Avnery *et al.* (2013) could be extended by conducting new screening experiments with a regional focus to inform farmer choice, modelling and breeding programmes and by using a stomatal uptake based modelling approach.

4.2.3 Breeding new varieties with multiple stress tolerance, including ~~of~~ ozone

The heterogeneity in variety response to ozone for soybean, wheat and rice (Fig. 1) has clearly shown ~~that there is clearly the~~ scope for breeding ~~of~~ ozone-tolerant varieties, and an ideotype for an ozone tolerant crop has been defined here (Fig.7). Ideally, the improved ozone response must not compromise the yield potential or other required agronomic characteristics (e.g. resistance to diseases, shattering, and lodging). Since this study has also shown that ozone stress can typically co-occur with stress caused by heat, pests and diseases, and to a lesser extent, aridity and nutrients, the breeding for ozone tolerance traits may cause potential synergies or trade-offs that also need to be considered (Fig. 7). Candidate traits for ozone tolerance were described in Section 3.42.

Traditional breeding approaches such as pedigree selection require extensive screening of a large number of plants in multiple locations over extended periods of time (Frei, 2015). Whilst feasible, experimentally maintaining designated ozone concentrations on a sufficiently large scale required for breeding (e.g. in large scale FACE (free air concentration exposure) experiments) seems economically unviable. Therefore, molecular breeding approaches such as marker assisted selection (MAS) appear to be more promising. Phenotypic variation in traits associated with ozone tolerance can be evaluated in smaller-scale controlled ozone

1 fumigation experiments and linked to genetic markers using mapping approaches, including
2 bi-parental quantitative trait locus (QTL) mapping (Frei *et al.*, 2008) and Genome-Wide
3 Association Study (GWAS, Ueda *et al.*, 2015). Theoretically, chromosomal fragments
4 associated with ozone tolerance traits can then be introgressed into recipient varieties using
5 marker assisted backcrossing without the need for large-scale fumigation experiments.

6 Although no large-scale marker-assisted breeding programs for ozone tolerance in crops have
7 been conducted to date, proof of concept has been shown for ozone tolerant rice breeding
8 lines carrying QTL for ozone tolerance (Chen *et al.*, 2011; Frei *et al.*, 2008, 2010) that have a
9 superior performance to the recipient varieties in terms of yield components (Wang *et al.*,
10 2014) and grain quality (Jing *et al.*, 2016). This example should encourage further breeding
11 efforts in rice and other crop species, specifically targeting widely grown mega-varieties of
12 crops grown in ozone-affected parts in the world. As an alternative strategy, traits
13 contributing to ozone tolerance could be incorporated into existing crop varieties through
14 genetic engineering. For example, rice and barley varieties engineered to contain enhanced
15 levels of ascorbate have been engineered and showed enhanced tolerance to a variety of
16 environmental stresses (Ali *et al.*, 2018).

17 Physiological trait modelling could also be used to understand how different traits intended to
18 confer tolerance for ozone, might influence crop physiology, growth and yield response under
19 a range of environmental conditions and stresses.

20 4.2.4 Reducing ozone uptake by strategic limitation of irrigation application

21 Ozone impacts on crops could be reduced by partial stomatal closure induced by reduced
22 irrigation, which could also save water use for irrigated crop production. In the rice growing
23 countries, in response to the increasing water demands by other sectors than agriculture,
24 alternate wetting and drying irrigation (AWD) has become popular in an attempt to reduce

1 water usage and methane emissions (Bouman *et al.*, 2007; Carrijo *et al.*, 2017). This
2 approach could also potentially be exploited to reduce ozone impacts on rice or other crops.
3 A comparison of two studies conducted about 30 km apart in the same city of China suggests
4 such a possibility. In Zhang *et al.* (2009), AWD with moderate water stress increased the
5 growth and yield of rice while reducing stomatal conductance compared to continuously
6 flooded crops, mostly resulting from a greater number of rice grains per panicle under AWD.
7 Interestingly, at a nearby site, elevated ozone reduced rice yield arising from a decrease in the
8 number of grains per panicle in two of the four varieties tested (Shi *et al.*, 2009). This
9 suggests that reduced ozone uptake could be an additional and unintended benefit of AWD
10 for farmers. The potential benefits of the AWD approach require further study.

11 | 4.2.5 Fertilizer application to compensate for crop yield losses

12 Crop loss from ozone exposure could potentially be counteracted by increasing the fertilizer
13 application rate (Cardoso-Vilhena & Barnes, 2001; Chen *et al.*, 2011). However, in addition
14 to the cost of fertilizer, recent analysis has indicated that this mitigation approach may be
15 associated with an aggravation of other environmental problems. It has been shown that the
16 nitrogen/protein yield of wheat is reduced by ozone at a certain level of nitrogen application
17 and this applies also to other nutrients like phosphorus and potassium (Broberg *et al.*, 2015;
18 2017). This means that the fraction of nitrogen applied which does not end up in the grain
19 could enhance other environmental problems (Di & Cameron 2002; Mosier *et al.* 1998) such
20 as nitrate leaching, conversion of fertilizer to N₂, emissions of N₂O and even NO, which
21 promotes further ozone formation, as shown for pasture (Sánchez-Martín *et al.*, 2017).
22 Adding nitrogen fertiliser to compensate for reductions in yield may also inadvertently
23 increase the stomatal conductance of leaves of crop plants thereby increasing ozone uptake
24 and subsequent damage (Mills *et al.*, 2016).

25 | 4.2.6 Chemical protection against ozone damage

1 There is scope for investigating the benefits of chemical protection against ozone damage.
2 The most successful antiozonant applied so far has been ethylenediurea (N-[2-(2-oxo-1-
3 imidazolidinyl)ethyl]-N'-phenylurea), abbreviated to EDU, first described by Carnahan *et al.*
4 (1978). This chemical is usually applied as a foliar spray or soil drench, and has been used
5 extensively in experiments and biomonitoring programmes to reduce the effects of ozone
6 pollution, including preventing visible ozone injury on the leaves and growth and yield
7 reductions (Agathokleous *et al.*, 2016; Feng *et al.*, 2010; Jiang *et al.*, 2018; Manning *et al.*,
8 2011; Pandey *et al.*, 2015; Rai *et al.*, 2015). A meta-analysis suggested that the antiozonant
9 activity of EDU is biochemical rather than biophysical (Feng *et al.*, 2010). Recent results
10 showed that EDU has no negative effects on plants at low O₃ concentration, but increases the
11 crop yield at high O₃ concentration (Ashrafuzzaman *et al.*, 2017). Whilst EDU has not yet
12 been evaluated for application at field scale, concerns have been raised about potential
13 toxicity to aquatic plants (Agathokleous *et al.*, 2016) and more research is needed to
14 determine if this chemical could be extensively used.

15 Other chemical protectants against ozone could be developed from a knowledge of plant
16 hormonal control of stomatal functioning and stress perception (Wilkinson *et al.*, 2011), and
17 could potentially provide multi-stress tolerance such as combined tolerance of ozone, heat
18 and drought stress. All three of these stresses induce synthesis of the crop stress hormone,
19 ethylene, and chemicals that inhibit ethylene perception such as 1-MCP (1-
20 methylcyclopropene) have the potential to reduce their effects (Wilkinson & Davies, 2010;
21 Wagg, 2012). Anti-transpirants that reduce stomatal aperture could also reduce ozone
22 effects by reducing ozone uptake in some species. However, there is a growing body of
23 knowledge that chronic exposure to ozone reduces the ability of stomata to respond to
24 abscisic acid under drought conditions, potentially leading to more rather than less ozone
25 uptake (Mills *et al.*, 2016; Wilkinson and Davies, 2009, 2010). An alternative chemical

protection approach has also been explored experimentally. Di-1-*p*-methene, a natural terpenic polymer derived from the resin of pine trees that mimics isoprene emissions from plants, has been shown to reduce visible injury in Pinto beans after exposure to 150 ppb of ozone for 4h (Francini *et al.*, 2010).

So far, chemical protection has only been explored at the experimental scale. Given the growing evidence presented here and elsewhere of the negative effects of the pollutant at the global scale, there is considerable scope for developing a chemical protectant against ozone damage, especially if it provides cross-tolerance against other co-occurring stresses.

5. Conclusions

This global-scale study shows that ozone is a very important stress, limiting yields of key crops and comparing in importance with other key stresses. For example in India, where food security concerns are particularly pressing, the mean YCS for effects of ozone on wheat of 3 falls in between those for nutrients and aridity (score 2) and for pests and diseases and heat stress (score 4). Globally, we show that the largest effects of ozone are often in areas already challenged by other stresses such as pests and diseases and heat, particularly in EAS, SAS and SEA. The global mean ozone yield gaps of 4.4 – 12.4 % identified here add up to 227 Tg of lost yield for soybean, wheat, rice and maize. We speculate that, ozone could at least partially, account for the unexplained yield gaps and stagnation in yield improvement seen in many areas of the world in recent years. Thus, international effort to reduce ozone pollution on a global scale would bring clear benefits for agriculture as well as for other types of vegetation, health, materials and climate change (Simpson *et al.*, 2014). However, it is likely to take many decades to achieve the required emission reductions, which is the only long-term solution for reducing the problems caused by tropospheric ozone. Meanwhile, the

global population is expected to grow significantly, which together with increasing real income levels, will see increasing demands placed on food production (Tilman *et al.*, 2011). Several interim solutions for closing the ozone yield gap have been outlined in this paper. These include: testing of current varieties for ozone sensitivity and selection of the most tolerant; crop breeding for multiple stress tolerance, including ozone; implementation of protective watering regimes such as AWD; and the development of chemical protection against ozone damage. Given the severity of ozone effects on staple food crops in areas of the world that are also challenged by other stresses, we recommend increased attention to the benefits that could be gained from taking mitigating action to reduce the ozone yield gap.

Acknowledgements

We thank the Natural Environment Research Council (NERC projects NEC05574 and NEC06476) for financial support and the Adlerbertska Foundation for supporting this study by funding Gina Mills' guest professorship at the University of Gothenburg (NERC project NEC05831). The Adlerbertska Foundation are also thanked for funding a workshop on the theme of this paper, attended by the authors, as part of the Guest Professorship funding for Gina Mills. This work has been partially funded by EMEP under UNECE, and the EU project ECLAIRE (project no. 282910). Computer time for EMEP model runs was supported by the Research Council of Norway through the NOTUR project EMEP (NN2890K) for CPU. We also wish to thank Edmar Teixeira for providing advice on calculating a heat stress index.

Data Accessibility Statement:

National-scale data are provided in the Supporting Information. Grid square values for ozone metrics for the LRTAP region can be downloaded from http://www.emep.int/mscw/mscw_data.html.

Statement on Competing Interests: none

The authors declare no competing interests.

References

Ainsworth EA (2008) Rice production in a changing climate: a meta-analysis of responses to elevated carbon dioxide and elevated ozone concentration. *Global Change Biology* 14, 1642-1650.

Ainsworth, EA (2016) Understanding and improving global crop response to ozone Pollution. *The Plant Journal* 90, 886–897.

Agathokleous E, Mouzaki-Paxinou A-C, Saitanis C *et al.* (2016) The first toxicological study of the antiozonant and research tool ethylene diurea (EDU) using a *Lemna minor* L. bioassay: Hints to its mode of action. *Environmental Pollution* 213, 996-1006.

Akhtar N, Yamaguchi M, Inada H *et al.* (2010) Effects of ozone on growth, yield and leaf gas exchange rates of four Bangladeshi cultivars of rice (*Oryza sativa* L.). *Environmental Pollution* 158, 2970-2976.

Ali B, Pantha S, Acharya R *et al.* Abiotic stress tolerance of transgenic cereal crops with enhanced ascorbate biosynthesis. *Plant Biotechnology Journal*, submitted.

- 1 Atlin GN, Cairns JE, Das B (2017) Rapid breeding and varietal replacement are critical to
2 adaptation of cropping systems in the developing world to climate change. *Global*
3 *Food Security* 12, 31-37.
- 4 Ashrafuzzaman M, Farzana AL, Holtkamp F *et al.* (2017) Diagnosing ozone stress and
5 differential tolerance in rice (*Oryza sativa* L.) with ethylenediurea (EDU).
6 *Environmental Pollution* 230, 339-350.
- 7 Asseng S, Cammarano D, Basso B (2017). Hot spots of wheat yield decline with rising
8 temperatures. *Global Change Biology* 23, 2464–2472.
- 9 Avnery S, Mauzerall DL, Liu J. & Horowitz LW (2011a) Global crop yield reductions due to
10 surface ozone exposure: 1. Year 2000 crop production losses and economic damage.
11 *Atmospheric Environment* 45, 2284-2296.
- 12 Avnery S, Mauzerall DL, Liu J. & Horowitz LW (2011b). Global crop yield reductions due
13 to surface ozone exposure: 2. Year 2030 potential crop production losses and
14 economic damage under two scenarios of O₃ pollution. *Atmospheric Environment* 45,
15 2297-2309.
- 16 Avnery S, Mauzerall DL, Fiore AM (2013) Increasing global agricultural production by
17 reducing ozone damages via methane emission controls and ozone-resistant cultivar
18 selection. *Global Change Biology* 19, 1285-1299.
- 19 Betzelberger AM, Gillespie KM, McGrath JM *et al.* (2010) Effects of chronic elevated ozone
20 concentration on antioxidant capacity, photosynthesis and seed yield of 10 soybean
21 cultivars. *Plant Cell and Environment* 33, 1569-1581.
- 22 Bhatia A, Ghosh A, Kumar V *et al.* (2011) Effect of elevated tropospheric ozone on methane
23 and nitrous oxide emission from rice soil in north India. *Agriculture, Ecosystems and*
24 *Environment* 144, 21–28.

Formatted: German (Germany)

- 1 Biswas DK, Xu H, Yang JC *et al.* (2009) Impacts of methods and sites of plant breeding on
2 ozone sensitivity in winter wheat cultivars. *Agriculture, Ecosystems & Environment*
3 134, 168-177.
- 4 Black VJ, Black CR, Roberts JA, Stewart CA (2000) Tansley Review No. 115 Impact of
5 ozone on the reproductive development of plants. *New Phytologist* 147, 421-447.
- 6 Bouman BAM, Lampayan RM, Tuong TP (2007) Water management in irrigated rice.
7 Coping with water scarcity. Manila, Philippines: International Rice Research Institute.
8 p 53
- 9 Broberg MC, Feng Z, Xin Y, Pleijel H (2015) Ozone effects on wheat grain quality - A
10 summary. *Environmental Pollution* 197, 203-213.
- 11 Broberg MC, Uddling J, Mills G, Pleijel H (2017) Fertilizer efficiency in wheat is reduced by
12 ozone pollution. *Science of the Total Environment* 607-608, 876-880.
- 13 Broberg MC, Feng Z, Xin Y, Pleijel H (2015) Ozone effects on wheat grain quality – a
14 summary. *Environmental Pollution* 197, 203-213.
- 15 Burkey KO, Carter TE (2009) Foliar resistance to ozone injury in the genetic base of U.S.
16 and Canadian soybean and prediction of resistance in descendent cultivars using
17 coefficient of parentage. *Field Crops Research* 111, 207-217.
- 18 Burkey KO, Miller JE, Fiscus EL (2005) Assessment of ambient ozone effects on vegetation
19 using snap bean as a bioindicator species. *Journal of Environmental Quality* 34,
20 1081–1086.
- 21 Burney J & Ramanathan V (2014) Recent climate and air pollution impacts on Indian
22 agriculture. *Proceedings of the National Academy of Sciences* 111, 16319-16324.
- 23 CABI (2005) Crop Protection Compendium, 2005 edition. Wallingford, UK.

- 1 Cardoso-Vilhena J, Barnes J (2001) Does nitrogen supply affect the response of wheat
2 (*Triticum aestivum* cv. Hanno) to the combination of elevated CO₂ and O₃? *Journal*
3 *of Experimental Botany* 52, 1901-1911.
- 4 Carnahan JE, Jenner EL, Wat EKW (1978) Prevention of ozone injury to plants by a new
5 protectant chemical. *Phytopathology* 68, 1225-1229.
- 6 Carrijo DR, Lundy ME, Linquist BA (2017) Rice yields and water use under alternate
7 wetting and drying irrigation: A meta-analysis. *Field Crops Research* 203, 173-180.
- 8 Challinor AJ, Ewert F, Arnold S *et al.* (2009) Crops and climate change: progress, trends, and
9 challenges in simulating impacts and informing adaptation, *Journal of Experimental*
10 *Botany*, 60, 2775–2789.
- 11 Challinor AJ, Wheeler TR, Craufurd PQ, Slingo JM (2005) Simulation of the impact of high
12 temperature stress on annual crop yields. *Agricultural and Forest Meteorology*,
13 135(1-4), 180-189.
- 14 Chang KL, Petropavlovskikh I, Cooper OR *et al.* (2017) Regional trend analysis of surface
15 ozone observations from monitoring networks in eastern North America, Europe and
16 East Asia, *Elementa: Science of the Anthropocene*. doi:
17 <http://doi.org/10.1525/elementa.243>
- 18 Chen CP, Frei M, Wissuwa M (2011a) The OzT8 locus in rice protects leaf carbon
19 assimilation rate and photosynthetic capacity under ozone stress. *Plant Cell and*
20 *Environment* 34, 1141-1149.
- 21 Chen J, Zeng Q, Zhu JG *et al.* (2011b) Nitrogen supply mitigates the effects of elevated [O₃]
22 on photosynthesis and yield in wheat. *Chinese Journal of Plant Ecology* 35, 523–530.
- 23 CLRTAP (2017) Chapter 3 "Mapping Critical Levels for Vegetation". LRTAP Convention
24 Modelling and Mapping Manual. <http://icpvegetation.ceh.ac.uk/>

- 1 Debona D, Rodrigues FÁ, Rios JA *et al.* (2014) Limitations to photosynthesis in leaves of
2 wheat plants infected by *Pyricularia oryzae*. *Phytopathology* 104, 34-39.
- 3 Dentener F, Guizzardi D. 2013. TDHTAP Work Package 2.1. Common set of regions.
4 Available at: [http://iek8wikis.iek.fz-](http://iek8wikis.iek.fz-juelich.de/HTAPWiki/WP2.1?action=AttachFile&do=view&target=_region_definition_document_v4.docx)
5 [juelich.de/HTAPWiki/WP2.1?action=AttachFile&do=view&target=_region_definition](http://iek8wikis.iek.fz-juelich.de/HTAPWiki/WP2.1?action=AttachFile&do=view&target=_region_definition_document_v4.docx)
6 [n_document_v4.docx](http://iek8wikis.iek.fz-juelich.de/HTAPWiki/WP2.1?action=AttachFile&do=view&target=_region_definition_document_v4.docx) Accessed 11 August, 2016.
- 7 Deryng D, Conway D, Ramankutty N *et al.* (2014) Global crop yield response to extreme
8 heat stress under multiple climate change futures. *Environmental Research Letters*
9 9(3), 034011.
- 10 Di HJ, Cameron KC (2002) Nitrate leaching in temperate agroecosystems: sources, factors
11 and mitigating strategies. *Nutrient Cycling in Agroecosystems* 64, 237 – 256.
- 12 Driedonks N, Rieu I, Vriezen, WH (2016) Breeding for plant heat tolerance at vegetative and
13 reproductive stages. *Plant Reproduction* 29, 67 – 79.
- 14 ECMWF. European Centre for Medium-Range Weather Forecasts European Centre for
15 Medium-Range Weather Forecasts. <http://www.ecmwf.int/> (accessed 2017).
- 16 Emberson LD, Ashmore MR, Simpson D *et al.* (2001) Modelling and Mapping Ozone
17 Deposition in Europe. *Water, Air, and Soil Pollution* 130, 577-582.
- 18 Emberson LD, Pleijel H, Ainsworth EA *et al.* (2018) Ozone effects on crops and
19 consideration in crop models. *Accepted by European Journal of Agronomy*.
- 20 FAO <http://www.fao.org/land-water/databases-and-software/crop-information/soybean/en/>
21 (accessed 2017).
- 22 FAOSTAT. Food and agriculture division of the United Nations statistics division.
23 <http://faostat3.fao.org/home/E> (accessed 2017).

- 1 Farooq M, Gogoi N, Barthakur S *et al.* (2017) Drought Stress in Grain Legumes during
 2 Reproduction and Grain Filling. *Journal of Agronomy and Crop Science* 203, 81 –
 3 102.
- 4 Feng ZZ, Kobayashi K (2009) Assessing the impacts of current and future concentrations of
 5 surface ozone on crop yield with meta-analysis. *Atmospheric Environment* 43, 1510-
 6 1519.
- 7 Feng ZZ, Kobayashi K, Ainsworth EA (2008) Impact of elevated ozone concentration on
 8 growth, physiology, and yield of wheat (*Triticum aestivum* L.): a meta-analysis.
 9 *Global Change Biology* 14, 2696-2708.
- 10 Feng Z, Liu X, Zhang F (2015) Air pollution effects on food security in China: Taking ozone
 11 as an example. *Frontiers of Agricultural Science and Engineering* 2, 152-158.
- 12 Feng ZZ, Pang J, Nouchi I *et al.* (2010a) Apoplastic ascorbate contributes to the differential
 13 ozone sensitivity in two varieties of winter wheat under fully open-air field
 14 conditions. *Environmental Pollution* 158, 3539-3545.
- 15 Feng Z, Wang S, Szantoi Z *et al.* (2010b) Protection of plants from ambient ozone by
 16 applications of ethylenediurea (EDU): a meta-analytic review. *Environmental*
 17 *Pollution* 158, 3236-3242.
- 18 Feng Z, Wang L, Pleijel H *et al.* (2016) Differential effects of ozone on photosynthesis of
 19 winter wheat among cultivars depend on antioxidative enzymes rather than stomatal
 20 conductance. *Science of the Total Environment* 572, 404–411.
- 21 Fischer G, Nachtergaele FO, Prieler S *et al.* (2012) Global Agro-ecological Zones
 22 Assessment for Agriculture (GAEZ v3.0). Model Documentation. IIASA, Laxenburg,
 23 Austria and FAO, Rome, Italy.

Formatted: Indent: Left: 0", Hanging: 0.5", Space Before: 6 pt, Line spacing:

Formatted: Font: (Default) Times New Roman, 12 pt, Italic

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Italic

Formatted: Font: (Default) Times New Roman, 12 pt, Italic

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Italic

1 Fiscus EL, Booker FL, Burkey KO (2005) Crop responses to ozone: uptake, modes of action,
2 carbon assimilation and partitioning. *Plant Cell & Environment* 28, 997-1011.

3 Francini A, Lorenzini G, Nali C (2011) The Antitranspirant Di1- p -menthene, a Potential
4 Chemical Protectant of Ozone Damage to Plants. *Water Air and Soil Pollution* 219,
5 459-472.

6 Frei M (2015) Breeding of ozone resistant rice: Relevance, approaches and challenges.
7 *Environmental Pollution* 197, 144-155.

8 Frei, M, Kohno, Y, Tietze S *et al.* (2012) The response of rice grain quality to ozone
9 exposure during growth depends on ozone level and genotype. *Environmental*
10 *Pollution* 163, 199-206.

11 Frei M, Tanaka JP, Chen CP, Wissuwa M (2010) Mechanisms of ozone tolerance in rice:
12 characterization of two QTLs affecting leaf bronzing by gene expression profiling and
13 biochemical analyses. *Journal of Experimental Botany* 61, 1405-1417.

14 Frei M, Tanaka JP, Wissuwa M (2008) Genotypic variation in tolerance to elevated ozone in
15 rice: dissection of distinct genetic factors linked to tolerance mechanisms. *Journal of*
16 *Experimental Botany* 59, 3741-3752.

17 Furher J (1994). The critical level for ozone to protect agricultural crops – An assessment of
18 data from European open-top chamber experiments. *Fuhrer J. & Achermann, B.,*
19 *(Eds). Critical Levels for Ozone. UNECE Workshop Report., 42 – 57.*

20 GAEZ. Global Agro-ecological Zones v3.0. <http://www.gaez.iiasa.ac.at/> (accessed 2017).

21 Gastal F, Lemaire G (2002) N uptake and distribution in crops: an agronomical and
22 ecophysiological perspective. *Journal of Experimental Botany* 53(370), 789-799.

Formatted: Font: Not Highlight

Formatted: Font: Not Highlight

Formatted: Font: Italic

Formatted: Font: Not Highlight

Formatted: Font: Italic, Not Highlight

Formatted: Font: Italic

- 1 Gill SS, Tuteja N (2010) Reactive oxygen species and antioxidant machinery in abiotic stress
2 tolerance in crop plants. *Plant Physiology and Biochemistry* 48, 909-930.
- 3 Gilliham M, Able JA, Roy SJ (2017a) Translating knowledge about abiotic stress tolerance to
4 breeding programmes. *The Plant Journal* 90, 898 – 917.
- 5 Gilliham M, Chapman S, Martin L *et al.* (2017b) The case for evidence-based policy to
6 support stress-resilient cropping systems. *Food and Energy Security* 6, 5 – 11.
- 7 Grassini P, Eskridge KM, Cassman KG (2013) Distinguishing between yield advances and
8 yield plateaus in historical crop production trends. *Nature Communications* 4, 2918 –
9 2929.
- 10 Harmens H, Hayes F, Mills G *et al.* (2018) Wheat yield responses to stomatal uptake of
11 ozone: Peak vs rising background ozone conditions. *Atmospheric Environment* 173, 1-
12 5.
- 13 Heagle AS (1989) Ozone and crop yield. *Annual Review of Phytopathology* 27, 397–423.
- 14 Heggestad HE (1991) Origin of Bel-W3, Bel-C and Bel-B tobacco varieties and their use as
15 indicators of ozone. *Environmental Pollution* 74, 264-291.
- 16 Hijmans RJ, Cameron SE, Parra JL *et al.* (2004) The WorldClim interpolated global
17 terrestrial climate surfaces, version 1.3. Available at <http://biogeo.berkeley.edu/>.
- 18 Hollaway MJ, Arnold SR, Challinor AJ, Emberson LD (2012) Intercontinental trans-
19 boundary contributions to ozone-induced crop yield losses in the Northern
20 Hemisphere. *Biogeosciences* 9, 271-292.
- 21 Huysmans M, Lema A S, Coll NS, Nowack MK (2017) Dying two deaths — programmed
22 cell death regulation in development and disease. *Current Opinion in Plant Biology*
23 35, 37-44.

Formatted: Font: 12 pt, Not Bold

Formatted: Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt, Not Bold, Italic

Formatted: Font: 12 pt, Not Bold

- 1 HWSD (2009) Harmonized World Soil Database (version 1.1).
2
3 FAO/IIASA/ISRIC/ISSCAS/JRC. FAO, Rome, Italy and IIASA, Laxenburg, Austria.
4
5 Jaswal S (2014) Challenges to Food Security in India. *IOSR Journal of Humanities and*
6 *Social Science* 19, 93-100.
7
8 Jiang LJ, Feng ZZ, Dai LL *et al.* (2018) Large variability in ambient ozone sensitivity across
9 19 ethylenediurea-treated Chinese cultivars of soybean is driven by total ascorbate.
10 *Journal of Environmental Sciences* 64, 10-22.
11
12 Jing LQ, Dombinov V, Shen SB *et al.* (2016) Physiological and genotype-specific factors
13 associated with grain quality changes in rice exposed to high ozone. *Environmental*
14 *Pollution* 210, 397-408.
15
16 Jordan DR, Hunt CH, Cruickshank AW *et al.* (2012) The relationship between the stay-green
17 trait and grain yield in elite sorghum hybrids grown in a range of environments. *Crop*
18 *Science* 52, 1153–1161.
19
20 Kangasjarvi J, Jaspers P, Kollist H (2005) Signalling and cell death in ozone-exposed plants.
21 *Plant Cell and Environment* 28, 1021-1036.
22
23 Klingberg J, Karlsson PE, Pihl Karlsson G (2012) Variation in ozone exposure in the
landscape of southern Sweden with consideration of topography and coastal climate.
Atmospheric Environment 47, 252-260.
24
25 Lal S, Venkataramani S, Naja M *et al.* (2017). Loss of crop yields in India due to surface
26 ozone: an estimation based on a network of observations. *Environmental Science and*
27 *Pollution Research* 24, 20972 – 20981.
28
29 Lesk C, Rowhani P, Ramankutty N (2016) Influence of extreme weather disasters on global
30 crop production. *Nature*, 529, 84-87.

Formatted: Font: Swedish (Sweden)

- 1 Lobell BD & Asseng S (2017) Comparing estimates of climate change impacts from process-
2 based and statistical crop models. *Environmental Research Letters*, 015001.
- 3 Lobell BD, Sibley A & Ortiz-Monasterio JI (2012) Extreme heat effects on wheat senescence
4 in India. *Nature Climate Change* 2, 186–189
- 5 Locato V, Gadaleta C, De Gara L, De Pinto MC (2008) Production of reactive species and
6 modulation of antioxidant network in response to heat shock: a critical balance for
7 cell fate. *Plant Cell and Environment* 31, 1606–1619.
- 8 Maas RP, Grennfelt RP (2016) Towards Cleaner Air. Scientific Assessment Report 2016.
9 EMEP Steering Body and Working Group on Effects of the Convention on Long-
10 Range Transboundary Air Pollution.
- 11 Manning WJ, Paoletti E, Sandermann H, Ernst (2011) Ethylenediurea (EDU): A research tool
12 for assessment and verification of the effects of ground level ozone on plants under
13 natural conditions. *Environmental Pollution* 159, 3283 – 3293.
- 14 McCormack ML, Dickie IA, Eissenstat DM et al. (2015) Redefining fine roots improves
15 understanding of below-ground contributions to terrestrial biosphere processes. *New*
16 *Phytologist* 207, 505–518.
- 17 McGrath JM, Betzelberger AM, Wang S et al (2015) An analysis of ozone damage to
18 historical maize and soybean yields in the United States. *Proceedings of the National*
19 *Academy of Sciences* 112(46), 14390–14395.
- 20 Mills G, Harmens H (2011) Ozone Pollution: A hidden threat to food security. Programme
21 Coordination Centre for the ICP Vegetation, Centre of Ecology & Hydrology,
22 Bangor, UK. <http://icpvegetation.ceh.ac.uk/>.
- 23 Mills G, Harmens H, Wagg S et al. (2016) Ozone impacts on vegetation in a nitrogen
24 enriched and changing climate. *Environmental Pollution* 208, 898 – 908.

Formatted: German (Germany)

- 1 Mills G, Hayes F, Simpson D *et al.* (2011) Evidence of widespread effects of ozone on crops
2 and (semi-)natural vegetation in Europe (1990–2006) in relation to AOT40- and flux-
3 based risk maps. *Global Change Biology* 17, 592 – 613.
- 4 Mills G, Pleijel H, Malley CS *et al.* (2018b) Tropospheric Ozone Assessment Report: Present
5 day tropospheric ozone distribution and trends relevant to vegetation. ~~Submitted to~~
6 ~~Accepted by~~ *Elementa: Science of the Anthropocene*.
- 7 Mills G, Sharps K, Simpson D *et al.* (2018a) Ozone pollution will compromise efforts to
8 increase global wheat production. ~~Accepted by~~ *Global Change Biology*.
9 <https://doi.org/10.1111/gcb.14157>.
- 10 Morgan PB, Ainsworth EA, Long SP (2003) How does elevated ozone impact soybean? A
11 meta-analysis of photosynthesis, growth and yield. *Plant Cell and Environment* 26,
12 1317-1328.
- 13 Mosier A, Kroeze C, Nevison C *et al.* (1998) Closing the global N₂O budget: nitrous oxide
14 emissions through the agricultural nitrogen cycle. *Nutrient Cycling in Agroecosystems*
15 52, 225–248.
- 16 Mueller, ND, Gerber JS, Johnston M *et al.* (2012) Closing yield gaps through nutrient and
17 water management. *Nature* 490, 254-257.
- 18 Mulchi CL, Lee E, Tuthill K, Olinick EV (1988) Influence of ozone stress on growth
19 processes, yields, and grain quality characteristics among soybean cultivars.
20 *Environmental Pollution* 53, 151-169.
- 21 Munne-Bosch S, Queval G, Foyer CH (2013) The Impact of Global Change Factors on
22 Redox Signaling Underpinning Stress Tolerance. *Plant Physiology* 161, 5-19.

Formatted: Font: (Default) Times New Roman, 12 pt

- 1 Oerke EC, Dehne H-W, Schönbeck F, Weber A (1994) Crop Production and Crop Protection.
2 Estimated losses in major food and cash crops. Elsevier Science, Amsterdam, the
3 Netherlands, 808 pp.
- 4 Oerke EC (2006) Crop losses to pests. *The Journal of Agricultural Science* 144, 31-43.
- 5 Osborne SA, Mills G, Hayes F *et al.* (2016) Has the sensitivity of soybean cultivars to ozone
6 pollution increased with time? An analysis of published dose–response data. *Global*
7 *Change Biology* 22, 3097 – 3111.
- 8 Pleijel H, Berglen Eriksen A, Danielsson H *et al.* (2006) Differential ozone sensitivity in an
9 old and a modern Swedish wheat cultivar – grain yield and quality, leaf chlorophyll
10 and stomatal conductance. *Environmental and Experimental Botany* 56, 63-71.
- 11 Pleijel H, Broberg MC, Uddling J, Mills G (2018). Current surface ozone concentrations
12 significantly decrease wheat growth, yield and quality. *Science of the Total*
13 *Environment* 613–614, 687–692.
- 14 Pleijel H, Danielsson H, Emberson L *et al.* (2007) Ozone risk assessment for agricultural
15 crops in Europe: Further development of stomatal flux and flux-response relationships
16 for European wheat and potato. *Atmospheric Environment* 41, 3022-3040.
- 17 Pleijel H, Danielsson H, Pihl Karlsson G *et al.* (2000). An ozone flux-response relationship
18 for wheat. *Environmental Pollution* 109, 453-462.
- 19 Pleijel H, Danielsson H, Simpson D, Mills G (2014) Have ozone effects on carbon
20 sequestration been overestimated? A new biomass response function for wheat.
21 *Biogeosciences* 11, 4521–4528.
- 22 Rai R, Agrawal M, Agrawal SB (2007) Assessment of yield losses in tropical wheat using
23 open top chambers, agriculture. *Atmospheric Environment* 41, 9543–9554.

Formatted: Font: Swedish (Sweden)

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

Ray DK, Ramankutty N, Mueller ND *et al.* (2012) Recent patterns of crop yield growth and stagnation. *Nature Communications* 3, 1293 – 1300.

Rebetzke GJ, Condon AG, Richards RA, Farquhar GD (2002) Selection for reduced carbon isotope discrimination increases aerial biomass and grain yield of rainfed bread wheat. *Crop Science* 42, 739–745.

Reynolds MP, Pierre CS, Saad ASI, Vargas M, Condon AG (2007) Evaluating potential genetic gains in wheat associated with stress-adaptive trait expression in elite genetic resources under drought and heat stress. *Crop Science* 47(S3), S172–S189.

Roche D (2015) Stomatal conductance is essential for higher yield of C3 crops. *Critical Reviews in Plant Science* 34, 429 – 453.

Sánchez-Martín L, Bermejo-Bermejo V, García-Torres L *et al.* (2017) Nitrogen soil emissions and belowground plant processes in Mediterranean annual pastures are altered by ozone exposure and N-inputs. *Atmospheric Environment* 165, 12–22.

Sarkar A, Agrawal SB (2010) Elevated ozone and two modern wheat cultivars: an assessment of dose dependent sensitivity with respect to growth, reproductive and yield parameters. *Environmental and Experimental Botany* 69, 328–337.

Semenov MA & Stratonovitch P (2013) Designing high-yielding wheat ideotypes for a changing climate. *Food and Energy Security* 2, 2048 – 3694.

Shi GY, Yang LX, Wang YX *et al.* (2009) Effects of elevated O₃ concentration on winter wheat and rice yields in the Yangtze River Delta, China. *Agriculture, Ecosystems and Environment* 131, 178–184.

Singh S, Agrawal SB (2011) Cultivar-Specific Response of Soybean (*Glycine max* L.) to ambient and elevated concentrations of ozone under open top chambers. *Water, Air and Soil Pollution* 217, 283–302.

Formatted: Swedish (Sweden)

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt, Italic

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: Italic

- 1 Simpson D, Arneth A, Mills G *et al.* (2014) Ozone — the persistent menace: interactions
2 with the N cycle and climate change. *Current Opinion in Environmental*
3 *Sustainability*, 9–10, 9-19
- 4 Simpson D, Benedictow A, Berge H *et al.* (2012) The EMEP MSC-W chemical transport
5 model – technical description. *Atmospheric Chemistry and Physics*, 12, 7825-
6 7865.
- 7 Simpson, D, Bergström R, Imhof H, Wind P (2017) Updates to the EMEP/MS-CW model,
8 2016-2017 Transboundary particulate matter, photo-oxidants, acidifying and
9 eutrophying components. Status Report 1/2017, The Norwegian Meteorological
10 Institute, Oslo, Norway, www.emep.int, 115-122.
- 11 Simpson D, Emberson L, Ashmore M, Tuovinen J (2007) A comparison of two different
12 approaches for mapping potential ozone damage to vegetation. A model study.
13 *Environmental Pollution* 146, 715-725.
- 14 Singh E, Rai R, Pandey B, Agrawal M (2017) Development of Resistance in Two Wheat
15 Cultivars Against Constant Fumigation of Ozone. *Proceedings of the National*
16 *Academy of Sciences, India Section B: Biological Sciences*, 1-14.
- 17 Siyiannis VF, Protonotarios VE, Zechmann B *et al.* (2012) Comparative spatiotemporal
18 analysis of root aerenchyma formation processes in maize due to sulphate, nitrate or
19 phosphate deprivation. *Protoplasma*, 249, 671-686.
- 20 Suzuki N, Koussevitzky S, Mittler R, Miller G (2012) ROS and redox signalling in the
21 response of plants to abiotic stress. *Plant Cell and Environment* 35, 259-270.
- 22 Tai APK & Martin MV (2017) Impacts of ozone air pollution and temperature extremes on
23 crop yields: Spatial variability, adaptation and implications for future food security.
24 *Atmospheric Environment* 169, 11 -21.

Formatted: German (Germany)

1 Tao F, Zhang Z, Zhang S. *et al.* (2016) Variability in crop yields associated with climate
2 anomalies in China over the past three decades. *Regional Environmental Change* 16,
3 1715 – 1723.

4

5 Tang H, Takigawa M, Liu G *et al.* (2013) A projection of ozone-induced wheat production
6 loss in China and India for the years 2000 and 2020 with exposure-based and flux-
7 based approaches. *Global Change Biology* 19, 2739-2752.

8 Teixeira EI, Fischer G, van Velthuisen H *et al.* (2013). Global hot-spots of heat stress on
9 agricultural crops due to climate change. *Agricultural and Forest Meteorology* 170,
10 206-215.

11 Tilman D, Balzer C, Hill J, Befort BL (2011) Global food demand and the sustainable
12 intensification of agriculture. *Proceedings of the National Academy of Sciences* 108,
13 20260-20264

14 Tomer R, Bhatia A, Kumar V *et al.* (2015) Impact of elevated ozone on growth, yield and
15 nutritional quality of two wheat species in Northern India. *Aerosol and Air Quality*
16 *Research* 15, 329-340.

17 Trabucco A & Zomer RJ (2009) Global Aridity Index (Global-Aridity) and Global Potential
18 Evapo-Transpiration (Global-PET) Geospatial Database. *CGIAR Consortium for*
19 *Spatial Information*. Published online, available from the CGIAR-CSI GeoPortal at:
20 <http://www.csi.cgiar.org>.

21 Ueda Y, Frimpong F, Qi Y *et al.* (2015a) Genetic dissection of ozone tolerance in rice (*Oryza*
22 *sativa* L.) by a genome-wide association study. *Journal of Experimental Botany* 66,
23 293-306.

Formatted: Font: Italic

Formatted: Pattern: Clear (White)

Formatted: Font: Italic

Formatted: Indent: First line: 0", Space Before: 0 pt, Line spacing: single, Pattern: Clear (White)

Formatted: Font: Italic

Formatted: Font: Italic

- 1 Ueda Y, Siddique S, Frei M (2015b) A Novel Gene, OZONE-RESPONSIVE APOPLASTIC
2 PROTEIN1, Enhances Cell Death in Ozone Stress in Rice. *Plant Physiology* 169,
3 873-889.
- 4 UNEP (United Nations Environment Programme), (1997) World atlas of desertification 2ED.
5 UNEP, London.
- 6 Van Dingenen R, Dentener FJ, Raes F *et al.* (2009) The global impact of ozone on
7 agricultural crop yields under current and future air quality legislation. *Atmospheric*
8 *Environment* 43, 604-618.
- 9 Van Doorn WG (2011) Classes of programmed cell death in plants, compared to those in
10 animals. *Journal of Experimental Botany* 62, 4749-4761.
- 11 Vollsnes AV, Kruse OMO, Eriksen AB *et al.* (2010) *In vivo* root growth dynamics of ozone
12 exposed *Trifolium subterraneum*. *Environmental and Experimental Botany* 69, 183-
13 188.
- 14 Wagg SK (2012) A mechanistic study of the implications of ozone and drought effects on
15 vegetation for global warming. PhD Thesis, University of Lancaster.
- 16 Wang Y, Frei M (2011) Stressed food - The impact of abiotic environmental stresses on crop
17 quality. *Agriculture Ecosystems & Environment* 141, 271-286.
- 18 Wang Y, Yang L, Hoeller M *et al.* (2014) Pyramiding of ozone tolerance QTLs OzT8 and
19 OzT9 confers improved tolerance to season-long ozone exposure in rice.
20 *Environmental and Experimental Botany* 104, 26-33.
- 21 Wassmann R, Jagadish SVK, Heuer S *et al.* (2009) Climate change affecting rice production:
22 the physiological and agronomic basis for possible adaptation strategies. *Advances in*
23 *Agronomy* 101, 59-122.

- 1 Wild O, Fiore AM, Shindell DT *et al.* (2012) Modelling future changes in surface ozone: a
2 parameterized approach. *Atmospheric Chemistry and Physics*, 12, 2037-2054.
- 3 Wilkinson S, Davies WJ (2009) Ozone suppresses soil drying- and abscisic acid (ABA)-
4 induced stomatal closure via an ethylene-dependent mechanism. *Plant, Cell and*
5 *Environment* 32, 949-959.
- 6 Wilkinson S, Davies WJ (2010) Drought, ozone, ABA and ethylene: new insights from cell
7 to plant to community. *Plant, Cell and Environment* 33, 510-525.
- 8 Wilkinson S, Mills G, Illidge R *et al.* (2012) How is ozone pollution reducing our food
9 supply? *Journal of Experimental Botany* 63, 527-536.
- 10 Wu L-B, Ueda Y, Lai S-K, Frei M (2017) Shoot tolerance mechanisms to iron toxicity in rice
11 (*Oryza sativa* L.). *Plant Cell and Environment* 40, 570-584.
- 12 Yendrek CR, Koester RP, Ainsworth EA (2015) A comparative analysis of transcriptomic,
13 biochemical, and physiological responses to elevated ozone identifies species-specific
14 mechanisms of resilience in legume crops. *Journal of Experimental Botany* 66, 7101-
15 7112.
- 16 Yendrek CR, Erice G, Montes CM *et al.* (2017) Elevated ozone reduces photosynthetic
17 carbon gain by accelerating leaf senescence of inbred and hybrid maize in a
18 genotype-specific manner. *Plant Cell and Environment* 40, 3088-3100.
- 19 Yuan XY, Calatayud V, Jiang LJ *et al.* (2015) Assessing the effects of ambient ozone in
20 China on snap bean genotypes by using ethylenediurea (EDU). *Environmental*
21 *Pollution* 205, 199-208.
- 22 Zhang H, Xue Y, Wang Z *et al.* (2009) An alternate wetting and moderate soil drying regime
23 improves root and shoot growth in rice. *Crop Science* 49, 2246-2260

- 1 Zhou B, Elazab A, Bort J *et al.* (2016) Agronomic and physiological responses of Chinese
2 facultative wheat genotypes to high-yielding Mediterranean conditions. *Journal of*
3 *Agricultural Science* 154, 870–889.
- 4 Zhu X, Feng Z, Sun T *et al.* (2011) Effects of elevated ozone concentration on yield of four
5 Chinese cultivars of winter wheat under fully open-air field conditions. *Global*
6 *Change Biology* 17, 2697–2706.
- 7 Zobel RW (2016) Fine roots – functional definition expanded to crop species? *New*
8 *Phytologist* 212, 310–312.
- 9 Zobel RW, Baligar VC, Kinraide TB (2007) Fine root diameters can change in response to
10 changes in nutrient concentrations. *Plant and Soil* 297, 243–254.
- 11

Figure legends

Fig. 1. Response functions for (a) soybean, (b) wheat, (c) rice and (d) maize derived from published data using the growing season ozone (7h mean, M7 in ppb) in the experiments. Data points are presented per cultivar/variety, with sources of data provided in the Supporting Information (Table S3). The response functions are: Soybean, $RY = -0.0050x + 1.001$ (r^2 (adj) = 0.625, $p < 0.001$); Wheat, $RY = -0.0048x + 0.96$ (r^2 (adj) = 0.547, $p < 0.001$); Rice, $RY = -0.0021x + 0.987$ (r^2 (adj) = 0.347, $p < 0.001$); and Maize, $RY = -0.0031x + 1.03$ (r^2 (adj) = 0.617, $p < 0.001$).

Fig. 2: The global effects of five biotic and abiotic stresses on soybean. All data are presented for the $1 \times 1^\circ$ grid squares where the mean production of soybean was > 500 tonnes (0.0005 Tg). (a) Presents the effects of ozone on crop production (thousand tonnes or 0.001 Tg per grid square), and (b) the percentage yield loss due to ozone, averaged for the period 2010-2012. In (d) to (h), the Yield Constraint Score (YCS) is presented per grid square on a scale of 1 to 5, where 5 is the highest level of stress (see Table 1) for ozone, pests and diseases, nutrients, heat stress and aridity, respectively, whilst (c) is the total YCS (YCS_{all}) calculated from the sum of each of these per grid square. The regional impacts are summarised in Table 2 for the five highest producing regions and Table S4 provides all country and regional means.

Fig. 3: The global effects of five biotic and abiotic stresses on wheat. All data are presented for the $1 \times 1^\circ$ grid squares where the mean production of wheat was > 500 tonnes (0.0005 Tg). (a) Presents the effects of ozone on crop production (thousand tonnes or 0.001 Tg per grid square), and (b) the percentage yield loss due to ozone, averaged for the period 2010-2012. In (d) to (h), the Yield Constraint Score (YCS) is presented per grid square on a scale of 1 to 5, where 5 is the highest level of stress (see Table 1) for ozone, pests and diseases, nutrients, heat stress and aridity, respectively, whilst (c) is the total YCS (YCS_{all}) calculated from the sum of each of these per grid square. The regional impacts are summarised in Table 2 for the five highest producing regions and Table S5 provides all country and regional means.

Fig. 4: The global effects of five biotic and abiotic stresses on rice. All data are presented for the $1 \times 1^\circ$ grid squares where the mean production of rice was > 500 tonnes (0.0005 Tg). (a) Presents the effects of ozone on crop production (thousand tonnes or 0.001 Tg per grid square), and (b) the percentage yield loss due to ozone, averaged for the period 2010-2012. In (d) to (h), the Yield Constraint Score (YCS) is presented per grid square on a scale of 1 to 5, where 5 is the highest level of stress (see Table 1) for ozone, pests and diseases, nutrients, heat stress and aridity, respectively, whilst (c) is the total YCS (YCS_{all}) calculated from the sum of each of these per grid square. The regional impacts are summarised in Table 2 for the five highest producing regions and Table S6 provides all country and regional means.

Comment [MGE9]: Provided here separately as a list

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Fig. 5: The global effects of five biotic and abiotic stresses on maize. All data are presented for the 1 x 1° grid squares where the mean production of maize was > 500 tonnes (0.0005 Tg). (a) Presents the effects of ozone on crop production (thousand tonnes or 0.001 Tg per grid square), and (b) the percentage yield loss due to ozone, averaged for the period 2010-2012. In (d) to (h), the Yield Constraint Score (YCS) is presented per grid square on a scale of 1 to 5, where 5 is the highest level of stress (see Table 1) for ozone, pests and diseases, nutrients, heat stress and aridity, respectively, whilst (c) is the total YCS (YCS_{all}) calculated from the sum of each of these per grid square. The regional impacts are summarised in Table 2 for the five highest producing regions and Table S7 provides all country and regional means.

Fig. 6: Yield Constraint Score (YCS) for five constraints on the yield of (a) wheat and (b) rice in the five Indian states with the highest production per crop. The bars represent the mean YCS per 1 x 1° grid square per state on a scale of 1 to 5, where 5 is the highest level of stress (see Table 1), rounded to the nearest integer. Note: The YCS for pests and disease is only available at the National Scale for India (score 4 for wheat and 5 for rice) and is presented here for information.

Fig. 7: An ideotype for an ozone-tolerant crop. ‘+’ indicates where there would be a benefit for other stresses of improving tolerance to ozone for the trait, whilst ‘-’ indicates a trade-off, and ‘0’ is no effect.