



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/140165/>

Version: Accepted Version

Article:

Buryak, A, Clader, E and Tessler, RJ (2019) Closed extended r-spin theory and the Gelfand–Dickey wave function. *Journal of Geometry and Physics*, 137. pp. 132-153. ISSN: 0393-0440

<https://doi.org/10.1016/j.geomphys.2018.11.007>

© 2018 Elsevier B.V. All rights reserved. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

CLOSED EXTENDED r -SPIN THEORY AND THE GELFAND–DICKEY WAVE FUNCTION

ALEXANDR BURYAK, EMILY CLADER, AND RAN J. TESSLER

ABSTRACT. We study a generalization of genus-zero r -spin theory in which exactly one insertion has a negative-one twist, which we refer to as the “closed extended” theory, and which is closely related to the open r -spin theory of Riemann surfaces with boundary. We prove that the generating function of genus-zero closed extended intersection numbers coincides with the genus-zero part of a special solution to the system of differential equations for the wave function of the r -th Gelfand–Dickey hierarchy. This parallels an analogous result for the open r -spin generating function in the companion paper [5] to this work.

1. INTRODUCTION

Witten’s conjecture [26], which was proposed in 1991 and soon proven by Kontsevich [18], states that the generating function for the integrals of the cotangent line classes $\psi_1, \dots, \psi_n \in H^2(\overline{\mathcal{M}}_{g,n})$ on the moduli space of curves is governed by the Korteweg–de Vries (KdV) hierarchy. At around the same time, Witten also proposed a generalization of his conjecture [27], in which the moduli space of curves is enhanced to the moduli space $\overline{\mathcal{M}}_{g,\{\alpha_1,\dots,\alpha_n\}}^{1/r}$ of r -spin structures. The latter is a natural compactification of the space of smooth marked curves $(C; z_1, \dots, z_n)$ with a line bundle S and an isomorphism

$$S^{\otimes r} \cong \omega_C \left(-\sum_{i=1}^n \alpha_i [z_i] \right),$$

where $\alpha_i \in \{0, 1, \dots, r-1\}$. This space admits a virtual fundamental class c_W , which is referred to as *Witten’s class* and is defined in genus zero by

$$c_W := e((R^1\pi_*\mathcal{S})^\vee),$$

with $\pi : \mathcal{C} \rightarrow \overline{\mathcal{M}}_{0,\{\alpha_1,\dots,\alpha_n\}}^{1/r}$ the universal curve and \mathcal{S} the universal r -spin structure; constructions of c_W in higher genus have now been given by a number of authors [22, 8, 19, 15, 7]. Let t_d^α and ε be formal variables, for $0 \leq \alpha \leq r-1$ and $d \geq 0$. The r -spin Witten conjecture states that if

$$F_{r,c}^{1,c}(t_*, \varepsilon) := \sum_{\substack{g \geq 0, n \geq 1 \\ 2g-2+n > 0}} \sum_{\substack{0 \leq \alpha_1, \dots, \alpha_n \leq r-1 \\ d_1, \dots, d_n \geq 0}} \frac{\varepsilon^{2g-2}}{n!} \left(r^{1-g} \int_{\overline{\mathcal{M}}_{g,\{\alpha_1,\dots,\alpha_n\}}^{1/r}} c_W \cap \psi_1^{d_1} \cdots \psi_n^{d_n} \right) t_{d_1}^{\alpha_1} \cdots t_{d_n}^{\alpha_n}$$

is the generating function of ψ -intersection numbers against Witten’s class, then $\exp(F_{r,c}^{1,c})$ becomes, after a certain rescaling of the variables t_d^α , a tau-function of the r -th Gelfand–Dickey (r -GD) hierarchy in the standard normalization of flows [13]. The superscript “ c ”, which stands for “closed”, is to distinguish this theory from the open theory discussed below. This result was proven by Faber–Shadrin–Zvonkine [14].

If one allows that exactly one of the indices α_i is equal to -1 and the rest lie in the range $\{0, 1, \dots, r-1\}$, then the space $\overline{\mathcal{M}}_{0,\{\alpha_1,\dots,\alpha_n\}}^{1/r}$ is still defined and $R^1\pi_*\mathcal{S}$ is still a vector bundle

The authors would like to thank Rahul Pandharipande and Jérémy Guéré for many fruitful discussions. The work of A. B. (Section 4) was supported by the grant no. 16-11-10260 of the Russian Science Foundation. E. C. was partially supported by NSF DMS Grant 1810969. R. T. was supported by a research grant from the Center for New Scientists of Weizmann Institute, and by Dr. Max Rössler, the Walter Haefner Foundation and the ETH Zürich Foundation.

in genus zero, so genus-zero r -spin intersection numbers can be defined exactly as above. This kind of intersection number was first considered in the paper [16]. We refer to this theory as *closed extended r -spin theory*, and we define a genus-zero generating function by

$$F_0^{\frac{1}{r}, \text{ext}}(t_*) := \sum_{n \geq 2} \sum_{\substack{0 \leq \alpha_1, \dots, \alpha_n \leq r-1 \\ d_1, \dots, d_n \geq 0}} \frac{1}{n!} \left(r \int_{\overline{\mathcal{M}}_{0, \{\alpha_1, \dots, \alpha_n, -1\}}^{1/r}} c_W \cap \psi_1^{d_1} \cdots \psi_n^{d_n} \right) t_{d_1}^{\alpha_1} \cdots t_{d_n}^{\alpha_n}.$$

In the current paper, we prove that $F_0^{\frac{1}{r}, \text{ext}}$ coincides with the genus-zero part of a special solution of the system of differential equations for the wave function of the r -GD hierarchy. The proof of this result proceeds by verifying that the closed extended r -spin correlators satisfy certain topological recursion relations that allow the entire theory to be recovered from just two initial conditions that can be explicitly calculated. We then verify that the genus-zero part of the special solution satisfies the same recursions and the same initial conditions.

The reason for our interest in the closed extended generating function arises from an intriguing connection to open r -spin theory, which is the generalization of r -spin theory to Riemann surfaces with boundary. In [5], we construct a moduli space $\overline{\mathcal{M}}_{0, k, l}^{1/r}$ of “graded r -spin disks” that generalizes $\overline{\mathcal{M}}_{0, n}^{1/r}$ to genus-zero curves C equipped with an involution that realizes C as two copies of a Riemann surface with boundary Σ , glued along their common boundary. The moduli space $\overline{\mathcal{M}}_{0, k, l}^{1/r}$ itself has boundary and is not necessarily canonically oriented, so one must prescribe boundary conditions for sections of bundles and specify relative orientations in order to ensure that integration of their relative top Chern class is well-defined. After carrying out this technical work, we obtain in [5] a definition of open r -spin correlators. When calculating recursions for open genus-zero correlators, the closed extended correlators appear naturally, and in fact, there is an intimate relation between the genus-zero sectors of the two theories that we call the open-closed correspondence; see Section 5 below. The origin of this correspondence is at present mysterious.

1.1. Plan of the paper. In Section 2, we recall the relevant background information on closed (non-extended) r -spin theory. Section 3 then generalizes the definitions to closed extended r -spin theory, proves the topological recursion relations, and calculates the two correlators that form the initial conditions for the potential $F_0^{\frac{1}{r}, \text{ext}}$. We turn to a detailed treatment of the integrable hierarchy in Section 4, which allows us to state the main result of the paper, Theorem 4.6, and to prove it. Finally, in Section 5, we explain the correspondence between closed extended and open r -spin theory.

2. BACKGROUND ON r -SPIN THEORY

We begin by reviewing the relevant background on the moduli space of r -spin structures and its intersection theory, referring the reader to [9, 17], among many other references, for more details.

Throughout what follows, fix an integer $r \geq 2$. An r -spin structure on a smooth marked curve $(C; z_1, \dots, z_n)$ of genus g is a line bundle L on C together with an isomorphism

$$(2.1) \quad L^{\otimes r} \cong \omega_{C, \log} := \omega_C \left(\sum_{i=1}^n [z_i] \right).$$

There is a smooth Deligne–Mumford stack $\mathcal{M}_{g, n}^{1/r}$ parameterizing such objects, equipped with a finite étale morphism to (indeed, a torsor structure over) the moduli space $\mathcal{M}_{g, n}$ of smooth curves. Some care must be taken in the compactification in order to preserve these properties of the moduli space of r -spin structures, and there are several ways to do so, as summarized in [12, Section 2.2]. In our case, we compactify by allowing orbifold structure.

More precisely, following [1], we define an *orbifold curve* as a one-dimensional Deligne–Mumford stack with a finite ordered collection of marked points and at worst nodal singularities such that

- (a) the only points with nontrivial isotropy are marked points and nodes;
- (b) all nodes are *balanced*—i.e., in the local picture $\{xy = 0\}$ at a node, the action of the distinguished generator of the isotropy group \mathbb{Z}_k is given by

$$(x, y) \mapsto (\zeta_k, \zeta_k^{-1}y),$$

where ζ_k is a primitive k th root of unity.

An orbifold curve is said to be *r -stable*, following [9], if the coarse underlying marked curve is stable and the isotropy group is \mathbb{Z}_r at every special point.

Let $(C; z_1, \dots, z_n)$ be an r -stable curve. An *r -spin structure* on C is an orbifold line bundle L together with an isomorphism as in (2.1). If $z_0 \in C$ is either a marked point or a branch of a node, then the *multiplicity* of L at z_0 is defined as the integer $m \in \{0, 1, \dots, r-1\}$ such that, in local coordinates (z, v) on the total space of L near z_0 , the action of the distinguished generator of the isotropy group \mathbb{Z}_r at z_0 is given by

$$(z, v) \mapsto (\zeta_r z, \zeta_r^m v).$$

A standard but crucial fact about the multiplicities is that they determine the relationship between L and its pushforward $|L|$ to the coarse underlying curve. Specifically, suppose that $C' \subset C$ is an irreducible component with special points $\{z_k\}$ at which the multiplicities of L are $\{m_k\}$. Then, if $\rho : C' \rightarrow |C'|$ is the natural map to the coarse underlying curve, we have

$$L|_{C'} = \rho^* \left(|L|_{|C'|} \right) \otimes \mathcal{O}_{C'} \left(\sum_k \frac{m_k}{r} [z_k] \right).$$

Given that $\omega_{C', \log} = \rho^* \omega_{|C'|, \log}$, we find that $|L|_{|C'|}$ satisfies the equation

$$(2.2) \quad \left(|L|_{|C'|} \right)^{\otimes r} \cong \omega_{|C'|, \log} \left(- \sum_k m_k [z_k] \right).$$

Using (2.2), one can prove (see, for example, the appendix of [11]) that there is an equivalence of categories between r -spin structures as above and orbifold line bundles L with an isomorphism

$$L^{\otimes r} \cong \omega_C \left(- \sum_{i=1}^n \mu_i [z_i] \right)$$

for which $\mu_i \in \{-1, 0, 1, \dots, r-2\}$ and the isotropy groups at all markings act trivially on the fiber of L . Finally, replacing L by $S := L(-\sum_{\mu_i=-1} [z_i])$, we find that there is a further equivalence with the category of orbifold line bundles satisfying

$$(2.3) \quad S^{\otimes r} \cong \omega_C \left(- \sum_{i=1}^n \alpha_i [z_i] \right)$$

with

$$\alpha_i \in \{0, 1, \dots, r-1\},$$

where again, the isotropy groups at all markings act trivially on the fiber of S . We view r -spin structures as in (2.3) in what follows, and we refer to the integers α_i as *twists*. When $\alpha_i = r-1$, we say that z_i is a *Ramond* marked point, and otherwise, it is said to be *Neveu–Schwarz*.

There is a proper, smooth Deligne–Mumford stack $\overline{\mathcal{M}}_{g,n}^{1/r}$ of dimension $3g-3+n$ parameterizing r -stable curves together with an orbifold line bundle S satisfying (2.3). It is equipped with a decomposition into open and closed substacks

$$\overline{\mathcal{M}}_{g, \{\alpha_1, \dots, \alpha_n\}}^{1/r} \subset \overline{\mathcal{M}}_{g,n}^{1/r}$$

on which S has twist α_i at z_i for each $i \in \{1, \dots, n\}$. Furthermore, it is equipped with a virtual fundamental class c_W from which a beautiful intersection theory can be defined. The construction of c_W , which we refer to as *Witten's class*, was first suggested by Witten in genus zero. Specifically, it is straightforward to check that in genus zero, if $\pi : \mathcal{C} \rightarrow \overline{\mathcal{M}}_{g,n}^{1/r}$ denotes the universal curve and \mathcal{S} the universal line bundle, then $R^0\pi_*\mathcal{S} = 0$ and hence $R^1\pi_*\mathcal{S}$ is a vector bundle. We define the *Witten bundle* as

$$\mathcal{W} := (R^1\pi_*\mathcal{S})^\vee = R^0\pi_*(\mathcal{S}^\vee \otimes \omega_\pi)$$

and set c_W to be its top Chern class:

$$c_W := e(\mathcal{W}).$$

Via the Riemann–Roch formula, one can check that the restriction of c_W to $\overline{\mathcal{M}}_{0,\{\alpha_1,\dots,\alpha_n\}}^{1/r}$ has complex codimension

$$\frac{-(r-2) + \sum_{i=1}^n \alpha_i}{r},$$

which is a non-negative integer if and only if $\overline{\mathcal{M}}_{0,\{\alpha_1,\dots,\alpha_n\}}^{1/r}$ is nonempty.

It is interesting and highly non-trivial to find the appropriate generalization of this definition to higher genus. Various constructions have now been given, by Polishchuk–Vaintrob [22], Chiodo [8], Mochizuki [19], Fan–Jarvis–Ruan [15], and Chang–Li–Li [7]. The result, in any of these cases, is a class on $\overline{\mathcal{M}}_{g,\{\alpha_1,\dots,\alpha_n\}}^{1/r}$ of complex codimension

$$(2.4) \quad e := \frac{(g-1)(r-2) + \sum_{i=1}^n \alpha_i}{r}.$$

All of these constructions have been shown to agree after pushforward to $\overline{\mathcal{M}}_{g,n}$ [20, Theorem 3], at least when all insertions are Neveu–Schwarz (which, by the Ramond vanishing explained below, is all that is needed).

One obtains correlators by integrating Witten's class against ψ -classes on the moduli space. Namely, for each $i \in \{1, \dots, n\}$, let \mathbb{L}_i be the cotangent line bundle to the coarse curve $|C|$ at the i th marked point. Then, in genus zero, we define closed r -spin correlators by

$$\left\langle \prod_{i=1}^n \tau_{d_i}^{\alpha_i} \right\rangle_0^{\frac{1}{r}} := r \int_{\overline{\mathcal{M}}_{0,\{\alpha_1,\dots,\alpha_n\}}^{1/r}} e \left(\mathcal{W} \oplus \bigoplus_{i=1}^l \mathbb{L}_i^{\oplus d_i} \right),$$

which is nonzero only if the equation

$$(2.5) \quad e + \sum_{i=1}^n d_i = 3g - 3 + n$$

is satisfied with $g = 0$.

Remark 2.1. Putting the coefficient r in front of the integral in the definition of the correlators $\left\langle \prod_{i=1}^n \tau_{d_i}^{\alpha_i} \right\rangle_0^{\frac{1}{r}}$ and, more generally, putting the rescaling coefficient r^{1-g} in the definition of the generating function $F^{\frac{1}{r},c}(t_*, \varepsilon)$ is a matter of tradition. This allows to present some results in a slightly more compact form.

A crucial fact that we require about these correlators is that they satisfy *Ramond vanishing*: if $\alpha_i = r-1$ for some i , then the correlator is zero. To prove this, suppose that S satisfies (2.3) and that $\alpha_1 = r-1$. Let \mathcal{S} be the universal line bundle on the universal curve $\pi : \mathcal{C} \rightarrow \overline{\mathcal{M}}_{0,\{\alpha_1,\dots,\alpha_n\}}^{1/r}$, let $\Delta_1 \subset \mathcal{C}$ be the divisor corresponding to the first marked point, and let $\tilde{\mathcal{S}} := \mathcal{S}(\Delta_1)$. Then there is an exact sequence

$$(2.6) \quad 0 \rightarrow R^0\pi_*\mathcal{S} \rightarrow R^0\pi_*\tilde{\mathcal{S}} \rightarrow \sigma_1^*\tilde{\mathcal{S}} \rightarrow R^1\pi_*\mathcal{S} \rightarrow R^1\pi_*\tilde{\mathcal{S}} \rightarrow 0,$$

where σ_1 is the section corresponding to the first marked point. We have $R^0\pi_*\tilde{\mathcal{S}} = 0$ and

$$(\sigma_1^*\tilde{\mathcal{S}})^{\otimes r} \cong \sigma_1^*\omega_{\pi,\log} \cong \mathcal{O}_{\overline{\mathcal{M}}_{0,\{\alpha_1,\dots,\alpha_n\}}^{1/r}},$$

which implies that $e(\sigma_1^*\tilde{\mathcal{S}}_1) = 0$. By multiplicativity of the Euler class in (2.6), this implies that $c_W = 0$.

3. CLOSED EXTENDED THEORY: GEOMETRY

Although we have thus far only defined the Witten class c_W under the assumption that all twists lie in the range $\{0, 1, \dots, r-1\}$, there exists a smooth Deligne–Mumford moduli stack $\overline{\mathcal{M}}_{g,\{\alpha_1,\dots,\alpha_n\}}^{1/r}$ parameterizing r -stable curves with a line bundle S satisfying (2.3) for *any* tuple of integers $\{\alpha_1, \dots, \alpha_n\}$. This observation was made by Jarvis–Kimura–Vaintrob [16], who studied precisely how the virtual class should vary when α_i is replaced by $\alpha_i + r$.

3.1. Definition of extended r -spin correlators. Suppose that $g = 0$,

$$(3.1) \quad \alpha_i \in \{-1, 0, \dots, r-1\}$$

for each i , and there is at most one i such that $\alpha_i = -1$. In this case, one still has $R^0\pi_*\mathcal{S} = 0$, so we can define $\mathcal{W} := (R^1\pi_*\mathcal{S})^\vee$ and set

$$c_W^{\text{ext}} := e((R^1\pi_*\mathcal{S})^\vee) = e(R^0\pi_*(\mathcal{S}^\vee \otimes \omega_\pi)).$$

When there is no i for which $\alpha_i = -1$, this simply recovers the definition of c_W given above. The same formula (2.4) (with $g = 0$) gives the complex codimension of c_W^{ext} , and there are correlators

$$\left\langle \prod_{i=1}^l \tau_{d_i}^{\alpha_i} \right\rangle_0^{\frac{1}{r}, \text{ext}} := r \int_{\overline{\mathcal{M}}_{0,\{\alpha_1,\dots,\alpha_n\}}^{1/r}} e \left(\mathcal{W} \oplus \bigoplus_{i=1}^l \mathbb{L}_i^{\oplus d_i} \right)$$

that vanish unless the dimension condition (2.5) is satisfied. We refer to these as *extended r -spin correlators*.

Remark 3.1. We caution the reader that the Ramond vanishing property does *not* hold for the extended r -spin correlators. For example, a nonvanishing extended correlator with an insertion of twist $r-1$ is calculated in Lemma 3.8.

3.2. Properties of the extended Witten class. Our goal for this section is to prove that the genus-zero extended r -spin correlators satisfy certain equations analogous to the string equation and topological recursion relations in Gromov–Witten theory. To do so, we must study how c_W^{ext} behaves under the inclusion of boundary divisors and forgetful morphisms.

Let us fix some notation. In general, boundary strata in $\overline{\mathcal{M}}_{g,\{\alpha_1,\dots,\alpha_n\}}^{1/r}$ are indexed by certain decorated graphs, in which each vertex v represents an irreducible component and is labeled with its genus $g(v)$, each half-edge h represents a half-node and is labeled with its twist $\alpha(h)$, and there are n numbered legs labeled with the twists $\alpha_1, \dots, \alpha_n$. We denote by $\vec{\alpha}(v)$ the tuple recording the twists at all half-edges incident to v , including the legs. Note that the elements of $\vec{\alpha}(v)$ lie in $\{-1, 0, 1, \dots, r-1\}$ and that the twist is -1 at each half-node on which the isotropy group acts trivially on the fiber of S .

Given Γ as above, let $\overline{\mathcal{M}}_\Gamma^{1/r} \subset \overline{\mathcal{M}}_{g,\{\alpha_1,\dots,\alpha_n\}}^{1/r}$ be the boundary stratum consisting of r -spin curves with decorated dual graph Γ . Let $\tilde{\Gamma}$ be the disconnected graph obtained by cutting all of the edges of Γ , and let

$$\overline{\mathcal{M}}_{\tilde{\Gamma}}^{1/r} := \prod_{v \in V(\Gamma)} \overline{\mathcal{M}}_{g(v), \vec{\alpha}(v)}^{1/r}$$

be the associated moduli space, where $V(\Gamma)$ is the vertex set of Γ . Unlike the moduli space of curves, the r -spin moduli space does not in general have a gluing map $\overline{\mathcal{M}}_{\tilde{\Gamma}}^{1/r} \rightarrow \overline{\mathcal{M}}_\Gamma^{1/r}$, because there is no canonical way to glue the fibers of S at the nodes. Nevertheless, letting $\overline{\mathcal{M}}_\Gamma$ and $\overline{\mathcal{M}}_{\tilde{\Gamma}}$

denote the moduli spaces of marked curves with dual graphs Γ and $\tilde{\Gamma}$, respectively, one has morphisms

$$(3.2) \quad \overline{\mathcal{M}}_{\tilde{\Gamma}}^{1/r} \xleftarrow{p} \overline{\mathcal{M}}_{\tilde{\Gamma}} \times_{\overline{\mathcal{M}}_{\Gamma}} \overline{\mathcal{M}}_{\Gamma}^{1/r} \xrightarrow{\tilde{\mu}} \overline{\mathcal{M}}_{\Gamma}^{1/r} \xrightarrow{\tilde{i}_{\Gamma}} \overline{\mathcal{M}}_{g, \{\alpha_1, \dots, \alpha_n\}}^{1/r}.$$

Let $g = 0$. Then for any decorated graph Γ of genus 0 we have $\overline{\mathcal{M}}_{\tilde{\Gamma}} = \overline{\mathcal{M}}_{\Gamma}$ and, therefore, $\overline{\mathcal{M}}_{\tilde{\Gamma}} \times_{\overline{\mathcal{M}}_{\Gamma}} \overline{\mathcal{M}}_{\Gamma}^{1/r} = \overline{\mathcal{M}}_{\Gamma}^{1/r}$. Then the map $\tilde{\mu}$ in (3.2) is the identity. Suppose now that the twists $\alpha_1, \dots, \alpha_n$ at the legs of Γ satisfy (3.1) with at most one i such that $\alpha_i = -1$, then we refer to Γ as a *genus-zero extended r -spin dual graph*. Note that even in this case, the vertex moduli spaces $\overline{\mathcal{M}}_{0, \vec{\alpha}(v)}^{1/r}$ may not themselves fit into the extended framework, since there may be more than one half-edge of twist -1 incident to a given vertex. Nevertheless, there is a consistent way to rectify the situation. Indeed, there is a unique function

$$\alpha' : H(\Gamma) \rightarrow \{-1, 0, 1, \dots, r-1\}$$

on the half-edge set $H(\Gamma)$ such that

- (a) $\alpha'(h) \equiv \alpha(h) \pmod{r}$, for all h ,
- (b) $\alpha'(h) = \alpha(h)$, if h is a leg,
- (c) for each edge $e = (h_1, h_2)$, such that $\alpha(h_1) = \alpha(h_2) = -1$, we have
 - (i) $\alpha'(h_1) = \alpha'(h_2) = r-1$, if $\alpha_1, \dots, \alpha_n \in \{0, \dots, r-1\}$,
 - (ii) exactly one $i \in \{1, 2\}$ with $\alpha'(h_i) = r-1$, if there exists $1 \leq j \leq n$, such that $\alpha_j = -1$,
- (d) for any edge e , each of the two connected components of the graph, obtained by cutting e , has at most one leg h for which $\alpha'(h) = -1$.

Let

$$(3.3) \quad \overline{\mathcal{M}}_{\tilde{\Gamma}}^{\text{ext}} := \prod_{v \in V(\Gamma)} \overline{\mathcal{M}}_{0, \vec{\alpha}'(v)}^{1/r},$$

where $\vec{\alpha}'(v)$ is the tuple recording the values of $\alpha'(h)$ at all of the half-edges incident to v . For each v , let $\mathcal{T}(v) \subset H(v)$ denote the subset of the incident half-edges for which $\alpha'(h) \neq \alpha(h)$. Then there is a morphism $\tau_v : \overline{\mathcal{M}}_{0, \vec{\alpha}(v)}^{1/r} \rightarrow \overline{\mathcal{M}}_{0, \vec{\alpha}'(v)}^{1/r}$ defined by sending an r -spin structure S with twists $\vec{\alpha}(v)$ to

$$S' := S \otimes \mathcal{O} \left(\sum_{h \in \mathcal{T}(v)} -[z_h] \right),$$

where z_h is the marked point corresponding to h ; it is straightforward to see that S' is an r -spin structure with twists $\vec{\alpha}'(v)$. The product of the morphisms τ_v defines

$$\tau : \overline{\mathcal{M}}_{\tilde{\Gamma}}^{1/r} \rightarrow \overline{\mathcal{M}}_{\tilde{\Gamma}}^{\text{ext}},$$

so we now have

$$\overline{\mathcal{M}}_{\tilde{\Gamma}}^{\text{ext}} \xleftarrow{q} \overline{\mathcal{M}}_{\Gamma}^{1/r} \xrightarrow{\tilde{i}_{\Gamma}} \overline{\mathcal{M}}_{0, \{\alpha_1, \dots, \alpha_n\}}^{1/r},$$

where $q := \tau \circ p$.

We can now state the decomposition property of the extended Witten class along nodes.

Lemma 3.2. *Let Γ be a genus-zero extended r -spin dual graph with two vertices v_1 and v_2 connected by a single edge e . Let $\overline{\mathcal{M}}_1$ and $\overline{\mathcal{M}}_2$ be the factors of $\overline{\mathcal{M}}_{\tilde{\Gamma}}^{\text{ext}}$ corresponding to the two vertices, and let $c_{W_1}^{\text{ext}}$ and $c_{W_2}^{\text{ext}}$ be the Witten classes on these two moduli spaces. Then*

$$(3.4) \quad q_{*l_{\Gamma}} \tilde{c}_W^{\text{ext}} = c_{W_1}^{\text{ext}} \boxtimes c_{W_2}^{\text{ext}}.$$

Before proving the lemma, let us present an alternative way to construct the extended Witten class. Note that for each $1 \leq i \leq n$ there is a canonical identification $\overline{\mathcal{M}}_{0,\{\alpha_1,\dots,\alpha_n\}}^{1/r} = \overline{\mathcal{M}}_{0,\{\alpha_1,\dots,\alpha_i+r,\dots,\alpha_n\}}^{1/r}$ defined by sending an r -spin structure S with twists $\alpha_1, \dots, \alpha_n$ to

$$S' := S \otimes \mathcal{O}(-[z_i]).$$

Moreover, this allows to identify the moduli spaces $\overline{\mathcal{M}}_{0,\{\alpha_1,\dots,\alpha_n\}}^{1/r}$ and $\overline{\mathcal{M}}_{0,\{\beta_1,\dots,\beta_n\}}^{1/r}$, if all the differences $\alpha_i - \beta_i$ are divisible by r . Denote by $\mathcal{S}_{\alpha_1,\dots,\alpha_n} \rightarrow \mathcal{C}$ the universal line bundle over the universal curve $\mathcal{C} \xrightarrow{\pi} \overline{\mathcal{M}}_{0,\{\alpha_1,\dots,\alpha_n\}}^{1/r}$.

Lemma 3.3. *For any $n \geq 2$ and an n -tuple $0 \leq \beta_1, \dots, \beta_n \leq r - 1$, the extended Witten class c_W^{ext} on $\overline{\mathcal{M}}_{0,\{-1,\beta_1,\dots,\beta_n\}}^{1/r}$ is equal to*

$$c_W^{\text{ext}} = c_{\text{rk}-1} \left((R^1 \pi_* \mathcal{S}_{r-1,\beta_1,\dots,\beta_n})^\vee \right) \in H^* \left(\overline{\mathcal{M}}_{0,\{r-1,\beta_1,\dots,\beta_n\}}^{1/r} \right) = H^* \left(\overline{\mathcal{M}}_{0,\{-1,\beta_1,\dots,\beta_n\}}^{1/r} \right),$$

where $\text{rk} = \text{rk} \left((R^1 \pi_* \mathcal{S}_{r-1,\beta_1,\dots,\beta_n})^\vee \right) = \frac{1+\sum \beta_i}{r}$.

Proof. Let us consider the exact sequence (2.6) over the moduli space $\overline{\mathcal{M}}_{0,\{r-1,\beta_1,\dots,\beta_n\}}^{1/r}$. Note that $\mathcal{S} = \mathcal{S}_{r-1,\beta_1,\dots,\beta_n}$ and $\tilde{\mathcal{S}} = \mathcal{S}_{-1,\beta_1,\dots,\beta_n}$. Therefore, by multiplicativity of the total Chern class in (2.6),

$$c_W^{\text{ext}} = e \left((R^1 \pi_* \mathcal{S}_{-1,\beta_1,\dots,\beta_n})^\vee \right) = c_{\text{rk}-1} \left((R^1 \pi_* \mathcal{S}_{r-1,\beta_1,\dots,\beta_n})^\vee \right),$$

which proves the lemma. \square

This lemma implies the following symmetry property of the extended Witten class.

Corollary 3.4. *Suppose $n \geq 1$ and $0 \leq \beta_1, \dots, \beta_n \leq r - 1$. Then, under the identification $\overline{\mathcal{M}}_{0,\{-1,r-1,\beta_1,\dots,\beta_n\}}^{1/r} = \overline{\mathcal{M}}_{0,\{r-1,-1,\beta_1,\dots,\beta_n\}}^{1/r}$, the extended Witten classes on these moduli spaces become equal. In other words,*

$$(3.5) \quad e \left((R^1 \pi_* \mathcal{S}_{-1,r-1,\beta_1,\dots,\beta_n})^\vee \right) = e \left((R^1 \pi_* \mathcal{S}_{r-1,-1,\beta_1,\dots,\beta_n})^\vee \right).$$

Proof. By Lemma 3.3, both sides of (3.5) are equal to $c_{\text{rk}-1} \left((R^1 \pi_* \mathcal{S}_{r-1,r-1,\beta_1,\dots,\beta_n})^\vee \right)$. \square

Proof of Lemma 3.2. Over $\overline{\mathcal{M}}_\Gamma^{1/r}$, there are two universal curves: namely, we define \mathcal{C}_Γ by the fiber diagram

$$\begin{array}{ccc} \mathcal{C}_\Gamma & \longrightarrow & \mathcal{C} \\ \pi \downarrow & & \downarrow \pi \\ \overline{\mathcal{M}}_\Gamma^{1/r} & \xrightarrow{\tilde{i}_\Gamma} & \overline{\mathcal{M}}_{0,\{\alpha_1,\dots,\alpha_n\}}^{1/r} \end{array}$$

and we define $\mathcal{C}_{\tilde{\Gamma}}$ by the fiber diagram

$$\begin{array}{ccc} \tilde{\mathcal{C}} & \longleftarrow & \mathcal{C}_{\tilde{\Gamma}} \\ \downarrow & & \downarrow \tilde{\pi} \\ \overline{\mathcal{M}}_{\tilde{\Gamma}}^{\text{ext}} & \xleftarrow{q} & \overline{\mathcal{M}}_\Gamma^{1/r} \end{array}$$

in which $\tilde{\mathcal{C}}$ is the universal curve over $\overline{\mathcal{M}}_{\tilde{\Gamma}}^{\text{ext}}$, induced by the product description (3.3). One can decompose $\mathcal{C}_{\tilde{\Gamma}} = \mathcal{C}_1 \sqcup \mathcal{C}_2$, and we let $\pi_i = \tilde{\pi}|_{\mathcal{C}_i}$, for $i = 1, 2$. Let

$$n : \mathcal{C}_{\tilde{\Gamma}} \rightarrow \mathcal{C}_\Gamma$$

be the universal normalization morphism.

Denote (by a slight abuse of notation) the pullback to \mathcal{C}_Γ of the universal line bundle on \mathcal{C} by \mathcal{S} , and let $\mathcal{S}_i = n^* \mathcal{S}|_{\mathcal{C}_i}$ for $i = 1, 2$. There is a normalization exact sequence

$$0 \rightarrow \mathcal{S} \rightarrow n_* n^* \mathcal{S} \rightarrow \mathcal{S}|_{\Delta_e} \rightarrow 0,$$

where $\Delta_e \subset \mathcal{C}_\Gamma$ picks out the node that corresponds to the edge e of Γ . The associated long exact sequence reads

$$(3.6) \quad 0 \rightarrow R^0 \pi_{1*} \mathcal{S}_1 \oplus R^0 \pi_{2*} \mathcal{S}_2 \rightarrow R^0 \pi_*(\mathcal{S}|_{\Delta_e}) \rightarrow R^1 \pi_* \mathcal{S} \rightarrow R^1 \pi_{1*} \mathcal{S}_1 \oplus R^1 \pi_{2*} \mathcal{S}_2 \rightarrow 0.$$

From here, we break the argument into three cases.

First, suppose that one (hence both) of the half-edges of e is Neveu–Schwarz. Then $R^0 \pi_*(\mathcal{S}|_{\Delta_e}) = 0$, since sections of an orbifold line bundle necessarily vanish at points where the isotropy group acts nontrivially on the fiber. Thus, we have

$$R^1 \pi_* \mathcal{S} \cong R^1 \pi_{1*} \mathcal{S}_1 \oplus R^1 \pi_{2*} \mathcal{S}_2$$

on $\overline{\mathcal{M}}_\Gamma^{1/r}$. This implies that $\tilde{i}_\Gamma^* c_W^{\text{ext}} = q^*(c_{W_1}^{\text{ext}} \boxtimes c_{W_2}^{\text{ext}})$, and the claim follows from the fact that $\deg q = 1$ [12, page 1348].

Next, suppose that one (hence both) of the half-edges of e is Ramond. Assume, without loss of generality, that the vertices v_1 and v_2 of Γ are incident to the legs marked by $\{1, \dots, n_1\}$ and $\{n_1 + 1, \dots, n\}$, respectively. Consider the universal line bundles $\mathcal{S}_{\alpha_1, \dots, \alpha_{n_1}, -1}$ and $\mathcal{S}_{-1, \alpha_{n_1+1}, \dots, \alpha_n}$ over $\tilde{\mathcal{C}}$, corresponding to the components $\overline{\mathcal{M}}_1$ and $\overline{\mathcal{M}}_2$, respectively. The pullbacks, via the map $\mathcal{C}_\Gamma \rightarrow \tilde{\mathcal{C}}$, of these line bundles to \mathcal{C}_1 and \mathcal{C}_2 , respectively, will be denoted by the same letters. Clearly,

$$\mathcal{S}_1 = \mathcal{S}_{\alpha_1, \dots, \alpha_{n_1}, -1}, \quad \mathcal{S}_2 = \mathcal{S}_{-1, \alpha_{n_1+1}, \dots, \alpha_n}.$$

Note that $R^0 \pi_*(\mathcal{S}|_{\Delta_e}) = \sigma_e^* \mathcal{S}$, where σ_e is the section of \mathcal{C}_Γ with image Δ_e . Since

$$(\sigma_e^* \mathcal{S})^{\otimes r} \cong \sigma_e^* \omega_{\pi, \log} \cong \mathcal{O}_{\overline{\mathcal{M}}_\Gamma^{1/r}},$$

we have $c_1(\sigma_e^* \mathcal{S}) = 0$.

Suppose, additionally, that $\alpha_1, \dots, \alpha_n \in \{0, \dots, r-1\}$. Then we have $R^0 \pi_{1*} \mathcal{S}_1 = R^0 \pi_{2*} \mathcal{S}_2 = 0$ and, by multiplicativity of the total Chern classes in (3.6), we find that

$$(3.7) \quad c_i((R^1 \pi_* \mathcal{S})^\vee) = c_i((R^1 \pi_{1*} \mathcal{S}_{\alpha_1, \dots, \alpha_{n_1}, -1})^\vee \oplus (R^1 \pi_{2*} \mathcal{S}_{-1, \alpha_{n_1+1}, \dots, \alpha_n})^\vee), \quad i \geq 0.$$

By the Ramond vanishing, the right-hand side of (3.4) is equal to zero. Note that $\text{rk}(R^1 \pi_* \mathcal{S}) = \text{rk}(R^1 \pi_{1*} \mathcal{S}_1 \oplus R^1 \pi_{2*} \mathcal{S}_2) + 1$, therefore, by (3.7), $\tilde{i}_\Gamma^* c_W^{\text{ext}} = e((R^1 \pi_* \mathcal{S})^\vee) = 0$.

Finally, suppose again that one (hence both) of the half-edges of e is Ramond, but there is a marked point of twist -1 . Without loss of generality, we can assume that $\alpha_1 = -1$. Let us write equation (3.7) for the n -tuple $r-1, \alpha_2, \dots, \alpha_n$ and for $i = \text{rk}(R^1 \pi_* \mathcal{S}_{r-1, \alpha_2, \dots, \alpha_n}) - 1$. Then, by Lemma 3.3, the left-hand side is equal to $\tilde{i}_\Gamma^* c_W^{\text{ext}}$. The right-hand side of (3.7) is equal to

$$\begin{aligned} e((R^1 \pi_{1*} \mathcal{S}_{r-1, \alpha_2, \dots, \alpha_{n_1}, -1})^\vee) e((R^1 \pi_{2*} \mathcal{S}_{-1, \alpha_{n_1+1}, \dots, \alpha_n})^\vee) &\stackrel{\text{by Corollary 3.4}}{=} \\ &= e((R^1 \pi_{1*} \mathcal{S}_{-1, \alpha_2, \dots, \alpha_{n_1}, r-1})^\vee) \cdot e((R^1 \pi_{2*} \mathcal{S}_{-1, \alpha_{n_1+1}, \dots, \alpha_n})^\vee) = \\ &= q^*(c_{W_1}^{\text{ext}} \boxtimes c_{W_2}^{\text{ext}}). \end{aligned}$$

Hence, $\tilde{i}_\Gamma^* c_W^{\text{ext}} = q^*(c_{W_1}^{\text{ext}} \boxtimes c_{W_2}^{\text{ext}})$ and using again that $\deg q = 1$ [12, page 1348] we see that the lemma is proved. \square

The other property of the extended Witten class that we require is its behavior under pullback via the forgetful map. There is only a forgetful map on the r -spin moduli space forgetting marked points whose twist equals zero. Let

$$\text{For}_{n+1}: \overline{\mathcal{M}}_{0, \{\alpha_1, \dots, \alpha_n, 0\}}^{1/r} \rightarrow \overline{\mathcal{M}}_{0, \{\alpha_1, \dots, \alpha_n\}}^{1/r}$$

be the map that forgets z_{n+1} and its orbifold structure and stabilizes the curve C as necessary.

Lemma 3.5. *Let c_W^{ext} be the Witten class for $\overline{\mathcal{M}}_{0, \{\alpha_1, \dots, \alpha_n\}}^{1/r}$, and let \tilde{c}_W^{ext} be the Witten class for $\overline{\mathcal{M}}_{0, \{\alpha_1, \dots, \alpha_n, 0\}}^{1/r}$. Then*

$$\tilde{c}_W^{\text{ext}} = \text{For}_{n+1}^* c_W^{\text{ext}}.$$

Proof. This follows immediately from the fact that the universal line bundle pulls back under the induced morphism $\widetilde{\text{For}}_{n+1} : \mathcal{C}' \rightarrow \mathcal{C}$ on the universal curves. \square

3.3. TRRs and string equation in extended r -spin theory. We are now prepared to state and prove the topological recursion relations satisfied by the extended r -spin correlators in genus zero. These follow from Lemmas 3.2 and 3.5, by essentially the same argument as given by Jarvis–Kimura–Vaintrob in [17]. We denote by $[n]$ the set $\{1, 2, \dots, n\}$.

Lemma 3.6. *For any i with $d_i > 0$ and any $j \neq k \in [n] \setminus \{i\}$, we have*

$$(3.8) \quad \left\langle \prod_{l \in [n]} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r}, \text{ext}} = \sum_{I \amalg J = [n] \setminus \{i\}} \sum_{\substack{\alpha = -1 \\ j, k \in J}}^{r-1} \left\langle \tau_0^\alpha \tau_{d_{i-1}}^{\alpha_i} \prod_{l \in I} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r}, \text{ext}} \left\langle \tau_0^{r-2-\alpha} \prod_{l \in J} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r}, \text{ext}}.$$

In particular, we have the following equations, where $K \subset [n]$ is the set of marked points whose twist is $r - 1$ and without loss of generality we can assume that $\alpha_1 = -1$:

(a) *(Neveu–Schwarz TRR) For any $i \in [n] \setminus (K \cup \{1\})$ with $d_i > 0$ and any $j \in [n] \setminus \{i, 1\}$ we have*

$$(3.9) \quad \left\langle \prod_{l \in [n]} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r}, \text{ext}} = \sum_{I \amalg J = [n] \setminus \{i\}} \sum_{\substack{\alpha = 0 \\ \{1, j\} \cup K \subseteq J}}^{r-2} \left\langle \tau_0^\alpha \tau_{d_{i-1}}^{\alpha_i} \prod_{l \in I} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r}, \text{ext}} \left\langle \tau_0^{r-2-\alpha} \prod_{l \in J} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r}, \text{ext}} + \\ + \sum_{I \amalg J = [n] \setminus \{i\}} \left\langle \tau_0^{-1} \tau_{d_{i-1}}^{\alpha_i} \prod_{l \in I} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r}, \text{ext}} \left\langle \tau_0^{r-1} \prod_{l \in J} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r}, \text{ext}}.$$

(b) *(Ramond TRR) For any $i \in K$ with $d_i > 0$ and any $j \in [n] \setminus \{1, i\}$ we have*

$$(3.10) \quad \left\langle \prod_{l \in [n]} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r}, \text{ext}} = \sum_{I \amalg J = [n] \setminus \{i\}} \left\langle \tau_0^{-1} \tau_{d_{i-1}}^{\alpha_i} \prod_{l \in I} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r}, \text{ext}} \left\langle \tau_0^{r-1} \prod_{l \in J} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r}, \text{ext}}.$$

(c) *(-1 TRR) Suppose that $d_1 > 0$. Then for any $j \neq k \in [n] \setminus \{1\}$ we have*

$$(3.11) \quad \left\langle \prod_{l \in [n]} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r}, \text{ext}} = \sum_{I \amalg J = [n] \setminus \{1\}} \sum_{\substack{\alpha = 0 \\ j, k \in J, K \subset I}}^{r-2} \left\langle \tau_0^\alpha \tau_{d_1-1}^{\alpha_1} \prod_{l \in I} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r}, \text{ext}} \left\langle \tau_0^{r-2-\alpha} \prod_{l \in J} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r}, \text{ext}} + \\ + \sum_{I \amalg J = [n] \setminus \{1\}} \left\langle \tau_0^{r-1} \tau_{d_1-1}^{\alpha_1} \prod_{l \in I} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r}, \text{ext}} \left\langle \tau_0^{-1} \prod_{l \in J} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r}, \text{ext}}.$$

Proof. Observe that the right-hand side of (3.8) is always defined. Indeed, while it could be the case that one of the two correlators in a summand has two insertions of twist -1 , it is always multiplied by a correlator with at least one insertion of twist $r - 1$ and no insertions of twist -1 ; thus, by Ramond vanishing in the usual closed theory, that summand of (3.8) vanishes.

The proof of (3.8) follows [17, Section 4.2]. Namely, on $\overline{\mathcal{M}}_{0,n}$, we apply the relation

$$\psi_i = \sum_{\Gamma} i_{\Gamma*}(1),$$

where the sum is over all single-edged graphs on which tail i is separated by the edge from tails j and k (see, for example, [28]). From here, we use the commutative diagram

$$(3.12) \quad \begin{array}{ccc} & \overline{\mathcal{M}}_{\Gamma}^{1/r} & \xrightarrow{\tilde{i}_{\Gamma}} \overline{\mathcal{M}}_{0, \{\alpha_1, \dots, \alpha_n\}}^{1/r} \\ & \swarrow q & \downarrow \tilde{\rho} \\ \overline{\mathcal{M}}_{\tilde{\Gamma}}^{\text{ext}} & & \downarrow \rho \\ & \searrow \rho' & \downarrow i_{\Gamma} \\ & \overline{\mathcal{M}}_{\Gamma} & \xrightarrow{i_{\Gamma}} \overline{\mathcal{M}}_{0,n} \end{array}$$

for each single-edged graph Γ . First of all, note that $i_{\Gamma}^* \rho_* = r \cdot \tilde{\rho}_* \tilde{i}_{\Gamma}^*$, which follows from the fact that $\deg \rho = \frac{1}{r}$ and $\deg \tilde{\rho} = \frac{1}{r^2}$. Combining this property with Lemma 3.2 shows that

$$\begin{aligned} r \cdot \rho_* (\psi_i \cap c_W^{\text{ext}}) &= r \sum_{\Gamma} \rho_* (\rho^* i_{\Gamma^*}(1) \cap c_W^{\text{ext}}) = \\ &= r \sum_{\Gamma} i_{\Gamma^*}(1) \cap \rho_* (c_W^{\text{ext}}) = \\ &= r \sum_{\Gamma} i_{\Gamma^*} (i_{\Gamma}^* \rho_* (c_W^{\text{ext}})) = \\ &= r^2 \sum_{\Gamma} i_{\Gamma^*} \tilde{\rho}_* \tilde{i}_{\Gamma}^* (c_W^{\text{ext}}) = \\ &= r^2 \sum_{\Gamma} i_{\Gamma^*} \rho'_* q_* \tilde{i}_{\Gamma}^* (c_W^{\text{ext}}) = \\ &= r^2 \sum_{\Gamma} i_{\Gamma^*} \rho'_* (c_{W_1}^{\text{ext}} \boxtimes c_{W_2}^{\text{ext}}). \end{aligned}$$

Finally, multiplying this equation by the remaining ψ -classes and integrating proves the claim. The other items are specializations of the general case, using that the extended theory agrees with the usual closed theory in the absence of an insertion of twist -1 , and that in the usual closed theory Ramond vanishing holds. \square

The string equation in the closed extended r -spin theory is exactly as usual:

Lemma 3.7. *We have*

$$\left\langle \tau_0^0 \prod_{i \in [n]} \tau_{d_i}^{\alpha_i} \right\rangle_0^{\frac{1}{r}, \text{ext}} = \begin{cases} \left\langle \sum_{\substack{i \in [n] \\ d_i > 0}} \tau_{d_i-1}^{\alpha_i} \prod_{j \neq i} \tau_{d_j}^{\alpha_j} \right\rangle_0^{\frac{1}{r}, \text{ext}}, & \text{if } n \geq 3, \\ \delta_{d_1, 0} \delta_{d_2, 0} \delta_{\alpha_1 + \alpha_2, r-2}, & \text{if } n = 2. \end{cases}$$

Proof. The string equation can either be deduced algebraically from the topological recursions above, or geometrically, by mimicking the proof of the ordinary string equation on $\overline{\mathcal{M}}_{0,n}$ and applying Lemma 3.5. \square

3.4. Base cases. In addition to the above relations, the proof that the extended r -spin correlators satisfy the equations for the wave function of the Gelfand–Dickey hierarchy requires two base cases. We collect these simple correlators in the following lemma.

Lemma 3.8. *We have*

$$\left\langle \tau_0^{-1} \tau_0^1 \tau_0^{r-2} \right\rangle_0^{1/r, \text{ext}} = 1$$

and

$$\left\langle \tau_0^{-1} \tau_0^1 \tau_0^{r-1} \tau_0^{r-1} \right\rangle_0^{1/r, \text{ext}} = -\frac{1}{r}.$$

Proof. In the first case, the moduli space is isomorphic to $B\mathbb{Z}_r$ and Witten’s class has rank zero, so the claim is immediate.

In the second case, Witten’s bundle has rank one, so

$$c_W^{\text{ext}} = c_1((R^1\pi_*\mathcal{S})^\vee) = -c_1(R^1\pi_*\mathcal{S}) = -\text{ch}_1(R^1\pi_*\mathcal{S}) = \text{ch}_1(R\pi_*\mathcal{S}).$$

The latter can be calculated via Chiodo’s Grothendieck–Riemann–Roch formula [10]. In our situation, Chiodo’s formula reads:

$$c_W^{\text{ext}} = \frac{B_2\left(\frac{1}{r}\right)}{2}\kappa_1 - \frac{B_2(0)}{2}\psi_1 - \frac{B_2\left(\frac{2}{r}\right)}{2}\psi_2 - \frac{B_2(1)}{2}\psi_3 - \frac{B_2(1)}{2}\psi_4 + 3\frac{rB_2\left(\frac{1}{r}\right)}{2}\tilde{i}_{\Gamma^*}(1),$$

where

$$B_2(x) = x^2 - x + \frac{1}{6}$$

is the second Bernoulli polynomial and Γ is any of the three one-edged graphs on $\overline{\mathcal{M}}_{0,\{-1,1,r-1,r-1\}}^{1/r}$, all of which yield the same divisor $\tilde{i}_{\Gamma^*}(1)$. From here, the claim is immediate from the fact that

$$\int_{\overline{\mathcal{M}}_{0,\{-1,1,r-1,r-1\}}^{1/r}} \kappa_1 = \int_{\overline{\mathcal{M}}_{0,\{-1,1,r-1,r-1\}}^{1/r}} \psi_i = \frac{1}{r},$$

for each i , and

$$\int_{\overline{\mathcal{M}}_{0,\{-1,1,r-1,r-1\}}^{1/r}} \tilde{i}_{\Gamma^*}(1) = \frac{1}{r^2},$$

in which the last integral is a consequence of the \mathbb{Z}_r scaling and the additional \mathbb{Z}_r ghost automorphisms on a nodal r -spin curve. \square

4. CLOSED EXTENDED THEORY: ALGEBRA

In this section we prove the main result of the paper, Theorem 4.6, which describes the function $F_0^{\frac{1}{r},\text{ext}}$ in terms of the r -th Gelfand–Dickey hierarchy.

Sections 4.1 and 4.2 contain the necessary background information on the KP hierarchy, its Gelfand–Dickey reduction, and their relation to the closed r -spin partition function. Then, in Section 4.3, we consider a special solution of the system of differential equations for the wave function of the r -GD hierarchy and discuss its main properties. In Section 4.4, we prove Theorem 4.6: the function $F_0^{\frac{1}{r},\text{ext}}$ coincides with the genus-zero part of the special solution. As a consequence of this result, in Section 4.5 we obtain a simple interpretation of the genus-zero part of the Lax operator of the r -GD hierarchy in terms of the potential $F_0^{\frac{1}{r},\text{ext}}$. Finally, in Section 4.6, we propose a conjecture about the structure of the closed extended r -spin correlators in all genera.

4.1. Brief review of the KP hierarchy. The material of this section is borrowed from the book [13].

Consider formal variables T_i for $i \geq 1$. A *pseudo-differential operator* A is a Laurent series

$$A = \sum_{n=-\infty}^m a_n(T_*, \varepsilon) \partial_x^n,$$

where $m \in \mathbb{Z}$, ∂_x is considered as a formal variable, and $a_n(T_*, \varepsilon)$ are formal power series in the variables T_i with coefficients that are complex Laurent polynomials in ε :

$$a_n(T_*, \varepsilon) \in \mathbb{C}[\varepsilon, \varepsilon^{-1}][[T_1, T_2, \dots]].$$

The non-negative and negative degree parts of the pseudo-differential operator A are defined by

$$A_+ := \sum_{n=0}^m a_n \partial_x^n \quad \text{and} \quad A_- := A - A_+,$$

and residue of the operator A is defined by

$$\text{res } A := a_{-1}.$$

The Laurent series

$$\widehat{A}(T_*, \varepsilon, z) := \sum_{n=-\infty}^m a_n(T_*, \varepsilon) z^n,$$

in which z is a formal variable, is called the *symbol* of the operator A .

The space of pseudo-differential operators is endowed with a structure of a non-commutative associative algebra, in which the multiplication is defined by the formula

$$\partial_x^k \circ f := \sum_{l=0}^{\infty} \frac{k(k-1)\dots(k-l+1)}{l!} \frac{\partial^l f}{\partial x^l} \partial_x^{k-l},$$

where $k \in \mathbb{Z}$, $f \in \mathbb{C}[\varepsilon, \varepsilon^{-1}][[T_1, T_2, \dots]]$, and the variable x is identified with T_1 . For any $r \geq 2$ and any pseudo-differential operator A of the form

$$A = \partial_x^r + \sum_{n=1}^{\infty} a_n \partial_x^{r-n},$$

there exists a unique pseudo-differential operator $A^{\frac{1}{r}}$ of the form

$$A^{\frac{1}{r}} = \partial_x + \sum_{n=0}^{\infty} \tilde{a}_n \partial_x^{-n},$$

such that $\left(A^{\frac{1}{r}}\right)^r = A$.

Consider the pseudo-differential operator

$$\mathcal{L} = \partial_x + \sum_{i \geq 1} u_i \partial_x^{-i}, \quad u_i \in \mathbb{C}[\varepsilon, \varepsilon^{-1}][[T_1, T_2, \dots]].$$

The *KP hierarchy* is the following system of partial differential equations for the power series u_i :

$$(4.1) \quad \frac{\partial \mathcal{L}}{\partial T_n} = \varepsilon^{n-1} [(\mathcal{L}^n)_+, \mathcal{L}], \quad n \geq 1.$$

For $n = 1$, the equation is equivalent to

$$\frac{\partial u_i}{\partial T_1} = \frac{\partial u_i}{\partial x}, \quad i \geq 1,$$

compatible with our identification of x with T_1 .

Remark 4.1. The factor ε^{n-1} is not included usually in the definition of the KP hierarchy (4.1). This rescaling is necessary, if we want to describe the function $\exp\left(F^{\frac{1}{r};c}(T_*, \varepsilon)\right)$ as a tau-function of the KP hierarchy.

Suppose an operator \mathcal{L} satisfies the system (4.1). Then there exists a pseudo-differential operator P of the form

$$(4.2) \quad P = 1 + \sum_{n \geq 1} p_n(T_*, \varepsilon) \partial_x^{-n},$$

satisfying $\mathcal{L} = P \circ \partial_x \circ P^{-1}$ and

$$(4.3) \quad \frac{\partial P}{\partial T_n} = -\varepsilon^{n-1} (\mathcal{L}^n)_- \circ P, \quad n \geq 1.$$

The operator P is called the *dressing operator*.

We can now introduce the notion of a tau-function. Denote by G_z the shift operator, which acts on a power series $f \in \mathbb{C}[\varepsilon, \varepsilon^{-1}][[T_1, T_2, \dots]]$ as follows:

$$G_z(f)(T_1, T_2, T_3, \dots) := f\left(T_1 - \frac{1}{z}, T_2 - \frac{1}{2\varepsilon z^2}, T_3 - \frac{1}{3\varepsilon^2 z^3}, \dots\right).$$

Let $P = 1 + \sum_{n \geq 1} p_n(T_*, \varepsilon) \partial_x^{-n}$ be the dressing operator of some operator \mathcal{L} satisfying the KP hierarchy (4.1). Then there exists a series $\tau \in \mathbb{C}[\varepsilon, \varepsilon^{-1}][[T_1, T_2, T_3, \dots]]$ with constant term $\tau|_{T_i=0} = 1$ for which

$$\widehat{P} = \frac{G_z(\tau)}{\tau}.$$

The series τ is called a *tau-function* of the KP hierarchy. The operator \mathcal{L} can be reconstructed from the tau-function τ by the following formula:

$$\text{res } \mathcal{L}^n = \varepsilon^{1-n} \frac{\partial^2 \log \tau}{\partial T_1 \partial T_n}, \quad n \geq 1.$$

Another important object associated to a solution of the KP hierarchy is the *wave function* (also called the Baker–Akhiezer function). Let P be the dressing operator of some operator \mathcal{L} satisfying the KP hierarchy (4.1) and let τ be the tau-function. Let

$$\xi(T_*, \varepsilon, z) := \sum_{k \geq 1} T_k \varepsilon^{k-1} z^k.$$

The wave function is defined by

$$w(T_*, \varepsilon, z) := \widehat{P} \cdot e^\xi = \frac{G_z(\tau)}{\tau} e^\xi \in \mathbb{C}[\varepsilon, \varepsilon^{-1}][[T_1, T_2, \dots]][[z, z^{-1}]].$$

It satisfies the equations

$$(4.4) \quad \frac{\partial w}{\partial T_n} = \varepsilon^{n-1} (\mathcal{L}^n)_+ w, \quad n \geq 1.$$

4.2. Gelfand–Dickey reduction and the closed r -spin partition function. Let $r \geq 2$. It is easy to see that the equation

$$(4.5) \quad (\mathcal{L}^r)_- = 0$$

is invariant with respect to the flows of the KP hierarchy (4.1). Therefore, it defines a reduction of the KP hierarchy that is called the *r -th Gelfand–Dickey hierarchy*. Let

$$L := \mathcal{L}^r = \partial_x^r + \sum_{i=0}^{r-2} f_i \partial_x^i.$$

Then all the coefficients u_i of the operator L can be expressed in terms of the functions f_0, f_1, \dots, f_{r-2} , and the Gelfand–Dickey hierarchy can be written as the following system of equations:

$$(4.6) \quad \frac{\partial L}{\partial T_n} = \varepsilon^{n-1} [(L^{n/r})_+, L], \quad n \geq 1.$$

Clearly, $\frac{\partial L}{\partial T_{dr}} = 0$ for any $d \geq 1$.

Suppose \mathcal{L} is a solution of the KP hierarchy satisfying the property (4.5). Then the dressing operator P and the tau-function τ can be chosen in such a way that $\frac{\partial P}{\partial T_{dr}} = 0$ and $\frac{\partial \tau}{\partial T_{dr}} = 0$ for any $d \geq 1$. In this case, the function τ is called a *tau-function* of the Gelfand–Dickey hierarchy.

Consider the generating series $F_r^{\frac{1}{r}, c}(t_*, \varepsilon)$ of the closed r -spin intersection numbers,

$$F_r^{\frac{1}{r}, c}(t_*, \varepsilon) = \sum_{\substack{g \geq 0, n \geq 1 \\ 2g-2+n > 0}} \frac{\varepsilon^{2g-2}}{n!} \sum_{\substack{0 \leq \alpha_1, \dots, \alpha_n \leq r-2 \\ d_1, \dots, d_n \geq 0}} \langle \tau_{d_1}^{\alpha_1} \dots \tau_{d_n}^{\alpha_n} \rangle_g^{\frac{1}{r}} t_{d_1}^{\alpha_1} \dots t_{d_n}^{\alpha_n}.$$

In [14], it is proven that the exponent $\tau_r^{\frac{1}{r},c} := e^{F_r^{\frac{1}{r},c}}$ becomes a tau-function of the Gelfand–Dickey hierarchy after the change of variables

$$(4.7) \quad T_k = \frac{1}{(-r)^{\frac{3k}{2(r+1)} - \frac{1}{2} - d} k!_r} t_d^\alpha, \quad 0 \leq \alpha \leq r-2, \quad d \geq 0,$$

where $k = \alpha + 1 + rd$ and

$$k!_r := \prod_{i=0}^d (\alpha + 1 + ri).$$

The corresponding solution L of the Gelfand–Dickey system (4.6) satisfies the initial condition

$$(4.8) \quad L|_{T_{\geq 2}=0} = \partial_x^r + \varepsilon^{-r} r x.$$

Let us prove a simple lemma describing the lowest-degree term in ε of the operator L .

Lemma 4.2. *Define a Poisson bracket $\{\cdot, \cdot\}$ in the ring $\mathbb{C}[[T_*]][z]$ by*

$$\begin{aligned} \{z, f\} &:= \partial_x f, & f &\in \mathbb{C}[[T_*]], \\ \{f_1, f_2\} &= \{z, z\} := 0, & f_1, f_2 &\in \mathbb{C}[[T_*]]. \end{aligned}$$

Then:

(a) *The functions f_i for $0 \leq i \leq r-2$ have the form*

$$f_i = \sum_{g \geq 0} f_i^{[g]} \varepsilon^{i-r+g}, \quad f_i^{[g]} \in \mathbb{C}[[T_*]].$$

(b) *Denote*

$$L_0 := \partial_x^r + \sum_{i=0}^{r-2} f_i^{[0]} \partial_x^i.$$

Then the operator L_0 is uniquely determined by the equations

$$\text{res } \widehat{L}_0^{n/r} = \frac{\partial^2 F_0^{1/r,c}}{\partial T_1 \partial T_n}, \quad 1 \leq n \leq r-1.$$

(c) *We have*

$$\frac{\partial \widehat{L}_0}{\partial T_a} = \left\{ \left(\widehat{L}_0^{\frac{a}{r}} \right)_+, \widehat{L}_0 \right\}.$$

Proof. Introduce the notation

$$w_i^{(j)} := \varepsilon^{1-i} \partial_x^j \frac{\partial^2 F_r^{\frac{1}{r},c}}{\partial T_1 \partial T_i}, \quad 1 \leq i \leq r-1, \quad j \geq 0,$$

$$f_i^{(j)} := \partial_x^j f_i, \quad 0 \leq i \leq r-2, \quad j \geq 0.$$

Then it is easy to see that

$$(4.9) \quad w_i = \frac{i}{r} f_{r-i-1} + P_i(f_{r-i}^{(*)}, \dots, f_{r-2}^{(*)}),$$

where P_i is a polynomial in $f_{r-i}^{(*)}, \dots, f_{r-2}^{(*)}$. Moreover, if we assign to $f_k^{(j)}$ degree $r-k+j$, then the polynomial P_i is homogeneous of degree $i+1$. The transformation (4.9) is clearly invertible, so we have

$$(4.10) \quad f_i = \frac{r}{i} w_{r-i-1} + Q_i(w_1^{(*)}, \dots, w_{r-i-2}^{(*)}),$$

where Q_i is a homogeneous polynomial of degree $r-i$ if we assign to $w_j^{(k)}$ degree $j+1+k$. It remains to note that

$$w_i = \varepsilon^{-i-1} \frac{\partial^2 F_0^{1/r,c}}{\partial T_1 \partial T_i} + O(\varepsilon^{-i})$$

and parts (a) and (b) of the lemma become clear.

Let us prove part (c), again using the homogeneity argument. We have

$$\left[\left(L^{\frac{a}{r}} \right)_+, L \right] = \sum_{i=0}^{r-2} R_i(f_*^{(*)}) \partial_x^i,$$

where a polynomial R_i has degree $a + r - i$. Let us assign to $f_i^{(j)}$ differential degree j and express a polynomial R_i as

$$R_i(f_*^{(*)}) = \sum_{j \geq 0} R_{i,j}(f_*^{(*)}),$$

where a polynomial $R_{i,j}$ has differential degree j . Clearly, $R_{i,0} = 0$ and for $R_{i,1}$ we have the formula

$$\sum_{i=0}^{r-2} R_{i,1} z^i = \left\{ \left(\widehat{L}^{\frac{a}{r}} \right)_+, \widehat{L} \right\}.$$

We have $R_{i,j} = O(\varepsilon^{i-r-a+2})$, for $j \geq 2$, and

$$R_{i,1} = \varepsilon^{i-r-a+1} \widetilde{R}_i(T_*) + O(\varepsilon^{i-r-a+2}),$$

where

$$\sum_{i=0}^{r-2} \widetilde{R}_i(T_*) z^i = \left\{ \left(\widehat{L}_0^{\frac{a}{r}} \right)_+, \widehat{L}_0 \right\}.$$

Part (c) of the lemma is proved. \square

4.3. Special solution of the equations for the wave function. Let L be the solution of the Gelfand–Dickey hierarchy (4.6) associated to the tau-function $\tau^{\frac{1}{r},c}$. Let $\Phi(T_*, \varepsilon)$ be the unique solution of the system of equations

$$(4.11) \quad \frac{\partial \Phi}{\partial T_n} = \varepsilon^{n-1} (L^{n/r})_+ \Phi$$

that satisfies the initial condition

$$\Phi|_{T_{\geq 2}=0} = 1.$$

Remark 4.3. In the case $r = 2$, the function Φ was first considered in [4], and the first author proved there that the logarithm of Φ coincides with the generating series of the intersection numbers on the moduli space of Riemann surfaces with boundary. Properties of the function Φ for general r were first studied in [2]. Note that the system of equations (4.11) for the function Φ coincides with the system of equations (4.4) for the wave function w of the KP hierarchy. In [2], the authors found an explicit formula for Φ in terms of the wave function w .

Denote $\phi := \log \Phi$ and consider the expansion

$$\phi = \sum_{g \in \mathbb{Z}} \varepsilon^{g-1} \phi_g, \quad \phi_g \in \mathbb{C}[[T_*]].$$

Let

$$(4.12) \quad T_{mr} = \frac{1}{(-r)^{\frac{m(r-2)}{2(r+1)}} m! r^m} t_{m-1}^{r-1}, \quad m \geq 1.$$

Define correlators $\langle \tau_{d_1}^{\alpha_1} \cdots \tau_{d_n}^{\alpha_n} \rangle_g^\phi$ by

$$\langle \tau_{d_1}^{\alpha_1} \cdots \tau_{d_n}^{\alpha_n} \rangle_g^\phi := \frac{\partial^n \phi_g}{\partial t_{d_1}^{\alpha_1} \cdots \partial t_{d_n}^{\alpha_n}} \Big|_{t_*^*=0}, \quad 0 \leq \alpha_1, \dots, \alpha_n \leq r-1, \quad d_1, \dots, d_n \geq 0.$$

Lemma 4.4. *The correlator $\langle \tau_{d_1}^{\alpha_1} \cdots \tau_{d_n}^{\alpha_n} \rangle_g^\phi$ can be non-zero only if*

$$(4.13) \quad g \geq 0,$$

$$(4.14) \quad \sum_{i=1}^n \left(\frac{\alpha_i}{r} + d_i - 1 \right) = \frac{(r+1)(g-1)}{r}.$$

Proof. Let us prove property (4.13). The function ϕ satisfies the following equations:

$$\frac{\partial \phi}{\partial T_n} = \varepsilon^{n-1} \frac{(L_{\frac{n}{r}})_+ e^\phi}{e^\phi}.$$

Therefore, it is sufficient to prove that for any function $\theta \in \mathbb{C}[\varepsilon, \varepsilon^{-1}][[T_*]]$ such that $\theta = O(\varepsilon^{-1})$, we have $\varepsilon^{n-1} \frac{(L_{\frac{n}{r}})_+ e^\theta}{e^\theta} = O(\varepsilon^{-1})$. From Lemma 4.2 it follows that the operator $L_{\frac{n}{r}}$ has the form $L_{\frac{n}{r}} = \sum_{i \leq n} R_i \partial_x^i$, $R_i \in \mathbb{C}[\varepsilon, \varepsilon^{-1}][[T_*]]$, where $R_i = O(\varepsilon^{i-n})$. By induction, it is easy to prove that

$$(4.15) \quad \frac{\partial_x^i e^\theta}{e^\theta} = i! \sum_{\substack{m_1, m_2, \dots \geq 0 \\ \sum j m_j = i}} \prod_{j \geq 1} \frac{(\partial_x^j \theta)^{m_j}}{(j!)^{m_j} m_j!}, \quad i \geq 0.$$

Since $\theta = O(\varepsilon^{-1})$, we have $\prod_{j \geq 1} (\partial_x^j \theta)^{m_j} = O(\varepsilon^{-i})$. Therefore, $R_i \frac{\partial_x^i e^\theta}{e^\theta} = O(\varepsilon^{-n})$ and

$$\varepsilon^{n-1} \sum_{i=0}^n R_i \frac{\partial_x^i e^\theta}{e^\theta} = O(\varepsilon^{-1}).$$

Property (4.13) is proved.

Let us prove property (4.14). Consider the linear differential operator

$$O := \sum_{\substack{0 \leq \alpha \leq r-1 \\ d \geq 0}} \left(\frac{\alpha}{r} + d - 1 \right) t_d^\alpha \frac{\partial}{\partial t_d^\alpha} - \frac{r+1}{r} \varepsilon \frac{\partial}{\partial \varepsilon}.$$

We have to prove that $O\phi = 0$ or, equivalently, $O\Phi = 0$. Using the Gelfand–Dickey equations (4.6), it is easy to show that

$$\left(z \frac{\partial}{\partial z} + O \right) \widehat{L} = r \widehat{L}.$$

Then, similarly, using equations (4.11) one can show that $O\Phi = 0$. The lemma is proved. \square

Lemma 4.5. *The function ϕ satisfies the string equation*

$$(4.16) \quad \left(\frac{\partial}{\partial T_1} - \sum_{i \geq 1} (i+r) T_{i+r} \frac{\partial}{\partial T_i} \right) \phi = r \varepsilon^{-1} T_r.$$

Proof. Equation (4.16) was proved in [2], but in order to make the paper more self-contained let us give a proof here. Denote by O the operator in the brackets on the left-hand side of (4.16). Using the Gelfand–Dickey equations (4.6), it is easy to show that $OL = \varepsilon^{-r} r$. Then, similarly, using equations (4.11) one can show that $O\Phi = (r \varepsilon^{-1} T_r) \Phi$. \square

4.4. Closed extended potential and the special solution. Consider the generating series $F_0^{\frac{1}{r}, \text{ext}}(t_*)$ of the closed extended r -spin intersection numbers in genus zero,

$$F_0^{\frac{1}{r}, \text{ext}}(t_*) = \sum_{n \geq 2} \frac{1}{n!} \sum_{\substack{0 \leq \alpha_1, \dots, \alpha_n \leq r-1 \\ d_1, \dots, d_n \geq 0}} \langle \tau_{d_1}^{\alpha_1} \cdots \tau_{d_n}^{\alpha_n} \tau_0^{-1} \rangle_0^{\frac{1}{r}, \text{ext}} t_{d_1}^{\alpha_1} \cdots t_{d_n}^{\alpha_n}.$$

The main result of our paper is the following theorem.

Theorem 4.6. *We have*

$$F_0^{\frac{1}{r}, \text{ext}}(t_*^{\leq r-2}, t_*^{r-1}) = \sqrt{-r} \phi_0 \left(t_*^{\leq r-2}, \frac{1}{\sqrt{-r}} t_*^{r-1} \right).$$

Before proving the theorem, let us formulate several auxiliary statements.

Lemma 4.7. *The function ϕ_0 satisfies the equations*

$$(4.17) \quad \frac{\partial \phi_0}{\partial T_n} = (\widehat{L}_0^n)_+ \Big|_{z=(\phi_0)_x}, \quad n \geq 1.$$

Proof. From Lemma 4.2 it follows that the operator $L^{\frac{n}{r}}$ has the form $L^{\frac{n}{r}} = \sum_{i \leq n} R_i \partial_x^i$, where $R_i = \sum_{j \geq 0} R_{i,j} \varepsilon^{i-n+j}$, $R_{i,j} \in \mathbb{C}[[T_*]]$. It is also clear that $\sum_{i \leq n} R_{i,0} z^i = \widehat{L}_0^{\frac{n}{r}}$. We have

$$\frac{\partial \phi}{\partial T_n} = \varepsilon^{n-1} \sum_{i=0}^n R_i \frac{\partial_x^i e^\phi}{e^\phi}.$$

Since $\phi = \varepsilon^{-1} \phi_0 + O(\varepsilon^0)$, for any numbers $m_1, m_2, \dots \geq 0$, satisfying $\sum j m_j = i$, we have

$$\prod_{j \geq 1} (\partial_x^j \phi)^{m_j} = \begin{cases} \varepsilon^{-i} (\phi_0)_x^i + O(\varepsilon^{-i+1}), & \text{if } m_1 = i \text{ and } m_2 = m_3 = \dots = 0, \\ O(\varepsilon^{-i+1}), & \text{otherwise.} \end{cases}$$

Using formula (4.15) we then get $R_i \frac{\partial_x^i e^\phi}{e^\phi} = R_{i,0} \varepsilon^{-n} (\phi_0)_x^i + O(\varepsilon^{-n+1})$ and

$$\varepsilon^{n-1} \sum_{i=0}^n R_i \frac{\partial_x^i e^\phi}{e^\phi} = \varepsilon^{-1} \sum_{i=0}^n R_{i,0} (\phi_0)_x^i + O(\varepsilon^0) = \varepsilon^{-1} (\widehat{L}_0^{\frac{n}{r}})_+ \Big|_{z=(\phi_0)_x} + O(\varepsilon^0).$$

The lemma is proved. \square

Proposition 4.8. *The function ϕ_0 satisfies the equations*

$$(4.18) \quad \frac{\partial^2 \phi_0}{\partial t_{p+1}^\alpha \partial t_q^\beta} = \sum_{\mu+\nu=r-2} \frac{\partial^2 F_0^{\frac{1}{r},c}}{\partial t_p^\alpha \partial t_0^\mu} \frac{\partial^2 \phi_0}{\partial t_0^\nu \partial t_q^\beta} + \frac{\partial \phi_0}{\partial t_p^\alpha} \frac{\partial^2 \phi_0}{\partial t_0^{r-1} \partial t_q^\beta}, \quad 0 \leq \alpha, \beta \leq r-1, \quad p, q \geq 0.$$

Proof. In the variables T_i equations (4.18) look as follows:

$$d \left(\frac{\partial \phi_0}{\partial T_{a+r}} \right) = \sum_{b=1}^{r-1} \frac{a+r}{b(r-b)} \frac{\partial^2 F_0^{\frac{1}{r},c}}{\partial T_a \partial T_b} d \left(\frac{\partial \phi_0}{\partial T_{r-b}} \right) + \frac{a+r}{r} \frac{\partial \phi_0}{\partial T_a} d \left(\frac{\partial \phi_0}{\partial T_r} \right), \quad a \geq 1.$$

By Lemma 4.7, this equation follows from

$$(4.19) \quad d \left(\widehat{L}_0^{\frac{a+r}{r}} \right)_+ = \sum_{b=1}^{r-1} \frac{a+r}{b(r-b)} \frac{\partial^2 F_0^{\frac{1}{r},c}}{\partial T_a \partial T_b} d \left(\widehat{L}_0^{\frac{r-b}{r}} \right)_+ + \frac{a+r}{r} \left(\widehat{L}_0^{\frac{a}{r}} \right)_+ d \widehat{L}_0, \quad a \geq 1.$$

Let

$$v_i := \text{res} \left(\widehat{L}_0^{\frac{i}{r}} \right), \quad 1 \leq i \leq r-1.$$

Clearly, the functions $f_0^{[0]}, \dots, f_{r-2}^{[0]}$ and v_1, \dots, v_{r-1} are related by an invertible polynomial transformation. Therefore, for any $i \in \mathbb{Z}$, the coefficients of the Laurent series \widehat{L}_0^i can be considered as polynomials in v_1, \dots, v_{r-1} . For $a \geq 1$ and $1 \leq b \leq r-1$ we have the following identity (see, for example, [13, Sections 3.5 and 3.6]):

$$\frac{\partial^2 F_0^{\frac{1}{r},c}}{\partial T_a \partial T_b} = \frac{b(r-b)}{a+r} \frac{\partial}{\partial v_{r-b}} \text{res} \widehat{L}_0^{\frac{a+r}{r}}.$$

We see that equation (4.19) is equivalent to

$$(4.20) \quad d \left(\widehat{L}_0^{\frac{a+r}{r}} \right)_+ = \sum_{b=1}^{r-1} \frac{\partial}{\partial v_b} \text{res} \left(\widehat{L}_0^{\frac{a+r}{r}} \right) d \left(\widehat{L}_0^{\frac{b}{r}} \right)_+ + \frac{a+r}{r} \left(\widehat{L}_0^{\frac{a}{r}} \right)_+ d \widehat{L}_0.$$

Let us express the left-hand side in the following way:

$$d\left(\widehat{L}_0^{\frac{a+r}{r}}\right)_+ = \frac{a+r}{r} \left(\widehat{L}_0^{\frac{a}{r}}\right)_+ \underline{d\widehat{L}_0} + \frac{a+r}{r} \left(\left(\widehat{L}_0^{\frac{a}{r}}\right)_- d\widehat{L}_0\right)_+.$$

Notice that the underlined term here cancels the underlined term on the right-hand side of (4.20). Therefore, equation (4.20) is equivalent to

$$(4.21) \quad \frac{a+r}{r} \left(\left(\widehat{L}_0^{\frac{a}{r}}\right)_- d\widehat{L}_0\right)_+ = \sum_{b=1}^{r-1} \frac{\partial}{\partial v_b} \operatorname{res} \left(\widehat{L}_0^{\frac{a+r}{r}}\right) d\left(\widehat{L}_0^{\frac{b}{r}}\right)_+.$$

We compute

$$\sum_{b=1}^{r-1} \frac{\partial}{\partial v_b} \operatorname{res} \left(\widehat{L}_0^{\frac{a+r}{r}}\right) d\left(\widehat{L}_0^{\frac{b}{r}}\right)_+ = \sum_{b=1}^{r-1} \frac{b}{r} \frac{\partial}{\partial v_b} \operatorname{res} \left(\widehat{L}_0^{\frac{a+r}{r}}\right) \left(\widehat{L}_0^{\frac{b-r}{r}} d\widehat{L}_0\right)_+.$$

We see that equation (4.21) follows from the property

$$(4.22) \quad \frac{a+r}{r} \left(\widehat{L}_0^{\frac{a}{r}}\right)_- - \sum_{b=1}^{r-1} \frac{b}{r} \frac{\partial}{\partial v_b} \operatorname{res} \left(\widehat{L}_0^{\frac{a+r}{r}}\right) \widehat{L}_0^{\frac{b-r}{r}} \in z^{-r-1} \mathbb{C}[f_0^{[0]}, \dots, f_{r-2}^{[0]}][[z^{-1}]].$$

Recall that (see, for example, [13, Section 3.5])

$$(4.23) \quad \frac{a+r}{r} \left(\widehat{L}_0^{\frac{a}{r}}\right)_- - \sum_{i=0}^{r-2} \frac{\partial}{\partial f_i^{[0]}} \operatorname{res} \left(\widehat{L}_0^{\frac{a+r}{r}}\right) z^{-i-1} \in z^{-r-1} \mathbb{C}[f_0^{[0]}, \dots, f_{r-2}^{[0]}][[z^{-1}]].$$

For any two elements $f, g \in \mathbb{C}[f_0^{[0]}, \dots, f_{r-2}^{[0]}][[z^{-1}]]$ let us write $f \equiv g$ if the difference $f - g$ lies in $z^{-r-1} \mathbb{C}[f_0^{[0]}, \dots, f_{r-2}^{[0]}][[z^{-1}]]$. Then, using identity (4.23), we can compute

$$\begin{aligned} \frac{a+r}{r} \left(\widehat{L}_0^{\frac{a}{r}}\right)_- &\equiv \sum_{i=0}^{r-2} \frac{\partial}{\partial f_i^{[0]}} \operatorname{res} \left(\widehat{L}_0^{\frac{a+r}{r}}\right) z^{-i-1} \equiv \sum_{b=1}^{r-1} \sum_{i=0}^{r-2} \frac{\partial}{\partial v_b} \operatorname{res} \left(\widehat{L}_0^{\frac{a+r}{r}}\right) \frac{\partial v_b}{\partial f_i^{[0]}} z^{-i-1} \equiv \\ &\equiv \sum_{b=1}^{r-1} \frac{b}{r} \frac{\partial}{\partial v_b} \operatorname{res} \left(\widehat{L}_0^{\frac{a+r}{r}}\right) \left(\widehat{L}_0^{\frac{b-r}{r}}\right)_- \equiv \sum_{b=1}^{r-1} \frac{b}{r} \frac{\partial}{\partial v_b} \operatorname{res} \left(\widehat{L}_0^{\frac{a+r}{r}}\right) \widehat{L}_0^{\frac{b-r}{r}}. \end{aligned}$$

Formula (4.22) is proved. This completes the proof of the proposition. \square

Lemma 4.9. *We have*

$$(4.24) \quad \langle \tau_0^1 \tau_0^{r-2} \rangle_0^\phi = \begin{cases} 1, & \text{if } r = 2, \\ \frac{1}{\sqrt{-r}}, & \text{if } r \geq 3, \end{cases} \quad \langle \tau_0^1 (\tau_0^{r-1})^2 \rangle_0^\phi = \begin{cases} 1, & \text{if } r = 2, \\ \frac{1}{\sqrt{-r}}, & \text{if } r \geq 3. \end{cases}$$

Proof. Let us prove the first equation in (4.24). We compute

$$\frac{\partial^2 \phi_0}{\partial T_2 \partial T_{r-1}} \Big|_{T_*=0} = \frac{\partial}{\partial T_{r-1}} \left((\phi_0)_x^2 + \frac{2}{r} f_{r-2}^{[0]} \right) \Big|_{T_*=0} = \frac{2}{r} \frac{\partial f_{r-2}^{[0]}}{\partial T_{r-1}} \Big|_{T_*=0}.$$

We proceed as follows:

$$\begin{aligned} \frac{\partial \widehat{L}_0}{\partial T_{r-1}} \Big|_{T_*=0} &= \left\{ \left(\widehat{L}_0^{\frac{r-1}{r}}\right)_+, \widehat{L}_0 \right\} \Big|_{T_*=0} = \left\{ \left((z^r + rx)^{\frac{r-1}{r}}\right)_+, z^r + rx \right\} \Big|_{x=0} \\ &= \{z^{r-1}, z^r + rx\} \Big|_{x=0} = r(r-1)z^{r-2}. \end{aligned}$$

Therefore, $\frac{\partial^2 \phi_0}{\partial T_2 \partial T_{r-1}} \Big|_{T_*=0} = 2(r-1)$. Applying changes of variables (4.7) and (4.12), we see that the first equation in (4.24) is true.

Now, let us prove the second equation in (4.24). First of all, we compute

$$\frac{\partial^2 \phi_0}{\partial T_1 \partial T_r} \Big|_{T_*=0} = \partial_x ((\phi_0)_x^r + rx) \Big|_{T_*=0} = r.$$

Therefore, we have

$$\left. \frac{\partial^3 \phi_0}{\partial T_2 \partial T_r^2} \right|_{T_*=0} = \left. \frac{\partial^2}{\partial T_r^2} \left((\phi_0)_x^2 + \frac{2}{r} f_{r-2}^{[0]} \right) \right|_{T_*=0} = 2 \left. \left(\frac{\partial^2 \phi_0}{\partial T_1 \partial T_r} \right)^2 \right|_{T_*=0} = 2r^2.$$

Using equations (4.7) and (4.12), we can see that the second equation in (4.24) is also true. \square

Proof of Theorem 4.6. We have seen (see Section 3.1 and Lemmas 3.6, 3.8) that the function $F_0^{\frac{1}{r}, \text{ext}}$ and the correlators $\langle \tau_0^{-1} \tau_{d_1}^{\alpha_1} \cdots \tau_{d_n}^{\alpha_n} \rangle_0^{\frac{1}{r}, \text{ext}}$ satisfy the following properties:

$$(4.25) \quad \langle \tau_0^{-1} \tau_{d_1}^{\alpha_1} \cdots \tau_{d_n}^{\alpha_n} \rangle_0^{\frac{1}{r}, \text{ext}} = 0 \quad \text{unless} \quad \frac{\sum \alpha_i - (r-1)}{r} + \sum d_i = n - 2,$$

$$(4.26) \quad \frac{\partial^2 F_0^{\frac{1}{r}, \text{ext}}}{\partial t_{p+1}^\alpha \partial t_q^\beta} = \sum_{\mu+\nu=r-2} \frac{\partial^2 F_0^{\frac{1}{r}, c}}{\partial t_p^\alpha \partial t_0^\mu} \frac{\partial^2 F_0^{\frac{1}{r}, \text{ext}}}{\partial t_0^\nu \partial t_q^\beta} + \frac{\partial F_0^{\frac{1}{r}, \text{ext}}}{\partial t_p^\alpha} \frac{\partial^2 F_0^{\frac{1}{r}, \text{ext}}}{\partial t_0^{r-1} \partial t_q^\beta}, \quad 0 \leq \alpha, \beta \leq r-1, \quad p, q \geq 0,$$

$$(4.27) \quad \langle \tau_0^{-1} \tau_0^1 \tau_0^{r-2} \rangle_0^{\frac{1}{r}, \text{ext}} = 1, \quad \langle \tau_0^{-1} \tau_0^1 (\tau_0^{r-1})^2 \rangle_0^{\frac{1}{r}, \text{ext}} = -\frac{1}{r}.$$

It is easy to see that for any non-zero complex constant C , the function $\tilde{\phi}_0$ defined by

$$\tilde{\phi}_0(t_*^{\leq r-2}, t_*^{r-1}) := C \phi_0(t_*^{\leq r-2}, C^{-1} t_*^{r-1})$$

also satisfies equations (4.18). Let $C = \sqrt{-r}$ and

$$\langle \tau_{d_1}^{\alpha_1} \cdots \tau_{d_n}^{\alpha_n} \rangle_0^{\tilde{\phi}} := \left. \frac{\partial^n \tilde{\phi}_0}{\partial t_{d_1}^{\alpha_1} \cdots \partial t_{d_n}^{\alpha_n}} \right|_{t_*^*=0}.$$

Lemmas 4.4, 4.9 and Proposition 4.8 imply that the function $\tilde{\phi}_0$ and the correlators $\langle \tau_{d_1}^{\alpha_1} \cdots \tau_{d_n}^{\alpha_n} \rangle_0^{\tilde{\phi}}$ also satisfy properties (4.25) – (4.27). Therefore, it is sufficient to check that these properties are enough to reconstruct the function $F_0^{\frac{1}{r}, \text{ext}}$.

Note that the dimension constraint in property (4.25) implies that $n \geq 2$. Therefore, by (4.26), it suffices to determine the primary correlators

$$(4.28) \quad \langle \tau_0^{-1} \tau_0^{\alpha_1} \cdots \tau_0^{\alpha_l} (\tau_0^{r-1})^k \rangle_0^{\frac{1}{r}, \text{ext}}, \quad 0 \leq \alpha_1, \dots, \alpha_l \leq r-2.$$

By (4.25), such a correlator can be non-zero only if

$$(4.29) \quad \sum_{i=1}^l (r - \alpha_i) + k = r + 1.$$

First of all, suppose that $l \leq 1$, so we consider the correlators

$$X_\alpha := \langle \tau_0^{-1} \tau_0^\alpha (\tau_0^{r-1})^{\alpha+1} \rangle_0^{\frac{1}{r}, \text{ext}}, \quad 0 \leq \alpha \leq r-1.$$

Suppose $0 \leq \gamma \leq r-2$ and consider the correlator

$$Y_\gamma := \langle \tau_0^{-1} \tau_1^1 \tau_0^\gamma (\tau_0^{r-1})^{\gamma+2} \rangle_0^{\frac{1}{r}, \text{ext}}.$$

Let us compute it using the topological recursion relation (4.26) in two different ways. Applying (4.26) with $\alpha = 1$, $p = q = 0$ and $\beta = \gamma$, we obtain

$$(4.30) \quad Y_\gamma = \binom{\gamma+2}{2} X_1 X_\gamma.$$

On the other hand, applying (4.26) with $\alpha = 1$, $p = q = 0$ and $\beta = r-1$, we get

$$(4.31) \quad Y_\gamma = \binom{\gamma+1}{2} X_1 X_\gamma + \langle \tau_0^1 \tau_0^\gamma \tau_0^{r-3-\gamma} \rangle_0^{\frac{1}{r}, \text{ext}} X_{\gamma+1}.$$

For $\gamma \leq r - 3$, we have $\langle \tau_0^1 \tau_0^\gamma \tau_0^{r-3-\gamma} \rangle_0^{\frac{1}{r}} = 1$ (see e.g. [20, Section 0.6]), and, by (4.27), $\langle \tau_0^1 \tau_0^{r-2} \tau_0^{-1} \rangle_0^{\frac{1}{r}, \text{ext}} = 1$. Therefore, equating the right-hand sides of (4.30) and (4.31), we obtain

$$X_{\gamma+1} = (\gamma + 1)X_1 X_\gamma, \quad 0 \leq \gamma \leq r - 2.$$

Since $X_1 = -\frac{1}{r}$, this immediately implies that $X_\alpha = (-1)^\alpha \frac{\alpha!}{r^\alpha}$, for any $0 \leq \alpha \leq r - 1$.

Consider now the correlator (4.28) with $l \geq 2$. Suppose that condition (4.29) is satisfied. Recall that by $[l]$ we denote the set $\{1, 2, \dots, l\}$. For a subset $I \subset [l]$, let

$$m_I := r + 1 - \sum_{i \in I} (r - \alpha_i), \quad A_I := \left\langle \tau_0^{-1} \left(\prod_{i \in I} \tau_0^{\alpha_i} \right) (\tau_0^{r-1})^{m_I} \right\rangle_0^{\frac{1}{r}, \text{ext}}.$$

Consider the correlator

$$(4.32) \quad Z := \left\langle \tau_0^{-1} \tau_1^{\alpha_1} \left(\prod_{i=2}^l \tau_0^{\alpha_i} \right) (\tau_0^{r-1})^{k+r} \right\rangle_0^{\frac{1}{r}, \text{ext}}.$$

Applying (4.26) with $\alpha = \alpha_1$, $p = q = 0$ and $\beta = \alpha_l$, we get

$$(4.33) \quad Z = \sum_{\substack{I \sqcup J = [l] \\ 1 \in I, l \in J}} \sum_{\mu + \nu = r - 2} \left\langle \left(\prod_{i \in I} \tau_0^{\alpha_i} \right) \tau_0^\mu \right\rangle_0^{\frac{1}{r}} \left\langle \tau_0^{-1} \tau_0^\nu \left(\prod_{j \in J} \tau_0^{\alpha_j} \right) (\tau_0^{r-1})^{k+r} \right\rangle_0^{\frac{1}{r}, \text{ext}} + \sum_{\substack{I \sqcup J = [l] \\ 1 \in I, l \in J}} \binom{r+k}{m_I} A_I A_J.$$

Note that the underlined term on the right-hand side of this equation vanishes, because otherwise we should have

$$\begin{aligned} r - \nu + \sum_{j \in J} (r - \alpha_j) + k + r = r + 1 &\Rightarrow \sum_{i \in I} (r - \alpha_i) = r + (r - \nu) \Rightarrow \\ &\Rightarrow \sum_{i \in I} (r - \alpha_i) \geq r + 2 \Rightarrow \sum_{i=1}^l (r - \alpha_i) \geq r + 2, \end{aligned}$$

which contradicts (4.29). On the other hand, applying to the correlator (4.32) relation (4.26) with $\alpha = \alpha_1$, $p = q = 0$, and $\beta = r - 1$, we obtain

$$(4.34) \quad \begin{aligned} Z &= \sum_{\substack{I \sqcup J = [l] \\ 1 \in I}} \binom{r+k-1}{m_I} A_I A_J = \\ &= \sum_{\substack{I \sqcup J = [l] \\ 1 \in I, l \in J}} \binom{r+k-1}{m_I} A_I A_J + \sum_{\substack{I \sqcup J = [l] \\ 1, l \in I, J \neq \emptyset}} \binom{r+k-1}{m_I} A_I A_J + \binom{r+k-1}{k} A_{[l]} \langle \tau_0^{-1} (\tau_0^{r-1})^{r+1} \rangle_0^{\frac{1}{r}, \text{ext}}. \end{aligned}$$

Note that here, by the same argument as above, we also do not have terms with closed r -spin correlators. Equating the right-hand side of (4.33) and expression (4.34) and using that $\langle \tau_0^{-1} (\tau_0^{r-1})^{r+1} \rangle_0^{\frac{1}{r}, \text{ext}} = (-1)^{r-1} \frac{(r-1)!}{r^{r-1}}$, we get

$$(4.35) \quad (-1)^{r-1} \frac{(r+k-1)!}{k! r^{r-1}} A_{[l]} = \sum_{\substack{I \sqcup J = [l] \\ 1 \in I, l \in J}} \binom{r+k-1}{m_I - 1} A_I A_J - \sum_{\substack{I \sqcup J = [l] \\ 1, l \in I, J \neq \emptyset}} \binom{r+k-1}{m_I} A_I A_J.$$

We see that for $l \geq 2$, this equation allows one to compute the primary correlator (4.28) in terms of primary correlators with smaller l . The theorem is proved. \square

4.5. Lax operator and the closed extended potential. Let us show that the operator $L_0|_{t_{\geq 1}^*=0}$ has a simple interpretation in terms of the function $F_0^{\frac{1}{r},\text{ext}}$. Define the primary closed extended potential in genus 0 by

$$\mathcal{F}_0^{\frac{1}{r},\text{ext}}(t_0^0, \dots, t_0^{r-1}) := F_0^{\frac{1}{r},\text{ext}} \Big|_{t_{\geq 1}^*=0}.$$

Proposition 4.10. *We have*

$$(4.36) \quad \frac{\partial \mathcal{F}_0^{\frac{1}{r},\text{ext}}}{\partial t_0^{r-1}} = \frac{1}{(-r)^{\frac{r-2}{2(r+1)}r}} \widehat{L}_0 \Big|_{z=(-r)^{\frac{1-2r}{2(r+1)}} t_0^{r-1}, t_{\geq 1}^*=0}.$$

Proof. Using Theorem 4.6 we see that equation (4.36) is equivalent to the equation

$$\frac{\partial \phi_0}{\partial T_r} \Big|_{T_{\geq r+1}=0} = \widehat{L}_0 \Big|_{z=rT_r, T_{\geq r+1}=0},$$

that follows from Lemma 4.7 and the fact that, by the string equation (4.16), $(\phi_0)_x|_{T_{\geq r+1}=0} = rT_r$. \square

4.6. Main conjecture. We conjecture that for any $b \geq 1$, $g, n \geq 0$, $0 \leq \alpha_1, \dots, \alpha_n \leq r-1$, and $d_1, \dots, d_n \geq 0$, there is a geometric construction of correlators

$$\langle \tau_{d_1}^{\alpha_1} \dots \tau_{d_n}^{\alpha_n} (\tau_0^{-1})^b \rangle_g^{\frac{1}{r},\text{ext}},$$

generalizing the construction for $b = 1$ and $g = 0$. Given such correlators, define a generating series $F_g^{\frac{1}{r},\text{ext}}(t_*)$ for any $g \geq 0$ by

$$F_g^{\frac{1}{r},\text{ext}}(t_*) := \sum_{\substack{h \geq 0, b \geq 1 \\ 2h+b-1=g}} \sum_{n \geq 0} \frac{1}{b!n!} \sum_{\substack{0 \leq \alpha_1, \dots, \alpha_n \leq r-1 \\ d_1, \dots, d_n \geq 0}} \langle \tau_{d_1}^{\alpha_1} \dots \tau_{d_n}^{\alpha_n} (\tau_0^{-1})^b \rangle_h^{\frac{1}{r},\text{ext}} t_{d_1}^{\alpha_1} \dots t_{d_n}^{\alpha_n}.$$

Conjecture 4.11. *For any $g \geq 0$,*

$$F_g^{\frac{1}{r},\text{ext}} = (-r)^{\frac{1-g}{2}} \phi_g \left(t_*^{\leq r-2}, \frac{1}{\sqrt{-r}} t_*^{r-1} \right).$$

5. OPEN–CLOSED CORRESPONDENCE

In the forthcoming work [5], the authors generalize the definition of the moduli space of r -spin structures and its Witten class to the setting of genus-zero surfaces with boundary, or *disks*. We summarize the construction of [5] in this section and explain the connection between open and closed extended theory.

5.1. Open r -spin theory. A Riemann surface with boundary is a tuple $(C, \phi, \Sigma, \{z_i\}, \{x_j\})$, where C is an orbifold curve equipped with an involution $\phi : C \rightarrow C$ that realizes $|C|$ as a union of two copies of the Riemann surface with boundary Σ , glued along their common boundary; we write

$$|C| = \Sigma \cup_{\partial\Sigma} \bar{\Sigma}.$$

Here, $z_1, \dots, z_n \in C$ are the *internal marked points* (whose images in $|C|$ lie in $\Sigma \setminus \partial\Sigma$, and each of which has a partner $\bar{z}_i := \phi(z_i)$ whose image lies in $\bar{\Sigma}$) and $x_1, \dots, x_m \in \partial\Sigma$ are the *boundary marked points*. We define a *graded r -spin structure* on a Riemann surface with boundary as an orbifold line bundle S on C together with an isomorphism

$$S^{\otimes r} \cong \omega_C \otimes \mathcal{O} \left(- \sum_{i=1}^n \alpha_i [z_i] - \sum_{i=1}^n \alpha_i [\bar{z}_i] - \sum_{j=1}^m (r-2) [x_j] \right),$$

an involution $\tilde{\phi} : S \rightarrow S$ lifting ϕ , and an additional structure that we refer to as a *grading*. (Roughly, a grading is an involution-invariant section of S on the complement of the special

points in $\partial\Sigma$ that changes sign at each boundary marked point, this notion was first defined for $r = 2$ in [23].) Here, we assume that the internal twists $\alpha_1, \dots, \alpha_n$ lie in the range $\{0, 1, \dots, r-1\}$. Note that we can re-express the genus in terms of the number b of boundary components of Σ and the genus g_c of the closed surface obtained from Σ by gluing a disk to each boundary component:

$$g = b + g_c - 1.$$

For example, a disk has $g = g_c = 0$ and $b = 1$.

In [5] we construct the moduli space $\overline{\mathcal{M}}_{0,m,\{\alpha_1,\dots,\alpha_n\}}^{1/r}$ parameterizing graded r -spin disks, and prove that it is nonempty exactly when a certain congruence condition on the twists α_i is satisfied. There should be no difficulty with constructing the all-genus generalization $\overline{\mathcal{M}}_{g,m,\{\alpha_1,\dots,\alpha_n\}}^{1/r}$ and this space is nonempty if and only if

$$(5.1) \quad e_o := \frac{(g+m-1)(r-2) + 2\sum_{i=1}^n \alpha_i}{r} \in \mathbb{N} \quad \text{and} \quad e_o \equiv 1 + m + g \pmod{2}.$$

There are line bundles \mathbb{L}_i for each $i = 1, \dots, n$, defined by the cotangent line to the orbifold curve C at the i th internal marked point. Furthermore, in genus zero, there is a real analogue of Witten's bundle,

$$\mathcal{W} := (R^0\pi_*(\mathcal{S}^\vee \otimes \omega_\pi))_+,$$

a real-rank- e_o bundle over $\overline{\mathcal{M}}_{0,m,\{\alpha_1,\dots,\alpha_n\}}^{1/r}$ whose fiber over $(C, \phi, \Sigma, \{z_i\}, \{x_j\}, S, \tilde{\phi})$ consists of $\tilde{\phi}$ -invariant sections of $\mathcal{S}^\vee \otimes \omega_C$.

Because $\overline{\mathcal{M}}_{0,m,\{\alpha_1,\dots,\alpha_n\}}^{1/r}$ has boundary, one must work with a relative version of the Chern classes of the cotangent line bundles and the Witten bundles. In [5] the requisite canonical boundary conditions are defined over the space $\overline{\mathcal{PM}}_{0,m,\{\alpha_1,\dots,\alpha_n\}}^{1/r}$, which is some canonical perturbation of the space $\overline{\mathcal{M}}_{0,m,\{\alpha_1,\dots,\alpha_n\}}^{1/r}$. From here, we define open r -spin correlators by

$$(5.2) \quad \left\langle \prod_{i=1}^n \tau_{d_i}^{\alpha_i} \sigma^m \right\rangle_0^{\frac{1}{r}, o} := \int_{\overline{\mathcal{PM}}_{0,m,\{\alpha_1,\dots,\alpha_n\}}^{1/r}} e \left(\mathcal{W} \oplus \bigoplus_{i=1}^n \mathbb{L}_i^{\oplus d_i}, s_{\text{canonical}} \right),$$

where $e(E, s)$ is the Euler class relative to the canonical boundary conditions s . Alternatively, the number in (5.2) can be defined as a weighted, signed count of the number of zeroes of a generic extension of $s_{\text{canonical}}$, defining this count to be zero unless $e_o + 2\sum_{i=1}^n d_i = m + 2n - 3 + 3g$, or in other words, unless

$$(5.3) \quad e_o + 2\sum_{i=1}^n d_i = m + 2n - 6 + 3g_c + 3b.$$

One of the key results of [5] is that these intersection numbers are independent of the specific choice of $s_{\text{canonical}}$.

We define a generating function for genus-zero open r -spin theory by

$$F_0^{\frac{1}{r}, o}(t_*, s) := \sum_{\substack{n, m \geq 0 \\ 2n+m-2 > 0}} \sum_{\substack{0 \leq \alpha_1, \dots, \alpha_n \leq r-1 \\ d_1, \dots, d_n \geq 0}} \frac{1}{n!m!} \left\langle \prod_{i=1}^n \tau_{d_i}^{\alpha_i} \sigma^m \right\rangle_0^{\frac{1}{r}, o} t_{d_1}^{\alpha_1} \dots t_{d_n}^{\alpha_n} s^m.$$

Remark 5.1. We caution the reader, that, similarly to the closed extended theory, the Ramond vanishing property doesn't hold for the open r -spin correlators.

Remark 5.2. In the case where $r = 2$ and all of the insertions are Neveu–Schwarz, open r -spin theory is equivalent to the intersection theory of disks constructed by Pandharipande, Solomon, and the third author in [21]. This is currently the only case in which we can extend the theory to higher genus [24], calculate all numbers [25], and prove the relationship to the wave function in all genus [4, 6].

5.2. Connection to closed extended theory. In [5, Theorem 1.3], we prove that the genus-zero open r -spin potential is related to the closed extended potential in the following way:

$$F_0^{\frac{1}{r},o}(t_*^0, t_*^1, \dots, t_*^{r-1}, s) = -\frac{1}{r} F_0^{\frac{1}{r},\text{ext}} \Big|_{t_d^{r-1} \mapsto t_d^{r-1-r\delta_{d,0}s}} + \frac{1}{r} F_0^{\frac{1}{r},\text{ext}}.$$

We currently do not know of a geometric explanation for the intimate relation between these two theories. Nevertheless, let us explore more explicitly what is known.

Heuristically, the dictionary between closed extended and open r -spin theory is given by

- (a) replacing a marked point with twist -1 by a boundary component, and
- (b) replacing a marked point with twist $r-1$ by a boundary marked point.

To make this more precise, we first observe that these exchanges are compatible with the rank-dimension constraints for the two theories. That is, replacing every boundary component in the open theory with a marked point of twist -1 converts equation (5.3) into equation (2.5), and replacing an internal marked point of twist $r-1$ by a boundary marked point leaves (5.3) invariant.¹ Moreover, at the level of intersection numbers, we have the relation

$$\left\langle \prod_{i=1}^n \tau_{d_i}^{\alpha_i} \sigma^m \right\rangle_0^{\frac{1}{r},o} = \begin{cases} 0, & \text{if } m = 0, \\ (-r)^{m-1} \left\langle \tau_0^{-1} \prod_{i=1}^n \tau_{d_i}^{\alpha_i} (\tau_0^{r-1})^m \right\rangle_0^{\frac{1}{r},\text{ext}} & \text{if } m \geq 1, \end{cases}$$

which realizes the above dictionary when there is a single boundary component.

Moreover, this dictionary matches the topological recursion relations in genus zero. In [5], we prove two topological recursion relations for open r -spin theory. First, for any $i \in [n]$ with $d_i > 0$ and any $j \in ([n] \setminus \{i\})$, we have

$$(5.4) \quad \left\langle \prod_{l \in [n]} \tau_{d_l}^{\alpha_l} \sigma^m \right\rangle_0^{\frac{1}{r},o} = \sum_{I \amalg J = [n] \setminus \{i\}} \sum_{\alpha=-1}^{r-2} \left\langle \tau_0^\alpha \tau_{d_{i-1}}^{\alpha_i} \prod_{l \in I} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r},\text{ext}} \left\langle \tau_0^{r-2-\alpha} \prod_{l \in J} \tau_{d_l}^{\alpha_l} \sigma^m \right\rangle_0^{\frac{1}{r},o} + \\ + \sum_{\substack{I \amalg J = [n] \setminus \{i\} \\ m_1 + m_2 = m}} \frac{m!}{m_1! m_2!} \left\langle \tau_{d_{i-1}}^{\alpha_i} \prod_{l \in I} \tau_{d_l}^{\alpha_l} \sigma^{m_1} \right\rangle_0^{\frac{1}{r},o} \left\langle \sigma \prod_{l \in J} \tau_{d_l}^{\alpha_l} \sigma^{m_2} \right\rangle_0^{\frac{1}{r},o}.$$

Second, if $m \geq 1$, then for any $i \in [n]$ with $d_i > 0$, we have

$$(5.5) \quad \left\langle \prod_{l \in [n]} \tau_{d_l}^{\alpha_l} \sigma^m \right\rangle_0^{\frac{1}{r},o} = \sum_{I \amalg J = [n] \setminus \{i\}} \sum_{\alpha=-1}^{r-2} \left\langle \tau_0^\alpha \tau_{d_{i-1}}^{\alpha_i} \prod_{l \in I} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r},\text{ext}} \left\langle \tau_0^{r-2-\alpha} \prod_{l \in J} \tau_{d_l}^{\alpha_l} \sigma^m \right\rangle_0^{\frac{1}{r},o} + \\ + \sum_{\substack{I \amalg J = [n] \setminus \{i\} \\ m_1 + m_2 = m-1}} \frac{(m-1)!}{m_1! m_2!} \left\langle \tau_{d_{i-1}}^{\alpha_i} \prod_{l \in I} \tau_{d_l}^{\alpha_l} \sigma^{m_1} \right\rangle_0^{\frac{1}{r},o} \left\langle \sigma \prod_{l \in J} \tau_{d_l}^{\alpha_l} \sigma^{m_2+1} \right\rangle_0^{\frac{1}{r},o}.$$

Under the above dictionary, every term on the right-hand side of (5.4) or (5.5) corresponds to a single term on the right-hand side of (3.9) or (3.10) with m Ramond marked points.

Since marked points of twist $r-1$ in closed extended theory may have descendants, one would expect the open-closed correspondence to generalize to that setting. In [3], the first author conjectured the precise equations that the open theory for $r=2$ should satisfy if it incorporates boundary descendants. The construction of boundary descendants when $r=2$ and all insertions are Neveu–Schwarz, was carried by Solomon and the third author, and will appear in the near future. It is also known how to construct these descendants for any r in

¹These observations are mostly numerical at this point, since the open theory is currently only defined for disks and the closed extended theory in genus zero with a single -1 twist.

genus zero, and that will also appear in a forthcoming work. The resulting topological recursion relations in genus zero are as follows. First, if $i \in [m]$, $b_i > 0$, and $j \in [n]$, then

$$(5.6) \quad \left\langle \prod_{h \in [m]} \sigma_{b_h} \prod_{l \in [n]} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r}, o} = \sum_{\substack{K_I \sqcup K_J = [m] \setminus \{i\} \\ I \sqcup J = [n] \\ j \in J}} \left\langle \sigma_{b_{i-1}} \prod_{h \in K_I} \sigma_{b_h} \prod_{l \in I} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r}, o} \left\langle \sigma \prod_{h \in K_J} \sigma_{b_h} \prod_{l \in J} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r}, o},$$

where σ_b corresponds to b descendants at a boundary marked point; and second, if $i \in [m]$, $b_i > 0$, and $j \in [m] \setminus \{i\}$, then

$$(5.7) \quad \left\langle \prod_{h \in [m]} \sigma_{b_h} \prod_{l \in [n]} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r}, o} = \sum_{\substack{K_I \sqcup K_J = [m] \setminus \{i\} \\ I \sqcup J = [n] \\ j \in K_J}} \left\langle \sigma_{b_{i-1}} \prod_{h \in K_I} \sigma_{b_h} \prod_{l \in I} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r}, o} \left\langle \sigma \prod_{h \in K_J} \sigma_{b_h} \prod_{l \in J} \tau_{d_l}^{\alpha_l} \right\rangle_0^{\frac{1}{r}, o}.$$

These two equations indeed transform to (3.10) under the open-closed dictionary.

Perhaps the most surprising effect of the open-closed correspondence is the -1 TRR (equation (3.11)). If we believe the dictionary, then this equation suggests, on the open side, the existence of “cotangent line classes” corresponding to a bundle \mathbb{L}_{boun} associated to a boundary component. These classes should satisfy the following equation for $h > 0$ and $i, j \in [n]$:

$$(5.8) \quad \left\langle \sigma_h^{\text{boun}} \prod_{p \in [n]} \tau_{d_p}^{\alpha_p} \prod_{q \in [m]} \sigma_{b_q} \right\rangle_0^{\frac{1}{r}, o} = \sum_{\substack{I \sqcup J = [n] \\ i, j \in J}} \sum_{\alpha=0}^{r-1} \left\langle \sigma_{h-1}^{\text{boun}} \tau_0^\alpha \prod_{p \in I} \tau_{d_p}^{\alpha_p} \prod_{q \in [m]} \sigma_{b_q} \right\rangle_0^{\frac{1}{r}, o} \left\langle \tau_0^{r-2-\alpha} \prod_{p \in J} \tau_{d_p}^{\alpha_p} \right\rangle_0^{\frac{1}{r}, \text{ext}} \\ + \sum_{\substack{I \sqcup J = [n] \\ K_I \sqcup K_J = [m] \\ i, j \in J}} \left\langle \sigma_{h-1}^{\text{boun}} \sigma \prod_{p \in I} \tau_{d_p}^{\alpha_p} \prod_{q \in K_I} \sigma_{b_q} \right\rangle_0^{\frac{1}{r}, o} \left\langle \prod_{p \in J} \tau_{d_p}^{\alpha_p} \prod_{q \in K_J} \sigma_{b_q} \right\rangle_0^{\frac{1}{r}, o},$$

where σ_h^{boun} corresponds to h copies of \mathbb{L}_{boun} . (There are analogous equations, also, if one or both of i, j lies in $[m]$.) Based on this hint, the first and the third authors have constructed a “class” that satisfies equation (5.8)—or, more precisely, a line bundle \mathbb{L}_{boun} and boundary conditions for which generic extensions give rise to (5.8). We leave the details of the construction, however, to future work.

REFERENCES

- [1] D. Abramovich and A. Vistoli. Compactifying the space of stable maps. *Journal of the American Mathematical Society*, 15(1):27–75, 2002.
- [2] M. Bertola and D. Yang. The partition function of the extended r -reduced Kadomtsev-Petviashvili hierarchy. *Journal of Physics. A. Mathematical and Theoretical*, 48(19), 2015.
- [3] A. Buryak. Equivalence of the open KdV and the open Virasoro equations for the moduli space of riemann surfaces with boundary. *Letters in Mathematical Physics*, 105(10):1427–1448, 2015.
- [4] A. Buryak. Open intersection numbers and the wave function of the KdV hierarchy. *Moscow Mathematical Journal*, 16(1):27–44, 2016.
- [5] A. Buryak, E. Clader, and R. J. Tessler. Open r -spin theory and the Gelfand–Dickey wave function. arXiv:1809.02536v1, 2018.
- [6] A. Buryak and R. J. Tessler. Matrix models and a proof of the open analog of Witten’s conjecture. *Communications in Mathematical Physics*, 353(3):1299–1328, 2017.
- [7] H.-L. Chang, J. Li, and W.-P. Li. Witten’s top Chern class via cosection localization. *Inventiones Mathematicae*, 200(3):1015–1063, 2015.
- [8] A. Chiodo. The Witten top Chern class via K -theory. *Journal of Algebraic Geometry*, 15(4):681–707, 2006.

- [9] A. Chiodo. Stable twisted curves and their r -spin structures. *Université de Grenoble. Annales de l'Institut Fourier*, 58(5):1635–1689, 2008.
- [10] A. Chiodo. Towards an enumerative geometry of the moduli space of twisted curves and r th roots. *Compositio Mathematica*, 144(6):1461–1496, 2008.
- [11] A. Chiodo and Y. Ruan. Landau–Ginzburg/Calabi–Yau correspondence for quintic three-folds via symplectic transformations. *Inventiones Mathematicae*, 182(1):117–165, 2010.
- [12] A. Chiodo and D. Zvonkine. Twisted r -spin potential and Givental's quantization. *Advances in Theoretical and Mathematical Physics*, 13(5):1335–1369, 2009.
- [13] L. A. Dickey. *Soliton equations and Hamiltonian systems. Second edition*, volume 26 of *Advanced Series in Mathematical Physics*. World Scientific Publishing Co., Inc., River Edge, NJ, 2003.
- [14] C. Faber, S. Shadrin, and D. Zvonkine. Tautological relations and the r -spin Witten conjecture. *Annales Scientifiques de l'École Normale Supérieure. Quatrième Série*, 43(4):621–658, 2010.
- [15] H. Fan, T. Jarvis, and Y. Ruan. The Witten equation, mirror symmetry and quantum singularity theory. *Annals of Mathematics*, 178(1):1–106, 2013.
- [16] T. J. Jarvis, T. Kimura, and A. Vaintrob. Gravitational descendants and the moduli space of higher spin curves. In *Advances in algebraic geometry motivated by physics (Lowell, MA, 2000)*, volume 276 of *Contemporary Mathematics*, pages 167–177. Amer. Math. Soc., Providence, RI, 2001.
- [17] T. J. Jarvis, T. Kimura, and A. Vaintrob. Moduli spaces of higher spin curves and integrable hierarchies. *Compositio Mathematica*, 126(2):157–212, 2001.
- [18] M. Kontsevich. Intersection theory on the moduli space of curves and the matrix Airy function. *Communications in Mathematical Physics*, 147(1):1–23, 1992.
- [19] T. Mochizuki. The virtual class of the moduli stack of stable r -spin curves. *Communications in Mathematical Physics*, 264(1):1–40, 2006.
- [20] R. Pandharipande, A. Pixton, and D. Zvonkine. Relations on $\overline{\mathcal{M}}_{g,n}$ via 3-spin structures. *Journal of the American Mathematical Society*, 28(1):279–309, 2015.
- [21] R. Pandharipande, J. P. Solomon, and R. J. Tessler. Intersection theory on moduli of disks, open KdV and Virasoro. arXiv:1409.2191, 2014.
- [22] A. Polishchuk and A. Vaintrob. Algebraic construction of Witten's top Chern class. In *Advances in algebraic geometry motivated by physics (Lowell, MA, 2000)*, volume 276 of *Contemporary Mathematics*, pages 229–249. Amer. Math. Soc., Providence, RI, 2001.
- [23] J. P. Solomon and R. J. Tessler. Graded spin surfaces and the open Arf invariant. In preparation.
- [24] J. P. Solomon and R. J. Tessler. Intersection Theory on the Moduli Space of Graded Riemann Surfaces with Boundary. In preparation.
- [25] R. J. Tessler. The combinatorial formula for open gravitational descendants. arXiv:1507.04951, 2015.
- [26] E. Witten. Two-dimensional gravity and intersection theory on moduli space. In *Surveys in differential geometry (Cambridge, MA, 1990)*, pages 243–310. Lehigh Univ., Bethlehem, PA, 1991.
- [27] E. Witten. Algebraic geometry associated with matrix models of two-dimensional gravity. In *Topological methods in modern mathematics (Stony Brook, NY, 1991)*, pages 235–269. Publish or Perish, Houston, TX, 1993.
- [28] D. Zvonkine. An introduction to moduli spaces of curves and their intersection theory. In *Handbook of Teichmüller theory. Volume III*, volume 17 of *IRMA Lectures in Mathematics and Theoretical Physics*, pages 667–716. Eur. Math. Soc., Zürich, 2012.

A. BURYAK:

SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LEEDS, LS2 9JT, UNITED KINGDOM, AND
 FACULTY OF MECHANICS AND MATHEMATICS, LOMONOSOV MOSCOW STATE UNIVERSITY, MOSCOW, GSP-1,
 119991, RUSSIAN FEDERATION

Email address: buryaksh@gmail.com

E. CLADER:

SAN FRANCISCO STATE UNIVERSITY, SAN FRANCISCO, CA 94132-1722, USA

Email address: eclader@sfsu.edu

R. J. TESSLER, INCUMBENT OF THE LILIAN AND GEORGE LYTTLE CAREER DEVELOPMENT CHAIR:

DEPARTMENT OF MATHEMATICS, WEIZMANN INSTITUTE OF SCIENCE, POB 26, REHOVOT 7610001, ISRAEL

Email address: ran.tessler@weizmann.ac.il