

This is a repository copy of *First degree cohomology of Specht modules and extensions of symmetric powers*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/141247/>

Version: Accepted Version

Article:

Donkin, Stephen and Geranios, Haralampos (2019) First degree cohomology of Specht modules and extensions of symmetric powers. *Advances in Mathematics*. pp. 618-701. ISSN 0001-8708

<https://doi.org/10.1016/j.aim.2019.01.022>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

This is a repository copy of *First degree cohomology of Specht modules and extensions of symmetric powers*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/113245/>

Monograph:

Donkin, Stephen and Geranios, Haralampos (Submitted: 2017) First degree cohomology of Specht modules and extensions of symmetric powers. Working Paper. (Submitted)

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

First degree cohomology of Specht modules and extensions of symmetric powers.

Stephen Donkin and Haralampos Geranios

Department of Mathematics, University of York, York YO10 5DD

`stephen.donkin@york.ac.uk, haralampos.geranios@york.ac.uk`

28 February 2017

Abstract

Let Σ_d denote the symmetric group of degree d and let K be a field of positive characteristic p . For $p > 2$ we give an explicit description of the first cohomology group $H^1(\Sigma_d, \mathrm{Sp}(\lambda))$, of the Specht module $\mathrm{Sp}(\lambda)$ over K , labelled by a partition λ of d .

We also give a sufficient condition for the cohomology to be non-zero for $p = 2$ and we find a lower bound for the dimension.

The cohomology of Specht modules has been considered in many papers including [12], [14], [17] and [23].

Our method is to proceed by comparison with the cohomology for the general linear group $G(n)$ over K and then to reduce to the calculation of $\mathrm{Ext}_{B(n)}^1(S^d E, K_\lambda)$, where $B(n)$ is a Borel subgroup of $G(n)$, where $S^d E$ denotes the d th symmetric power of the natural module E for $G(n)$ and K_λ denotes the one dimensional $B(n)$ -module with weight λ .

The main new input is the description of module extensions by: extension sequences, coherent triples of extension sequences and coherent multi-sequences of extension sequences, and the detailed calculation of the possibilities for such sequences. These sequences arise from the action of divided powers elements in the negative part of the hyperalgebra of $G(n)$.

Our methods are valid also in the quantised context and we aim to treat this in a separate paper.

Introduction

Let K be an algebraically closed field and let p be the characteristic of K . Let n be a positive integer. We write $G(n)$ for the general linear group over K of degree n , write $B(n)$ for the Borel subgroup consisting of lower triangular matrices and $T(n)$ for the maximal torus consisting of diagonal

matrices. The weights of a $G(n)$ -module or a $B(n)$ -module (or $T(n)$ -module) will be computed with respect to $T(n)$.

We set $X(n) = \mathbb{Z}^n$ and let $X^+(n)$ be the set of dominant weights, i.e., the set of $\lambda = (\lambda_1, \dots, \lambda_n) \in X(n)$ such that $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$. We write $\Lambda(n)$ for the set of polynomial weights, i.e., the set of $\lambda = (\lambda_1, \dots, \lambda_n) \in X(n)$ such that $\lambda_1, \dots, \lambda_n \geq 0$ and we write $\Lambda^+(n)$ for the set of dominant polynomial weights, i.e., $X^+(n) \cap \Lambda(n)$. The degree of a polynomial weight $\lambda = (\lambda_1, \dots, \lambda_n)$ is $\lambda_1 + \dots + \lambda_n$. For $d \geq 0$ we write $\Lambda(n, d)$ for the set of polynomial weights of degree d and $\Lambda^+(n, d)$ for $\Lambda(n, d) \cap X^+(n)$, the set of polynomial dominant weights of degree d . For $1 \leq r \leq n$ we define $\epsilon_r = (0, \dots, 0, 1, 0, \dots, 0) \in X(n)$ (with 1 appearing in the r th position).

We write E for the natural module and $S^d E$ for its d th symmetric power, for $d \geq 0$. For $\lambda \in X(n)$ we write K_λ for the one dimensional $B(n)$ -module with weight λ . For each $\lambda \in \Lambda^+(n)$ we have the induced module $G(n)$ -module $\nabla(\lambda) = \text{ind}_{B(n)}^{G(n)} K_\lambda$. In particular we have $\nabla(d, 0, \dots, 0) = S^d E$. Moreover, by a result of Cline, Parshall, Scott, van der Kallen, see e.g., [16] II, 4.7 Corollary a) we have the following.

Theorem 0.1. *For $i \geq 0$ and $\lambda, \mu \in \Lambda^+(n)$ we have*

$$\text{Ext}_{G(n)}^i(\nabla(\mu), \nabla(\lambda)) = \text{Ext}_{B(n)}^i(\nabla(\mu), K_\lambda)$$

and in particular

$$\text{Ext}_{G(n)}^i(S^d E, \nabla(\lambda)) = \text{Ext}_{B(n)}^i(S^d E, K_\lambda)$$

if $\lambda \in \Lambda^+(n, d)$.

We denote by Σ_d the symmetric group of degree d . For each partition λ of d we have the Specht module $\text{Sp}(\lambda)$ over K , for Σ_d .

Recall that, for $n \geq d$ and $\lambda, \mu \in \Lambda^+(n, d)$ we have $\text{Ext}_{G(n)}^i(\nabla(\mu), \nabla(\lambda)) = \text{Ext}_{\Sigma_d}^i(\text{Sp}(\mu), \text{Sp}(\lambda))$, for $i = 0$ and $p \neq 2$ and for $i = 1$ and $p \neq 2, 3$, by [5], Proposition 10.5 (i) and Proposition 10.6 (i), or [13]. Our interest in this paper is in degrees 0 and 1 so the relevant result for us is the following.

Proposition 0.2. *For $\lambda \in \Lambda^+(n, d)$ we have:*

- (i) $\text{Hom}_{G(n)}(S^d E, \nabla(\lambda)) = H^0(\Sigma_d, \text{Sp}(\lambda))$; and
- (ii) $\dim H^1(\Sigma_d, \text{Sp}(\lambda)) \geq \dim \text{Ext}_{G(n)}^1(S^d E, \nabla(\lambda))$, with equality if $p \neq 2$.

This follows in most cases by the above results or the paper [17] of Kleshchev and Nakano. We also give a proof, via the arguments of [5], Section 10, in Appendix I.

The detailed description of $\text{Ext}_{B(n)}^1(S^d E, K_\lambda)$ is quite complicated in general. However there are a couple of striking consequences that we now give. Throughout the analysis the notions of what we call James pairs and James

partitions are of paramount importance. We suppose that K has prime characteristic p . Let (a, b) be a two part partition. Let $v = \text{val}_p(a + 1)$ be the p -adic valuation of $a + 1$. Let β denote the p -adic length of b , i.e., β is the non-negative integer such that $b = \sum_{i=0}^{\beta} b_i p^i$, with $0 \leq b_i < p$ and $b_{\beta} \neq 0$. We call (a, b) a James pair if $\beta < v$ (equivalently $b < p^v$, equivalently $\binom{a+i}{i} = 0$, modulo p , for $1 \leq i \leq b$). We call a partition $\lambda = (\lambda_1, \dots, \lambda_n)$ of length $n \geq 2$ a James partition if $(\lambda_r, \lambda_{r+1})$ is James for all $1 \leq r < n$.

Let $\lambda = (\lambda_1, \dots, \lambda_n)$ be a James partition and let $l_i = \text{len}_p(\lambda_i)$ and $v_i = \text{val}_p(\lambda_i + 1)$, for $1 \leq i \leq n$. We define the *segments* of λ to be the equivalence classes of $\{1, \dots, n\}$ for the relation $r = s$ if and only if $l_r = l_s$. For $1 \leq r, s \leq n$ we define r and s to be adjacent if they belong to the same segment or if $1 < r < n$, $s = r + 1$, and $r + 1$ is the only element in its segment and $\lambda_r = p^{v_{r-1}} - 1$. We define the p -segments of λ to be the equivalent classes of $\{1, \dots, n\}$ for the equivalence relation generated by adjacency.

The following, obtained in Section 7, is a complete description of the dimensions of the extension spaces for James partitions.

Theorem 0.3. *Let λ be a James partition or degree d of length $n \geq 2$ and let $N \geq n$.*

- (i) *If $l_1 = l_2$ then $\dim \text{Ext}_{B(N)}^1(S^d E, K_{\lambda})$ is the number of p -segments of $\{1, \dots, n\}$.*
- (ii) *If $l_1 > l_2$ then $\dim \text{Ext}_{B(N)}^1(S^d E, K_{\lambda})$ is one less than the number of p -segments of $\{1, \dots, n\}$.*

In the non-James case the results are not so easy to describe but the following general result, Proposition 9.4(i) and (ii), is significant.

Proposition 0.4. *Let $\lambda = (\lambda_1, \dots, \lambda_n)$ be a non-James partition of length n and let $N \geq n$. Let r be minimal such that $(\lambda_r, \lambda_{r+1})$ is not James.*

- (i) *If $n \geq r + 2$ then there is an injective linear map from $\text{Ext}_{B(N)}^1(S^d E, K_{\lambda})$ to $\text{Ext}_{B(N)}^1(S^{d_0} E, K_{\mu})$, where μ is the 3-part partition $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ and d_0 is the degree of μ ;*
- (ii) *if $r = n - 1$ then there is an injective linear map from $\text{Ext}_{B(N)}^1(S^d E, K_{\lambda})$ to $\text{Ext}_{B(N)}^1(S^{d_0} E, K_{\mu})$, where μ is the 2-part partition $(\lambda_{n-1}, \lambda_n)$ and d_0 is the degree of μ .*

This leads to the following, see Theorem 12.1.

Theorem 0.5. *Let λ be a non-James partition or degree d of length $n \geq 2$ and let $N \geq n$. Then we have*

$$\dim \text{Ext}_{B(N)}^1(S^d E, K_{\lambda}) \leq 1.$$

In Theorems 12.30 and 12.31 we describe all non-James, non-split partitions, i.e. partitions λ with $\dim \text{Ext}_{B(N)}^1(S^d E, K_{\lambda}) = 1$ (where d is the degree of λ and N is at least the number of parts of λ).

Via Proposition 0.2, these results also give the dimensions of the first cohomology spaces of Specht modules for $p \neq 2$.

We mention some of the other highlights of the paper.

It is important for us to establish a close relationship between modules for the hyperalgebra of $B(n)$ and rational $B(n)$ -modules. What we do is in fact to give some version of Verma's Conjecture for $B(n)$ -modules. Since the arguments also give a proof of Verma's Conjecture (for all semisimple, simply connected algebraic groups) we include, in Section 1, a new proof (for earlier proofs see the papers by Sullivan, [21] and Cline, Parshall and Scott, [2]).

We give, in Section 3, a simple proof of James's Theorem on the fixed points of Specht modules, [15], 24.4 Theorem, via our formalism in terms of divided powers operators.

Our investigation into the nature of coherent multi-sequence starts in Section 5, where we determine all extension sequences for two part partitions. As a corollary we recover the result of Erdmann, [9], (3.6) Theorem, describing the first extension groups between Weyl modules for $SL_2(K)$.

As a by-product of our approach to extension spaces via coherent multi-sequences, we give, in Section 9, a short proof of the main result of Weber, [23], Theorem 1.2, which says that $H^1(\Sigma_d, Sp(\lambda)) = 0$ if the partition λ contains two distant non-James pairs.

In Section 13, using our main result on extensions, we give a counterexample to a conjecture of Hemmer on the cohomology of Specht modules.

We now describe the layout of the paper. In Section 1 we describe the link between modules for the hyperalgebra and rational modules, which is the starting point of our investigation. In Section 2 this is taken further to describe relations between the divided powers of operators and to give a description of the hyperalgebra by generators and relations.

In Section 3 we give, as a warm-up exercise in the use of the divided powers operations, a proof of James's Theorem on invariants of Specht modules.

In Section 4 we define extension multi-sequences make explicit the link between certain extensions of $B(n)$ -modules and coherent multi-sequences.

In Section 5 we determine the basic extension sequences, i.e, those sequence arising via two part partitions. The result is that a two part partition falls into one of three types: it is either split, or James, or what we call pointed. This trichotomy is the basis of the analysis for 3-part partitions which we give in later sections.

In Section 6 we give the definition of a coherent triple in terms of certain relations between basic extension sequences. These relations are used many times in the sections which follow, particularly to determine extension groups for 3-part partitions.

In Section 7 we first determine all coherent triples for a James 3-part partition. We then use this to determine the dimension of $\text{Ext}_{B(n)}^1(S^d E, K_\lambda)$

(and hence $H^1(\Sigma_d, \text{Sp}(\lambda))$), for any James partition λ . We find in particular that if λ is a James partitions of length n then the extension space has dimension at most $n - 1$, and that this bound is achieved.

In Section 8 we calculate the extension sequence for 3 part partitions (a, b, c) in which (a, b) is James and (b, c) is not.

In Section 9, we use the results given so far to describe various general results including Proposition 0.4 above, that the cohomology group, for a non-James partition λ is embedded in the cohomology for a certain 3 part or 2 part partition made up of consecutive parts of λ .

In Section 10 we describe all coherent triples for 3 part partitions (a, b, c) in which (a, b) is split.

In Section 11 we describe all coherent triples for 3 part partitions (a, b, c) in which (a, b) is pointed.

In Section 12 we obtain our main result for extension groups. The final result turns out to be much simpler than one might have anticipated, in view of the complexity coming from the analysis of coherent triples.

In Section 13 we comment on some questions raised by Hemmer, [12].

In Appendix I we give a proof of Proposition 0.2, based on [5], Section 10.

Finally, in Appendix II, we list together, for the convenience of the reader, the complete set of relations for coherent multi-sequences that are used throughout.

1 Divided powers operators, hyperalgebras and Verma's Conjecture.

In calculating extensions of $B(n)$ -modules it is technically much simpler to work with the action of the hyperalgebra, more precisely with $(\text{hy}(U(n)), T(n))$ -modules, where $\text{hy}(U(n))$ is the hyperalgebra of the group $U(n)$ of lower unitriangular matrices in $G(n)$, than the group $B(n)$ itself. In this section we describe the hyperalgebra of $U(n)$. We shall also need a version of Verma's Conjecture relating rational modules and modules for the hyperalgebra of a semisimple group. In fact the approach we take here leads to a new proof, which we include here, of the usual version of Verma's Conjecture.

Let G be a linear algebraic group over K . We write $K[G]$ for the coordinate algebra of G . Then $K[G]$ is naturally a Hopf algebra, in particular a coalgebra, and the full linear dual $K[G]^*$ is thus naturally an associative K -algebra. If V is a rational G -module then V is naturally a $K[G]^*$ -module. The action is given as follows. If V has basis $\{v_i \mid i \in I\}$ and the elements

$f_{ij} \in K[G]$, are defined by the formulas $gv_i = \sum_{j \in I} f_{ji}(g)v_j$, for $g \in G$, $i, j \in I$, then, for $\gamma \in K[G]^*$, we have $\gamma(v_i) = \sum_{j \in J} \gamma(f_{ji})v_j$.

Let M be the augmentation ideal of $K[G]$, i.e., $M = \{f \in K[G] \mid f(1) = 0\}$. Then the hyperalgebra (or algebra of distributions) $\text{hy}(G)$ is defined by

$$\text{hy}(G) = \{\gamma \in K[G]^* \mid \gamma(M^r) = 0 \text{ for some } r \geq 0\}.$$

Then $\text{hy}(G)$ is a Hopf subalgebra of the Hopf dual of $K[G]$. For a left $\text{hy}(G)$ -module V we have the space of “fixed points”

$$H^0(\text{hy}(G), V) = \{v \in V \mid Xv = \epsilon(X)v \text{ for all } X \in \text{hy}(G)\}$$

where $\epsilon : \text{hy}(G) \rightarrow K$ is the augmentation map of the Hopf algebra $\text{hy}(G)$. i.e., the map given by $\epsilon(X) = X(1_{K[G]})$, for $X \in \text{hy}(G)$.

Let e_{ij} , $1 \leq i, j \leq n$, be the matrix units of the algebra $M(n)$ of $n \times n$ matrices with entries in K and let I_n denote the identity matrix. For $1 \leq r, s \leq n$ with $r \neq s$, we define $u(r, s)(\xi) = I_n + \xi e_{rs} \in G(n)$, for $\xi \in K$, and define $U(r, s)$ to be the one dimensional unipotent subgroup $\{u(r, s)(\xi) \mid \xi \in K\}$ of $G(n)$.

The coordinate algebra $K[U(r, s)]$ is the free polynomial algebra in one variable $x(r, s)$, where $x(r, s)(u(r, s)(\xi)) = \xi$, for $\xi \in K$. The hyperalgebra $\text{hy}(U(r, s))$ has K -basis $X(r, s)_i$, $i \geq 0$, where $X(r, s)_i(x(r, s)^j) = \delta_{ij}$ (the Kronecker delta), for $i, j \geq 0$. We shall call these basis elements the *divided powers operators*.

If V is a rational $U(r, s)$ -module and $v \in V$ and $v_0, v_1, \dots \in V$ are such that

$$u(r, s)(\xi)v = \sum_{i \geq 0} \xi^i v_i$$

for $\xi \in K$, then $X(r, s)_i v = v_i$. From the fact that $u(r, s)(\nu)u(r, s)(\xi) = u(r, s)(\nu + \xi)$, for $\nu, \xi \in K$, we deduce that

$$X(r, s)_i X(r, s)_j v = \binom{i+j}{i} X(r, s)_{i+j} v$$

for $v \in V$, $i, j \geq 0$. Taking $V = K[G(n)]$, the left regular module and evaluating at 1, we deduce that

$$X(r, s)_i X(r, s)_j = \binom{i+j}{i} X(r, s)_{i+j}$$

for all $i, j \geq 0$.

One easily deduces that a linear map between $U(r, s)$ -modules is a module homomorphism if and only if it commutes with the action of all $X(r, s)_i$.

We write $U(n)$ for the unipotent radical of $B(n)$, i.e., the group of all lower unitriangular matrices.

Suppose that $1 \leq s < r \leq n$. Restriction of functions $K[U(n)]$ to $K[U(r, s)]$ is a surjection of Hopf algebras which induces a monomorphism of Hopf algebras $\text{hy}(U(r, s)) \rightarrow \text{hy}(U(n))$, by which we identify $\text{hy}(U(r, s))$ with a Hopf subalgebra of $\text{hy}(U(n))$.

Furthermore multiplication

$$U(2, 1) \times U(3, 1) \times U(3, 2) \times \cdots \times U(n, n-1) \rightarrow U(n)$$

is an isomorphism of varieties and induces K -space isomorphism

$$\text{hy}(U(2, 1)) \otimes \text{hy}(U(3, 1)) \otimes \text{hy}(U(3, 2)) \otimes \cdots \otimes \text{hy}(U(n, n-1)) \rightarrow \text{hy}(U(n)).$$

Hence the elements

$$X(2, 1)_{i(2,1)} X(3, 1)_{i(3,1)} X(3, 2)_{i(3,2)} \cdots X(n, n-1)_{i(n,n-1)} \quad (*)$$

with $i(2, 1), i(3, 1), i(3, 2), \dots, i(n, n-1) \geq 0$, form a K -basis of $\text{hy}(U(n))$ (a version of the Poincaré-Birkhoff-Witt Theorem).

Lemma 1.1. (i) A linear map between $U(n)$ -modules is a $U(n)$ -module homomorphism if and only if it commutes with the action of all $X(r+1, r)_i$, with $1 \leq r < n$, $i > 0$.

(ii) A linear map between $B(n)$ -modules is a $B(n)$ -module homomorphism if and only if it is a $T(n)$ -module homomorphism and commutes with the action of all $X(r+1, r)_i$, with $1 \leq r < n$, $i > 0$.

Proof. Let V and W be $B(n)$ -modules and let $\phi : V \rightarrow W$ be a linear map. Certainly if ϕ is a $B(n)$ -module map then it is a $k[B(n)]^*$ -module map and hence commutes with the action of all $X(r+1, r)_i$, $1 \leq r < n$, $i > 0$. Suppose, on the other hand that ϕ commutes with the action of all $X(r+1, r)_i$. Then, for $v \in V$ and $\xi \in K$, we have

$$\begin{aligned} \phi(u(r+1, r)(\xi)v) &= \phi\left(\sum_{i \geq 0} \xi^i X(r+1, r)_i v\right) = \sum_{i \geq 0} \xi^i X(r+1, r)_i \phi(v) \\ &= u(r+1, r)(\xi)\phi(v). \end{aligned}$$

Hence ϕ is a module homomorphism for the groups $U(r+1, r)$, $1 \leq r < n$, and hence for the group they generate, namely $U(n)$. This proves (i) and now (ii) is clear. \square

By a similar, but simpler argument, one also has the following.

Lemma 1.2. (i) A subspace of a $U(n)$ -module is a $U(n)$ -submodule if and only if it is stabilised by all divided power operators $X(r+1, r)_i$, for $1 \leq r < n$, $i > 0$.

- (ii) A $U(n)$ -module is trivial if and only if all divided power operators, $X(r+1, r)_i$, for $1 \leq r < n$, $i > 0$, act as zero.
- (iii) A subspace of a $B(n)$ -module is a $B(n)$ -submodule if and only if it is a $T(n)$ -submodule and stabilised by all divided power operators $X(r+1, r)_i$, for $1 \leq r < n$, $i > 0$.

We shall consider the category of $(\text{hy}(U(n)), T(n))$ -modules. By a $(\text{hy}(U(n)), T(n))$ -module we mean a vector space V , say, which is a $\text{hy}(U(n))$ -module and $T(n)$ -module in such a way that $X(r, s)_i V^\alpha \subseteq V^{\alpha + i(\epsilon_r - \epsilon_s)}$, for all $\alpha \in X(n)$, $1 \leq r, s \leq n$ with $r > s$ and $i > 0$. If V and W are $(\text{hy}(U(n)), T(n))$ -modules then a map $\phi : V \rightarrow W$ is a morphism of $\text{hy}(U(n))$ -modules and $T(n)$ -modules. If V is a $B(n)$ -module then we write V^{hy} , for V regarded as a $T(n)$ -module and $\text{hy}(U(n))$ -module as above. Thus we have an exact functor $V \mapsto V^{\text{hy}}$, from finite dimensional $B(n)$ -modules to finite dimensional $(\text{hy}(U(n)), T(n))$ -modules. We shall need that this is an equivalence of categories.

In fact the argument we shall give is valid for an arbitrary reductive group over K and we give it in that context. So now let G be a reductive group over K . Let T be a maximal torus and let $W = N_G(T)/T$ be the Weyl group (where $N_G(T)$ is the normaliser of T in G). Let $X(T)$ be the character group of T . Then W acts naturally on $X(T)$. We pick a positive definite symmetric W -invariant bilinear form on $\mathbb{R} \otimes_{\mathbb{Z}} X(T)$. Now T acts naturally on the Lie algebra $\text{Lie}(G)$ of G and the set of Φ of non-zero weights is a root system in the \mathbb{R} -span, E say, of Φ . We choose a set of positive roots $\Phi^+ = \{\alpha_1, \dots, \alpha_N\}$ (where $N = |\Phi^+|$). We have the usual partial order on $X(T)$. Thus for $\lambda, \mu \in X(T)$ we write $\lambda \geq \mu$ if $\lambda - \mu$ is a sum of positive roots.

Remark 1.3. We shall use the following elementary fact. If μ is a weight of a B -module V then there exists a one dimensional submodule isomorphic to K_λ for some $\lambda \in X(T)$ with $\lambda \leq \mu$. (This follows for example from [4], (1.4.4), or [16], II, 4.9 Lemma.)

The height of an element $\mu \in X(T)$ with $\mu \geq 0$ is defined by the condition that μ is a sum of $\text{ht}(\mu)$ simple roots. For $\mu \geq 0$ we define $P(\mu)$ to be the number of ways of writing μ as a sum of positive roots, i.e., the number of tuples $(m_\alpha)_{\alpha \in \Phi^+}$ such that $\mu = \sum_{\alpha \in \Phi^+} m_\alpha \alpha$.

For $\alpha \in \Phi$ let U_α be the corresponding root subgroup. Let G_{add} be the additive group of K , regarded as an algebraic group, in the natural way. For $\alpha \in \Phi$, we choose an isomorphism of algebraic groups $\phi_\alpha : G_{\text{add}} \rightarrow U_\alpha$ such that $t\phi_\alpha(\xi)t^{-1} = \phi_\alpha(\alpha(t)\xi)$, for $\xi \in K$.

There are uniquely determined elements $X_{\alpha, i}$, $i \geq 0$, of the hyperalgebra of U_α such that if V is a rational U_α -module and $v \in V$ then $\phi_\alpha(\xi)v = \sum_{i \geq 0} \xi^i X_{\alpha, i} v$. We denote by B the negative Borel subgroup of G and denote by U be the unipotent radical of B . Identifying $\text{hy}(U_\alpha)$ with a subalgebra

of $\text{hy}(U)$, for $\alpha \in \Phi^-$, we have, by the argument of (*) above, that $\text{hy}(U)$ has a K -basis consisting of the elements

$$X_{-\alpha_1, i_1} X_{-\alpha_2, i_2} \cdots X_{-\alpha_N, i_N}$$

with i_1, \dots, i_N running over the non-negative integers.

For $\lambda \in X(T)$ we denote by K_λ the one dimensional B -module on which T acts with weight λ . We shall use the fact that if $\text{Ext}_B^1(K_\lambda, K_\mu) \neq 0$, for $\lambda, \mu \in X(T)$, then $\lambda > \mu$ (see e.g. [16]).

We shall need a version of Lemma 1.2 in the more general situation. We leave the details to the reader.

Lemma 1.4. (i) *A subspace of a U -module is a U -submodule if and only if it is stabilised by all divided power operators $X_{-\alpha, i}$, for $\alpha \in \Phi^+$, $i > 0$.*
(ii) *A U -module is trivial if and only if all divided power operators, $X_{-\alpha, i}$, for $\alpha \in \Phi^+$, $i > 0$, act as zero.*
(iii) *A subspace of a B -module is a B -submodule if and only if it is a T -submodule and stabilised by all divided power operators $X_{-\alpha, i}$, for $\alpha \in \Phi^+$, $i > 0$.*

By a $(\text{hy}(U), T)$ -module we mean a vector space V , say, which is a T -module and $\text{hy}(U)$ -module in such a way that $X_{-\alpha, i} V^\lambda \subseteq V^{\lambda - i\alpha}$, for all $\lambda \in X(T)$, $\alpha \in \Phi^+$, $i \geq 0$. If V is a B -module we write V^{hy} for the corresponding $(\text{hy}(U), T)$ -module. We write $\text{mod}(B)$ for the category of finite dimensional rational B -module and $\text{mod}(\text{hy}(U), T)$ for the category of finite dimensional $(\text{hy}(U), T)$ -modules. We have, as above, an exact functor $F_B : \text{mod}(B) \rightarrow \text{mod}(\text{hy}(U), T)$, sending $V \in \text{mod}(B)$ to $V^{\text{hy}} \in \text{mod}(\text{hy}(U), T)$.

Proposition 1.5. *The functor $F_B : \text{mod}(B) \rightarrow \text{mod}(\text{hy}(U), T)$ is an equivalence of categories.*

Proof. It is enough to prove that for any $V \in \text{mod}(\text{hy}(U), T)$ there exists $Z \in \text{mod}(B)$ such that $Z = V^{\text{hy}}$.

We first produce a B -module that will be of great use to us. Multiplication $T \times U \rightarrow B$ is an isomorphism of varieties. We put

$$A = \{f \in K[B] \mid f(tu) = f(u) \text{ for all } t \in T, u \in U\}.$$

Then A is a left B -module summand of $K[B]$ and has trivial socle K . Thus A is the injective hull of K , as a B -module. The map $\pi : A \rightarrow K[U]$, given by restriction of functions is an isomorphism of K -algebra. The group T acts on U by conjugation and induces on $K[U]$ the structure of a left T -module. For $f \in A$, $t \in T$, $u \in U$, we have

$$\pi(tf)(u) = f(ut) = f(tt^{-1}ut) = (tf)(t^{-1}ut) = f(t^{-1}ut) = (t\pi(f))(u).$$

Hence π is an isomorphism of T -modules. Let $M = \{f \in K[U] \mid f(1) = 0\}$. For $r \geq 0$, the r th power M^r of the ideal M is a T -submodule of $K[U]$. By

complete reducibility we have that $K[U]$ and $\bigoplus_{r \geq 0} M^r/M^{r+1}$ are isomorphic as T -modules. Now $K[U]$ is a polynomial algebra so that the natural map from the r th symmetric power $S^r(M/M^2)$ to M^r/M^{r+1} is an isomorphism, for $r \geq 0$. Furthermore M/M^2 is T -module isomorphic to $\text{Lie}(U)^*$, the dual of the Lie algebra of U . The weights of $\text{Lie}(U)$ are $-\alpha_1, \dots, -\alpha_N$, and hence the weights of $\text{Lie}(U)^*$ are $\alpha_1, \dots, \alpha_N$, each occurring with multiplicity one. It follows that for $\mu \in X(T)$, we have $\dim K[U]^\mu$ is $P(\mu)$, if $\mu \geq 0$ and is 0 otherwise.

We regard $K[U]$ as a B -module via the isomorphism $\pi : A \rightarrow K[U]$. We fix $r \geq 0$. Let Q_r be the largest B -submodule of $K[U]$ with weights of height at most r . We claim that $\dim Q_r^\mu = P(\mu)$, for $\mu \in X(T)$ with $\text{ht}(\mu) \leq r$. Suppose that $\nu \in X(T)$ and K_ν appears in the socle of $K[U]/Q_r$. Then there is a submodule H of $K[U]$ containing Q_r such that H/Q_r is isomorphic to K_ν . Since H is not contained in Q_r we have $\text{ht}(\nu) > r$. Now if τ is any weight of $K[U]/Q_r$ then we must have $\tau \geq \nu$ for some $\nu \in X(T)$ such that K_ν appears in the socle of $K[U]/Q_r$, by Remark 1.3. Hence we have $\text{ht}(\tau) \geq \text{ht}(\nu) > r$. Hence we have $\dim Q_r^\tau = \dim K[U]^\tau = P(\tau)$, for all $\tau \geq 0$ with $\text{ht}(\tau) \leq r$.

We now show that any finite dimensional $(\text{hy}(B), T)$ -module V embeds in a module of the form Z^{hy} , for some B -module Z . Suppose V_1, V_2 are non-zero $(\text{hy}(B), T)$ -submodules such that $V_1 \cap V_2 = 0$. Thus V embeds in $V/V_1 \oplus V/V_2$ so that if V/V_1 embeds in Z_1^{hy} and V/V_2 embeds in Z_2^{hy} , for B -modules Z_1, Z_2 then V embeds in $Z_1^{\text{hy}} \oplus Z_2^{\text{hy}} = (Z_1 \oplus Z_2)^{\text{hy}}$. Hence we may suppose that V has simple socle L , say. Then L is isomorphic to K_λ^{hy} , for some $\lambda \in X(T)$. If $V \otimes L^*$ embeds in Z^{hy} , for some B -module Z then, since V is isomorphic to $(V \otimes L^*) \otimes L$, it embeds in $Z^{\text{hy}} \otimes K_\lambda^{\text{hy}} = (Z \otimes K_\lambda)^{\text{hy}}$. Hence, replacing V by $V \otimes L^*$, we may assume that socle of V is a one dimensional trivial module. Thus all weights of V are ≥ 0 and we choose r such that $\text{ht}(\mu) \leq r$, for every weight μ of V .

The dual module $H = V^*$ has simple head the one dimensional simple module. In particular H is generated by a weight vector h_0 , say, of weight 0. Hence H is spanned by the elements $X_{-\alpha_1, i_1} \dots X_{-\alpha_N, i_N} h_0$ such that $\text{ht}(i_1 \alpha_1 + \dots + i_N \alpha_N) \leq r$. We write I_r for the K -span of the elements $X_{-\alpha_1, i_1} \dots X_{-\alpha_N, i_N}$ such that $\text{ht}(i_1 \alpha_1 + \dots + i_N \alpha_N) > r$. We have a $\text{hy}(U)$ -module surjection $\phi : \text{hy}(U) \rightarrow H$, $\phi(X) = Xh_0$, for $X \in \text{hy}(U)$. Let J_H be the kernel of ϕ , i.e., the annihilator of h_0 . Then we have $I_r \subseteq J_H$ and $\text{hy}(U)/J_H \cong H$, in particular we have

$$\dim H \leq \dim \text{hy}(U)/I_r = \sum_{\mu} P(\mu)$$

where the sum is over all $\mu \in X(T)$ with $\mu \geq 0$ and $\text{ht}(\mu) \leq r$.

This applies in particular to Q_r^* and since $\dim Q_r^* = \dim \text{hy}(U)/I_r$ we have that I_r is the annihilator of an element of weight zero in Q_r^* , in particular

I_r is a left ideal and, regarding $\text{hy}(U)/I_r$ as a $(\text{hy}(U), T)$ -module, in such a way that $X_{-\alpha_1, i_1} \dots X_{-\alpha_N, i_N} + I_r$ has weight $-(i_1 \alpha_1 + \dots + i_N \alpha_N)$, we have $Q_r^* \cong \text{hy}(U)/I_r$. Hence we have that $H = \text{hy}(U)/J$ is an epimorphic image of Q_r^* and so V embeds in Q_r^{hy} .

Thus we have that a $(\text{hy}(U), T)$ -module V may be identified with a submodule of a module of the form Z^{hy} , for some B -module Z . But then, for $\alpha \in \Phi^+$, we have $\phi_{-\alpha}(\xi)v = \sum_{i \geq 0} \xi^i X_{-\alpha, i} v$, for $\xi \in K$, $v \in V$. Hence V is a B -submodule of Z and $V = V_0^{\text{hy}}$, where V_0 is the space V , regarded as a B -submodule of Z . □

Verma's Conjecture

We note that in fact the above Proposition gives a new proof of the so-called Verma's conjecture. For other proofs see [21] and [2]. We take G to be semisimple and simply connected. Then a G -module V is naturally a $\text{hy}(G)$ -module giving an exact functor $F_G : \text{mod}(G) \rightarrow \text{mod}(\text{hy}(G))$, where $\text{mod}(\text{hy}(G))$ denotes the category of finite dimensional $\text{hy}(G)$ -modules. Less formally, for a G -module V , we will say also that V is regarded as a $\text{hy}(G)$ -module. The action of $X_{\alpha, i}$, for $\alpha \in \Phi$, $i \geq 0$, on $v \in V$, is given by the formula $x_{\alpha}(\xi)v = \sum_{i \geq 0} \xi^i x_{\alpha, i} v$. Similarly a rational T -module may be regarded as a $\text{hy}(T)$ -module. The set $\text{Hom}_{K\text{-alg}}(\text{hy}(T), K)$, of K -algebra homomorphisms from $\text{hy}(T)$ to K , is naturally a group (since $\text{hy}(T)$ is a Hopf algebra). For $\lambda \in X(T)$, one has a k -algebra homomorphism $\lambda' : \text{hy}(T) \rightarrow K$, given by $h v = \lambda(h) v$, for $v \in T_{\lambda}$. Then map $X(T) \rightarrow \text{Hom}_{K\text{-alg}}(\text{hy}(T), K)$, sending λ to λ' is an injective homomorphism. We say that a K -algebra homomorphism from $\text{hy}(T)$ to K is integral if it has the form λ' for some $\lambda \in X(T)$. We identify $\lambda \in X(T)$ with $\lambda \in \text{Hom}_{K\text{-alg}}(\text{hy}(T), K)$ and in this way identify $X(T)$ with the group of integral K -algebra homomorphisms from $\text{hy}(T)$ to K . For a finite dimensional $\text{hy}(G)$ -module V we have $V = \bigoplus_{\lambda \in X(T)} V^{\lambda}$, where $V^{\lambda} = \{v \in V \mid h v = \lambda(h) v \text{ for all } h \in H\}$. Furthermore if V is simple then, it follows from the classification of finite dimensional simple modules, that V is isomorphic to $F_G(M)$, for some finite dimensional G -module M . For further details see e.g., [11].

We shall say that a finite dimensional $\text{hy}(G)$ -module V is rational if it is isomorphic to $F_G(M)$, for some rational G -module M . This is equivalent to the statement that the vector space V may be given the structure of a rational G -module in such a way that V^{λ} is the λ weight space for V , regarded as a T -module and $x_{\alpha}(\xi)v = \sum_{i \geq 0} \xi^i X_{\alpha, i} v$, for $\alpha \in \Phi$, $\xi \in K$, $v \in V$.

We shall need the following.

Lemma 1.6. (i) For a finite dimensional $\text{hy}(G)$ -module V we have $H^0(\text{hy}(G), V) = H^0(\text{hy}(U), V)^T$.

(ii) For finite dimensional $\text{hy}(G)$ -modules V_1, V_2 we have $\text{Hom}_{\text{hy}(G)}(V_1, V_2) = \text{Hom}_{\text{hy}(U)}(V_1, V_2)^T$.

Proof. (i) Certainly $H^0(\text{hy}(G), V) \leq H^0(\text{hy}(U), V)^T$. Let U^+ be the unipotent radical of the positive Borel subgroup G . Then we have that the multiplication map $\text{hy}(U^+) \otimes \text{hy}(T) \otimes \text{hy}(U) \rightarrow \text{hy}(G)$ is a vector space isomorphism. Let $v_0 \in H^0(\text{hy}(U), V)^T$. Then we have that $\text{hy}(G)v_0 = \text{hy}(U^+)\text{hy}(T)v_0$ and $\text{hy}(T)$ acts by scalar multiplication on Kv_0 so that $\text{hy}(G)v_0 = \text{hy}(U^+)v_0$. Moreover $\text{hy}(U^+)$ is spanned by elements of the form $X_{\alpha_1, i_1} \cdots X_{\alpha_N, i_N}$ so that the module $\text{hy}(G)v_0$ has only weights ≥ 0 . Hence the only composition factor of $\text{hy}(G)v_0$ is the trivial module and so $\text{hy}(G)v_0$ is trivial. Hence $v_0 \in H^0(\text{hy}(G), V)$ and $H^0(\text{hy}(G), V) = H^0(\text{hy}(U), V)^T$.

(ii) This follows from (i) by identifying $V_1^* \otimes V_2$ with $\text{Hom}_K(V_1, V_2)$ in the usual way. □

Theorem 1.7. (*Verma's Conjecture*) *The functor $F_G : \text{mod}(G) \rightarrow \text{mod}(\text{hy}(G))$ is an equivalence of categories.*

Proof. It is enough to prove that a finite dimensional $\text{hy}(G)$ -module is rational. Let V be a finite dimensional $\text{hy}(G)$ -module. We regard V as a $(\text{hy}(U), T)$ -module. By Proposition 1.5 the space V , regarded as a $(\text{hy}(U), T)$ -module has the structure of a B -module, i.e., there is a B -module action on V such that $x_\alpha(\xi)v = \sum_{i \geq 0} \xi^i X_{\alpha, i}v$, for $\alpha \in \Phi^+$, $\xi \in K$, $v \in V$. We write \bar{V} for the K -space V regarded as a B -module via this action.

The induction functor $\text{ind}_B^G : \text{mod}(B) \rightarrow \text{mod}(G)$ is left exact and we have the right derived functors $R^i \text{ind}_B^G : \text{mod}(B) \rightarrow \text{mod}(G)$, for $i \geq 0$. If M is a G -module then we have that $R^1 \text{ind}_B^G M = 0$ and, by [22], Exercise 2.1.4 for example, the evaluation map $\text{ev}_M : \text{ind}_B^G M \rightarrow M$ is a B -module isomorphism.

We claim that the evaluation map $\text{ev}_{\bar{V}} : \text{ind}_B^G \bar{V} \rightarrow \bar{V}$ is an isomorphism. If V is irreducible then it has the form $F_G(M)$ for some simple G -module M and then \bar{V} is isomorphic to the restriction of M to B , and hence $\text{ev}_{\bar{V}} : \text{ind}_B^G \bar{V} \rightarrow \bar{V}$ is an isomorphism. Now assume that V is not irreducible, let L be an irreducible $\text{hy}(G)$ -submodule and let $Q = V/L$. Then we have a commutative diagram

$$\begin{array}{ccccccc} 0 & \rightarrow & \text{ind}_B^G \bar{L} & \rightarrow & \text{ind}_B^G \bar{V} & \rightarrow & \text{ind}_B^G \bar{Q} \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \rightarrow & \bar{L} & \rightarrow & \bar{V} & \rightarrow & \bar{Q} \rightarrow 0 \end{array}$$

of B -modules, with rows exact (since $R^1 \text{ind}_B^G \bar{L} = 0$), where the vertical maps are the evaluation maps. We may assume, by induction on dimension, that the first and third vertical maps are isomorphisms and hence, so too is the middle map $\text{ev}_{\bar{V}} : \text{ind}_B^G \bar{V} \rightarrow \bar{V}$.

This proves the claim and therefore that the action of B on \overline{V} extends to a G -module action. Let Z be the vector space V regarded as a G -module via this action. Then we have $Z^\lambda = V^\lambda$, for $\lambda \in X(T)$, and $x_{-\alpha}(\xi)z = \sum_{i \geq 0} \xi^i X_{-\alpha, i} z$, for $\alpha \in \Phi^+$, $\xi \in K$, $z \in Z$. Hence, as a $(\text{hy}(U), T)$ -module, Z is precisely V . The identity map $Z \rightarrow V$ is an isomorphism of $(\text{hy}(U), T)$ -modules and hence, by Lemma 1.6(ii), an isomorphism of $\text{hy}(G)$ -modules. Thus we have $V = F_G(Z)$ and we are done. \square

2 Generators and Relations

To make further progress we give generators and relations for the hyperalgebra $\text{hy}(U(n))$ of $U(n)$. First fix $1 \leq r, s \leq n$, with $r \neq s$. Then, for $\eta, \zeta \in K$, we have $u(r, s)(\eta)u(r, s)(\zeta) = u(r, s)(\eta + \zeta)$. Hence, for a rational G -module V and $v \in V$, we have

$$u(r, s)(\eta)u(r, s)(\zeta)v = u(r, s)(\eta + \zeta)v$$

and hence

$$\sum_{i \geq 0, j \geq 0} \eta^i \zeta^j X(r, s)_i X(r, s)_j v = \sum_{k \geq 0} (\eta + \zeta)^k X(r, s)_k v.$$

Thus, for $i, j \geq 1$, we have

$$X(r, s)_i X(r, s)_j v = \binom{i+j}{i} X(r, s)_{i+j} v.$$

Applying this to the regular module $K[G]$ and evaluating at 1, we obtain

$$X(r, s)_i X(r, s)_j = \binom{i+j}{i} X(r, s)_{i+j}.$$

Since we are interested in the hyperalgebra of $U(n)$ we are interested in the operators $X(r, s)_i$ with $n \geq r > s \geq 1$, and $i > 0$. For ease of calculation we will find it useful to set $Y(r, s)_i = X(s, r)_i$, for $1 \leq r < s \leq n$, $i > 0$. We thus have

$$Y(r, s)_i Y(r, s)_j = \binom{i+j}{i} Y(r, s)_{i+j} \quad (1)$$

We now fix $1 \leq r < s < t \leq n$. Then it is easy to check that we have

$$u(t, s)(\eta)u(s, r)(\zeta) = u(s, r)(\zeta)u(t, r)(\eta\zeta)u(t, s)(\eta)$$

for $\eta, \zeta \in K$. Hence, for any G -module V and any $v \in V$ we have

$$u(t, s)(\eta)u(s, r)(\zeta)v = u(s, r)(\zeta)u(t, r)(\eta\zeta)u(t, s)(\eta)v$$

and therefore

$$\sum_{j, i \geq 0} \eta^j \zeta^i X(t, s)_j X(s, r)_i v = \sum_{h, i, j \geq 0} \zeta^{i+h} \eta^{j+h} X(s, r)_i X(t, r)_h X(t, s)_j v.$$

Hence for all $i, j \geq 0$ we have

$$X(t, s)_j X(s, r)_i v = \sum_{h=0}^{\min\{i, j\}} X(s, r)_{i-h} X(t, r)_h X(t, s)_{j-h} v$$

for all $v \in V$. Applying this to the regular module $K[G]$ and evaluating at the identity we have

$$X(t, s)_j X(s, r)_i = \sum_{h=0}^{\min\{i, j\}} X(s, r)_{i-h} X(t, r)_h X(t, s)_{j-h}.$$

Hence we have

$$Y(s, t)_j Y(r, s)_i = \sum_{h=0}^{\min\{i, j\}} Y(r, s)_{i-h} Y(r, t)_h Y(s, t)_{j-h} \quad (2)$$

for $1 \leq r < s < t \leq n$ and $i, j \geq 1$.

Moreover, we have $u(t, r)(\zeta)u(s, r)(\eta) = u(s, r)(\eta)u(t, r)(\zeta)$ and $u(t, r)(\zeta)u(t, s)(\eta) = u(t, s)(\eta)u(t, r)(\zeta)$ from which we deduce that

$$Y(r, s)_i Y(r, t)_j = Y(r, t)_j Y(r, s)_i \quad (3)$$

and

$$Y(s, t)_i Y(r, t)_j = Y(r, t)_j Y(s, t)_i \quad (4)$$

for all $1 \leq r < s < t \leq n$, $i, j > 0$.

Now suppose that $1 \leq q, r, s, t \leq n$ are distinct. Then we have $u(q, r)(\eta)u(s, t)(\zeta) = u(s, t)(\zeta)u(q, r)(\eta)$ and by the argument above we obtain

$$Y(q, r)_j Y(s, t)_i = Y(s, t)_i Y(q, r)_j \quad (5)$$

for all $1 \leq q, r, s, t \leq n$, with q, r, s, t distinct and $q < r$, $s < t$ and $i, j \geq 1$.

Now it is easy to see that relations (1)-(5) may be used to express any monomial in $Y(r, s)_i$ (with $1 \leq r, s \leq n$, $r \neq s$, $i \geq 1$) as a linear combination of monomials of the form

$$Y(1, 2)_{i(1,2)} Y(1, 3)_{i(1,3)} Y(2, 3)_{i(2,3)} \dots Y(n-1, n)_{i(n-1,n)}$$

with $i(1, 2), \dots, i(n-1, n)$ non-negative integers.

Hence we have:

Proposition 2.1. *For the generators $Y(r, s)_i$, $1 \leq r < s \leq n$, $i \geq 1$ the relations (1)-(5) above form a set of defining relations for $\text{hy}(U(n))$.*

3 James's Theorem on invariants of Specht modules.

We are interested in the degree 1 cohomology of Specht modules. However, we should mention that the degree 0 situation, namely the invariants of the Specht modules, was solved by James, see [15], 24.4 Theorem. As a warm-up exercise we deduce this from a result of $B(n)$ -module homomorphisms using the divided powers operators.

It is convenient to describe some notation and recall Lucas's formula at this point. In this section, for $a, b \geq 0$, the binomial coefficient $\binom{a}{b}$ will be understood to lie in the field K (of positive characteristic p), i.e., we write simply $\binom{a}{b}$ for $\binom{a}{b} 1_K$.

Notation: We write a non-negative integer a in base p as $a = \sum_{i \geq 0} p^i a_i$, with $0 \leq a_i < p$. If a is positive its p length $\text{len}_p(a)$ is the largest non-negative integer l such that $a_l \neq 0$ and its p -adic valuation $\text{val}_p(a)$ is the maximum positive integer v such that p^v divides a .

We recall that for a positive integers a and b written in base p as $a = \sum_{i \geq 0} a_i p^i$ and $b = \sum_{i \geq 0} b_i p^i$ we have

$$\binom{a}{b} = \binom{a_0}{b_0} \binom{a_1}{b_1} \dots \binom{a_m}{b_m} \quad (\text{Lucas's Formula})$$

where m is the maximum of $\text{len}_p(a)$ and $\text{len}_p(b)$.

For a non-negative integer a we define \bar{a} by the formula $a = a_0 + p\bar{a}$. By Lucas's formula, for non-negative integers $a = a_0 + p\bar{a}$, $b = b_0 + p\bar{b}$ we have $\binom{a}{b} = \binom{a_0}{b_0} \binom{\bar{a}}{\bar{b}}$.

Remark 3.1. For non-negative integers a, b , written in base p as above, we have $\binom{a+b}{b} \neq 0$ if and only if $a_i + b_i \leq p - 1$ for all $i \geq 0$. To see this we write $a = a_0 + p\bar{a}$, $b = b_0 + p\bar{b}$, as above. If $a_0 + b_0 \geq p$ then $(a+b)_0 < b_0$ and $\binom{a+b}{b} = 0$ by Lucas's Formula. If $a_0 + b_0 \leq p - 1$ then $\binom{(a+b)_0}{b_0} \neq 0$ and $\binom{a}{b} = \binom{(a+b)_0}{b_0} \binom{\bar{a} + \bar{b}}{\bar{b}}$ so the result follows by induction.

We first describe a condition that will be of crucial importance to us for the rest of the paper.

Definition 3.2. A two part partition (a, b) , with $b > 0$, will be called a James partition if $\binom{a+i}{i} = 0$ for all $1 \leq i \leq b$. A partition $\lambda = (\lambda_1, \dots, \lambda_n)$ of length n will be called a James partition if $(\lambda_r, \lambda_{r+1})$ is James, for all $1 \leq r < n$. We adopt the convention that a partition of length one is James.

By a two part partition we mean a partition with precisely two non-zero parts.

Remark 3.3. A two part partition (a, b) is a James partition if and only if $b < p^{\text{val}_p(a+1)}$. The James condition is that for all $1 \leq m \leq b$ we have $\binom{a+m}{m} = 0$, i.e., writing $m = \sum_{i \geq 0} m_i p^i$, we have $a_i + m_i > p - 1$ for some i . This can only happen if $a_0 = \dots = a_{\text{len}_p(b)} = p - 1$, i.e., $\text{val}_p(a+1) \geq \text{len}_p(b)$, i.e., $b < p^{\text{val}_p(a+1)}$.

We establish some notation involving the symmetric powers of the natural module that will be in force for rest of the paper. The module E has the basis e_1, \dots, e_n , where e_i is the column vector of length n with 1 in the i th position and zeroes elsewhere. For $\alpha = (\alpha_1, \dots, \alpha_n) \in \Lambda(n, d)$ we set $e^\alpha = e_1^{\alpha_1} e_2^{\alpha_2} \dots e_n^{\alpha_n} \in S^d E$.

Let $1 \leq s, t \leq n$ with $s \neq t$. We calculate the effect of $X(s, t)_i$ on $S^d E$. We have

$$u(s, t)(\xi)e_i = \begin{cases} e_i + \xi e_s, & \text{if } i = t; \\ e_i, & \text{if } i \neq t. \end{cases}$$

Since $U(n)$ acts on $S(E)$ by algebra automorphisms we have

$$u(s, t)(\xi)e^\alpha = e_1^{\alpha_1} \dots e_{t-1}^{\alpha_{t-1}} (e_t + \xi e_s)^{\alpha_t} e_{t+1}^{\alpha_{t+1}} \dots e_n^{\alpha_n}$$

which is

$$\sum_{i=0}^{\alpha_t} \binom{\alpha_t}{i} \xi^i e^{\alpha + i(\epsilon_s - \epsilon_t)}.$$

Hence we get

$$X(s, t)_i e^\alpha = \begin{cases} \binom{\alpha_t}{i} e^{\alpha + i(\epsilon_s - \epsilon_t)}, & \text{if } i \leq \alpha_t; \\ 0, & \text{otherwise.} \end{cases}$$

in particular, for $1 \leq r < n$, $i > 0$, we have

$$Y(r, r+1)_i e^\alpha = \begin{cases} \binom{\alpha_r}{i} e^{\alpha - i(\epsilon_r - \epsilon_{r+1})}, & \text{if } i \leq \alpha_r; \\ 0, & \text{otherwise.} \end{cases}$$

Let λ be a partition of degree d with at most n parts. We consider the space $\text{Hom}_{B(n)}(S^d E, K_\lambda)$. We have the largest trivial $U(n)$ -module quotient $H_0(U(n), S^d E)$. Thus $H_0(U(n), S^d E) = S^d E/N$, where N is the smallest $U(n)$ submodule of $S^d E$ with trivial quotient. Thus, by Lemma 1.2 (iii), we have

$$H_0(U(n), S^d E) = S^d E/N$$

where

$$N = \sum_{\substack{1 \leq r < n, i > 0, \\ \alpha \in \Lambda(n, d)}} KY(r, r+1)_i e^\alpha.$$

The weight spaces in $S^d E$, and hence in N and $S^d E/N$ are at most one dimensional. Thus N is the K -span of the e^β with $(\beta_1, \dots, \beta_n) \in \Lambda(n, d)$ such that e^β is a scalar multiple of $X(r, r+1)_i e^\alpha$, for some $\alpha = (\alpha_1, \dots, \alpha_n) \in \Lambda(n, d)$ and $1 \leq r < n$, $i > 0$. By considering weights we see that we must have $\alpha = \beta + i(\epsilon_r - \epsilon_{r+1})$ and $Y(r, r+1)_i e^\alpha = \binom{\beta_r + i}{i} e^\beta$. Hence $e^\beta \in N$ if and only if we have there exists some $1 \leq r < n$ and $0 < i \leq \beta_{r+1}$ such that $\binom{\beta_r + i}{i} \neq 0$,

Hence $H_0(U(n), S^d E)$ has K -basis $e^\mu + N$, where μ runs through the set of James partitions of n of degree r . Now we have $\text{Hom}_B(S^d E, K_\lambda) = \text{Hom}_{T(n)}(H_0(U(n), S^d E), K_\lambda)$, and so by Proposition 0.1 we obtain the following result.

Proposition 3.4. *For a partition λ of degree d we have*

$$\text{Hom}_B(S^d E, K_\lambda) = \begin{cases} K, & \text{if } \lambda \text{ is a James partition;} \\ 0, & \text{otherwise} \end{cases}$$

or

$$\text{Hom}_{G(n)}(S^d E, \nabla(\lambda)) = \begin{cases} K, & \text{if } \lambda \text{ is a James partition;} \\ 0, & \text{otherwise.} \end{cases}$$

By Proposition 0.2, we have James's result, [15], 24.4 Theorem:

Proposition 3.5. *For a partition λ of degree d we have*

$$H^0(\Sigma_d, \text{Sp}(\lambda)) = \begin{cases} K, & \text{if } \lambda \text{ is a James partition;} \\ 0, & \text{otherwise.} \end{cases}$$

4 Extension sequences and module extensions

We now wish to examine the possible $B(n)$ -module extensions of $S^d E$ by a one dimensional module.

We fix a partition λ with degree d and having at most n parts. Let V be an extension of $S^d E$ by K_λ . Thus V contains a one dimensional $B(n)$ -submodule V_∞ isomorphic to K_λ and V/V_∞ is isomorphic to $S^d E$. We fix $0 \neq v_\infty \in V_\infty$ and an isomorphism $\theta : V/V_\infty \rightarrow S^d E$. Note that θ is determined up to multiplication by a non-zero scalar since

$$\text{End}_{B(n)}(S^d E, S^d E) = \text{End}_{G(n)}(S^d E, S^d E) = K.$$

For $\alpha \in \Lambda(n, r)$, $\alpha \neq \lambda$ there is a unique element v_α of weight α such that $\theta(v_\alpha + V_\infty) = e^\alpha$. We also choose an element $v_\lambda \in V$ of weight λ such that $\theta(v_\lambda + V_\infty) = e^\lambda$.

Let $1 \leq r < s \leq n$. Suppose $\alpha \in \Lambda(n, r)$, $0 < i \leq \alpha_r$. Then we have

$$Y(r, s)_i e^\alpha = \binom{\alpha_r}{i} e^{\alpha - i(\epsilon_r - \epsilon_s)}$$

by Section 2 and hence, applying θ to $Y(r, s)_i v_\lambda + V_\infty$ we see that there exist scalars $y(r, s)_i \in K$ such that

$$Y(r, s)_i v_\alpha = \begin{cases} \binom{\alpha_r}{i} v_{\alpha - i(\epsilon_r - \epsilon_s)} + y(r, s)_i v_\infty, & \text{if } \alpha = \lambda + i(\epsilon_r - \epsilon_s); \\ \binom{\alpha_r}{i} v_{\alpha - i(\epsilon_r - \epsilon_s)}, & \text{if } \alpha \neq \lambda + i(\epsilon_r - \epsilon_s). \end{cases}$$

The scalars $y(r, s)_i$ are of paramount importance for our analysis of extensions of symmetric powers.

Definition 4.1. We call the sequence $(y(r, s)_i)_{1 \leq i \leq \lambda_s}$ the (r, s) -extension sequence determined by the V, v_∞ and θ . We call the collection of sequences $(y(r, s)_i)_{1 \leq r < s \leq n, 1 \leq i \leq \lambda_s}$, the extension multi-sequence determined by V, v_∞ and θ .

Of course we can take for V the split extension $V = S^d E \oplus K_\lambda$ and $V_\infty = K_\lambda = K v_\infty$. We take $\theta : V/V_\infty \rightarrow S^d E$ to be the natural map (induced by the projection from $V = S^d E \oplus V_\infty$ onto the first summand). We take $v_\lambda = e^\lambda - v_\infty$. Then, for $0 < i \leq \lambda_s$, we have

$$\begin{aligned} Y(r, s)_i v_{\lambda + i(\epsilon_r - \epsilon_s)} &= Y(r, s)_i e^{\lambda + i(\epsilon_r - \epsilon_s)} \\ &= \binom{\lambda_r + i}{i} e^\lambda = \binom{\lambda_r + i}{i} v_\lambda + \binom{\lambda_r + i}{i} v_\infty \end{aligned}$$

Definition 4.2. *The extension multi-sequence produced in this way will be called the standard multi-sequence for λ and denoted $(y(r, s)_i^{\text{st}})$. Specifically we have $y(r, s)_i = \binom{\lambda_r + i}{i}$, for $1 \leq r < s \leq n$ and $1 \leq i \leq \lambda_s$. A multiple of the standard multi-sequence will be called a standard multi-sequence.*

We continue with an analysis of extension multi-sequences arising from a general module extension V .

Suppose $i, j \geq 1$ and $i + j \leq \lambda_s$. Then by Section 2, (1), we have

$$Y(r, s)_i Y(r, s)_j v_{\lambda + (i+j)(\epsilon_r - \epsilon_s)} = \binom{i+j}{i} Y(r, s)_{i+j} v_{\lambda + (i+j)(\epsilon_r - \epsilon_s)}$$

i.e.,

$$\binom{\lambda_r + i + j}{j} Y(s, t)_i v_{\lambda + i(\epsilon_s - \epsilon_t)} = \binom{i+j}{i} Y(s, t)_{i+j} v_{\lambda + (i+j)(\epsilon_s - \epsilon_t)}$$

and taking the coefficients of v_∞ we get:

$$(E) \quad \binom{\lambda_r + i + j}{j} y(r, s)_j = \binom{i+j}{i} y(r, s)_{i+j}, \quad i, j \geq 1, i+j \leq \lambda_s.$$

It will be important to know all sequences satisfying this relation (E) and we classify them in the next section.

Now we consider a triple $1 \leq r < s < t \leq n$ and consider the relations that exist between the extension sequences $(y(r, s)_i)$, $(y(s, t)_j)$ and $(y(r, t)_k)$. We take $1 \leq i \leq \lambda_s$, $1 \leq k \leq \lambda_t$. By Section 2, (3), we have

$$Y(r, s)_i Y(r, t)_k v_{\lambda + (i+k)\epsilon_r - i\epsilon_s - k\epsilon_t} = Y(r, t)_k Y(r, s)_i v_{\lambda + (i+k)\epsilon_r - i\epsilon_s - k\epsilon_t}$$

i.e.,

$$\binom{\lambda_r + i + k}{k} Y(r, s)_i v_{\lambda + i(\epsilon_r - \epsilon_s)} = \binom{\lambda_r + i + k}{i} Y(r, t)_k v_{\lambda + k(\epsilon_r - \epsilon_t)}$$

and taking the coefficients of v_∞ we get:

$$(T1) \quad \binom{\lambda_r + i + k}{k} y(r, s)_i = \binom{\lambda_r + i + k}{i} y(r, t)_k, \quad 1 \leq i \leq \lambda_s, 1 \leq k \leq \lambda_t.$$

We take now $j, k \geq 1$ with $j + k \leq \lambda_t$. By Section 2, (4), we have

$$Y(s, t)_j Y(r, t)_k v_{\lambda + k\epsilon_r + j\epsilon_s - (j+k)\epsilon_t} = Y(r, t)_k Y(s, t)_j v_{\lambda + k\epsilon_r + j\epsilon_s - (j+k)\epsilon_t}$$

i.e.,

$$\binom{\lambda_r + k}{k} Y(s, t)_j v_{\lambda + j(\epsilon_s - \epsilon_t)} = \binom{\lambda_s + j}{j} Y(r, t)_k v_{\lambda + k(\epsilon_r - \epsilon_t)}$$

and taking the coefficients of v_∞ we get:

$$(T2) \quad \binom{\lambda_r + k}{k} y(s, t)_j = \binom{\lambda_s + j}{j} y(r, t)_k, \quad j, k \geq 1, j + k \leq \lambda_t.$$

We now take $1 \leq i \leq j \leq \lambda_t$ and use relation Section 2, (2). We have

$$\begin{aligned} & Y(s, t)_j Y(r, s)_i v_{\lambda+i(\epsilon_r-\epsilon_s)+j(\epsilon_s-\epsilon_t)} \\ &= \sum_{h=0}^i Y(r, s)_{i-h} Y(r, t)_h Y(s, t)_{j-h} v_{\lambda+i(\epsilon_r-\epsilon_s)+j(\epsilon_s-\epsilon_t)} \end{aligned}$$

i.e.,

$$\begin{aligned} & \binom{\lambda_r + i}{i} Y(s, t)_j v_{\lambda+j(\epsilon_s-\epsilon_t)} \\ &= \sum_{h=0}^i \binom{\lambda_s - i + j}{j - h} Y(s, t)_{i-h} Y(r, t)_h v_{\lambda+i(\epsilon_r-\epsilon_s)+h(\epsilon_s-\epsilon_t)} \\ &= \sum_{0 \leq h < i} \binom{\lambda_s - i + j}{j - h} \binom{\lambda_r + i}{h} Y(r, s)_{i-h} v_{\lambda+(i-h)(\epsilon_r-\epsilon_s)} \\ & \quad + \binom{\lambda_s - i + j}{j - i} Y(r, t)_i v_{\lambda+i(\epsilon_r-\epsilon_t)} \end{aligned}$$

and taking coefficients of v_∞ we get

$$\begin{aligned} (T3a) \quad & \binom{\lambda_r + i}{i} y(s, t)_j \\ &= \sum_{0 \leq h < i} \binom{\lambda_s - i + j}{j - h} \binom{\lambda_r + i}{h} y(r, s)_{i-h} + \binom{\lambda_s - i + j}{j - i} y(r, t)_i, \\ & \quad \text{for } 1 \leq i \leq j \leq \lambda_t. \end{aligned}$$

Now suppose that $1 \leq j \leq \lambda_t$, $j < i \leq \lambda_s + j$. Then we have $\lambda + (\epsilon_r - \epsilon_s) + j(\epsilon_s - \epsilon_t) \in \Lambda(n, r)$ and, using Section 2, (2) we have

$$\begin{aligned} & Y(s, t)_j Y(r, s)_i v_{\lambda+i(\epsilon_r-\epsilon_s)+j(\epsilon_s-\epsilon_t)} \\ &= \sum_{h=0}^j Y(r, s)_{i-h} Y(r, t)_h Y(s, t)_{j-h} v_{\lambda+i(\epsilon_r-\epsilon_s)+j(\epsilon_s-\epsilon_t)} \end{aligned}$$

i.e.,

$$\begin{aligned} & \binom{\lambda_r + i}{i} Y(s, t)_j v_{\lambda+j(\epsilon_s-\epsilon_t)} \\ &= \sum_{h=0}^i \binom{\lambda_s - i + j}{j - h} Y(s, t)_{i-h} Y(r, t)_h v_{\lambda+i(\epsilon_r-\epsilon_s)+h(\epsilon_s-\epsilon_t)} \\ &= \sum_{h=0}^i \binom{\lambda_s - i + j}{j - h} \binom{\lambda_r + i}{h} Y(r, s)_{i-h} v_{\lambda+(i-h)(\epsilon_r-\epsilon_s)} \end{aligned}$$

and taking coefficients of v_∞ we get

$$(T3b) \quad \binom{\lambda_r + i}{i} y(s, t)_j = \sum_{h=0}^j \binom{\lambda_s - i + j}{j - h} \binom{\lambda_r + i}{h} y(r, s)_{i-h},$$

for $1 \leq j \leq \lambda_t, j < i \leq \lambda_s + j$.

Solutions to the equations (T1)-(T3b) (i.e., what will be called coherent triples) will be analysed extensively in subsequence sections. There remains one further equation, coming from the commuting relation Section 2,(5). So suppose $1 \leq q, r, s, t \leq n$ and distinct and $q < r, s < t$. Then, for $1 \leq i \leq \lambda_t$, $1 \leq j \leq \lambda_r$, we obtain

$$Y(q, r)_j Y(s, t)_i v_{\lambda + i(\epsilon_s - \epsilon_t) + j(\epsilon_q - \epsilon_r)} = Y(s, t)_i Y(q, r)_j v_{\lambda + i(\epsilon_s - \epsilon_t) + j(\epsilon_q - \epsilon_r)}$$

i.e.,

$$\binom{\lambda_s + i}{i} Y(q, r)_j v_{\lambda + j(\epsilon_q - \epsilon_r)} = \binom{\lambda_q + j}{j} Y(s, t)_i v_{\lambda + i(\epsilon_s - \epsilon_t)}$$

and taking coefficients of v_∞ we obtain the “commuting condition”,

$$(C) \quad \binom{\lambda_s + i}{i} y(q, r)_j = \binom{\lambda_q + j}{j} y(s, t)_i,$$

for $1 \leq q, r, s, t \leq n$ distinct with $q < r, s < t$.

Proposition 4.3. *Let λ be a partition and let $(y(r, s)_i)$ be an extension multi-sequence for λ . Then the elements $y(r, s)_i \in K$ satisfy the equations (E),(T1)-(T3b) and (C) above. Conversely, given elements $y(r, s)_i \in K$, $1 \leq r < s \leq n$, $1 \leq i \leq \lambda_s$, satisfying (E),(T1)-(T3b) and (C) then $(y(r, s)_i)$ is an extension multi-sequence for some extension of $S^d E$ by K_λ .*

Proof. We have already proved the first statement, in deriving (E),(T1)-(T3b) and (C) above form an extension of $S^d E$ by K_λ .

Now suppose that we have elements $y(r, s)_i \in K$, $1 \leq r < s \leq n$, $1 \leq i \leq \lambda_s$, satisfying (E),(T1)-(T3b) and (C). We take a vector space $V(y)$ with basis elements $v_\alpha(y)$, $\alpha \in \Lambda(n, d)$, and $v_\infty(y)$. We make $V(y)$ into a T -module giving $v_\alpha(y)$ weight α , for $\alpha \in \Lambda(n, d)$, and giving $v_\infty(y)$ weight λ . We define K -linear endomorphisms $\hat{Y}(r, s)_i$, of $V(y)$, for $1 \leq r < s \leq n$, $1 \leq i \leq \lambda_s$, by

$$\hat{Y}(r, s)_i v_\alpha(y) = \begin{cases} \binom{\alpha_r}{i} v_{\alpha - (\epsilon_r - \epsilon_s)}(y) + y(r, s)_i v_\infty(y), & \text{if } i \leq \alpha_s, \alpha = \lambda + i(\epsilon_r - \epsilon_s); \\ \binom{\alpha_r}{i} v_{\alpha - (\epsilon_r - \epsilon_s)}, & \text{if } i \leq \alpha_s, \alpha \neq \lambda + i(\epsilon_r - \epsilon_s); \\ 0, & \text{otherwise} \end{cases}$$

and making $\hat{Y}(r, s)_i$ act as 0 on $v_\infty(y)$. Then the relations (E),(T1)-(T3b) and (C) guarantee that the endomorphisms $\hat{Y}(r, s)_i$ satisfy the relations (1)-(5) of Section 2. Hence by Proposition 2.1, there is a $hy(U)$ -module action on

$V(y)$ with $Y(r, s)_i$ acting via $\hat{Y}(r, s)_i$, $1 \leq r < s \leq n$, $1 \leq i \leq \lambda_s$. Moreover the T action makes $V(y)$ into a $(\text{hy}(U), T)$ -module and hence, by Proposition 1.5, a B -module. Clearly the extension multi-sequence associated to $V(y)$, with $v_\alpha = v_\alpha(y)$, $\alpha \in \Lambda(n, d)$ and $v_\infty = v_\infty(y)$ is $(y(r, s)_i)$. \square

Notation: It follows from Proposition 4.3 that the set of all extension multi-sequences for a partition λ forms a K -vector space, we denote this space by $E(\lambda)$.

We are now going to produce a K -space surjection $E(\lambda) \rightarrow \text{Ext}^1(S^d E, K_\lambda)$, whose kernel is the space of the standard coherent multi-sequence. For K_λ we take the field K with T acting with weight λ and $Y(r, s)_i$ acting as 0, for $1 \leq r < s \leq n$, $1 \leq i \leq \lambda_s$. For $y \in E(\lambda)$ let $V(y)$ be the B -module constructed via y , as above. We have a short exact sequence of B -modules

$$0 \rightarrow K_\lambda \xrightarrow{i_y} V(y) \xrightarrow{\pi_y} S^d E \rightarrow 0$$

where $i_y(1) = v_\infty(y)$ and $\pi_y(v_\alpha(y)) = e^\alpha$, for $\alpha \in \Lambda(n, d)$.

Starting with any B -module extension

$$0 \rightarrow K_\lambda \xrightarrow{i} V \xrightarrow{\pi} S^d E \rightarrow 0$$

we have $v_\infty = i(1)$ and $v_\alpha \in V$ such that $\pi(v_\alpha) = e^\alpha$ and we get an extension multi-sequence $y = (y(r, s)_i)$ given by

$$Y(r, s)_i v_{\lambda + i(\epsilon_r - \epsilon_s)} = \binom{\lambda_r + i}{i} v_\lambda + y(r, s)_i v_\infty$$

for $1 \leq i \leq \lambda_s$. Now it is easy to see that the B -module map $\phi : V \rightarrow V(y)$ such that $\phi(v_\alpha) = v_\alpha(y)$, $\alpha \in \Lambda(n, d)$, and $\phi(v_\infty) = v_\infty(y)$, induces an isomorphisms of extensions (in the sense of for example [1], Section I.5, or [20], Chapter 7) between the sequences $0 \rightarrow K_\lambda \xrightarrow{i} V \xrightarrow{\pi} S^d E \rightarrow 0$ and $0 \rightarrow K_\lambda \xrightarrow{i_y} V(y) \xrightarrow{\pi_y} S^d E \rightarrow 0$. In particular every extension of $S^d E$ by K_λ is equivalent to one arising via an element of $E(\lambda)$.

Now suppose that $y, y' \in E(\lambda)$ give short exact sequences $0 \rightarrow K_\lambda \xrightarrow{i_y} V(y) \xrightarrow{\pi_y} S^d E \rightarrow 0$ and $0 \rightarrow K_\lambda \xrightarrow{i_{y'}} V(y') \xrightarrow{\pi_{y'}} S^d E \rightarrow 0$. If these sequences are isomorphic then there is a B -module isomorphism $\phi : V(y) \rightarrow V(y')$ such that the diagram

$$\begin{array}{ccccccccc} 0 & \rightarrow & K_\lambda & \xrightarrow{i_y} & V(y) & \xrightarrow{\pi_y} & S^d E & \rightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \rightarrow & K_\lambda & \xrightarrow{i_{y'}} & V(y') & \xrightarrow{\pi_{y'}} & S^d E & \rightarrow & 0 \end{array}$$

commutes (where the first and third vertical maps are the identities and the middle vertical map is ϕ). Commutativity implies that $\phi(v_\infty(y)) = v_\infty(y')$ and also that $\phi(v_\alpha(y)) = v_\alpha(y')$, for $\lambda \neq \alpha \in \Lambda(n, d)$. Moreover, we must have $\phi(v_\lambda(y)) = v_\lambda(y') + cv_\infty(y')$, for some $c \in K$. The condition for ϕ to be a module homomorphism reduces to the condition

$$\phi(Y(r, s)_i v_{\lambda+i(\epsilon_r-\epsilon_s)}(y)) = Y(r, s)_i \phi(v_{\lambda+i(\epsilon_r-\epsilon_s)}(y))$$

i.e.,

$$\phi\left(\binom{\lambda_r+i}{i} v_\lambda(y) + y(r, s)_i v_\infty(y)\right) = Y(r, s)_i v_{\lambda+i(\epsilon_r-\epsilon_s)}(y')$$

i.e.,

$$\binom{\lambda_r+i}{i} (v_\lambda(y') + cv_\infty(y')) + y(r, s)_i v_\infty(y') = \binom{\lambda_r+i}{i} v_\lambda(y') + y'(r, s)_i v_\infty(y')$$

i.e.,

$$c \binom{\lambda_r+i}{i} + y(r, s)_i = y'(r, s)_i$$

for all $1 \leq r < s \leq n$, $1 \leq i \leq \lambda_s$. So we have shown the following result.

Lemma 4.4. *Every extension $0 \rightarrow K_\lambda \rightarrow V \rightarrow S^d E \rightarrow 0$ is equivalent to one of the form $0 \rightarrow K_\lambda \xrightarrow{i_y} V(y) \xrightarrow{\pi_y} S^d E \rightarrow 0$. Moreover, for $y, y' \in E(\lambda)$ the associated short exact sequences $0 \rightarrow K_\lambda \xrightarrow{i_y} V(y) \xrightarrow{\pi_y} S^d E \rightarrow 0$ and $0 \rightarrow K_\lambda \xrightarrow{i_{y'}} V(y') \xrightarrow{\pi_{y'}} S^d E \rightarrow 0$ are equivalent if and only if $y' - y$ is a multiple of the standard multi-sequence.*

We now follow a standard procedure (cf. [1], Section I.5, or [20], Chapter 7) to relate extensions of $S^d E$ by K_λ to elements of $\text{Ext}_B^1(S^d E, K_\lambda)$ and from this produce a linear epimorphism $E(\lambda)$ to $\text{Ext}_B^1(S^d E, K_\lambda)$ with kernel consisting of the standard coherent multi-sequences.

We fix an injective module I containing K_λ . Let $Q = I/K_\lambda$ and let $\nu : I \rightarrow Q$ be the natural map. Applying $\text{Hom}_B(S^d E, -)$ to the short exact sequence $0 \rightarrow K_\lambda \rightarrow I \rightarrow Q \rightarrow 0$ we obtain an exact sequence

$$\text{Hom}_B(S^d E, I) \rightarrow \text{Hom}_B(S^d E, Q) \rightarrow \text{Ext}_B^1(S^d E, K_\lambda)$$

so we may identify $\text{Ext}_B^1(S^d E, K_\lambda)$ with $\text{Hom}_B(S^d E, Q)/\text{Hom}_B(S^d E, Q)_0$, where

$$\text{Hom}_B(S^d E, Q)_0 = \{\nu \circ \theta \mid \theta \in \text{Hom}_B(S^d E, I)\}.$$

We write $E(S^d E, K_\lambda)$ for the set of equivalence classes of B -module extensions

$$0 \rightarrow K_\lambda \xrightarrow{i} V \rightarrow S^d E \xrightarrow{\pi} 0 \quad (\mathcal{E}).$$

For such an extension we have, by injectivity, a homomorphism $\theta : V \rightarrow I$ such that $\theta \circ i : K_\lambda \rightarrow I$ is inclusion. We obtain a homomorphism $\bar{\theta} : S^d E \rightarrow Q$, given by $\bar{\theta}(\pi(v)) = \theta(v) + K_\lambda$, $v \in V$, and hence an element $\bar{\theta} + \text{Hom}_B(S^d E, Q)_0$ of $\text{Hom}_B(S^d E, Q)/\text{Hom}_B(S^d E, Q)_0$. One checks that this element depends only on the equivalence class $[\mathcal{E}]$ of \mathcal{E} . Hence we have a map $\Phi : E(S^d E, K_\lambda) \rightarrow \text{Hom}_B(S^d E, Q)/\text{Hom}_B(S^d E, Q)_0$ and, for extensions \mathcal{E} and \mathcal{F} we have $\Phi([\mathcal{E}]) = \Phi([\mathcal{F}])$ if and only if \mathcal{E} and \mathcal{F} are equivalent.

Now for $y \in E(\lambda)$ we have the extension $0 \rightarrow K_\lambda \xrightarrow{i_y} V(y) \xrightarrow{\pi_y} S^d E \rightarrow 0$, which we now call $\mathcal{E}(y)$. Hence we get a map $\Psi : E(\lambda) \rightarrow \text{Hom}_B(S^d E, Q)/\text{Hom}_B(S^d E, Q)_0$, defined by $\Psi(y) = \Phi([\mathcal{E}(y)])$. It is easy to check that $\Phi([\mathcal{E}(y+y')]) = \Phi([\mathcal{E}(y)]) + \Phi([\mathcal{E}(y')])$, for $y, y' \in E(\lambda)$ and that $\Phi([\mathcal{E}(cy)]) = c\Phi([\mathcal{E}(y)])$, for $y \in E(\lambda)$, $y \in E(\lambda)$. Hence we have a K -linear map $\Psi : E(\lambda) \rightarrow \text{Hom}_B(S^d E, Q)/\text{Hom}_B(S^d E, Q)_0$. It is the composite of two surjections and hence surjective. Moreover, for $y \in E(\lambda)$ we have $\Psi(y) = 0$ if and only if $\Phi([\mathcal{E}(y)]) = 0$, i.e., if and only if $\mathcal{E}(y)$ is split, which by Lemma 1.3, is if and only if y is a standard coherent multi-sequence. Since $\text{Hom}_B(S^d E, Q)/\text{Hom}_B(S^d E, Q)_0$ is isomorphic to $\text{Ext}^1(S^d E, K_\lambda)$ we have shown the following.

Proposition 4.5. (i) *There is a surjective K -linear map from $E(\lambda)$ to $\text{Ext}^1(S^d E, K_\lambda)$ with kernel spanned by the standard multi-sequence.*
(ii) *We have*

$$\dim \text{Ext}_B^1(S^d E, K_\lambda) = \begin{cases} \dim E(\lambda), & \text{if } \lambda \text{ is James} \\ \dim E(\lambda) - 1, & \text{if } \lambda \text{ is not James.} \end{cases}$$

This prompts the following definition.

Definition 4.6. *We shall say that a partition λ is split if every coherent multi-sequence for λ is a multiple of the standard sequence.*

Thus a partition λ is split if and only if we have $\text{Ext}_{B(N)}^1(S^d E, K_\lambda) = 0$ (where d is the degree of λ , where N is at least the number of parts of λ and E is the natural $G(N)$ -module).

5 Extension sequences for two part partitions

Let $\lambda = (\lambda_1, \dots, \lambda_n)$ be a partition of length at least 2. We fix $1 \leq r < s \leq n$ and set $a = \lambda_s$, $b = \lambda_t$. According to Section 4, (E), we have

$$\binom{a+i+j}{j} y(r, s)_i = \binom{i+j}{j} y(r, s)_{i+j}$$

for all $i, j \geq 1$ with $i + j \leq b$.

Definition 5.1. Let (a, b) be a two part partition. By an extension sequence for (a, b) we mean a sequence $(x_i) = (x_1, \dots, x_b)$ of elements of K such that

$$\binom{a+i+j}{j} x_i = \binom{i+j}{j} x_{i+j}$$

for all $i, j \geq 1$ with $i + j \leq b$.

Since a general extension multi-sequence is a family of extension sequences it is important to thoroughly understand the possible extension sequences. We describe all such in this section. We then specialise to the case of $n = 2$ to recover the main result of Erdmann's paper, [9]

We now make some elementary observations on extension sequences.

Remark 5.2. Suppose (x_i) is an extension sequence for a two part partition (a, b) and $c + sp^k \leq b$, with $0 < c < p^k$. Then we have

$$\binom{a+c+sp^k}{c} x_{sp^k} = \binom{a+c+sp^k}{sp^k} x_c = \binom{c+sp^k}{sp^k} x_{c+sp^k}$$

and moreover, we have $\binom{c+sp^k}{sp^k} = 1$, by Lucas's formula. So we deduce

$$x_{c+sp^k} = \binom{a+c+sp^k}{c} x_{sp^k} = \binom{a+c+sp^k}{sp^k} x_c.$$

Lemma 5.3. Suppose that (a, b) is a two part partition. An extension sequence for (a, b) which is 0 at all powers of p is identically 0.

Proof. Suppose x_1, \dots, x_b is such a sequence. By induction it is enough to prove that $x_b = 0$ and we may assume that $b = sp^k$, for some $1 < s \leq p-1$. Taking $i = (s-1)p^k$, $j = p^k$ in the defining relation we get $\binom{sp^k}{p^k} x_{sp^k} = 0$, and $\binom{sp^k}{p^k} = s \neq 0$ so that $x_{sp^k} = 0$, as required. \square

Remark 5.4. Let (a, b) be a two part partition. We have the standard extension sequence (x_i) for (a, b) given by $x_i^{\text{st}} = \binom{a+i}{i}$, $1 \leq i \leq b$.

Remark 5.5. A two part partition (a, b) is James if and only if the standard extension sequence for (a, b) is the zero sequence.

Remark 5.6. There is a non-zero extension sequence for a James partition (a, b) . To see this consider the sequence of integers $N_i = \binom{a+i}{i}$, $1 \leq i \leq b$. Note that these integers satisfy the condition

$$\binom{a+i+j}{j} N_i = \binom{i+j}{i} N_{i+j}$$

for all $i, j \geq 1$ with $i + j \leq b$. Let h be the largest positive integer such that p^h divides all N_i and set $N'_i = N_i/p^h$, for $1 \leq i \leq b$. Then we also have

$$\binom{a+i+j}{j}_{N'_i} = \binom{i+j}{i}_{N'_{i+j}}$$

for all $i, j \geq 1$ with $i + j \leq b$ so that taking N'_1, \dots, N'_b modulo p we obtain a non-zero extension sequence. We call this the *canonical extension sequence*. We shall explore this in greater depth (for James partitions of any length) in Section 7. In particular we have $E(a, b) \neq 0$.

It is easy to describe the extension sequences for a James partition.

Lemma 5.7. *Let (a, b) be a two part James partition and let $\beta = \text{len}_p(b)$. A sequence x_1, \dots, x_b is an extension sequence if and only if*

$$x_{p^\beta} = 2x_{2p^\beta} = \dots = b_\beta x_{b_\beta p^\beta}$$

and $x_i = 0$ when i is not a multiple of p^β .

Proof. Let $k \leq \beta$ and suppose that $1 < s \leq b_k$. Then, taking $i = (s-1)p^k$ and $j = p^k$ in the defining condition for an extension sequence, we have

$$\binom{a+sp^k}{p^k} x_{(s-1)p^k} = \binom{sp^k}{(s-1)p^k} x_{sp^k}.$$

Now $k < \text{val}_p(a+1)$ so we have $a_k = p-1$ and therefore $(a+sp^k)_k = s-1$ and hence, by Lucas's formula, $(s-1)x_{(s-1)p^k} = sx_{sp^k}$ and so we get

$$x_{p^k} = 2x_{2p^k} = \dots = b_k x_{b_k p^k}. \quad (*)$$

We now apply the defining condition with $i = (p-1)p^{\beta-1}$ and $j = p^{\beta-1}$. We get

$$\binom{a+p^\beta}{p^{\beta-1}} x_{(p-1)p^{\beta-1}} = \binom{p^\beta}{(p-1)p^{\beta-1}} x_{p^\beta} = 0.$$

But $\binom{a+p^\beta}{p^{\beta-1}} = (a+p^\beta)_{\beta-1} = p-1 \neq 0$ so that $x_{(p-1)p^{\beta-1}} = 0$ and so by (*), $x_{sp^{\beta-1}} = 0$ for $1 \leq s \leq p-1$. But now $p^\beta - 1 < b$ and $(a, p^\beta - 1)$ is James so we may assume, by induction, that $x_i = 0$ for $1 \leq i < p^\beta$. Now Remark 5.2 gives that in fact $x_i = 0$ for all $i \leq b$, with i not a multiple of p^β . Furthermore the space of extension sequences for (a, b) is at most one dimensional, by (*). However, we know, by Remark 5.6, that there is a non-zero extension sequence, it follows that the space of such sequences is one dimensional and is as described in the statement of the Lemma. \square

Remark 5.8. *Lemma 5.7 implies that an extension sequence for a two part James partition is a multiple of the canonical sequence.*

Definition 5.9. A two part partition (a, b) will be called a *pointed partition* if b can be written in the form $b = \hat{b} + p^\beta$, with $\beta = \text{len}_p(b)$ with $\hat{b} < p^{\text{val}_p(a+1)}$ and $\beta > \text{val}_p(a+1)$. In this case we call the extension sequence (x_i^{pt}) , $1 \leq i \leq b$, defined by

$$x_i^{\text{pt}} = \begin{cases} 1, & \text{if } i = p^\beta; \\ 0, & \text{otherwise.} \end{cases}$$

the point sequence for (a, b) .

Remark 5.10. Note that the point sequence is non-standard since, with the above notation, the p^v th term in the standard sequence $\binom{a+p^v}{p^v} = a_v + 1$ is non-zero, where $v = \text{val}_p(a+1)$.

Lemma 5.11. Suppose that (a, b) is a two part partition which is not James. Let $v = \text{val}_p(a+1)$ and $\beta = \text{len}_p(b)$. Then (a, b) is split unless we may write $b = \hat{b} + p^\beta$, where $\hat{b} < p^v < p^\beta$ and in this case $E(a, b)$ is spanned by the standard sequence and the point sequence.

Proof. First assume that $b = \hat{b} + p^\beta$, with $\hat{b} < p^v < p^\beta$. We check that the point sequence x_1, \dots, x_b is indeed an extension sequence. If $\hat{b} = 0$ then we need only to check the defining relation for $i + j = p^\beta$ and in that case $\binom{i+j}{i} = 0$ and $x_i = 0$ (for $i < p^\beta$) so the relation holds.

So we suppose that $\hat{b} > 0$. We may assume inductively that x_1, \dots, x_{b-1} is an extension sequence and again we need only to check the defining relations with $i + j = b$. Moreover $x_b = 0$, as $b \neq p^\beta$, and so we must check that $\binom{a+b}{j} x_i$ is zero for $b = i + j$. Since $x_i = 0$ for $i \neq p^\beta$ we may assume that $i = p^\beta$, $j = \hat{b}$ and we must check that $\binom{a+b}{\hat{b}} = 0$. We choose k minimal such that $\hat{b}_k \neq 0$. Then $a_k = p - 1$ (since $\hat{b} < p^v$) and hence $(a+b)_k = b_k - 1$ so that $\binom{a+b}{\hat{b}_k} = 0$ and $\binom{a+b}{\hat{b}} = 0$, by Lucas's Formula.

We now prove that (a, b) is split unless it has the above form and in that case the space of extensions sequences is spanned by those described above. If $b = 1$ then (a, b) is not pointed and (since (a, b) is not James) we have $v = 0$ and $x_1^{\text{st}} \neq 0$ so that an extension sequence x_1 is a multiple of the standard sequence.

We now consider the case $b > 1$. Let x_1, \dots, x_b be an extension sequence for (a, b) .

Assume that $(a, b-1)$ is not James. If $(a, b-1)$ is split then x_1, \dots, x_{b-1} is a standard sequence and hence, subtracting a standard sequence for (a, b) we may assume that $x_i = 0$ for $i < b$. Hence, by Lemma 5.3 (x_i) is the zero sequence unless $b = p^\beta$. But we have $p^\beta - 1 \geq p^v$ so that $\beta > v$ and (a, b) is of the required form and the extension sequence is also of the required form. If $(a, b-1)$ is not split we may assume inductively that we can write $b-1 = c + p^t$ with $c < p^v < p^t$. Moreover, after subtracting a linear combination of the standard solution and the point solution for $(a, b-1)$,

we may assume that $x_i = 0$ for all $i < b$. If b is not a power of p this implies that x_b is also zero by Remark 5.2. Hence we can assume that $b = p^\beta$. Then we get $p^{\beta-1} - 1 < p^v < p^{\beta-1}$, and this is not possible.

It remains to consider the case in which $(a, b-1)$ is James. So we have $b-1 < p^v$ and, since (a, b) is not James, $b = p^v$. By Lemma 5.7 we have

$$x_{p^{v-1}} = 2x_{2p^{v-1}} = \cdots = (p-1)x_{(p-1)p^{v-1}}$$

and $x_i = 0$ for $i < b$ and i not divisible by p^{v-1} .

Taking $i = (p-1)p^{v-1}$, $j = p^{v-1}$ in the defining relation we get

$$\binom{a+p^v}{p^{v-1}} x_{(p-1)p^{v-1}} = 0.$$

Moreover $a_{v-1} = p-1$ so that $\binom{a+p^v}{p^{v-1}} = p-1$, by Lucas's Formula, and $x_{(p-1)p^{v-1}} = 0$. Hence we have $x_i = 0$ for $0 < i < b = p^v$. Hence the space of extensions sequences for (a, b) is at most one dimensional. Hence the space of extension sequences consists of standard solutions and (a, b) is split. \square

We now take $n = 2$ and consider the extensions for the general linear group $G(2)$, over K . Let $(r, s), (t, u) \in \Lambda^+(n, r)$ with $r+s = t+u$. Then we have $\text{Ext}_{G(2)}^1(\nabla(r, s), \nabla(t, u)) = 0$ unless $(r, s) > (t, u)$. Then have $s < u$. Moreover, we have $\nabla(r, s) = \nabla(r-s, 0) \otimes D^{\otimes s}$ and $\nabla(t, u) = \nabla(t-s, u-s) \otimes D^{\otimes s}$, where D is the one dimensional determinant module. Hence we have

$$\text{Ext}_{G(2)}^1(\nabla(r, s), \nabla(t, u)) = \text{Ext}_{G(2)}^1(\nabla(r-s, 0), \nabla(t-s, u-s))$$

i.e.,

$$\text{Ext}_{G(2)}^1(\nabla(r, s), \nabla(t, u)) = \text{Ext}_{G(2)}^1(S^d E, \nabla(t-s, u-s)).$$

where $d = r-s$. Hence, from Lemmas 5.7 and 5.11, we have the following.

Corollary 5.12. *Let $(r, s), (t, u)$ be two part partitions of the same degree. Then*

$$\text{Ext}_{G(2)}^1(\nabla(r, s), \nabla(t, u)) = \begin{cases} K, & \text{if } (t-s, u-s) \text{ is James or pointed;} \\ 0, & \text{otherwise} \end{cases}$$

i.e.,

$$\text{Ext}_{G(2)}^1(\nabla(r, s), \nabla(t, u)) = \begin{cases} K, & \text{if } u-s < p^v \\ & \text{or } u-s-p^l < p^v < p^l; \\ 0, & \text{otherwise} \end{cases}$$

where $v = \text{val}_p(t-s+1)$ and $l = \text{len}_p(u-s)$.

We now consider extensions for $\mathrm{SL}_2(K)$, as in [9]. For $r \geq 0$ we write simply $\nabla(r)$ for the r th symmetric power of the natural representation. We have $\mathrm{Ext}_{\mathrm{SL}_2(K)}^1(\nabla(r), \nabla(s)) = 0$ unless $r - s = 2m$ for some positive integer m . Moreover, in this case we have $\mathrm{Ext}_{\mathrm{SL}_2(K)}^1(\nabla(r), \nabla(s)) = \mathrm{Ext}_{G(2)}^1(\nabla(r, 0), \nabla(s + m, m))$. Hence we obtain the following result of Erdmann, [9], (3.6) Theorem.

Corollary 5.13. *For $r, s \geq 0$ we have*

$$\mathrm{Ext}_{\mathrm{SL}_2(K)}^1(\nabla(r), \nabla(s)) = \begin{cases} K, & \text{if } r - s = 2m \text{ is positive even and } m < p^v \\ & \text{or } m - p^l < p^v < p^l; \\ 0, & \text{otherwise.} \end{cases}$$

where $v = \mathrm{val}_p(s + m + 1)$ and $l = \mathrm{len}_p(m)$.

In fact A. Parker has given a precise recursive description of all higher extensions $\mathrm{Ext}_{\mathrm{SL}_2(K)}^i(\nabla(r), \nabla(s))$, $r, s \geq 0$, in [19], Section 5 (in the equivalent dual formulation for Weyl modules).

6 Coherent Triples

In the sections that follow, particularly Sections 8, 10, and 11 we shall be working extension sequences for 3 part partitions. We here abstract the notion of a coherent triple from the relations given in Section 4.

Definition 6.1. *We fix a three part partition $\lambda = (a, b, c)$. By a coherent triple of extension sequences for λ we mean a triple $((x_i)_{1 \leq i \leq b}, (y_j)_{1 \leq j \leq c}, (z_k)_{1 \leq k \leq c})$ of extension sequences satisfying the following relations:*

$$(T1) \quad \binom{a+i+k}{k} x_i = \binom{a+i+k}{i} z_k, \quad 1 \leq i \leq b, 1 \leq k \leq c;$$

$$(T2) \quad \binom{a+k}{k} y_j = \binom{b+j}{j} z_k, \quad 1 \leq j, k \leq c, j+k \leq c;$$

$$(T3a) \quad \binom{a+i}{i} y_j = \sum_{s=0}^{i-1} \binom{b+j-i}{j-s} \binom{a+i}{s} x_{i-s} + \binom{b+j-i}{j-i} z_i, \quad 1 \leq i \leq j \leq c;$$

$$(T3b) \quad \binom{a+i}{i} y_j = \sum_{s=0}^j \binom{b+j-i}{j-s} \binom{a+i}{s} x_{i-s}, \quad 1 \leq j \leq c, j < i \leq b+j.$$

Thus if $\lambda = (\lambda_1, \dots, \lambda_n)$ is a partition of length n and $(y(t, u)_i)$ is a coherent multi-sequence for λ then for $1 \leq q < r < s \leq n$ we may extract the coherent triple $(x_i), (y_j), (z_k)$ for $(\lambda_q, \lambda_r, \lambda_s)$ given by $x_i = y(q, r)_i$, $y_j = y(r, s)_j$, and $z_k = y(q, s)_k$, for $1 \leq i \leq r$, $1 \leq j, k \leq \lambda_s$.

Remark 6.2. In any coherent triple $((x_i), (y_j), (z_k))$ the extension sequence (z_k) is determined by (x_i) and (y_j) . (Take $i = j$ in relation (3a).)

We have the space $E(\lambda)$ of all coherent triples for $\lambda = (a, b, c)$. It will emerge (as a result of explicit calculation) that in fact $E(\lambda)$ is at most two dimensional.

Note that for $\lambda = (a, b, c)$ the standard multi-sequence for λ gives the standard triple $x_i^{\text{st}} = \binom{a+i}{i}$, $1 \leq i \leq b$, $y_j^{\text{st}} = \binom{b+j}{j}$, $1 \leq j \leq c$, $z_k^{\text{st}} = \binom{a+k}{k}$, $1 \leq k \leq c$.

7 Extensions dimensions for James partitions

Recall that for a partition λ we write $E(\lambda)$ for the space of extension multi-sequences for λ . In this section we determine the dimension of $E(\lambda)$, for λ a James partition. Note that, by Proposition 4.5 this is the dimension of $\text{Ext}_{B(N)}^1(S^d E, K_\lambda)$ (where d is the degree of λ , N is at least the number of parts of λ and E is the natural $G(N)$ -module).

Definition 7.1. A partition λ will be called constrained if $\text{Ext}_{B(N)}^1(S^d E, K_\lambda)$ is at most one dimensional (where d, N and E are as above).

For a James partition λ , this is equivalent to the condition that $E(\lambda)$ is at most one dimensional.

We extend the notion of the definition of the canonical multi-sequence defined in Section 5 for a two part James partition to an arbitrary James partition $\lambda = (\lambda_1, \dots, \lambda_n)$ of length $n \geq 2$.

Definition 7.2. Let $\lambda = (\lambda_1, \dots, \lambda_n)$ be a James partition with $n \geq 2$ parts. We define the James index $\text{JI}(\lambda)$ to be the largest positive integer k such that p^k divides all integers $\binom{\lambda_i + j}{j}$, for $1 \leq i < n$ and $1 \leq j \leq \lambda_{i+1}$.

Definition 7.3. Let $\lambda = (\lambda_1, \dots, \lambda_n)$ be a James partition of length n then we have the canonical coherent multi-sequence $(y^{\text{can}}(r, s)_i)$, where $y^{\text{can}}(r, s)_i$, for $1 \leq r < s \leq n$, $1 \leq i \leq \lambda_s$, is obtained by taking $\binom{\lambda_r + i}{i}/p^{\text{JI}(\lambda)}$ modulo p .

We begin by deriving a concrete description of the canonical multi-sequence.

Let (a, b) be a James 2-part partition. We put $v = \text{val}_p(a+1)$, $w = \text{val}_p(b)$. We write $a = (p^v - 1) + p^v A$ and $b = p^w B$. Thus we have

$$\binom{a+b}{b-1} = \binom{(p^v - 1) + p^v A + p^w B}{p^w B - 1} = \binom{p^w B - 1 + p^v(A+1)}{p^w B - 1}$$

and hence

$$\binom{a+b}{b-1} \equiv 1 \pmod{p} \quad (1)$$

by Lucas's Formula.

Now we have

$$\binom{a+b}{b} = \frac{a+1}{b} \binom{a+b}{b-1}$$

which shows that $\text{val}_p\left(\binom{a+b}{b}\right) \geq v - w$ and also gives

$$p^w B \binom{a+b}{b} = p^v(A+1) \binom{a+b}{b-1}$$

so that

$$B \left[\frac{\binom{a+b}{b}}{p^{v-w}} \right] = (A+1) \binom{a+b}{b-1}$$

so from (1) we get

$$b_w \left[\frac{\binom{a+b}{b}}{p^{v-w}} \right] \equiv a_v + 1 \pmod{p}. \quad (2)$$

In particular $\binom{a+b}{b}/p^{v-w}$ is non-zero modulo p so that

$$\text{val}_p \binom{a+b}{b} = v - w \quad (3).$$

We now assume that $\lambda = (\lambda_1, \dots, \lambda_n)$ is a partition of length $n \geq 2$. We adopt the following notation: $v_i = \text{val}_p(\lambda_i + 1)$, for $1 \leq i < n$, and $l_i = \text{len}_p(\lambda_i)$ for $2 \leq i \leq n$. In the light of (2),(3) we now have the following description of the James index.

Proposition 7.4. *Let $\lambda = (\lambda_1, \dots, \lambda_n)$ be a James partition with $n \geq 2$ parts. Then $\text{JI}(\lambda)$ is the minimum of $v_1 - l_2, \dots, v_{n-1} - l_n$. Moreover we have*

$$y^{\text{can}}(r, s)_i = \begin{cases} ((\lambda_r)_{v_r} + 1)/t, & \text{if } v_r - l_s = \text{JI}(\lambda) \text{ and } i = tp^{l_s}; \\ 0, & \text{otherwise.} \end{cases}$$

We fix a three part James partition (a, b, c) and study the corresponding coherent triples. We set $v = \text{val}_p(a+1)$, $w = \text{val}_p(b+1)$ and $\beta = \text{len}_p(b)$, $\gamma = \text{len}_p(c)$.

We specialise the above to the case of (a, b, c) . (For a two part partition the canonical sequence, at least up to multiplication by scalars, is given by Lemma 5.8.)

Proposition 7.5. *Let $\lambda = (a, b, c)$. Then $\text{JI}(\lambda)$ is the minimum of $v - \beta$ and $w - \gamma$. Moreover the canonical coherent triple $(x_i^{\text{can}}), (y_j^{\text{can}}), (z_k^{\text{can}})$ for λ is given as follows:*

$$x_i^{\text{can}} = \begin{cases} (a_v + 1)/t, & \text{if } v - \beta = \text{JI}(\lambda) \text{ and } i = tp^\beta \text{ for some } 1 \leq t \leq b_\beta; \\ 0, & \text{otherwise} \end{cases}$$

$$y_j^{\text{can}} = \begin{cases} (b_w + 1)/t, & \text{if } w - \gamma = \text{JI}(\lambda) \text{ and } j = tp^\gamma \text{ for some } 1 \leq t \leq c_\gamma; \\ 0, & \text{otherwise} \end{cases}$$

$$z_k^{\text{can}} = \begin{cases} (a_v + 1)/t, & \text{if } v - \gamma = \text{JI}(\lambda) \text{ and } k = tp^\gamma \text{ for some } 1 \leq t \leq c_\gamma; \\ 0, & \text{otherwise.} \end{cases}$$

Remark 7.6. Suppose $\beta > \gamma$. Then $v = \text{len}_p(b + p^\gamma)$ if and only if $b = p^v - 1$. Certainly $v = \text{len}_p(b + p^\gamma)$ if $b = p^v - 1$. Now suppose $v = \text{len}_p(b + p^\gamma)$. We write $b = (p^w - 1) + p^w B$. Then we have $b + p^\gamma = (p^\gamma - 1) + p^w(B + 1)$ so that $\text{len}_p(B + 1) = v - w$. If B does not have the form $p^k - 1$ for some k then also $\text{len}_p(B) = v - w$ and $\text{len}_p(b) = v$, which is impossible since $\beta < v$ by the James condition. Hence we have $B = p^k - 1$ and $\text{len}_p(B + 1) = k = v - w$ so that $b = (p^w - 1) + p^w(p^{v-w} - 1) = p^v - 1$.

We note that in checking the coherence conditions for a multi-sequence $(y(r, s)_i)$ of extension sequences for a James partition sequence λ it is sufficient to check coherent of all relevant triples since in the commuting relation (C), for a commuting quadruple, both sides are zero (by the James condition).

Lemma 7.7. (i) If $\beta = \gamma$ then λ is constrained and

$$x_{p^\beta}^{\text{can}} = z_{p^\gamma}^{\text{can}} = (a_{\beta+1} + 1)y_{p^\gamma}^{\text{can}}.$$

(ii) If $\beta > \gamma$ and $(a, b + p^\gamma)$ is not James then λ is constrained and $(y_j^{\text{can}}) = 0$, $(z_k^{\text{can}}) = 0$.

(iii) If $\beta > \gamma$ and $(a, b + p^\gamma)$ is James then $\dim E(\lambda) = 2$ and in any coherent triple we have $(z_k) = 0$ and (x_i) and (y_j) are multiples of the canonical sequences, for (a, b) and (b, c) .

Proof. We leave it to the reader to check that in (i) and (ii) the canonical triples are as described.

(i) We have $w \leq \beta + 1 \leq v$ so that $v - \beta \geq w - \gamma$. Hence (y_k^{can}) is non-zero and, subtracting a multiple of the canonical sequence from an arbitrary coherent triple we can obtain a coherent triple $(x_i), (y_j), (z_k)$ in which $(y_j) = 0$. Now $a_\beta = p - 1$ so that, taking $i = k = p^\beta$ in (T1), we obtain $x_{p^\beta} = z_{p^\beta}$.

We note that $\beta = \gamma < w$ implies that $b = p^{\beta+1} - 1$. We take $j = p^\beta$, $i = p^{\beta+1}$ in relation (T3b). The only non-zero term x_{i-s} is obtained by taking $s = p^\beta$ and we obtain $\binom{a+p^{\beta+1}}{p^\beta} x_{(p-1)p^\beta} = 0$. Moreover, we have $\binom{a+p^{\beta+1}}{p^\beta} = (p-1)$, by Lucas's Formula. So we have $x_{(p-1)p^\beta} = 0$ and hence $(x_i) = 0$ and therefore also $(z_k) = 0$. Hence $(x_i), (y_j), (z_k)$ is the zero triple and every coherent triple is a multiple of the canonical triple.

(ii) We have $\text{len}_p(b + p^\gamma) \leq \beta + 1$ and, since (a, b) is James but $(a, b + p^\gamma)$ is not James we have $\text{len}_p(b + p^\gamma) = \beta + 1 \geq v$. Hence $\beta + 1 = v$ and so $b = p^v - 1$, by Remark 7.6 and so $\beta + 1 = v = w$. Hence $v - \beta < w - \gamma$, $v - \gamma$ so that $(x_i^{\text{can}}) \neq 0$. Subtracting a multiple of the canonical triple from an arbitrary coherent triple we can thus obtain a coherent triple $(x_i), (y_j), (z_k)$ in which $(x_i) = 0$. Taking $i = j = p^\gamma$ in (T1) we obtain $z_{p^\gamma} = 0$ and hence the sequence $(z_k) = 0$. Taking $j = p^\gamma$, $i = p^{\beta+1}$ in (T3b) we get $\binom{a+p^{\beta+1}}{p^{\beta+1}} y_{p^\gamma} = 0$, i.e., $(a_v + 1)y_{p^\gamma} = 0$, therefore $y_{p^\gamma} = 0$ and hence the sequence (y_j) is 0. Hence $(x_i), (y_j), (z_k)$ is the zero triple and every coherent triple is a multiple of the canonical triple.

(iii) We show first that in this case we have that in any coherent triple $(z_k) = 0$. Since (a, c) is James it is enough to show that $z_{p^\gamma} = 0$. Now this follows by (T1) for $i = k = p^\gamma$, using the fact that $x_{p^\gamma} = 0$, since $\beta > \gamma$ and (a, b) is James. Therefore $\dim E(\lambda) \leq 2$ and so it is enough to show that there is a coherent triple $(x_i), (y_j), (z_k)$ in which (x_i) non-zero and $(y_j) = 0$, $(z_k) = 0$ and one in which $(x_i) = 0$, (y_j) is non-zero and $(z_k) = 0$.

We first check that the triple $(x_i), (y_j), (z_k)$ with

$$x_i = \begin{cases} 1/t, & \text{if } i = tp^\beta, 1 \leq t \leq b_\beta; \\ 0, & \text{otherwise} \end{cases}$$

and $(y_j) = 0$, $(z_k) = 0$ is coherent.

For (T1) we require $\binom{a+tp^\beta+k}{k} = 0$, for $1 \leq k \leq c$, $1 \leq t \leq b_\beta$. This is true since $(a + tp^\beta, c)$ is James, as $\text{val}_p(a + tp^\beta + 1) \geq \beta > \gamma$. In (T2) both sides

are 0. For (T3a) we require

$$\sum_{s=0}^{i-1} \binom{b+j-i}{j-s} \binom{a+i}{s} x_{i-s} = 0$$

for $1 \leq i \leq j \leq c$. However, if in this equation $x_{i-s} \neq 0$ then we have $i-s = tp^\beta$, for some $1 \leq t \leq b_\beta$ and so $i \geq p^\beta$ and this is impossible since $i \leq c$ and $\gamma < \beta$. Hence (T3a) holds.

For (T3b) we require

$$\sum_{s=0}^j \binom{b+j-i}{j-s} \binom{a+i}{s} x_{i-s} = 0$$

for $1 \leq j \leq c$, $j < i \leq b+j$. Now, x_{i-s} is zero unless $i-s = tp^\beta$ and then $\binom{a+i}{s} = \binom{a+tp^\beta+s}{s}$ is zero for $s \neq 0$ (as $(a+tp^\beta, c)$ is James). Hence the condition to be checked is $\binom{b+j-tp^\beta}{j} = 0$ for $1 \leq j \leq c$, and this is true since $(b-tp^\beta, c)$ is James.

It remains to check that the triple $(x_i), (y_j), (z_k)$ with $(x_i) = 0$, with

$$y_j = \begin{cases} 1/t, & \text{if } j = tp^\gamma, 1 \leq t \leq c_\gamma; \\ 0, & \text{otherwise} \end{cases}$$

and $(z_k) = 0$, is coherent.

Certainly (T1) is satisfied since both sides are 0. Also in (T2) both sides are zero since (a, c) and (b, c) are James. We also have that (T3a) holds since both sides are 0 again (since (a, c) is James). Finally, (T3b) reduces to the condition that $\binom{a+i}{i} = 0$ for $tp^\gamma < i \leq b+tp^\gamma$, $1 \leq t \leq c_\gamma$. Now if $a_h + i_h \geq p$ for some $0 \leq h < v$ then $\binom{a+i}{i} = 0$ so we may assume that $i = p^v I$, for some $I \neq 0$. We get

$$p^v I \leq b + tp^\gamma < p^{\beta+1} - 1 + p^w = (p^w - 1) + p^{\beta+1}.$$

This gives $v \leq \beta+1$ but $\beta < v$ (since λ is James) and so $v = \beta+1$ and $I = 1$. Hence, $\text{len}_p(b + p^\gamma) = \beta + 1 = v$. But this is impossible since $(a, b + p^\gamma)$ is James.

□

Lemma 7.8. *If $l_2 = \dots = l_n$ then λ is constrained.*

Proof. We proceed by induction on n . Certainly the result holds for $n = 2, 3$, by Lemmas 5.8 and Lemma 7.7(i). Now assume $n \geq 4$ and the result holds for shorter partitions. Note that $\text{JI}(\lambda_1, \dots, \lambda_n) = v_{n-1} - l$, where $l = l_2 = \dots = l_n$ and so $y^{\text{can}}(n-1, n)_{p^l} \neq 0$.

By subtracting a multiple of the canonical multi-sequence from an arbitrary coherent multi-sequence we can obtain a coherent multi-sequence

$(y(r, s)_i)$ in which $(y(n-1, n)_i) = 0$. By applying the inductive hypothesis to the partition $(\lambda_2, \dots, \lambda_n)$ we obtain $(y(r, s)_i) = 0$ for all $2 \leq r < s \leq n$. In particular the sequences $(y(2, 3)_i), (y(3, 4)_i), \dots, (y(n-1, n)_i)$ are zero. But now applying Lemma 7.7(i) to the coherent triple $(y(1, 2)_i), (y(2, 3)_j), (y(1, 3)_k)$ we obtain that $(y(1, 2)_i) = 0$. Hence the sequences $(y(1, 2)_i), (y(3, 4)_i), \dots, (y(n-1, n)_i)$ are all zero and $(y(r, s)_i)$ is the zero multi-sequence. Hence every coherent multi-sequence is a multiple of the canonical multi-sequence and we are done. \square

Note that for $1 \leq m \leq n$ we have a natural linear map $E(\lambda) \rightarrow E(\lambda_1, \dots, \lambda_m)$, which we call restriction, sending a coherent multi-sequence $(y(r, s)_i)$, $1 \leq r < s \leq n$, $1 \leq i \leq \lambda_s$, to the coherent multi-sequence $(y(r, s)_i)$, $1 \leq r < s \leq m$, $1 \leq i \leq \lambda_s$.

Lemma 7.9. *Suppose that $1 \leq m \leq n-2$ and that $l_{m+1} > l_{m+2}$. Then restriction $E(\lambda) \rightarrow E(\lambda_1, \dots, \lambda_{m+1})$ is surjective.*

Proof. Let $(y(r, s)_i)$ ($1 \leq r < s \leq m+1$) be a coherent multi-sequence for $(\lambda_1, \dots, \lambda_{m+1})$. We define a multi-sequence $(\hat{y}(r, s)_i)$, $1 \leq r < s \leq n$, $1 \leq i \leq \lambda_s$, for λ by

$$\hat{y}(r, s)_i = \begin{cases} y(r, s)_i, & \text{if } 1 \leq r < s \leq m+1, 1 \leq i \leq \lambda_s; \\ 0, & \text{otherwise.} \end{cases}$$

We claim that $(\hat{y}(r, s)_i)$ is coherent. To do this we check that for all $1 \leq r < s < t \leq n$, the triple $(\hat{y}(r, s)_i), (\hat{y}(s, t)_j), (\hat{y}(r, t)_k)$ is coherent.

If $1 \leq r < s < t \leq m+1$ this is true since $(y(r, s)_i)$ is a coherent multi-sequence. Suppose now that $1 \leq r < s \leq m+1$ and $m+1 < t \leq n$. Then the sequences $(\hat{y}(s, t)_j)$ and $(\hat{y}(r, t)_k)$ are identically 0. Moreover we have $l_s \geq l_{m+1} > l_{m+2} \geq l_t$ so that $l_s > l_t$ and the triple $(\hat{y}(r, s)_i), (\hat{y}(s, t)_j), (\hat{y}(r, t)_k)$ is coherent, by Lemma 7.7(ii) and (iii). Finally, if $s > m+1$ then $(\hat{y}(r, s)_i), (\hat{y}(s, t)_j), (\hat{y}(r, t)_k)$ is the zero triple (which is coherent).

Thus $(\hat{y}(r, s)_i)$ is a coherent multi-sequence mapping to $(y(r, s)_i)$ and the restriction map is surjective. \square

Lemma 7.10. *If $1 \leq m \leq n-2$ and $l_{m+2} = \dots = l_n$ then the kernel of restriction $E(\lambda) \rightarrow E(\lambda_1, \dots, \lambda_{m+1})$ has dimension at most 1.*

Proof. In addition to the restriction map $E(\lambda) \rightarrow E(\lambda_1, \dots, \lambda_{m+1})$ we also have the natural map $\rho : E(\lambda) \rightarrow E(\lambda_{m+1}, \dots, \lambda_n)$, taking a coherent multi-sequence $(y(r, s)_i)$, $1 \leq r < s \leq n$, $1 \leq i \leq \lambda_s$ for λ to the coherent multi-sequence $(\bar{y}(r, s)_i)$, $1 \leq r < s \leq n-m$, $1 \leq i \leq \lambda_{m+s}$, for $(\lambda_{m+1}, \dots, \lambda_n)$, where $\bar{y}(r, s)_i = y(m+r, m+s)_i$, for $1 \leq r < s \leq n-m$, $1 \leq i \leq \lambda_{m+s}$.

We obtain an injective map $E(\lambda) \rightarrow E(\lambda_1, \dots, \lambda_{m+1}) \oplus E(\lambda_{m+1}, \dots, \lambda_n)$. But $\dim E(\lambda_{m+1}, \dots, \lambda_n) = 1$, by Lemma 7.8 and the kernel of restriction $E(\lambda) \rightarrow E(\lambda_1, \dots, \lambda_{m+1})$ embeds in $E(\lambda_{m+1}, \dots, \lambda_n)$ so we are done. \square

The next result essentially determines the dimension of the extension spaces for James partitions.

Proposition 7.11. *Suppose that $1 \leq m \leq n-2$ and $l_{m+1} > l_{m+2} = \dots = l_n$. Then we have*

$$\dim E(\lambda) = \begin{cases} \dim E(\lambda_1, \dots, \lambda_{m+1}), & \text{if } m+2 = n \text{ and } \lambda_{m+1} = p^{v_m} - 1; \\ \dim E(\lambda_1, \dots, \lambda_{m+1}) + 1, & \text{otherwise.} \end{cases}$$

Proof. First suppose that $m+2 = n$ and $\lambda_{m+1} = p^{v_m} - 1$. Let $(y(r, s)_i)$ be a coherent multi-sequence in the kernel of restriction $E(\lambda) \rightarrow E(\lambda_1, \dots, \lambda_{m+1})$. Thus the sequences $(y(1, 2)_i), (y(2, 3)_i), \dots, (y(m, m+1)_i)$ are identically 0. We have the coherent triple $(y(m, m+1)_i), (y(m+1, m+2)_j), (y(m, m+2)_k)$. Moreover, $((y(m, m+1)_i)$ is identically 0 and $l_{m+1} > l_{m+2}$ so that, by Lemma 7.7(ii), $(y(m+1, m+2)_j)$ is 0. Hence the sequences $(y(1, 2)_i), (y(2, 3)_i), \dots, (y(n-1, n)_i)$ are 0 and so $(y(r, s)_i)$ is the zero multi-sequence. Thus the kernel of the restriction is 0 in this case and by Lemma 7.9, we have $\dim E(\lambda) = \dim E(\lambda_1, \dots, \lambda_{m+1})$.

We next consider the case $v_m > \text{len}_p(\lambda_{m+1} + p^{l_{m+2}})$. We have the canonical multi-sequence $(y^{\text{can}}(r, s)_i)$ for $(\lambda_{m+1}, \dots, \lambda_n)$ and we make a multi-sequence $(y(r, s)_i)$ for λ by putting $y(m+r, m+s)_i = y^{\text{can}}(r, s)_i$, for $1 \leq r < s \leq n-m$, $1 \leq i \leq \lambda_s$, and all other terms 0. We check that this is a coherent multi-sequence and hence the kernel of restriction $E(\lambda) \rightarrow E(\lambda_1, \dots, \lambda_{m+1})$ is not zero.

We have to check that each triple (r, s, t) with $1 \leq r < s < t \leq n$ gives a coherent triple of extension sequences $(y(r, s)_i), (y(s, t)_j), (y(r, t)_k)$.

If $t \leq m$ then this is the zero triple (which is coherent).

Suppose next that $1 \leq r < s \leq m$, $m+1 \leq t$. Again $(y(r, s)_i), (y(s, t)_j), (y(r, t)_k)$ is the zero triple and so coherent.

Suppose now that $1 \leq r \leq m$, $m+1 \leq s < t \leq n$. Then the sequences $(y(r, s)_i)$ and $(y(r, t)_k)$ are identically 0 and since we have

$$v_r \geq v_m > \text{len}_p(\lambda_{m+1} + p^{l_{m+2}}) \geq \text{len}_p(\lambda_s + p^{l_t})$$

the triple $(y(r, s)_i), (y(s, t)_j), (y(r, t)_k)$ is coherent by Lemma 7.7(iii).

Finally if $m+1 \leq r < s < t \leq n$ then we may write $r = r_0 + m$, $s = s_0 + m$, $t = t_0 + m$ and

$$((y(r, s)_i), (y(s, t)_j), (y(r, t)_k)) = ((y^{\text{can}}(r_0, s_0)_i), (y^{\text{can}}(s_0, t_0)_j), (y^{\text{can}}(r_0, t_0)_k))$$

which is certainly coherent. Hence $(y(r, s)_i)$ is a coherent multi-sequence in the kernel of restriction $E(\lambda) \rightarrow E(\lambda_1, \dots, \lambda_{m+1})$ and so by Lemmas 7.9 and 7.10 we have $\dim E(\lambda) = \dim E(\lambda_1, \dots, \lambda_{m+1}) + 1$.

It remains to consider the case $m+2 < n$ and $\lambda_{m+1} = p^{v_m} - 1$. Once again we use the canonical multi-sequence $(y^{\text{can}}(r, s)_i)$ for $(\lambda_{m+1}, \dots, \lambda_n)$ to define the multi-sequence $(y(r, s)_i)$ for λ with $y(m+r_0, m+s_0)_i = y^{\text{can}}(r_0, s_0)_i$, for $1 \leq r_0 < s_0 \leq n-m$, $1 \leq i \leq \lambda_{m+s_0}$, and all other terms 0.

We need to check that (r, s, t) , with $1 \leq r < s < t \leq n$, gives a coherent triple $((y(r, s)_i), (y(s, t)_j), (y(r, t)_k))$.

If $1 \leq r < s \leq m$ then this is the zero triple.

Now suppose that $1 \leq r \leq m$, $s = m+1 < t \leq n$. Then $(y(r, m+1)_i)$ and $(y(r, t)_k)$ are identically 0. Now $l_{m+2} < l_{m+1}$ so that $\lambda_{m+2} \neq \lambda_{m+1}$ so that $\lambda_{m+2} < p^{v_m} - 1$ and so $v_{m+2} < v_m$. Thus we have

$$v_{m+1} - l_t = v_m - l_t > v_{m+2} - l_t \geq v_{n-1} - l_n = \text{JI}(\lambda_{m+1}, \dots, \lambda_n).$$

Writing $t = m + t_0$ we thus have that $(y^{\text{can}}(1, t_0)_i)$ is the zero sequence, by Proposition 7.4 and hence $(y(m+1, t)_i)$ is the zero sequence and $((y(r, m+1)_i), (y(m+1, t)_j), (y(r, t)_k))$ is the zero triple, and hence coherent.

Now suppose that $r \leq m$, $m+1 < s < t \leq n$. Again we have that the sequences $(y(r, s)_i)$ and $(y(r, t)_k)$ are zero. We claim that $v_r > \text{len}_p(\lambda_s + p^{l_t})$ and hence, by Lemma 7.7(iii), the triple $((y(r, s)_i), (y(s, t)_j), (y(r, t)_k))$ is coherent. We have $v_r \geq v_m$ and $\text{len}_p(\lambda_s + p^{l_t}) \leq \text{len}_p(\lambda_{m+2} + p^{l_t})$ so it is enough to check that $v_{m+1} > \text{len}_p(\lambda_{m+2} + p^{l_t})$. If this fails then we have $\lambda_{m+2} = p^{v_{m+1}} - 1 = p^{v_m} - 1$ and this is not possible since $l_{m+1} > l_{m+2}$.

Finally if $m+1 \leq r < s < t \leq n$ then we may write $r = m+r_0$, $s = m+s_0$, $t = m+t_0$ and

$$((y(r, s)_i), (y(s, t)_j), (y(r, t)_k)) = ((y^{\text{can}}(r_0, s_0)_i), (y^{\text{can}}(s_0, t_0)_j), (y^{\text{can}}(r_0, t_0)_k))$$

which is certainly coherent.

Thus the kernel of restriction $E(\lambda) \rightarrow E(\lambda_1, \dots, \lambda_{m+1})$ is not zero and by Lemmas 7.8 and 7.9 we have $\dim E(\lambda) = \dim E(\lambda_1, \dots, \lambda_{m+1}) + 1$. \square

Definition 7.12. Let $\lambda = (\lambda_1, \dots, \lambda_n)$ be a James partition of length n . We define the segments of λ to be the equivalence classes of $\{1, \dots, n\}$ for the relation $r = s$ if and only if $l_r = l_s$. For $1 \leq r, s \leq n$ we define r and s to be adjacent if they belong to the same segment or if $1 < r < n$, $s = r+1$ and $r+1$ is the only element in its segment and $\lambda_r = p^{v_{r-1}} - 1$. We define the p -segments of λ to be the equivalent classes of $\{1, \dots, n\}$ for the equivalence relation generated by adjacency.

For a James partition λ we define $e(\lambda) = \dim E(\lambda)$.

Corollary 7.13. Let $\lambda = (\lambda_1, \dots, \lambda_n)$ be a James partition of length $n \geq 2$.
(i) If $l_1 = l_2$ then $e(\lambda)$ is the number of p -segments of $\{1, \dots, n\}$.

(ii) If $l_1 > l_2$ then $e(\lambda)$ is one less than the number of p -segments of $\{1, \dots, n\}$.

Proof. We argue by induction on n .

(i) If $l_1 = l_2 = \dots = l_n$ then the result is true by Lemma 7.8, in particular the result holds for $n = 2$. Suppose now that there are at least 2 segments and let $1 \leq r \leq n$ be such that $l_r > l_{r+1} = \dots = l_n$. Then $r \geq 2$. If $n - r \geq 2$ then we have $e(\lambda) = e(\lambda_1, \dots, \lambda_r) + 1$, by Proposition 7.11, and inductively we have that this is one more than the number of p -segments in $(\lambda_1, \dots, \lambda_r)$, i.e., the number of p -segments in λ . So we may assume that $r = n - 1$. If $\lambda_r \neq p^{v_{r-1}} - 1$ then by Proposition 7.11, we have $e(\lambda) = e(\lambda_1, \dots, \lambda_{n-1}) + 1$, i.e., one more than the number of p -segments in $(\lambda_1, \dots, \lambda_{n-1})$ and this is the number of p -segments in λ . If $\lambda_r = p^{v_{r-1}} - 1$ then, by Proposition 7.11 we have $e(\lambda) = e(\lambda_1, \dots, \lambda_{n-1})$, the number of p -segments in $(\lambda_1, \dots, \lambda_{n-1})$, and this is also the number of p -segments in λ .

(ii) If $n = 2$ we have $e(\lambda) = 1$ and λ has two p -segments so the result is correct. If λ has two segments then we have $l_1 > l_2 = \dots = l_n$ so λ also has two p -segments and the result is correct, by Lemma 7.8. Now suppose that λ has at least three p -segments and that $1 \leq r < n$ is such that $l_r > l_{r+1} = \dots = l_n$. Then we have $r \geq 2$. If $n - r \geq 2$ then, by Proposition 7.11, we have $e(\lambda) = e(\lambda_1, \dots, \lambda_r) + 1$, which is the number of p -segments of $(\lambda_1, \dots, \lambda_r)$ and this is one less than the number of p -segments of λ , as required. So we may suppose that $r = n - 1$. Again if $\lambda_r \neq p^{v_{r-1}} - 1$ then Proposition 7.11 gives $e(\lambda) = e(\lambda_1, \dots, \lambda_{n-1}) + 1$, which is the number of p -segments of $(\lambda_1, \dots, \lambda_{n-1})$ and hence one less than the number of p -segments of λ . If $\lambda_r = p^{v_{r-1}} - 1$ then Proposition 7.11 gives $e(\lambda) = e(\lambda_1, \dots, \lambda_{n-1})$ which is one less than the number of p -segments of $(\lambda_1, \dots, \lambda_{n-1})$ and hence also one less than the number of p -segments of λ . □

8 Three part partitions (a, b, c) with (a, b) James and (b, c) not James

We fix a three part partition $\lambda = (a, b, c)$.

We set $\beta = \text{len}_p(b)$, $\gamma = \text{len}_p(c)$, $v = \text{val}_p(a + 1)$ and $w = \text{val}_p(b + 1)$. We have the standard coherent triple of extension sequences given (x_i) , (y_j) , (z_k) , given by $x_i^{\text{st}} = \binom{a+i}{i}$, $y_j^{\text{st}} = \binom{b+j}{j}$, $z_k^{\text{st}} = \binom{a+k}{k}$, for $1 \leq i \leq b$, $1 \leq j, k \leq c$.

Lemma 8.1. *Assume that (a, b) is James and (b, c) is not James. Then λ is non-split if and only if (b, c) is pointed and $v > \text{len}_p(b + p^\gamma)$. In this case*

$E(\lambda)$ is spanned by the standard triple and the triple with $(x_i) = 0$, $(z_k) = 0$ and (y_j) given by

$$y_j = \begin{cases} 1, & \text{if } j = p^\gamma; \\ 0, & \text{otherwise.} \end{cases}$$

Proof. We claim that in any coherent triple $(x_i), (y_j), (z_k)$ we have $(x_i) = 0$ and $(z_k) = 0$. Since (a, b) and (a, c) are James it is enough to show that $x_{p^\beta} = 0$ and that $z_{p^\gamma} = 0$. Suppose first that $w < \beta$. Now $\beta \geq \gamma \geq w$ (since (b, c) is not James) so we may apply relation (T3b) in the case $j = p^w$, $i = p^\beta$. We get

$$0 = \binom{b + p^w - p^\beta}{p^w} x_{p^\beta}$$

(since $x_i = 0$ for $i < p^\beta$). By Lucas's Formula we get $(b_w + 1)x_{p^\beta} = 0$ so that $x_{p^\beta} = 0$ and $(x_i) = 0$. Also by (T3a) for $i = j = p^\gamma$ we have that $z_p^\gamma = 0$ and so $(z_k) = 0$.

We now suppose $\beta \leq w$. Since $\beta \geq \gamma \geq w$ this implies $\beta = w$ and $\beta = \gamma$. Since (a, b) is James we have $a_\beta = p - 1$ and (T1) with $i = k = p^\beta$ gives $x_{p^\beta} = z_{p^\beta}$. But (T3a), with $i = j = p^\beta$ gives $0 = \binom{b}{p^\beta} x_{p^\beta} + z_{p^\beta} = 0$, i.e., $b_\beta x_{p^\beta} + z_{p^\beta} = 0$ and hence $(b_\beta + 1)x_{p^\beta} = 0$. Now $b_\beta = b_w \neq p - 1$ so that $(b_\beta + 1) \neq 0$ and so $x_{p^\beta} = 0$ and (x_i) and (z_k) are the zero sequences.

Assume that (b, c) is split. Then we can subtract a multiple of the standard triple from an arbitrary coherent triple for λ to obtain a coherent triple $(x_i), (y_j), (z_k)$ in which $(y_j) = 0$. But, then by the claim, $(x_i) = 0$ and $(z_k) = 0$ and hence $(x_i), (y_j), (z_k)$ is the zero triple. Hence $E(\lambda)$ consists of multiples of the standard triple and λ is split.

Now suppose that (b, c) is pointed. After subtracting a multiple of the standard triple from any coherent triple we can obtain a coherent triple with $(x_i) = 0$, $(z_k) = 0$ and $y_j = 0$ for $j \neq p^\gamma$.

We consider the triple of extension sequences given by $(x_i) = 0$, $(z_k) = 0$ and

$$y_j^{\text{pt}} = \begin{cases} 1, & \text{if } j = p^\gamma; \\ 0, & \text{otherwise.} \end{cases}$$

Then (T1), (T2), (T3a) hold trivially for this triple and so it is coherent if and only if (T3b) holds, i.e., if and only if $\binom{a+i}{i} y_j^{\text{pt}} = 0$ for all $1 \leq j \leq c$, $j < i \leq b + j$, i.e., if and only if $\binom{a+i}{i} = 0$ for all $p^\gamma < i \leq b + p^\gamma$, i.e., if and only if $a \geq b + p^\gamma$ and $(a, b + p^\gamma)$ is James, i.e., if and only if $v > \text{len}_p(b + p^\gamma)$. It follows that $E(\lambda)$ is spanned by the standard solution if $v = \text{len}_p(b + p^\gamma)$ and by the standard solution and the point solution just constructed if $v > \text{len}_p(b + p^\gamma)$. \square

9 General results for n -rows

Let $\lambda = (\lambda_1, \dots, \lambda_n)$ be a partition of length $n \geq 4$. Let $(y(r, s)_i)$, be a coherent extension multi-sequence for λ . We shall use the commuting relation, Section 4, (C), in the following special case.

For $1 \leq r < s < n$ with $s \neq r + 1$ we have

$$(C') \quad \binom{\lambda_s + j}{j} y(r, r+1)_i = \binom{\lambda_r + i}{i} y(s, s+1)_j, \quad 1 \leq i \leq \lambda_{r+1}, 1 \leq j \leq \lambda_{s+1}.$$

We shall use the notation $v_i = \text{val}_p(\lambda_i + 1)$, for $1 \leq i \leq n-1$ and $l_i = \text{len}_p(\lambda_i)$, for $2 \leq i \leq n$.

Lemma 9.1. *Suppose that $1 \leq r < n$ and $(\lambda_r, \lambda_{r+1})$ is not James. Let $(y(t, u)_i)$ be a coherent multi-sequence for λ and suppose that $(y(r, r+1)_i)$ is either zero or the point extension sequence. Then, for $1 \leq s < n$ with $s \neq r-1, r, r+1$ the extension sequence $(y(s, s+1)_j)$ is zero.*

Proof. If $(\lambda_r, \lambda_{r+1})$ is pointed and $(y(r, r+1)_i)$ is the point sequence we have $v_r < l_{r+1}$ so in all cases we have $y(r, r+1)_{p^{v_r}} = 0$. We have the commuting relation

$$\binom{\lambda_s + j}{j} y(r, r+1)_{p^{v_r}} = \binom{\lambda_r + p^{v_r}}{p^{v_r}} y(s, s+1)_j$$

for $1 \leq j \leq \lambda_{s+1}$. The left hand side is 0 and λ_r has v_r th component in the base p expansion different from $p-1$ so Lucas's Formula $\binom{\lambda_r + p^{v_r}}{p^{v_r}}$ is non-zero and therefore $(y(s, s+1)_j) = 0$. □

We give now a short proof of Weber's main theorem, [23] Theorem 1.2.

Lemma 9.2. *Let $\lambda = (\lambda_1, \dots, \lambda_n)$ be a partition with $n \geq 5$. Assume that there exist two pairs of consecutive rows of λ , say $(\lambda_r, \lambda_{r+1})$ and $(\lambda_s, \lambda_{s+1})$, with $r+2 < s < n$ which are not James pairs. Then λ is split.*

Proof. After subtracting a standard multi-sequence from an arbitrary coherent multi-sequence we can obtain a multi-sequence $(y(t, u)_i)$ in which either $(y(r, r+1)_i) = 0$ (if $(\lambda_r, \lambda_{r+1})$ is split) or $(y(r, r+1)_i)$ is a multiple of the point sequence (if $(\lambda_r, \lambda_{r+1})$ is pointed). By the above Lemma we have $(y(t, t+1)_j) = 0$ for $t \neq r-1, r, r+1$. In particular we have $(y(s, s+1)_j) = 0$. Applying the above Lemma again with s in place of r we get $(y(t, t+1)_j) = 0$ if $t \neq s-1, s, s+1$. Thus we get $(y(t, t+1)_j) = 0$ for all $1 \leq t < n$ and hence $(y(t, u)_i)$ is the zero multi-sequence for λ . We have shown that we can obtain the zero multi-sequence by subtracting from an arbitrary coherent multi-sequence a multiple of the standard multi-sequence. Hence $E(\lambda)$ consists of multiples of the standard multi-sequence and λ is split. □

We formulate this result in the following way.

Remark 9.3. Let $\lambda = (\lambda_1, \dots, \lambda_n)$ be a partition which is not James. Let r be minimal such that $(\lambda_r, \lambda_{r+1})$ is not James. If λ is not split then either:
(i) the partitions $(\lambda_1, \dots, \lambda_r)$ and $(\lambda_{r+3}, \dots, \lambda_n)$ (if $n \geq r+3$) are James; or
(ii) $(\lambda_1, \dots, \lambda_{n-1})$ is James but $(\lambda_{n-1}, \lambda_n)$ is not.

We show now that whether or not λ is split depends heavily on the triple $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ in the first case and on the pair $(\lambda_{n-1}, \lambda_n)$ in the second case.

Proposition 9.4. *Let $\lambda = (\lambda_1, \dots, \lambda_n)$ be a partition which is not James and $r \leq n-1$ minimal such that $(\lambda_r, \lambda_{r+1})$ is not a James pair.*

(i) If $r \leq n-2$, then the restriction map $E(\lambda) \rightarrow E(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ is injective.

Equivalently, if $(y(t, u)_i)$ is a coherent multi-sequence in which the extension sequences $(y(r, r+1)_i)$ and $(y(r+1, r+2)_i)$ are zero then $(y(t, u)_i)$ is identically zero. In particular if $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ is split then λ is split.

(ii) If $r = n-1$, then the restriction map $E(\lambda) \rightarrow E(\lambda_{n-1}, \lambda_n)$ is injective.

Equivalently, if $(y(t, u)_i)$ is a coherent multi-sequence in which the extension sequence $(y(n-1, n)_i)$ is zero then $(y(t, u)_i)$ is identically zero. In particular if $(\lambda_{n-1}, \lambda_n)$ is split then λ is split.

(iii) If $r = n-1$, we also have that the restriction map $E(\lambda) \rightarrow E(\lambda_{n-2}, \lambda_{n-1}, \lambda_n)$ is injective.

Equivalently, if $(y(t, u)_i)$ is a coherent multi-sequence in which the extension sequences $(y(n-2, n-1)_i)$ and $(y(n-1, n)_i)$ are zero then $(y(t, u)_i)$ is identically zero. In particular if $(\lambda_{n-2}, \lambda_{n-1}, \lambda_n)$ is split then λ is split.

Proof. We give the details of the proof of (i). The proofs of (ii) and (iii) are similar. Let $(y(t, u)_i)$ be a coherent multi-sequence for λ in which $(y(r, r+1)_i)$ and $(y(r+1, r+2)_i)$ are zero. By Lemma 9.1 we have then that $(y(t, t+1)_i) = 0$ for $t \neq r-1$. Moreover, by Lemma 8.1 we get that $(y(r-1, r)_i) = 0$, since $(\lambda_{r-1}, \lambda_r)$ is James. Hence we have $(y(t, t+1)_i) = 0$ for all $1 \leq t < n$ and $(y(t, u)_i)$ is the zero multi-sequence. This proves everything. \square

We finish this section with the case where in which the first pair of rows of λ which is not a James pair is the last two rows.

Proposition 9.5. *Suppose $n \geq 3$ and that $\lambda = (\lambda_1, \dots, \lambda_n)$ is not a James partition, but $(\lambda_1, \dots, \lambda_{n-1})$ is James.*

The partition λ is non-split if and only if $(\lambda_{n-1}, \lambda_n)$ is pointed and $v_{n-2} > \text{len}_p(\lambda_{n-1} + p^{l_n})$. In this case $E(\lambda)$ is spanned by the standard multi-sequence and the point sequence $(y^{\text{pt}}(r, s)_i)$, given by

$$y^{\text{pt}}(r, s)_i = \begin{cases} 1, & \text{if } (r, s) = (n-1, n), i = p^{l_n}; \\ 0, & \text{otherwise.} \end{cases}$$

Proof. Let λ be non-split. By Proposition 9.4(iii) and Lemma 8.1 we can assume that $(\lambda_{n-1}, \lambda_n)$ is pointed and $v_{n-2} > \text{len}_p(\lambda_{n-1} + p^{l_n})$. It remains to prove that the point multi-sequence $(y^{\text{pt}}(r, s)_i)$, as described in the statement of the lemma, is coherent.

Notice that for (s, t) with $s < t < n - 1$ the pairs (s, t) and $(n - 1, n)$ are related via the commuting relation (C). Since for any such (s, t) the partition (λ_s, λ_t) is James and $(y(s, t)_i) = 0$ we have that both sides in the commuting relation are zero and so this relation is satisfied.

Hence it remains to check the coherence conditions for the triples $(\lambda_q, \lambda_{n-1}, \lambda_{n-2})$ with $q \leq n - 2$. For $q \leq n - 2$ we have that $v_q \geq v_{n-2} > \text{len}_p(\lambda_{n-1} + p^{l_n})$. Therefore by Lemma 8.1 our multi-sequence satisfies the coherent conditions for all these triples. \square

Therefore, from now on, we need only consider a partition λ which contains a non-split triple $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ such that $(\lambda_r, \lambda_{r+1})$ is not James but $(\lambda_1, \dots, \lambda_r)$ and $(\lambda_{r+3}, \dots, \lambda_n)$ are James. We shall take up our analysis in these terms in Section 12, after dealing with the remaining cases of 3-part partitions in Sections 10 and 11.

10 Partitions (a, b, c) with (a, b) split.

We continue our analysis of $E(\lambda)$ for a 3 part partition $\lambda = (a, b, c)$. We have already dealt with the case in which λ is James, in Section 7 and with the case where λ is not James and (a, b) is James in Section 8. So we assume from now on that (a, b) is not James. In this section we consider the case in which (a, b) is split.

As usual we write $\beta = \text{len}_p(b)$ and $\gamma = \text{len}_p(c)$ for p -adic lengths of b and c and $v = \text{val}_p(a + 1)$ and $w = \text{val}_p(b + 1)$ for the p -adic evaluations of $a + 1$ and $b + 1$.

Since (a, b) is not James we have $\beta \geq v$. Moreover since (a, b) is split, it is not pointed.

Lemma 10.1. *If (a, c) is James, i.e., $\gamma < v$, then $\lambda = (a, b, c)$ is split.*

Proof. Let $(x_i), (y_j), (z_k)$ be a coherent triple for $\lambda = (a, b, c)$. Since (a, b) is split after subtracting a multiple of the standard triple we obtain a coherent triple, which we also denote $(x_i), (y_j), (z_k)$, in which $(x_i) = 0$. By Remark 6.2 It is enough to show that $(y_j) = 0$.

Since $\gamma < v$ we have that $c < p^v$ and so we may take $i = p^v$ in (T3b) to get

$$\binom{a+p^v}{p^v} y_j = 0$$

and $\binom{a+p^v}{p^v} = a_v + 1 \neq 0$ so that $y_j = 0$ for $j \leq c$.

We have shown that from any coherent triple we may subtract a multiple of the standard coherent triple to obtain the zero triple, i.e., all coherent triples are standard, $E(\lambda)$ is one dimensional, and λ is split. \square

We therefore assume now on that $\gamma \geq v$, i.e., (a, c) is not James.

Lemma 10.2. *If $(a + p^v, b)$ is not James then $\lambda = (a, b, c)$ is split.*

Proof. We must show that every coherent triple is standard. After subtracting a multiple of the standard triple from an arbitrary coherent triple we obtain a coherent triple $(x_i), (y_j), (z_k)$ with $(x_i) = 0$ and it is enough to prove that this triple is zero.

Taking $i = k = p^t < p^v$ in relation (T1) we get

$$0 = \binom{a + p^t + p^t}{p^t} z_{p^t}$$

and, since $a_t = p - 1$ (as $t < v$) we get $(a + p^t + p^t)_t = 1$ and hence, by Lucas's Formula, $\binom{a+p^t+p^t}{p^t} = 1$ and $z_{p^t} = 0$. Now suppose $v \leq t \leq \gamma$. From (T1), we get

$$0 = \binom{a + i + p^t}{p^t} z_{p^t}$$

for $1 \leq i \leq b$. If $t = v$ then we have $\binom{a+p^v+i}{p^v} \neq 0$ for some $1 \leq i \leq b$ (since $(a + p^v, b)$ is not James) and we get $z_{p^v} = 0$. If $v < t$ then for $i = p^v$ we have $\binom{a+i+p^t}{p^t} = \binom{a+p^v+p^t}{p^t} = a_v + 1 \neq 0$, and again $z_{p^t} = 0$. Hence $(z_k) = 0$.

For $1 \leq t < v$, choosing $i = p^v$, $j = p^t$ in (T3b) we get $y_{p^t} = 0$. For $v \leq t \leq \gamma$ we take $i = p^v$, $j = p^t$ in (T3a) to get $y_{p^t} = 0$. Hence $(y_j) = 0$ and we are done. \square

Remark 10.3. (i) Therefore from now on we can assume that $\text{val}_p(a + p^v + 1) > \beta$ (i.e. $(a, b + p^v)$ is James), which means that $a_t = p - 1$ for $0 \leq t \leq \beta, t \neq v$ and $a_v = p - 2$.

(ii) Recall that we are assuming that $\gamma \geq v$.

These assumptions will be in force for the rest of this section.

Lemma 10.4. *Assume that (a, b) is split. Under the assumptions of Remark 10.3, in any coherent triple $(x_i), (y_j), (z_k)$ for $\lambda = (a, b, c)$ we have:*

- (i) $x_i = 0$ for $i \neq p^v$; and
- (ii) $z_k = 0$ for $k \neq p^v$.

Proof. Suppose $1 \leq i \leq b$. If $\binom{a+i}{i} \neq 0$ then we have $a_h + i_h \leq p - 1$ for all $0 \leq h \leq \beta$, by Remark 3.1. But by Remark 10.3 the only possibility is $i = p^v$. This proves that $x_i^{\text{st}} = 0$, for $1 \leq i \leq b$, $i \neq p^v$ and also proves that $z_k^{\text{st}} = 0$, for $1 \leq k \leq c$, $k \neq p^v$. Moreover, since (a, b) is split (x_i) is a multiple of the standard sequence and we have (i).

Since (a, c) is not James, it is either split or pointed. If (a, c) is split then (z_k) is a multiple of the standard sequence and hence $z_k = 0$ for $k \neq p^v$. If (a, c) is pointed then (z_k) is a linear combination of the sequences (z_k^{st}) and (z_k^{pt}) . Now $z_k^{\text{st}} = z_k^{\text{pt}} = 0$ for $k \neq p^v, p^\gamma$. If $\gamma = v$ we are done. If $\gamma > v$ we take $i = j = p^\gamma$ in (T1) to get

$$\binom{a + p^\gamma + p^\gamma}{p^\gamma} z_{p^\gamma} = 0.$$

Now $a_\gamma = p - 1$ and we get $\binom{a + p^\gamma + p^\gamma}{p^\gamma} = 1$ so that $z_{p^\gamma} = 0$ and we are done. \square

Remark 10.5. Assume that $(a + p^v, b)$ is James. If $(x_i), (y_j), (z_k)$ is a triple of extension sequences for λ and $x_i = 0$ for $i \neq p^v$ and $z_k = 0$ for $k \neq p^v$ then relation (T1) is satisfied. In fact both sides of the equation (T1) are zero. If $x_i \neq 0$ then $i = p^v$ and in that case $\binom{a+i+k}{k} = 0$ (since $(a + p^v, b)$ is James). This shows that the left hand side is 0. The argument also shows that the right hand side is 0.

Remark 10.6. Assume that (a, b) is split. Under the assumptions of Remark 10.3, we have by Lemma 10.4 that in any coherent triple $(x_i), (y_j), (z_k)$ for $\lambda = (a, b, c)$, $x_i = 0$ for $i \neq p^v$. Here we give a very simple expression of the relations (T3a) and (T3b) for these partitions. If a term $\binom{a+i}{s} x_{i-s} \neq 0$ appearing in the right hand side is non-zero then we have $i - s = p^v$ so that $\binom{a+i}{s} = \binom{a+p^v+s}{s} \neq 0$. Since $s \leq c$ and $(a + p^v, c)$ is James the only possibility is $s = 0$. Hence the equations take the following forms

$$\binom{a+i}{i} y_j = \binom{b+j-i}{j} x_i + \binom{b+j-i}{j-i} z_i \quad (*)$$

for $1 \leq i \leq j \leq c$, and

$$\binom{a+i}{i} y_j = \binom{b+j-i}{j} x_i \quad (**)$$

for $1 \leq j \leq c$, $j < i \leq b + j$.

It will be easier sometimes to use these simplified forms of (T3a) and (T3b) in order to prove that certain partitions λ are not split.

Lemma 10.7. *If (a, b) is split and (b, c) is James then $\lambda = (a, b, c)$ is non-split if and only if $\gamma = v$, $c_v = 1$ and $\text{len}_p(b + p^v) < \text{val}_p(a + p^v + 1)$.*

In this case $E(\lambda)$ is spanned by the standard triple and the triple $(x_i), (y_j), (z_k)$ with $(x_i) = 0$, $y_{p^v} = 1$, $z_{p^v} = -1$ and $y_k = z_k = 0$ for $k \neq p^v$.

Proof. Assume that λ is non-split. Then there exists a non-standard coherent triple $(x_i), (y_j), (z_k)$ for λ and, subtracting a multiple of the standard triple we obtain a non-zero coherent triple, also denoted $(x_i), (y_j), (z_k)$, in which $(x_i) = 0$. Taking $i = j = p^v$ in (T3a) we obtain, via Remark 10.3, that $(p-1)y_{p^v} = z_{p^v}$, i.e., $y_{p^v} + z_{p^v} = 0$. If $\gamma > v$ then $y_{p^v} = 0$ since (b, c) is James and so $z_{p^v} = 0$. Moreover, taking $i = p^v$ and $j = p^\gamma$ in (T3a) we get that $y_{p^\gamma} = 0$ and so λ is split. Therefore we must have that $v = \gamma$.

Now if $p^v < tp^v \leq c$ then, taking $i = p^v$, $j = tp^v$ in (T3a) we get $-y_{tp^v} = 0$ since (b, c) is James. But (y_j) is an extension sequence for the James partition (b, c) so this is only possible if $(y_j) = 0$. But then $(x_i), (y_j), (z_k)$ is the zero triple and λ is split. Hence (if λ is non-split) we must have $c_v = 1$.

Moreover, we have $y_k = z_k = 0$ for $k \neq p^v$ and $y_{p^v} + z_{p^v} = 0$. Hence λ is non-split if and only if $E(\lambda)$ is spanned by the standard coherent triple and the triple $(x_i), (y_j), (z_k)$ with $(x_i) = 0$, $y_{p^v} = 1$, $z_{p^v} = -1$ and $y_k = z_k = 0$ for $k \neq p^v$.

It remains to determine when this triple is coherent. The condition (T1) is satisfied by Remark 10.5. The condition (T2) is that $\binom{a+k}{k} = 0$ for $p^v + k \leq c$ (using the fact that (b, c) is James). This implies that $k < p^v$, since $c_v = c_\gamma = 1$ and so (T2) holds.

We now consider (T3a). For $i = j$ it is enough to check (T3a) for $i = j = p^v$ and we have already done this. For $i < j$ the right hand side is always zero (since (b, c) is James) and it is enough to take $j = p^v$. But then $\binom{a+i}{i} = 0$ so that (T3a) holds.

Now (T3b) holds if and only if we have

$$\binom{a+i}{i} = 0$$

for $p^v < i \leq p^v + b$. We claim that this is exactly the condition for $(a+p^v, b+p^v)$ to be James, i.e., $\text{len}_p(b+p^v) < \text{val}_p(a+p^v+1)$. First assume that this condition holds. Then we get $a_t = p-1$ for all $p^v < p^t \leq p^v + b$ and so $(a+p^v)_t = p-1$ for all $p^t \leq b+p^v$, i.e., $(a+p^v, b+p^v)$ is James. We leave it to the reader to check the reverse and thus complete the proof. \square

Therefore from now on we consider cases in which $\gamma \geq w$ (i.e., (b, c) is not James). Recall that we also always assuming that $\gamma \geq v$.

Remark 10.8. Assume that $\gamma \geq w$ (and so (b, c) is not James). Then after subtracting a multiple of a standard solution from an arbitrary coherent triple we can get a coherent triple $(x_i), (y_j), (z_k)$ in which $y_j = 0$ for $j \neq p^\gamma$ (with y_{p^γ} also 0 if (b, c) is split).

Lemma 10.9. *If $\gamma \geq w > v$ then $\lambda = (a, b, c)$ is split.*

Proof. By Remark 10.8, we can subtract a standard triple from an arbitrary coherent triple to obtain a coherent triple $(x_i), (y_j), (z_k)$ in which $y_j = 0$ for $j \neq p^\gamma$. Also by Lemma 10.4, we have $x_i = 0$ for $i \neq p^v$ and $z_k = 0$ for $k \neq p^v$. Hence, it is enough to prove that $x_{p^v} = z_{p^v} = y_{p^\gamma} = 0$.

First notice that $y_{p^w} = 0$. This is certainly true if $\gamma > w$. If $\gamma = w$ then (b, c) is split and so we have again that $y_{p^w} = 0$.

Setting now $i = p^v$ and $j = p^w$ in relation (T3a) we have that,

$$0 = \binom{b + p^w - p^v}{p^w} x_{p^v} + \binom{b + p^w - p^v}{p^w - p^v} z_{p^v}.$$

Moreover, we have that $\binom{b + p^w - p^v}{p^w} = (b_w + 1)$. Also, $\binom{b + p^w - p^v}{p^w - p^v} = 0$ as $(b - p^v + p^w)_v = p - 2$ and $(p^w - p^v)_v = p - 1$. Hence we conclude that $(b_w + 1)x_{p^v} = 0$ and so $x_{p^v} = 0$. Now taking $i = j = p^v$ in (T3a) we get that $z_{p^v} = 0$, since $y_{p^v} = x_{p^v} = 0$.

Finally, for $i = p^v$ and $j = p^\gamma$ in (T3a) we get again that $\binom{a + p^v}{p^v} y_{p^\gamma} = 0$, and since $a_v = p - 2$ we obtain $-y_{p^\gamma} = 0$ and we are done, i.e., we have shown that $E(\lambda)$ is spanned by the standard triple. \square

Therefore we can assume from now on that $\gamma \geq v \geq w$.

Lemma 10.10. *If $\gamma > v > w$ then $\lambda = (a, b, c)$ is split.*

Proof. Again after subtracting a standard triple from an arbitrary coherent triple we can obtain a coherent triple $(x_i), (y_j), (z_k)$ in which $y_j = 0$ for $j \neq p^\gamma$. Also by Lemma 10.4 we have that $x_i = 0$ for $i \neq p^v$ and $z_k = 0$ for $k \neq p^v$ and so it is enough to prove that $x_{p^v} = z_{p^v} = y_{p^\gamma} = 0$.

Since $\gamma > v > w$ we have that $p^v + p^w \leq c$. Applying relation (T2) with $j = p^w$ and $k = p^v$, we have

$$0 = \binom{b + p^w}{p^w} z_{p^v} = (b_w + 1)z_{p^v},$$

and so $z_{p^v} = 0$.

Now taking $i = p^v$ and $j = p^w$ in relation (T3b) we get

$$0 = \binom{b + p^w - p^v}{p^w} x_{p^v} = (b_w + 1)x_{p^v}$$

and so $x_{p^v} = 0$.

Finally, taking $i = p^v$ and $j = p^\gamma$ in (T3a) we get $\binom{a + p^v}{p^v} y_{p^\gamma} = 0$, and since $a_v = p - 2$, we have that $-y_{p^\gamma} = 0$. Hence $E(\lambda)$ is spanned by the standard triple. \square

Therefore we are left with the following three cases $\gamma = v = w$, $\gamma = v > w$ and $\gamma > v = w$.

Note that if $\gamma = v = w$ then (b, c) is split.

Lemma 10.11. *Suppose that (a, b) is split and that (b, c) is split. Then $\lambda = (a, b, c)$ is non-split if and only if $(a + p^v, b)$ is James and :*

(i) *if $b_v = 0$ then $\text{val}_p(b - p^v + 1) > \gamma$ (in particular $b - p^v \geq c$);*

(ii) *if $b_v \neq 0$ then $\gamma = v = w$ and $c_\gamma = 1$.*

If λ is non-split then $E(\lambda)$ is spanned by the standard triple and the triple $(x_i), (y_j), (z_k)$ with $(y_j) = 0$, $x_i = 0$ for $i \neq p^v$, $z_k = 0$ for $k \neq p^v$, $x_{p^v} = 1$ and $z_{p^v} = -b_v$.

Proof. Assume that λ is non-split. Let $(x_i), (y_j), (z_k)$ be a non-standard coherent triple for λ . By subtracting a multiple of the standard triple we obtain a non-zero coherent triple, also denoted $(x_i), (y_j), (z_k)$, in which $(y_j) = 0$ and by Lemma 10.4 $x_i = 0$ for $i \neq p^v$ and $z_k = 0$ for $k \neq p^v$. We apply (T3a) in the form of (*) of Remark 10.6, with $i = j = p^v$ (which we may do since $\gamma \geq v$) and we get $\binom{b}{p^v} x_{p^v} + z_{p^v} = 0$. Hence we must have that $x_{p^v} \neq 0$ and, after scaling, we may assume that $x_{p^v} = 1$ and $z_{p^v} = -b_v$. Now we just need to check the conditions for this triple to be coherent.

We check (T3b) in the form of (**) of Remark 10.6. This is clearly satisfied unless $i = p^v$, in which case we have that (T3b) will be satisfied if and only if, for all $1 \leq j < p^v$ we have $\binom{b - p^v + j}{j} = 0$, i.e., $(b - p^v, p^v - 1)$ is James (in particular $b \geq 2p^v - 1$). We now assume this. Notice that this gives directly that $w \geq v$ and since $v \geq w$ we have that $v = w$.

Condition (T1) is satisfied by Remark 10.5.

We consider the cases: (i) $b_v = 0$ and; (ii) $b_v \neq 0$.

Case (i). Suppose $b_v = 0$. The condition (T2) is satisfied trivially. In (T3a) (in the form of (*) of Remark 10.6) we can assume that $i = p^v$ and the condition is that $\binom{b + j - p^v}{j} = 0$ for all $p^v \leq j \leq c$. However, we also have that $(b - p^v, p^v - 1)$ is James and $\binom{b + p^v - p^v}{p^v} = b_v = 0$ so the condition is precisely that $(b - p^v, c)$ is James, i.e., $\text{val}_p(b - p^v + 1) > \gamma$ (in particular $b - p^v \geq c$).

Case (ii). Suppose $b_v \neq 0$. Then (T2) gives $\binom{b + j}{j} = 0$ for $1 \leq j \leq c - p^v$, i.e., $(b, c - p^v)$ is James. Note that this implies $\text{len}_p(c - p^v) < \text{len}_p(c)$ (since (b, c) is not James) and since $\gamma \geq v = w$ we get that $v = \gamma$ and $c_\gamma = 1$. We now consider (T3a), in the form of (*) of Remark 10.6. Once again we only need to consider the case $i = p^v$ and the condition is then $\binom{b + j - p^v}{j} - \binom{b + j - p^v}{j - p^v} b_v = 0$ for $p^v \leq j \leq c$. The condition holds if $j = p^v$. Writing $j = p^v + t$, with $1 \leq t \leq c - p^v \leq p^v - 1$ the condition is $\binom{b + t}{t + p^v} = \binom{b + t}{t} b_v$, for $1 \leq t \leq c - p^v$ or, since $(b, c - p^v)$ is James, $\binom{b + t}{t + p^v} = 0$, for $1 \leq t \leq c - p^v$. So let $1 \leq t \leq c - p^v$. Hence $\text{len}_p(t) \leq \text{len}_p(c - p^v) < v$ (since $\gamma = v$ and $c_v = 1$). Now we have $\binom{b + t}{t_i} = 0$, for some $0 \leq i < v$, by Lucas's Formula and the fact that

$(b, c - p^v)$ is James. But then, since $i < v$ we have $(t + p^v)_i = t_i$ and so $\binom{b+t}{t+p^v} = 0$, by Lucas's Formula again. \square

We have dealt with the situation in which (b, c) is split and we move on to the case in which (b, c) is pointed.

Lemma 10.12. *Assume that (b, c) is pointed and $\gamma = v > w$. Then $\lambda = (a, b, c)$ is non-split if and only if $\text{len}_p(b + p^v) < \text{val}_p(a + p^v + 1)$. In that case $E(\lambda)$ is spanned by the standard triple and the triple $(x_i), (y_j), (z_k)$ with $(x_i) = 0$, $y_j = z_k = 0$ for $j, k \leq c$ and $j, k \neq p^v$, and $y_{p^v} = 1$, $z_{p^v} = -1$.*

Proof. Assume that λ is not split. Since (b, c) is pointed we may write $c = \hat{c} + p^\gamma$, with $\hat{c} < p^w < p^\gamma$. After subtracting a standard triple from an arbitrary coherent triple and using Lemma 10.4 we can obtain a coherent triple $(x_i), (y_j), (z_k)$ in which $x_i = 0$, $y_j = 0$ and $z_k = 0$ for $i, j, k \neq p^v$. Moreover, taking $i = p^v$, $j = p^w$, in the form $(**)$ of (T3b) from Remark 10.6, we get

$$0 = \binom{b + p^w - p^v}{p^w} x_{p^v} = (b_w + 1) x_{p^v}$$

and so $x_{p^v} = 0$.

Taking $i = j = p^v$ in the form $(*)$ of (T3a) from Remark 10.6, we get

$$\binom{a + p^v}{p^v} y_{p^v} = z_{p^v}$$

and, since $a_v = p - 2$, we get $y_{p^v} + z_{p^v} = 0$. Hence λ is non-split if and only if the triple $(x_i), (y_j), (z_k)$ with $(x_i) = 0$, $y_j = z_k = 0$ for $j, k \leq c$ and $j, k \neq p^v$, and $y_{p^v} = 1$, $z_{p^v} = -1$ is coherent.

Certainly condition (T1) is satisfied. Note that in (T2) the left hand side is 0 since we either have $j \neq p^v$ or $1 \leq k \leq \hat{c} < p^v$. Similarly the right hand side is 0 so that (T2) is satisfied.

We now consider (T3a) in the form $(*)$ from Remark 10.6. For $i = j$ we need only to consider $i = j = p^v$, and this we have already done. So we assume that $i < j$. We may assume that $i = p^v$ or $j = p^v$. For $i = p^v < j \leq c$ since $\gamma = v$ we have $j = p^v + t$ with $1 \leq t \leq \hat{c}$. So the condition reads $0 = \binom{b+j-p^v}{j-p^v} = \binom{b+t}{t}$ i.e., (b, \hat{c}) is James, but this is true by hypothesis, since $\hat{c} < p^w$. For $1 \leq i < j = p^v$ both sides are 0 and the condition holds.

Finally, we consider (T3b) in the form $(**)$ from Remark 10.6. The condition is simply that $\binom{a+i}{i} = 0$ for all $p^v < i \leq b + p^v$. Now we already have $\binom{a+p^v+p^h}{p^h} = a_h + 1 = 0$ for $1 \leq h < v$ and $\binom{a+p^v+p^v}{p^v} = a_v + 2 = 0$. Moreover for $p^v < p^h \leq p^v + b$ we have $\binom{a+p^h}{p^h} = \binom{a+p^v+p^h}{p^h}$, so the condition implies that $\binom{a+p^v+t}{t} = 0$ for all $t = p^h \leq b + p^v$, i.e., $(a + p^v, b + p^v)$ is James. Moreover it is easy to reverse the argument to show that if $(a + p^v, b + p^v)$ is James then (T3b) holds.

□

Lemma 10.13. *Assume that (b, c) is pointed and $\gamma > v = w$. Then $\lambda = (a, b, c)$ is non-split if and only if $(a + p^v, b)$ is James and either:*

(i) $\text{val}_p(b - p^v + 1) > \gamma$; or

(ii) $\text{val}_p(b - p^v + 1) = \gamma$ and $\text{len}_p(b + p^\gamma) < \text{val}_p(a + p^v + 1)$.

If λ is non-split then $E(\lambda)$ is spanned by the standard triple and the triple $(x_i), (y_j), (z_k)$ with $(z_k) = 0$, $x_i = 0$ for $i \neq p^v$, $x_{p^v} = 1$, $y_j = 0$ for $j \neq p^\gamma$ and $y_{p^\gamma} = -b_\gamma$.

Proof. Assume that λ is not split. We write $c = \hat{c} + p^\gamma$ with $\hat{c} < p^v < p^\gamma$.

By Lemma 10.4, from an arbitrary coherent triple we may subtract a multiple of the standard triple to obtain a coherent triple $(x_i), (y_j), (z_k)$ with $x_i = 0$ for $i \neq p^v$, $z_k = 0$ for $k \neq p^v$ and $y_j = 0$ for $j \neq p^\gamma$.

We show first that $z_{p^v} = 0$. Indeed, since $v < \gamma$ the relation (T2) applies for $j = k = p^v$ and we get

$$0 = \binom{b + p^v}{p^v} z_{p^v} = (b_v + 1) z_{p^v}$$

and since $v = w$ we have $b_v + 1 \neq 0$ and so $z_{p^v} = 0$.

Further, taking $i = p^v$, $j = p^\gamma$ in relation (T3a), in the form (*) from Remark 10.6, we get

$$\binom{a + p^v}{p^v} y_{p^\gamma} = \binom{b + p^\gamma - p^v}{p^\gamma} x_{p^v} \quad (\dagger).$$

Since $a_v = p - 2$ the left hand side is $-y_{p^\gamma}$. Thus if $x_{p^v} = 0$ then $y_{p^\gamma} = 0$ and $(x_i), (y_j), (z_k)$ is the zero triple. Hence if λ is non-split there exists a non-standard triple with $x_{p^v} \neq 0$.

Now taking $i = p^v \leq j < p^\gamma$ in (T3a), in the form (*) from Remark 10.6, we deduce that if λ is non-split then $\binom{b - p^v + j}{j} = 0$. Moreover, if $1 \leq j < i = p^v$ in (T3b), in the form (**) from Remark 10.6, we have $y_j = 0$ and so $\binom{b - p^v + j}{j} = 0$. Thus, if λ is non-split, then $\binom{b - p^v + j}{j} = 0$ for all $j < p^\gamma$, i.e., $\text{val}_p(b - p^v + 1) \geq \gamma$. We now assume that this is the case.

But now (†) gives $-y_{p^\gamma} = b_\gamma x_{p^v}$. Thus λ is non-split if and only if the triple $(x_i), (y_j), (z_k)$, with $(z_k) = 0$, $x_i = 0$ for $i \neq p^v$, $x_{p^v} = 1$, $y_j = 0$ for $j \neq p^\gamma$ and $y_{p^\gamma} = -b_\gamma$, is coherent, and in this case $E(\lambda)$ is spanned by the standard triple and this triple. It remains to check when this triple is coherent.

Certainly (T1) is satisfied. Condition (T2) is satisfied, since either $y_j = 0$ or $j = p^\gamma$ and then $\binom{a + k}{k} = 0$ as $1 \leq k \leq \hat{c} < p^v$.

In equation (T3a), in the form (*) from Remark 10.6, the left hand side is 0 unless $i = p^v$ and $j = p^\gamma$ and we have already checked this case. Thus we require that in all other cases the right hand side is 0. As usual, it is enough to check the case in which $i = p^v$ and the condition is then (if λ is

non-split) $\binom{b+j-p^v}{j} = 0$ for all $p^v \leq j \leq c$, $j \neq p^\gamma$. We have $\binom{b-p^v+j}{j} = 0$ for $1 \leq j < p^\gamma$, since we are assuming $\text{val}_p(b-p^v+1) \geq \gamma$. Hence the condition is that for all $1 \leq t \leq \hat{c}$ we have $\binom{b-p^v+p^\gamma+t}{p^\gamma+t} = 0$. However, we may write $b = p^v - 1 + p^\gamma B$, for some $B \geq 1$, and so $\binom{b-p^v+p^\gamma+t}{p^\gamma+t} = \binom{t-1+p^\gamma B+p^\gamma}{t+p^\gamma}$, which from Lucas's Formula is $\binom{t-1}{t}(B+1)$ and therefore 0. Hence (T3a) is satisfied.

Finally we consider (T3b) in the form (**) from Remark 10.6. If the right hand side is non-zero then $i = p^v$ and in this case the right hand side is $\binom{b+j-p^v}{j}$, with $j < p^v$. But $\text{val}_p(b-p^v+1) \geq \gamma$ so that $\binom{b+j-p^v}{j} = 0$. Hence the right hand side is always zero. If $\text{val}_p(b-p^v+1) > \gamma$ then $b_\gamma = 0$ and so $y_{p^\gamma} = 0$. Thus in this case the left hand side is also 0 and the coherence conditions are satisfied. This proves part (i).

So we can assume that $\text{val}_p(b-p^v+1) = \gamma$. As the right hand side in (T3b) is always 0 the condition to be checked is $\binom{a+i}{i}y_j = 0$ for $1 \leq j < i \leq b+j$, i.e., $\binom{a+i}{i} = 0$ for $p^\gamma < i \leq b+p^\gamma$. If $\binom{a+i}{i} \neq 0$ then we have $a_h + i_h \leq p-1$ for all $h \geq 0$, by Remark 3.1. Hence by Remark 10.3 we have $i = i_v p^v + p^{\beta+1}I$, for some I and $i_v = 0$ or 1. Since $p^\gamma < i$ we must have $I \neq 0$. But also $i \leq b+p^\gamma$ gives $p^{\beta+1}I \leq p^{\beta+1} - 1 + p^\gamma < 2p^{\beta+1}$ and so $I = 1$. Hence (T3b) is not satisfied if and only if $p^{\beta+1} \leq b+p^\gamma$ and $\binom{a+p^{\beta+1}}{p^{\beta+1}} \neq 0$, i.e. $\text{len}_p(b+p^\gamma) = \text{val}_p(a+p^v+1)$. So we must have that $\text{len}_p(b+p^\gamma) < \text{val}_p(a+p^v+1)$ and this proves part (ii). \square

We summarise our findings from this section in the following result.

Proposition 10.14. *Let (a, b, c) be a partition with (a, b) split. Then $\lambda = (a, b, c)$ is non split if and only if $(a+p^v, b)$ is James and one of the following holds.*

- (i) *We have $\gamma \geq v = w$ and $\text{val}_p(b-p^v+1) > \gamma$. In this case $E(\lambda)$ is spanned by the standard triple and the coherent triple $(x_i), (y_j), (z_k)$ with $x_{p^v} = 1, x_i = 0$ for $i \neq p^v$ and $(y_j) = 0, (z_k) = 0$.*
- (ii) *We have $\gamma = v = w$, $b_v \neq 0$, and $c_v = 1$. In this case $E(\lambda)$ is spanned by the standard triple and the coherent triple $(x_i), (y_j), (z_k)$ with $x_{p^v} = 1, x_i = 0$ for $i \neq p^v$, with $(y_j) = 0$ and $z_{p^v} = -b_v, z_k = 0$ for $k \neq p^v$.*
- (iii) *We have $\gamma > v = w$, $\text{val}_p(b-p^v+1) = \gamma$, $c = \hat{c} + p^\gamma$ with $\hat{c} < p^v$ and $\text{len}_p(b+p^\gamma) < \text{val}_p(a+p^v+1)$. In this case $E(\lambda)$ is spanned by the standard triple and the coherent triple $(x_i), (y_j), (z_k)$ with $x_{p^v} = 1, x_i = 0$ for $i \neq p^v$, with $y_{p^v} = -b_\gamma, y_j = 0$ for $j \neq p^\gamma$ and with $(z_k) = 0$.*
- (iv) *We have $\gamma = v < w$, $c_\gamma = 1$ and $\text{len}_p(b+p^v) < \text{val}_p(a+p^v+1)$. In this case $E(\lambda)$ is spanned by the standard triple and the coherent triple $(x_i), (y_j), (z_k)$ with $(x_i) = 0$, with $y_{p^v} = 1, y_j = 0$ for $j \neq p^v$ and with $z_{p^v} = -1, z_k = 0$ for $k \neq p^v$.*
- (v) *We have $\gamma = v > w$, $c = \hat{c} + p^v$ with $\hat{c} < p^w$ and $\text{len}_p(b+p^v) < \text{val}_p(a+p^v+1)$. In this case $E(\lambda)$ is spanned by the standard triple and the*

coherent triple $(x_i), (y_j), (z_k)$ with $(x_i) = 0$, with $y_{p^v} = 1, y_j = 0$ for $j \neq p^v$ and with $z_{p^v} = -1, z_k = 0$ for $k \neq p^v$.

Proof. Part (i) follows by Lemma 10.11 (i) and Lemma 10.13 (i). Part (ii) is Lemma 10.11 (ii) and part (iii) is Lemma 10.13 (ii). Now (iv) is Lemma 10.7 and (v) is Lemma 10.12. \square

11 Partitions (a, b, c) with (a, b) pointed.

We continue with our analysis of $E(\lambda)$ for a 3-part partition $\lambda = (a, b, c)$. We deal here with the remaining case in which (a, b) is pointed.

As usual we write $\beta = \text{len}_p(b)$ and $\gamma = \text{len}_p(c)$ for the p -adic lengths of b and c and $v = \text{val}_p(a + 1)$ and $w = \text{val}_p(b + 1)$ for the p -adic evaluations of $a + 1$ and $b + 1$.

Since (a, b) is pointed and may write $b = \hat{b} + p^\beta$ with $\hat{b} < p^v < p^\beta$. Notice that here we have that $w \leq v$ since $b_v = 0$.

Lemma 11.1. *If $\gamma < w$, i.e., (b, c) is James, then $\lambda = (a, b, c)$ is non-split and $E(\lambda)$ is spanned by the standard triple and the triple $(x_i), (y_j), (z_k)$ with $x_{p^\beta} = 1, x_i = 0$ for $i \neq p^\beta$ and $(y_j) = 0, (z_k) = 0$.*

Proof. Let $(x_i), (y_j), (z_k)$ be a coherent triple for λ . Now (a, b) is pointed and by subtracting a multiple of the standard triple, we obtain a coherent triple, which we also denote $(x_i), (y_j), (z_k)$, in which $x_i = 0$ for $i \neq p^\beta$.

We claim that $(y_j) = 0$ and $(z_k) = 0$. Now (b, c) is James and since $\gamma < w \leq v$ so is (a, c) . Thus it is enough to prove that $y_{p^\gamma} = z_{p^\gamma} = 0$.

From (T1) with $i = k = p^\gamma$ we get

$$0 = \binom{a + p^\gamma + p^\gamma}{p^\gamma} z_{p^\gamma}$$

and since $\gamma < v$ we have $a_\gamma = p - 1$ so $\binom{a + p^\gamma + p^\gamma}{p^\gamma} = 1$ and $z_{p^\gamma} = 0$.

From (T3b) with $i = p^v$ and $j = p^\gamma$ we get

$$\binom{a + p^v}{p^v} y_{p^\gamma} = 0$$

and so $(a_v + 1)y_{p^\gamma} = 0$. Since $a_v + 1 \neq 0$ we get $y_{p^\gamma} = 0$ as required.

Thus we will be done provided that we show that the triple $(x_i), (y_j), (z_k)$ described in the statement of the Lemma is coherent. For (T1) we require

$(a+p^\beta+k) = 0$ for $1 \leq k \leq c$, i.e., $(a+p^\beta, c)$ is James. But we have $\text{val}_p(a+p^\beta+1) = v \geq w > \gamma$, so this is indeed the case.

In (T2) both sides are 0.

We have $c < p^w < p^\beta$ so that $x_i = 0$ for $1 \leq i \leq c$ and therefore (T3a) holds.

A summand on the right hand side of (T3b) can only be non-zero if $i-s = p^\beta$ and moreover, since $(a+p^\beta+s) = 0$, for $1 \leq s \leq c$ (as $(a+p^\beta, c)$ is James) so we may restrict attention to the case $s = 0$. Hence the condition (T3b) is that $(b+j-p^\beta) = 0$ for $j \leq c$. But $\text{val}_p(b+1-p^\beta) = \text{val}_p(b+1) > \gamma$ so that $(b-p^\beta, c)$ is James and the condition is satisfied. \square

Lemma 11.2. *If $w \leq \gamma < v$ then $\lambda = (a, b, c)$ is split.*

Proof. Subtracting a multiple of the standard triple from an arbitrary coherent triple for λ we obtain a coherent triple $(x_i), (y_j), (z_k)$ with $x_i = 0$ for $i \neq p^\beta$. Since $\gamma < v$ the partition (a, c) is James so that, to prove $(z_k) = 0$, it suffices to prove that $z_{p^\gamma} = 0$. Now $\gamma < v < \beta$ so, we may apply (T1) with $i = k = p^\gamma$, which gives

$$0 = \binom{a+p^\gamma+p^\gamma}{p^\gamma} z_{p^\gamma}$$

and, again since $\gamma < v$, we have $a_\gamma = p-1$ so that $\binom{a+p^\gamma+p^\gamma}{p^\gamma} = 1$ and $z_{p^\gamma} = 0$.

Taking $i = p^v$ in (T3b) we get $\binom{a+p^v}{p^v} y_j = 0$ for $1 \leq j \leq c$ (as $c < p^v$). Thus we have $(a_v+1)y_j = 0$ and therefore $y_j = 0$ for $1 \leq j \leq c$, i.e., $(y_j) = 0$.

Finally, taking $i = p^\beta, j = p^w$ in (T3b) we get

$$0 = \binom{b+p^w-p^\beta}{p^w} x_{p^\beta} = (b_w+1)x_{p^\beta}$$

and since $b_w+1 \neq 0$ we get $x_{p^\beta} = 0$ and therefore $(x_i) = 0$. We have shown that all triples in $E(\lambda)$ are standard, i.e., λ is split. \square

We therefore assume from now on that $\gamma \geq v$ and so $w \leq v \leq \gamma \leq \beta$.

Lemma 11.3. *If $\text{val}_p(a+p^v+1) < \beta$ then $\lambda = (a, b, c)$ is split.*

Proof. Since (a, b) is pointed we may subtract a standard triple from an arbitrary coherent triple to obtain a coherent triple $(x_i), (y_j), (z_k)$ in which $x_i = 0$ for $i \neq p^\beta$. We show first that $(z_k) = 0$. By Lemma 5.3 it is enough to show that $z_{p^t} = 0$ for $t \leq \gamma$. For $t < v$ this follows directly from relation (T1) with $i = k = p^t$ since $a_t = p-1$. Taking $k = p^v$ in relation (T1) we have that

$$0 = \binom{a+i+p^v}{i} z_{p^v}$$

for $1 \leq i < p^\beta$. Since $\text{val}_p(a + p^v + 1) < \beta$ there is some $1 \leq i < p^\beta$ such that $\binom{a+i+p^v}{i} \neq 0$ and hence $z_{p^v} = 0$. Now suppose that $t > v$. Taking $k = p^t$ and $i = p^v$ in (T1) we have

$$\binom{a+p^v+p^t}{p^v} z_{p^t} = 0$$

and since $\binom{a+p^v+p^t}{p^v} = a_v + 1 \neq 0$ we conclude that $z_{p^t} = 0$ and hence $(z_k) = 0$.

We now show that $(x_i) = 0$. Setting $i = p^\beta$ and $k = p^v$ in (T1) we have

$$\binom{a+p^\beta+p^v}{p^v} x_{p^\beta} = 0$$

i.e., $(a_v + 1)x_{p^\beta} = 0$ so that $x_{p^\beta} = 0$ and hence $(x_i) = 0$.

Finally, setting $i = p^v$ in (T3a) and (T3b) we get that $(a_v + 1)y_j = 0$ for $1 \leq j \leq c$ and hence $(y_j) = 0$.

We have shown that all triples in $E(\lambda)$ are standard, i.e., λ is split. \square

Remark 11.4. (i) Therefore from now on we can assume that $\text{val}_p(a + p^v + 1) \geq \beta$, which means that $a_t = p - 1$ for $0 \leq t < \beta$, $t \neq v$ and $a_v = p - 2$.

(ii) Recall that we are also assuming $w \leq v \leq \gamma \leq \beta$.

These assumptions will be in force for the rest of this section.

Lemma 11.5. *Assume that (a, b) is pointed. Under the assumptions of Remark 11.4, in any coherent triple $(x_i), (y_j), (z_k)$ for $\lambda = (a, b, c)$ we have:*

- (i) $x_i = 0$ for $i \neq p^v, p^\beta$;
- (ii) $z_k = 0$ for $k \neq p^v, p^\beta$;
- (iii) $z_k = 0$ for $k \neq p^v$, if $\gamma < \beta$;

Proof. Let $1 \leq i \leq b$. If $\binom{a+i}{i} \neq 0$ then we have $a_h + i_h \leq p - 1$ for all $0 \leq h \leq \beta$. Hence we must have $i_v = 0$ or 1 and $i_h = 0$ for $0 \leq h < \beta$, $h \neq v$. Moreover, $b_\beta = 1$ so we must have $i = p^v, p^\beta$ or $i = p^v + p^\beta$. However, the last case is not possible since $b = \hat{b} + p^\beta < p^v + p^\beta$. This proves (i) since (x_i) is a linear combination of the standard sequence and the point sequence. It also proves $z_k^{\text{st}} = 0$ for $k \neq p^v, p^\beta$ since $c \leq b$. Hence also $z_k = 0$ for $k \neq p^v, p^\beta$ if (a, c) is split.

Now $v \leq \gamma$ so (a, c) is not James and hence is pointed if it is not split. In this case $v < \gamma$. If $\gamma = \beta$ then since (z_k) is a linear combination of the standard extension sequence and the point sequence we also get $z_k = 0$ for $k \neq p^v, p^\beta$. Now suppose $v < \gamma < \beta$. Applying (T1) with $i = j = p^\gamma$ we get

$0 = \binom{a+p^\gamma+p^\gamma}{p^\gamma} z_{p^\gamma}$. But $a_\gamma = p-1$ so this gives $z_{p^\gamma} = 0$. This proves (ii) and (iii). \square

We consider now the situation in which $\beta = \gamma$.

Lemma 11.6. *Assume that $\beta = \gamma$. Then $E(\lambda)$ is spanned by the standard triple and triples $(x_i), (y_j), (z_k)$ in which $x_i = 0$ for $i \neq p^v$, $y_j = 0$ for $j \neq p^\beta$ and $z_k = 0$ for $k \neq p^\beta$.*

Proof. Since $v < \gamma = \beta$ we have that (b, c) is not James and hence is either split or pointed. Hence after subtracting a multiple of the standard triple from an arbitrary coherent triple for λ we can obtain a coherent triple $(x_i), (y_j), (z_k)$ in which $y_j = 0$ for $j \neq p^\gamma$. By Lemma 11.5 we have $x_i = 0$ for $i \neq p^v, p^\beta$ and $z_k = 0$ for $k \neq p^v, p^\beta$.

We now show that $z_{p^v} = x_{p^\beta} = 0$. Since $v < \beta$ we have $p^v + p^v \leq c$ and we may apply (T2) with $j = k = p^v$ to obtain

$$\binom{b+p^v}{p^v} z_{p^v} = 0.$$

Since (a, b) is pointed we have $b_v = 0$ so that $\binom{b+p^v}{p^v} = 1$ and hence $z_{p^v} = 0$.

Now taking $i = p^\beta$, $k = p^v$ in (T1) we obtain

$$\binom{a+p^\beta+p^v}{p^v} x_{p^\beta} = 0.$$

Hence $(a_v + 1)x_{p^\beta} = 0$ and so $x_{p^\beta} = 0$ and we are done. \square

Lemma 11.7. *If $\beta = \gamma$ and $w < v$ then $\lambda = (a, b, c)$ is split.*

Proof. Let $(x_i), (y_j), (z_k)$ be a coherent triple for λ . By Lemma 11.6, we can subtract a standard triple to obtain a coherent triple, which we also denote $(x_i), (y_j), (z_k)$ in which $x_i = 0$ for $i \neq p^v$, $y_j = 0$ for $j \neq p^\beta$ and $z_k = 0$ for $k \neq p^\beta$.

Taking $i = p^v$ and $j = p^w$ in relation (T3b) we get

$$0 = \binom{b+p^w-p^v}{p^w} x_{p^v}$$

and since $\binom{b+p^w-p^v}{p^w} = b_w + 1 \neq 0$ we get $x_{p^v} = 0$.

Now taking $k = p^\beta$ and $i = p^v$ in (T1) we have

$$0 = \binom{a+p^v+p^\beta}{p^v} z_{p^\beta} = (a_v + 1)z_{p^\beta}$$

and since $a_v + 1 \neq 0$, we get $z_{p^\beta} = 0$.

Finally, taking $i = p^v$ and $j = p^\beta$ in (T3a) we get $(a_v + 1)y_{p^\beta} = 0$, and so $y_{p^\beta} = 0$. Hence $(x_i), (y_j), (z_k)$ is the zero triple and $E(\lambda)$ contains only standard triples, i.e., λ is split. \square

Lemma 11.8. *Suppose $\beta = \gamma$. If $\lambda = (a, b, c)$ is non-split then either $\text{val}_p(a + p^v + 1) > \beta$ or $\text{val}_p(a + p^v + p^\beta + 1) > \beta$.*

Proof. By Lemma 11.3 we have that if (a, b, c) is non-split then we may write $a = (p^\beta - 1) - p^v + p^\beta a'$. We shall assume that $a_\beta \neq p - 1, p - 2$ and show that $\lambda = (a, b, c)$ is split. By Lemma 11.6 it is enough to prove if $(x_i), (y_j), (z_k)$ is a coherent triple for λ in which $x_i = 0$ for $i \neq p^v$, $y_j = 0$ for $j \neq p^\beta$ and $z_k = 0$ for $k \neq p^\beta$ then this triple is 0.

Taking $i = k = p^\beta$ is (T1) we get

$$0 = \binom{a + p^\beta + p^\beta}{p^\beta} z_{p^\beta} = (a_\beta + 2)z_{p^\beta}$$

and since $a_\beta \neq p - 2$ we obtain $z_{p^\beta} = 0$.

Now by (T1) again, this time with $i = p^v$ and $k = p^\beta$, we have

$$\binom{a + p^v + p^\beta}{p^\beta} x_{p^v} = 0$$

and so $(a_\beta + 1)x_{p^v} = 0$. Since $a_\beta \neq p - 1$ we have $x_{p^v} = 0$.

Finally, taking $i = p^v$ and $j = p^\beta$ in (T3a) we get that $(a_v + 1)y_{p^\beta} = 0$ so that $y_{p^\beta} = 0$ and we are done. \square

We now describe the non-split triples for the case $\beta = \gamma$.

Lemma 11.9. *Suppose $\beta = \gamma$. Then $\lambda = (a, b, c)$ is non-split triple if and only if $v = w$, so $b = p^v - 1 + p^\beta$, $c = \hat{c} + p^\beta$ with $\hat{c} < p^v$, and one of the following holds.*

(i) $\text{len}_p(b + p^\beta) < \text{val}_p(a + p^v + 1)$.

In that case $E(\lambda)$ is spanned by the standard triple and the triple $(x_i), (y_j), (z_k)$, with $x_{p^v} = 1$, $x_i = 0$ for $i \neq p^v$, with $y_{p^\beta} = -1$, $y_j = 0$ for $j \neq p^\beta$, and $(z_k) = 0$.

(ii) $\text{len}_p(b + p^\beta) < \text{val}_p(a + p^v + p^\beta + 1)$.

In that case $E(\lambda)$ is spanned by the standard triple and the triple $(x_i), (y_j), (z_k)$ with $x_{p^v} = 1$, $x_i = 0$ for $i \neq p^\beta$, with $y_{p^\beta} = -1$, $y_j = 0$ for $j \neq p^\beta$ and $z_{p^\beta} = 1$, $z_k = 0$ for $k \neq p^\beta$.

Proof. By Lemma 11.7 we have that if λ is not split then $v = w$ and since (a, b) is pointed we have $b = p^v - 1 + p^\beta$. By Lemma 11.6 we can subtract a multiple of a standard triple from a coherent triple to obtain a triple

$(x_i), (y_j), (z_k)$ in which $x_i = 0$ for $i \neq p^v$, $y_j = 0$ for $j \neq p^\beta$ and $z_k = 0$ for $k \neq p^\beta$.

By Lemma 11.8 we have either $\text{val}_p(a + p^v + 1) > \beta$ or $\text{val}_p(a + p^v + p^\beta + 1) > \beta$.

Case (i) : $\text{val}_p(a + p^v + 1) > \beta$.

We note that equations (T3a) and (T3b) in this case take a simplified form. If a term $\binom{a+i}{s}x_{i-s} \neq 0$ appearing in the right hand side is non-zero then we have $i - s = p^v$ so that $\binom{a+i}{s} = \binom{a+p^v+s}{s} \neq 0$. Since $s \leq c$ and $(a + p^v, c)$ is James (as $\text{val}_p(a + p^v + 1) > \beta$) the only possibility is $s = 0$. Hence the equations take the following forms

$$\binom{a+i}{i}y_j = \binom{b+j-i}{j}x_i + \binom{b+j-i}{j-i}z_i \quad (*)$$

for $1 \leq i \leq j \leq c$, and

$$\binom{a+i}{i}y_j = \binom{b+j-i}{j}x_i \quad (**)$$

for $1 \leq j \leq c$, $j < i \leq b + j$.

We have $a_\beta = p - 1$ and so for $i = k = p^\beta$ in relation (T1) we get

$$0 = \binom{a + p^\beta + p^\beta}{p^\beta}z_{p^\beta} = z_{p^\beta}.$$

Now taking $i = p^v$ and $j = p^\beta$ in (T3a) we obtain

$$-y_{p^\beta} = \binom{b + p^\beta - p^v}{p^\beta}x_{p^v}$$

and since $b - p^v = p^\beta - 1$ we have that $\binom{b+p^\beta-p^v}{p^\beta} = 1$ and so we get that $-y_{p^\beta} = x_{p^v}$. Thus our analysis of case (i) will be complete if we check the conditions for the triple $(x_i), (y_j), (z_k)$, with $x_{p^v} = 1$, $x_i = 0$ for $i \neq p^v$, and $y_{p^\beta} = -1$, $y_j = 0$ for $j \neq p^\beta$, and $(z_k) = 0$ to be coherent.

Equation (T1) says that $\binom{a+p^v+k}{k} = 0$, for $1 \leq k \leq c$, and this is true since $\text{val}_p(a + p^v + 1) > \beta = \gamma$ so that $(a + p^v, c)$ is James.

Equation (T2) says that $\binom{a+k}{k} = 0$ for $k \geq 1$ and $p^\beta + k \leq c$. For such k we have $p^\beta + k \leq b = \hat{b} + p^\beta$, hence $k \leq \hat{b} < p^v$ and therefore $\binom{a+k}{k} = 0$.

Now (T3a), in the form (*), says

$$\binom{a+i}{i}y_j = \binom{b+j-i}{j}x_i$$

for $1 \leq i \leq j \leq c$. We may assume that $i = p^v$ or $j = p^\beta$. For $i = p^v$, writing $b = p^v - 1 + p^\beta$ this becomes

$$-y_j = \binom{p^\beta - 1 + j}{j}$$

for $p^v \leq j \leq c$. Both sides are 0 for $j < p^\beta$ and 1 for $j = p^\beta$. For $j > p^\beta$ we may write $j = t + p^\beta$, with $0 < t < p^v$ and then

$$\binom{p^\beta - 1 + j}{j} = \binom{p^\beta - 1 + t + p^\beta}{t + p^\beta} = \binom{t - 1 + p^\beta + p^\beta}{t + p^\beta}$$

which is 0. Hence (T3a) holds in case $i = p^v$. Now suppose $j = p^\beta$. The condition is then

$$-\binom{a + i}{i} = \binom{b + p^\beta - i}{p^\beta} x_i$$

for $1 \leq i \leq p^\beta$. We have already checked this for $i = p^v$ so the condition is that $\binom{a+i}{i} = 0$ for $1 \leq i \leq p^\beta$, $i \neq p^v$. This holds for $i \neq p^\beta$ by applying Lemma 11.5 (i) to the standard triple. For $i = p^\beta$ the condition is that $\binom{a+p^\beta}{p^\beta} = 0$ and this is true as $a_\beta = p - 1$. Hence (T3a) holds.

We now consider (T3b) in the form (**). We may assume $i = p^v$ or $j = p^\beta$. For $i = p^v$ the condition is

$$\binom{a + p^v}{p^v} y_j = \binom{b + j - p^v}{j}$$

for $1 \leq j < p^v$ or, since $y_j = 0$, that $\binom{b+j-p^v}{j} = 0$, for $1 \leq j < p^v$. But $\binom{b+j-p^v}{j} = \binom{p^\beta-1+j}{j}$ so this is certainly true.

For $j = p^\beta$ the condition is

$$-\binom{a + i}{i} = \binom{b + p^\beta - i}{p^\beta} x_i$$

for $p^\beta < i \leq b + p^\beta$. For such i we have $x_i = 0$ so the condition is

$$\binom{a + i}{i} = 0$$

for $p^\beta < i \leq b + p^\beta$. Now if this condition holds then, for $p^v < p^t \leq b + p^\beta$ we have $\binom{a+p^v+p^t}{p^t} = \binom{a+p^t}{p^t} = 0$. But we also have $\binom{a+p^v+p^t}{p^t} = 0$ for $p^t \leq p^v$ so that $(a + p^v, b + p^\beta)$ is James. Suppose conversely that $(a + p^v, b + p^\beta)$ is James and that $\binom{a+i}{i} \neq 0$ for some $p^v < i \leq b + p^\beta$. Then we have $a_h + i_h \leq p - 1$ for all $h \geq 0$ and so $i_v = 0$ or 1 and $i_h = 0$ for $h \neq v$. But then i is 0 or p^v , neither of which is possible since $p^v < i$. Hence (T3b) holds if and only if $(a + p^v, b + p^\beta)$ is James, i.e. $\text{len}_p(b + p^\beta) < \text{val}_p(a + p^v + 1)$.

Case (ii): $\text{val}_p(a + p^v + p^\beta + 1) > \beta$.

In this case we have $a_\beta = p - 2$. Applying (T1) with $i = p^v$, $k = p^\beta$ gives

$$\binom{a + p^v + p^\beta}{p^\beta} x_{p^v} = \binom{a + p^v + p^\beta}{p^v} z_{p^\beta}$$

i.e., $-x_{p^v} = -z_{p^\beta}$ so that $x_{p^v} = z_{p^\beta}$. Taking $i = p^v$, $j = p^\beta$ in (T3a) we get

$$\binom{a + p^v}{p^v} y_{p^\beta} = \binom{b + p^\beta - p^v}{p^\beta} x_{p^v}$$

i.e., $-y_{p^\beta} = \binom{p^\beta + p^\beta - 1}{p^\beta} x_{p^v}$ and hence $-y_{p^\beta} = x_{p^v}$. Thus our analysis of Case (ii) will be complete if we check the conditions for the triple $(x_i), (y_j), (z_k)$, with $x_{p^v} = 1$, $x_i = 0$ for $i \neq p^v$, and $y_{p^\beta} = -1$, $y_j = 0$ for $j \neq p^\beta$, and $z_{p^\beta} = 1$, $z_k = 0$ for $k \neq p^\beta$, to be coherent.

In checking (T1) we may assume that $i = p^v$ or $j = p^\beta$. For $i = p^v$ the condition is that $\binom{a + p^v + k}{k} = 0$ for all $1 \leq k \leq c$, $k \neq p^\beta$ (we have already checked the case $i = p^v$, $k = p^\beta$). However, if $\binom{a + p^v + k}{k} \neq 0$ then we have $(a + p^v)_h + k_h \leq p - 1$ for all $h \geq 0$. This gives $k_h = 0$ for all $h \neq \beta$ and $k = p^\beta$, which is not the case. It remains to check $k = p^\beta$, $1 \leq i \leq c$, $i \neq p^v$. The condition is

$$0 = \binom{a + i + p^\beta}{i}.$$

If $\binom{a + i + p^\beta}{i} \neq 0$ then $(a + p^\beta)_h + i_h \leq p - 1$ for all h . This implies $i_h = 0$ for $h \neq v$ and so $i = p^v$ which is not the case.

We now consider (T2). Suppose $1 \leq j, k \leq c$ and $j + k \leq c$. If $\binom{a + k}{k} y_j \neq 0$ then $j = p^\beta$, but then $p^\beta + k \leq c \leq b$ gives $k < p^v$ so that $\binom{a + k}{k} = 0$. Thus the left hand side of (T2) is 0 in all cases and, similarly so is the right hand side.

We now consider (T3a). We notice again that if a term $\binom{a + i}{s} x_{i-s}$ appearing in the right hand side is non-zero then we have $i - s = p^v$ so that $\binom{a + i}{s} = \binom{a + p^v + s}{s} \neq 0$. Since $\text{val}_p(a + p^v + p^\beta + 1) > \beta$ this can only happen if $s = 0$ or $s = p^\beta$. However for the latter case this would force $i = p^v + p^\beta$ and as $i \leq j \leq c < p^v + p^\beta$ this is illegal. Hence the only possible case is when $s = 0$ and so (T3a) takes again the form

$$\binom{a + i}{i} y_j = \binom{b + j - i}{j} x_i + \binom{b + j - i}{j - i} z_i$$

for $1 \leq i \leq j \leq c$, and

Suppose $i = p^v$. Then the condition is

$$\binom{a + p^v}{p^v} y_j = \binom{b + j - p^v}{j}$$

i.e.,

$$-y_j = \binom{p^\beta - 1 + j}{j}$$

for $p^v \leq j \leq c$. This is clear for $j < p^\beta$ and for $j = p^\beta$. For $p^\beta < j \leq c$ we require $\binom{p^\beta - 1 + j}{j} = 0$. This is clear by writing $j = p^\beta + t$, with $1 \leq t < p^v$, and appealing to Lucas's Formula.

Assume now that $i = p^\beta$. Then the condition is $\binom{a+p^\beta}{p^\beta} y_j = \binom{b+j-p^\beta}{j-p^\beta}$, or $-y_j = \binom{p^v-1+j}{j-p^\beta}$ for $p^\beta \leq j \leq c$. This is certainly true for $j = p^\beta$ and so we require $\binom{p^v-1+j}{j-p^\beta} = 0$ for $p^\beta < j \leq c$. Again this becomes clear on writing $j = p^\beta + t$, $1 \leq t < p^v$, and using Lucas's Formula.

Now suppose $j = p^\beta$. The condition is

$$-\binom{a+i}{i} = \binom{b+p^\beta-i}{p^\beta} x_i + \binom{b+p^\beta-i}{p^\beta-i} z_i$$

for $1 \leq i \leq p^\beta$. If $i < p^v$ then both sides are 0. If $i = p^v$ the equation is $1 = \binom{p^\beta-1+p^\beta}{p^\beta}$, which is true. For $p^v < i < p^\beta$ both sides are 0. For $i = p^\beta$ both sides of the equation are 1.

We now consider (T3b).

Let $j < p^\beta$. Then the left hand side in this relation is 0. Now, we have that if a term $\binom{a+i}{s} x_{i-s}$ appearing in the right hand side is non-zero then we have $i-s = p^v$ so that $\binom{a+i}{s} = \binom{a+p^v+s}{s} \neq 0$. Since $\text{val}_p(a+p^v+p^\beta+1) > \beta$ this can only happen for $s = 0$, $s = p^\beta$ or $s = p^v + p^\beta$. However, we also have that $s \leq j < p^\beta$ and so this leaves $s = 0$ as the only option and so for $j < p^\beta$, (T3b) takes the form

$$0 = \binom{b+j-i}{j} x_i$$

for $1 \leq j < p^\beta$ and $j < i \leq b+j$. Now for $i \neq p^v$ the right hand side is 0 and so it remains to check this relation for $i = p^v$ where we get

$$0 = \binom{b+j-p^v}{j} = \binom{p^\beta-1+j}{j}$$

for $1 \leq j < p^\beta$, which is clearly true.

Assume now that $j > p^\beta$. Then the left hand side again is 0 and if a term $\binom{a+i}{s} x_{i-s}$ appearing in the right hand side is non-zero then we have $i-s = p^v$ so that $\binom{a+i}{s} = \binom{a+p^v+s}{s} \neq 0$. Since $\text{val}_p(a+p^v+p^\beta+1) > \beta$ this can only happen for $s = 0$, $s = p^\beta$ or $s = p^v + p^\beta$. Now since $s \leq c < p^v + p^\beta$ we have that the last case is impossible. Moreover, if $s = 0$ that would force $i = p^v$ which is illegal since $p^\beta < j < i$. Hence this leaves $s = p^\beta$ as the only option and so for $j > p^\beta$, (T3b) takes the form

$$0 = \binom{b+j-i}{j-p^\beta} x_{i-p^\beta}$$

for $1 < p^\beta < j \leq c$ and $j < i \leq b+j$. Now for $i \neq p^v + p^\beta$ the right hand side is 0 and so it remains to check this relation for $i = p^v + p^\beta$ where we get

$$0 = \binom{b+j-p^v-p^\beta}{j-p^\beta}$$

for $p^\beta < j \leq c$. But such j has the form $j = p^\beta + t$ with $0 < t < p^v$ and so we have

$$0 = \binom{b+t-p^v}{t} = \binom{p^\beta-1+t}{t}$$

and since $0 < t < p^v$ this is clearly true.

Hence, it remains to consider the case $j = p^\beta$. Making the same remarks as before for the right hand side of the relation we have that the relation takes the form

$$-\binom{a+i}{i} = x_{i-p^\beta}$$

for $p^\beta < i \leq b + p^\beta$. This is true for $i = p^v + p^\beta$ and so the condition remaining to be checked is $\binom{a+i}{i} = 0$ for $p^\beta < i \leq b + p^\beta$. One easily check (cf. the final step of Case (i)) that this is exactly the condition for $(a + p^v + p^\beta, b + p^\beta)$ to be James, i.e. $\text{len}_p(b + p^\beta) < \text{val}_p(a + p^v + p^\beta + 1)$. \square

We continue with the cases in which $\beta > \gamma$.

Lemma 11.10. *Assume that $v < \gamma < \beta$. Then $E(\lambda)$ is spanned by standard triples and triples $(x_i), (y_j), (z_k)$ in which $x_i = 0$ for $i \neq p^v$, $(y_j) = 0$ and $(z_k) = 0$.*

Proof. Since (a, b) is pointed we may write $b = \hat{b} + p^\beta$ with $\hat{b} < p^v < p^\beta$, in particular we have $b_v = 0$ and $b_\gamma = 0$. Since $\gamma > v$ we can subtract a multiple of a standard triple from an arbitrary coherent triple to obtain a coherent triple $(x_i), (y_j), (z_k)$ in which $y_j = 0$ for $j \neq p^\gamma$. By Lemma 11.5 we have $x_i = 0$ for $i \neq p^v, p^\beta$ and $z_k = 0$ for $k \neq p^v$. Since $v < \gamma$ we have $p^v + p^v \leq c$ and so (T2) with $j = k = p^v$ gives $\binom{b+p^v}{p^v} z_{p^v} = 0$, i.e. $z_{p^v} = 0$ so that the sequence (z_k) is 0. Now taking $i = p^\beta$, $k = p^v$ in (T1) gives $\binom{a+p^\beta+p^v}{p^v} x_{p^\beta} = 0$ i.e., $(a_v + 1)x_{p^\beta} = 0$ so that $x_{p^\beta} = 0$. Finally, taking $i = p^v$ and $j = p^\gamma$ in (T3a) we get

$$-y_{p^\gamma} = \binom{b+p^\gamma-p^v}{p^\gamma} x_{p^v}$$

i.e., $-y_{p^\gamma} = b_\gamma x_{p^v}$ and since $b_\gamma = 0$ we get $y_{p^\gamma} = 0$ and therefore the sequence (y_j) is 0. \square

Lemma 11.11. *If $w < v < \gamma < \beta$ then $\lambda = (a, b, c)$ is split.*

Proof. By Lemma 11.10 we can subtract a standard triple to obtain a coherent triple $(x_i), (y_j), (z_k)$ with $x_i = 0$ for $i \neq p^v$, $(y_j) = 0$ and $(z_k) = 0$. Taking $i = p^v$ and $j = p^w$ in (T3b) gives $0 = \binom{b+p^w-p^v}{p^w} x_{p^v}$ i.e., $0 = x_{p^v}(b_w + 1)$ and so $x_{p^v} = 0$. Hence $(x_i), (y_j), (z_k)$ is the zero triple and $E(\lambda)$ contains only standard triples, i.e., λ is split. \square

Lemma 11.12. *If $w = v < \gamma < \beta$ then $\lambda = (a, b, c)$ is non-split and $E(\lambda)$ is spanned by standard triples and the triple $(x_i), (y_j), (z_k)$ in which, $x_{p^v} = 1, x_i = 0$ for $i \neq p^v$, $(y_j) = 0$ and $(z_k) = 0$*

Proof. Since (a, b) is pointed and $w = v$ we have $b = p^v - 1 + p^\beta$. This implies that $(b - p^v, c)$ is James and that $b_v = 0$. By Lemma 11.10 it is enough to prove that the given triple is coherent. Condition (T1) reduces to the statement that $\binom{a+p^v+k}{k} = 0$ for $1 \leq k \leq c$ and this is true since $(a + p^v, c)$ is James. Condition (T2) is satisfied since all terms are 0.

Condition (T3a) holds provided that

$$0 = \binom{b+j-p^v-s}{j-s} \binom{a+p^v+s}{s}$$

for $i = p^v + s \leq j \leq c$. If $s \neq 0$ then $\binom{a+p^v+s}{s} = 0$ since $(a + p^v, c)$ is James and so (T3a) reduces to the condition that

$$0 = \binom{b+j-p^v}{j}$$

i.e., $\binom{p^{\beta-1}+j}{j} = 0$, for $p^v \leq j \leq c$, which holds since $(p^\beta - 1, c)$ is James.

Finally, for (T3b) we require

$$0 = \binom{b+j-p^v-s}{j-s} \binom{a+p^v+s}{s}$$

for $1 \leq j \leq c$, $j < p^v + s \leq b + j$, and with $s \leq j$. Again this reduces to $0 = \binom{b+j-p^v}{j}$ i.e., $\binom{p^{\beta-1}+j}{j} = 0$, and this holds since $(p^\beta - 1, c)$ is James. \square

We finish off with the last cases where $\gamma = v \geq w$.

Lemma 11.13. *If $w = v = \gamma$ then $\lambda = (a, b, c)$ is non-split and $E(\lambda)$ is spanned by standard triple and the triple $(x_i), (y_j), (z_k)$ in which $x_{p^v} = 1$, $x_i = 0$ for $i \neq p^v$, $(y_j) = 0$ and $(z_k) = 0$.*

Proof. Since $\gamma = w$ the partition (b, c) is split. Subtracting a standard triple from an arbitrary coherent triple we can obtain a triple $(x_i), (y_j), (z_k)$ in which $(y_j) = 0$. Further, by Lemma 11.5, we have $x_i = 0$ for $i \neq p^v, p^\beta$ and $z_k = 0$ for $k \neq p^v$.

Taking $i = j = p^v$ in (T3a) we have $\binom{b}{p^v}x_{p^v} + z_{p^v} = 0$ and $b_v = b_w = 0$ (since (a, b) is pointed) and hence $z_{p^v} = 0$. Now taking $i = p^\beta$, $k = p^v$ in (T1) we get $\binom{a+p^\beta+p^v}{p^v}x_{p^\beta} = 0$, i.e. $(a_v + 1)x_{p^\beta} = 0$ and so $x_{p^\beta} = 0$.

It remains to check that the triple $(x_i), (y_j), (z_k)$ described in the statement of the lemma is coherent. The condition (T1) reduces to the statement that $\binom{a+p^v+k}{k} = 0$ for all $1 \leq k \leq c$, and this is true since $(a + p^v, c)$ is James. Condition (T2) is satisfied trivially.

We now consider (T3a). We check that $0 = \sum_{s=0}^{i-1} \binom{b+j-i}{j-s} \binom{a+i}{s} x_{i-s}$, for $1 \leq i \leq j \leq c$. Now x_{i-s} is zero unless $i - s = p^v$ and then $\binom{a+i}{s} = \binom{a+p^v+s}{s}$ is zero for $s \neq 0$ (as $(a + p^v, c)$ is James). Hence the condition to be checked is $\binom{b+j-p^v}{j} = 0$, for $p^v \leq j \leq c$. But, since $v = \gamma$ we have $b = p^v - 1 + p^\beta$ and $(b - p^v, c)$ is James so that $\binom{b+j-p^v}{j} = 0$, for $p^v \leq j \leq c$, as required.

Similarly, condition (T3b) reduces to the condition that $\binom{b+j-p^v}{j} = 0$ for all $1 \leq j < p^v$ and this is also guaranteed by the fact that $(b - p^v, c)$ is James. □

Lemma 11.14. *If $w < v = \gamma$ and (b, c) is split then $\lambda = (a, b, c)$ is split.*

Proof. Subtracting a standard triple from an arbitrary coherent triple for λ we can obtain a coherent triple $(x_i), (y_j), (z_k)$ in which $(y_j) = 0$. By Lemma 11.5 we have $x_i = 0$ for $i \neq p^v, p^\beta$ and $z_k = 0$ for $k \neq p^v$.

Taking $i = p^v$ and $j = p^w$ in (T3b) we have

$$0 = \binom{b+p^w-p^v}{p^w} x_{p^v} = (b_w + 1)x_{p^v}$$

and hence $x_{p^v} = 0$.

Taking $i = p^\beta$ and $j = p^w$ in (T3b) we have

$$0 = \binom{b+p^w-p^\beta}{p^w} x_{p^\beta} = (b_w + 1)x_{p^\beta}$$

and so $x_{p^\beta} = 0$.

Taking $1 \leq i = j \leq c$ in (T3a) we now obtain $z_i = 0$, for all $1 \leq i \leq c$. Hence $(x_i), (y_j), (z_k)$ is the 0 triple and every coherent triple is standard, i.e., λ is split. □

Lemma 11.15. *If $w < v = \gamma$ and (b, c) is pointed then $\lambda = (a, b, c)$ is non-split if and only if $\beta < \text{val}_p(a + p^v + 1)$.*

In that case $E(\lambda)$ is spanned by the standard solutions and the triple $(x_i), (y_j), (z_k)$ with $(x_i) = 0$, $y_{p^v} = 1$, $y_j = 0$ for $j \neq p^v$, $z_{p^v} = -1$ and $z_k = 0$ for $k \neq p^v$.

Proof. Assume that λ is non-split. By Lemma 11.5 we may subtract, from an arbitrary coherent triple, a standard triple to obtain a coherent triple $(x_i), (y_j), (z_k)$ in which $x_i = 0$ for $i \neq p^v, p^\beta$, $y_j = 0$ for $j \neq p^\beta$ and $z_k = 0$ for $k \neq p^v$.

Taking $i = p^v$ and $j = p^w$ in (T3b) we obtain

$$0 = \binom{b + p^w - p^v}{p^w} x_{p^\beta} = (b_w + 1) x_{p^v}$$

and so $x_{p^v} = 0$.

Again by (T3b), with $i = p^\beta$ and $j = p^w$ we have

$$0 = \binom{b + p^w - p^\beta}{p^w} x_{p^\beta} = (b_w + 1) x_{p^\beta},$$

so that $x_{p^\beta} = 0$ and hence $(x_i) = 0$.

Now by (T3a) with $i = j = p^v$ we get $-y_{p^v} = z_{p^v}$.

By Remark 11.4 we may write $a = (p^\beta - 1) - p^v + p^\beta a'$. We show that $a_\beta = p - 1$. If $a_\beta \neq p - 1$ then for $i = p^\beta$ and $k = p^v$ in relation (T1) we get

$$0 = \binom{a + p^v + p^\beta}{p^\beta} z_{p^v} = (a_\beta + 1) z_{p^v}$$

i.e. $z_{p^v} = 0$ and so λ is split. Hence we have that $a_\beta = p - 1$ and so $(a + p^v, b)$ is James, i.e. $\beta < \text{val}_p(a + p^v + 1)$.

It remains to prove that the triple specified in the statement of the Lemma is coherent.

For (T1) we require $\binom{a+i+p^v}{i} = 0$, for $1 \leq i \leq c$, and this is true since $(a + p^v, b)$ is James.

We now consider (T2). If $\binom{a+k}{k} y_j \neq 0$, with $1 \leq j, k \leq c$, $j + k \leq c$, then we must have $j = p^v = p^\gamma$ and so $p^\gamma + k < p^w + p^\gamma$ and $k < p^w < p^v$ so that $\binom{a+k}{k} = 0$. Hence the left hand side of (T2) is zero for all relevant values of j and k . Similarly, if $\binom{b+j}{j} z_k \neq 0$ with $1 \leq j, k \leq c$, $j + k \leq c$, then $k = p^v = p^\gamma$ and so $j < p^w$ and $\binom{b+j}{j} = 0$. Hence the right hand side of (T2) is also zero for all relevant values of j and k .

We now consider (T3a). If $i = j = p^v$ then the equation reads $\binom{a+p^v}{p^v} = -1$, which is true. If $i = p^v$ and $p^v < j \leq c$ then the condition is $\binom{b+j-p^v}{j-p^v} = 0$. Writing $t = j - p^v = j - p^\gamma$ we have $t < p^w$ and so $\binom{b+j-p^v}{j-p^v} = \binom{b+t}{t} = 0$. For $i \neq p^v$ then condition is $\binom{a+i}{i} = 0$, and this is true.

Finally we consider the condition (T3b). The condition is that $\binom{a+i}{i} = 0$ for $p^v < i \leq b + p^v$. But if $\binom{a+i}{i} \neq 0$ then $a_h + i_n \leq p - 1$ for all $h \geq 0$. But then $i_h = 0$ for $h \neq v$ and $i_v = 1$. Hence $i = p^v$. But $p^v < i$ so this is not possible. □

We gather together the main results of this section in the following result.

Proposition 11.16. *Let $\lambda = (a, b, c)$ be a partition with (a, b) pointed, so that b has the form $b = \hat{b} + p^\beta$ with $\hat{b} < p^v < p^\beta$. Then (a, b, c) is non split if and only if it has one of the following forms:*

(i) $\beta > \gamma \geq v = w$ and $\text{val}_p(a + p^v + 1) \geq \beta$. In this case $E(\lambda)$ is spanned by the standard triple and the coherent triple $(x_i), (y_j), (z_k)$ with $x_{p^v} = 1, x_i = 0$ for $i \neq p^v$ and $(y_j) = 0, (z_k) = 0$.

(ii) $\beta = \gamma > v = w$ and $\text{len}_p(b + p^\beta) < \text{val}_p(a + p^v + 1)$. In this case $E(\lambda)$ is spanned by the standard triple and the coherent triple $(x_i), (y_j), (z_k)$ with $x_{p^v} = 1, x_i = 0$ for $i \neq p^v$, with $y_{p^\beta} = -1, y_j = 0$ for $j \neq p^\beta$ and with $(z_k) = 0$.

(iii) $\beta = \gamma > v = w$ and $\text{len}_p(b + p^\beta) < \text{val}_p(a + p^v + p^\beta + 1)$. In this case $E(\lambda)$ is spanned by the standard triple and the coherent triple $(x_i), (y_j), (z_k)$ with $x_{p^v} = 1, x_i = 0$ for $i \neq p^v$, with $y_{p^\beta} = -1, y_j = 0$ for $j \neq p^\beta$ and with $z_{p^\beta} = 1, z_k = 0$ for $k \neq p^\beta$.

(iv) $v \geq w > \gamma$. In this case $E(\lambda)$ is spanned by the standard triple and the coherent triple $(x_i), (y_j), (z_k)$ with $x_{p^\beta} = 1, x_i = 0$ for $i \neq p^\beta$ and $(y_j) = 0, (z_k) = 0$.

(v) $\gamma = v > w$, $\text{val}_p(a + p^v + 1) > \beta$ and $c = \hat{c} + p^w$ with $\hat{c} < p^w$. In this case $E(\lambda)$ is spanned by the standard triple and the coherent triple $(x_i), (y_j), (z_k)$ with $(x_i) = 0$, with $y_{p^v} = -1, y_j = 0$ for $j \neq p^v$ and with $z_{p^v} = -1, z_k = 0$ for $k \neq p^v$.

Proof. Part (i) follows by Lemmas 11.3, 11.12 and 11.13. Part (ii) is Lemma 11.9 (i) and part (iii) is Lemma 11.9 (ii). Now (iv) is Lemma 11.1 and (v) is Lemma 11.15. □

12 Non-split partitions

12.1 A partition is either James or constrained.

Recall that a partition λ is constrained if $\text{Ext}_{B(N)}^1(S^d E, K_\lambda)$, has dimension at most 1 where $d = |\lambda|$, N is at least the number of parts of λ and E is the natural $G(N)$ -module. Equivalently λ is constrained if $E(\lambda)$ is spanned by the standard multi-sequence and at most one other multi-sequence, so if λ is not James, the condition is that $E(\lambda)$ has dimension at most 2.

We have already dealt with James partitions, in Section 7, so the following, which is a consequence of Propositions 10.14 and 11.16, is very reassuring.

Theorem 12.1. *A partition is either James or constrained.*

Proof. We may assume that λ is not James. Let $1 \leq r < n$ be minimal such that $(\lambda_r, \lambda_{r+1})$ is not James. If $r = n - 1$ then the result follows by Proposition 9.5. Hence we may assume that $r < n - 1$ and then by Proposition 9.4(i) we have that the restriction map $E(\lambda) \rightarrow E(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ is injective. Now by Propositions 10.14 and 11.16 we have that the partition $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ is constrained and we are done. \square

Remark 12.2. From now on $\lambda = (\lambda_1, \dots, \lambda_n)$ denotes a partition of length $n \geq 4$ which is not James and non-split. We let $1 \leq r < n$ be minimal such that $(\lambda_r, \lambda_{r+1})$ is not James. Since we have done already the case in which $r = n - 1$, in Proposition 9.5, we also assume that $r < n - 1$. Moreover, by Lemma 9.2 and Proposition 9.4(i) we know that $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ is non-split and that (if $n \geq r + 3$) $(\lambda_{r+3}, \dots, \lambda_n)$ is James. These properties will be our standard assumptions in this section and from now on we will assume that λ has always such a form without further reference. In this section we give a complete description of the non-split, non-James partitions.

We note that our results so far do not give any information for the pair $(\lambda_{r+2}, \lambda_{r+3})$ (for $n \geq r + 3$). This pair may or may not be James. We will investigate these two possibilities separately, starting with the case in which $(\lambda_{r+2}, \lambda_{r+3})$ is not a James partition.

12.2 The case in which $(\lambda_{r+2}, \lambda_{r+3})$ is not James.

We state now the general result for the case in which $(\lambda_{r+2}, \lambda_{r+3})$ is not a James partition. As usual we set $v_i = \text{val}_p(\lambda_i + 1)$, $1 \leq i \leq n - 1$, and $l_i = \text{len}_p(\lambda_i)$, for $2 \leq i \leq n$.

We assume throughout this subsection that $n > r + 2$ and that $(\lambda_{r+2}, \lambda_{r+3})$ is not James.

Proposition 12.3. *Suppose that λ is not James and $r < n - 2$ is minimal such that $(\lambda_r, \lambda_{r+1})$ is not a James partition. Assume further that $(\lambda_{r+2}, \lambda_{r+3})$ is not James and that $(\lambda_{r+3}, \dots, \lambda_n)$ is James. Then the partition λ is non-split if and only if the quadruple $(\lambda_r, \lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$ satisfies the following conditions:*

- (i) $\lambda_r = (p^{l_{r+1}+1} - 1) - p^{v_r} + p^{l_{r+1}+1}\lambda'_r$, for some $\lambda'_r \geq 0$; and
- (ii) $\lambda_{r+1} = (p^{v_r+1} - 1) - p^{v_r} + p^{v_r+1}\lambda'_{r+1}$, $(\lambda_{r+1})_{v_r} \neq 0$ and $\lambda'_{r+1} \geq 0$; and
- (iii) $\lambda_{r+2} = p^{v_r} - 1 + p^{v_r}$; and
- (iv) $\lambda_{r+3} = \hat{\lambda}_{r+3} + p^{v_r}$, with $0 \leq \hat{\lambda}_{r+3} < p^{v_r}$.

If these conditions hold then $E(\lambda)$ is spanned by the standard multi-sequence and a non-zero multi-sequence $(y(t, u)_i)$ satisfying

$$y(r, r + 2)_{p^{v_r}} = y(r + 1, r + 3)_{p^{v_r}} = -y(r + 1, r + 2)_{p^{v_r}} = -y(r, r + 3)_{p^{v_r}}$$

and $y(s, t)_i = 0$ for all the other choices of s, t and i .

Note that since we require in condition (ii) that $\lambda_{r+1} = (p^{v_r+1} - 1) - p^{v_r} + p^{v_r+1}\lambda'_{r+1}$ with $(\lambda_{r+1})_{v_r} \neq 0$, this case does not appear in characteristic 2. In particular, in characteristic 2 the assumption that $(\lambda_{r+2}, \lambda_{r+3})$ is not James forces λ to split.

We give a proof of this proposition in a series of Lemmas. Since $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ is non-split and $(\lambda_r, \lambda_{r+1})$ is not James, the three part partition must have one of the forms described in Proposition 10.14 or Proposition 11.16. Though there are many possibilities the remarkable fact is that only in a single case, namely that described in Proposition 10.14(ii), is it possible for λ to be non-split.

Our method is to work through all these possibilities in turn, but leaving the case described in Proposition 10.14(ii) until last. The consideration of all these split cases occupies Lemmas 12.5 to 12.13. The final exceptional case is dealt with in Lemma 12.14.

We assume until further notice that $n \geq r + 3$. Before we embark on Lemmas 12.5 to 12.14, we prove the following useful result that we will use repeatedly.

Lemma 12.4. *The restriction map $E(\lambda) \rightarrow E(\lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$ is injective, equivalently if $(y(t, u)_i)$ is a coherent multi-sequence in which the extension sequences $(y(r + 1, r + 2)_i)$ and $(y(r + 2, r + 3)_i)$ are zero then $(y(t, u)_i)$ is identically zero. In particular if $(\lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$ is split then λ is split.*

Proof. Let $(y(t, u)_i)$ be a coherent multi-sequence in which $(y(r + 1, r + 2)_i)$ and $(y(r + 2, r + 3)_i)$ are zero. By the commuting relation (C) for the pairs $(r, r + 1)$ and $(r + 2, r + 3)$, and using the facts that $(y(r + 2, r + 3)_j) = 0$ and $(\lambda_{r+2}, \lambda_{r+3})$ is not James we deduce that $(y(r, r + 1)_i) = 0$. Hence we have that $(y(r, r + 1)_i) = 0$ and $(y(r + 1, r + 2)_j) = 0$ and so by Proposition 9.4 (i) λ splits. \square

The Lemmas 12.5 to 12.8 which follow correspond to the cases in which $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ has the form described in Proposition 10.14 (i), (iii), (iv), (v). We therefore adopt as standing assumptions that $(\lambda_r, \lambda_{r+1})$ is split and $l_{r+1} < \text{val}_p(\lambda_r + p^{v_r} + 1)$.

Our strategy is the same in all cases, namely we show that by subtracting a standard multi-sequence from an arbitrary coherent multi-sequence for λ we can obtain a coherent multi-sequence $(y(t, u)_i)$ in which the extension sequences $(y(r + 1, r + 2)_i)$ and $(y(r + 2, r + 3)_j)$ are zero. Thus, by Lemma 12.4, $(y(t, u)_i)$ is identically zero and hence any coherent multi-sequence is standard.

Our detailed analysis begins with the case in which $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ has the form described in Proposition 10.14(i).

Lemma 12.5. *Suppose $l_{r+2} \geq v_r = v_{r+1}$, with $l_{r+1} < \text{val}_p(\lambda_r + p^{v_r} + 1)$ and $\text{val}_p(\lambda_{r+1} - p^{v_r} + 1) > l_{r+2}$. Then λ is split.*

Proof. By Proposition 10.4(i), by subtracting a standard multi-sequence from an arbitrary multi-sequence for λ we can obtain a multi-sequence $(y(t, u)_i)$ in which the sequences $(y(r+1, r+2)_j) = 0$ and $(y(r, r+2)_k) = 0$ and that $y(r, r+1)_i = 0$ for $i \neq p^{v_r}$. It remains to prove that $(y(r+2, r+3)_i) = 0$.

We first show $(y(r+1, r+3)_k) = 0$. The pairs $(r, r+2)$ and $(r+1, r+3)$ are related via the commuting relation (C), and so we have

$$\binom{\lambda_r + k}{k} y(r+1, r+3)_i = \binom{\lambda_{r+1} + i}{i} y(r, r+2)_k.$$

Since $(y(r, r+2)_k) = 0$ we have

$$\binom{\lambda_r + k}{k} y(r+1, r+3)_i = 0$$

for $1 \leq k \leq \lambda_{r+2}$. Now as $l_{r+2} \geq v_r$ we can choose $k = p^{v_r}$ and we have that $\binom{\lambda_r + p^{v_r}}{p^{v_r}} \neq 0$. Therefore, we deduce that $(y(r+1, r+3)_i) = 0$.

We consider $(\lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$. We have the coherent triple $(x_i), (y_j), (z_k)$ with $x_i = y(r+1, r+2)_i, y_j = y(r+2, r+3)_j, z_k = y(r+1, r+3)_k$, for $1 \leq i \leq \lambda_{r+2}, 1 \leq j, k \leq \lambda_{r+3}$

Since $y((r+1, r+3)_k) = 0$ and $(y(r+1, r+2)_j) = 0$ we have $(x_i) = 0$ and $(z_k) = 0$. Since $v_r = v_{r+1}$, by setting $i = p^{v_r}$ in relations (T3a) and (T3b) we deduce that $(y_j) = 0$, i.e. $(y(r+2, r+3)_j) = 0$ and we are done. \square

We now treat the case in which $(\lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$ has the form described in Proposition 10.14(iii).

Lemma 12.6. *Suppose $l_{r+2} > v_r = v_{r+1}$, with $\text{len}_p(\lambda_{r+1} + p^{l_{r+2}}) < \text{val}_p(\lambda_r + p^{v_r} + 1)$, $\text{val}_p(\lambda_{r+1} - p^{v_r} + 1) = l_{r+2}$ and $\lambda_{r+2} = \hat{\lambda}_{r+2} + p^{l_{r+2}}$, with $\hat{\lambda}_{r+2} < p^{v_r}$. Then λ is split.*

Proof. By Proposition 10.4(iii), by subtracting a standard multi-sequence from an arbitrary multi-sequence for λ we can obtain a multi-sequence $(y(t, u)_i)$ in which $(y(r, r+2)_k) = 0, y(r, r+1)_i = 0$ for $i \neq p^{v_r}, y(r+1, r+2)_j = 0$ for $j \neq p^{l_{r+2}}$ and

$$y(r+1, r+2)_{p^{l_{r+2}}} + (\lambda_{r+1})_{l_{r+2}} y(r, r+1)_{p^{v_r}} = 0. \quad (*)$$

Hence, by Lemma 12.4, it suffices to show that $y(r+1, r+2)_{p^{l_{r+2}}} = 0$ and $(y(r+2, r+3)_j) = 0$. Notice though that if $(y(r+2, r+3)_j) = 0$, then by the commuting relation for the pairs $(r, r+1)$ and $(r+2, r+3)$ (and the fact that $(\lambda_{r+2}, \lambda_{r+3})$ is not James) we get that $(y(r, r+1)_i) = 0$, in particular

$y(r, r+1)_{p^{v_r}} = 0$. Then from (*) we deduce directly that $y(r+1, r+2)_{p^{l_{r+2}}} = 0$. Therefore, it is enough to show that $(y(r+2, r+3)_j) = 0$. We show first that $(y(r, r+3)_k) = 0$. The pairs $(r+1, r+2)$ and $(r, r+3)$ are related via the commuting relation (C), and

$$\binom{\lambda_{r+1} + j}{j} y(r, r+3)_k = \binom{\lambda_r + k}{k} y(r+1, r+2)_j.$$

Now since $l_{r+2} > v_r = v_{r+1}$ and $y(r+1, r+2)_{p^{v_r}} = 0$, setting $j = p^{v_r}$ we get directly that $y(r, r+3)_k = 0$.

We consider now $(\lambda_r, \lambda_{r+2}, \lambda_{r+3})$. We have the coherent triple $(x_i), (y_j), (z_k)$ with $x_i = y(r, r+2)_i, y_j = y(r+2, r+3)_j, z_k = y(r, r+3)_k$, for $1 \leq i \leq \lambda_{r+2}, 1 \leq j, k \leq \lambda_{r+3}$. Moreover, we have $(x_i) = 0, (z_k) = 0$. Setting $i = p^{v_r}$ in relations (T3a) and (T3b) we obtain $(y_j) = 0$, i.e., $y(r+2, r+3)_j = 0$ for all $1 \leq j \leq \lambda_{r+3}$, and we are done. \square

We now treat the case in which $(\lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$ has the form described in Proposition 10.14(iv).

Lemma 12.7. *Suppose $l_{r+2} = v_r < v_{r+1}$, with $\text{len}_p(\lambda_{r+1} + p^{l_{r+2}}) < \text{val}_p(\lambda_r + p^{v_r} + 1)$ and $\lambda_{r+2} = \hat{\lambda}_{r+2} + p^{l_{r+2}}$ with $\hat{\lambda}_{r+2} < p^{v_r}$. Then λ is split.*

Proof. By Proposition 10.4(iv), by subtracting a standard multi-sequence from an arbitrary multi-sequence for λ we can obtain a multi-sequence $(y(t, u)_i)$ in which $(y(r, r+1)_i) = 0$ and $y(r+1, r+2)_j = y(r, r+2)_k = 0$ for $j, k \neq p^{v_r}$, with

$$y(r+1, r+2)_{p^{v_r}} + y(r, r+2)_{p^{v_r}} = 0.$$

Since $(y(r, r+1)_i) = 0$ (and $(\lambda_r, \lambda_{r+1})$ is not James) from the commuting relation (C) for the pairs $(r, r+1)$ and $(r+2, r+3)$ we get directly that $(y(r+2, r+3)_j) = 0$. Hence it remains to prove that $(y(r+1, r+2)_i) = 0$. We consider $(\lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$. Since $l_{r+2} < v_{r+1}$, we have that $(\lambda_{r+1}, \lambda_{r+2})$ is a James pair. Since $(\lambda_{r+2}, \lambda_{r+3})$ is not a James partition we get by Lemma 8.1 that in any coherent triple for $(\lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$ we have $(y(r+1, r+2)_i) = 0$, and we are done. \square

We now treat the case in which $(\lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$ has the form described in Proposition 10.14(v).

Lemma 12.8. *Suppose $l_{r+2} = v_r > v_{r+1}$, with $\text{len}_p(\lambda_{r+1} + p^{l_{r+2}}) < \text{val}_p(\lambda_r + p^{v_r} + 1)$ and $\lambda_{r+2} = \hat{\lambda}_{r+2} + p^{l_{r+2}}$, with $\hat{\lambda}_{r+2} < p^{v_{r+1}}$. Then λ is split.*

Proof. By Proposition 10.4(v), by subtracting a standard multi-sequence from an arbitrary multi-sequence for λ we can obtain a multi-sequence $(y(t, u)_i)$ in which $(y(r, r+1)_i) = 0$, and $y(r+1, r+2)_j = y(r, r+2)_k = 0$ for $j, k \neq p^{v_r}$ and

$$y(r+1, r+2)_{p^{v_r}} + y(r, r+2)_{p^{v_r}} = 0.$$

Since $(y(r, r+1)_i) = 0$ (and $(\lambda_r, \lambda_{r+1})$ is not James) the commuting relation (C) for the pairs $(r, r+1)$ and $(r+2, r+3)$ gives directly that $(y(r+2, r+3)_j) = 0$. Hence it remains to prove that $y(r+1, r+2)_{p^{v_r}} = 0$, or equivalently by the equation above, that $y(r, r+2)_{p^{v_r}} = 0$.

We show first that $(y(r, r+3)_k) = 0$. By the commuting relation (C) for the pairs $(r, r+3)$ and $(r+1, r+2)$ we have,

$$\binom{\lambda_{r+1} + j}{j} y(r, r+3)_k = \binom{\lambda_r + k}{k} y(r+1, r+2)_j.$$

Setting $j = p^{v_{r+1}}$ and using the fact that $y(r+1, r+2)_{p^{v_{r+1}}} = 0$, as $v_r > v_{r+1}$, we get that $(y(r, r+3)_k) = 0$.

We consider now $(\lambda_r, \lambda_{r+2}, \lambda_{r+3})$ and the corresponding coherent triple $(y(r, r+2)_i), (y(r+2, r+3)_j), (y(r, r+3)_k)$. Setting $i = p^{v_r}$ and $j = p^{v_{r+2}}$ in relation (T3b) we get

$$0 = \binom{\hat{\lambda}_{r+2} + p^{v_{r+2}}}{p^{v_{r+2}}} y(r, r+2)_{p^{v_r}}$$

where $\lambda_{r+2} = \hat{\lambda}_{r+2} + p^{v_r}$ with $\hat{\lambda}_{r+2} < p^{v_{r+1}}$. Now $v_{r+1} < v_r = l_{r+2}$ and we have $v_{r+2} = \text{val}_p(\hat{\lambda}_{r+2} + 1)$ so that $\binom{\hat{\lambda}_{r+2} + p^{v_{r+2}}}{p^{v_{r+2}}} \neq 0$ and therefore $y(r, r+2)_{p^{v_r}} = 0$. So we are done. \square

The Lemmas 12.9 to 12.13 which follow correspond to the cases in which $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ has the form described in Proposition 11.16 (i)-(v). We therefore adopt as standing assumption that $(\lambda_r, \lambda_{r+1})$ is pointed.

We use the same strategy as in the above cases, namely we show that by subtracting a standard multi-sequence from an arbitrary coherent multi-sequence for λ we can obtain a coherent multi-sequence $(y(t, u)_i)$ in which the extension sequences $(y(r+1, r+2)_i)$ and $(y(r+2, r+3)_j)$ are zero and hence, by Lemma 12.4, $(y(t, u)_i)$ is identically zero and hence any coherent multi-sequence is standard.

The first case is that in which $(\lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$ has the form described in Proposition 11.16(i).

Lemma 12.9. *Suppose that $l_{r+1} > l_{r+2} \geq v_r = v_{r+1}$ with $\text{val}_p(\lambda_r + p^{v_r} + 1) \geq l_{r+1}$ and $\lambda_{r+1} = p^{v_r} - 1 + p^{l_{r+1}}$. Then λ is split.*

Proof. The proof for this case is completely analogous to the proof of Lemma 12.5. \square

In the next case $(\lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$ has the form described in Proposition 11.16(ii).

Lemma 12.10. *Suppose that $l_{r+1} = l_{r+2} > v_r = v_{r+1}$ with $\text{len}_p(\lambda_{r+1} + p^{l_{r+2}}) < \text{val}_p(\lambda_r + p^{v_r} + 1)$ and $\lambda_{r+1} = p^{v_r} - 1 + p^{l_{r+1}}$. Then λ is split.*

Proof. The proof is completely analogous to the proof of Lemma 12.6. \square

The next case, in which $(\lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$ has the form described in Proposition 11.16(iii), requires some additional argument.

Lemma 12.11. *Suppose that $l_{r+1} = l_{r+2} > v_r = v_{r+1}$ with $\text{len}_p(\lambda_{r+1} + p^{l_{r+2}}) < \text{val}_p(\lambda_r + p^{v_r} + p^{l_{r+1}} + 1)$ and $\lambda_{r+1} = p^{v_r} - 1 + p^{l_{r+1}}$. Then λ is split.*

Proof. By Proposition 11.16(iii), by subtracting a standard multi-sequence from an arbitrary multi-sequence for λ we can obtain a multi-sequence $(y(t, u)_i)$ in which $y(r, r+1)_i = 0$ for $i \neq p^{v_r}$, and $y(r+1, r+2)_j = 0$ for $j \neq p^{l_{r+1}}$, and $y(r, r+2)_k = 0$ for $k \neq p^{l_{r+1}}$ and that

$$y(r, r+1)_{p^{v_r}} = -y(r+1, r+2)_{p^{l_{r+1}}} = y(r, r+2)_{p^{l_{r+1}}}.$$

As usual we have to prove that $(y(r+1, r+2)_i) = 0$ and $(y(r+2, r+3)_j) = 0$.

We show first that $(y(r+2, r+3)_j) = 0$. By the commuting relation (C) for the pairs $(r, r+1)$ and $(r+2, r+3)$, we get that

$$\binom{\lambda_r + i}{i} y(r+2, r+3)_k = \binom{\lambda_{r+2} + k}{k} y(r, r+1)_i.$$

Since $(\lambda_r)_{l_{r+1}} = p-2$ and $y(r, r+1)_{p^{l_{r+1}}} = 0$, setting $i = p^{l_{r+1}}$ in the relation above we get that $y(r+2, r+3)_k = 0$.

Now by the same relation and the fact that $(\lambda_{r+2}, \lambda_{r+3})$ is not James we also get that $(y(r, r+1)_i) = 0$. Since now $y(r, r+1)_{p^{v_r}} = -y(r+1, r+2)_{p^{l_{r+1}}}$ and $y(r+1, r+2)_j = 0$ for $j \neq p^{l_{r+1}}$ we get directly that the extension sequence $(y(r+1, r+2)_j)$ is 0 and we are done. \square

We next consider the case in which $(\lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$ has the form described in Proposition 11.16(iv).

Lemma 12.12. *Suppose that $l_{r+2} < v_{r+1} \leq v_r < l_{r+1}$ and $\lambda_{r+1} = \hat{\lambda}_{r+1} + p^{l_{r+1}}$ with $0 \leq \hat{\lambda}_{r+1} < p^{v_r}$. Then λ is split.*

Proof. By Proposition 11.16(iv), by subtracting a standard multi-sequence from an arbitrary multi-sequence for λ we can obtain multi-sequence $(y(t, u)_i)$ in which $(y(r+1, r+2)_i) = 0$ and $(y(r, r+2)_k) = 0$ and that $y(r, r+1)_i = 0$ for $i \neq p^{l_{r+1}}$. Hence it remains to prove that $(y(r+2, r+3)_j) = 0$. This is straightforward. Considering the commuting relation (C) for $(r, r+1)$ and $(r+2, r+3)$ we have

$$\binom{\lambda_r + i}{i} y(r+2, r+3)_j = \binom{\lambda_{r+2} + j}{j} y(r, r+1)_i.$$

Now setting $i = p^{v_r}$, and since $y(r, r+1)_{p^{v_r}} = 0$, we deduce that the extension sequence $(y(r+2, r+3)_j)$ is zero and we are done. \square

In the next case $(\lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$ has the form described in Proposition 11.16(v).

Lemma 12.13. *Suppose $l_{r+1} > l_{r+2} = v_r > v_{r+1}$ with $l_{r+1} < \text{val}_p(\lambda_r + p^{v_r} + 1)$, $\lambda_{r+1} = \hat{\lambda}_{r+1} + p^{l_{r+1}}$ with $0 \leq \hat{\lambda}_{r+1} < p^{v_r}$, and $\lambda_{r+2} = \hat{\lambda}_{r+2} + p^{v_r}$ with $0 \leq \hat{\lambda}_{r+2} < p^{v_{r+1}}$. Then λ is split.*

Proof. The proof for this case is entirely analogous to the proof of Lemma 12.8. \square

Thus we are left with the case in which the non-split triple $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ corresponds to the case of Proposition 10.14(ii). We show here that under certain assumptions this triple provides a non-split partition λ and this finishes the proof of Proposition 12.3. More precisely, we prove the following.

Lemma 12.14. *Suppose that $l_{r+2} = v_r = v_{r+1}$, with $l_{r+1} < \text{val}_p(\lambda_r + p^{v_r} + 1)$, $(\lambda_{r+1})_{v_r} \neq 0$ and $\lambda_{r+2} = \hat{\lambda}_{r+2} + p^{v_r}$ with $0 \leq \hat{\lambda}_{r+2} < p^{v_r}$. Then λ is non-split if and only if:*

- (i) $\lambda_{r+1} = (p^{v_{r+1}} - 1) - p^{v_r} + p^{v_{r+1}} \lambda'_{r+1}$, $(\lambda_{r+1})_{v_r} \neq 0$ and $\lambda'_{r+1} \geq 0$; and
- (ii) $\lambda_{r+2} = (p^{v_r} - 1) + p^{v_r}$; and
- (iii) $\lambda_{r+3} = \hat{\lambda}_{r+3} + p^{v_r}$ with $0 \leq \hat{\lambda}_{r+3} < p^{v_r}$.

In that case $E(\lambda)$ is spanned by the standard multi-sequence and a non-zero multi-sequence $(y(t, u)_i)$ in which

$$y(r, r+2)_{p^{v_r}} = y(r+1, r+3)_{p^{v_r}} = -y(r+1, r+2)_{p^{v_r}} = -y(r, r+3)_{p^{v_r}}.$$

and $y(t, u)_i = 0$ for all other choices of t, u and i .

Proof. We first emphasise that since $v_r = v_{r+1}$ and $\lambda_{r+1} = p^{v_r} - 1 + p^{v_r} \lambda'_{r+1}$ with $(\lambda_{r+1})_{v_r} \neq 0$, the characteristic of K is odd. We show now that if λ_{r+1} or λ_{r+3} does not have the form described in (i) and (iii) then $(\lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$ is split and so by Lemma 12.4, we get that λ is split.

First notice that since $l_{r+2} = v_{r+1}$ we have that $(\lambda_{r+1}, \lambda_{r+2})$ is split. Now by Remark 10.3 (i) and (ii), or directly by Proposition 10.14, we deduce that if λ is non-split then we may write $\lambda_{r+1} = (p^{v_r+1} - 1) - p^{v_r} + p^{v_r+1}\lambda'_{r+1}$ and $l_{r+3} \geq v_r$. Since now $\lambda_{r+2} = \hat{\lambda}_{r+2} + p^{v_r}$ we get that $\lambda_{r+3} = \hat{\lambda}_{r+3} + p^{v_r}$ with $\hat{\lambda}_{r+3} < p^{v_r}$. So we have that if λ is non-split then λ_{r+1} and λ_{r+3} must have the form of (i) and (iii) described in the statement.

We show now that if λ_{r+2} is not as in (ii) of the statement then λ is split. By the description of its parts so far, we have that if $(\lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$ is a non-split triple then it corresponds to one of the cases (ii) or (v) of Proposition 10.14, where $v_{r+2} = v_{r+1}$ or $v_{r+2} < v_{r+1}$ respectively. We show that if $v_{r+2} < v_{r+1}$ then λ is split. Indeed, if $v_{r+2} < v_{r+1}$ then the triple $(\lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$ corresponds to case of Proposition 10.14(v). Thus we can subtract a multiple of the standard multi-sequence to obtain one in which $(y(r+1, r+2)_i) = 0$, and $y(r+1, r+3)_k = y(r+2, r+3)_j = 0$ for $j, k \neq p^{v_r}$, and

$$y(r+2, r+3)_{p^{v_r}} = -y(r+1, r+3)_{p^{v_r}}.$$

Considering the commuting relation (C) for the pairs $(r, r+1)$ and $(r+2, r+3)$ we have

$$\binom{\lambda_{r+2} + j}{j} y(r, r+1)_i = \binom{\lambda_r + i}{i} y(r+2, r+3)_j.$$

Setting $j = p^{v_{r+2}}$ we have $\binom{\lambda_{r+2} + j}{j} \neq 0$ and $y(r+2, r+3)_{p^{v_{r+2}}} = 0$, since $v_{r+2} < v_r$. Therefore, we have $(y(r, r+1)_i) = 0$. We have deduced that $(y(r, r+1)_i) = 0$ and $(y(r+1, r+2)_j) = 0$ and so by Proposition 9.4(i) we get that λ is split.

Hence, from now on we can assume that $(\lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$ is a non-split triple, corresponding to Proposition 10.14(ii), where $v_{r+2} = v_{r+1}$ and since $v_r = v_{r+1}$ we have that $\lambda_{r+2} = (p^{v_r} - 1) + p^{v_r}$, as in the statement.

We prove that in this case, λ is a non-split partition and we have a non-zero coherent multi-sequence of the given form.

We consider first the triple $(\lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$. By Proposition 10.14(ii) we can assume that after subtracting a multiple of the standard multi-sequence we have $(y(r+2, r+3)_j) = 0$, $y(r+1, r+3)_k = y(r+1, r+2)_i = 0$ for $i, k \neq p^{v_r}$, and

$$y(r+1, r+3)_{p^{v_r}} = -y(r+1, r+2)_{p^{v_r}}.$$

We make the following simple observation. Since the pairs $(\lambda_{r+3}, \lambda_t)$ with $t > r+3$ are James partitions we have

$$l_t < v_{r+3} \leq v_r = v_{r+1} = v_{r+2}$$

and so all the pairs (λ_s, λ_t) with $t > r+3$ are James. Moreover, all the pairs (λ_s, λ_t) with $s < r$ are James. Therefore, the only pairs of rows of λ which are not James are those in the quadruple $(\lambda_r, \lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$.

We deal first with the coherent conditions involving $y(r, r+2)_{p^{v_r}}$. For s, t both different from $r, r+2$ we have that the pairs $(r, r+2)$ and (s, t) are related via the commuting relation (C). If we assume further that $(s, t) \neq (r+1, r+3)$ we have that for such a pair (λ_s, λ_t) is James and $(y(s, t)_i) = 0$. This gives directly that both sides in the commuting relation are zero and so this relation is satisfied. Hence, it remains to investigate the relations between the pairs $(\lambda_r, \lambda_{r+2})$ and $(\lambda_{r+1}, \lambda_{r+3})$ and those relations coming from the triples $(\lambda_q, \lambda_r, \lambda_{r+2})$ with $q < r$ and $(\lambda_r, \lambda_{r+2}, \lambda_t)$ with $t > r+2$.

The pairs $(r, r+2)$ and $(r+1, r+3)$ are related via the commuting relation (C) and we have,

$$\binom{\lambda_{r+1} + i}{i} y(r, r+2)_{p^{v_r}} = \binom{\lambda_r + p^{v_r}}{p^{v_r}} y(r+1, r+3)_i = -y(r+1, r+3)_i$$

where the last equality follows from the fact that $(\lambda_r + p^{v_r}, \lambda_{r+1})$ is James. Now notice that since $\lambda_{r+1} = (p^{v_r+1} - 1) - p^{v_r} + p^{v_r+1} \lambda'_{r+1}$ and $l_{r+3} = v_r$, we also have that for $i \leq l_{r+3}$, $\binom{\lambda_{r+1} + i}{i} \neq 0$ if and only if $i = p^{v_r}$, in which case $\binom{\lambda_{r+1} + p^{v_r}}{p^{v_r}} = -1$. Hence, here we recover the relation $y(r, r+2)_{p^{v_r}} = y(r+1, r+3)_{p^{v_r}}$ that we have in the multi-sequence.

We move on to the triples $(\lambda_q, \lambda_r, \lambda_{r+2})$ with $q < r$ and $(\lambda_r, \lambda_{r+2}, \lambda_t)$ with $t > r+2$. For the triple $(\lambda_q, \lambda_r, \lambda_{r+2})$ we have that $y(r, r+2)_j$ is identified with y_j in the relations (T1)-(T3b). Notice that for this triple we have that $x_i^{\text{st}} = 0$, $z_k^{\text{st}} = 0$ and $y_j^{\text{st}} = 0$ for $j \neq p^{v_r}$ with $y_{p^{v_r}}^{\text{st}} = -1$. Therefore, the relations associated with this triple are satisfied by our multi-sequence.

For the triple $(\lambda_r, \lambda_{r+2}, \lambda_t)$ we have that $y(r, r+2)_i$ is identified with x_i in the relations (T1)-(T3b). If $t > r+3$ we have that since (λ_r, λ_t) and $(\lambda_{r+2}, \lambda_t)$ are James, the standard multi-sequence for this triple is $y_j^{\text{st}} = 0$, $z_k^{\text{st}} = 0$ and $x_i^{\text{st}} = 0$ for $i \neq p^{v_r}$ with $x_{p^{v_r}}^{\text{st}} = -1$. So again, the relations associated with this triple are satisfied by our multi-sequence.

If $t = r+3$ then we have the triple $(\lambda_r, \lambda_{r+2}, \lambda_{r+3})$. This triple corresponds to the case (ii) of Proposition 10.14. Here $y(r, r+2)_{p^{v_r}}$ and $y(r, r+3)_{p^{v_r}}$ are both involved in the relations of this triple. Also, $y(r, r+2)_i$ is identified with x_i and $y(r, r+3)_k$ is identified with z_k in the relations (T1)-(T3b). Now by Proposition 10.14(ii) we get that $y(r, r+2)_{p^{v_r}} = -y(r, r+3)_{p^{v_r}}$ is a solution for the relations of this triple and this is exactly what we have in our multi-sequence.

We now deal with $y(r+1, r+3)_{p^{v_r}}$. For s, t both different from $r+1, r+3$ we have that the pairs $(r+1, r+3)$ and (s, t) are related via the commuting relation (C). If we assume further that $(s, t) \neq (r, r+2)$ then for such a pair (λ_s, λ_t) is James and $(y(s, t)_i) = 0$. This gives directly that both sides

in the commuting relation are zero and so this relation is satisfied. Hence, it remains to investigate the relations between the pairs $(\lambda_{r+1}, \lambda_{r+3})$ and $(\lambda_r, \lambda_{r+2})$ and those relations coming from the triples $(\lambda_q, \lambda_{r+1}, \lambda_{r+3})$ with $q \leq r$ and $(\lambda_{r+1}, \lambda_{r+3}, \lambda_t)$ with $t > r + 3$.

For the pairs $(\lambda_{r+1}, \lambda_{r+3})$ and $(\lambda_r, \lambda_{r+2})$ we have already seen that our multi-sequence satisfies the commuting relation associated with these pairs in our previous consideration. Hence, we can move on to the triples

$(\lambda_q, \lambda_{r+1}, \lambda_{r+3})$ with $q \leq r$ and $(\lambda_{r+1}, \lambda_{r+3}, \lambda_t)$ with $t > r + 3$.

For the triple $(\lambda_q, \lambda_{r+1}, \lambda_{r+3})$ we have that $y(r+1, r+3)_j$ is identified with y_j in relations (T1)-(T3b). Notice that if $q < r$, then $x_i^{\text{st}} = 0$, $z_k^{\text{st}} = 0$ and $y_j^{\text{st}} = 0$ for $j \neq p^{v_r}$ with $y_{p^{v_r}}^{\text{st}} = -1$. Therefore, the relations associated with this triple are satisfied by our multi-sequence. If $q = r$ then $(\lambda_r, \lambda_{r+1}, \lambda_{r+3})$ corresponds to the case of Proposition 10.14(ii). By Proposition 10.14(ii) we have that any solution for this triple has the form $x_i = gx_i^{\text{st}} + hx'_i$, for some scalars g, h , with $x'_i = 0$ for $i \neq p^{v_r}$ and $x'_{p^{v_r}} = 1$, $y_j = gy_j^{\text{st}}$ and $z_k = gz_k^{\text{st}} + hz'_k$, with $z'_k = 0$ for $k \neq p^{v_r}$ and $z'_{p^{v_r}} = 2$. Also, notice that $x_i^{\text{st}} = y_j^{\text{st}} = z_k^{\text{st}} = 0$ for $i, j, k \neq p^{v_r}$ and $x_{p^{v_r}}^{\text{st}} = y_{p^{v_r}}^{\text{st}} = z_{p^{v_r}}^{\text{st}} = -1$. Therefore setting $h = g$ we obtain a solution of the form $x_i = 0$, $y_j = z_k = 0$ for $j, k \neq p^{v_r}$ and $y_{p^{v_r}} = -z_{p^{v_r}}$. But this is exactly the solution in our multi-sequence, since here $y_{p^{v_r}}$ is $y(r+1, r+3)_{p^{v_r}}$ and $z_{p^{v_r}}$ is $y(r, r+3)_{p^{v_r}}$. Therefore, we are done.

For the triple $(\lambda_{r+1}, \lambda_{r+3}, \lambda_t)$ now we have that $y(r+1, r+3)_i$ is identified with x_i in the relations (T1)-(T3b). The standard coherent triple satisfies $y_j^{\text{st}} = 0$, $z_k^{\text{st}} = 0$ and $x_i^{\text{st}} = 0$ for $i \neq p^{v_r}$ with $x_{p^{v_r}}^{\text{st}} = -1$. Therefore, the relations associated with this triple are satisfied by our multi-sequence.

The considerations for $y(r+1, r+2)_{p^{v_r}}$, and $y(r, r+3)_{p^{v_r}}$ are identical with those of the last two cases and we leave them to the reader to check. \square

Example 12.15. *One can easily verify that for $p = 3$ the partition $\lambda = (1, 1, 1, 1)$ satisfies the conditions of Lemma 12.14 and that for $n \geq 4$ we have $\text{Ext}_{B(n)}^1(S^4 E, K_\lambda) = H^1(\Sigma_4, \text{Sp}(1, 1, 1, 1)) = K$, where $\text{Sp}(1, 1, 1, 1)$ is the sign representation for Σ_4 .*

12.3 The case in which $(\lambda_{r+2}, \lambda_{r+3})$ is James.

We consider now the case in which the pair $(\lambda_{r+2}, \lambda_{r+3})$ is a James partition. We have the following result.

Proposition 12.16. *Suppose λ is a partition which is not James and $r < n - 1$ minimal such that $(\lambda_r, \lambda_{r+1})$ is not James. Assume further that $(\lambda_{r+2}, \dots, \lambda_n)$ is a James partition. Then λ is non-split if and only if one of the following holds.*

(i) $(\lambda_{r+1}, \lambda_{r+2})$ is not James and $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ is a non-split triple.

(ii) $(\lambda_{r+1}, \lambda_{r+2})$ is James, $(\lambda_r, \lambda_{r+1})$ splits and $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ is a non-split triple, with the extra condition that $l_{r+3} < l_{r+2}$ if $p = 2$ and $n \geq r + 3$.
(iii) $(\lambda_{r+1}, \lambda_{r+2})$ is James, $(\lambda_r, \lambda_{r+1})$ is pointed and there is no $q < r$ with $v_q = \text{len}_p(\lambda_r + p^{l_{r+1}})$.

In all these cases a non-standard multi-sequence $(y(t, u)_i)$ is obtained from a non-standard coherent triple for $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ with $(y(t, u)_i) = 0$ for $(t, u) \neq (r, r + 1), (r + 1, r + 2)$ and $(r, r + 2)$.

We note that for $p \geq 3$ the condition $l_{r+3} < l_{r+2}$ in case (ii) of Proposition 12.16 is automatic, coming from the fact that $(\lambda_{r+2}, \lambda_{r+3})$ is James and the form of the non split triple $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ with $(\lambda_r, \lambda_{r+1})$ split and $(\lambda_{r+1}, \lambda_{r+2})$ James, which is the case (iv) in Proposition 10.14.

We give a proof of Proposition 12.16 in a series of Lemmas corresponding to the forms of the non-split triples of Propositions 10.14 and 11.16. Before we embark on the proof of Lemmas 12.18 to 12.28 we make the following remark that will be used repeatedly.

Remark 12.17. Let λ be a partition which is not James and assume that r is minimal such that $(\lambda_r, \lambda_{r+1})$ not James. Assume further that $(\lambda_{r+2}, \dots, \lambda_n)$ is a James partition. Then for every pair (λ_s, λ_t) with $s \neq r, r + 1$ the partition (λ_s, λ_t) is James.

Indeed, since $(\lambda_1, \dots, \lambda_r)$ and $(\lambda_{r+2}, \dots, \lambda_n)$ are James partitions, we have $l_{q+1} < v_q$ for $q \leq r - 1$ and $q \geq r + 2$. Now, since $s \neq r, r + 1$ we have that $s \leq r - 1$ or $s \geq r + 2$. Also, for $t > s$ we have that $l_t \leq l_{s+1} < v_s$ and so (λ_s, λ_t) is James.

We state and prove now the following ten Lemmas. Recall that λ is a partition which is not James, $1 \leq r < n - 1$ is minimal such that $(\lambda_r, \lambda_{r+1})$ is not James and $(\lambda_{r+2}, \dots, \lambda_n)$ is James.

The Lemmas 12.18 to 12.23 which follow correspond to the cases in which $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ is as described in Proposition 10.14, (i)-(v). We therefore adopt as standing assumptions that $(\lambda_r, \lambda_{r+1})$ is split and $l_{r+1} < \text{val}_p(\lambda_r + p^{v_r} + 1)$.

We first treat the case in which $(\lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$ has the form described in Proposition 10.14(i).

Lemma 12.18. Suppose $l_{r+2} \geq v_r = v_{r+1}$, with $l_{r+1} < \text{val}_p(\lambda_r + p^{v_r} + 1)$ and $\text{val}_p(\lambda_{r+1} - p^{v_r} + 1) > l_{r+2}$. Then λ is non-split and $E(\lambda)$ is spanned by the standard multi-sequence and a multi-sequence $(y(t, u)_i)$ with $y(r, r + 1)_{p^{v_r}} \neq 0$ and $y(t, u)_i = 0$ for all other choices of t, u and i .

Proof. Let $(y(t, u)_i)$ be a multi-sequence as in the statement. It suffices to prove that this is coherent. By Proposition 10.14(i) the sequence satisfies the coherence conditions for the triple $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$.

Consider (s, t) with $s < t$, and s and t different from r and $r + 1$. Then, $(r, r + 1)$ and (s, t) are related via the commuting relation (C). Since $(y(s, t)_j) = 0$ the condition is

$$\binom{\lambda_s + j}{j} y(r, r + 1)_i = 0.$$

However, since $s \neq r, r + 1$ we have that (λ_s, λ_t) is a James partition by Remark 12.17 and so $\binom{\lambda_s + j}{j} = 0$ for $1 \leq j \leq \lambda_t$. Hence, this relation is satisfied.

Therefore we can assume that s or t are equal to r or $r + 1$ and consider the triples $(\lambda_q, \lambda_r, \lambda_{r+1})$ and $(\lambda_r, \lambda_{r+1}, \lambda_t)$.

We consider the triple $(\lambda_q, \lambda_r, \lambda_{r+1})$ first. Here, $y(r, r + 1)_j$ is identified with y_j in the relations (T1)-(T3b). Since (λ_q, λ_r) and $(\lambda_r + p^{v_r}, \lambda_{r+1})$ are James partitions we have that the standard solution for this triple is $x_i^{\text{st}} = z_k^{\text{st}} = 0$ and $y_j^{\text{st}} = 0$ for $j \neq p^{v_r}$ with $y_{p^{v_r}}^{\text{st}} = -1$. Therefore the coherence conditions are satisfied for this triple.

We consider now the triple $(\lambda_r, \lambda_{r+1}, \lambda_t)$. Here $y(r, r + 1)_i$ is identified with x_i in the relations (T1)-(T3b). But as $l_t \leq l_{r+2}$ and any value of $y(r, r + 1)_{p^{v_r}}$ (and all other values 0) is coherent for $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ so we get directly that the coherence conditions are satisfied also for $(\lambda_r, \lambda_{r+1}, \lambda_t)$. \square

We next consider the case in which $(\lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$ has the form described in Proposition 10.14(ii).

Lemma 12.19. *Suppose that $l_{r+2} = v_r = v_{r+1}$, with $l_{r+1} < \text{val}_p(\lambda_r + p^{v_r} + 1)$, $(\lambda_{r+1})_{v_r} \neq 0$ and $\lambda_{r+2} = \hat{\lambda}_{r+2} + p^{v_r}$ with $0 \leq \hat{\lambda}_{r+2} < p^{v_r}$. Then λ is non-split.*

Moreover $E(\lambda)$ is spanned by the standard multi-sequence and a non-zero multi-sequence $(y(t, u)_i)$ such that $y(r, r + 1)_i = y(r, r + 2)_k = 0$ for $i, k \neq p^{v_r}$, with

$$y(r, r + 2)_{p^{v_r}} + (\lambda_{r+1})_{v_r} y(r, r + 1)_{p^{v_r}} = 0$$

and $(y(s, t)_j) = 0$ for $(s, t) \neq (r, r + 1), (r, r + 2)$.

Proof. By Remark 12.17, (λ_s, λ_t) is a James partition for $s \neq r, r + 1$. We show that in this case we also have that for $t \geq r + 3$ the pairs (λ_r, λ_t) and $(\lambda_{r+1}, \lambda_t)$ are James.

First notice that since $(\lambda_{r+1})_{v_r} \neq 0$ and $v_r = v_{r+1}$, the characteristic of K is odd. Now since $(\lambda_{r+2}, \lambda_{r+3})$ is James and $\lambda_{r+2} = \hat{\lambda}_{r+2} + p^{v_r}$ with $\hat{\lambda}_{r+2} < p^{v_r}$, (and since $\text{char}(K) \neq 2$), we have that $\lambda_{r+3} \leq \hat{\lambda}_{r+2} < p^{v_r}$. Therefore, $l_{r+3} < v_r = v_{r+1}$ and so $(\lambda_r, \lambda_{r+3})$ and $(\lambda_{r+1}, \lambda_{r+3})$ are James partitions. Moreover for $t > r + 3$ we have $l_t \leq l_{r+3}$ and so (λ_r, λ_t) and $(\lambda_{r+1}, \lambda_t)$ are also James partitions. Therefore, we deduce that the only pairs which are not James come from $(\lambda_r, \lambda_{r+1})$, $(\lambda_{r+1}, \lambda_{r+2})$ and $(\lambda_r, \lambda_{r+2})$.

Let $(y(t, u)_i)$ be a multi-sequence as in the statement of the Lemma. By Proposition 10.14(ii) the coherence condition is satisfied for $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$.

We deal first with coherence conditions involving $y(r, r+1)_{p^{v_r}}$. For a pair (s, t) , with $s < t$, and both s and t different from r and $r+1$ we have that $(r, r+1)$ and (s, t) are related via the commuting relation (C). Since $(y(s, t)_j) = 0$ and (λ_s, λ_t) is a James partition we have though that this relation becomes a tautology.

Therefore, it remains to consider the triples of the form $(\lambda_q, \lambda_r, \lambda_{r+1})$ with $q < r$ and $(\lambda_r, \lambda_{r+1}, \lambda_t)$ with $t > r+2$.

For the triple $(\lambda_q, \lambda_r, \lambda_{r+1})$ with $q < r$ we have that $y(r, r+1)_j$ is identified with y_j in the relations (T1)-(T3b). Here the standard solution for this triple has the form $x_i^{\text{st}} = z_k^{\text{st}} = 0$ and $y_j^{\text{st}} = 0$ for $j \neq p^{v_r}$ with $y_{p^{v_r}}^{\text{st}} = -1$. Therefore, the relations associated with this triple are satisfied by our multi-sequence.

For the triple $(\lambda_r, \lambda_{r+1}, \lambda_t)$ with $t > r+2$ we have that $y(r, r+1)_i$ is identified with x_i in the relations (T1)-(T3b). Here the standard solution for this triple has the form $x_i^{\text{st}} = 0$ for $i \neq p^{v_r}$, $x_{p^{v_r}}^{\text{st}} = -1$ and $y_j^{\text{st}} = z_k^{\text{st}} = 0$. Therefore, the relations associated with this triple are satisfied by our multi-sequence.

We now deal with coherence conditions involving $y(r, r+2)_{p^{v_r}}$. For a pair (s, t) with $s < t$ and both s and t different from r and $r+2$ we have that $(r, r+2)$ and (s, t) are related via the commuting relation (C). As (λ_s, λ_t) is James and $(y(s, t)_j) = 0$ for these values of (s, t) we have directly that both sides in the commuting relation are 0.

Therefore it remains to consider the triples $(\lambda_q, \lambda_r, \lambda_{r+2})$ with $q < r$ and $(\lambda_r, \lambda_{r+2}, \lambda_t)$ with $t > r+2$.

For the triple $(\lambda_q, \lambda_r, \lambda_{r+2})$ with $q < r$ we have that $y(r, r+2)_j$ is identified with y_j in the relations (T1)-(T3b). Here, the standard solution for this triple is $x_i^{\text{st}} = z_k^{\text{st}} = 0$ and $y_j^{\text{st}} = 0$ for $j \neq p^{v_r}$ with $y_{p^{v_r}}^{\text{st}} = -1$. Therefore, the relations associated with this triple are satisfied by our multi-sequence.

For the triple $(\lambda_r, \lambda_{r+2}, \lambda_t)$ we have that $y(r, r+2)_i$ is identified with x_i in the relations (T1)-(T3b). Notice that since $l_t < v_r$ we have that the standard solution for this triple is $x_i^{\text{st}} = 0$ for $i \neq p^{v_r}$ and $x_{p^{v_r}}^{\text{st}} = -1$ and $y_j^{\text{st}} = z_k^{\text{st}} = 0$. Hence, again the relations associated with this triple are satisfied by our multi-sequence. □

We next consider the case in which $(\lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$ has the form described in Proposition 10.14(iii).

Lemma 12.20. *Suppose $l_{r+2} > v_r = v_{r+1}$, with $\text{len}_p(\lambda_{r+1} + p^{l_{r+2}}) < \text{val}_p(\lambda_r + p^{v_r} + 1)$, $\text{val}_p(\lambda_{r+1} - p^{v_r} + 1) = l_{r+2}$ and $\lambda_{r+2} = \hat{\lambda}_{r+2} + p^{l_{r+2}}$, with $\hat{\lambda}_{r+2} < p^{v_r}$. Then λ is non-split.*

The space $E(\lambda)$ is spanned by the standard multi-sequence and a non-zero multi-sequence $(y(t, u)_i)$ such that $y(r, r+1)_i = 0$ for $i \neq p^{v_r}$, and

$y(r+1, r+2)_j = 0$ for $j \neq p^{l_{r+2}}$ with

$$y(r+1, r+2)_{p^{l_{r+2}}} + (\lambda_{r+1})_{l_{r+2}} y(r, r+1)_{p^{v_r}} = 0$$

and $(y(s, t)_j) = 0$ for $(s, t) \neq (r, r+1), (r+1, r+2)$.

Proof. By Remark 12.17 we have that (λ_s, λ_t) is a James partition for $s \neq r, r+1$. We show that in this case we also have that for $t \geq r+3$, the pairs (λ_r, λ_t) and $(\lambda_{r+1}, \lambda_t)$ are James.

First notice that since $l_{r+2} > v_r$ and $\lambda_{r+2} = \hat{\lambda}_{r+2} + p^{l_{r+2}}$ with $\hat{\lambda}_{r+2} < p^{v_r}$ our assumption that $(\lambda_{r+2}, \lambda_{r+3})$ is James gives directly that $l_{r+3} < v_r = v_r + 1$ and so we have that $(\lambda_r, \lambda_{r+3})$ and $(\lambda_{r+1}, \lambda_{r+3})$ are James partitions. Now, for $t \geq r+3$ we have $l_t \leq l_{r+3}$ and so (λ_r, λ_t) and $(\lambda_{r+1}, \lambda_t)$ are also James partitions. Therefore, we deduce that the only pairs which are not James come from $(\lambda_r, \lambda_{r+1}), (\lambda_{r+1}, \lambda_{r+2})$ and $(\lambda_r, \lambda_{r+2})$.

Let $(y(t, u)_i)$ be a multi-sequence as in the statement of the Lemma. By Proposition 10.14(iii) the coherence condition is satisfied for $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$.

We deal first with coherence conditions involving $y(r, r+1)_{p^{v_r}}$. For a pair (s, t) , with $s < t$ and both s and t different from r and $r+1$ we have $(r, r+1)$ and (s, t) are related via the commuting relation (C). Since $(y(s, t)_j) = 0$ and (λ_s, λ_t) is a James partition we get directly that this relation becomes a tautology.

Therefore, it remains to consider the triples of the form $(\lambda_q, \lambda_r, \lambda_{r+1})$ with $q < r$ and $(\lambda_r, \lambda_{r+1}, \lambda_t)$ with $t > r+2$.

For the triple $(\lambda_q, \lambda_r, \lambda_{r+1})$ with $q < r$ we have that $y(r, r+1)_j$ is identified with y_j in the relations (T1)-(T3b). Here the standard solution for this triple has the form $x_i^{\text{st}} = z_k^{\text{st}} = 0$ and $y_j^{\text{st}} = 0$ for $j \neq p^{v_r}$ with $y_{p^{v_r}}^{\text{st}} = -1$. Therefore the coherence conditions are satisfied for this triple.

For the triple $(\lambda_r, \lambda_{r+1}, \lambda_t)$ with $t > r+2$ we have that $y(r, r+1)_i$ is identified with x_i in the relations (T1)-(T3b). The standard solution for this triple is $x_i^{\text{st}} = 0$ for $i \neq p^{v_r}$ and $x_{p^{v_r}}^{\text{st}} = -1$ and $y_j^{\text{st}} = z_k^{\text{st}} = 0$. So, again the coherence conditions are satisfied for this triple.

We now deal with coherence conditions involving $y(r+1, r+2)_{p^{l_{r+2}}}$. For a pair (s, t) with $s < t$ and s and t both different from $r+1$ and $r+2$ we have that $(r+1, r+2)$ and (s, t) are related via the commuting relation (C). As (λ_s, λ_t) is a James partition and $(y(s, t)_j) = 0$ for such (s, t) , this relation becomes a tautology.

Therefore, it remains to consider the triples $(\lambda_q, \lambda_{r+1}, \lambda_{r+2})$ with $q < r$ and $(\lambda_{r+1}, \lambda_{r+2}, \lambda_t)$ with $t > r+2$.

For the triple $(\lambda_q, \lambda_{r+1}, \lambda_{r+2})$ with $q < r$ we have that $y(r+1, r+2)_j$ is identified with y_j in the relations (T1)-(T3b). Notice now that the partitions $(\lambda_q, \lambda_{r+1}, \lambda_{r+2})$ satisfies the properties of the case of Lemma 8.1 and so we get that the coherence conditions are satisfied for this triple.

For the triple $(\lambda_{r+1}, \lambda_{r+2}, \lambda_t)$ we have that $y(r+1, r+2)_i$ is identified with x_i in the relations (T1)-(T3b). This triple now satisfies the properties

of the case of Proposition 11.16(iv) and we have that $y(r+1, r+2)_{p^{l_{r+2}}} \neq 0$, with all the other values 0, is coherent for this triple. \square

We next consider the case in which $(\lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$ has the form described in Proposition 10.14 (iv).

Lemma 12.21. *Suppose $l_{r+2} = v_r < v_{r+1}$, with $\text{len}_p(\lambda_{r+1} + p^{l_{r+2}}) < \text{val}_p(\lambda_r + p^{v_r} + 1)$ and $\lambda_{r+2} = \hat{\lambda}_{r+2} + p^{l_{r+2}}$ with $\hat{\lambda}_{r+2} < p^{v_r}$. Then*

(i) *For $p \geq 3$ the partition λ is non-split.*

(ii) *For $p = 2$ the partition λ is non-split if and only if $l_{r+3} < l_{r+2}$.*

In the non split case $E(\lambda)$ is spanned by the standard multi-sequence and a non-zero multi-sequence $(y(t, u)_i)$ satisfying $y(r+1, r+2)_j = y(r, r+2)_k = 0$ for $j, k \neq p^{v_r}$ with

$$y(r+1, r+2)_{p^{v_r}} + y(r, r+2)_{p^{v_r}} = 0$$

and $(y(s, t)_j) = 0$ for $(s, t) \neq (r+1, r+2), (r, r+2)$.

Proof. Assume first that $l_{r+3} < l_{r+2}$. Notice that this is always the case if $\text{char}(K) \geq 3$, since $(\lambda_{r+2}, \lambda_{r+3})$ is James and $\lambda_{r+2} = \hat{\lambda}_{r+2} + p^{v_r}$, with $\hat{\lambda}_{r+2} < p^{v_r}$.

By Remark 12.17, (λ_s, λ_t) is a James partition for $s \neq r, r+1$. Now, for $t \geq r+3$ we have $l_t \leq l_{r+3} < v_r < v_{r+1}$, and so we get that (λ_r, λ_t) and $(\lambda_{r+1}, \lambda_t)$ are also James pairs for $t \geq r+3$. Therefore, the only pairs which are not James come from $(\lambda_r, \lambda_{r+1})$ and $(\lambda_r, \lambda_{r+2})$.

Let $(y(t, u)_i)$ be a multi-sequence as in the statement of the Lemma. By Proposition 10.14 (iv) the coherence condition is satisfied for $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$.

We deal first with coherence conditions involving $y(r+1, r+2)_{p^{v_r}}$. For a pair (s, t) , with $s < t$, and both s and t different from $r+1$ and $r+2$ we have $(r+1, r+2)$ and (s, t) are related via the commuting relation (C). Since $(y(s, t)_j) = 0$ and (λ_s, λ_t) is a James partition for such (s, t) we get directly that this relation becomes a tautology.

Therefore, it remains to consider the triples of the form $(\lambda_{r+1}, \lambda_{r+2}, \lambda_t)$ with $t > r+2$ and $(\lambda_q, \lambda_{r+1}, \lambda_{r+2})$ with $q < r$.

For the triple $(\lambda_{r+1}, \lambda_{r+2}, \lambda_t)$ with $t > r+2$ we have that $y(r+1, r+2)_i$ is identified with x_i in the relations (T1)-(T3b). Now, since $l_{r+2} = v_r < v_{r+1}$ and $(\lambda_{r+2}, \lambda_t)$ is a James partitions we get that the partition $(\lambda_{r+1}, \lambda_{r+2}, \lambda_t)$ is James. Moreover, as $l_t \leq l_{r+3} < l_{r+2}$ we have by Lemma 7.7 that the coherence conditions are satisfied for this triple.

For the triple $(\lambda_q, \lambda_{r+1}, \lambda_{r+2})$ with $q < r$ we have that $y(r+1, r+2)_j$ is identified with y_j in the relations (T1)-(T3b). Again we have that this partition is James. Also, since $\text{len}_p(\lambda_{r+1} + p^{l_{r+2}}) < \text{val}_p(\lambda_r + p^{v_r} + 1)$ we have that $\text{len}_p(\lambda_{r+1} + p^{l_{r+2}}) < v_q$. Therefore, if $l_{r+2} = l_{r+3}$ we get that the coherence conditions are satisfied by Lemma 7.7(i) and if $l_{r+2} > l_{r+3}$ the conditions are satisfied by Lemma 7.7(iii).

We deal now with the coherence conditions involving $y(r, r+2)_{p^{v_r}}$. For a pair (s, t) with $s < t$, and both s and t different from r and $r+2$ we have that $(r, r+2)$ and (s, t) are related only via the commuting relation (C). Since $(y(s, t)_j) = 0$ and (λ_s, λ_t) is a James partition for such (s, t) we get directly that this relation becomes a tautology.

Therefore it remains to consider the triples $(\lambda_r, \lambda_{r+2}, \lambda_t)$ with $t > r+2$ and $(\lambda_q, \lambda_r, \lambda_{r+2})$ with $q < r$.

For the triple $(\lambda_r, \lambda_{r+2}, \lambda_t)$ we have that $y(r, r+2)_i$ is identified with x_i in the relations (T1)-(T3b). The standard solution for this triple is $x_i^{\text{st}} = 0$ for $i \neq p^{v_r}$ and $x_{p^{v_r}}^{\text{st}} = -1$ and $y_j^{\text{st}} = z_k^{\text{st}} = 0$. So the coherence conditions are satisfied.

For the triple $(\lambda_q, \lambda_r, \lambda_{r+2})$ with $q < r$ we have that $y(r, r+2)_j$ is identified with y_j in the relations (T1)-(T3b). Now here we have that the standard solution for this triple has the form $x_i^{\text{st}} = z_k^{\text{st}} = 0$ and $y_j^{\text{st}} = 0$ for $j \neq p^{v_r}$ with $y_{p^{v_r}}^{\text{st}} = -1$ and so again we are fine.

We show now that if $l_{r+3} = l_{r+2}$ then λ is split. Notice that since $(\lambda_{r+2}, \lambda_{r+3})$ is James and $\lambda_{r+2} = \hat{\lambda}_{r+2} + p^{l_{r+2}}$, this is valid only in characteristic 2.

We consider an arbitrary coherent multi-sequence for λ . By Proposition 10.14(iv) we have that after subtracting a multiple of the standard solution we can assume that $(y(r, r+1)_i) = 0$, $y(r+1, r+2)_j = 0$ for $j \neq 2^{v_r}$ and $y(r, r+2)_k = 0$ for $k \neq 2^{v_r}$, with $y(r+1, r+2)_{2^{v_r}} + y(r, r+2)_{2^{v_r}} = 0$. By Proposition 9.4(i) it is enough to show that $(y(r, r+1)_i)$ and $(y(r+1, r+2)_j)$ are zero. Hence it remains to prove that $y(r+1, r+2)_{2^{v_r}} = 0$. Now the pairs $(r, r+3)$ and $(r+1, r+2)$ are related via the commuting relation (C) and since $(\lambda_{r+1}, \lambda_{r+2})$ is James we have that

$$\binom{\lambda_r + i}{i} y(r+1, r+2)_j = 0, \text{ with } i \leq \lambda_{r+3}.$$

As $l_{r+3} = l_{r+2} = v_r$ we can set $i = 2^{v_r}$ and we have $\binom{\lambda_r + 2^{v_r}}{2^{v_r}} \neq 0$. Therefore, $y(r+1, r+2)_j = 0$ and we are done. \square

Example 12.22. We give a concrete example to describe the failure of the above result for the situation when $l_{r+3} = l_{r+2} = v_r < v_{r-1}$ in characteristic 2. It is easy to see that the partition $(2, 1, 1, 1)$ satisfies all the assumptions of Lemma 12.20 with $v_1 = l_3 = l_4 = 0$ and $v_2 = 1$. By Proposition 10.14(iv) we have that $\text{Ext}_{B(n)}^1(S^4 E, K_{(2,1,1)}) = K$. However $\text{Ext}_{B(n)}^1(S^5 E, K_{(2,1,1,1)}) = 0$, for $n \geq 5$, since the partition $(2, 1, 1, 1)$ has 2-core $(2, 1)$ and so $\text{Ext}_{B(n)}^1(S^5 E, K_{(2,1,1,1)}) = \text{Ext}_{G(n)}^1(S^5 E, \nabla(2, 1, 1, 1)) = 0$.

We next consider the case in which $(\lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$ has the form described in Proposition 10.14(v).

Lemma 12.23. *Suppose $l_{r+2} = v_r > v_{r+1}$, with $\text{len}_p(\lambda_{r+1} + p^{l_{r+2}}) < \text{val}_p(\lambda_r + p^{v_r} + 1)$ and $\lambda_{r+2} = \hat{\lambda}_{r+2} + p^{l_{r+2}}$, with $\hat{\lambda}_{r+2} < p^{v_{r+1}}$. Then λ is non-split.*

Moreover $E(\lambda)$ is spanned by the standard multi-sequence and a non-zero multi-sequence $(y(t, u)_i)$ satisfying $y(r+1, r+2)_j = y(r, r+2)_k = 0$ for $j, k \neq p^{v_r}$, with

$$y(r+1, r+2)_{p^{v_r}} + y(r, r+2)_{p^{v_r}} = 0$$

and $(y(s, t)_j) = 0$ for $(s, t) \neq (r+1, r+2), (r, r+2)$.

Proof. By Remark 12.17, (λ_s, λ_t) is a James partition for $s \neq r, r+1$. We show that in this case we also have that (λ_r, λ_t) and $(\lambda_{r+1}, \lambda_t)$, with $t \geq r+3$ are James.

Since $(\lambda_{r+2}, \lambda_{r+3})$ is James, $\lambda_{r+2} = \hat{\lambda}_{r+2} + p^{l_{r+2}}$, with $\hat{\lambda}_{r+2} < p^{v_{r+1}}$, and $v_r > v_{r+1}$ we must have that $\lambda_{r+3} < p^{v_{r+1}}$. Hence $l_t \leq l_{r+3} < v_{r+1} < v_r$ and so (λ_r, λ_t) and $(\lambda_{r+1}, \lambda_t)$ are James partitions. Therefore, the only pairs which are not James come from $(\lambda_r, \lambda_{r+1})$, $(\lambda_{r+1}, \lambda_{r+2})$ and $(\lambda_r, \lambda_{r+2})$.

Let $(y(t, u)_i)$ be a multi-sequence as in the statement of the Lemma. By Proposition 10.14(v) the coherence condition is satisfied for $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$.

We first deal with the coherence conditions involving $y(r+1, r+2)_{p^{v_r}}$. For a pair (s, t) , with $s < t$, and both s and t different from $r+1$ and $r+2$ we have that $(r+1, r+2)$ and (s, t) are related via the commuting relation (C). Since $(y(s, t)_j) = 0$ and (λ_s, λ_t) is a James partition for such (s, t) we have that this relation becomes a tautology.

Therefore, it remains to consider the triples of the form $(\lambda_{r+1}, \lambda_{r+2}, \lambda_t)$ with $t > r+2$ and $(\lambda_q, \lambda_{r+1}, \lambda_{r+2})$ with $q < r$.

For a triple $(\lambda_{r+1}, \lambda_{r+2}, \lambda_t)$ with $t > r+2$ we have that $y(r+1, r+2)_i$ is identified with x_i in the relations (T1)-(T3b). Notice that $(\lambda_{r+1}, \lambda_{r+2})$ is a pointed partition and that $l_t < v_{r+1}$. Hence this triple has the form described in Proposition 11.16(iv) and the coherence condition holds for our multi-sequence.

For the triple $(\lambda_q, \lambda_{r+1}, \lambda_{r+2})$ with $q < r$ we have that $y(r+1, r+2)_j$ is identified with y_j in the relations (T1)-(T3b). Now, since $(\lambda_q, \lambda_{r+1})$ is James and $(\lambda_{r+1}, \lambda_{r+2})$ is a pointed partition with $\text{len}_p(\lambda_{r+1} + p^{v_r}) < \text{val}_p(\lambda_r + p^{v_r} + 1) \leq v_s$ we have that this is the case appearing in Lemma 8.1 and this gives directly that the coherence conditions hold.

We now deal with the coherence conditions involving $y(r, r+2)_{p^{v_r}}$.

For a pair (s, t) , with $s < t$, and both s and t different from r and $r+2$ we have that $(r, r+2)$ and (s, t) are related via the commuting relation (C). As (λ_s, λ_t) is a James partition and $(y(s, t)_j) = 0$ for these values of (s, t) we have directly that both sides in the commuting relation are 0.

Therefore it remains to consider the triples $(\lambda_r, \lambda_{r+2}, \lambda_t)$ with $t > r+2$ and $(\lambda_q, \lambda_r, \lambda_{r+2})$ with $q < r$.

For the triple $(\lambda_r, \lambda_{r+2}, \lambda_t)$ we have that $y(r, r+2)_i$ is identified with x_i in the relations (T1)-(T3b). Notice that since $l_t < v_r$ and $(\lambda_r + p^{v_r}, \lambda_{r+1})$ is James, the standard solution for this triple is $x_i^{\text{st}} = 0$ for $i \neq p^{v_r}$ and $x_{p^{v_r}}^{\text{st}} = -1$ and $y_j^{\text{st}} = z_k^{\text{st}} = 0$. So, our multi-sequence satisfies the conditions for this triple.

For the triple $(\lambda_q, \lambda_r, \lambda_{r+2})$ with $q < r$ we have that $y(r, r+2)_j$ is identified with y_j in the relations (T1)-(T3b). Here, the standard solution for this triple is $x_i^{\text{st}} = z_k^{\text{st}} = 0$ and $y_j^{\text{st}} = 0$ for $j \neq p^{v_r}$ with $y_{p^{v_r}}^{\text{st}} = -1$ and so again the coherence conditions for our multi-sequence are satisfied. \square

The remaining five Lemmas correspond to the cases in which $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ has the form described in Proposition 11.16 (i)-(v). We therefore adopt as the standing assumption that $(\lambda_r, \lambda_{r+1})$ is pointed.

We begin with the case in which $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ has the form described in Proposition 11.16(i).

Lemma 12.24. *Suppose $l_{r+1} > l_{r+2} \geq v_r = v_{r+1}$, with $\text{val}_p(\lambda_r + p^{v_r} + 1) \geq l_{r+1}$ and $\lambda_{r+2} = p^{v_r} - 1 + p^{l_{r+1}}$. Then λ is non-split.*

The space $E(\lambda)$ is spanned by the standard multi-sequence and a non-zero multi-sequence $(y(t, u)_i)$ satisfying $y(r, r+1)_i = 0$ for $i \neq p^{v_r}$ and the extension sequence $(y(t, u)_j) = 0$ for $(t, u) \neq (r, r+1)$.

Proof. Let $(y(t, u)_i)$ be a multi-sequence as in the statement of the Lemma. By Proposition 11.16(i), the coherence conditions are satisfied for $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$.

For (s, t) with $s < t$, and both s and t different from r and $r+1$, we have that $(r, r+1)$ and (s, t) are related via the commuting relation (C). Now, since (λ_s, λ_t) is James by Remark 12.17 and $(y(s, t)_j) = 0$ for such (s, t) we have that both sides are 0.

Hence it remains to consider the triples $(\lambda_r, \lambda_{r+1}, \lambda_t)$ with $t > r+2$ and $(\lambda_q, \lambda_r, \lambda_{r+1})$ with $s < r$.

For the triple $(\lambda_r, \lambda_{r+1}, \lambda_t)$, we have that $y(r, r+1)_i$ is identified with x_i in the relations (T1)-(T3b). But as $l_t \leq l_{r+2}$ and any value of $y(r, r+1)_{p^{v_r}}$ (and all other values 0) is coherent for $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ then we get directly that the coherence conditions are satisfied also for $(\lambda_r, \lambda_{r+1}, \lambda_t)$.

For the triple $(\lambda_q, \lambda_r, \lambda_{r+1})$, we have that $y(r, r+1)_i$ is identified with y_j in the relations (T1)-(T3b). Since (λ_q, λ_r) is James and $\text{len}_p(\lambda_r + p^{v_r}) = l_r$ It is a trivial check to see that our solution here satisfies the relations (T1)-(T3b) for this triple. \square

We next consider the case in which $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ has the form described in Proposition 11.16(ii).

Lemma 12.25. *Suppose $l_{r+1} = l_{r+2} > v_r = v_{r+1}$, with $\text{len}_p(\lambda_{r+1} + p^{l_{r+2}}) < \text{val}_p(\lambda_r + p^{v_r} + 1)$ and $\lambda_{r+1} = p^{v_r} - 1 + p^{l_{r+1}}$. Then λ is non-split.*

The space $E(\lambda)$ is spanned by the standard multi-sequence and a non-zero multi-sequence $(y(t, u)_i)$ satisfying $y(r, r+1)_i = 0$ for $i \neq p^{v_r}$ and $y(r+1, r+2)_j = 0$ for $j \neq p^{l_{r+1}}$ with

$$y(r, r+1)_{p^{v_r}} + y(r+1, r+2)_{p^{l_{r+1}}} = 0$$

and $(y(t, u)_j) = 0$ for $(t, u) \neq (r+1, r+2), (r, r+1)$.

Proof. By Remark 12.17 we have that (λ_s, λ_t) is a James partition for $s \neq r, r+1$. We show that in this case we also have that (λ_r, λ_t) and $(\lambda_{r+1}, \lambda_t)$, with $t \geq r+3$, are also James. Notice that since $\lambda_{r+2} = \hat{\lambda}_{r+2} + p^{l_{r+1}}$ with $\hat{\lambda}_{r+2} < p^{v_r}$, $l_{r+1} > v_r$ and $(\lambda_{r+2}, \lambda_{r+3})$ is James we get that $\lambda_{r+3} \leq \hat{\lambda}_{r+2} < p^{v_r}$. Therefore we have that the pairs (λ_r, λ_t) and $(\lambda_{r+1}, \lambda_t)$ with $t \geq r+3$ are also James partitions. Hence, the only pairs of rows of λ which are not James come from $(\lambda_r, \lambda_{r+1})$, $(\lambda_{r+1}, \lambda_{r+2})$, and $(\lambda_r, \lambda_{r+2})$.

Let $(y(t, u)_i)$ be a multi-sequence as in the statement of the Lemma. By Proposition 11.16(ii), the coherence conditions are satisfied for $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$.

We first deal with the coherence conditions involving $y(r, r+1)_{p^{v_r}}$. For (s, t) with $s < t$, and both s and t different from r and $r+1$, we have that $(r, r+1)$ and (s, t) are related via the commuting relation (C). Now, since (λ_s, λ_t) is James and $(y(s, t)_j) = 0$ for such (s, t) we have both sides of the relation are zero.

Hence it remains to consider the triples $(\lambda_r, \lambda_{r+1}, \lambda_t)$ with $t > r+2$ and $(\lambda_q, \lambda_r, \lambda_{r+1})$ with $q < r$.

For the triple $(\lambda_r, \lambda_{r+1}, \lambda_t)$, we have that $y(r, r+1)_i$ is identified with x_i in the relations (T1)-(T3b). Since $\text{len}_p(\lambda_{r+1} + p^{l_{r+2}}) < \text{val}_p(\lambda_r + p^{v_r} + 1)$ and $l_t < v_r$ we have that the standard solution for this triple is $y_j^{\text{st}} = z_k^{\text{st}} = 0$ and $x_i^{\text{st}} = 0$ for $i \neq p^{v_r}$ with $x_{p^{v_r}}^{\text{st}} = -1$. Hence, the coherence conditions for our multi-sequence are satisfied.

For the triple $(\lambda_q, \lambda_r, \lambda_{r+1})$, we have that $y(r, r+1)_i$ is identified with y_j in the relations (T1)-(T3b). Here the standard solution for this triple is $x_i^{\text{st}} = z_k^{\text{st}} = 0$ and $y_j^{\text{st}} = 0$ for $i \neq p^{v_r}$ with $y_{p^{v_r}}^{\text{st}} = -1$. So, the relations associated with this triple are satisfied by our multi-sequence.

We now deal with the coherence conditions involving $y(r+1, r+2)_{p^{l_{r+1}}}$. Again for (s, t) with $s < t$, and both s and t different from $r+1$ and $r+2$, we have that $(r, r+1)$ and (s, t) are related only via the commuting relation (C). Now, since (λ_s, λ_t) is James and $(y(s, t)_j) = 0$ for such (s, t) , both sides in this relation are 0.

Hence it remains to consider the triples $(\lambda_r, \lambda_{r+1}, \lambda_t)$ with $t > r+2$ and $(\lambda_q, \lambda_r, \lambda_{r+1})$ with $q < r$.

For the triple $(\lambda_{r+1}, \lambda_{r+2}, \lambda_t)$, we have that $y(r+1, r+2)_i$ is identified with x_i in the relations (T1)-(T3b). Now $l_t < v_r = v_{r+1}$ and so this triple

is of the form described in Proposition 11.16(iv). Thus we get directly that the coherence conditions for our multi-sequence are satisfied.

For the triple $(\lambda_q, \lambda_{r+1}, \lambda_{r+2})$, we have that $y(r+1, r+2)_j$ is identified with y_j in the relations (T1)-(T3b). Here this triple has the form appearing in Lemma 8.1 and so our multi-sequence satisfies the conditions for triple. \square

We next consider the case in which $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ has the form described in Proposition 11.16(iii).

Lemma 12.26. *Suppose $l_{r+1} = l_{r+2} > v_r = v_{r+1}$, with $\text{len}_p(\lambda_{r+1} + p^{l_{r+2}}) < \text{val}_p(\lambda_r + p^{v_r} + p^{l_{r+1}} + 1)$ and $\lambda_{r+1} = p^{v_r} - 1 + p^{l_{r+1}}$. Then λ is non-split.*

The space $E(\lambda)$ is spanned by the standard multi-sequence and a non-zero multi-sequence $(y(t, u)_i)$ satisfying $y(r, r+1)_i = 0$ for $i \neq p^{v_r}$, $y(r+1, r+2)_j = y(r, r+2)_k = 0$ for $j, k \neq p^{l_{r+1}}$ with

$$y(r, r+1)_{p^{v_r}} = -y(r+1, r+2)_{p^{l_{r+1}}} = y(r, r+2)_{p^{l_{r+1}}}$$

and $y(t, u)_j = 0$ for all other choices of t, u and j .

Proof. By Remark 12.17 we have that (λ_s, λ_t) is a James partition for $s \neq r, r+1$. We show that in this case we also have that (λ_r, λ_t) and $(\lambda_{r+1}, \lambda_t)$, with $t \geq r+3$, are James. We have that since $\lambda_{r+2} = \hat{\lambda}_{r+2} + p^{l_{r+1}}$ with $\hat{\lambda}_{r+2} < p^{v_r}$, $l_{r+1} > v_r$ and $(\lambda_{r+2}, \lambda_{r+3})$ is James then $\lambda_{r+3} \leq \hat{\lambda}_{r+2} < p^{v_r}$. Therefore all the pairs (λ_r, λ_t) and $(\lambda_{r+1}, \lambda_t)$ with $t \geq r+3$ are James. Hence the only pairs of rows of λ which are not James partitions come from $(\lambda_r, \lambda_{r+1})$, $(\lambda_{r+1}, \lambda_{r+2})$, and $(\lambda_r, \lambda_{r+2})$.

Let $(y(t, u)_i)$ be a multi-sequence as in the statement of the Lemma. By Proposition 11.16(iii), the coherence conditions are satisfied for $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$.

We first deal with the coherence conditions involving $y(r, r+1)_{p^{v_r}}$. For (s, t) with $s < t$, and both s and t different from r and $r+1$, we have that $(r, r+1)$ and (λ_s, λ_t) are related via the commuting relation (C). Now, since (λ_s, λ_t) is James and $(y(s, t)_j) = 0$ for such (s, t) , both sides of the relation are 0.

Hence it remains to consider the triples $(\lambda_r, \lambda_{r+1}, \lambda_t)$ with $t > r+2$ and $(\lambda_q, \lambda_r, \lambda_{r+1})$ with $q < r$.

For the triple $(\lambda_r, \lambda_{r+1}, \lambda_t)$, we have that $y(r, r+1)_i$ is identified with x_i in the relations (T1)-(T3b). The standard solution for this triple is $y_j^{\text{st}} = z_k^{\text{st}} = 0$ and $x_i^{\text{st}} = 0$ for $i \neq p^{v_r}, p^{l_{r+1}}$, with $x_{p^{v_r}}^{\text{st}} = x_{p^{l_{r+1}}}^{\text{st}} = -1$. Also $(\lambda_r, \lambda_{r+1}, \lambda_t)$ satisfies the properties of the triple of Proposition 11.16(iv) and so we have that in a coherent triple, $x_i = hx_i^{\text{st}} + gx_i^{\text{pt}}$, with $h, g \in K$. Setting $h = g$ we have that any value of $x_{p^{v_r}} = y(r, r+1)_{p^{v_r}}$ (with all the other values being 0) is a solution for this triple and we are done.

For the triple $(\lambda_q, \lambda_r, \lambda_{r+1})$, we have that $y(r, r+1)_i$ is identified with y_j in the relations (T1)-(T3b). Since (λ_q, λ_r) is James and $\text{len}_p(\lambda_r + p^{v_r}) = l_r$ it is a routine to check that our multi-sequence satisfies the coherence conditions for this triple.

We now deal with the coherence conditions involving $y(r+1, r+2)_{p^{l_{r+1}}}$. Again for (s, t) with $s < t$, and both s and t different from $r+1$ and $r+2$, we have that $(r, r+1)$ and (s, t) are related only via the commuting relation (C). Now, since (λ_s, λ_t) is James and $(y(s, t)_j) = 0$ for such (s, t) , both sides of the relation are zero.

Hence it remains to consider the triples $(\lambda_{r+1}, \lambda_{r+2}, \lambda_t)$ with $t > r+2$ and $(\lambda_q, \lambda_{r+1}, \lambda_{r+2})$ with $q < r$.

For the triple $(\lambda_{r+1}, \lambda_{r+2}, \lambda_t)$, we have that $y(r+1, r+2)_i$ is identified with x_i in the relations (T1)-(T3b). Notice that $l_t < v_r = v_{r+1}$ (since (λ_r, λ_t) is James) and so we are in the case described in Proposition 11.16(iv) and we get directly that any value of $y(r+1, r+2)_{p^{l_{r+1}}}$ (with all the other values being 0) satisfies the coherence relations.

Also the triple $(\lambda_q, \lambda_r, \lambda_{r+2})$ corresponds to the case appearing in Lemma 8.1 and so again we get directly that the coherence conditions are satisfied for this triple.

We leave to the reader to check that the same considerations take care the case of $y(r, r+2)_{p^{l_{r+1}}}$. □

We now consider the case in which $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ has the form described in Proposition 11.16(iv).

Lemma 12.27. *Suppose $l_{r+2} < v_{r+1} \leq v_r$, and $\lambda_{r+1} = \hat{\lambda}_{r+1} + p^{l_{r+1}}$ with $0 \leq \hat{\lambda}_{r+1} < p^{v_r}$. Then λ is non-split if and only if there is no $q < r$ with $v_q = \text{len}_p(\lambda_r + p^{l_{r+1}})$.*

In that case $E(\lambda)$ is spanned by the standard multi-sequence and a non-zero multi-sequence $(y(t, u)_i)$ satisfying $y(r, r+1)_i = 0$ for $i \neq p^{l_{r+1}}$ and $(y(t, u)_j) = 0$ for $(t, u) \neq (r, r+1)$.

Proof. We show first that if there exists a $q < r$ with $v_q = \text{len}_p(\lambda_r + p^{l_{r+1}})$ then λ splits. Assume that such a q exists.

Let $(y(t, u)_i)$ be a coherent multi-sequence. Considering $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ we obtain by Proposition 11.16(iv) that $(y(r+1, r+2)_i) = 0$. Also for such q we have by Lemma 8.1 that $(\lambda_q, \lambda_r, \lambda_{r+1})$ splits. Hence we can subtract a standard multi-sequence to get a coherent multi-sequence (also denoted $(y(t, u)_i)$) in which $(y(r, r+1)_i) = 0$. Now by Proposition 9.4(i) we get directly that λ is split.

We now assume that $v_q > \text{len}_p(\lambda_r + p^{l_{r+1}})$ for every $1 \leq q < r$. Let $(y(t, u)_i)$ be a multi-sequence as in the statement of the Lemma. By Proposition 11.16(iv), the multi-sequence satisfies the coherence conditions for the triple $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$. By our usual arguments for the pairs related via the

commuting relation (C) we obtain an equation in which both sides are zero and the coherence conditions are satisfied.

Hence it remains to consider the triples $(\lambda_q, \lambda_r, \lambda_{r+1})$ with $q < r$ and $(\lambda_{r+1}, \lambda_{r+2}, \lambda_t)$ with $t > r + 2$.

The triple $(\lambda_q, \lambda_r, \lambda_{r+1})$ corresponds to the case appearing in Lemma 8.1 and since $v_q > \text{len}_p(\lambda_r + p^{l_{r+1}})$ for every $q < r$ we have by Lemma 8.1 that the coherence conditions are satisfied.

Also for the triple $(\lambda_r, \lambda_{r+1}, \lambda_t)$, $y(r, r + 1)_i$ is identified with x_i in the relations (T1)-(T3b). But as $l_t \leq l_{r+2}$ and any value of $y(r, r + 1)_{p^{v_r}}$ (and all other values 0) is coherent for $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ then we get directly that the coherence conditions are satisfied also for $(\lambda_r, \lambda_{r+1}, \lambda_t)$. \square

The final case corresponds to the situation in which $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ has the form described in Proposition 11.16(v).

Lemma 12.28. *Suppose $l_{r+2} = v_r > v_{r+1}$ with $l_{r+1} < \text{val}_p(\lambda_r + p^{v_r} + 1)$, $\lambda_{r+1} = \hat{\lambda}_{r+1} + p^{l_{r+1}}$ with $0 \leq \hat{\lambda}_{r+1} < p^{v_r}$, and $\lambda_{r+2} = \hat{\lambda}_{r+2} + p^{v_r}$ with $0 \leq \hat{\lambda}_{r+2} < p^{v_{r+1}}$. Then λ is non-split.*

The space $E(\lambda)$ is spanned by the standard multi-sequence and a non-zero multi-sequence $(y(t, u)_i)$ satisfying $y(r + 1, r + 2)_j = y(r, r + 2)_k = 0$ for $j, k \neq p^{v_r}$ with

$$y(r + 1, r + 2)_{p^{v_r}} + y(r, r + 2)_{p^{v_r}} = 0$$

and $(y(t, u)_j) = 0$ for $(t, u) \neq (r + 1, r + 2), (r, r + 2)$.

Proof. The considerations here are identical with those of Lemma 12.23 and the proof follows in the same way. \square

12.4 Summary

We gather together now our results from Corollary 7.13 and Propositions 9.5, 12.3 and 12.16 in the following theorems to describe all the non-split partitions. We remind to the reader that for a partition $\lambda = (\lambda_1, \dots, \lambda_n)$ we have $v_i = \text{val}_p(\lambda_i + 1)$ and $l_i = \text{len}(\lambda_i)$ for $1 \leq i \leq n$. Also, a partition λ is James if and only if $v_i > l_{i+1}$ for $1 \leq i \leq n - 1$.

Theorem 12.29. *A James partition is non-split if and only if it has length $n \geq 2$. The dimension of the extension group is bounded by $n - 1$, and is described explicitly in Corollary 7.13.*

Theorem 12.30. *Suppose that $p \geq 3$ and $\lambda = (\lambda_1, \dots, \lambda_n)$ is a non-James partition. Let $1 \leq r < n$ be minimal such that $(\lambda_r, \lambda_{r+1})$ is not James. Then λ is non-split if and only if one of the following holds.*

(i) We have $r < n-1$, $(\lambda_{r+1}, \lambda_{r+2})$ is not James, $(\lambda_r, \lambda_{r+1})$ and $(\lambda_{r+1}, \lambda_{r+2})$ are the only non-James consecutive pairs of rows of λ and $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ is non-split.

(iii) We have $r < n-1$, $(\lambda_r, \lambda_{r+1})$ is the only non-James consecutive pair of rows of λ , $(\lambda_r, \lambda_{r+1})$ is split and $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ is non-split.

(iii) We have $r < n$, $(\lambda_r, \lambda_{r+1})$ is the only non-James consecutive pair of rows of λ , $(\lambda_r, \lambda_{r+1})$ is pointed and there is no $1 \leq q < r$ with $v_q = \text{len}_p(\lambda_r + p^{l_{r+1}})$.

(iv) We have $r < n-2$, $(\lambda_{r+3}, \dots, \lambda_n)$ is James and the quadruple

$(\lambda_r, \lambda_{r+1}, \lambda_{r+2}, \lambda_{r+3})$ satisfies the following conditions:

$\lambda_r = (p^{l_{r+1}+1} - 1) - p^{v_r} + p^{l_{r+1}+1} \lambda'_r$, for some $\lambda'_r \geq 0$; and

$\lambda_{r+1} = (p^{v_r+1} - 1) - p^{v_r} + p^{v_r+1} \lambda'_{r+1}$, $(\lambda_{r+1})_{v_r} \neq 0$ and $\lambda'_{r+1} \geq 0$; and

$\lambda_{r+2} = p^{v_r} - 1 + p^{v_r}$; and

$\lambda_{r+3} = \hat{\lambda}_{r+3} + p^{v_r}$, with $0 \leq \hat{\lambda}_{r+3} < p^{v_r}$.

In all cases the dimension of the space of extensions is 1.

Theorem 12.31. Suppose that $p = 2$ and $\lambda = (\lambda_1, \dots, \lambda_n)$ is a non-James partition. Let $1 \leq r < n$ be minimal such that $(\lambda_r, \lambda_{r+1})$ is not James. Then λ is non-split if and only if one of the following holds.

(i) We have $r < n-1$, $(\lambda_{r+1}, \lambda_{r+2})$ is not James, $(\lambda_r, \lambda_{r+1})$ and $(\lambda_{r+1}, \lambda_{r+2})$ are the only non-James consecutive pairs of rows of λ and $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ is non-split.

(ii) We have $r < n-1$, $(\lambda_r, \lambda_{r+1})$ is the only non-James consecutive pair of rows of λ , $(\lambda_r, \lambda_{r+1})$ is a split pair, $(\lambda_r, \lambda_{r+1}, \lambda_{r+2})$ is non-split and if $r < n-2$ then $l_{r+3} < l_{r+2}$.

(iii) We have $r < n$, $(\lambda_r, \lambda_{r+1})$ is the only non-James consecutive pair of rows of λ , $(\lambda_r, \lambda_{r+1})$ is pointed and there is no $q < r$ with $v_q = \text{len}_p(\lambda_r + p^{l_{r+1}})$.

In all cases the dimension of the space of extensions is 1.

13 Further Remarks

In this section we highlight some consequences of our main result and give counterexamples to a conjecture of D. Hemmer in each characteristic and answer a problem that he states in the last section of [12].

Corollary 13.1. Let $p \geq 3$ and $\lambda = (\lambda_1, \dots, \lambda_n)$ a partition of degree d with at least three parts. Then $H^1(\Sigma_{pd}, \text{Sp}(p\lambda)) = 0$.

Proof. This follows immediately by our main results, Theorem 12.29-31. However, we give here an alternative proof using some of our results from the earlier sections. First notice that since $(p\lambda_1, p\lambda_2)$ is not James we get by Proposition 9.4(i) that it is enough to show that $\mu = (p\lambda_1, p\lambda_2, p\lambda_3)$ splits. If $(p\lambda_1, p\lambda_2)$ is split then by Lemma 10.2, μ splits, since $(p\lambda_1 + 1, p\lambda_2)$ is not James.

If $(p\lambda_1, p\lambda_2)$ is pointed then by Lemma 11.3, μ splits, since $\text{val}_p((p\lambda_1 + 1) + 1) = 0$ and $\lambda_2 > 0$. So we are done. \square

Remark 13.2. The above result is not true for $n = 2$. More precisely, by Lemma 5.12 we have that for a partition $\lambda = (\lambda_1, \lambda_2)$ of degree d with $\lambda_2 > 0$, the first cohomology group $H^1(\Sigma_{pd}, \text{Sp}(p\lambda)) \neq 0$ if and only if $p\lambda$ is pointed, i.e. $\lambda_2 = p^t$ for some $t > 0$. In this case we have that the first cohomology group is 1 dimensional. Of course this also follows from the work of Erdmann [9].

Remark 13.3. In [12], Hemmer states that for a partition $\lambda = (\lambda_1, \dots, \lambda_n)$ of degree d there is an isomorphism of vector spaces $H^1(\Sigma_{p^c d}, \text{Sp}(p^c \lambda)) \cong H^1(\Sigma_{p^{c+1} d}, \text{Sp}(p^{c+1} \lambda))$ for $c \geq 1$, [12] Theorem 6.5.8.

If λ has length 2 this follows by our remark above, observing first that for $c \geq 1$ the partition $p^c \lambda$ is pointed if and only if $p\lambda$ is pointed. For $n \geq 3$, Corollary 13.1 gives a much stronger result, namely that

$H^1(\Sigma_{pd}, \text{Sp}(p\lambda)) = 0$ for all the partitions λ with at least three parts.

We note that though Hemmer's statement is true his argument is problematic. More precisely, [12] Corollary 6.4 is not correct and it is heavily used in what follows. This applies also to other parts of the paper (see for example Lemma 6.5.2). The source of these difficulties is his statement that for two partitions λ and μ with length at most n and $\lambda > \mu$ one has that $\lambda_i - \lambda_{i+1} \geq \mu_i - \mu_{i+1}$ for every $1 \leq i \leq n - 1$. This is not true as one may see in simple examples. This statement is used in the argument given for Corollary 6.4. The reader can also check that for $p = 3$ and $G = \text{GL}_5(K)$ the partitions $\lambda = (23, 11, 2)$ and $\mu = (18, 9, 9)$ satisfy the conditions of Corollary 6.4 and that $\lambda - \mu - 2\rho$ is a weight of the Steinberg module St_1 contrary to the statement of Corollary 6.4.

13.4 Counterexamples to Hemmer's Conjecture. In the same paper, the author conjectures, [12], Conjecture 8.2.1, that if $p \geq 3$ and λ is a partition of degree d with $\lambda \neq (d)$ such that the simple module $L(\lambda)$ for $\text{GL}_N(K)$ (with $N \geq d$) appears as a composition factor of $S^d(E)$, then

$$H^1(\Sigma_d, \text{Sp}(\lambda)) \neq 0.$$

Here we give a counterexample to this conjecture in each characteristic for $n \geq 2$. By Krop's description of the composition factors of the symmetric

powers in [18] (see also [8], Remark 4.13) we have that for $p\lambda = p(p-1, \dots, p-1)$, the partition with all n parts equal to $p(p-1)$, the simple module $L(\lambda)$ is a composition factor of $S^d(E)$, where $d = (p^2 - p)(n-1)$.

Now by Corollary 13.1 we have that $H^1(\Sigma_{pd}, \text{Sp}(p\lambda)) = 0$ for $n \geq 3$. Moreover for $n = 2$, since $p \geq 3$, we have $\lambda_2 \neq p^t$ and so by Remark 13.2 we have again that $H^1(\Sigma_{pd}, \text{Sp}(p\lambda)) = 0$. Thus $p\lambda = p(p-1, \dots, p-1)$ is a counterexample to this conjecture for $n \geq 2$ and $p \geq 3$.

13.5 Another Problem. We conclude by answering a question raised in the last section of Hemmer's paper. In [12], Lemma 8.2.3, the author makes the following observation, via James's description of the fixed points of the Specht modules. Given a partition $\lambda = (\lambda_1, \dots, \lambda_n)$ of degree d and $a \geq \lambda_1$ with $\text{len}_p(\lambda_1) < \text{val}_p(a+1)$ (i.e. (a, λ_1) is James) one has that $H^0(\Sigma_d, \text{Sp}(\lambda)) \cong H^0(\Sigma_{a+d}, \text{Sp}(\mu))$, where $\mu = (a, \lambda_1, \dots, \lambda_n)$. It is then asked, [12], Problem 8.2.4, whether one has such an isomorphism for cohomology in higher degree.

Note that in the case of $i = 1$, using Lemma 7.7 and Proposition 7.11 (or directly by Corollary 7.13), one can verify that

$$\dim H^1(\Sigma_d, \text{Sp}(p^n - 1, p^{n-1} - 1, \dots, p^2 - 1, 1)) = n - 1$$

and

$$\dim H^1(\Sigma_{d'}, \text{Sp}(p^{n+1} - 1, p^n - 1, \dots, p^2 - 1, 1)) = n$$

(where d is the degree of $(p^n - 1, p^{n-1} - 1, \dots, p^2 - 1, 1)$ and d' is the degree of $(p^{n+1} - 1, p^n - 1, \dots, p^2 - 1, 1)$).

Appendix I: Comparing Cohomology

We fix integers $n > 0$, $d \geq 0$, with $n \geq d$. We shall use the notation of [7], taking $q = 1$ where appropriate.

We have the Schur algebra $S(n, d)$ and an idempotent $e \in S = S(n, d)$ with eSe identified with the group algebra $K\Sigma_d$, as in [10], and [7]. For a partition λ of d we write $M(\lambda)$ for the corresponding permutation module and $Y(\lambda)$ for the corresponding Young module. The argument comparing extensions for $G(n)$ -modules and $K\Sigma_d$ -modules is based on Section 10 of [5].

For a finite dimensional algebra A , we write $\text{mod}(A)$ for the category of finite dimensional left A -modules.

We have the Schur functor $f : \text{mod}(S) \rightarrow \text{mod}(eSe)$, given on objects by $fV = eV$, $V \in \text{mod}(S)$, as in [10], Chapter 6. Then f is exact and we have the (right exact) functor $g : \text{mod}(eSe) \rightarrow \text{mod}(S)$, given on objects by $gX = Se \otimes_{eSe} X$, as in [5], Chapter 6. We have $\text{Hom}_S(gX, Y) = \text{Hom}_{eSe}(X, fY)$, for $X \in \text{mod}(eSe)$, $Y \in \text{mod}(S)$, by Frobenius reciprocity. We write $L^i g : \text{mod}(eSe) \rightarrow \text{mod}(S)$ for the left derived functors and simply Lg for the first derived functor of g . Now g takes projective modules to projective modules so the factorisation $\text{Hom}_{eSe}(-, fY) = \text{Hom}_S(-, Y) \circ g$ gives rise to a Grothendieck spectral sequence, with second page $\text{Ext}_S^i(L^j gX, Y)$, converging to $\text{Ext}_{eSe}^*(X, fY)$, for $X \in \text{mod}(eSe)$, $Y \in \text{mod}(S)$. In particular we have the 5-term exact sequence

$$\begin{aligned} \text{Ext}_{eSe}^2(X, fY) &\rightarrow \text{Ext}_S^2(gX, Y) \rightarrow \text{Hom}_S(LgX, Y) \\ &\rightarrow \text{Ext}_{eSe}^1(X, fY) \rightarrow \text{Ext}_S^1(gX, Y) \rightarrow 0. \end{aligned}$$

For $\alpha = (\alpha_1, \dots, \alpha_n) \in \Lambda(n, d)$ we write $S^\alpha E$ for $S^{\alpha_1} E \otimes \dots \otimes S^{\alpha_n} E$, the tensor product of symmetric powers of the natural module E . For $\lambda \in \Lambda^+(n, d)$ we write $I(\lambda)$ for the injective hull of the simple module $L(\lambda)$ with highest weight λ , in the polynomial category.

We shall need the following.

Lemma For $\lambda \in \Lambda^+(n, r)$ we have $gY(\lambda) = I(\lambda)$ and $gM(\lambda) = S^\lambda E$.

Proof. By unitriangularity of decompositions of the modules $S^\lambda E$ as a direct sum of $I(\mu)$ s (see e.g. [7], 2.1 (8)) and the Krull-Schmidt Theorem it is enough to show that $gM(\lambda) = S^\lambda E$. For this we argue as follows. For $\alpha \in \Lambda(n, d)$ we have $fS^\alpha E = M(\alpha)$ (see e.g., Section 9, (1)(i) of [5]) and hence

$$\text{Hom}_S(gM(\lambda), S^\alpha E) = \text{Hom}_{eSe}(M(\lambda), fS^\alpha E) = \text{Hom}_{eSe}(M(\lambda), M(\alpha))$$

by Frobenius reciprocity. But this is independent of characteristic (since $M(\lambda)$ and $M(\alpha)$ are permutation modules). Hence, by [7], 2.1 (8) the weight

space dimension $\dim(gM(\lambda))^\alpha$ is independent of characteristic. But in characteristic 0 the functors f and g are inverse equivalences of categories and $fS^\lambda E = M(\lambda)$ so we have $\dim gM(\lambda) = \dim S^\lambda E$ (in all characteristics).

Now for $\lambda = (1^d)$ we have $M(1^d) = eSe$ so that

$$gM(1^d) = Se \otimes_{eSe} eSe = Se = E^{\otimes d}.$$

Now for arbitrary λ we have the natural epimorphism $E^{\otimes d} \rightarrow S^\lambda E$ and so the epimorphism $eSe \rightarrow M(\lambda)$ and so the epimorphism $E^{\otimes d} \rightarrow gM(\lambda)$. Now we have a commutative diagram

$$\begin{array}{ccc} Se \otimes_{eSe} M(1^d) & \rightarrow & E^{\otimes d} \\ \downarrow & & \downarrow \\ Se \otimes_{eSe} M(\lambda) & \rightarrow & S^\lambda E \end{array}$$

where the horizontal maps are induced from the inclusions $M(1^d) \rightarrow E^{\otimes d}$ and $M(\lambda) \rightarrow S^\lambda E$. The top map is an isomorphism and the right map is surjective, hence the bottom map is surjective. Since $\dim Se \otimes_{eSe} M(\lambda) = \dim S^\lambda E$ we have an isomorphism $gM(\lambda) \rightarrow S^\lambda E$. □

For $X = M(\mu)$ and $Y = \nabla(\lambda)$ the 5-term exact sequence takes the form

$$\begin{aligned} \text{Ext}_{\Sigma_d}^2(M(\mu), \text{Sp}(\lambda)) &\rightarrow \text{Ext}_{G(n)}^2(S^\mu E, \nabla(\lambda)) \rightarrow \text{Hom}_{G(n)}(LgM(\mu), \nabla(\lambda)) \\ &\rightarrow \text{Ext}_{\Sigma_d}^1(M(\mu), \text{Sp}(\lambda)) \rightarrow \text{Ext}_{G(n)}^1(S^\mu E, \nabla(\lambda)) \rightarrow 0. \end{aligned}$$

On the other hand if we take $X \in \text{mod}(eSe)$ and Y an injective S -module then the spectral sequence degenerates and we have

$\text{Ext}_{eSe}^i(X, fY) = \text{Hom}_S(L^i gX, Y)$. Hence we have $LgX = 0$ provided that $\text{Ext}_{eSe}^1(X, fY) = 0$, for all injective modules Y . Now an injective S -module is a direct summand of a direct sum of copies of the symmetric powers $S^\alpha E$, $\alpha \in \Lambda^+(n, d)$ and so we have $LgX = 0$ provided that $\text{Ext}_{\Sigma_d}^1(X, M(\tau)) = 0$ for all $\tau \in \Lambda^+(n, d)$. For $X = M(d)$, the trivial module, we have $\text{Ext}_{\Sigma_d}^1(X, M(\tau)) = H^1(\Sigma_d, M(\tau))$, which by Shapiro's Lemma, is $H^1(\Sigma_\tau, K)$, where $\Sigma_\tau = \Sigma_{\tau_1} \times \Sigma_{\tau_2} \times \dots$ (and $\tau = (\tau_1, \tau_2, \dots)$). Now if $p > 2$ we have $H^1(\Sigma_\tau, K) = 0$ and hence $LgM(d) = 0$. Hence, from the above 5-term exact sequence we have $\text{Ext}_{\Sigma_d}^1(M(d), \text{Sp}(\lambda)) = \text{Ext}_{G(n)}^1(S^d E, \nabla(\lambda))$. Of course in all characteristics we have a surjection $\text{Ext}_{\Sigma_d}^1(M(d), \text{Sp}(\lambda)) \rightarrow \text{Ext}_{G(n)}^1(S^d E, \nabla(\lambda))$, from the 5-term exact sequence.

Proposition For $\lambda \in \Lambda^+(n, d)$ we have:

- (i) $\text{Hom}_{G(n)}(S^d E, \nabla(\lambda)) = H^0(\Sigma_d, \text{Sp}(\lambda))$; and
- (ii) $\dim H^1(\Sigma_d, \text{Sp}(\lambda)) \geq \dim \text{Ext}_{G(n)}^1(S^d E, \nabla(\lambda))$, with equality if $p \neq 2$.

The first assertion comes directly from the Lemma and Frobenius reciprocity and the second from the discussion above.

Appendix II: The Relations

For the convenience of the reader, we collect together the relations that are used a great deal throughout the paper.

Definition 13.4. *Let (a, b) be a two part partition. By an extension sequence for (a, b) we mean a sequence $(x_i) = (x_1, \dots, x_b)$ of elements of K such that*

$$\binom{a+i+j}{j} x_i = \binom{i+j}{j} x_{i+j}.$$

Definition 13.5. *We fix a three part partition $\lambda = (a, b, c)$. By a coherent triple of extension sequences for λ we mean a triple $((x_i)_{1 \leq i \leq b}, (y_j)_{1 \leq j \leq c}, (z_k)_{1 \leq k \leq c})$ of extension sequences satisfying the following relations:*

$$(T1) \quad \binom{a+i+k}{k} x_i = \binom{a+i+k}{i} z_k, \quad 1 \leq i \leq b, 1 \leq k \leq c;$$

$$(T2) \quad \binom{a+k}{k} y_j = \binom{b+j}{j} z_k, \quad 1 \leq j, k \leq c, j+k \leq c;$$

$$(T3a) \quad \binom{a+i}{i} y_j = \sum_{s=0}^{i-1} \binom{b+j-i}{j-s} \binom{a+i}{s} x_{i-s} + \binom{b+j-i}{j-i} z_i, \quad 1 \leq i \leq j \leq c;$$

$$(T3b) \quad \binom{a+i}{i} y_j = \sum_{s=0}^j \binom{b+j-i}{j-s} \binom{a+i}{s} x_{i-s}, \quad 1 \leq j \leq c, j < i \leq b+j.$$

In any coherent multi-sequence $(y(t, u)_i)$ for a partition $\lambda = (\lambda_1, \dots, \lambda_n)$ we have the following commuting relations:

$$(C) \quad \binom{\lambda_s + i}{i} y(q, r)_j = \binom{\lambda_q + j}{j} y(s, t)_i, \\ \text{for } 1 \leq q, r, s, t \leq n \text{ distinct with } q < r, s < t.$$

In particular for $1 \leq r < s < n$ with $s > r + 1$ we have

$$(C') \quad \binom{\lambda_s + j}{j} y(r, r+1)_i = \binom{\lambda_r + i}{i} y(s, s+1)_j, \quad 1 \leq i \leq \lambda_{r+1}, 1 \leq j \leq \lambda_{s+1}.$$

References

- [1] Maurice Auslander, Idun Reiten and Sverre O. Smalø, *Representation Theory of Artin Algebras*, *Cambridge Tracts in Mathematics* **36**, CUP 1995
- [2] Edward Cline, Brian Parshall and Leonard Scott, *Cohomology, hyperalgebras and representations*, *Journal of Algebra*, **63**, 98-123, (1980)
- [3] R. Dipper and S. Donkin, *Quantum SL_n* , *Proc. Lond. Math. Soc.* **63**, 165-211, (1991).
- [4] S. Donkin, *Rational Representations of Algebraic Groups: Tensor Products and Filtrations*, *Lecture Notes in Math.* 1140, Springer 1985.
- [5] S. Donkin, *Tilting modules for algebraic groups and finite dimensional algebras*, in: *Handbook of Tilting Theory*, *London Mathematical Society Lecture Note Series* **332**, pp. 215-257. Edited by L. A. Hügel, D. Happel and H. Krause, Cambridge University Press 2007.
- [6] S. Donkin, *Standard Homological Properties for Quantum GL_n* , *J. Algebra*, **181**, 235-266, 1996.
- [7] S. Donkin, *The q -Schur algebra*, *LMS Lecture Notes* 253, Cambridge University Press 1998.
- [8] S. Donkin and H. Geranios, *Composition factors of a tensor product of truncated symmetric powers*, *Advances in Mathematics* **285**, 394-433, 2015.
- [9] K. Erdmann, *Ext^1 for Weyl modules of $SL_2(K)$* , *Math. Z* **218**, 447-459 (1995).
- [10] K. Erdmann, J. A. Green and M. Schocker, *Polynomial Representations of GL_n , Second Edition with an Appendix on Schensted Correspondence and Littelmann Paths*, *Lecture Notes in Mathematics* 830, Springer 2007.
- [11] W. Haboush, *Central differential operators on split semi-simple groups over fields of positive characteristic* In: *Séminaire d'Algèbre P. Dubreil et M.-P. Malliavain, Proc. 1979*," *Lecture Notes in Mathematics* **795**, pp.35-83, Springer 1980, Berlin, Heidelberg, New York.
- [12] D. J Hemmer. *Cohomology and generic cohomology of Specht modules for the symmetric group*, *Journal of Algebra*, **322** (2009), 1498-1515.
- [13] D. J. Hemmer and D. K. Nakano, *Specht filtrations for Hecke algebras of type A* , *J. London Math. Soc.* **69**, (2004), 623-638.

- [14] D. J. Hemmer and D. K. Nakano, *On the cohomology of Specht modules*, Journal of Algebra, **306** (2006), 191-200.
- [15] G. D. James, *The Representation Theory of the Symmetric Groups*, Lecture Notes in Mathematics 682, Springer 1970.
- [16] Jens Carsten Jantzen, *Representations of Algebraic Groups*, second ed., Math. Surveys Monogr., vol 107, Amer. Math. Soc., 2003.
- [17] Alexander S. Kleshchev and Daniel K. Nakano, *On comparing the cohomology of the general linear and symmetric groups*, Pacific Journal of Mathematics, **201**, 339-355, 2001.
- [18] L. Krop, *On the representations of the full matrix semigroup on homogeneous polynomials, I*, J. Algebra, **99**, 284-300, 1986.
- [19] Alison E. Parker, *Higher extensions between modules for SL_2* , Advances in Mathematics, **209**, 381-405, (2007).
- [20] Joseph J. Rotman, *An Introduction to Homological Algebra*, Second Edition, Springer 2008.
- [21] J. B. Sullivan, *Simply connected groups, the hyperalgebra, and Verma's conjecture*, Amer. J. Math. **100**, 1015-1019, (1978).
- [22] W. van der Kallen, *Lectures on Frobenius Splittings and B-modules*, Tata Institute Lecture Notes on Mathematics and Physics, Springer 1993
- [23] Christian Weber, *First degree cohomology of Specht modules over fields of odd prime characteristic*, J. Algebra **392**, 23-41, 2013.