

This is a repository copy of *A role for the OsHKT 2;1 sodium transporter in potassium use efficiency in rice.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/142604/>

Version: Accepted Version

Article:

Hartley, Thomas Noel, Thomas, Alice Sarah and Maathuis, Franciscus Johannes Maria orcid.org/0000-0001-6033-6428 (2019) A role for the OsHKT 2;1 sodium transporter in potassium use efficiency in rice. Jexbot.

<https://doi.org/10.1093/jxb/erz113>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Title: **A role for the OsHKT 2;1 sodium transporter in potassium use efficiency in rice.**

Running Title: Genome-wide association study of potassium use efficiency

Authors: Tom N. Hartley, Alice S Thomas, Frans J. M. Maathuis

Address: Department of Biology, University of York, York, YO1 5DD, UK

Email Addresses: tnh500@york.ac.uk; ast535@york.ac.uk; frans.maathuis@york.ac.uk

Tel number: 44 1904 328652, Orcid: 0000-0001-6033-6428 (F Maathuis, corres. author)

DOS 09/02/2019

2 tables

7 figures (figure 5 in colour in print)

WORD COUNT: 3910

With 5 and 2 supplementary tables and figures, respectively.

Highlight

Genome-wide association studies were used to analyse potassium use efficiency in rice. Novel associations were found along with a role for sodium replacement via the OsHKT2;1 sodium transporter.

Abstract

Increasing the potassium use efficiency (KUE) of crops is important for agricultural sustainability. However, a greater understanding of this complex trait is required to develop new, high KUE cultivars. To this end, a genome-wide association study (GWAS) was applied to diverse rice (*Oryza sativa* L.) genotypes grown under potassium stressed and replete conditions. Using high stringency criteria, the genetic architecture of KUE was uncovered, together with the breadth of physiological responses to low-potassium stress. Specifically, 3 quantitative trait loci (QTLs) were identified, which contained over 90 candidate genes. Of these, the sodium transporter gene *OsHKT2;1* emerged as a key factor that impacts on KUE based on (i) the correlation between shoot Na^+ and KUE, and (ii) higher levels of *HKT2;1* expression in high KUE lines.

Key Words

Fertiliser use, GWAS, *HKT2;1*, potassium, potassium use efficiency, rice, sustainable agriculture, sodium.

Abbreviations

GWAS: genome-wide association study

KUE: potassium use efficiency

QTLs: Quantitative trait loci

SNP: single nucleotide polymorphism

Introduction

K^+ is the most abundant cation in most plants. It is an essential cofactor for many enzymes and has a dominant role in turgor provision and water homeostasis (Maathuis, 2009). The large amounts of K^+ that are required by plants is typically sustained by application of K^+ fertiliser in agronomic contexts. Global demand for potassium fertilisers is currently over 30 million tonnes annually and steadily increasing (FAO, 2017). And though there are ample K^+ reserves, production and application of K^+ fertiliser has important environmental influence: Potash fertilisers contribute to agricultural energy use and greenhouse gas emissions (Brentup and Pallière, 2008; Camargo *et al.*, 2013). In 2016, over 95% of potash was produced in the northern hemisphere (USGS, 2017), exacerbating deleterious environmental consequences through transportation-related emissions. Agriculture is also implicated in adding to atmospheric K^+ deposition (Allen *et al.*, 2010). Taken together, judicious use of potash fertilisers clearly forms an important part of future sustainable agriculture.

At the same time, deficiency for potassium in agricultural soils is widespread and rapidly increasing in areas such as the Australian wheat belt and Chinese rice paddies (Römheld and Kirkby, 2010). Under-fertilisation sometimes results from agricultural malpractice, but is more commonly due to economic considerations, with the cost of K^+ fertiliser purchase and application proving insurmountable. A sustainable solution to mitigate the economic and environmental consequences of growing K^+ demand, while meeting food demand, is to develop crops with higher potassium use efficiency (KUE).

In order to increase crop KUE, knowledge of its genetic underpinnings is important to inform targeted improvement. Studies have been conducted with a range of species and have led to the identification of quantitative trait loci (QTLs) associated with plant responses to potassium deficiency (*e.g.* Wu *et al.*, 1998; Prinzenberg *et al.*, 2010; Kong *et al.*, 2013; Zhao *et al.*, 2014). Similarly, transcriptomics studies (*e.g.* Armengaud *et al.*, 2004; Wang *et al.*, 2012) in low K^+ conditions point to genes that encode membrane proteins involved in transport and other proteins for transcriptional regulation. Genes for such proteins can therefore be seen as putative targets for crop improvements (Shin,

2014; Wang and Wu, 2015), but a more complete understanding of the genetic underpinnings of KUE is still required.

In rice, QTLs for several traits, including potassium uptake and tissue potassium concentration in salt- and non-stressed plants, have been reported (Koyama *et al.*, 2001; Lin *et al.*, 2004; Garcia-Oliveria *et al.*, 2009). Furthermore, QTLs in the context of potassium deficiency have been published (Wu *et al.*, 1998; Miyamoto *et al.*, 2012; Fang *et al.*, 2015), although little overlap in the identified regions was apparent. However, both Miyamoto *et al.* (2012) and Fang *et al.* (2015) described associations in a large (~7 Mb) QTL on chromosome 6 that were linked with shoot sodium, potassium, and calcium concentrations.

The detection of QTLs and genes related to agriculturally important traits in rice has been aided in recent years by genome-wide association studies (GWAS) which typically yield much higher resolution than conventional QTL mapping approaches. Studies have examined abiotic stresses such as aluminium (Famoso *et al.*, 2014) and salt (Kumar *et al.*, 2015; Campbell *et al.*, 2017; Patishtan *et al.*, 2017) and were able to detect novel loci as well as gene candidates. However, the response of rice to potassium deficiency has yet to be examined using GWAS. In this study, the genetic architecture of low potassium stress was explored using the Rice Diversity Panel 1 (Zhao *et al.*, 2011; Eizenga *et al.*, 2014) and in doing so, novel QTLs were detected as well as some which co-localised with those in the prior literature. From this, putative targets for crop improvement were proposed.

Materials and Methods

Plant Growth and Germplasm

Five seeds from each of 324 rice (*Oryza sativa*) cultivars (see Supplementary Table 1 for a full list of accessions) were germinated in sand flooded with distilled water for two weeks prior to transfer to hydroponic treatments. Seedlings were placed in 9 L boxes which contained a nutrient solution adapted from Yoshida *et al.* (1976) which consisted of: (in mM) 1.4 NH_4NO_3 , 0.3 NaH_2PO_4 , 1 CaCl_2 , 1.6 $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, and 0.2 $\text{Na}_2\text{O}_3\text{Si}$ and (in μM) 9.5 MnCl_2 , 0.07 $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24}$, 18 H_3BO_3 , 0.15 ZnSO_4 , 0.16 CuSO_4 , 71 citric

acid monohydrate. Potassium was added as KCl to a final concentration of 0.1 (low K⁺ or LK treatment) or 1 mM (high K⁺ or HK treatment). Nutrient solutions were changed weekly. One seedling from each cultivar was placed in each treatment and growth trials were replicated five times. Plants were grown in a glasshouse for four weeks (or as indicated in the text) with 12 hour day and night periods with temperatures of 32 and 28 °C in the day and night respectively. The relative humidity was maintained between 50 and 60%. For detailed growth experiments on IR64, plants were grown as described above in the presence of 0.01, 0.1, 0.5, 1 or 5 mM K⁺ (added as KCl) and a total amount of 3 mM Na⁺.

Tissue Cation Analysis

Sampled plants were separated into roots and shoots, and their fresh weights were recorded before being oven dried at 80 °C for three days. Tissues were then re-weighed before potassium and sodium concentrations were determined after extraction in 20 mM CaCl₂ for 24 hours. Cation concentrations were measured using a flame photometer (Sherwood Scientific, Cambridge, Cambridgeshire, UK).

Trait Measurement

Briefly, each rice genotype was grown in potassium deficient (0.1 mM) and replete (1 mM) nutrient solutions (see above). Relative growth rate (RGR) was calculated as $[\ln(FW_{\text{end}}) - \ln(FW_{\text{start}})] / (t_{\text{end}} - t_{\text{start}})$, where FW is the whole plant fresh weight. Potassium and sodium tissue concentrations were measured as described above. Phenotype data were based on five biological replicates and least squares means were calculated from raw data. Cultivars with fewer than three replicates were excluded from the analysis. Two different KUE metrics were used: KUE-RGR (defined as the percentage reduction in RGR between LK and HK conditions) and KUE-K (defined as RGR at LK treatment divided by shoot K⁺ concentration at LK treatment). The latter trait examines the K⁺ utilisation, while KUE-RGR can be influenced by both the uptake and utilisation of K⁺.

Genome-wide Association Studies

GWAS was carried out using R 3.3.3 and the *GenABEL* R package (Aulchenko *et al.*, 2007) for KUE metrics, RGR, and tissue cation concentrations. SNPs with a minor allele

frequency < 0.05 and a call rate < 0.9 were excluded from analyses to minimise the risk of spurious associations. Mixed linear models were used for analyses to control for the population structure present in rice (Zhao *et al.*, 2011) which can also induce spurious associations between traits and genetic loci. The top three principal components for population structure were included as fixed effects if this resulted in a model with a genomic inflation factor (Devlin and Roeder, 1999) nearer unity. Previous work has found that the use of mixed models with principal components as covariates to be successful in limiting the occurrence of false signals (Zhao *et al.*, 2011; Kumar *et al.*, 2015; Patish-tan *et al.*, 2017). Associations between SNPs and genotypes were declared significant if their P-value was $< 1 \times 10^{-5}$ (Crowell *et al.*, 2016) and the false discovery rate (Benja-mini and Hochberg, 1995) was less than 10%.

Identification of Quantitative Trait Loci and Candidate Genes

A minimum of two significant associations within a 200 kbp window was required for a significant association to be considered as a QTL to minimise the risk of false positives. This genomic region window size was chosen because linkage disequilibrium in rice de-clines rapidly over this distance (Zhao *et al.*, 2011; McCouch *et al.*, 2016) and genes that are proximal to associations can be considered more credible candidates for influ-encing the trait in question. QTLs which overlapped were grouped into a single QTL. Genes within QTLs were sourced from found using the the Rice Genome Annotation Project website (http://rice.plantbiology.msu.edu/pub/data/Eukaryot-ic_Projects/o_sativa/annotation_dbs/pseudomolecules/version_7.0/). Candidate genes were found among these genes, with those with products relating to transport, signalling, and transcription considered to be more credible candidates. Co-localisation of signifi-cantly associated SNPs and genes within QTLs was examined using the Rice Diversity Allele Finder (<http://rs-bt-mccouch4.biotech.cornell.edu/AF/>). Such co-localisation with a gene could indicate relevance to the trait and non-synonymous SNPs could lead to changes that ultimately alter KUE.

HKT2;1 Expression Analysis

Seeds for the following cultivars germinated (where 'L' indicates low KUE and 'H' indicates high KUE): Cybonnet (L), Dom Sufid (L), Edith (L), Padi Kasalle (L), Tox 782-20-1 (L), 116 (H), Sathi (H), Saturn (H), Ghati Kamma Nangarhar (H), Wanica (H). Plants were grown as described above on an adapted Yoshida nutrient solution containing (in mM) 2.9 NH₄NO₃, 0.3 H₃PO₄, 0.01 KCl, 1 CaCl₂·2H₂O and 1.6 MgSO₄ (micronutrients as described above). Medium was adjusted to pH 5.6 using methyl glucamine and supplemented with either 0 mM NaCl or 1 mM NaCl. Plants were grown for four weeks after which roots from the three plants of each cultivar were pooled and frozen in liquid nitrogen. The root samples were ground to a powder in liquid nitrogen and total RNA was extracted using a Nucleospin RNA Plant and Fungi kit (Macherey-Nagel Bioanalysis). cDNA was synthesised using a Superscript II reverse transcriptase kit (Invitrogen) with oligo dT primers. Quantitative polymerase chain reactions (qPCR) were performed using the QuantStudio 3 (Thermo Fisher) system and Fast SYBR green master mix (Thermo Fisher) using 5'CTCCATCGACTGCTCACTCA3' and 5'GGACAGTGCAAATGTTGTGCG3' as forward and reverse HKT2;1 specific primers. The expression of Elongation Factor 1 alpha was used as an internal control with 5'CACATTGCCGTCAAGTTTGC3' and 5'CCATACCAGCATCACCGTTC3' forward and reverse primers respectively. Data are presented as the average of three biological replications.

Results and Discussion

Influence of Potassium Stress on Growth and Tissue Cation Concentrations

Lowering the medium K⁺ concentration from 1 (HK) to 0.1 (LK) mM had a substantial effect on growth and tissue cation levels. Fig. 1a shows that the mean final mass of LK plants was approximately 40% of that achieved by HK plants. However, at the tissue level, plant growth was not affected uniformly. For example, root to shoot mass ratio was significantly higher in the LK treatment compared to the HK treatment (data not shown). Furthermore, Fig. 1b shows that rice cultivars vary greatly in their growth response to LK. The RGR reduction ranged from 30% to -5% when comparing LK and HK growth data. In other words, the relative growth rates of some lines declines by nearly a

third between LK and HK conditions, while others were not at all or only little affected, irrespective of a 10-fold change in medium K^+ concentration.

As expected, both root and shoot K^+ concentrations were lower in the LK treatment. Across the cultivars, the average shoot potassium concentration declined from 686 to 154 $\mu\text{mol gDW}^{-1}$ between the HK and LK conditions, while the root concentrations declined from 198 to 59 $\mu\text{mol gDW}^{-1}$ (Fig. 2). Shoot potassium concentrations were consistently greater than those of roots. In combination, the growth and tissue K^+ data show that the LK conditions were effective in causing stress which reduced rice growth, likely arising from insufficient tissue K^+ levels. Indeed, many previous studies have shown a strong link between tissue K^+ and growth across several plant species (*e.g.* Asher and Ozanne, 1967; Fageria, 1976; Spear *et al.*, 1978).

While low tissue K^+ is strongly linked with reduced RGR between treatments, the association is less clear within a specific treatment: In both LK and HK treatments only weak non-significant correlations were derived between tissue K^+ and growth. Such seemingly contradictory outcomes can be explained by the existence of considerable (genetic) variation in the sensitivity of cultivars when exposed to declining levels of tissue K^+ .

Table 1 shows growth and tissue cation data for the ten highest and lowest ranking rice cultivars for KUE. KUE-RGR is a measure for the relative growth reduction when changing from HK to LK conditions (RGR_LK/RGR_HK) and differed significantly between cultivars (one-way ANOVA, $P < 0.01$). KUE-K denotes the utilisation of K^+ (amount of growth per unit K^+ ; $\text{RGR_LK/shoot } K\text{_LK}$) and this too, varied significantly between cultivars (one-way ANOVA, $P < 0.001$) with a 5-6 fold difference between the lowest and highest values (Suppl. Table 4). Interestingly, KUE_RGR and LK shoot $[\text{Na}^+]$ showed a highly significant negative correlation ($r = -0.385$, $P < 0.001$; Figure 3) and similar, but weaker, negative correlations were found between KUE-RGR and HK shoot $[\text{Na}^+]$, LK root $[\text{Na}^+]$ and HK root $[\text{Na}^+]$ respectively (Suppl. Fig. 1). Such evidence points to a potential beneficial effect of Na^+ in rice shoots when potassium is limiting, and this may be the result of replacement of K^+ by Na^+ . However, in contrast to KUE-RGR, KUE-K did not correlate significantly with either root or shoot levels of Na^+ . Indeed, very little overlap between the KUE-K and KUE-RGR was apparent with only two cultivars (GSOR 117

and 142) emerging as high KUE lines irrespective of the KUE definition (see Suppl. Table 4). The lack of similarity between KUE-RGR and KUE-K emphasises the different phenomena these metrics describe: while KUE-K is determined by high growth rates and low shoot $[K^+]$ (e.g. ~ 90 mM and ~ 250 mM in high and low KUE-K lines respectively, see Table 1), KUE-RGR expresses how well growth is maintained by cultivars in the face of a shortage of K^+ . Though both approaches are valuable in an agronomic context one may be more suitable for optimising local requirements such as soil nutrient status or availability of K fertiliser. The wide variability in either parameter suggests there is a large scope to enhance these traits.

Genome-wide Association Studies of Potassium Stress

In order to better understand which mechanisms contribute to KUE, GWAS was applied to the growth, cation, and KUE data (Supplementary Table 2). Based on the stringency criteria outlined in the Methods section, a total of four association signals was detected; one each for KUE-K (defined as RGR/shoot K), RGR at LK treatment, shoot $[Na^+]$ and root $[Na^+]$ at LK treatment (Fig. 4; Table 2). Furthermore, the two sodium-related signals co-localised at a position approximately 29.5 Mbp along chromosome 6 and had the same significantly associated SNPs.

The three independent QTLs subsumed a total of 86 unique genes (Suppl. Table 5) and 8 significantly associated SNPs (Table 2). Interrogation of the Rice Diversity Allele Finder (<http://rs-bt-mccouch4.biotech.cornell.edu/AF/>) showed that the two SNPs belonging to the RGR_LK association were synonymous and were located in the coding region of a putative retrotransposon protein (LOC_Os01g39640). One of the KUE-K associations was a synonymous SNP in the intron of another putative retrotransposon protein (LOC_Os01g59580), and both SNPs the Na^+ -related signal were synonymous and located in the coding region of the gene for OsHKT2;1 (LOC_Os06g48810), a sodium transporter.

QTLs repeatedly found across different studies can help to identify robust candidates for crop improvement. The positions of QTLs identified in this study were therefore compared against those previously reported (Figure 5). Though it is noted that many previ-

ous studies had relatively low resolution, leading to QTLs that span many Mbp (e.g. Fang et al., 2015), an overlap was found for the chromosome 1 RGR-K signal which is positioned at the beginning of a ~10 Mbp QTL described by Fang et al. (2015). The tissue Na⁺ associated signals on chromosome 6 found in this study were previously described by Miyamoto *et al.* (2012) who identified a 6.4 Mbp region on chromosome 6 related to sodium uptake and, using a map based cloning strategy, isolated a 100 kb chromosomal region that contained HKT2;1.

Putative Drivers of KUE

Out of the 86 genes covered by the significant association signals, the 42 annotated genes were further evaluated to identify potential drivers of KUE. Gene ontology analysis is problematic with a sample of this size and it is therefore not surprising that no enriched functional class was discovered. In addition to HKT2;1, three further genes (OsCML1 - Calmodulin-related calcium sensor protein; OsSub52 - Putative Subtilisin homologue; OsHKT2;4 - Na⁺ transporter) were previously shown to respond transcriptionally to low K⁺ conditions (Shankar et al., 2013) suggesting they may play a role in K⁺ homeostasis. Furthermore, on the basis of functional annotations the list contains a large proportion (>10%) of genes that are involved in 'disease resistance' (n=7) and in 'RNA translation' (n=5), pointing to a potential role of these processes in establishing KUE. There is a well documented link between K⁺ deficiency and disease (e.g. Davis et al., 2018); Rice diseases like brown leaf spot, scab and stem rot are generally not problematic in K⁺ replete fields but can easily overwhelm K⁺-deficient rice. It is not directly obvious how disease impacts on KUE but LK treatment could (transcriptionally or otherwise) prime plants and thus make them more disease resilient. Improved resilience could alter KUE via generic growth effects. Ribosomal functioning is frequently mentioned as an example process that requires high levels (>100 mM) of K⁺ (e.g. Maathuis, 2009). Similar to disease resistance, the link between RNA translation and KUE may be convoluted but more efficient ribosomal constituents and enzymes involved in translation could improve growth and/or allow plants to adequately synthesise proteins at lower cytoplasmic K⁺ levels. In contrast to the above, the connection between Na⁺ and K⁺ (and hence between Na⁺ and KUE) is well established (e.g. Maathuis and Amtmann,

1999). Thus the appearance of two putative Na⁺ transporters, in combination with significant signals in the root Na⁺ and shoot Na⁺ traits, strongly suggest that Na⁺ transport is an important contributing factor in KUE.

HKT2;1 Plays a Role in KUE via Shoot Sodium

The cation transport category contains two 'high affinity K transporters'. HKT2;1 and HKT2;4 are part of significant association signals when either root or shoot Na⁺ concentration was used as trait. HKT2;4 (Os06g48800) is located in the plasma membrane and expressed in the peripheral layers of rice roots and in the shoot vasculature (Sassie et al., 2012). Members of subgroup II HKTs typically perform K:Na cotransport but in heterologous systems HKT2;4 was shown to move K⁺ without the need for Na⁺ (Horie et al., 2011). Thus, HKT2;4 could be involved in K⁺ (re)distribution, for example between root and shoot. However, its loss of function did not generate a K⁺-dependent phenotype, though this could be due to functional redundancy with, for example, the very similar HKT2;3 (Horie et al., 2011).

In contrast to HKT2;4, HKT2;1 strongly discriminates against K⁺ and, in a physiological context, is believed to exclusively function as a Na⁺ transporter (Horie et al., 2007; Miyamoto et al., 2012). This would fit in with the observation that HKT2;1 is associated with tissue Na⁺ phenotypes (Suppl Table 5). Earlier work by Horie et al, (2007) showed that HKT2;1 is mostly expressed in rice roots and that expression is induced during low K⁺ conditions. Furthermore, HKT2;1 was previously identified in a QTL associated with high Na⁺ accumulation in K-deficient rice plants (Miyamoto et al., 2012). Thus, HKT2;1 has been identified in multiple QTL studies and is transcriptionally regulated in a K⁺ dependent manner. It therefore forms a high confidence candidate that impacts on KUE via the replacement of non essential K⁺ by the physico-chemically similar monovalent Na⁺.

Na⁺ behaves as a beneficial nutrient for K⁺-starved glycophytes when present at moderate concentration (e.g. Maathuis, 2013). Substitution of K⁺ by Na⁺ in such conditions could make a valuable contribution to maintaining non-critical functions of K⁺, such as turgor generation, and thus contribute to KUE. Detailed growth experiments with one of

the cultivars (IR64) show that there is a clear negative correlation between external K^+ levels and tissue Na^+ , for both roots and shoots (Fig. 6). In addition, our physiological data suggest that raised root and shoot Na^+ has a positive effect on KUE: Fig. 3 shows that both root and shoot levels of Na^+ negatively correlate with KUE-RGR but that this is clearly more significant for shoot Na^+ in the LK treatment. This phenomenon also becomes clear when overall tissue cation composition is compared between high and low KUE lines (Table 1). In HK conditions, shoot K^+ ($\sim 650 \text{ } \mu\text{mol gDW}^{-1}$) and shoot Na^+ ($\sim 50 \text{ } \mu\text{mol gDW}^{-1}$) generate a K:Na ratio of around 10-18, and is similar for high and low KUE accessions (Table 1), using either KUE definition. But LK treatment causes a dramatic change in the K:Na ratio to less than one of around 0.7 and 0.3 in low and high KUE lines respectively, reflecting the greater capacity of high KUE cultivars to exploit Na^+ as a K^+ replacement.

Since there is a clear positive impact of Na^+ on KUE-RGR it is imperative to identify the molecular mechanisms involved. Our GWAS studies identified HKT2;1 as a potential causative agent for Na^+ dependent variation in KEU. There is considerable allelic variation in the HKT2;1 coding sequence which contains 5 non-synonymous SNPs that are located in the cytoplasmic N terminal and at the end of the 1st and 6th transmembrane spans (Oomen et al., 2012). Extensive measurements on oocytes that heterologously express HKT2;1 showed that neither of the amino acid substitutions has a significant effect on HKT2;1 functional properties (Oomen et al., 2012). However, the HKT2;1 promoter region contains a large number (>50) of polymorphisms (e.g <http://snp-seek.irri.org/>), many of which are located in transcription factor binding domains (e.g. PlantPan2; <http://plantpan2.itps.ncku.edu.tw/>) and consequently could affect expression levels. We therefore tested whether HKT2;1 expression levels differed between five high and five low KUE lines grown on 0.01 mM K^+ and with or without 1 mM Na^+ . Figure 7 shows that in these very low K^+ grown plants, the average expression level of HKT2;1 in both low and high KUE lines is induced in the presence of Na^+ (1 mM) as was reported previously (Horie et al., 2007). However, in both conditions, HKT2;1 expression levels were more than two fold higher in high KUE lines, a difference that was highly significant in the minus NaCl condition ($p=0.015$) but less so in the plus NaCl treatment ($p=0.066$).

Although no significant association signals were detected, further Na⁺ transporters may be involved in tissue K⁺ substitution by Na⁺: For example, OsHKT1;5 is involved in shoot Na⁺ exclusion by retrieving Na⁺ from the xylem stream and via phloem recirculation (Kobayashi et al., 2017). Downregulation of this mechanism during low K⁺ conditions could therefore augment K⁺ substitution. Other HKTs such as OsHKT2;2, which is primarily root located and could mediate uptake of both K⁺ and Na⁺ (Oomen et al., 2012), is another potential contributor.

Conclusions

A clearer picture of the physiological and molecular underpinnings of KUE variability would be extremely useful in developing high KUE crops. Differences in KUE can be achieved through various mechanisms including: an altered cellular K⁺ distribution, especially between vacuole and cytoplasm; tissue K⁺ distribution, i.e. preferential allocation of K⁺ to the most sensitive tissue such as translocation to the shoot; changes in K⁺ uptake capacity, especially at low external K⁺; changes in K⁺ supply such as enhancing available soil K⁺ via root exudation; and the functional replacement of K⁺ with other ions such as Na⁺ and Ca²⁺. The relative contribution of these mechanisms is largely unknown and may depend on plant species, developmental stage and soil properties.

In this study, KUE was explored using a rice diversity panel. Variation in KUE was found to be considerable and the underlying genetic architecture was examined. By deliberately applying high stringency criteria KUE-related high resolution QTLs were discovered that identified K⁺ substitution by Na⁺ as a likely component of KEU in low K⁺ conditions. Although it is likely that multiple Na⁺ and K⁺ transporters play a role in this process, OsHKT2;1 emerged as the prime suspect responsible for increased Na⁺ uptake. This transporter and other identified candidates could serve as breeding targets to improve crop performance during low K⁺ conditions.

Supplementary Data

Suppl. Tables: spreadsheet Tables containing extended genotype and phenotype data.

Suppl. Figure 1: correlations between growth and tissue Na⁺ concentrations.

Suppl. Figure 2: all Manhattan plots of GWAS analyses

Acknowledgements: This work was supported by a BBSRC doctoral grant to TNH

References

Allen AG, Cardoso AA, Wiatr AG., Machado CMD, Paterlini WC, Baker J. 2010. Influence of intensive agriculture on dry deposition of aerosol nutrients. *Journal of the Brazilian Chemical Society* **21**, 87-97.

Armengaud P, Breitling R, Amtmann A. 2004. The potassium-dependent transcriptome of *Arabidopsis* reveals a prominent role of jasmonic acid in nutrient signaling. *Plant Physiology* **136**, 2556–2576.

Asher CJ, Ozanne PG. 1967. Growth and potassium content of plants in potassium cultures maintained at constant potassium concentrations. *Soil Science* **103**, 155-161.

Aulchenko YS, Ripke S, Isaacs A, van Duijn CM. 2007. GenABEL: An R library for genome-wide association analysis. *Bioinformatics* **23**, 1294–1296.

Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society, Series B (Methodological)* **57**, 289–300.

Brentup, F, Pallière, C. 2008. Energy efficiency and greenhouse gas emissions in European nitrogen fertilizer production and use. York: International Fertiliser Society.

Camargo GGT, Ryan MR, Richard TL. 2013. Energy use and greenhouse gas emissions from crop production using the Farm Energy Analysis Tool. *BioScience* **63**, 263-273.

Campbell MT, Bandillo N, Al Shiblawi FRA, et al. 2017. Allelic variants of OsHKT1;1 underlie the divergence between indica and japonica subspecies of rice (*Oryza sativa*) for root sodium content. *PLoS Genetics* **13**, DOI: 10.1371/journal.pgen.1006823

375 **Ciceri D, Manning DAC, Allanore A.** 2015. Historical and technical developments of
376 potassium resources. *Science of The Total Environment* **502**, 590–601.

377 **Crowell S, Korniliev P, Falcão A, Ismail A, Gregorio G, Mezey J, McCouch S.** 2016.
378 Genome-wide association and high-resolution phenotyping link *Oryza sativa* panicle
379 traits to numerous trait-specific QTL clusters. *Nature Communications* **7**, DOI:
380 10.1038/ncomms10527

381 **Davis JL, Armengaud P, Larson TR, Graham IA, White PJ, Newton AC, Amtmann**
382 **A.** (2018). Contrasting nutrient–disease relationships: Potassium gradients in barley
383 leaves have opposite effects on two fungal pathogens with different sensitivities to jas-
384 monic acid. *Plant Cell Environment* **41**, 2357–2372.

385 **Devlin B, Roeder K.** 1999. Genomic control for association studies. *Biometrics* **55**,
386 997–1004.

387 **Eizenga GC, Ali ML, Bryant RJ, Yeater KM, McClung AM, McCouch SR.** 2014. Reg-
388 istration of the Rice Diversity Panel 1 for Genomewide Association Studies. *Journal of*
389 *Plant Registrations* **8**, DOI: 10.3198/jpr2013.03.0013crmp

390 **Fageria NK.** 1976. Influence of potassium concentration on growth and potassium up-
391 take by rice plants. *Plant and Soil* **44**, 567–573.

392 **Famoso AN, Zhao K, Clark RT, Tung C-W, Wright MH, Bustamante C, Kochian L V,**
393 **McCouch SR.** 2011. Genetic architecture of aluminum tolerance in rice (*Oryza sativa*)
394 determined through genome-wide association analysis and QTL mapping. *PLoS Genet-*
395 *ics* **7**, DOI: 10.1371/journal.pgen.1002221

396 **Fang Y, Wu W, Zhang X, Jiang H, Lu W, Pan J, Hu J, Guo L, Zeng D, Xue D.** 2015.
397 Identification of quantitative trait loci associated with tolerance to low potassium and re-
398 lated ions concentrations at seedling stage in rice (*Oryza sativa* L.). *Plant Growth Regu-*
399 *lation* **77**, 157–166.

400 **FAO.** 2017. World fertilizer trends and outlook to 2020. Rome: FAO.

401 **Garcia-Oliveira AL, Tan L, Fu Y, Sun C.** 2009. Genetic identification of quantitative
402 trait loci for contents of mineral nutrients in rice grain. *Journal of Integrative Plant Biol-*
403 *ogy* **51**, 84–92.

404 **Horie T, Brodsky DE, Costa A, Kaneko T, Lo Schiavo F, Katsuhara M, Schroeder**
405 **JL.** 2011. K⁺ Transport by the OsHKT2;4 Transporter from Rice with Atypical Na⁺ Trans-

port Properties and Competition in Permeation of K^+ over Mg^{2+} and Ca^{2+} Ions. *Plant Physiology* **156**, 1493–1507.

Horie T, Costa A, Kim TH, Han MJ, Horie R, Leung HY, Miyao A, Hirochika H, An G, Schroeder JI. 2007. Rice OsHKT2;1 transporter mediates large Na^+ influx component into K^+ -starved roots for growth. *EMBO Journal* **26**, 3003–3014.

Kong F-M, Guo Y, Liang X, Wu C-H, Wang Y-Y, Zhao Y, Li S-S. 2013. Potassium (K) effects and QTL mapping for K efficiency traits at seedling and adult stages in wheat. *Plant and Soil* **373**, 877–892.

Kobayashi NI, Yamaji N, Yamamoto H, Okubo K, Ueno H, Costa A, Tanoi K, Matsumura H, Fujii-Kashino M, Horiuchi T, Al Nayef M, Shabala S, An G, Ma JF, Horie T. 2017. OsHKT1;5 mediates Na^+ exclusion in the vasculature to protect leaf blades and reproductive tissues from salt toxicity in rice. *Plant Journal* **91**, 657–670.

Koyama ML, Levesley A, Koebner RM, Flowers TJ, Yeo a R. 2001. Quantitative trait loci for component physiological traits determining salt tolerance in rice. *Plant Physiology* **125**, 406–422.

Kumar V, Singh A, Mithra SVA, et al. 2015. Genome-wide association mapping of salinity tolerance in rice (*Oryza sativa*). *DNA Research* **22**, 133–145.

Lin HX, Zhu MZ, Yano M, Gao JP, Liang ZW, Su WA, Hu XH, Ren ZH, Chao DY. 2004. QTLs for Na^+ and K^+ uptake of the shoots and roots controlling rice salt tolerance. *Theoretical and Applied Genetics* **108**, 253–260.

Maathuis FJM. 2009. Physiological functions of mineral macronutrients. *Current Opinion Plant Biology* **12**, 250-258.

Maathuis FJM. 2013. Signalling and regulation of sodium fluxes in plants. *Journal Experimental Botany* **65**, 849-858.

Maathuis FJM, Amtmann, A. 1999. K^+ nutrition and Na^+ toxicity: The basis of cellular K^+/Na^+ ratios. *Annual Botany* **84**, 123-133.

Manning, DAC. 2010. Mineral sources of potassium for plant nutrition. *Agronomy for Sustainable Development* **30**, 281-294

McCouch SR, Wright MH, Tung CW, et al. 2016. Open access resources for genome-wide association mapping in rice. *Nature Communications* **7**, DOI: 10.1038/ncomms10532

- Miyamoto T, Ochiai K, Takeshita S, Matoh T.** 2012. Identification of quantitative trait loci associated with shoot sodium accumulation under low potassium conditions in rice plants. *Soil Science and Plant Nutrition* **58**, 728–736.
- Oomen RJFJ, Benito B, Sentenac H, Rodríguez-Navarro A, Talón M, Véry AA, Domingo C.** 2012. HKT2;2/1, a K⁺-permeable transporter identified in a salt-tolerant rice cultivar through surveys of natural genetic polymorphism. *Plant Journal* **71**, 750–762.
- Patishtan J, Hartley TN, Fonseca de Carvalho R, Maathuis FJM.** 2017. Genome-wide association studies identify rice salt-tolerance markers. *Plant, Cell and Environment* **41**, 970–982.
- Prinzenberg AE, Barbier H, Salt DE, Stich B, Reymond M.** 2010. Relationships between Growth, Growth Response to Nutrient Supply, and Ion Content Using a Recombinant Inbred Line Population in *Arabidopsis*. *Plant Physiology* **154**, 1361–1371.
- Römheld V, Kirkby EA.** 2010. Research on potassium in agriculture: needs and prospects. *Plant and Soil* **335**, 155–180.
- Sassi A, Mieulet D, Khan I, Moreau B, Gaillard I, Sentenac H, Véry AA.** 2012. The Rice Monovalent Cation Transporter OsHKT2;4: Revisited Ionic Selectivity. *Plant Physiology* **160**, 498–510.
- Shin R.** 2014. Strategies for improving potassium use efficiency in plants. *Molecules and cells* **37**, 575–84.
- Spear SN, Edwards DG, Asher CJ.** 1978. Response of cassava, sunflower, and maize to potassium concentration in solution I. growth and plant potassium concentration. *Field Crops Research* **1**, 375–389.
- USGS.** 2017. Mineral commodity summaries 2017. Washington DC: USGS.
- Wang C, Chen HF, Hao QN, Sha AH, Shan ZH, Chen LM, Zhou R, Zhi HJ, Zhou XA.** 2012. Transcript profile of the response of two soybean genotypes to potassium deficiency. *PLoS ONE* **7**, DOI: 10.1371/journal.pone.0039856
- Wang Y, Wu W-H.** 2015. Genetic approaches for improvement of the crop potassium acquisition and utilization efficiency. *Current Opinion in Plant Biology* **25**, 46–52.
- Wu P, Ni JJ, Luo AC.** 1998. QTLs underlying rice tolerance to low-potassium stress in rice seedlings. *Crop Science* **38**, 1458–1462.
- Yoshida S, Forno, Douglas A, Cock JH, Gomez KA.** 1976. Laboratory manual for physiological studies of rice. Los Banos: The International Rice Research Institute.

Zhao K, Tung C-W, Eizenga GC, et al. 2011. Genome-wide association mapping re-

Table 1: Growth and tissue cation concentrations for high and low KUE accessions

veals a rich genetic architecture of complex traits in *Oryza sativa*. Nature Communica-
tions **2**, DOI: 10.1038/ncomms1467

Zhao Y, Li X, Zhang S, Wang J, Yang X, Tian J, Hai Y, Yang X. 2014. Mapping QTLs
for potassium-deficiency tolerance at the seedling stage in wheat (*Triticum aestivum* L.).
Euphytica **198**, 185–198.

KUE_K

KUE_RGR

	low KUE	high KUE		low KUE	high KUE
RGR	0.088	0.11	RGR	n.d.	n.d.
DW HK (g)	0.45	0.79	DW HK (g)	0.72	0.42
DW LK (g)	0.26	0.47	DW LK (g)	0.29	0.34
ShootK HK (mM)	656	646	ShootK HK (mM)	713	626
ShootK LK (mM)	244	86	ShootK LK (mM)	136	135
ShootNa HK (mM)	47	40	ShootNa HK (mM)	39	66
ShootNa LK (mM)	352	232	ShootNa LK (mM)	197	369
RootK HK (mM)	253	184	RootK HK (mM)	236	170
RootK LK (mM)	57	52	RootK LK (mM)	53	59
RootNa HK (mM)	92	67	RootNa HK (mM)	79	91
RootNa LK (mM)	104	148	RootNa LK (mM)	140	200
ShootK:Na ratio (HK)	14	16.2	ShootK:Na ratio (HK)	18.3	9.5
ShootK:Na ratio (LK)	0.69	0.37	ShootK:Na ratio (LK)	0.69	0.37

483

484

485

486

487

488

489

490

491

Table 2: Summary of quantitative trait loci identified in GWAS				
Trait	Description	Chr	Position	Significant SNP Positions
RGR_LK	Relative growth rate at low K treatment	1	22,260,180 - 22,463,799	22,360,180; 22,361,410; 22,361,482; 22,363,799
RGR_K	K use efficiency defined as RGR/shoot K concentration at LK treatment	1	34,344,598 - 34,563,159	34,444,598; 34,463,159
NaR_LK	Root Na concentration at low K treatment	6	29,440,164 - 29,640,591	29,540,164; 29,540,591
NaS_LK	Shoot Na concentration at low K treatment	6	29,440,164 - 29,640,591	29,540,164; 29,540,591

493

494

495

496

497

498

499

500

501

502

503 **Figure Legends**

Figure 1: Responses of rice genotypes to potassium stress. **a)** Mean plant dry mass of cultivars when grown in the presence of 0.1 (LK) and 1 (HK) mM potassium. **b)** Relative plant dry mass (dry mass LK/ dry mass HK). **c)** Reduction in relative growth rate (RGR) in LK compared to HK conditions.

Figure 2: Distribution of root (top two panels) and shoot (bottom two panels) K^+ concentration across the diversity panel for plants grown on LK (0.1 mM) and HK (1 mM) K^+ medium.

Figure 3: Significant ($p < 0.05$) correlation between RGR reduction and shoot tissue Na^+ concentration of plants grown on LK medium.

Figure 4: Manhattan plots for traits (RGR at LK, KUE-K, root $[Na^+]$ at LK and shoot $[Na^+]$ at LK) that generated significant association signals (arrows) using criteria as explained in the Methods. Note that 'shoot Na ' and 'root Na ' trait data associate with the same locus on chromosome 6.

Figure 5: Co-incidence of previously described QTLs and loci identified in this study related to low K^+ growth in the rice genome. Each bar represents a chromosome and previously reported QTLs are marked in white (Wu et al., 1998), yellow (Miyamoto et al., 2012) or red (Fang et al., 2015). Triangles indicate the position of QTLs derived from this study.

Figure 6: Reducing levels of medium K^+ drastically increases Na^+ concentrations in both roots and shoots of rice cultivar IR64. Plants were grown hydroponically for 7 weeks in the presence of varying K^+ levels and 3 mM NaCl. Error bars show SD of three biological replicates.

Figure 7: qPCR analysis of HKT2;1 expression in roots of 5 high KUE cultivars (GSOR 54, 109, 133, 357 and 366, see Suppl Table 1) and 5 low KUE rice cultivars (GSOR 42, 115, 276, 377 and 401). Plants were grown for 4 weeks in medium containing 0.01 mM K^+ supplemented with 0 or 1 mM NaCl. Data are means for 3 biological replicates with error bars denoting SD.






Fig 2




Fig 3




Fig 4




Fig 5




Fig 6




Fig 7

