



This is a repository copy of *Parents' experiences of having an excessively crying baby and implications for enhancing support services*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/143364/>

Version: Accepted Version

Article:

Garratt, R., Bamber, D., Powell, C. et al. (6 more authors) (2019) Parents' experiences of having an excessively crying baby and implications for enhancing support services. *Journal of Health Visiting*, 7 (3). pp. 132-140. ISSN 2050-8719

10.12968/johv.2019.7.3.132

This document is the Accepted Manuscript version of a Published Work that appeared in final form in *Journal of Health Visiting*, copyright © MA Healthcare, after peer review and technical editing by the publisher. To access the final edited and published work see <https://doi.org/10.12968/johv.2019.7.3.132>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Title: Parents' experiences of having an excessively crying baby and implications for enhancing support services

Introduction

During their first four months, approximately 20% of infants cry for long periods for no apparent reason (Alvarez, 2004; St James-Roberts & Halil, 1991). Traditionally, this has been attributed to gastro-intestinal disorders and referred to as 'infant colic' (Illingworth, 1954; Wessel et al, 1954), but more recent evidence suggests most of these infants are healthy and go on to develop normally (Lehtonen, 2001; Stifter & Braungart, 1992). The crying, which tends to peak at around 1-2 months of age (Barr, 1990; St James-Roberts & Halil, 1991), usually resolves spontaneously by five months and is now thought to be linked to normal development (Barr & Gunnar, 2000; Barr et al, 2005; St James-Roberts et al, 2013).

Whilst much research has focused on the causes of excessive infant crying, there is growing recognition of the need to understand its impact on parental well-being. The crying, and in particular the 'unsoothable' crying bouts often associated with it (Fujiwara et al, 2011), cause considerable alarm and distress to parents. The impact of the crying also depends partly on parents' coping abilities. Factors such as depression and anxiety and influence how parents interpret and respond to infant crying, whilst social isolation may also increase its impact (Laurent & Ablow, 2012; Pearson et al, 2010). As a result, the actual duration of the crying may not be as significant in determining its impact as parents' subjective experience of it. In this context, the term "excessive crying" is used to indicate crying judged by parents to be 'too much', rather than an objective measure of its duration.

Excessive crying has also been found to impact on a range of outcomes for both parent and infant, including premature termination of breastfeeding (Howard et al, 2006), over-feeding (Stifter et al, 2011), parental distress, anxiety and depression (Kurth et al, 2011; Murray & Cooper, 2001; Stifter et al, 2003), poor parent-child relationships (Kurth et al, 2011; Oldbury & Adams, 2015; Tabuchi & Shimada, 2008), problems with long-term child development (Wolke et al, 2002) and, in a small number of cases, infant abuse (Barr et al, 2006; Fairbrother et al, 2015).

In other studies, parents reported feeling unprepared, challenging their expectations of parenthood and leading to feelings of disappointment, both in themselves and their baby (Cox & Roos, 2008; Ellett et al, 2009; Megel et al, 2011; Nash et al, 2008). The perceived stigma and shame of the excessive crying led to self-imposed isolation, reducing support and external contact (Ellett & Swenson, 2005; Megel et al, 2011). Parents described the crying as overwhelming and frustrating, evoking anger and resentment (Gaffney et al, 2008; Landgren & Hallström, 2010), sometimes including thoughts of aggression towards their baby (Fairbrother et al, 2015; Levitsky & Cooper, 2000).

Despite these negative outcomes and experiences, there are currently no evidence-based services in the UK to support parents with an excessively crying baby. As a result, parents consult books, magazines or websites, which often provide conflicting advice (Catherine et al, 2008). They also seek help from health professionals, but as only around 5% of such infants are found to be unwell (Freedman et al, 2009), this represents a significant potentially avoidable cost to health services (Morris et al, 2001). The “Surviving Crying” study (Long et al, 2018) represented a first step towards developing and evaluating an intervention to support parents worried about their baby’s excessive crying. To inform the development of the intervention, the first

stage of the study aimed to verify and provide a more detailed understanding of existing qualitative findings within a UK health service context. To achieve this, focus groups were held to consult with parents with recent experience of an excessively crying baby. This paper reports the findings from these focus groups and the implications for practice.

Methods

Aim

To explore the experiences of UK parents with a baby who had previously cried excessively, and to obtain their views on what support they would have wanted.

Design

This was the qualitative aspect of the development phase of a larger mixed methods feasibility study (Long et al, 2018).

Ethical approval

Study public registration no.ISRCTN84975637; ethical approval provided by De Montfort University (13450) and the National Research Ethics Committee East Midlands (Nottingham) (project ID 152836; NRES: 14/EM/1202).

Recruitment

Parents who had previously had an excessively crying baby were identified by Health Visitors/Community Public Health Nurses (HV/CPHNs) using the following criteria:

- The crying had taken place in the first 6 months and was now resolved;
- The baby had no underlying health problems at the time of the crying;

- Parents were English speaking, or supported by an English speaker.

HV/CPHNs obtained consent from the parents for a researcher to contact them. The study was also publicised through the National Childbirth Trust and in local community settings, enabling parents to contact the research team directly. On receipt of an expression of interest, the researchers telephoned parents to explain the study and confirm eligibility. Eligible parents were sent a letter and information sheet and then followed up to confirm their interest and availability. To facilitate participation, parents were able to bring their babies and/or older children. In order to obtain a diversity of views within the time constraints of the study, a sample size of 20 was set.

Data Collection

Data were collected between April-August 2015, using a semi-structured format to guide discussion. Having obtained written consent, parents were asked open-ended questions about their experiences of having an excessively crying baby: what they found challenging; what had helped them to manage; family, community and health service support; and what resources should be provided as part of routine NHS care.

Whilst the initial intention was to collect all data through focus groups, on a number of occasions only one parent attended for the focus group and in these instances an interview was conducted using the same format. Groups were facilitated by two researchers, whilst two others wrote observations and provided informal childcare when needed. Roles were shared between two researchers for interviews. All parents received shopping vouchers in acknowledgement of their participation.

Four focus groups of 2 - 5 parents and six individual interviews were held, a total of ten events. Sessions were audio-recorded for later verbatim transcription and anonymisation.

Data Analysis

Thematic analysis was used to analyse the data (Braun & Clarke, 2014) supported by NVivo10 software (International Pty Ltd, 2012). One researcher conducted a brief preliminary analysis of the transcripts to produce a summary of key themes, which was agreed by the team. A second researcher then undertook a full analysis, reading all transcripts several times and conducting a line-by-line analysis of the content. Data were initially coded to a large number of themes, which were refined by grouping similar themes under broader headings. These themes were shared with the research team and, following discussion, refined and agreed.

The findings informed the development of the intervention resources, and the draft resources were shared with parents and HV/CPHNs to ensure the team had accurately understood parents' needs and the current context of healthcare provision.

Results

Eighteen mothers and two fathers (both partners of participating mothers) took part (Table 1 gives demographic details). Most participants were white British, with high education levels, and married or living with a partner. Nearly half of the babies were second or third-born, with eight of the 18 families having other children who had not cried excessively. Only one parent had experienced two excessively crying babies.

Three key themes were identified: Disrupted expectations and experiences of parenthood; Stigma and social isolation; and Seeking support and validation of experience.

1. Disrupted Expectations and Experiences of Parenthood

Participants reflected that prior to their baby's birth, their expectations were positive, with the majority having little or no awareness of the phenomenon of excessive crying. They were therefore shocked and unprepared for the experience.

I think the, the expectation that you have after you have been carrying your unborn child and then feeling this is going to be wonderful, and the screaming *just doesn't stop, and you think to yourself "it's going to stop, it will just be the first week" and it didn't for a long, long time.* (P1P19)

Parents tried many strategies to comfort their babies but these often had no effect. Their inability to soothe their babies frequently produced feelings of inadequacy, frustration and anxiety, and eroded their confidence in their parenting ability:

I would say it made you feel like you are no good at it [being a parent], like you are helpless really, because you try everything. I expected when you have a baby that when you pick them up and you comfort them they are going to stop crying. But when she carried on crying you try and feed them, you change them, there is nothing wrong with them you just feel like useless.
(P1P7)

Whilst some experiences were linked to being a new parent, participants who already had other children also experienced shock and disrupted expectations. They recounted being taken aback by the difference from their previous experiences (even

in one instance where their previous baby had cried excessively), and how this undermined their belief in their skills as a parent.

Because this is, he's my third and the other two haven't been like it, people think "oh well you should, you know, you know what you are doing", but when they are totally different you feel at a bit of a loss. (P1P14)

Managing their baby's crying also often resulted in parents being physically exhausted, and when the crying continued at night, sleep deprivation further undermined their ability to cope. Even basic self-care could be compromised:

I wasn't looking after myself, I wanted to but I just couldn't because, everything was so consumed with the crying that I just didn't, I just almost neglected myself. (P1P13)

The experience of managing the crying also created tensions in relationships sometimes, particularly at night when parents were most tired. During the day, many mothers were alone at home with their baby and some felt that the strain of this situation was not fully understood or acknowledged by their partner, creating friction between the couple:

I don't think he really understood what it was like to be at home all day on your own with a crying baby. And whilst he never came home and said "well it's a mess, where is my dinner?" you still have to do those things. He'd come in and be, "oh I am so tired, I have been at work", and expect to have a chill out and she would be there screaming the place down. I started to resent him. (P1P16)

In contrast, others described examples of a positive shared parental approach:

He would do the night feeds and then I would get up at 6. With the crying to *be fair he never lost his temper once, he didn't shout*. In the middle of the *night he would wake up and say "oh give her to me"*. And he slept downstairs for three months with her so that I got sleep. (P1P5)

2. Stigma and Social Isolation

Even where parents were managing to work together to cope with the excessive crying, its unexpected nature and other people's reactions posed significant challenges. Parents perceived that both they and their baby were somehow different from others who appeared to have a 'normal baby'.

"The year I had my baby there were five of us that all went to school that all had our babies within a few months. All of their babies were happy, they never particularly cried. So we were very much the odd ones out". (P1P16).

Negative comparisons between their situation and that of others often created a sense of isolation and loneliness. Whilst parents recognised that meeting friends or attending groups could be beneficial, the all-consuming task of trying to soothe their baby, combined with physical exhaustion, made going out both physically and mentally challenging. Furthermore, sometimes their sense of isolation and not being "normal" was only compounded by these experiences:

"I went to them baby massage classes but I only went for one...because he was crying the whole time we were there. I couldn't do a lot with him because I just thought it's just gonna be embarrassing. Especially cos' you'd have all the mums there and they'd be like "oh my baby is like so", "he's brilliant, sleeps 8 hours, happy to just sit there", but that don't make you feel better." (P1P17)

Discussing their crying baby at parent and baby groups was seen as taboo, as was admitting to peers that they were finding the experience of parenthood difficult and struggling to cope:

It's not the kind of thing where you come to a baby group ...and you say "do you know what, I have got a really troublesome baby". You just don't because you are conditioned to think I have got to go and talk about how lovely my baby is. (P1P3)

More generally, when parents did take their babies out, they often experienced intolerance and criticism from others, which reinforced their perceptions of being inadequate parents, further undermining their confidence:

And you always get people who think they are doing good by coming up and "oh what's the matter with her?" and actually it just makes you feel worse as a parent. (P1P6)

I remember going in (supermarket) with her and she was absolutely screaming... and this old lady said to me, "why can't you just shut your baby up, do they have to cry all the way round the shop?" (P1P5)

Feelings of failure and inadequacy affected parents' willingness to seek support, even in some cases from partners or close family:

It's sort of like admitting defeat isn't it....you feel like you have been defeated by this little child and it's actually admitting that to your family. Even my other half I never told him how bad I actually felt. (P1P5).

Parents' reluctance to ask for help was also associated with not wanting to "inflict *[baby] on other people*" (P1P1) and this hesitancy was sometimes confirmed when friends and family were unwilling to help or avoided visiting because of the crying:

Nobody would even babysit for me or anything either. My *mum couldn't cope with her because she didn't sleep. My mum would just say "I can't do it".* We never ever got a break because nobody could deal with her because she was so hard. (P1P5)

Notably, for some, a reluctance to seek help was fuelled by fears of being judged by others, particularly health professionals, who they feared would take their baby away.

"But you do think that don't you? "Oh my god they (health professionals) are going to come round and take her off me". (P1P1)

3. Seeking Support and Validation of Experience

Feelings of inadequacy and difference also impacted more widely on parents' help-seeking experiences and interactions with health professionals. Parents described repeatedly going through mental checklists of routine causes of crying such as nappy changing, feeding and temperature. When they had exhausted these options, they became concerned their baby was unwell. Many reported repeated internet searching, particularly of professionally endorsed websites, to find other explanations for the excessive crying, but struggling to find specific information:

"Yeah, (I used the internet)a lot, when I was trying to look on Google, some stuff came up but it wasn't stuff that gave you proper advice or anything on what to do." (P1P10)

In addition to reliable factual information, online peer support was appreciated by some, particularly given the difficulties experienced in trying to go out. Parent forums offered important validation and reassurance as they discovered they were not the only ones dealing with an excessively crying baby.

“I was just looking for what other people had been through and what they’d said; the whole act of finding and reading what other parents have been through helped me - I felt like I wasn’t on my own.” (P1P18)

Whilst online information and peer support provided some help, for most parents this was not sufficient to allay their concerns. Despite their reluctance, they therefore sought advice and reassurance from health professionals, particularly doctors and HV/CPHNs, and described experiencing a wide range of responses.

In most cases no physical cause for the crying was found, but as it continued, many parents returned repeatedly to health professionals, becoming convinced that there was something seriously wrong:

“It (crying) went on and on for two weeks straight it was, he just cried so much. I remember going to the doctors’ and then the midwife was coming round and I asked, “what’s he crying for?”, And I just kept looking at him and thought “what’s wrong with him?”, why is he crying?”. (P1P12)

A few parents felt their concerns were not taken seriously by health professionals or that they were considered to be exaggerating, which increased their frustration and distress.

“You are banging your head against a brick wall, you are knocking on everyone’s door and you are not getting anywhere. So not only did I feel

undervalued *from a mother's point of view* [and] feel that what I was doing was wrong, but as an individual, as an adult I felt like nobody was listening to me, nobody cared, nobody is taking me seriously, it was a nightmare.”

(P1P11)

In other instances, health professionals were supportive, but parents questioned their knowledge and skills in relation to excessive crying:

“I found that the health visitor was helpful, she was quite sympathetic. She *didn't have the answers that I felt that I needed, and she didn't have probably the experience of babies that cry*. But it just made it just nicer to talk to somebody who was sympathetic to talk to. (P1P8)

or found that the support offered did not fully meet their needs or expectations:

“I used to think she (health visitor) should come to my house and see what I *am going through. She never came to my house, she'd ring me and she was just trying to tell me things to do over the phone.*” (P1P7).

For a few parents, lack of continuity of support was an additional difficulty, and the need to repeatedly explain their situation made them less likely to share their anxieties:

“It's a lot of work, because we have had so many health visitors it's like starting from scratch all over again. He has three already, he's only 15 months old and not one of them knows him. So when they ring I just say “yes I am fine”. I just don't bother now.” (P1P5)

Other parents described examples of highly supportive care. Home visits from HV/PCHNs were particularly valued, allowing a health professional to see the crying

at first-hand, and in doing so validating parents experience. Continuity of care was also seen as important, enabling the development of a much-valued relationship:

I knew you were only meant to see them (health visitors) routinely at 6 weeks, so for her to offer to come back the next week, I took that as, *“oh that’s really nice, like really”*, so when she came back the second week it was lovely, she’d come, she’d come to see how I was. (P1P1)

“My health visitor was really supportive, even now she’s really supportive. She was really nice, really good, even like now she is really good, and she always like rings me to see how I am. She has given me her number and said if you ever feel down, if you ever want to talk to anyone ring me and, I go to the groups and she is there sometimes and she has a chat and she is really helpful.” (P1P9)

Discussion

Participants’ accounts clearly described how having an excessively crying baby had presented significant emotional and physical challenges. Whilst the crying had now resolved, in some cases more than a year previously, many parents still became distressed recalling the difficulties they had faced. The excessive crying had shattered their preconceptions of parenting a new baby and, because it was not a phenomenon they were previously aware of, led them to question whether there was something wrong. When no medical problem could be identified, many parents felt a sense of personal inadequacy and failure, leading them to withdraw from social contact and be reluctant to seek help. The focus group findings concur with other studies (Landgren & Hallström, 2010; Megel et al, 2011), where parents attempted to maintain a façade of having the ‘perfect baby’ by removing themselves from public

scrutiny. Lack of understanding and support from others, including health professionals in some instances, increased their distress and isolation.

Stigma theory (Goffman, 1990) provides an insight into the behaviour of those who perceive themselves, or are perceived by others, not to fulfil the characteristics associated with socially constructed identity categories. Whether consciously or not, individuals are usually very aware of the expectations of society, and of how others might perceive them as failing to make “the standard”, resulting in feelings of shame and low esteem (Goffman, 1990). In relation to parenting, there are expectations that it should be done ‘well’ and without undue distress or difficulty (Maushart, 1997; Phoenix et al, 1991). Such notions are reinforced by widespread media representations of smiling parents with perfectly behaved infants (Kaplan, 1993). However, as was evident for the participants, these expectations of parenthood frequently meet reality with significant dissonance (Belsky et al, 1986; Guendouzi, 2005; Lawrence et al, 2007). Goffman (Goffman, 1990) suggests that when there is doubt as to whether individuals fulfil the socially required characteristics, or they possess an attribute that makes them different from others, this lack or attribute is regarded as a stigma and can have a discrediting effect, being seen as a shortcoming or failing, with individuals being viewed as different or ‘other’. These experiences were evident in participants’ perceptions of themselves and their babies as ‘different’, with these feelings being reinforced by other people’s reactions. Their perceptions of themselves being inadequate parents lead them to be reluctant to seek help or admit they were not coping as they felt they should be.

Implications for practice

This study has highlighted how unprepared parents were for having an excessively

crying baby and how prior knowledge would have helped them cope. Concerns about 'doing something wrong' or being a 'bad parent' resulted in an initial reluctance to seek help which was often only overcome as the crying continued. These are important considerations for health professionals supporting parents in the antenatal and immediate postnatal period. Parents need timely, accessible and evidence-based information about excessive infant crying, acknowledgement of the challenges of managing the crying, and reassurance that finding it difficult is not an indicator of poor parenting.

Other research into initiatives to reduce perceived health-related stigma has identified the importance of education and counselling in raising awareness and modifying behavioural and cognitive responses to experiences (Heijnders & Van der Meij, 2006). However, because stigma is socially constructed (Corrigan, 2005), interventions need to be aimed not only at those directly affected, but also the wider population (Heijnders & Van der Meij, 2006). Increasing awareness of excessive crying and its impact could therefore enable health professionals to support parents more effectively, increase public tolerance, and help prospective parents re-fashion their expectations, reducing feelings of inadequacy. Consideration also needs to be given to offering internet-based group support, as this can allow individuals experiencing stigma to interact with others in a relatively anonymous fashion, providing a way to belong which may not otherwise be available (Parker Oliver et al, 2017).

Some parents reported disappointment at the level of care and support received from health professionals. The schedule of HV/CPHN contacts reported in this study was generally reflective of the universal service delivery of the Healthy Child Programme (DOH, 2009; 2011; PHE, 2016), including an antenatal home visit, new

birth visit and further contact at 6 weeks and 4 months. Specific enhanced care packages can be offered, including support for unsettled infants, but this study indicates the need for a more specialised package to support parents with excessively crying babies. Specific recommendations are listed in Table 2.

Limitations

Participants were mostly white middle class educated women supported by a partner, and future research needs to explore the perspectives of fathers, ethnic minorities, single parents, and vulnerable and disadvantaged families.

During the time when focus group participants had their excessively crying baby, substantial organisational changes to the NHS HV/CPHN services were being undertaken which may have influenced experiences of health visiting services, including disruptions to the continuity of provision.

CONCLUSION

Parents reported that having an excessively crying baby was an unexpected and distressing experience that undermined their confidence in their parenting skills and led them to isolate themselves. Participants felt that more information before or soon after their baby's birth might have helped them to cope better. Understanding the experience of excessive crying and the perceived stigma associated with it can enable development of better support for parents, including the education and training of health professionals that care for them. The findings highlight a currently unmet need in the National Health Service.

Key points

- Parents were unprepared for and distressed by their baby's excessive crying

- Feelings of failure made parents reluctant to disclose concerns and seek help
- Some experienced a lack of understanding from health professionals and general public
- Specific resources and services are needed to help parents cope with excessively crying babies
- Greater public awareness of excessive crying could increase tolerance and support

Financial support

This research was funded by the National Institute for Health Research HTA Programme Grant Number: 12/150/04

Department of Health and Social Care Disclaimer

This report presents independent research commissioned by the National Institute for Health Research (NIHR) HTA Programme. The views and opinions expressed by authors in this publication are those of the authors and do not necessarily reflect those of the NHS, the NIHR, MRC, CCF, NETSCC, the HTA Programme or the Department of Health and Social Care.

Acknowledgements

The authors thank the families who participated, the NIHR HTA Programme for its support, the staff of Leicestershire Partnership NHS Trust (LPT) for their assistance, and our Steering Committee and Management Group for their guidance. In particular, the study could not have taken place without the help of the following LPT staff: Gail Melvin, Research Manager; Joanna McGarr, Trainee Research Assistant; Lynn Hartwell, Research Nurse; and the many health visitors who recruited parents

and provided feedback. We thank our other UK collaborators: Dr Elaine Boyle, University of Leicester; Charlie Owen, University College London; Sally Rudge, Counselling Psychologist & CBT Practitioner; Rachel Plachcinski, National Childbirth Trust; Jan and John Bullen, Cry-Sis. Four university or charity research groups gave permission to evaluate their parental support materials during Stage 1 of the study and we are grateful to them and their institutions for this support: Ron and Marilyn Barr for the PURPLE Crying Programme; Harriet Hiscock and Fallon Cook of Murdoch Children's Research Institute for Cry Baby; Jane Fisher and Heather Rowe, Monash University, for What Were We Thinking; Denise Coster and Peter Richards of the National Society for Prevention of Cruelty to Children, for Coping with Crying.

REFERENCES

- Alvarez, M. (2004) Caregiving and early infant crying in a Danish community. *Journal of Developmental & Behavioural Paediatrics*, 25(2), 91-98.
- Barr, R. (1990) The normal crying curve: what do we really know? *Developmental Medicine and Child Neurology*, 32(4), 356-362.
- Barr, R. & Gunnar, M. (2000) Colic: the 'transient responsivity' hypothesis, in Barr, R., Hopkins, B. & Green, J. (eds), *Crying as a sign, a symptom and a signal*. London: MacKeith Press, 41-66.
- Barr, R., Paterson, J., MacMartin, L., Lehtonen, L. & Young, S. (2005) Prolonged unsoothable crying bouts in infants with and without colic. *Journal of Developmental and Behavioural Paediatrics*, 26(1), 14-22.
- Barr, R., Trent, R. & Cross, J. (2006) Age-related incidence curve of hospitalized shaken baby syndrome cases: convergent evidence for crying as trigger to shaking. *Child Abuse & Neglect* 30(1), 7-16.
- Belsky, J., Ward, M. & Rovine, M. (1986) Prenatal expectations, postnatal experiences, and the transition to parenthood, in Ashmore, R. & Brodzinsky, B. (eds), *Thinking about the family*. Hillsdale, NJ: Erlbaum, 139-145.
- Braun, V. & Clarke, V. (2014) Using thematic analysis in psychology. *Qualitative research in psychology*, 3(2), 77-101.

- Catherine, N., Ko, J. & Barr, R. (2008) Getting the word out: Advice on crying and colic in popular parenting magazines. *Journal of Developmental and Behavioral Pediatrics*, 29(6), 508-511.
- Corrigan, P. (2005) *On the stigma of mental illness: Practical strategies for research and social change* American Psychological Association.
- Cox, L. & Roos, V. (2008) The experiences of first-time mothers with colic infants who seek help from medical professionals. *Health SA Gesondheid* 13(1), 4-13.
- DOH (2009) *Healthy Child Programme - Pregnancy and the First Five Years of Life*. London: Available online:
https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/167998/Health_Child_Programme.pdf [Accessed
- DOH (2011) *Health Visitor Implementation Plan - A Call to Action*. London: Available online:
https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/213718/dh_131064.pdf [Accessed
- Ellett, M., Appleton, M. & Sloan, R. (2009) Out of the abyss of colic: a view through the father's eyes. *American Journal of Maternal Child Nursing*, 34(3), 164-171.
- Ellett, M. & Swenson, M. (2005) Living with a colicky infant. *Gastroenterology Nursing* 28(1), 19-25.
- Fairbrother, N., Barr, R., Pauwels, J., Brant, R. & Green, J. (2015) Maternal thoughts of harm in response to infant crying: an experimental analysis. *Archives of Women's Mental Health*, 18(3), 447-455.
- Freedman, S., Al-Harthy, N. & Thull-Freedman, J. (2009) The crying infant: diagnostic testing and frequency of serious underlying disease. *Pediatrics*, 123(3), 841-848.
- Fujiwara, T., Barr, R., Brant, R. & Barr, M. (2011) Infant distress at five weeks of age and caregiver frustration. *The Journal of Pediatrics*, 159(3), 425-430.
- Gaffney, K., Beckwitt, A. & Friesen, M. (2008) Mothers' reflections about infant irritability and postpartum tobacco use. *Birth*, 35(1), 66-72.
- Goffman, E. (1990) *Stigma: Notes on the management of spoiled identity*. London: Penguin Books.
- Guendouzi, J. (2005) "I feel quite organised this morning": How mothering is achieved through talk. *Sexualities, Evolution and Gender*, 7(1), 17-35.
- Heijnders, M. & Van der Meij, S. (2006) The fight against stigma: An overview of stigma-reduction strategies and interventions. *Psychology, Health and Medicine*, 11(3), 353-363.
- Howard, C., Lanphear, N., BP, L., Eberly, S. & Lawrence, R. (2006) Parental responses to infant crying and colic: the effect on breastfeeding duration. *Breastfeeding Medicine*, 1(3), 146-155.
- Illingworth, R. (1954) Three month colic. *Archives of Diseases in Childhood*, 29, 165-174.

International Pty Ltd, Q. (2012) NVivo 10.

Kaplan, E. (1993) *Women and film: both sides of the camera*. London: Routledge.

Kurth, E., Kennedy, H., Spichiger, E., Hosli, I. & Stutz, E. (2011) Crying babies, tired mothers: what do we know? A systematic review. *Journal of Midwifery*, 27(2), 187-194.

Landgren, K. & Hallström, I. (2010) Parents' experience of living with a baby with infantile colic-a phenomenological hermeneutic study. *Scandinavian Journal of Caring Sciences*, 25(2), 317-324.

Laurent, H. & Ablow, J. (2012) A cry in the dark: Depressed mothers show reduced neural activation to their own infant's cry. *Social Cognitive and Affective Neuroscience*, 7(2), 125-134.

Lawrence, E., Nylen, K. & Cobb, R. (2007) Prenatal expectations and marital satisfaction over transition to parenthood. *Journal of Family Psychology*, 21(2), 155-164.

Lehtonen, L. (2001) From colic to toddlerhood in Barr, R., St James-Roberts, I. & Keefe, M. (eds), *New Evidence on unexplained early infant crying: Its origins, nature and management*. New Brunswick, NH: Johnson & Johnson Pediatric Institute, 259-271.

Levitsky, S. & Cooper, R. (2000) Infant colic syndrome: Maternal fantasies of aggression and infanticide. *Clinical Paediatrics*, 39(7), 395-400.

Long, J., Powell, C., Bamber, D., Garratt, R., Brown, J., Dyson, S. & St James-Roberts, I. (2018) Development of materials to support parents whose babies cry excessively: findings and health service implications. *Primary Healthcare Research and Development*, 19(4), 320-332.

Maushart, S. (1997) *The mask of motherhood, how mothering changes everything and why we pretend it doesn't*. Sydney, Australia: Random House.

Megel, M., Wilson, M., Bravo, K., McMahon, N. & Towne, A. (2011) Baby lost and found: mothers' experiences of infants who cry persistently. *Journal of Pediatric Health Care*, 25(3), 144-152.

Morris, S., James-Roberts, I., Sleep, J. & Gillham, P. (2001) Economic evaluation of strategies for managing crying and sleeping problems. *Archives of Disease in Childhood*, 84(1), 15-19.

Murray, L. & Cooper, P. (2001) The impact of irritable infant behavior on maternal mental state: a longitudinal study and a treatment trial, in Barr, R., St James-Roberts, I. & Keefe, M. (eds), *New evidence on unexplained early infant crying: its origins, nature and management*. Skillman, NJ: Johnson & Johnson Pediatric Institute, 149-164.

Nash, C., Morris, J. & Goodman, B. (2008) A study describing mothers' opinions of the crying behaviour of infants under one year of age. *Child Abuse Review* 17(3), 191-200.

Oldbury, S. & Adams, K. (2015) The impact of infant crying on the parent-infant relationship. *Community Practitioner* 88(3), 29-34.

Parker Oliver, D., Patal, S., Benson, J., Gage, A., Washington, K., Kruse, R. & Demiris, G. (2017) The Effect of Internet Group Support for Caregivers on Social Support, Self-Efficacy, and Caregiver Burden: A Meta-Analysis. *Telemedicine and e-Health*, 23(8), 621-629.

Pearson, R., Cooper, R., Penton-Voak, I., Lightman, S. & Evans, J. (2010) Depressive symptoms in early pregnancy disrupt attentional processing of infant emotion. *Psychological medicine*, 40(4), 621-631.

PHE (2016) Best start in life and beyond: Improving public health outcomes for children, young people and families. Guidance to support the commissioning of the Healthy Child Programme 0-19 years: Health Visiting and School Nurse Service. Commissioning Guide 1: Background information and commissioning and service model. London:

Phoenix, A., Woollett, A. & Lloyd, E. (1991) *Motherhood, meanings, practices and ideologies*. Newbury Park, California: Sage.

St James-Roberts, I., Alvarez, M. & Hovish, K. (2013) Emergence of a developmental explanation for prolonged crying in 1 to 4 month old infants: review of the evidence. *Journal of Paediatric Gastroenterology and Nutrition*, 57(Suppl 1), S30-36.

St James-Roberts, I. & Halil, T. (1991) Infant crying patterns in the first year: normal community and clinical findings. *Journal of Child Psychology and Psychiatry*, 32(6), 951-968.

Stifter, C., Anzman-Frasca, S., Birch, A. & Voegtline, K. (2011) Parent use of food to soothe infant/toddler distress and child weight status. An exploratory study. *Appetite*, 57(3), 693-699.

Stifter, C., Bono, M. & Spinrad, T. (2003) Parent characteristics and conceptualizations associated with the emergence of infant colic. *Journal of Reproductive and Infant Psychology*, 21(4), 309-322.

Stifter, C. & Braungart, J. (1992) Infant colic: a transient condition with no apparent effects. *Journal of Applied Developmental Psychology*, 13(4), 447-462.

Tabuchi, N. & Shimada, K. (2008) Maternal characteristics and feelings of distress resulting from the crying of high-risk infants. *Japan Journal of Nursing Science* 5(2), 99-108.

Wessel, M., Cobb, J., Jackson, E., Harris, G. & Detwiler, A. (1954) Paroxymal fussing in infancy, sometimes called "colic". *Paediatrics*, 14(5), 421-433.

Wolke, D., Rizzo, P. & Woods, S. (2002) Persistent infant crying and hyperactivity problems in middle childhood. *Pediatrics*, 109(6), 1054-1060.