



This is a repository copy of *Animal mobility in Chalcolithic Portugal: Isotopic analyses of cattle from the sites of Zambujal and Leceia*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/144121/>

Version: Accepted Version

Article:

Wright, E. orcid.org/0000-0002-0770-3898, Waterman, A.J., Peate, D.W. et al. (3 more authors) (2019) Animal mobility in Chalcolithic Portugal: Isotopic analyses of cattle from the sites of Zambujal and Leceia. *Journal of Archaeological Science: Reports*, 24. pp. 804-814. ISSN 2352-409X

<https://doi.org/10.1016/j.jasrep.2019.02.005>

Article available under the terms of the CC-BY-NC-ND licence
(<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

34 The Portuguese Chalcolithic (c. 3000-1900BC) was a pivotal time of social and economic
35 change with evidence of increasing social complexity resulting in the formation of hierarchical
36 settlements. With these changes came the emergence of long-distance exchange networks and
37 more complex population movements and interactions. Domesticated animals would have played
38 an important role in these emerging economies, and it is assumed that animals migrated with,
39 and were exchanged by, humans as part of these new networks. While direct evidence of these
40 networks is still limited in this region, new methodologies have the potential to expand our
41 knowledge of animal mobility and exchange. This study uses $^{87}\text{Sr}/^{86}\text{Sr}$ ratios in tooth enamel to
42 identify potential non-local animals at these two settlements. Results indicate that Leceia may
43 have had a higher proportion of non-local animals than Zambujal and had a wider catchment area
44 for its stock, suggesting variations in settlement economies across relatively short distances in
45 this region. These results have important implications for our understanding of animal
46 management at Portuguese Chalcolithic sites, and the involvement of animals in the emerging
47 economies of the time.

48

49 Keywords: Iberia; Zambujal; Leceia; Copper Age; strontium isotopes; mobility; fauna

50

51

52 **1. Introduction**

53

54 The Portuguese Chalcolithic (c. 3000 - 1900BC) was a time of significant social and economic
55 change, with evidence of the expansion of complex settlements and exchange networks (cf.
56 Cardoso, 2007; Gonçalves, 1999; Lillios, 1995; Uerpmann, 1995; Valera et al.,
57 2017). Domesticated animals played an important role in these economies, and the
58 archaeological record demonstrates that people commonly raised domesticated cattle, pigs, sheep
59 and goats for meat and secondary products (Harrison 1985; Valente and Carvalho, 2014). It is
60 assumed that domesticated animals moved with humans, and were sold or exchanged by people
61 as part of these new networks. However, direct evidence of these exchanges is still limited in this
62 region. In the last decade, radiogenic isotope studies ($^{87}\text{Sr}/^{86}\text{Sr}$) have provided a means of
63 identifying migrant human and animals in the Iberian peninsula (Carvalho et al., 2016; Díaz-del-
64 Río, 2017; Diaz-Zorita Bonilla, 2013; Díaz-Zorita Bonilla et al., 2018; Waterman et al., 2014)

65 and elsewhere in the world (cf. Knudson et al., 2016; Madgwick et al., 2017; Price et al., 2015;
66 Zhao, 2015). In this paper we outline the results of new strontium isotopic analyses from cattle
67 recovered from the Chalcolithic fortified settlement sites of Castro do Zambujal (Zambujal) and
68 Leceia (both in the Estremadura region) and use this data to identify migrant animals within
69 these settlements.

70

71 **1.1 Social complexity and exchange networks in Chalcolithic Portugal**

72 The Chalcolithic period in Portugal saw the rise of large ditched-enclosed, fortified hilltop, and
73 walled settlement types. The archaeological record for this time suggests that with the rise of
74 these complex settlements, long and short distance exchange networks emerged (Cardoso, 2003;
75 Gonçalves, 2000; 2001; Jorge, 2000). Alongside these important socio-economic changes there
76 is variation in site type and function, especially between regions (e.g. Cardoso, 2007; Gonçalves
77 et al., 2013); with the establishment of new large fortified sites, mostly in the Estremadura
78 region. The appearance of new types of material culture also in this region, made of materials
79 such as copper, slate, variscite, amphibolite and ivory, has provided strong evidence of the
80 establishment of long-distance exchange networks (e.g. Cardoso and Carvalhosa, 1995; Cardoso
81 and Schuhmacher, 2012; Cardoso et al., 2013; Gauß, 2015; Lillios, 1997; Müller et al., 2007;
82 Odriozola et al., 2010; Odriozola et al., 2013; Roberts, 2008; Schuhmacher et al., 2009;
83 Schuhmacher, 2012; Schuhmacher, 2017). Archaeological data suggests that some of these
84 materials may have been travelling from areas of southern Iberia, or even northern Africa, into
85 the Estremadura region (Schumacher et al., 2009). Using strontium isotopes, a number of studies
86 have now found direct evidence of human migrants into the Estremadura region. The isotopic
87 signature of these migrants indicates that some of them may have also travelled in from areas of
88 southern Iberia (such as the Alentejo region) possibly as part of these exchange networks
89 (Carvalho et al., 2016; Waterman et al., 2014). It is likely that these migrants were bringing
90 domesticated animals with them as they entered new regions, but to date no study focusing on
91 animal mobility in the Estremadura region has been completed.

92

93 **1.2 Animal husbandry in Chalcolithic Portugal**

94 The variations between site types across the region are clearly reflected in the animal remains
95 recovered at settlement sites (Valente and Carvalho, 2014). Fortified sites in the Estremadura

96 region tend to yield large faunal assemblages, dominated by the three main domesticated animals
97 (cattle, sheep/goat and pig), with evidence for the use of secondary products (i.e. milk, wool, and
98 traction) (Cardoso and Detry, 2002; von den Driesch and Boessneck, 1976), whereas smaller
99 unfortified sites have yielded larger proportions of wild species, indicating a different approach
100 to animal exploitation, as part of a more mobile type of existence (e.g. Cabaço, 2010; Correia,
101 2015; Davis and Mataloto, 2012; Moreno-Garcia and Sousa, 2013; Valente 2013). The two sites
102 included in this study, Leceia and Zambujal, are two of the largest well-known fortified sites in
103 the Estremadura region (Kunst, 2017; Becker and Flade-Becker, 2017). Both of their
104 assemblages show a preponderance of domestic species, with only small proportions of wild
105 species. The use of secondary products, as seen at Portuguese fortified sites such as these,
106 reflects the intensification in animal husbandry seen across Iberia at this time. This
107 intensification has long been linked to the formation of networks for exchanging animals and
108 animal products (e.g. Harrison 1985).

109

110 **2. Strontium isotope ratios and landscapes**

111 The strontium isotope signature of a geographic region is controlled by the nature of the
112 underlying geology (rock lithology) and permeates its landscape and groundwater. This signature
113 is absorbed into the biological tissues of local plants and animals (Faure and Powell, 1972; Gilli
114 et al., 2009). In animals, strontium isotopes are incorporated into tooth and bone through
115 ingestion of water and food. This is due to a physiological process in which Sr substitutes for
116 calcium in the mineral component (hydroxyapatite) of hard tissues (Bentley, 2006; Ericson,
117 1985; Sealy et al., 1991; Schroeder et al., 1972:496). Unlike carbon, nitrogen and oxygen
118 isotopes that are used in many archaeological studies of prehistoric diet, once incorporated into
119 biological tissues, strontium isotopes do not fractionate (i.e. no change in $^{87}\text{Sr}/^{86}\text{Sr}$), when passed
120 from prey to consumer. Therefore, an organism's strontium isotope signature directly reflects the
121 bioavailable strontium in its environmental range, rather than its trophic level (Graustein, 1989).
122 Due to how Sr infiltrates biological hard tissues during formation, humans and animals residing
123 in the same territorial ranges and consuming only local plants and animals, should exhibit similar
124 $^{87}\text{Sr}/^{86}\text{Sr}$ signatures (Tommasini, 2018). In contrast, humans and animals should exhibit
125 differences in strontium isotope ratios between regions that are geologically distinctive. When
126 sufficient geologic heterogeneity is present across regional landscapes, humans and animals can

127 migrate between areas with significant divergences in local $^{87}\text{Sr}/^{86}\text{Sr}$ values. If animals or
128 humans die and are interred in a location with a marked difference in local $^{87}\text{Sr}/^{86}\text{Sr}$ values from
129 their own biological tissues, they can be recognized as migrant individuals. This method for
130 identifying migrants has been used productively in many archaeological studies of human and
131 animal migration patterns (cf. Bentley, 2006; Crowley et. al, 2017; Price et al., 2002; Price et al.,
132 2012).

133
134 Because using $^{87}\text{Sr}/^{86}\text{Sr}$ values to identify nonlocal humans and animals requires geologic
135 diversity over reasonably traversable distances, geologically homogeneous regions may limit this
136 method's effectiveness. Portugal and Western Spain exhibit marked differences in geological
137 lithologies, thereby making this region an excellent location to use strontium isotope
138 fingerprinting to study human and animal migration in prehistory. Nonetheless, we must keep in
139 mind that this methodology can only provide a minimum estimate of mobility (minimum number
140 of migrants, MNM), as it is not possible to distinguish between humans and animals who
141 originate from settlements that share similar bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ values due to similar
142 underlying geology. Additionally, it is important to note that this method assumes that local
143 populations only consumed local foods and drank from local water sources, as consumption of
144 large amounts of foreign foods can change $^{87}\text{Sr}/^{86}\text{Sr}$ values (Burton and Hahn, 2016).

145

146 **2.1 Archaeological sites and regional geology**

147 The prehistoric settlement sites of Leceia and Zambujal lie in the Estremadura region of Portugal
148 (Figure 1). The Estremadura is a historically-defined province in the southwestern region of
149 Portugal which encompasses both the Lisbon and Setúbal peninsulas and extends westward to
150 the Atlantic coast. Both Leceia and Zambujal lie close to the coast and to (former) estuaries and
151 interior waterways and, thus, would have been key places for both coastal and interior trade
152 networks.

153

154 **2.1.1 Zambujal**

155 Zambujal is one of the most prominent and well-known prehistoric fortified settlements in
156 Portugal (Sangmeister and Schubart, 1981). It was occupied from the Chalcolithic period until
157 the early Bronze Age (c. 2900-1700 cal BC), and was subject to several important building

158 phases with a series of walls and other fortifications (Arnold and Kunst, 2011; Kunst, 2010;
159 Kunst, 2018; Sangmeister and Schubart, 1981). Geomagnetic prospections, archaeological
160 surveys and excavations between 1994 and 2013 have indicated that the fortified settlement was
161 larger than previously thought, and is currently estimated at c. 26 hectares (Kunst and Uerpmann,
162 2002; Kunst et al. 2013; Kunst 2017a; Becker and Flade-Becker, 2017). Material culture
163 recovered from Zambujal suggests that it was permanently settled throughout the 3rd millennium
164 BC and that craft and metal production took place there, likely at a household scale (Müller et
165 al., 2007; Gauß, 2015; Kunst et al., 2016). Additionally, as evidenced by the recovery of raw
166 materials with origins outside of the region --such as copper, amphibolite, ivory and gold -
167 Zambujal was an important center of local and regional exchange networks, (Kunst, 1995;
168 Sangmeister and Schubart, 1981; Uerpmann, 1995; Uerpmann and Uerpmann, 2003).

169

170 By the end of the excavations that took place from 1964 to 1973, over 150,000 faunal specimens
171 had been recovered, representing over 95 species. Domestic cattle, pigs and caprines dominate,
172 but wild boar, aurochs and red deer are also present, albeit in comparatively small numbers (von
173 den Driesch and Boessneck, 1976; 1981). Caprines and pigs were the main focus of the domestic
174 assemblage, with smaller numbers of cattle.

175

176 **2.1.2 Leceia**

177 The fortified settlement of Leceia (c. 3500-2200 cal BC), had earlier origins than Zambujal,
178 during the late Neolithic period, but was also abandoned earlier - before the Early Bronze Age.
179 Excavations at the site between 1983 and 2002 revealed a complex settlement fortified by a
180 defensive series of stone walls and towers (Cardoso, 1994; 1997; 2000; 2010). Leceia has
181 received less attention than Zambujal, but its location (elevated above the river Barcarena), its
182 size, and its rich material culture, suggest it was an important regional center for agriculture,
183 material goods, production and trade (Cardoso, 2000). In contrast to Zambujal, there is poor
184 evidence for extractive metallurgy. Neither smelting slags or copper ore were found, but the
185 presence of many copper artefacts, including some which are unfinished, suggest that some
186 copper working was undertaken here (Müller and Cardoso, 2008). Other imported materials,
187 such as amphibolite, have also been recovered, indicating that the site was linked to broader
188 exchange networks (Cardoso and Carvalhosa, 1995; Cardoso, 2004). Faunal remains recovered

189 from Leceia were studied by Cardoso and Detry (2002), and were dominated by domestic cattle,
190 pigs and caprines throughout the period of occupation. Red deer represented the main wild
191 species but, unlike at Zambujal, no aurochs or wild boar were identified. Cattle, pigs and
192 caprines were present in equal proportions during the late Neolithic period but an increase in
193 caprines along with a decrease in cattle was seen during the Chalcolithic period, bringing the
194 overall proportions of these animals broadly in line with that seen at Zambujal. In contrast to
195 Zambujal no isotopic work on either human or faunal remains has previously been undertaken.

196

197 **2.1.3 Regional Geology**

198 The sites of Zambujal and Leceia both lie in the geologically diverse landscape of the
199 Estremadura region of Portugal (Figure 2). The area occupies a portion of the Lusitanian Basin,
200 which is a northern Atlantic basin that was created during a late Triassic rifting phase. In the
201 southeast, this basin connects to the Alentejo and the Algarve Basins and in the north and east it
202 abuts the Late Paleozoic Hercynian basement rocks of the Iberian Meseta (Cunha and dos Reis,
203 1995; Wilson, 1988). The Lusitanian Basin, mainly composed of Cretaceous and Jurassic
204 sediments with northern pockets of Triassic sediments, is geologically younger than other parts
205 of Portugal and Spain with a heterogeneous mix of lithologies including sandstones and
206 limestones, clays, marls, and some volcanic rocks (Azerêdo et al., 2002; Wilson, 1988: See
207 Waterman, 2012 for more detail). Because it is a carbonate-dominated Mesozoic landscape,
208 Lusitanian Basin sediments should have $^{87}\text{Sr}/^{86}\text{Sr}$ close to marine values (0.707-0.710: e.g.
209 Schneider et al., 2009). Additionally, as a coastal region, seawater rainfall and sea spray can be
210 incorporated into the terrestrial food chain which may also contribute to $^{87}\text{Sr}/^{86}\text{Sr}$ values that are
211 close to that of seawater (0.709) (Bentley, 2006). Slightly higher $^{87}\text{Sr}/^{86}\text{Sr}$ values should be found
212 in some parts due to variations in clastic deposits, and local water analyses have recorded
213 $^{87}\text{Sr}/^{86}\text{Sr}$ ratios of 0.709-0.711 (Voerkelius et al., 2010). In contrast, the older Palaeozoic
214 Hercynian basement metamorphic and granitic rocks of the Portuguese interior should generally
215 have more radiogenic values ($^{87}\text{Sr}/^{86}\text{Sr} > 0.713$: e.g. Bea et al., 2003).

216

217 **3. Materials and Methods**

218 **3.1 Sampled materials**

219 For this study 16 cattle teeth from Leceia and 27 cattle teeth from Zambujal were analyzed.
220 (Table 1). The Leceia samples come from material that is housed in the Centro de Estudos
221 Arqueológicos do Concelho de Oeiras/Câmara Municipal de Oeiras (Bacarena, Portugal),
222 previously studied by Cardoso and Detry (2002). The Zambujal samples are from the von den
223 Driesch and Boessneck faunal assemblage which is currently housed at the Leonel Trindade
224 Municipal Museum (Torres Vedras, Portugal). The Zambujal samples selected are dominated by
225 specimens dated to the early Chalcolithic (c. 3000-2500 cal BC), whereas the majority of our
226 specimens from Leceia were dated to the Full/Late Chalcolithic (c. 2500-2200 cal BC).

227

228 Left third molars were prioritized to ensure that each sample was from a separate individual, and
229 one enamel slice per tooth was taken, in order to maximize the number of individuals being
230 investigated. A transversal slice of enamel was cut from the base of the protoconid using a
231 diamond cutter disc coupled to a dentist drill (following the method outlined in Minniti et al.,
232 2014). Only fully formed teeth, with closed roots and which were in wear, were used. This made
233 the location independent from wear, and ensured that the samples were unaffected by any
234 potential age bias. Samples of both enamel and dentine were taken for 18 animals in order to
235 examine any intra-individual variation indicative of relationships between mobility and life
236 history. In order to test for potential effects of sample diagenesis 10 samples were divided during
237 the wet chemistry process and one portion underwent additional washes of acetic acid before
238 further processing.

239

240 **3.2 Determining Local Values**

241 The most established method for estimating the local $^{87}\text{Sr}/^{86}\text{Sr}$ range for a region is by using the
242 mean of sampled local faunal or human remains ± 2 s.d. to account for the upper and lower limits
243 of the range. Because nonlocal outlier samples can skew the local estimate, tooth and bone
244 samples are best taken from animals with very limited geographic ranges (e.g. rabbits) (Bentley
245 et al., 2004; Price et al., 2002). Plants, water and soil samples can also provide local estimates,
246 but may not provide as accurate of an account of the local bioavailable range as animal tissues.
247 For this study local small fauna (rabbits) and plants were collected and analyzed to help
248 determine the local bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ isotope composition at Leceia and to reaffirm the local
249 range for Zambujal that was presented in Waterman et al., (2014).

250

251 **3.3 Wet Chemistry and Mass Spectrometry**

252 All chemical processing of the samples was carried out at the University of Iowa Department of
253 Earth & Environmental Sciences clean laboratory. Details of the laboratory protocol used for this
254 analysis followed the procedures outlined in Waight et al., (2002), see Waterman et al., (2014)
255 for a full description. $^{87}\text{Sr}/^{86}\text{Sr}$ ratios were measured using a Nu Plasma HR multicollector
256 inductively-coupled-plasma mass-spectrometer (MC-ICP-MS) in the Department of Geology at
257 the University of Illinois at Urbana-Champaign. Samples were introduced to the machine using a
258 Nu Instruments DSN-100 desolvator system equipped with a nebulizer with an aspiration rate
259 near 0.1 mL min^{-1} . The samples were alternately run with standards (SRM 987, SCS coral and
260 E&A) using a sample-standard-bracketing measurement protocol wherein standards were run
261 every 3-5 samples (Rehkämper et al. 2004). The ^{88}Sr beam intensities for all samples and
262 standards ranged from 4 to 12 V (100 ppb solutions). Masses of ^{83}Kr to ^{88}Sr were measured
263 during a single cycle comprised of 2 blocks of 25 scans (5 s integration per scan) with a 40 s
264 baseline determination using ESA-deflected signals. Instrumental mass bias was internally
265 normalized to an $^{86}\text{Sr}/^{88}\text{Sr}$ ratio of 0.11940 and then corrected ratios were normalized to the
266 NIST SRM 987 international standard value of 0.710268 (which had a reproducibility of \pm
267 0.000013; 2 s.d., n=47) to correct for day-to-day variability. No corrections were necessary for
268 Sr introduced as part of sample production as procedural blanks were $< 100 \text{ pg Sr}$.

269

270 **4. Results**

271 Results are presented in Tables 1-2 and Figure 3. The 10 separated samples subjected to a more
272 rigorous regiment of acid washes showed negligible differences in Sr values to the control
273 samples (0.00019-0.00003), suggesting diagenesis was not a major concern for the dental
274 enamel. Both sites yielded results with a relatively wide range of enamel values. At Zambujal
275 enamel samples had an $^{87}\text{Sr}/^{86}\text{Sr}$ range of 0.7054 to 0.7127, while at Leceia the range was wider,
276 from 0.7046 to 0.7179. For Leceia, based upon the sampled leaves and small fauna, the local
277 $^{87}\text{Sr}/^{86}\text{Sr}$ value range is estimated to be 0.7067-0.7077. For Zambujal the $^{87}\text{Sr}/^{86}\text{Sr}$ local range
278 was previously defined as 0.709-0.7115 (Waterman et al., 2014). The small fauna and leaves
279 tested in this analysis fell right below the lower end of this spectrum, thus we have adjusted the
280 local range for Zambujal slightly to 0.7085-0.7115. This range fits the majority of the heavily

281 clustered Zambujal fauna in this study. The dentine samples had a much smaller range than the
282 enamel at both sites. As dentine is more likely to be contaminated with the Sr isotope signature
283 from the local soil, or to remodel to the local Sr isotope signature after movement into a new
284 area, this more limited value range was expected. However, whilst at Zambujal all of the dentine
285 values fall within the calculated local range, at Leceia the dentine range is much wider and some
286 samples deviate from the calculated local range. This is an intriguing pattern which will be
287 discussed further below.

288

289 **4.1 Enamel values**

290 The cattle enamel samples from Zambujal and Leceia show some divergence in the $^{87}\text{Sr}/^{86}\text{Sr}$
291 ratio with more specimens from each site clustering around the defined local ranges (Figure 3).
292 Using the nonparametric Mann-Whitney U test (due to the non-normal distribution of the data),
293 the differences between the sites are found to be extremely statistically significant ($p=0.000$)
294 (Table 3). These differences are also demonstrated using box plots in Figure 4. Density plots
295 (Figure 5) also show that cattle from Leceia display a wider range of values than Zambujal,
296 despite having a smaller sample size, indicating that Leceia had a wider catchment area for its
297 cattle.

298

299 At Zambujal a number of specimens plot outside of its local range. Those falling above are in the
300 range of 0.712-0.713, but those below are more spread out from 0.705-0.708. Some of these fall
301 within the local range calculated for Leceia (0.706-0.708), but one specimen falls even below
302 this range (0.705). At Leceia a number of specimens have values that are higher than the site's
303 local range. Most of these fall into the Zambujal range, but there is also one very high value
304 (0.7179), well above the local range calculated for either site. There are also some specimens that
305 fall in the region of the lowest values from Zambujal (<0.705), lower than the range from either
306 site. These results highlight the possibility that cattle were being moved between these two sites,
307 but also that some of them must have been brought in from further afield.

308

309 **4.2 Dentine and enamel pairs**

310 For six cattle from Zambujal and three from Leceia both enamel and dentine were sampled from
311 the same animals (this was also attempted for four other individuals, but one sample failed in

312 each case). These dentine and enamel sets were compared in order to look for evidence of
313 lifetime mobility (Table 2, Figure 6).

314

315 At Zambujal, all the cattle for which both dentine and enamel samples were taken exhibited very
316 low $^{87}\text{Sr}/^{86}\text{Sr}$ variation between samples (0.0001-0.0012). Additionally, all of these values fell
317 within the local range for Zambujal. None of these individuals, therefore, show clear evidence of
318 having been moved into Zambujal from outside of the region between the time of enamel
319 formation and dentine remodeling.

320

321 At Leceia two of the three sampled cattle exhibited relatively consistent enamel and dentine Sr
322 values. However, these values (on both the enamel and dentine) are all lower than the calculated
323 Leceia local range. For the third animal the enamel value was higher than local Leceia range
324 (0.7084) (close to the low-end of the Zambujal range), while the dentine value was below the
325 local Leceia range (0.7051) (similar in value from the first two cattle). This suggests that this
326 animal grew up in a different region from where it last lived. Additionally one other dentine
327 sample from Leceia (without an associated enamel sample due to machine error) exhibits a Sr
328 value which is fairly high (0.7122), matching some enamel samples from Zambujal. This is a
329 very high value to have been found in this area, and could potentially result from some kind of
330 contamination.

331

332 **5. Discussion**

333 The results from this study indicate that cattle were being moved into, out of, and within the
334 Estremadura region during the Chalcolithic period. This movement likely occurred most
335 commonly within the Estremadura region, and potentially between Leceia and Zambujal
336 themselves, which were two of the most prominent sites in the region. However, the data
337 gathered here suggests that some movement from more distant regions was also occurring.

338

339 In general we can organize the Sr values for the tested cattle into 4 broad groups. These groups
340 account for all the sampled cattle with the exception of the outlying sample from Leceia with a
341 very high value of 0.7179. These groups are:

- 342 1. Cattle in the range of 0.7085-0.7115 that match local values for Zambujal and sites in the
343 Zambujal region (from Waterman et al., 2014),
- 344 2. Cattle in the range of 0.706-0.708 that match local values from Leceia.
- 345 3. Cattle in the range of 0.712-0.713 which fall above the values for Zambujal.
- 346 4. Cattle in the range of 0.704 -0.706 which fall below local values for Leceia

347

348 In the Iberian Peninsula $^{87}\text{Sr}/^{86}\text{Sr}$ isotope ratio ranges are available for some geological areas
349 based on water, soil, and plant data (e.g. Freitas et al., 2003; Moita et al., 2009; Schneider et al.,
350 2009; Villaseca et al., 2009; Voerkelius et al., 2010), and predictions about likely $^{87}\text{Sr}/^{86}\text{Sr}$
351 isotope ratio ranges can be made for other areas based upon the local geologic lithologies. While
352 detailed maps of bioavailable $^{87}\text{Sr}/^{86}\text{Sr}$ isotope ratios have not yet been completed for Portugal
353 and Spain, a number of archaeological studies, focusing on individual archaeological sites or
354 regions, have been completed in the last decade. These studies provide us with local bioavailable
355 $^{87}\text{Sr}/^{86}\text{Sr}$ isotope range for a number of places in central and south-west Iberia that we can
356 compare with the data from Zambujal and Leceia (Table 4, Figure 7). By examining these
357 ranges, the diversity of local and regional geology, and the archaeological evidence for exchange
358 networks in the region, we can begin to identify possible cattle origin and movement patterns in
359 the Zambujal and Leceia regions.

360

361 Groups 1 and 2: These groups are composed of animals local to Zambujal and Leceia with
362 $^{87}\text{Sr}/^{86}\text{Sr}$ isotope values of c. 0.707-0.712. These values are consistent with the underlying
363 geology of most of the Estremadura region—mainly Mesozoic or Cenozoic sediments with small
364 pockets of other lithologies. These are the most common types of underlying geology across
365 Europe, and are present in many parts of Iberia. Local value ranges from most sites in the
366 Estremadura region including Leceia and Zambujal (this study) the Zambujal region (Waterman
367 et al., 2014), Bom Santo (Carvalho et al., 2016), and (part of the range of) Rego de Murta
368 (Waterman et al., 2013) all fall into this range. These values are also present in some parts of the
369 Alentejo region, such as at the site of Monte de Cegonha (Saragoça et al. 2016), as well as at a
370 number of Spanish sites, including Valencina-Castilleja (Diaz-Zorita-Bonilla, 2013), and sites
371 near Madrid (Díaz-del-Río et al., 2017). We are therefore unable to rule out the possibility that
372 some of the sampled cattle in this group are, in fact, non-local but from a region with similar

373 local $^{87}\text{Sr}/^{86}\text{Sr}$ isotope values. However, the most parsimonious approach to this dataset is to
374 assume these are animals raised locally and, as they are most numerous at both sites, that most
375 cattle were raised and consumed locally during this time period.

376

377 Group 3: 'Non-local' animals with values between 0.712 and 0.713. These are likely to be from
378 an area with underlying Middle to Upper Palaeozoic sediments (Voerkelius et al., 2010). There
379 are multiple areas in Iberia which have this kind of geology, including in southern Portugal in
380 both the eastern Alentejo and Algarve regions, but also in southern Spain as far away as
381 Andalusia. In south-west Iberia the most prominent area with this kind of geology is the Ossa
382 Morena Zone (OMZ) located in southern areas of Portugal and south-west Spain. The range of
383 values here fits with those from La Pijotilla in south-west Spain, which is in this geological zone,
384 but also at some sites near Madrid (Díaz-del-Río et al., 2017). However, in closer proximity the
385 burial sites of Rego de Murta, located (170km) north and east of Zambujal have a calculated
386 local $^{87}\text{Sr}/^{86}\text{Sr}$ isotope of ~0.711-0.713 (based on small fauna) but many humans and some larger
387 fauna recovered from the burial had Sr isotope values in the 0.713-0.714 range. This suggests
388 that this region could also be a place of origin for animals in the Group 2 category.

389

390 Group 4: 'Non-local' animals with values which fall below 0.707. These are likely to be from an
391 area with basaltic volcanic rocks such as the Quaternary and Tertiary volcanic rocks or in regions
392 of basic Palaeozoic volcanic rocks. There are some small areas of the southern Estremadura
393 which have the potential for yielding Sr isotope values this low. These are located to both the
394 east and west of Leceia. No Sr isotope values are currently available from these areas, so these
395 can only be considered as a possibility.

396

397 These groups and places of possible origin cover all of the cattle except the migrant animal with
398 the very high Sr isotope value at 0.7179. This animal is likely to be from an area with underlying
399 Lower Palaeozoic sediments (Voerkelius et al., 2010). Some areas of the OMZ also have this
400 kind of geology, including parts of Alentejo. Perdigões, which is in this region has the highest
401 local values of any of the sites presented, with a local range of up to 0.7135-0.7145 (Žalaitė et
402 al., 2018). However, the region around Perdigões has very varied geology, and as part of that
403 study baseline samples were taken from a 10 km radius around the site in order to account for

404 this. Some of the values that resulted from this were even higher –up to 0.7184, which indicates
405 that such high values are possible in this area. Waterman et al. (2014) found one human from the
406 Cova da Moura burial near Zambujal to have a Sr isotope value even higher than the outlier
407 animal (0.720), and the Waterman et al. (2013) study on humans from the Rego da Murta burials
408 in the Ribatejo region of Portugal found three individuals who have Sr isotope values in the
409 0.717-0.720 range. This suggests that another potential place of origin may be from older
410 geologic formations northeast of the Estremadura region.

411
412 The presence of raw materials and artefacts at Zambujal and Leceia provide more evidence of
413 the links between the Estremadura and other regions. We know that variscite, slate, amphibolite,
414 arsenical copper ore, and other materials from the Alentejo region (OMZ) commonly made their
415 way into the Estremadura during the Chalcolithic period. Provenance studies based on lead
416 isotope analysis of artefacts from both Zambujal and Leceia have also indicated that their raw
417 materials come from ore deposits found in the OMZ (Gauß, 2013; Gauß, 2015; Müller and
418 Cardoso, 2008). This increases the likelihood that at least some of the individuals from our result
419 group 3 originate from this region.

420
421 Finally, some attention should be given to the differences in catchments between these two
422 settlement sites, and why Leceia may have received livestock from a wider area than Zambujal.
423 This is particularly interesting in the light of further isotopic work we have been undertaking in
424 parallel to this study, which has indicated differences in cattle diet between the two sites (Wright
425 et al., in prep). One possibility is that Leceia was geographically better located for contact with
426 outside regions than Zambujal. It is located slightly closer to the Alentejo region, for example,
427 and may have been a first point of contact for people travelling up into the Estremadura region
428 from the south. Leceia is also located much closer to geology that could potentially yield low Sr
429 isotope values, although this must remain a tentative suggestion until more Sr isotope mapping
430 of the region is undertaken. A second option is that these patterns could be related to a temporal
431 trend. As the sample from Leceia is dominated with specimens from Full/Late Chalcolithic
432 layers, whereas the majority of the Zambujal sample is earlier in date. This could be reflecting an
433 increase in cattle mobility through time during the Chalcolithic period, through increased use and

434 consolidation of exchange networks in south-west Iberia. Larger datasets from more sites are
435 needed to be able to investigate this further.

436

437 An alternative explanation may be related to differences in husbandry practices between the two
438 settlements. Perhaps Zambujal was more effective at breeding and keeping its own herds than
439 Leceia was, so it had less need to incorporate more livestock from outside regions. This is
440 something that needs further investigation. The local environment surrounding each of these
441 settlements needs to be examined more closely in terms of suitability for cattle production, and
442 more detailed attention needs to be given to the differences in the faunal assemblages between
443 the two sites, tasks which are beyond the scope of this paper.

444

445 **6. Conclusion**

446 This paper provides the results of one of the first strontium isotope studies focusing on cattle
447 remains, animal mobility, and social organization in southern Portugal. Using data on cattle from
448 the expansive Portuguese Chalcolithic (3000/2900-2000 BC) settlement sites of Zambujal and
449 Leceia (Estremadura, Portugal), we provide evidence that cattle were circulating through the
450 region with non-local animals being documented at both sites. Results indicate that cattle at
451 Leceia had a wider catchment area for its stock than Zambujal, with more migrant animals.
452 Domesticated animals would have played an important role in these emerging economies of
453 these sites, and these exchange networks likely overlap with human mobility and the exchange of
454 other trade goods. Thus, these findings have important implications for our understanding of
455 long and short distant trade and regional economic integration. We suggest that it is likely that
456 cattle with non-local Sr values higher than the $^{87}\text{Sr}/^{86}\text{Sr}$ local range for Zambujal range may have
457 originated in the Ossa Morena Zone as other evidence of direct exchange links with prominent
458 sites in the Alentejo region, such as Perdigões, are documented. However, the origin for the
459 highest $^{87}\text{Sr}/^{86}\text{Sr}$ values (>0.718) are still being investigated.

460

461 Future studies exploring the involvement of animals in exchange networks in Chalcolithic south-
462 west Iberia will require larger Sr datasets, alongside regionally focused zooarchaeological work
463 comparing animal husbandry regimes. While considerably more Sr isotope values are available
464 for prehistoric sites in south west Iberia than were just a few years ago, further research is needed
465 to provide regional baseline maps. To this end, the Australian National University/Griffith
466 University Strontium Basemap Project currently being undertaken by colleagues has great
467 potential for furthering our interpretation of this data set in the future.

468

469 **Acknowledgements**

470 This work was funded through an Archaeological Institute of America Archaeology in Portugal
471 Fellowship. Thanks to Silvia Valenzuela-Lamas for useful discussions on Iberian prehistoric
472 animal mobility, and to Mount Mercy Students Matthew Purl and Nicholas Wilson for their help
473 processing samples at the University of Iowa. Also thanks to Craig Lundstrom, Tom Johnson,
474 and Gideon Bartov of the University of Illinois at Urbana-Champaign for their help with the
475 Mass Spectrometry and to the Museu Municipal of Torres Vedras and the Centro de Estudos
476 Arqueológicos do Concelho de Oeiras/Câmara Municipal de Oeiras for allowing access to the
477 material included in this study. Finally, thank you to the two reviewers for their useful comments
478 and suggestions.

479

480 **References Cited**

481 Arnold, F., Kunst, M., 2011. Zur Rekonstruktion kupferzeitlicher Befestigungsanlagen auf der
482 Iberischen Halbinsel–Turm B von Zambujal (Torres Vedras, Lisboa, Portugal). Madrider
483 Mitteilungen 52, 36-86 and Farbtaf. 1 and Taf. 1-12.

484

485 Azerêdo, A.C., Wright, V.P., Ramalho, M.M., 2002. The Middle-Late Jurassic forced regression
486 and disconformity in central Portugal: Eustatic, tectonic and climatic effects on a carbonate ramp
487 system. *Sedimentology* 49 (6), 1339-1370.

488

489 Bea, F., Montero, P., Zinger, T., 2003. The nature, origin, and thermal influence of the granite
490 source layer of central Iberia. *Journal of Geology* 111, 579-595.

491

492 Beard, B.L., Johnson, C.M., 2000. Strontium isotope composition of skeletal material can
493 determine the birth place and geographic mobility of humans and animals. *Journal of Forensic*
494 *Sciences* 45, 1049–1061.

495

496 Becker, H., Flade-Becker, A.-S., 2017. Zambujal 2013, Teil 2. Magnetische Prospektionen.
497 *Madridrer Mitteilungen* 58, 31-56.

498

499 Bentley, R.A., 2006. Strontium isotopes from the earth to the archaeological skeleton: A review.
500 *Journal of Archaeological Method and Theory* 13, 135-187. [https://doi.org/10.1007/s10816-006-](https://doi.org/10.1007/s10816-006-9009-x)
501 [9009-x](https://doi.org/10.1007/s10816-006-9009-x)

502

503 Bentley, R.A., Price, T.D., Stephan, E., 2004. Determining the ‘local’ $^{87}\text{Sr}/^{86}\text{Sr}$ range for
504 archaeological skeletons: a case study from Neolithic Europe. *Journal of Archaeological Science*
505 31, 365–375. <https://doi.org/10.1016/j.jas.2003.09.003>

506

507 Burton, J.H., Hahn, R., 2016. Assessing the “local” $^{87}\text{Sr}/^{86}\text{Sr}$ ratio for humans. In: Grupe, G.,
508 McGlynn, G.C., (Eds.) *Isotopic Landscapes in Bioarchaeology*. Springer, Berlin, Heidelberg.
509 p113-121

510

511 Cabaço, N., 2010. Restos faunísticos em contexto do Neolítico final do Sector Q do recinto dos
512 Perdigões (Reguengos de Monsaraz). *Apontamentos de Arqueologia e Património* 5, 27–30.

513

514 Cardoso, J.L., 1994. Leceia 1983-1993. Escavações do povoado pré-histórico, *Estudos*
515 *Arqueológicos de Oeiras no especial*.

516

517 Cardoso, J.L., 1997. O povoado de Leceia sentinela do Tejo no terceiro milénio antes Cristo.
518 *Instituto Português de Museus*.

519

520 Cardoso, J.L., 2000. The Fortified Site of Leceia (Oeiras) in the Context of the Chalcolithic in
521 Portuguese Estramadura. *Oxford Journal of Archaeology* 19(1), 37–55.

522

523 Cardoso, J.L., 2004. Polished stone artefacts at the prehistoric settlement of Leceia (Oeiras).
524 *Madrider Mitteilungen. Mainz.* 45: 1-32.

525

526 Cardoso, J.L., 2003. O comércio de matérias-primas de origem geológica dos meados do VI
527 milénio a.C. aos finais do III milénio a.C., no Ocidente peninsular: breve ensaio. *Boletim da*
528 *Sociedade de Geografia de Lisboa* 121, 91-106.

529

530 Cardoso, J.L., 2007. *Pré-História de Portugal*. Universidade Aberta, Lisboa.

531

532 Cardoso, J.L., 2010. Povoado pré-histórico de Leceia (Oeiras): evolução arquitectónica do
533 sistema defensivo e das técnicas construtivas correlativas. *Transformação e Mudança no centro*
534 *e sul de Portugal: o 4.º e o 3.º milénios a.n.e.* Colóquio Internacional (Cascais, 2005). *Actas* 43-
535 63.

536

537 Cardoso, J.L., Carvalhosa, A.B., 1995. Estudos petrográficos de artefactos de pedra polida do
538 povoado pré-histórico de Leceia (Oeiras). *Análise de proveniências. Estudos Arqueológicos de*
539 *Oeiras* 5, 123-151.

540

541 Cardoso, J.L., Detry, C., 2002. Estudo Arqueozoológico Dos Restos de Ungulados Do Povoado
542 Préhistórico de Leceia (Oeiras). *Estudos Arqueológicos de Oeiras* 10, 131–82.

543

544 Cardoso, J., Schuhmacher, T.X., 2012. Marfiles calcolíticos en Portugal. Estado de la cuestión.
545 *Elfenbeinstudien. Marfil y elefantes en la Península Ibérica y el Mediterráneo occidental.* *Actas*
546 *del coloquio internacional en Alicante el 26 y 27 noviembre 2008.* Darmstadt/Mainz :Verlag
547 *Philipp von Zabern, 95-110 (Iberia Archaeologica Band 16 Faszikel 1).*

548

- 549 Cardoso, J.L., Vilstrup, J.T., Eisenmann, V., Orlando, L., 2013. First evidence of *Equus asinus*
550 L. in the Chalcolithic disputes the Phoenicians as the first to introduce donkeys into the Iberian
551 Peninsula. *Journal of Archaeological Science* 40, 4483–4490.
552 <https://doi.org/10.1016/j.jas.2013.07.010>
553
- 554 Carvalho, A.F., Alves-Cardoso, F., Gonçalves, D., Granja, R., Cardoso, J.L., Dean, R.M., Gibaja,
555 J.F., Masucci, M.A., Arroyo-Pardo, E., Fernández-Domínguez, E., Petchey, F., Douglas Price,
556 T., Mateus, J.E., Queiroz, P.F., Callapez, P., Pimenta, C., Regala, F.T., 2016. The Bom Santo
557 Cave (Lisbon, Portugal): Catchment, Diet, and Patterns of Mobility of a Middle Neolithic
558 Population. *European Journal of Archaeology* 19, 187–214.
559 <https://doi.org/10.1179/1461957115Y.0000000014>.
560
- 561 Correia, F., 2015. O Castro da Columbeira (Bombarral): A Exploração dos Recursos Faunísticos
562 no Calcolítico Estremenho (Masters Thesis). Universidade do Algarve.
563
- 564 Crowley, B.E., Miller, J.H., Bataille, C.P., 2017. Strontium isotopes ($^{87}\text{Sr}/^{86}\text{Sr}$) in terrestrial
565 ecological and palaeoecological research: empirical efforts and recent advances in continental-
566 scale models. *Biological Reviews* 92(1), 43-59.
567
- 568 Cunha P.P, dos Reis R.P., 1995. Cretaceous sedimentary and tectonic evolution of the northern
569 sector of the Lusitanian Basin (Portugal). *Cretaceous Research* 16, 155-170.
570
- 571 Davis, S., Mataloto, R., 2012. Animal Remains from Chalcolithic São Pedro (Redondo,
572 Alentejo): Evidence for a Crisis in the Mesolithic. *Revista Portuguesa de Arqueologia* 15, 47–85.
573
- 574 Díaz-del-Río, P., Waterman, A.J., Thomas, J.T., Peate, D.W., Tykot, R.H., Martínez-Navarrete,
575 M.I., Vicent, J.M., 2017. Diet and mobility patterns in the Late Prehistory of central Iberia
576 (4000–1400 cal BC): the evidence of radiogenic ($^{87}\text{Sr}/^{86}\text{Sr}$) and stable ($\delta^{18}\text{O}$, $\delta^{13}\text{C}$) isotope ratios.
577 *Archaeological and Anthropological Sciences* 9, 1439–1452. [https://doi.org/10.1007/s12520-](https://doi.org/10.1007/s12520-017-0480-y)
578 [017-0480-y](https://doi.org/10.1007/s12520-017-0480-y)
579

- 580 Diaz-Zorita-Bonilla, M. 2013. The Copper Age in South-West Spain: A Bioarchaeological
581 approach to prehistoric social organisation. PhD Thesis. Durham University
582
- 583 Díaz-Zorita Bonilla, M., 2013. Bioarqueología de las prácticas funerarias del yacimiento de la
584 Edad del Cobre de Valencina de la Concepción-Castilleja de Guzmán: revisión de las
585 investigaciones (State of the art of the bioarchaeology of the funerary practices at the
586 archaeological site of Valencina de la Concepción). In: García Sanjuán, L., Hurtado Pérez, V.,
587 Vargas Jiménez, J. M., Ruiz Moreno, T. and Cruz-Auñón Briones, R. (Eds): Valencina
588 Prehistórica. Actas del Congreso Conmemorativo del Descubrimiento de La Pastora (1860-
589 2010). Sevilla, Universidad de Sevilla.
- 590
- 591 Díaz-Zorita Bonilla, M., Beck, J., Bocherens, H., Díaz-del-Río, P., 2018. Isotopic evidence for
592 mobility at large-scale human aggregations in Copper Age Iberia: the mega-site of Marroquíes.
593 *Antiquity*, 92(364), 991-1007.
- 594
- 595 Driesch, von den, A., Boessneck, J., 1976. Die Fauna vom Castro do Zambujal (Fundmaterial der
596 Grabungen von 1966 bis 1973 mit Ausnahme der Zwingerfunde). In: Studien über frühe
597 Tierknochenfunde von der Iberischen Halbinsel 5. München, Madrid: Institut für
598 Palaeoanatomie, Domestikationsforschung und Geschichte der Tiermedizin der Universität
599 München. Deutsches Archäologisches Institut Abteilung Madrid.
- 600
- 601 Driesch, von den, A., Boessneck J., 1981. Die fauna von Zambujal. In: Sangmeister E, Schubart
602 H, (Eds). Zambujal, die grabungen 1964 bis 1973. Mainz am Rhein: P. Von Zabern. p302-314.
603
- 604 Ericson J.E., 1985. Strontium isotope characterization in the study of prehistoric human ecology.
605 *Journal of Human Evolution* 14(5), 503-514.
- 606
- 607 Faure G, Powell J.L., 1972. Strontium isotope geology. New York, N.Y: Springer-Verlag.
608
- 609 Freitas, M.C., Andrade, C., Rocha, F., Tassinari, C., Munha, J.M., Cruces, A., Vidinha, J., da
610 Silva, C.M., 2003. Lateglacial and Holocene environmental changes in Portuguese coastal

- 611 lagoons 1: the sedimentological and geochemical records of the Santo André coastal area.
612 Holocene 13, 433-446.
613
- 614 Gauß, R., 2013. The development of metallurgy on the Iberian Peninsula. Technological and
615 social patterns of a long-term innovation process. In: Burmeister, S., Hansen, S., Kunst, M.,
616 Müller-Scheeßel, N. (Eds.), Metal Matters: Innovative Technologies and Social Change in
617 Prehistory and Antiquity, Menschen-Kulturen-Traditionen, Studien aus den Forschungsclustern
618 des Deutschen Archäologischen Instituts 12, Rahden/Westf.:Marie Leidorf, p209-229.
619
- 620 Gauß, R., 2015. Zambujal und die Anfänge der Metallurgie in der Estremadura (Portugal).
621 Technologie der Kupfergewinnung, Herkunft des Metalls und soziokulturelle Bedeutung der
622 Innovation. Iberia Archaeologica 15, 1. Ernst Wasmuth Verlag, Tübingen, Berlin.
623
- 624 Gilli, A., Hodell, D.A., Kamenov, G.D., Brenner, M., 2009. Geological and archaeological
625 implications of strontium isotope analysis of exposed bedrock in the Chicxulub crater basin,
626 northwestern Yucatan, Mexico. Geology 37(8), 723-726.
627
- 628 Gonçalves, V.S., 2000/2001. O Trigo, a Cobre, a Lã e o Leite: Um Guia Bibliográfico e Uma
629 Curta Introdução Às Sociedades Camponesas Da Primeira Metade Do 3º Milénio No Centro e
630 Sul de Portugal. Zephyrus 53–54, 273–92.
631
- 632 Goncalves, V.S., Sousa, A.C., Costeira, C., 2013. Walls, gates and towers. Fortified settlements
633 in the south and centre of Portugal: some notes about violence and walls in the 3rd millenium
634 BCe. Cuadernos de Prehistoria y Arqueologia de la Universidad de Granada 23, 35-97.
635
- 636 Graustein, W.C., 1989. $^{87}\text{Sr}/^{86}\text{Sr}$ ratios measure the sources and flow of strontium in terrestrial
637 ecosystems. In: Rundel, P, Ehleringer, J.R., Nagy, K.A. (Eds.) Stable isotopes in ecological
638 research Springer, New York. p491-512.
639
- 640 Harrison, R. J. 1985., The “Policultivo Ganadero,” or the Secondary Products Revolution in
641 Spanish Agriculture, 5000-1000 BC. Proceedings of the Prehistoric Society 51, 75–102.

642

643 Jorge, S.O. 2000., Domesticating the Land: The First Agricultural Communities in Portugal.
644 *Journal of Iberian Archaeology* 2, 43–98.

645

646 Knudson, K.J., Stanish, C., Cerna, M.C.L., Faull, K.F., Tantaleán, H., 2016. Intra-individual
647 variability and strontium isotope measurements: A methodological study using $^{87}\text{Sr}/^{86}\text{Sr}$ data
648 from Pampa de los Gentiles, Chincha Valley, Peru. *Journal of Archaeological Science: Reports*
649 5, 590-597.

650

651 Kunst, M., 1995. Central Places and Social Complexity in the Iberian Copper Age. In: Lillios,
652 K.T. (Eds.) *The Origins of Complex Societies in Late Prehistoric Iberia*. International
653 *Monographs in Prehistory* Ann Arbor. p32-43.

654

655 Kunst, M., 2010. Zambujal, A dinâmica da sequência construtiva. In: Gonçalves, V.S., Sousa,
656 A.C. (Eds.) *Transformação e Mudança no Centro e Sul de Portugal: o 4.º e o 3.º milénios a.n.e.*,
657 *Actas do Colóquio Internacional (Cascais, 4-7 Outubro 2005)*, Coleção Cascais Tempos
658 Antigos 2, Cascais, 131-153.

659

660 Kunst, M., Morán, E., Parreira, R., 2013. Zambujal (Torres Vedras, Lisboa): relatório das
661 escavações de 2002. *Revista Portuguesa de Arqueologia* 16, 103-131.

662

663 Kunst, M., 2017a. Zambujal 2013, Teil 1. Eine kupferzeitliche befestigte Groß-Siedlung?
664 Vermessungsarbeiten und Prospektionen, mit einem Anhang von Anika Herb, *Madriider*
665 *Mitteilungen* 58, 1-30

666

667 Kunst, M., 2017b. Campaniforme em Zambujal (Torres Vedras). In: Gonçalves, V.S. (Ed.) *Sinos*
668 *e Taças junto ao Oceano e mais longe. Aspectos da presença campaniforme na Península Ibérica*,
669 *Bells and bowls near the ocean and far away. About beakers in the Iberian Peninsula*, *Estudos &*
670 *memórias* 10, UNIARQ, Lisboa, 194-213.

671

- 672 Kunst, M., Dambeck, R., Thiemeyer, H., 2016. Climatic and Environmental Changes on the
673 Iberian Peninsula: The Sizandro and Alcabrichel Project and some Considerations on
674 Neolithisation and the Beginning of the Bell-Beaker Phenomenon. In: Reindel, M., Bartl, K.,
675 Lüth, F., Benecke, N. (Eds.), Palaeoenvironment and the Development of Early Settlements.
676 Proceedings of the International Conferences Palaeoenvironment and the Development of Early
677 Societies (Şanlıurfa/Turkey, 5-7 October 2012), The Development of Early Settlement in Arid
678 Regions (Aqaba/Jordan, 12-15 November 2013), Menschen-Kulturen-Traditionen. Studien aus
679 den Forschungsclustern des Deutschen Archäologischen Instituts 14, Rahden/Westfalen: Marie
680 Leidorf, p51-62.
- 681
- 682 Kunst, M., Uerpamm, H.-P., 2002. Zambujal (Torres Vedras, Lisboa): relatório das escavações
683 de 1994 e 1995, Revista Portuguesa de Arqueologia 5 (1), 67-120.
- 684
- 685
- 686 Lillios, K.T., 1995. The Origins of Complex Societies in Late Prehistoric Iberia. International
687 Monographs in Prehistory, Ann Arbor.
- 688
- 689 Lillios, K.T., 1997. Amphibolite Tools of the Portuguese Copper Age (3000–2000 BC): A
690 Geoarchaeological Approach to Prehistoric Economics and Symbolism. Geoarchaeology 12(2),
691 137–163.
- 692
- 693 Madgwick, R., Lewis, J., Grimes, V., Guest, P., 2017. On the hoof: exploring the supply of
694 animals to the Roman legionary fortress at Caerleon using strontium ($^{87}\text{Sr}/^{86}\text{Sr}$) isotope analysis.
695 Archaeological and Anthropological Sciences 11(1), 223-225.
- 696
- 697 Minniti, C., Valenzuela-Lamas, S., Evans, J., Albarella, U., 2014. Widening the market.
698 Strontium isotope analysis on cattle teeth from Owslebury (Hampshire, UK) highlights changes
699 in livestock supply between the Iron Age and the Roman period. Journal of Archaeological
700 Science 42, 305–314. <https://doi.org/10.1016/j.jas.2013.10.008>
- 701
- 702 Moita, P., Santos J.F., Pereira M.F., 2009. Layered granitoids: Interaction between continental

703 crust recycling processes and mantle-derived magmatism. Examples from the Évora Massif
704 (Ossa Morena Zone, southwest Iberia, Portugal). *Lithos* 111, 125–141.

705

706 Moreno-García, M., Sousa, A.C., 2013. A exploração de recursos Final, faunísticos no Penedo
707 do Lexim (Mafra) durante no Neolítico Final. In: Gonçalves, V.S., Diniz, M., Sousa, A.C. (Eds.)
708 5° Congresso do Neolítico Peninsular. Cascais: Câmara Municipal de Cascais, p67–76.

709

710 Müller, R., Cardoso, J., 2008. The origin and use of copper at the Chalcolithic fortification of
711 Leceia, Portugal. *Madrider Mitteilungen* 49, 64–93.

712

713 Müller, R., Goldenberg, G., Bartelheim M., Kunst, M., Pernicka, E., 2007. Zambujal and the
714 beginnings of metallurgy in southern Portugal. In: La Niece, S., Hook, D., Craddock, P (Eds).
715 *Metals and mines studies in Archaeometallurgy*. London: Archetype. p15-26.

716 Nelson B.K., DeNiro, M.J., Schoeninger M.J., DePaolo D.J., Hare, P.E., 1986. Effects of
717 diagenesis on strontium, carbon, nitrogen, and oxygen concentration and isotopic composition in
718 bone. *Geochim Cosmochim* 50, 1941–1949.

719

720 Odriozola, C.P., Linares-Catela, J.A., Hurtado-Pérez. V., 2010. Variscite Source and Source
721 Analysis: Testing Assumptions at Pico Centeno (Encinasola, Spain). *Journal of Archaeological*
722 *Science* 37(12), 3146–3157. <https://doi.org/10.1016/j.jas.2010.07.016>.

723

724 Odriozola, C., Villalobos García, R., Boaventura, R., Sousa, A.C., Martínez-Blanes, J. M., 2013.
725 Las producciones de adorno personal en rocas verdes del SW peninsular: los casos de Leceia,
726 Moita da ladra y Penha Verde. *Estudios Arqueológicos de Oeiras* 20, 605-622.

727

728 Price, T.D., Burton, J.H., Bentley, R.A., 2002. The characterization of biologically available
729 strontium isotope ratios for the study of prehistoric migration. *Archaeometry* 44, 117–135.

730

731 Price, T.D., Burton, J., Cucina, A, Zabala, P., Frei, R., Tykot, R.H., Tiesler, V., 2012. Isotopic
732 studies of human skeletal remains from a Sixteenth to Seventeenth Century AD churchyard in
733 Campeche, Mexico: diet, place of origin, and age. *Current Anthropology* 53(4), 396-433.

- 734
- 735 Price, T.D., Burton, J.H., Fullagar, P. D., Wright, L. E., Buikstra, J. E., Tiesler, V., 2015.
736 Strontium isotopes and the study of human mobility among the Ancient Maya. In: Cucina, A
737 (Ed): Archaeology and bioarchaeology of population movement among the Prehispanic Maya.
738 Springer, Cham. p119-132.
- 739
- 740 Rehkämper, M., Wombacher, F., Aggarwal, J.K., 2004. Stable isotope analysis by. Multiple
741 Collector ICP-MS. In: de Groot, P.A. (Ed.), Handbook of Stable Isotope Analytical Techniques.
742 Elsevier, p692-725.
- 743
- 744 Roberts, B., 2008. Migration, Craft Expertise and Metallurgy: Archaeological Review from
745 Cambridge 23(2), 27–45.
- 746
- 747 Sangmeister, E., Schubart, H. 1981. Zambujal, Die Grabungen 1964 bis 1973. Madrider Beiträge
748 5,1. Mainz: Philipp von Zabern.
- 749
- 750 Saragoça, P., Maurer, A.-F., Šoberl, L., Lopes, M. da C., Alfenim, R., Leandro, I., Umbelino, C.,
751 Fernandes, T., Valente, M.J., Ribeiro, S., Santos, J.F., Janeiro, A.I., Barrocas, C.D., 2016. Stable
752 isotope and multi-analytical investigation of Monte da Cegonha: A Late Antiquity population in
753 southern Portugal. Journal of Archaeological Science: Reports 9, 728–742.
- 754
- 755 Sealy, J.C., van der Merwe, N.J., Sillen, A., Kruger, F.J., Krueger, H.W., 1991. $^{87}\text{Sr}/^{86}\text{Sr}$ as a
756 Dietary Indicator in Modern and Archaeological Bone. Journal of Archaeological Science 18,
757 399–416.
- 758
- 759 Schneider, S., Fürsich, F.T., Werner, W., 2009. Sr-isotope stratigraphy of the Upper Jurassic of
760 central Portugal (Lusitanian Basin) based on oyster shells. International Journal of Earth
761 Sciences. 98(8), 1949-1970.
- 762
- 763 Schroeder H.A., Nason A.P., Tipton I.H., 1972. Essential metals in man: strontium and barium.
764 Journal of Chronic Diseases 25, 491–517.

765

766 Schuhmacher, T.X., 2012. Die Elfenbeinobjekte des Chalkolithikums und der Frühen Bronzezeit
767 auf der Iberischen Halbinsel. Interdisziplinäre Studien zu Herkunft, Austausch, Verarbeitung und
768 sozialer Bedeutung von Elfenbein. *Iberia Archaeologica* 16(2). Darmstadt, Mainz: Philipp von
769 Zabern.

770

771 Schuhmacher, T.X., 2017. Ivory Exchange Networks in the Chalcolithic of the Western
772 Mediterranean. In: Bartelheim, M., Bueno Ramírez, P., Kunst, M. (Eds.), *Key Resources and*
773 *Socio-Cultural Developments in the Iberian Chalcolithic*, *RessourcenKulturen* 6, Tübingen,
774 p291-312.

775

776 Schuhmacher, T.X., Cardoso, J.L., Banerjee, A., 2009. Sourcing African ivory in Chalcolithic
777 Portugal. *Antiquity* 83: 983–997.

778

779 Tommasini S., Tesciona, I., Casalini, M., Braschi, E., Avanzinelli, R., Conticelli, S., 2018
780 Strontium Isotopes in Biological Material: A Key Tool for the Geographic Traceability of Foods
781 and Humans Beings. In: Gupta D., Walther C. (Eds.) *Behaviour of Strontium in Plants and the*
782 *Environment*. Springer, Cham. p145-166

783

784 Uerpmann, H.-P., 1995. Observações sobre a ecologia e economia do Castro do Zambujal. In:
785 Kunst, M. (Ed.), *Origens, Estruturas e Relações das Culturas Calcolíticas da Península Ibérica*,
786 *Actas das I Jornadas Arqueológicas de Torres Vedras 3-5 Abril 1987*, *Trabalhos de Arqueologia*
787 *7*, Lisboa: IPPA, p47-53.

788

789 Uerpmann, H-P., Uerpmann, M., 2003. Zambujal: Die Stein und Beinartefakte aus den
790 Grabungen 1964 bis 1973. *Madrider Beiträge Band 5, Zambujal Teil 4*. Mainz am Rhein: Philip
791 von Zabern.

792

793 Valente, M.J. 2013. Moinho de Valadares, Mourão. Estudo da fauna mamalógica das sondagens
794 1, 2 e 3 (Campanha de 1999). In: Valera, A.C. (Ed.), *As sociedades agropastoris na margem*
795 *esquerda do Guadiana (2 a metade do IV aos inícios do II milénio AC)*. EDIA/DRCALEN.

796 p353-360.

797

798 Valente, M.J., Carvalho, A.F., 2014. Zooarchaeology in the Neolithic and Chalcolithic of
799 Southern Portugal. *Environmental Archaeology* 19(3), 226-240.

800

801 Valera, A.C., Simão, I., Nunes, T., do Pereiro, T., Costa, C., 2017. Neolithic ditched enclosures
802 in Southern Portugal (4th millennium BC): New data and new perspectives. *Estudos do*
803 *Quaternário/Quaternary Studies* 17, 57-76.

804

805 Villaseca, C., Bellido, F., Pérez-Soba, C., Billström, K., 2009. Multiple crustal sources for post-
806 tectonic I type granites in the Hercynian Iberian Belt. *Mineralogy and Petrology* 96, 197–211.

807

808 Voerkelius, S., Lorenz Gesine, D., Rummel, S., Quételet Christophe, R., Heiss, G., Baxter, M.,
809 Brach-Papa, C., Deters-Itzelsberger, P., Hoelzl, S., Hoogewerff, J., Ponzevera, E., Van
810 Bocxstaele, M., Ueckermann, H., 2010. Strontium isotopic signatures of natural mineral waters,
811 the reference to a simple geological map and its potential for authentication of food. *Food*
812 *Chemistry* 118, 933–940. <https://doi.org/10.1016/j.foodchem.2009.04.125>.

813

814 Waight, T.E., Baker, J.A., Peate, D.W., 2002. Sr isotope ratio measurements by double focusing
815 MC-ICPMS: techniques, observations and pitfalls. *International Journal of Mass Spectrometry*
816 221, 229–244.

817

818 Waterman, A.J. 2012. *Marked in Life and Death: Identifying biological markers of social*
819 *differentiation in late prehistoric Portugal*. PhD Thesis. University of Iowa.

820

821 Waterman, A., Figueiredo, A., Thomas, J., Peate, D., 2013. Identifying migrants in the late
822 Neolithic burials of the antas of Rego da Murta (Alvaiazere, Portugal) using strontium isotopes.
823 *Antropo* 0, 190–196.

824

825 Waterman, A.J., Peate, D. W., Silva, A.M, Thomas, J.T., 2014. In *Search of Homelands: Using*
826 *Strontium Isotopes to Identify Biological Markers of Mobility in Late Prehistoric Portugal*.

827 Journal of Archaeological Science 42, 119–27. <https://doi.org/10.1016/j.jas.2013.11.004>.

828

829 Wilson, R.C.L., 1988. Mesozoic development of the Lusitanian Basin, Portugal. *Revista de la*
830 *Sociedad Geológica de España* 1, 393–407.

831

832 Žalaitė, I., Maurer, A.F., Grimes, V., Silva, A.M., Ribeiro, S., Santos, J.F., Barrocas Dias, C.,
833 Valera, A.C., 2018. Diet and mobility of fauna from Late Neolithic–Chalcolithic site of
834 Perdigões, Portugal. *Journal of Archaeological Science: Reports* 19, 674–685.
835 <https://doi.org/10.1016/j.jasrep.2018.03.033>

836

837 Zhao, C. 2015. Strontium isotope analysis and human mobility from late Neolithic to early
838 Bronze Age in the central plain of China. *CAA2015*, 783.

839

840 **List of Tables**

841 Table 1: $^{87}\text{Sr}/^{86}\text{Sr}$ results from all samples

842

843 Table 2: Results from enamel and dentine pairs

844

845 Table 3: Results of the Mann-Whitney U test for differences in $^{87}\text{Sr}/^{86}\text{Sr}$ values between sites

846

847 Table 4: $^{87}\text{Sr}/^{86}\text{Sr}$ generalized `local` values from other sites across central and south west Iberia

848

849 **List of Figures**

850 Figure 1: Location of the two sites

851

852 Figure 2: Geological map of the study area

853

854 Figure 3: Plot of $^{87}\text{Sr}/^{86}\text{Sr}$ ratios from cattle teeth at Zambujal and Leceia, alongside ratios taken from
855 locally collected rabbit and plants. The generalised local range for Zambujal is shown in green, and for
856 Leceia in blue.

857

858 Figure 4: Box-plot comparing $^{87}\text{Sr}/^{86}\text{Sr}$ enamel values between the two sites

859

860 Figure 5: Density plots comparing $^{87}\text{Sr}/^{86}\text{Sr}$ enamel values between the two sites

861

862 Figure 6: Plot showing $^{87}\text{Sr}/^{86}\text{Sr}$ values for enamel and dentine pairs

863

864 Figure 7: Sites listed in table with generalized $^{87}\text{Sr}/^{86}\text{Sr}$ ranges. 1. Leceia, 2. Zambujal, 3. Bom Santo, 4.

865 Perdigões, 5. La Pijotilla, 6. Rego da Murta I and II, 7. Valencina de la conception, 8. Madrid-region

866 sites. 9. Monte da Cegonha