



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/144187/>

Version: Accepted Version

---

**Article:**

Tierney, GJ, Gildea, K, Krosshaug, T et al. (2019) Analysis of ball carrier head motion during a rugby union tackle without direct head contact: A case study. *International Journal of Sports Science & Coaching*, 14 (2). pp. 190-196. ISSN: 1747-9541

<https://doi.org/10.1177/1747954119833477>

---

© 2019, The Author(s). This is an author produced version of a paper published in *International Journal of Sports Science & Coaching*. Reprinted by permission of SAGE Publications.

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



# International Journal of Sports Science & Coaching

## Analysis of ball carrier head motion during a rugby union tackle without direct head contact: a case study

Journal:	<i>International Journal of Sports Science &amp; Coaching</i>
Manuscript ID	SPO-18-0181.R2
Manuscript Type:	Original Article
Date Submitted by the Author:	31-Oct-2018
Complete List of Authors:	Tierney, Gregory; University of Dublin Trinity College, Gildea, Kevin; University of Dublin Trinity College Krosshaug, Tron; Oslo Sports Trauma Research Centre, The Norwegian School of Sport Sciences, Oslo, Norway Simms, Ciaran; University of Dublin Trinity College, Centre for Bioengineering
Keywords:	rugby union, concussion, tackle, motion analysis, biomechanics
Abstract:	Rugby union players can be involved in many tackles per game. However, little is known of the regular head loading environment associated with tackling in rugby union. In particular, the magnitude and influencing factors for head kinematics during the tackle are poorly understood. Accordingly, the goal of this study was to measure head motion of a visually unaware ball carrier during a real game tackle to the upper trunk with no direct head contact, and compare the kinematics with previously reported concussive events. Model-Based Image-Matching was utilised to measure ball carrier head linear and angular velocities. Ball carrier componential maximum change in head angular velocities of 38.1, 20.6 and 13.5 rad/s were measured for the head local X (coronal plane), Y (sagittal plane) and Z (transverse plane) axes respectively. The combination of a high legal tackle height configuration and visually unaware ball carrier can lead to kinematics similar to average values previously reported for concussive direct head impacts.

SCHOLARONE™  
Manuscripts

1  
2  
3     **Analysis of ball carrier head motion during a rugby union tackle**  
4     **without direct head contact: a case study**  
5  
6

7     **Gregory J. Tierney<sup>[1]</sup>**  
8  
9     gtierne@tcd.ie  
10  
11     **Kevin Gildea<sup>[1]</sup>**  
12     kgildea@tcd.ie  
13  
14     **Tron Krosshaug<sup>[2]</sup>**  
15     tron.krosshaug@nih.no  
16  
17     **Ciaran K. Simms<sup>[1]</sup>**  
18     csimms@tcd.ie  
19  
20  
21  
22     <sup>[1]</sup> Trinity Centre for Bioengineering, Trinity College Dublin, Ireland  
23  
24     <sup>[2]</sup> Oslo Sports Trauma Research Centre, The Norwegian School of Sport Sciences, Oslo, Norway  
25  
26  
27     **Short Title:** High ball carrier head kinematics in a rugby union tackle  
28  
29  
30     **Corresponding Author:**  
31     Gregory Tierney  
32     Trinity Centre for Bioengineering  
33     Trinity College Dublin  
34     Dublin 2  
35     Ireland  
36     gtierne@tcd.ie  
37     +353 89 2381688  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Abstract**

Rugby union players can be involved in many tackles per game. However, little is known of the regular head loading environment associated with tackling in rugby union. In particular, the magnitude and influencing factors for head kinematics during the tackle are poorly understood. Accordingly, the goal of this study was to measure head motion of a visually unaware ball carrier during a real game tackle to the upper trunk with no direct head contact, and compare the kinematics with previously reported concussive events. Model-Based Image-Matching was utilised to measure ball carrier head linear and angular velocities. Ball carrier componential maximum change in head angular velocities of 38.1, 20.6 and 13.5 rad/s were measured for the head local X (coronal plane), Y (sagittal plane) and Z (transverse plane) axes respectively. The combination of a high legal tackle height configuration and visually unaware ball carrier can lead to kinematics similar to average values previously reported for concussive direct head impacts.

## Introduction

Head injuries remain a considerable concern in rugby union. An emerging field in brain injury research is the study of repeated sub-concussive head impacts,<sup>1-3</sup> defined as “head impacts that do not result in symptoms typically used to define concussion”.<sup>4</sup> The definition of sub-concussive impacts presents challenges as a lower threshold for sub-concussive impacts has not been established. However, for practical purposes, events that result in less than 10g head acceleration have generally not been considered in rugby union head kinematic studies.<sup>5 6</sup> In sports such as boxing and soccer, sub-concussive head impacts have been linked with acute changes in brain function,<sup>2</sup> structural changes in white matter,<sup>1</sup> biomarkers of neuronal injury,<sup>3</sup> and short term cognitive impairments.<sup>7</sup> However, a recent systematic review on sub-concussion found the term to be inconsistently used, poorly defined and misleading and this must be considered when reviewing the abovementioned studies.<sup>8</sup>

Head linear and angular kinematics are linked to brain injury.<sup>9-13</sup> Studies generally focus on the magnitude of a single hit.<sup>9-13</sup> For example, McIntosh et al.<sup>9</sup> found that concussive direct head impacts cause on average 103g of peak resultant head linear acceleration and a change in head angular velocity of 33 rad/s. However, there is an emerging concept of neuronal vulnerability to injury due to repetitive sub-concussive loading if the time between hits is sufficiently small (up to 24 hours).<sup>4 14 15</sup> Merchant-Borna et al.<sup>4</sup> argues that injury thresholds should not be based on a single hit but that the number and magnitude of hits and the time between hits should also be considered. In extreme cases, rugby union players can be involved in over 30 tackles per game.<sup>16</sup> However, little is known of the regular head loading environment associated with rugby union. In particular, the magnitude and influencing factors for head kinematics during regular rugby union play without any direct head contact are poorly understood, even though a recent video review found that 1 in 7 head injury assessments in rugby union could not be associated with a specific collision event.<sup>17</sup>

During one season of an amateur rugby union team, King et al.<sup>5</sup> recorded 181 impacts (0.9% of total impacts) over 95g (head linear acceleration concussion injury threshold utilised for comparison by King et al.<sup>5</sup>) and 4452 impacts (21.5% of total impacts) over 5500 rad/s<sup>2</sup> (head rotational acceleration concussion injury threshold utilised for comparison by King et al.<sup>5</sup>). However, no diagnosed concussive head impacts were included in this dataset. King et al.<sup>5</sup> reported that inertial head loading (no direct head contact) most likely accounted for a

high proportion of these large head kinematic values recorded. However, no further assessment was provided. In rugby union, the ball carrier can be visually unaware of an approaching tackler, i.e. not anticipating the tackle,<sup>18</sup> and failing to brace may result in a higher susceptibility to injury<sup>18</sup> and could lead to higher inertial head loading. Direct measurement of head kinematics during tackling with on-field measurement devices remains challenging.<sup>5</sup> An alternative approach is to use Model-Based Image-Matching (MBIM).<sup>19 20</sup> This approach uses multiple camera views of an impact and a computerised skeletal model to extract six degree-of-freedom head kinematics directly from video. Accordingly, the goal of this exploratory study is to use MBIM to measure head kinematics of a visually unaware ball carrier during a real game shoulder tackle<sup>21</sup> to the upper trunk and to compare to average concussion kinematics values reported in the literature for direct head impacts.<sup>9</sup> The approach provides a case specific example to the body of evidence on the regular head loading environment in rugby union.

## Methods

Similar to a previous video analysis study on knee injuries in rugby union,<sup>22</sup> video search engines (e.g. YouTube) were utilised to identify suitable clips. The three selection criteria utilised were 1) contact was to the upper trunk (Figure 1) of the ball carrier;<sup>23</sup> 2) The ball carrier was visually unaware of the tackle (tackler approaching outside the ball carrier's peripheral vision);<sup>24 25</sup> 3) there was a minimum of three synchronisable camera views of the tackle event available for MBIM reconstruction.<sup>19</sup> Although Tierney et al.<sup>24 25</sup> found that the ball carrier is visually unaware during roughly one-third of side-on tackles, only one tackle event satisfied all criteria (due to criterion #3). In this single tackle event, the ball carrier had just passed the ball and was impacted around the left scapula by the tackler (Figure 2). The player received on-field medical attention but was not immediately removed from play (the player was substituted roughly 45 minutes after the event) and the subsequent medical history is unknown. The tackle was reviewed by the on-field referee/video referees and was deemed legal play as the tackler had committed to the tackle before the ball was passed. The video data was freely available online and no medical data was obtained/reported in this study, so ethical permission was not required, similar to other rugby union video analysis studies.<sup>22 26 27</sup>





Figure 1. The body regions of the ball carrier.

MBIM has been previously described in detail by Krosshaug and Bahr.<sup>28</sup> Briefly, this method uses a multibody skeleton model to estimate human body joint angle time-histories from multiple camera views of human movement. For every video frame, the skeleton model joint angles are manually adjusted to match the segment position and orientation of the model with respect to the target athlete in the multiple camera views. The MBIM technique has been validated previously for six degree of freedom tracking of the pelvis,<sup>28</sup> hip,<sup>28</sup> knee,<sup>28</sup> ankle<sup>29</sup> and head.<sup>19</sup> The MBIM analysis has been shown to be repeatable by both a single researcher and multiple researchers for six degree of freedom head motion data (Intra-class Correlation Coefficients (ICC) greater than 0.9 for six degree of freedom head displacement measurement).<sup>19</sup> The method has previously shown Root Mean Square (RMS) errors of less than 20 mm for linear displacement and less than 0.04 rad for rotational displacement for reconstructing head motion in a vehicle cadaver head-windscreen impact.<sup>19</sup> However, RMS errors up to 5.38 rad/s were reported for the MBIM method for measuring angular velocity

1  
2  
3 during direct head impacts.<sup>19</sup> The MBIM method is considered suitable for measuring componential angular  
4 velocity during indirect head impacts (i.e. inertial head kinematics as a result of an impact to the body) for  
5 which lower frequency head motion is typically associated with.<sup>30</sup>  
6  
7


8  
9 In this case, matching was conducted on synchronised video of three camera views of the tackle. Each video  
10 had a resolution of 720p and frame rate of 25 fps. One researcher performed the matching using 3-D  
11 animation software Poser 4 and Poser Pro Pack (Curious Labs Inc, Santa Cruz, California). The surroundings  
12 were built in a virtual environment based on the dimensions of the rugby field. The videos were imported into  
13 the Poser workspace background and the surroundings were then matched to the background video footage  
14 for every camera view. As the camera locations were unknown, this was achieved by manually adjusting the  
15 camera positioning tool which contains three translational and three rotational degrees of freedom, as well as  
16 variable focal length. Since the cameras were moving during the impact, the environment matching was  
17 conducted for each individual video frame. A skeleton model from Zygote Media Group Inc (Provo, Utah, US)  
18 was manipulated to fit the player's head for each video frame (Figure 2). Linear closing speed estimates were  
19 also calculated by tracking the players' pelvises using the MBIM approach utilised by Krosshaug and Bahr.<sup>28</sup>  
20  
21




53  
54 Figure 2. (a) A time lapse of the upper trunk active shoulder tackle with the MBIM matching for one camera  
55 view and (b) the MBIM matching for three camera views at time t=80ms.  
56  
57

58 This approach yielded head successive rotation angles of order yaw-pitch-roll (or Z-Y-X see Figure 3<sup>19</sup>) and  
59 linear position measurements every 40ms. The time derivatives of the yaw, pitch and roll angles were  
60

1  
2  
3 calculated using the Matlab gradient function and hence the components of the body local head angular  
4 velocity (Figure 3) every 40ms were calculated.<sup>19</sup> The same method was used to calculate the head and pelvis  
5 linear velocity components in the global coordinate system (Figure 4). The maximum change in head  
6 componential angular velocity values were compared to the average concussion values reported in the  
7 literature for unhelmeted sports.<sup>9</sup> The comparison was not conducted for the maximum change in head linear  
8 velocity results, as componential data is not available for unhelmeted sports.<sup>31</sup> **The MBIM analysis for this case**  
9 took roughly 60 hours to complete.  
10  
11  
12  
13  
14  
15  
16




35 Figure 3. The local axes of the head.  
36  
37




1  
2  
3 Figure 4. The global coordinate system utilised for the MBIM method.  
4  
5  
6

## Results

9 The componential change in head linear velocity is mostly in the global Y direction (lateral for the player)  
10 (Figure 5). The componential head angular kinematics from this case are similar to the reported average values  
11 for concussive direct head impacts in unhelmeted sports such as rugby and Australian rules football <sup>9</sup> (Figure  
12 6). For the X (coronal plane) and Y (sagittal plane) components, the maximum change in head angular velocity  
13 is greater than in average concussive cases. The linear and angular velocity values for each time frame can be  
14 seen in Appendix A. Although it was a side-on tackle,<sup>24 25 32</sup> the resultant tackler closing speed was 5.5 m/s (-2.5  
15 m/s, 4.9 m/s and -0.6 m/s in the global X, Y and Z direction, respectively). The resultant ball carrier closing  
16 speed was 3.1 m/s (-0.6 m/s, -3.0 m/s and -0.3 m/s in the global X, Y and Z direction, respectively).  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32



1  
2  
3 Figure 5. The componential change in head linear velocity results for the ball carrier about the global  
4 coordinate system (Figure 4).  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14



37  
38 Figure 6. The maximum change in head angular velocity results about the head local coordinate system (Figure  
39 3) from inertial head loading in this study compared to the corresponding average concussion values from  
40 direct head impacts reported in the literature.<sup>9</sup>  
41  
42

#### 43 Discussion

44  
45 This case study used the MBIM method to measure head kinematics of a visually unaware ball carrier during a  
46 rugby union shoulder tackle to the upper trunk. Componential head angular velocities were similar to the  
47 average values reported for concussive direct head impacts in unhelmeted sports (Figure 6).<sup>9</sup> Although the  
48 long term medical outcome of this case is unknown, these results support the finding that legal shoulder  
49 tackles to the upper trunk where the ball carrier is visually unaware is a concern for inertial head loading.<sup>30</sup> A  
50 conclusion regarding injury risk associated with these tackles requires correlation with injury data and this  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 should be a focus of future work. Longitudinal studies considering blood biomarkers, medical imaging,  
4  
5 concussion history, medical reports, injury data and overall head impact exposure would be of benefit.  
6  
7

8 In this case study, the ball carrier was impacted by the tackler just after passing the ball. The tackle was  
9 subsequently reviewed by the on-field referee/video referees and regarded as legal play as the tackler was  
10 committed to the tackle before the ball was passed. It could be considered difficult for a ball carrier to protect  
11 themselves/brace when impacted from behind and without the ball in their hands. Further work should look at  
12 these types of tackles to examine their incidence as well as propensity for injury and high head kinematics. **This**  
13  
14 **will build an evidence base for coaches to develop pre-contact tackle strategies such as ensuring ball carriers**  
15  
16 **pass to end off in a protective position.**  
17  
18

19 Recent studies have proposed a biomechanical justification for tackling lower around the mid/lower trunk  
20 (Figure 1) in rugby union, for which this case study provides further support.<sup>30 33</sup> Upper body tackles,<sup>26 27</sup> especially when primary contact is with the ball carrier's upper trunk,<sup>23</sup> are the main cause of direct head  
21 impacts for the tackler. A recent study identified that tackling at the upper trunk has a high propensity to  
22 result in tackler direct head impacts and emphasised the importance of tackling lower risk body regions such  
23 as the mid and lower trunk.<sup>34</sup> For the ball carrier, studies using multibody models and staged tackles in a  
24 motion analysis laboratory have found that higher tackle heights, particularly to the upper trunk but below the  
25 legal limit (line of the shoulders), can result in significantly higher inertial head kinematics in comparison to  
26 tackles below the upper trunk.<sup>30 33</sup> Thus, if contact in this case had been made below the upper trunk of the  
27 ball, the ball carrier's inertial head kinematics would likely have been reduced, potentially by over 50%.<sup>33</sup> The  
28 energy transmitted during an impact is attenuated along a damped/deformable linkage system through  
29 viscous dissipation.<sup>35</sup> Thus, the head kinematics resulting from an impact to the body are inversely related to  
30 the distance of the impact from the head. The overall ball carrier angular momentum about the point of  
31 contact is conserved in the tackle. This results in a lower rotational inertia above the point of contact when the  
32 tackle height is greater and hence greater head rotations. **This highlights the importance for coaches to**  
33  
34 **encourage tackling at lower risk body regions such as the mid- and lower trunk.**  
35  
36

#### 37 Limitations

38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Root mean square errors up to 5.38 rad/s and 1.29m/s were measured for componential angular and linear  
4 velocity reconstruction in the MBIM validation study, respectively.<sup>19</sup> This should be considered when  
5 interpreting the current results. Linear and angular acceleration were not measured using the MBIM method  
6 as the sample frequency (video frame rate) was too low (25 Hz) and head acceleration measures typically  
7 require 1000 Hz sampling frequency.<sup>5</sup> The frame rate of 25 fps could be considered low for inertial head  
8 angular velocity measurement. It is possible that higher frequency head motion was unmeasured. The time  
9 duration associated with the head kinematics measured in this study (Appendix A) are much longer than those  
10 typically associated with concussive direct head impacts (peak values usually measured within 54 ms).<sup>36</sup> Only  
11 one case was analysed in this current study. Much larger sample sizes were utilised by McIntosh et al.<sup>9</sup> and  
12 this should be considered when interpreting the kinematic result comparisons. The study would have  
13 benefited from access to immediate and follow up medical data, if applicable, from this case.  
14  
15

16 The MBIM approach is currently a time-consuming process as it requires manual frame-by-frame matching.  
17 This case took approximately 60 hours to complete. Further work should look at automating/semi-automating  
18 the MBIM technique. The sample size for this study was only one due to the selection criteria required for the  
19 MBIM technique. Ideally, for comparison, the MBIM method could have been conducted on a tackle with  
20 similar closing speeds that occurred lower down on the body of an unaware ball carrier.<sup>37</sup> **The closing speeds in  
21 this case study were similar to the average closing speeds measured in elite level rugby union.<sup>38</sup>** Access to  
22 multiple camera view synchronised video footage directly from the sports broadcaster could have allowed  
23 more cases to be analysed.  
24  
25

## 43 Conclusion

44

45 Model-Based Image-Matching was utilised to measure maximum changes in head linear and angular velocities  
46 of a visually unaware ball carrier during a real game tackle to the upper trunk. Even though no direct  
47 head/neck contact occurred in this case, the head angular kinematics were similar to the average values  
48 previously reported for concussive direct head impacts in unhelmeted sports. The combination of a high legal  
49 tackle height configuration and a visually unaware ball carrier can lead to high inertial head kinematics.  
50 Previous work indicates that a lower tackle height would reduce this. Further biomechanical and clinical  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 collaborative research is required to conclude on the long-term effects of tackles generating high head  
4  
5 rotational velocity changes.  
6  
7

## 8 Declaration of Conflicting Interests 9

10  
11 The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or  
12  
13 publication of this article.  
14  
15

## 16 Funding 17

18  
19 The author(s) received funding from the Irish Research Council for this article.  
20  
21  
22  
23  
24  
25  
26

## 27 References 28

- 30 1. Bazarian JJ, Zhu T, Blyth B, et al. Subject-specific changes in brain white matter on diffusion tensor imaging  
31  
32 after sports-related concussion. *Magn Reson Imaging* 2012;30(2):171-80.  
33  
34 2. Breedlove EL, Robinson M, Talavage TM, et al. Biomechanical correlates of symptomatic and asymptomatic  
35  
36 neurophysiological impairment in high school football. *J Biomech* 2012;45(7):1265-72.  
37  
38 3. Neselius S, Brisby H, Theodorsson A, et al. CSF-biomarkers in Olympic boxing: diagnosis and effects of  
39  
40 repetitive head trauma. *PLoS one* 2012;7(4):e33606.  
41  
42 4. Merchant-Borna K, Asselin P, Narayan D, et al. Novel method of weighting cumulative helmet impacts  
43  
44 improves correlation with brain white matter changes after one football season of sub-concussive  
45  
46 head blows. *Ann Biomed Eng* 2016;44(12):3679-92.  
47  
48 5. King D, Hume PA, Brughelli M, et al. Instrumented mouthguard acceleration analyses for head impacts in  
49  
50 amateur rugby union players over a season of matches. *Am J Sports Med* 2015;43(3):614-24.  
51  
52 6. King DA, Hume PA, Gissane C, et al. Similar head impact acceleration measured using instrumented ear  
53  
54 patches in a junior rugby union team during matches in comparison with other sports. *J Neurosurg  
55  
56 Pediatr* 2016;18(1):65-72.  
57  
58  
59  
60

- 1  
2  
3 7. McAllister T, Flashman L, Maerlender A, et al. Cognitive effects of one season of head impacts in a cohort of  
4 collegiate contact sport athletes. *Neurology* 2012;78(22):1777-84.  
5  
6 8. Mainwaring L, Pennock K, Mylabathula S, et al. Subconcussive head impacts in sport: a systematic review of  
7 the evidence. *Int J Psychophysiol* 2018. doi: 10.1016/j.ijpsycho.2018.01.007  
8  
9 9. McIntosh AS, Patton DA, Fréchède B, et al. The biomechanics of concussion in unhelmeted football players  
10 in Australia: a case-control study. *BMJ Open* 2014;4(5) doi: 10.1136/bmjopen-2014-005078  
11  
12 10. Frechede B, McIntosh AS. Numerical reconstruction of real-life concussive football impacts. *Med Sci Sports  
13 Exerc* 2009;41(2):390-8  
14  
15 11. Takhounts EG, Craig MJ, Moorhouse K, et al. Development of brain injury criteria (BrIC). *Stapp Car Crash J*  
16  
17 2013;57:243-66.  
18  
19 12. Takhounts EG, Hasija V, Ridella S, et al. Kinematic rotational brain injury criterion (BRIC). In: Proceedings of  
20 the 22nd Enhanced Safety of Vehicles Conference, 2011.  
21  
22 13. Newman J, Barr C, Beusenberg MC, et al. A new biomechanical assessment of mild traumatic brain injury.  
23  
24 Part 2: Results and conclusions. In: *IRCOBI Conference Proceedings*, 2000.  
25  
26 14. Effgen G, Gill E, Morrison B. A model of repetitive, mild traumatic brain injury and a novel pharmacological  
27 intervention to block repetitive injury synergy. In: *IRCOBI Conference Proceedings*, 2012.  
28  
29 15. Slemmer JE, Weber JT. The extent of damage following repeated injury to cultured hippocampal cells is  
30 dependent on the severity of insult and inter-injury interval. *Neurobiol Dis* 2005;18(3):421-31.  
31  
32 16. Deutsch M, Kearney G, Rehrer N. Time-motion analysis of professional rugby union players during match-  
33 play. *J Sports Sci* 2007;25(4):461-72.  
34  
35 17. Tucker R, Raftery M, Fuller GW, et al. A video analysis of head injuries satisfying the criteria for a head  
36 injury assessment in professional Rugby Union: a prospective cohort study. *Br J Sports Med* 2017;  
37 51(15):1147-1151.  
38  
39 18. Hendricks S, O'Connor S, Lambert M, et al. Video analysis of concussion injury mechanism in under-18  
40 rugby. *BMJ Open Sport Exerc Med* 2016;2(1):e000053.  
41  
42 19. Tierney GJ, Joodaki H, Krosshaug T, et al. Assessment of model-based image-matching for future  
43 reconstruction of unhelmeted sport head impact kinematics. *Sports Biomech* 2018;17(1):33-47.  
44  
45 20. Tierney GJ, Krosshaug T, Wilson F, et al. An Assessment of a Novel Approach for Determining the Player  
46 Kinematics in Elite Rugby Union Players. In: *IRCOBI Conference Proceedings*, 2015.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 21. Fuller CW, Ashton T, Brooks JH, et al. Injury risks associated with tackling in rugby union. *Br J Sports Med*  
4 2010;44(3):159-67.  
5  
6 22. Montgomery C, Blackburn J, Withers D, et al. Mechanisms of ACL injury in professional rugby union: a  
7 systematic video analysis of 36 cases. *Br J Sports Med* 2016; doi:10.1136/bjsports-2016-096425.  
8  
9 23. Tierney GJ, Simms CK. The effect of intended primary contact location on tackler head impact risk. *In:*  
10 *IRCOBI Conference Proceedings*, 2017.  
11  
12 24. Tierney GJ, Denvir K, Farrell G, et al. The effect of technique on tackle gainline success outcomes in elite  
13 level rugby union. *Int J Sports Sci Coach* 2018;13(1):16-25.  
14  
15 25. Tierney GJ, Denvir K, Farrell G, et al. Does player time-in-game affect tackle technique in elite level rugby  
16 union? *J Sci Med Sport* 2018;21(2):221-25.  
17  
18 26. Tierney GJ, Denvir K, Farrell G, et al. The Effect of Tackler Technique on Head Injury Assessment Risk in Elite  
19 Rugby Union. *Med Sci Sports Exerc* 2018;50(3):603-08.  
20  
21 27. Tierney GJ, Lawler J, Denvir K, et al. Risks associated with significant head impact events in elite rugby  
22 union. *Brain Inj* 2016;30(11):1350-61.  
23  
24 28. Krosshaug T, Bahr R. A model-based image-matching technique for three-dimensional reconstruction of  
25 human motion from uncalibrated video sequences. *Journal Biomech* 2005;38(4):919-29.  
26  
27 29. Mok K-M, Fong DT-P, Krosshaug T, et al. An ankle joint model-based image-matching motion analysis  
28 technique. *Gait & posture* 2011;34(1):71-75.  
29  
30 30. Tierney GJ, Simms CK. The effects of tackle height on inertial loading of the head and neck in Rugby Union:  
31 A multibody model analysis. *Brain Inj* 2017;31(13-14):1925-31.  
32  
33 31. McIntosh AS, McCrory P, Comerford J. The dynamics of concussive head impacts in rugby and Australian  
34 rules football. *Medicine and science in sports and exercise* 2000;32(12):1980-4.  
35  
36 32. Tierney GJ, Simms CK. Can tackle height influence tackle gainline success outcomes in elite level rugby  
37 union? *Int J Sports Sci Coach* 2018;13(3):415-20.  
38  
39 33. Tierney GJ, Richter C, Denvir K, et al. Could lowering the tackle height in rugby union reduce ball carrier  
40 inertial head kinematics? *J Biomech* 2018; doi: 10.1016/j.jbiomech.2018.02.023.  
41  
42 34. Tierney GJ, Simms CK. Can tackle height influence head injury assessment risk in elite rugby union? *J Sci  
43 Med Sport* 2018; doi: 10.1016/j.jsams.2018.05.010.  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 35. Kim W, Voloshin AS, Johnson SH. Modeling of heel strike transients during running. *Human Movement*  
4  
5 *Science* 1994;13(2):221-44.  
6

7  
8 36. Frechede B, McIntosh A. Use of MADYMO's human facet model to evaluate the risk of head injury in  
9 impact. In: Proceedings of the 21st ESV conference, 2007.  
10

11  
12 37. Tierney GJ, Denvir K, Farrell G, et al. Does ball carrier technique influence tackler head injury assessment  
13 risk in elite rugby union? *J Sports Sci* 2018; doi: 10.1080/02640414.2018.1494952.  
14

15  
16 38. Hendricks S, Karpul D, Nicolls F, et al. Velocity and acceleration before contact in the tackle during rugby  
17 union matches. *J Sports Sci* 2012;30(12):1215-1224.  
18

19  
20  
21 **Appendix A.** The ball carrier head linear and angular velocity values for each time frame  
22

Time (ms)	Linear velocity (m/s)			Angular velocity (rad/s)		
	X	Y	Z	X	Y	Z
0	-0.9	-3.7	0.1	0.2	0.1	-0.4
40	-0.9	-3.5	0.1	0.3	0.2	-1.1
80	-0.9	-0.5	0.3	-13.3	-8.5	3.9
120	-0.8	3.4	0.2	-10.0	-9.7	-9.6
160	-0.1	2.9	-0.9	14.2	4.7	2.7
200	-1.2	3.4	-1.4	24.7	10.9	-7.6
240	-2.9	5.6	-1.4	16.0	5.1	-5.1