



UNIVERSITY OF LEEDS

This is a repository copy of *The impact of New Labour's English health inequalities strategy on geographical inequalities in infant mortality: a time-trend analysis*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/144239/>

Version: Accepted Version

Article:

Robinson, T, Brown, H, Norman, PD orcid.org/0000-0002-6211-1625 et al. (3 more authors) (2019) The impact of New Labour's English health inequalities strategy on geographical inequalities in infant mortality: a time-trend analysis. *Journal of Epidemiology and Community Health*, 73 (6). pp. 564-568. ISSN 0143-005X

<https://doi.org/10.1136/jech-2018-211679>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

The impact of New Labour’s English health inequalities strategy on geographical inequalities in infant mortality: a time trend analysis

Corresponding author:

Professor Clare Bamba

Professor of Public Health

Institute of Health & Society

Newcastle University

Baddiley-Clark Building

Richardson Road

Newcastle upon Tyne

NE2 4AX

clare.bambra@ncl.ac.uk

Co-authors:

Dr Tomos Robinson, Institute of Health & Society, Newcastle University, Newcastle, UK

Dr Heather Brown, Institute of Health & Society, Newcastle University, Newcastle, UK

Dr Paul Norman, School of Geography, University of Leeds, Leeds, UK

Dr Lorna Fraser, Department of Health Sciences, University of York, York, UK

Dr Ben Barr, Institute of Psychology, Health & Society, University of Liverpool, Liverpool, UK

Keywords

Infant Mortality, Health Inequalities, Epidemiology, Health Policy, Geography

Word Count

3322

Abstract

Background: The English health inequalities strategy (1999-2010) aimed to reduce health inequalities between the most deprived local authorities and the rest of England. The multifaceted strategy included increased investment in health care, the early years, education and neighbourhood renewal. The objective of this study was to investigate whether the strategy was associated with a reduction in geographic inequalities in the infant mortality rate (IMR).

Methods: We used segmented regression analysis to measure inequalities in the IMR between the most deprived local authorities and the rest of England before, during and after the health inequalities strategy period.

Results: Before the strategy was implemented (1983-1999), absolute inequalities in the IMR increased between the most deprived local authorities and the rest of England at a rate of 0.033 annually (95% CI 0.003 to 0.063). During the strategy period (1999-2010), absolute inequalities decreased at a rate of -0.115 annually (95% CI -0.168 to -0.058). After the strategy period ended (2011-2017), absolute inequalities increased at a rate of 0.034 annually (95% CI -0.114 to 0.103). Relative inequalities also decreased during the strategy period.

Conclusion: The English health inequalities strategy was associated with a decline in absolute and relative geographical inequalities in the IMR. This research adds to the evidence base showing that the English health inequalities strategy was at least partially effective in reducing health inequalities, and that current austerity policies may undermine these gains.

Text for Summary Box

What is already known on this subject?

Introduced between 1999-2010, the English health inequalities strategy was one of the most ambitious government strategies of its kind, resulting in large increases in levels of public spending on a range of social programmes including a substantial increase in expenditure on the National Health Service. Since it came to an end, a number of studies have attempted to evaluate whether the English health inequalities strategy was a success or not. The results thus far are mixed.

What does this study add?

In this study, we add to the empirical literature by analysing the association between the health inequalities strategy and both absolute and relative inequalities in the IMR, using high quality subnational data from 323 local authorities and segmented regression analysis. Our results show that both absolute and relative inequalities reduced during the strategy period.

Introduction

The infant mortality rate (IMR), defined as the number of deaths under the age of 1 per 1,000 live births, can be seen as a key indicator of the environment in which babies are born [1] and, more broadly, can serve as a proxy for the health of a population [2]. After years of steady improvement, the English IMR has begun to increase in recent years [3]. England, like many other countries, also has substantial inequalities in IMRs according to geographical area and socioeconomic group [4]. For example, national figures from 2016 published by the Office for National Statistics (ONS) have shown that the IMR was 5.9 per 1,000 live births in the most deprived areas and 2.6 in the least deprived areas [3].

Following the election of the New Labour government in 1997, there was a clear attempt to use government policy levers to reduce health inequalities. Consequently, the UK became the first European country in which policy-makers systematically and explicitly attempted to reduce inequalities in health, focusing specifically on: supporting families, engaging communities in tackling deprivation, improving prevention, increasing access to health care and tackling the underlying social determinants of health [5].

This comprehensive, multi-faceted programme to reduce health inequalities in England was one of the most ambitious strategies of its kind. For example, the policy included large increases in levels of public spending on a range of social programmes, the introduction of the national minimum wage, area-based interventions such as the Health Action Zones, and a substantial increase in expenditure on the NHS. The latter was targeted at more deprived neighbourhoods when, after 2001, a “health inequalities weighting” was added to the way in which NHS funds were geographically distributed, so that areas of higher deprivation received more funds per head to reflect higher health need [6].

Furthermore, the then government made tackling health, social and educational inequalities a public service priority by setting public service agreement (PSA) targets. The key targets of the Labour government’s health inequalities strategy were to: 1) reduce the relative gap in life expectancy at birth between the most deprived local authorities and the English average by 10% by 2010; and 2) cut relative inequalities by occupational class in the IMR by 10%. However, with a change in government and the establishment of the Conservative-Liberal Democrat coalition in 2010, this strategy came to an end. The change of government also led to an introduction of austerity measures intended to reduce the national deficit. These measures were realised through a structured and significant reduction in levels of public spending, including substantial cuts to funding for local authorities, real term reductions to the NHS budget, cuts to the education sector and various reductions across the welfare system [7]. The scale of the cut is typified by the closure of as many as 1,000 Sure Start children’s centres since 2010 [8].

A number of studies have subsequently attempted to evaluate whether the English health inequalities strategy was a success or not. Overall, the findings are mixed. Prominent early studies [5, 7, 9] concluded that although the strategy can be considered a partial success, it

failed to reach its ambitious targets. However, these studies were published shortly after the end of the strategy period (2010-2011), by which time it is unlikely that the impact had been fully realised. The studies also could not take into account the trend in inequality after the strategy period had ended.

A more recent study [10] used a modified difference-difference approach to compare trends in health inequalities in England both over time and between other European countries. The study found no detectable effect of the English health inequalities strategy on health inequalities, as compared to trends in Finland, the Netherlands and Italy. However, although the empirical studies mentioned above and others [11, 12, 13] found no effect or increases in inequalities during the strategy period, other studies have found that social inequalities in unemployment, child poverty, housing quality, educational attainment and mortality amenable to health care decreased during the strategy period [14, 15, 16, 17].

Most recently, Barr *et al* [18] have shown using time trend regression analysis that geographical health inequalities marginally decreased during the English health inequalities strategy period, reversing a previously increasing trend. There was also evidence that inequalities had started to increase again following the end of the strategy period. However, these previous studies have largely focused on the first part of the health inequalities strategy, which was aimed at reducing inequalities in life expectancy. No previous study has instead examined the effect of the strategy on its second aim- to reduce inequalities in the IMR. We therefore investigated whether the English health inequalities strategy was associated with a decrease in inequalities in the IMR between the most disadvantaged local authorities and the rest of England, compared with trends both before and after the strategy period.

Methods

Data

We used data from the UK Data Archive and the ONS on the annual number of births and infant deaths in local authority (LA) areas across England between 1983 and 2017. These data were mapped to 323 lower tier local authorities, excluding the Isles of Scilly, City of London and Rutland due to their small population sizes. The outcome variable, the IMR, was calculated as the ratio of number of deaths under the age of 1 per LA per year to the number of live births per LA per year, multiplied by 1,000 [19].

We used the Townsend index of material deprivation to identify the most deprived local authorities in England [20]. This census-based index is made up of four variables (unemployment, non-car ownership, non-home ownership and household overcrowding), which can be used to generate a total score for a particular geographical area. All input data were adjusted to the current LA boundaries [21]. This measure has previously been used in a variety of settings, including as an input to resource allocation mechanisms and as a measure of deprivation in academic studies [22]. Although the original New Labour health

inequalities strategy target was to reduce the gap in the IMR “between the routine and manual groups and the population as a whole”, analysing this gap is not possible with the data available to us. In this study, we have taken a geographical approach, as this is the best available proxy, and also one which fits within the broad aims of the health inequalities strategy.

The strategy was introduced to bring those from the lowest socioeconomic groups to the same level as the rest of the country. To identify those local authorities in the most deprived quintile of the Townsend index (compared to other areas), we used 1991 Census data to represent the situation before the start of the strategy period. We checked the robustness of the results to deprivation measured in each of the 1981, 2001 and 2011 Censuses and an alternative measures of deprivation - the Index of Multiple Deprivation (IMD). These results are shown in Supplementary Appendix 1.

Empirical analyses

The analyses focused on comparing changes in the IMR between the 20% most deprived local authorities and the rest of England. A priori, we expected the level of inequality to decrease during the health inequalities policy period. To estimate the changes in the level of absolute inequality over time, we estimated fixed effects (FE) segmented linear regression models, including marginal spline terms to take account of the time trends, with breakpoints at the beginning and end of the health inequalities strategy period. We interacted these time trends with a dummy variable indicating the deprived group of local authorities, allowing the change in the IMR to vary by the level of deprivation. This segmented regression model therefore provided an estimate of the trend in the absolute difference in the IMR between the most deprived local authorities and the rest of England before, during and after the health inequalities strategy period. The full model specification is presented in Supplementary Appendix 2.

Robustness tests

To fully assess the robustness of our findings, we subjected our empirical analysis to a large number of robustness tests. We estimated models using different breakpoints to the main analysis (Supplementary Appendix 3). We estimated Random Effects models rather than FE models (Supplementary Appendix 4). We included linear spline terms rather than marginal spline terms (Supplementary Appendix 5). To account for outliers, we estimated models with the top 1%, top 5%, bottom 1% and bottom 5% of IMR observations trimmed respectively (Supplementary Appendix 6). Rather than FE models with the IMR in each local authority area as the dependent variable, we estimated Poisson regression models using the number of deaths in each local authority area per year as the dependent variable and the number of births in each local authority area as the offsetting variable (Supplementary Appendix 7). We used different specifications of deprivation, specifically a categorical measure comparing the 5 quintiles of deprivation (Supplementary Appendix 8) and a

continuous measure of deprivation (Supplementary Appendix 9). Finally, we included a measure of ethnicity in the RE model (Supplementary Appendix 10), as it has previously been shown that ethnicity may be a significant risk factor for infant mortality [23]. Ethnicity measures could not be included in the FE models, as data were not available in a consistent manner across all census years.

Results

IMR trends for those in the most deprived local authorities and the rest of England are shown graphically in Figure 1. For both the most deprived local authorities and the rest of England there has been a decline in the IMR over time. For the most deprived local authorities, the average IMR has decreased from 10.69 deaths per 1,000 live births in 1983 to 4.41 deaths per 1,000 live births in 2017, while for the rest of England this rate has decreased from 9.45 deaths per 1,000 live births in 1983 to 3.32 deaths per 1000 live births in 2017.

[FIGURE 1 ABOUT HERE]

Trends in the absolute and relative gap between the most deprived areas and the rest of English are shown graphically in Figure 2 and Figure 3 respectively. Before the start of the strategy period (1983-1999), absolute inequality in the IMR between the most deprived areas and the rest of England had increased from 1.10 in 1983 to 1.43 in 1999. During the strategy period (2000-2010), there was evidence of a decrease in absolute inequalities in the IMR, from 1.75 in 2000 to 1.26 in 2010. After the strategy period (2011-2017), absolute inequalities increased from 0.94 in 2011 to 0.97 in 2017. There was also evidence of a decrease in relative inequalities during the strategy period. Before the start of the strategy period, the relative gap in the IMR between the deprived local authorities and the rest of England increased from 1.16 in 1983 to 1.28 in 1999. During the strategy period, the relative gap in the IMR decreased from 1.37 in 2000 to 1.35 in 2010. After the strategy period, the relative gap in the IMR decreased from 1.22 in 2011 to 1.19 in 2017.

[FIGURES 2 AND 3 ABOUT HERE]

The results from the preferred FE segmented regression models (Table 1) show the trend in absolute infant mortality inequalities estimated before, during and after the strategy period. Before the strategy, the absolute gap in the IMR between the most deprived local authorities and the rest of England increased at an average of 0.033 per year (95% CI 0.003 to 0.063). During the strategy, the gap in the IMR between the most deprived local authorities and the rest of England decreased at an average rate of -0.113 (95% CI -0.168 to -0.058) per year. After the strategy, the gap in the IMR increased at an average rate of 0.034, however these estimates were not statistically precise (95% CI -0.114 to 0.103).

[TABLE 1 ABOUT HERE]

Results from the various robustness tests implied that the trends in absolute inequalities before and after the strategy period shown in the main analysis were associated with a substantial level of uncertainty. For example, when we used linear splines rather than linear marginal splines (as shown in Supplementary Appendix 5), the gap in the IMR between the most deprived local authorities and the rest of England was shown to have decreased after the policy period at an average rate of -0.046 (95% CI -0.096 to 0.003). Furthermore, when we trimmed the bottom 5% of the IMR observations (as shown in Supplementary Appendix 6), the trend of inequality before the policy period was lower than in the main analysis (0.014), and not as statistically precise (95% CI -0.016 to 0.043).

Although it appears that the estimates of the trends before and after the strategy period may depend on the exact specification of the empirical model (both in terms of the magnitude of association and statistical precision), the estimates of the changes of inequality during the strategy period remained relatively stable in all robustness checks. In FE models with a binary measure of deprivation, the upper bound for the trend of inequality during the strategy period from the robustness checks was -0.126 (95% CI -0.186 to -0.067) (Supplementary Appendix 3), while the lower bound was -0.080 (95% CI -0.116 to -0.055) (Supplementary Appendix 5).

Discussion

Our study has shown that geographical inequalities in infant mortality declined in both absolute and relative terms during the English health inequalities strategy period, a time in which the government specifically introduced policies aimed at reducing socioeconomic inequalities in infant mortality. This is in contrast to a previous trend of increasing geographical inequalities in both absolute and relative terms. There was no conclusive evidence that the reduction in geographical inequalities has continued after the strategy period.

The results from this study differ from several previous studies which have shown that the English health inequalities strategy did not fully meet its ambitions [5, 7, 9, 10, 11, 12, 13]. Instead, our study is more in line with the recent work of Barr *et al* [18], which found that socioeconomic inequalities in both male and female life expectancy decreased during the English health inequalities strategy period, and have also begun to increase again after the strategy period. The results are also in line with relatively recent evidence from the USA [24], which has shown that when taking account of the overall downward trend in infant mortality, the IMR is an average of 3% lower during Democratic administrations compared to Republican administrations. Democrats, like New Labour in the 2000s, are more likely to expand public expenditure on health care and welfare, whilst Republicans have tended to decrease expenditure in these areas. Other research from the USA has found that welfare and public health expansion in the 1960s and 1970s following the civil rights acts led to reductions in inequalities in IMR between African-Americans and White Americans [25].

Although we found a decrease in the level of inequality in the IMR during the English health inequalities strategy period, we cannot conclusively say that this is causal relationship, and there are other factors that may explain the decrease in absolute inequality during this period. For example, for a substantial duration of the strategy period (up to the 2008 financial crash), the UK encountered relative macroeconomic stability, with annual growth in GDP and lower rates of unemployment. Recent research from the USA has argued that there may be a significant relationship between macroeconomic conditions and infant mortality, mediated through factors such as the availability of good quality prenatal care [26]. Indeed, infant mortality can be seen to be directly caused by a number of factors, including immaturity, congenital abnormalities, intrapartum causes and sudden infant deaths [27], which in turn have a number of complex and interacting risk factors, including smoking [28], maternal obesity in pregnancy [29], maternal age [11] and ethnicity [23]. Although some of these risk factors are modifiable (smoking and obesity) and may be affected by health care, public health and social policy, others are not.

Strengths and Limitations

There are several strengths in our analysis. Firstly, we have analysed changes over time within local authorities, which, unlike cross sectional analysis, enabled us to control for time constant differences between local authorities. Secondly, we conducted an extensive range of robustness checks to check that our main results were not driven by the model specification used in the main analysis. However, despite these advantages, we must note some important limitations. Firstly, we were not able to establish a true causal relationship between the English health inequalities strategy and IMR. For example, it may be the case that the observed decrease in inequalities in the IMR was due to a broader government strategy rather than the health inequalities strategy itself. Secondly, as with the majority of research in this subject area, our analysis used area-based measures, which are potentially subject to the “ecological fallacy”, which assumes that all individuals residing in a certain geographical area have similar characteristics. Thirdly, we were unable to differentiate neonatal and post-neonatal deaths. Our results also may have differed if we had used infant survival rather than infant mortality as our outcome measure [30] or occupational class instead of an area measure of deprivation. However, births and deaths data by occupational class were not available to us with appropriate time increments to assess the English health inequalities strategy. Finally, as we were only able to measure the level of deprivation every 10 years (in line with the national Census), regression to the mean may have impacted our results.

Policy Implications

The results from this study have important implications for current health policy, especially given the recent increase in infant mortality in England in recent years. Since 2010, successive Conservative led English governments have introduced a programme of austerity. This has included substantial cuts to funding for local authorities, real terms reductions to

the NHS budget, cuts to the education sector and various reductions across the welfare system. Our analysis though suggests that it is *increases* in public spending on health care and welfare that are associated with decreases in inequalities in the IMR. Current policies are arguably going in the wrong direction and may squander some of the gains made in the health inequalities strategy period. Indeed, our results suggest that the English health inequalities strategy was effective at reducing both absolute and relative inequalities in the IMR, and this is something that should therefore be learnt from by current and future governments.

Although the reduction in relative inequalities was shown to be small in magnitude, it has been argued by a number of prominent health inequalities researchers that an overall decrease in the level of mortality will inevitably result in an increase in relative inequalities [31], and therefore any policy intervention that either maintains or decreases the level of relative inequalities can be considered at least a partial success. However, it has also been argued that implementing interventions that decrease absolute inequalities but do not substantially decrease relative inequalities may result in groups at the lower end of the social hierarchy at a significant comparative disadvantage. As argued in the Marmot Review [11], focusing solely on the most disadvantaged in society “will not reduce health inequalities sufficiently”. This implies that in order to reduce health inequalities in health (including inequalities in infant mortality), population wide strategies should be used in combination with evidence based interventions specifically targeted at those areas of higher deprivation, that have a higher health need: with increased levels of infant mortality: a proportionate universal approach.

Conclusion

The multifaceted English health inequalities strategy implemented between 1999 and 2010 was associated with a decrease in geographical inequalities in the IMR between the most and less deprived English local authorities. These results imply that government policies specifically introduced to decrease inequalities in health may be beneficial, and that their discontinuation as a result of austerity may see inequalities increasing again.

References

1. Macfarlane A, Mugford M. Birth counts. Vol. 1. Statistics of pregnancy and childbirth. Vol. 2. Statistics of pregnancy and childbirth: tables. HMSO; 1984.
2. Reidpath DD, Allotey P. Infant mortality rate as an indicator of population health. *J Epidemiol Community Health* 2003 May 1;57(5):344-6.
3. Office for National Statistics. Child mortality in England and Wales: 2016. <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriage/deaths/bulletins/childhoodinfantandperinatalmortalityinenglandandwales/2016> (accessed 24th July 2018).
4. Norman P, Gregory I, Dorling D, Baker A. Geographical trends in infant mortality: England and Wales, 1970-2006. *Health Stat Q.* 2008 Dec 1(40):18.
5. Mackenbach JP. The English strategy to reduce health inequalities. *Lancet.* 2011 Jun 11;377(9782):1986-8.
6. Bambra C. *Health divides: where you live can kill you.* Policy Press; 2016.
7. Mackenbach JP. Can we reduce health inequalities? An analysis of the English strategy (1997–2010). *J Epidemiol Community Health.* 2011 Jul 1;65(7):568-75.
8. The Sutton Trust. 1,000 Children’s Centres ‘Lost’ Since 2009. <https://www.suttontrust.com/newsarchive/1000-childrens-centres-closed-since-2009/> (accessed 10 September 2018).
9. Mackenbach JP. Has the English strategy to reduce health inequalities failed?. *Soc Sci Med.* 2010 Oct 1;71(7):1249-53.
10. Hu Y, van Lenthe FJ, Judge K, Lahelma E, Costa G, de Gelder R, et al. Did the English strategy reduce inequalities in health? A difference-in-difference analysis comparing England with three other European countries. *BMC Public Health.* 2016 Dec;16(1):865.
11. Marmot M, Allen J, Goldblatt P, Boyce T, McNeish D, Grady M. *Fair society, healthy lives. The Marmot Review.* 2010.
12. Department of Health. *Tackling health inequalities: 2007 status report of the program for action.* London. 2007. http://webarchive.nationalarchives.gov.uk/+/www.dh.gov.uk/en/Publicationsandstatistics/Publications/DH_083471.(Accessed 24th July 2018).
13. Department of Health. *Tackling health inequalities: 10 years on.* London.2009. http://webarchive.nationalarchives.gov.uk/20130107105354/http://www.dh.gov.uk/prod_consum_dh/groups/dh_digitalassets/documents/digitalasset/dh_098934.pdf. (Accessed 24th July 2018).
14. Bambra C. Reducing health inequalities: new data suggest that the English strategy was partially successful. *J Epidemiol Community Health.* 2012 Jul 1;66(7):662
15. Barr B, Bambra C, Whitehead M. The impact of NHS resource allocation policy on health inequalities in England 2001-11: longitudinal ecological study. *BMJ.* 2014 May 27;348:g3231.

16. Buck D, Dixon A. Improving the allocation of health resources in England. London: Kings Fund. 2013
17. Asaria M, Ali S, Doran T, Ferguson B, Fleetcroft R, Goddard M, et al. How a universal health system reduces inequalities: lessons from England. *J Epidemiol Community Health*. 2016 70:637-643.
18. Barr B, Higgerson J, Whitehead M. Investigating the impact of the English health inequalities strategy: time trend analysis. *BMJ*. 2017 26;358:j3310.
19. Handbook of Vital Statistics Systems and Methods, Volume 1: Legal, Organisational and Technical Aspects, United Nations Studies in Methods, Glossary, Series F, No. 35, United Nations, New York 1991.
20. Townsend P, Phillimore P, Beattie A. Health and deprivation: inequality and the North. Routledge; 1988.
21. Norman PD, Darlington-Pollock F. The changing geography of deprivation in Great Britain: Exploiting small area census data, 1971 to 2011.
22. Senior M. Deprivation indicators. The census data system. 2002:123-39.
23. Moser K, Stanfield KM, Leon DA. Birthweight and gestational age by ethnic group, England and Wales 2005: introducing new data on births. *Health Stat Q*. 2008 Oct 1;39:22-31.
24. Rodriguez JM, Bound J, Geronimus AT. US infant mortality and the President's party. *Int J Epidemiol*. 2013 Dec 31;43(3):818-26.
25. Krieger N, Chen JT, Coull B, Waterman PD, Beckfield J. The unique impact of abolition of Jim Crow laws on reducing inequities in infant death rates and implications for choice of comparison groups in analyzing societal determinants of health. *Am J Public Health*. 2013 Dec;103(12):2234-44.
26. Orsini C, Avendano M. Macro-economic conditions and infant health: a changing relationship for black and white infants in the United States. *PloS one*. 2015 May 14;10(5):e0123501.
27. Gray R, Hollowell J, Brocklehurst P, Graham H, Kurinczuk J. Health inequalities infant mortality target: technical background. Inequalities in infant mortality project briefing paper 2. 2009
28. Johansson AL, Dickman PW, Kramer MS, Cnattingius S. Maternal smoking and infant mortality: does quitting smoking reduce the risk of infant death?. *Epidemiology*. 2009 Jul 1:590-7.
29. Meehan S, Beck CR, Mair-Jenkins J, Leonardi-Bee J, Puleston R. Maternal obesity and infant mortality: a meta-analysis. *Pediatrics*. 2014 Apr 1:peds-2013.
30. Kjellsson, G., Gerdtham, U.G. and Petrie, D. Lies, damned lies, and health inequality measurements: understanding the value judgments. *Epidemiology* 2015. 26(5), p.673.
31. Scanlan JP. Race and mortality. *Society*. 2000 Jan 1;37(2):29-35.

Figure 1




Figure 2




Figure 3




Table 1

Table 1- Trend in absolute inequalities in the IMR between the most deprived local authorities and the rest of England, before, during and after the English HI strategy period. Trend is shown as the annual increase or decrease (minus values) in the absolute gap in the IMR (defined as the number of deaths under one per 1000 births in the local authority area).	
Period	Annual change in absolute gap in the IMR between the most deprived LAs and the rest of England [95% CI]and the rest of England [95% CI]
Before (1983-1999)	0.034 [0.001 to 0.067]
During (2000-2010)	-0.116 [-0.178 to -0.053]
After (2011-2017)	0.042 [-0.042 to 0.125]
N=11,305 (323 Local Authorities x 35 Years), Overall R ² =0.3857	
Estimates based on a fixed effects regression model using LA panel dataset of IMR from 1983 to 2017. Break points at 1999 and 2010.	