



This is a repository copy of *The evolutionary status of Cataclysmic Variables: Eclipse modelling of 15 systems*.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/144934/>

Version: Accepted Version

Article:

McAllister, M., Littlefair, S.P., Parsons, S.G. et al. (13 more authors) (2019) The evolutionary status of Cataclysmic Variables: Eclipse modelling of 15 systems. *Monthly Notices of the Royal Astronomical Society*, 486 (4). pp. 5535-5551. ISSN 0035-8711

<https://doi.org/10.1093/mnras/stz976>

This is a pre-copyedited, author-produced PDF of an article accepted for publication in *Monthly Notices of the Royal Astronomical Society* following peer review. The version of record M McAllister, S P Littlefair, S G Parsons, V S Dhillon, T R Marsh, B T Gaensicke, E Breedt, C Copperwheat, M J Green, C Knigge, D I Sahman, Martin J Dyer, P Kerry, R P Ashley, P Irawati, S Rattanasoon, *The evolutionary status of Cataclysmic Variables: Eclipse modelling of 15 systems*, *Monthly Notices of the Royal Astronomical Society* is available online at: <https://doi.org/10.1093/mnras/stz976>.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

The evolutionary status of Cataclysmic Variables: Eclipse modelling of 15 systems.

M. McAllister¹, S. P. Littlefair¹, S. G. Parsons¹, V. S. Dhillon^{1,2}, T. R. Marsh³
 B. T. Gänsicke³, E. Breedt⁴, C. Copperwheat⁵, M. J. Green³, C. Knigge⁶
 D. I. Sahman¹, Martin J. Dyer¹, P. Kerry¹, R. P. Ashley³, P. Irawati⁷, S. Rattanasoon⁷

¹Dept of Physics and Astronomy, University of Sheffield, Sheffield, S3 7RH, UK

²Instituto de Astrofísica de Canarias (IAC), E-38200, La Laguna, Tenerife, Spain

³Dept of Physics, University of Warwick, Coventry, CV4 7AL, UK

⁴Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge, CB3 0HA, UK

⁵Astrophysics Research Institute, Liverpool John Moores University, IC2, Liverpool Science Park, L3 5RF, UK

⁶School of Physics & Astronomy, University of Southampton, Southampton SO17 1BJ, UK

⁷National Astronomical Research Institute of Thailand, 191 Siriphanich Bldg., Huay Kaew Road, Chiang Mai 50200, Thailand

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

We present measurements of the component masses in 15 Cataclysmic Variables (CVs) - 6 new estimates and 9 improved estimates. We provide new calibrations of the relationship between superhump period excess and mass ratio, and use this relation to estimate donor star masses for 225 superhumping CVs. With an increased sample of donor masses we revisit the implications for CV evolution. We confirm the high mass of white dwarfs in CVs, but find no trend in white dwarf mass with orbital period. We argue for a revision in the location of the orbital period minimum of CVs to 79.6 ± 0.2 min, significantly shorter than previous estimates. We find that CV donors below the gap have an intrinsic scatter of only $0.005 R_{\odot}$ around a common evolutionary track, implying a correspondingly small variation in angular momentum loss rates. In contrast to prior studies, we find that standard CV evolutionary tracks - without additional angular momentum loss - are a reasonable fit to the donor masses just below the period gap, but that they do not reproduce the observed period minimum, or fit the donor radii below $0.1 M_{\odot}$.

Key words: binaries: close – binaries: eclipsing – stars: evolution – novae, cataclysmic variables – white dwarfs – stars: dwarf novae

1 INTRODUCTION

Cataclysmic Variables (CVs) are close binary stars in which a white dwarf is accreting material from a low-mass donor star. Without angular momentum loss (AML) from the system, mass transfer could not be sustained; thus it is the AML that drives the secular evolution of CVs. The currently accepted picture of CV evolution is that CVs evolve from long to short periods under the influence of AML caused by magnetic braking. A reduction in AML due to magnetic braking is thought to arise when the donor becomes fully convective. This causes the CV to become detached, and is the cause of the dearth of CVs in the 2–3 hour orbital period range; the CV *period gap*. When the CV resumes mass transfer, AML is driven by gravitational radiation and the mass transfer rate is lower. The CV evolves slowly through a *period minimum*, which arises because the thermal timescale of the donor becomes comparable to the mass loss timescale, and the donor begins to expand in response to mass loss, which leads to a widening of the orbit.

This long-standing picture has survived for over 35 years (Rappaport et al. 1982, 1983) despite the fact that it struggles to explain the observed value of the period minimum (Gänsicke et al. 2009), the scarcity of known post-period-minimum systems (Hernández Santisteban et al. 2018) and the average high white dwarf mass in CVs (Zorotovic et al. 2011). Modifications to the standard model exist that can potentially explain some of these issues. The orbital period minimum problem can be solved with an additional source of AML for short period systems (Patterson 1998; Knigge et al. 2011), and AML in nova outbursts may cause CVs with low-mass white dwarfs to be unstable, explaining the high average white dwarf mass (Schreiber et al. 2016; Nelemans et al. 2016). However, it remains to be seen if those modifications can correctly describe the observed properties of known CVs. In particular, the mass and radius of the donor star is a sensitive probe of the secular evolution. This is because the radius of the donor star in a CV can be inflated from the main-sequence value, by some amount that depends upon

the mass-loss history of the donor. In particular, the donor radius is more likely to track the long-term average mass loss rate than other physical properties of the CV such as the accretion light or the effective temperature of the accreting white dwarf (see [Knigge et al. 2011](#), and references within).

One of the best methods of measuring donor masses and radii is to model the primary eclipse. During primary eclipse, the white dwarf and accretion disc are occulted, along with the bright spot, located where the accretion stream impacts the outer rim of the disc. The path of the gas stream is determined by the mass ratio, and so the detailed shape of the primary eclipse contains enough information to derive extremely precise masses that are consistent with conventional spectroscopic methods (see [Tulloch et al. 2009](#); [Copperwheat et al. 2010](#); [Savoury et al. 2012](#), for example). The photometric method has the advantage that it does not rely on detection of the light from the donor star, which is often invisible given the much brighter white dwarf and accretion disc, particularly for CVs with shorter orbital periods. It does however require high quality lightcurves of the eclipses, which occur on timescales of minutes. With this in mind, our group has been acquiring high quality lightcurves of eclipsing CVs with the high time-resolution instruments ULTRACAM ([Dhillon et al. 2007](#)) and ULTRASPEC ([Dhillon et al. 2014](#)). Here we present the analysis of 15 systems, and review the evolutionary status of CV systems in light of the results.

1.1 Systems selected for eclipse modelling

The 15 systems modelled in this paper are listed in [Table 1](#). CTCV 1300, DV UMa, SDSS 1152, SDSS 1501 have existing mass determination from eclipse modelling of ULTRACAM data ([Savoury et al. 2011](#)), whilst Z Cha, OY Car, IY UMa, GY Cnc and SDSS 1006 have existing mass determinations in the literature from various methods ([Wood et al. 1986](#); [Wade & Horne 1988](#); [Wood & Horne 1990](#); [Thorstensen 2000](#); [Steehls et al. 2003](#); [Southworth et al. 2009](#); [Copperwheat et al. 2012](#)). The existing mass determinations have large associated errors, and we re-analyse them here in the light of new data, and an updated modelling approach (see [McAllister et al. 2017a](#), for details). The remaining 6 systems have no existing donor mass estimates, and were chosen from the eclipsing CVs observed with ULTRACAM/ULTRASPEC to date; the primary reason for their selection was an eclipse shape suitable for modelling, with visible white dwarf and bright spot eclipses.

2 OBSERVATIONS AND DATA REDUCTION

The observations in this paper span a range of dates from May 2003 to Feb 2017. All data were taken with the triple-band fast camera ULTRACAM, or the single-band fast camera ULTRASPEC. ULTRACAM data were taken on three telescopes; the 4.2-m William Herschel Telescope (WHT) situated at the Roque de los Muchachos Observatory on La Palma, Spain, the 8.2-m Very Large Telescope (VLT) at Paranal, Chile, and the 3.5-m New Technology Telescope (NTT) located at La Silla, Chile. All ULTRASPEC data

were taken using the 2.4-m Thai National Telescope (TNT), located on Doi Inthanon in Thailand. All observations were obtained using the Sloan Digital Sky Survey (SDSS) filter set, with the exception of some of the ULTRASPEC observations, which use the KG5 filter. This filter is described in detail in [Hardy et al. \(2017\)](#); it is a broadband filter encompassing the SDSS u' , g' and r' passbands. For a full journal of observations, see [Table C1](#).

Data reduction was carried out using the ULTRACAM pipeline reduction software (see [Dhillon et al. 2007](#)). One or more nearby, photometrically stable comparison stars were used to correct for transparency variations during observations. If the comparison stars have tabulated SDSS magnitudes, we used these to transform the photometry into the $u' g' r' i' z'$ standard system ([Smith et al. 2002](#)), otherwise observations of standard stars from the nearest photometric night were used. Photometry was corrected for extinction using the median extinction coefficients for each observatory, as derived from long duration time-series taken with ULTRACAM and ULTRASPEC.

3 METHODS

3.1 Orbital Ephemerides

Updated orbital ephemerides for the CVs in this paper were calculated, and are shown in [Table 1](#). Mid-eclipse times were determined by averaging the time of white dwarf ingress and egress, as determined by locating the minima and maxima of a smoothed lightcurve derivative. Mid eclipse times were corrected to the Solar System Heliocentre or Barycentre using ASTROPY ([The Astropy Collaboration et al. 2018](#)). The correction used was decided upon a system-to-system basis, and depended on previous mid-eclipse times and ephemerides in the literature. Heliocentric times are recorded in Coordinated Universal Time (UTC), Barycentric times in Barycentric Dynamical Time (TDB). Mid-eclipse times for each individual eclipse observed are presented in [Table C1](#).

3.2 Eclipse light-curve modelling

The model used to fit the eclipse light curve is described by [Savoury et al. \(2011\)](#). The important assumptions in the model are that the bright spot lies on the ballistic trajectory from the donor, the white dwarf follows a theoretical mass-radius relation and that the white dwarf is unobscured. The model has recently received two major improvements, as outlined in [McAllister et al. \(2017a\)](#). The model now has the ability to fit multiple lightcurves simultaneously whilst sharing parameters that do not change; such as the mass ratio q , the white dwarf eclipse width $\Delta\Phi$ and the white dwarf radius, scaled by the binary separation R_1/a . In addition, the model now has a statistical treatment of flickering using Gaussian Processes (GPs) that makes the uncertainty estimates for these parameter robust in the presence of flickering. For each system we either fit all the individual eclipses, or averaged several eclipses in the same filter. Averaging eclipses can ease convergence of the model, by reducing the number of free parameters, but it is not suitable when the lightcurve features change between eclipses, for example due to a changing accretion disc radius.

Object	Right Ascension	Declination	Out-of eclipse Mag. (g')	T_0 (MJD)	P_{orb} (d)	N_{ecl}	Add. Ecl. Times
CTCV J1300–3052	13 00 29.05	−30 52 57.1	18.6	54262.099166(18) ^h	0.0889406998(17)	4	1
DV UMa	09 46 36.65	+44 46 45.1	18.7	52782.973948(10) ^h	0.0858526308(7)	4	2,3,4
SDSS J115207.00+404947.8	11 52 07.01	+40 49 48.0	19.5	55204.101279(6) ^h	0.0677497026(3)	7	5
SDSS J150137.22+550123.4	15 01 37.24	+55 01 23.5	19.0	56178.870444(8) ^h	0.05684126603(21)	12	–
CSS080623 J140454–102702	14 04 53.97	−10 27 02.3	19.5	55329.234631(13) ^h	0.059578971(3)	10	–
CSS110113 J043112–031452	04 31 12.45	−03 14 51.6	19.5	55942.014642(15) ^h	0.0660508707(18)	12	–
GY Cnc	09 09 50.55	+18 49 47.5	16.7	55938.263734(22) ^b	0.175442399(6)	12	–
IY UMa	10 43 56.73	+58 07 31.9	17.1	56746.6395010(9) ^h	0.07390892818(21)	10	8
OY Car	10 06 22.07	−70 14 04.6	15.6	55353.996477(3) ^h	0.06312092545(24)	7	–
SDSS J090103.94+480911.0	09 01 03.94	+48 09 11.0	19.5	55942.116358(8) ^h	0.0778805321(5)	10	9
SDSS J100658.40+233724.4	10 06 58.42	+23 37 24.6	18.6	56682.72973(5) ^h	0.185913107(13)	11	7,10
SSS130413 J094551–194402	09 45 51.00	−19 44 00.8	16.7	56683.673971(12) ^h	0.0657692903(12)	17	6
SSS100615 J200331–284941	20 03 31.27	−28 49 41.3	19.6	56873.023625(5) ^h	0.0587045(4)	3	–
V713 Cep	20 46 38.70	+60 38 02.8	18.5	56176.936402(7) ^h	0.0854185080(12)	15	11
Z Cha	08 07 27.75	−76 32 00.7	15.6	53498.011471(4) ^h	0.0744992631(3)	14	–

Table 1. Ephemerides for the CVs modelled in this paper. T_0 is the mid-eclipse time of cycle 0, P_{orb} is the orbital period, while N_{ecl} is the total number of eclipses obtained. References for additional eclipse times: (1) Tappert et al. (2004), (2) Howell et al. (1988), (3) Patterson et al. (2000), (4) Nogami et al. (2001), (5) Southworth et al. (2010), (6) Thorstensen et al. (2016), (7) Woudt (priv. comm.), (8) Coppejans (priv. comm.), (9) Dillon et al. (2008), (10) Southworth et al. (2007), (11) Bours (priv. comm.).

^hHeliocentric times in HMJD(UTC), ^bBarycentric times in BMJD(TDB).

Eclipse averaging was used for six systems: CSS080623, CSS110113, DV UMa, SDSS 0901, SDSS 1152, SSS100615. All systems have multiple eclipse light curves observed close together in time (e.g. during the same observing run), and contain only low amplitude flickering. When selecting eclipses for the construction of each average eclipse, great care was taken to exclude any eclipses with differing disc radius/flux and/or bright spot shape/flux changes. Firstly, only eclipses obtained during the same observing run were considered for each average eclipse. Secondly, before averaging, all eclipses were phase-folded and overlaid, with any differing eclipses removed from consideration. An average eclipse was created for each available wavelength band, typically $u'g'r'$ or $u'g'i'$. As both CSS080623 and SDSS 0901 have multiple eclipses from two separate observing runs, two average eclipses in each wavelength band were created. For the remaining nine systems, we did not average lightcurves prior to fitting.

In general, the majority of eclipses showing a clear bright spot ingress feature were selected for modelling. However, for systems with many high signal-to-noise eclipses containing very clear bright spot eclipse features (e.g. OY Car and Z Cha), only six were selected. In these cases, the inclusion of additional eclipses had an insignificant effect on the system parameter values and errors, and did not justify the resulting increased model complexity and computational time. This approach was also taken with SSS130413 and V713 Cep, two systems with moderately clear bright spot features.

4 RESULTS

4.1 Simultaneous Eclipse Light Curve Modelling

For each of the 15, the chosen eclipses were fit with the CV eclipse model, with GPs used to model the flickering component. The binary model contains two possible versions

of the bright spot (see Savoury et al. 2011, for details). A more complex bright spot model was used for all but three systems (SDSS 1501, SSS100615, V713 Cep). The simple bright spot was used in these three systems due to each containing a weak bright spot component in their eclipse light curves. The typical phase range of the eclipse light curves modelled was -0.10 to 0.15 , however an extended phase range was used for a number of systems. The phase range was increased for systems with a prominent bright spot (e.g. CTCV 1300, GY Cnc, SDSS 1006) in addition to SDSS 1501 (tenuous bright spot component) and V713 Cep (combination of heavy flickering post-eclipse and significant disc contribution).

Posterior probability distributions of all parameters in the binary model were estimated using a Markov Chain Monte Carlo (MCMC) approach. The full results of all eclipse fits are shown in Figures A1–A15. Figure 1 shows an example g' -band eclipse light curve fit for each system. In addition to the most probable fit of the eclipse model (blue line), a blue band is plotted that covers 1σ from the mean of a random sample (size 1000) of the MCMC chain. The grey points represent the actual eclipse light curves, while the black points are the result of subtracting the GP's posterior mean (itself shown, $\pm 1\sigma$, by the red band covering the residuals below each plot). Also plotted are the separate components of the eclipse model: white dwarf (purple), bright spot (red), accretion disc (yellow) and donor (green).

4.2 System Parameters

Once the parameters of the binary model are estimated, the system parameters can be found. A full discussion can be found in McAllister et al. (2017a). In brief, this involves an iterative procedure where the white dwarf spectral energy distribution (SED) – measured from the eclipse depth of the white dwarf – is fit by white dwarf atmosphere models (Bergeron et al. 1995). This yields estimates of the white





Figure 1. Eclipse model fits to *g*-band light curves of 15 CVs. The lightcurves are shown in grey points; see Section 4.1 for full details of what is plotted. The system name is displayed in the top-right corner of each plot, along with whether the eclipse is an individual (Ind) or average (Ave) eclipse. For individual eclipses the cycle number is shown, while for average eclipses the month and year of the eclipses is shown. See Appendix A for a complete set of eclipse plots.

dwarf temperature T_1 and distance d . The values of q , $\Delta\Phi$ and R_1/a from the binary model, combined with Kepler's third law and a temperature-corrected mass-radius relationship for the white dwarf, are used to calculate the posterior probability distribution functions (PDFs) of the system parameters:

- mass ratio, q ;
- white dwarf mass, M_1 ;
- white dwarf radius, R_1 ;
- white dwarf $\log g$;
- donor mass, M_2 ;

- donor radius, R_2 ;
- binary separation, a ;
- inclination, i .

System parameter values (see Table 2) were then obtained from the peak of each posterior PDF, with errors from the 67% confidence level. The results of the white dwarf SED fits are shown in Figure 2, and the resulting T_1 and d values for each system are also displayed in Table 2. Note that the white dwarf flux fitting was not carried out for either IY UMa or SDSS 1006, due to the lack of u' -band eclipses in their eclipse model fits. Thankfully, precise measurements

Figure 1. *Continued.*

of T_1 for both IY UMa¹ (18000 ± 1000 K) and SDSS 1006 (16000 ± 1000 K) from spectral fitting are given in Pala et al. (2017).

As a sanity check on our white dwarf atmosphere fitting, we can compare the derived distances with the parallaxes found in Gaia DR2 (Lindgren et al. 2018). We naively converted our distances to parallaxes, and compared to the parallaxes in Gaia DR2. The results are perfectly consistent with Gaussian statistics; the parallaxes of all but 4 out of

15 CVs agree within 1 standard deviation, whilst the most discrepant CV (GY Cnc) has a 2σ discrepancy between our derived distance and the Gaia DR2 parallax. This gives us confidence on our distance estimates and also their uncertainties.

¹ IY UMa entered outburst between the observations of Pala et al. (2017) and this work, so this T_1 measurement may be slightly lower than T_1 of the white dwarf in the eclipse light curves.

In section 5 we discuss the implications of the measured system parameters for CV evolution. However, before then, we discuss some remarkable aspects of the data for two individual systems.




Figure 2. White dwarf fluxes for 13 CVs, showing the white dwarf fluxes from the eclipse model fits (blue) and white dwarf atmosphere predictions (red), at wavelengths corresponding to u' (355.7 nm), g' (482.5 nm), $KG5$ (507.5 nm), r' (626.1 nm) and i' (767.2 nm) filters. The name of each system is displayed in the top-right corner of each plot.




Figure 2. *Continued.*




Figure 3. O–C diagram for all 12 available ULTRACAM eclipses of SDSS 1501, spanning ~ 8 yrs. The vertical dashed line corresponds to Sep 2010, when SDSS 1501 was reportedly observed in superoutburst. The y-axis covers ± 15 s.

4.3 White Dwarf Flux and Orbital Period Variations in SDSS 1501

The eclipses of SDSS 1501 are white dwarf dominated, but some show faint bright spot features. There are a total of 15 available ULTRACAM eclipses of SDSS 1501, obtained during observing runs in 2004 (one eclipse), 2006 (eight), 2010 (two) and 2012 (one) (see Table C1 for further details). However, only the single eclipses from 2004 and 2012 show signs of a bright spot eclipse, so both² were chosen for simultaneous eclipse modelling described above.

It became apparent that there was an appreciable increase in white dwarf flux across all three ($u'g'r'$) bands between the 2004 and 2012 eclipses. For this reason, model atmosphere fitting to the white dwarf fluxes was carried out separately for each eclipse, as shown in Figure 2. The resulting d and T_1 for each eclipse are shown in Table 2. The white dwarf in 2012 appears marginally hotter, but note the 1.6σ discrepancy in d , which should of course remain constant. Both distances are formally consistent with a formal inference of the distance from Gaia DR2, including a weak distance prior (Bailer-Jones et al. 2018). However, the most likely distance from Gaia DR2 is 340 pc; favouring the 2012 distance estimate. The white dwarf flux fitting was repeated for both eclipses, but this time with d held fixed at 360 pc. This now gives $T_1(2004) = 12100 \pm 300$ K and $T_1(2012) = 15800 \pm 300$ K, a much larger increase of 3700 K.

Such a large discrepancy in T_1 indicates that the white dwarf in SDSS 1501 underwent a period of enhanced accretion between 2004 and 2012, most likely a superoutburst. According to vsnet-alert 12169³, the superoutburst occurred in Sep 2010, with the observer claiming to have observed SDSS 1501 in outburst in addition to obtaining part of a superhump. Unfortunately, there is not enough coverage of this outburst to determine a superhump period.

² Each ULTRACAM eclipse is in three bands ($u'g'r'$), giving six individual eclipses for modelling.

³ <http://ooruri.kusastro.kyoto-u.ac.jp/mailarchive/vsnet-alert/12169>

In addition to the white dwarf flux variations, SDSS 1501 also exhibits small orbital period variations. The white dwarf-dominated SDSS 1501 eclipses enable very precise mid-eclipse times to be obtained. We show the mid-eclipse times – after the subtraction of a linear ephemeris – in Figure 3. The orbital period of SDSS 1501 appears to depart from linearity by approximately ± 7 s over the ~ 8 yr ULTRACAM observational baseline. Such variations are not uncommon in CVs, and are thought to be caused by a magnetically-driven process within the donor. However, they are not observed in CVs with donors of spectral type later than M6 (Bours et al. 2016), due to magnetic activity in the donor decreasing with later spectral types. SDSS 1501’s donor mass obtained through eclipse modelling is substellar ($0.061 \pm 0.004 M_{\odot}$), strongly indicating a spectral type later than M6, and so the observation of period variations is surprising.

A logical deduction from looking at Figure 3 is that the superoutburst from Sep 2010 (dashed line) may have caused the observed change in orbital period, as the ephemeris appears approximately linear up until this point. In this scenario, the 2012 eclipse occurs ~ 21 s later than expected, implying an increase in SDSS 1501’s orbital period of 0.0016 s ($\Delta P_{\text{orb}}/P_{\text{orb}} = 3.2 \times 10^{-7}$). It is not clear how the superoutburst could have caused such a large change in the orbital period. If some fraction of the disc mass was ejected during superoutburst, we would expect $\Delta P_{\text{orb}}/P_{\text{orb}} = 2M_{\text{ej}}/(M_1 + M_2)$, where M_{ej} is the mass ejected. This implies ejected masses of $10^{-7} M_{\odot}$, and disc masses in excess of this. A period change might be induced by a change in the quadrupole moment of the white dwarf and disc, due to the disc draining onto the white dwarf. In this case, Applegate (1992) gives

$$\Delta P_{\text{orb}}/P_{\text{orb}} \approx -\frac{9\Delta Q}{M_1 a^2},$$

where ΔQ is the change in quadrupole moment. We can obtain an order-of-magnitude estimate for ΔQ if we approximate the disc as a ring of mass M_{d} and radius $a/3$, and assume that during superoutburst the disc completely drains onto the white dwarf, giving $\Delta Q \approx -M_{\text{d}}a^2/9$. Therefore $\Delta P_{\text{orb}}/P_{\text{orb}} \approx M_{\text{d}}/M_1$, again implying disc masses of order $10^{-7} M_{\odot}$. With SDSS 1501’s system parameters known (Table 2), the pre-outburst white dwarf temperature can be used to determine a medium-term average mass transfer rate for SDSS 1501 (Townesley & Bildsten 2003, 2004; Townesley & Gänsicke 2009) of $\dot{M} = 9.3 \times 10^{-11} M_{\odot} \text{ yr}^{-1}$. Period minimum systems are observed to have superoutburst cycles of order 20–30 yrs. Therefore the required disc masses are unrealistic, and the Sep 2010 superoutburst is not (at least not fully) responsible for the period variations exhibited by SDSS 1501. Another possible cause of the period variations is the presence of a third body within the system, however additional precise mid-eclipse timings are required in order to investigate this further.

4.4 Observed Low State of V713 Cep

The ULTRACAM/ULTRASPEC data archive contains a total of 15 V713 Cep eclipses, with two ULTRACAM eclipses (cycle nos. 11 [$u'g'r'$] and 3655 [$u'g'i'$]) showing clear bright-spot features suitable for eclipse modelling. A feature of




Figure 4. g' -band eclipse light curve (24 Jun 2015, cycle no. 11955) of V713 Cep during a low state.

these two eclipses is a notable disc contribution (see Figure A14), which is seen in all other V713 Cep eclipses in the archive, with the exception of one. The ULTRACAM $u'g'r'$ eclipse of 24 Jun 2015 (cycle no. 11955, g' -band eclipse shown in Figure 4) contains no obvious signs of either a disc or bright spot eclipse, and at first glance resembles an eclipse of a detached, non-accreting binary. However, on closer inspection there are signs of flickering outside of white dwarf eclipse, as well as a very slight curvature inside eclipse. These two features are both evidence for the presence of an – albeit considerably diminished – accretion disc. A dwindling accretion disc and no sign of a bright spot indicates that the secondary has stopped supplying the disc with material and the system is in what is known as a ‘low state’.

Low states are relatively common phenomena for both magnetic CVs and a subgroup of novalike (NL) CVs called VY Scl stars, however they appear to be very rare (and unexpected) for DNe below the period gap. In fact, there is only one other documented occurrence in the literature – an extended (> 2 yrs) low state of IR Com (Manser & Gänsicke 2014). Given the rarity of low states in DNe, it is notable that IR Com and V713 Cep have similar orbital periods, just at the lower edge of the period gap. With only one eclipse of V713 Cep obtained during its low state, it is not known exactly how long this low state lasted. An upper limit of 403 days can be estimated based on the timings of other ULTRACAM eclipses, and therefore it was significantly shorter than the low state of IR Com.

5 DISCUSSION

With the new and revised system parameters obtained in this work, we now discuss what impact these results may have on the current understanding of CVs and their evolution. In what follows, we combine the parameters presented here with a compilation of reliable parameters for 46 CVs from the literature. This compilation is presented in Table B1.

It has been shown that there is a significant discrepancy between the mean white dwarf mass in the field and that within CVs. Zorotovic et al. (2011) obtained a mean CV white dwarf mass of $0.82 \pm 0.03 M_{\odot}$, and an intrinsic scatter of white dwarf masses of $\sigma = 0.15 M_{\odot}$. With the

Parameter	CSS080623	CSS110113	CTCV 1300	DV UMa
q	0.114 ± 0.005	0.105 ± 0.006	0.233 ± 0.004	$0.172^{+0.002}_{-0.007}$
M_1 (M_\odot)	0.710 ± 0.019	$1.00^{+0.04}_{-0.01}$	0.717 ± 0.017	1.09 ± 0.03
R_1 (R_\odot)	$0.0117^{+0.0001}_{-0.0004}$	0.0080 ± 0.0003	0.01133 ± 0.00021	0.0072 ± 0.0004
M_2 (M_\odot)	0.081 ± 0.005	0.105 ± 0.007	$0.166^{+0.006}_{-0.003}$	$0.187^{+0.003}_{-0.012}$
R_2 (R_\odot)	0.1275 ± 0.0024	0.149 ± 0.003	$0.2111^{+0.0025}_{-0.0014}$	$0.215^{+0.001}_{-0.005}$
a (R_\odot)	0.593 ± 0.005	$0.711^{+0.009}_{-0.003}$	0.805 ± 0.007	$0.889^{+0.006}_{-0.012}$
K_1 (km s^{-1})	50.8 ± 2.3	$51.1^{+2.9}_{-2.4}$	86.4 ± 1.4	$76.1^{+0.9}_{-2.9}$
K_2 (km s^{-1})	449^{+1}_{-6}	487 ± 3	371 ± 3	444 ± 4
i ($^\circ$)	80.76 ± 0.19	79.94 ± 0.19	$86.9^{+0.5}_{-0.2}$	$83.29^{+0.29}_{-0.10}$
$\log g$	$8.15^{+0.01}_{-0.04}$	8.63 ± 0.03	8.186 ± 0.019	8.77 ± 0.04
T_1 (K)	15500 ± 1700	14500 ± 2200	11000 ± 1000	17400 ± 1900
d (pc)	550 ± 60	430 ± 60	340 ± 40	380 ± 40
Parameter	GY Cnc	IY UMa	OY Car	SDSS 0901
q	$0.448^{+0.014}_{-0.021}$	$0.146^{+0.009}_{-0.001}$	$0.1065^{+0.0009}_{-0.0029}$	$0.182^{+0.009}_{-0.004}$
M_1 (M_\odot)	0.881 ± 0.016	$0.955^{+0.013}_{-0.028}$	$0.882^{+0.011}_{-0.015}$	$0.752^{+0.024}_{-0.018}$
R_1 (R_\odot)	$0.00976^{+0.00021}_{-0.00018}$	$0.0087^{+0.0003}_{-0.0001}$	$0.00957^{+0.00018}_{-0.00012}$	$0.01105^{+0.00022}_{-0.00029}$
M_2 (M_\odot)	$0.394^{+0.016}_{-0.022}$	0.141 ± 0.007	$0.093^{+0.004}_{-0.001}$	0.138 ± 0.007
R_2 (R_\odot)	$0.446^{+0.006}_{-0.009}$	0.1770 ± 0.0028	$0.1388^{+0.0018}_{-0.0003}$	0.182 ± 0.003
a (R_\odot)	1.429 ± 0.012	$0.765^{+0.004}_{-0.009}$	0.662 ± 0.003	0.739 ± 0.007
K_1 (km s^{-1})	125 ± 4	66^{+4}_{-1}	50.4 ± 0.9	73 ± 3
K_2 (km s^{-1})	278.0 ± 2.4	453 ± 3	475.9 ± 2.1	401 ± 3
i ($^\circ$)	$77.06^{+0.29}_{-0.18}$	$84.9^{+0.1}_{-0.5}$	$83.27^{+0.10}_{-0.13}$	$81.4^{+0.1}_{-0.3}$
$\log g$	8.40 ± 0.019	8.54 ± 0.03	$8.422^{+0.017}_{-0.013}$	$8.228^{+0.022}_{-0.025}$
T_1 (K)	25900 ± 2300	–	18600^{+2800}_{-1600}	14900 ± 2000
d (pc)	320 ± 30	–	90 ± 5	600 ± 70

Table 2. System parameters for the 15 eclipsing systems analysed in this paper.

updated sample of CV masses now available, we can revise the mean white dwarf mass in CVs, following the procedure outlined in Appendix B of [Knigge \(2006\)](#), to $0.81 \pm 0.02 M_\odot$ ($\sigma = 0.13 M_\odot$), entirely consistent with [Zorotovic et al's](#) value.

One way to explain the presence of high white dwarf masses in CVs is through white dwarf mass growth through steady accretion across the lifetime of a CV. Since CVs evolve to shorter orbital periods over their lives, this requires the observation of higher white dwarf masses in systems with lower orbital periods. To test this, $\langle M_1 \rangle$ was re-calculated for 31 systems below the period gap ($P_{\text{orb}} \sim 2.15$ hrs), giving $\langle M_1 \rangle(\text{below gap}) = 0.81 \pm 0.02 M_\odot$ ($\sigma = 0.10 M_\odot$), and for 16 systems above the gap ($P_{\text{orb}} \sim 3.18$ hrs), giving $\langle M_1 \rangle(\text{above gap}) = 0.82 \pm 0.02 M_\odot$ ($\sigma = 0.10 M_\odot$). We therefore see no evidence for white dwarf mass growth in CVs. While white dwarf mass growth in CVs appears doubtful,

further precise white dwarf masses from systems at long period (> 3 hrs) are required before it can be entirely dismissed.

5.1 Testing the Validity of the Empirical CAML Model

An alternative explanation for the high white dwarf mass in CVs was proposed by [Schreiber et al. \(2016\)](#). The authors put forward an empirical consequential angular momentum loss (eCAML) model, which produces a dynamical stability limit on q , causing systems with low-mass white dwarfs to become unstable to mass transfer. These systems consequently merge, removing them from the CV population. The eCAML model is attractive as it can simultaneously explain the low observed space density of CVs ([Belloni et al. 2018](#)) and the origin of isolated low-mass white dwarfs ([Zorotovic & Schreiber 2017](#)).

The top-left plot of Figure 2 in [Schreiber et al. \(2016\)](#)

Parameter	SDSS 1006	SDSS 1152	SDSS 1501	SSS100615
q	0.46 ± 0.03	$0.153^{+0.015}_{-0.011}$	0.084 ± 0.004	0.095 ± 0.004
M_1 (M_\odot)	0.82 ± 0.11	0.62 ± 0.04	$0.723^{+0.017}_{-0.013}$	0.88 ± 0.03
R_1 (R_\odot)	0.0102 ± 0.0013	0.0129 ± 0.0006	$0.01142^{+0.00016}_{-0.00022}$	0.0095 ± 0.0003
M_2 (M_\odot)	0.37 ± 0.06	$0.094^{+0.016}_{-0.009}$	0.061 ± 0.004	0.083 ± 0.005
R_2 (R_\odot)	$0.457^{+0.022}_{-0.026}$	0.147 ± 0.006	$0.1129^{+0.0025}_{-0.0016}$	$0.1276^{+0.0028}_{-0.0024}$
a (R_\odot)	1.46 ± 0.07	0.627 ± 0.014	0.574 ± 0.004	0.628 ± 0.007
K_1 (km s^{-1})	124 ± 9	62 ± 5	$39.5^{+2.2}_{-1.3}$	$46.5^{+2.2}_{-1.7}$
K_2 (km s^{-1})	270 ± 13	402 ± 7	468 ± 3	493 ± 5
i ($^\circ$)	$83.1^{+1.2}_{-0.7}$	82.6 ± 0.5	$83.89^{+0.20}_{-0.27}$	85.1 ± 0.3
$\log g$	8.33 ± 0.13	8.01 ± 0.05	$8.182^{+0.016}_{-0.019}$	8.43 ± 0.03
T_1 (K)	–	15900 ± 2000	13400 ± 1100 (2004) 14900 ± 1000 (2012)	13600 ± 1500
d (pc)	–	610 ± 80	400 ± 30 (2004) 338 ± 21 (2012)	350 ± 30

Parameter	SSS130413	V713 Cep	Z Cha
q	$0.169^{+0.011}_{-0.006}$	$0.246^{+0.006}_{-0.014}$	0.189 ± 0.004
M_1 (M_\odot)	0.84 ± 0.03	$0.703^{+0.012}_{-0.015}$	0.803 ± 0.014
R_1 (R_\odot)	$0.0102^{+0.0006}_{-0.0002}$	$0.01173^{+0.00020}_{-0.00015}$	0.01046 ± 0.00017
M_2 (M_\odot)	$0.140^{+0.012}_{-0.008}$	$0.176^{+0.007}_{-0.018}$	0.152 ± 0.005
R_2 (R_\odot)	0.163 ± 0.004	$0.208^{+0.002}_{-0.005}$	0.1820 ± 0.0020
a (R_\odot)	$0.680^{+0.007}_{-0.011}$	0.781 ± 0.006	0.734 ± 0.005
K_1 (km s^{-1})	75 ± 4	91^{+2}_{-5}	$78.4^{+1.4}_{-1.8}$
K_2 (km s^{-1})	443^{+3}_{-7}	$367.6^{+2.6}_{-2.3}$	$413.2^{+2.5}_{-2.0}$
i ($^\circ$)	82.5 ± 0.3	81.7 ± 0.3	80.44 ± 0.11
$\log g$	8.35 ± 0.04	$8.147^{+0.017}_{-0.014}$	8.304 ± 0.016
T_1 (K)	24000 ± 3000	17000^{+6000}_{-3000}	16300 ± 1400
d (pc)	240 ± 40	320 ± 30	103 ± 6

Table 2. Continued.

was updated to take into account the results of this work (Figure 5). This plot is in M_2 vs q parameter space, with regions (grey) that are theoretically prohibited due to constraints put on M_1 . The dark grey prohibited region in the bottom right of Figure 5 is an upper mass limit on M_1 , resulting from the Chandrasekhar mass limit of a white dwarf ($1.44 M_\odot$). The light grey prohibited region is a lower mass limit on M_1 and is a consequence of the dynamical stability limit on q supplied by the eCAML model. Also plotted in Figure 5 are systems with measured M_2 and q , either from this work (green points) or elsewhere (black/blue points; see Table B1). These systems with measured system parameters provide a test of the eCAML model, as all should lie within the valid region (white). Any systems lying inside the prohibited dynamically unstable region would compromise the credibility of the model.

All systems modelled in this work lie comfortably within the valid region of Figure 5, along with the vast majority of other systems. Two appear to (just) violate the dynamical instability constraint, namely SDSS 0756+0858

(Tovmassian et al. 2014) and DQ Her (Horne et al. 1993), however both systems could feasibly be stable under the eCAML model after taking into account their uncertainties. This outcome offers support to the validity of the eCAML model as a solution to the CV white dwarf mass problem, however a much larger sample of systems with precise system parameters is necessary in order to provide a more stringent test of the model.

5.2 Reviewing the Properties of the Period Spike

The period spike is a feature of the orbital period distribution which is expected to occur as systems “pile-up” near the orbital period minimum due to the long evolutionary timescale. It was finally observed by Gänsicke et al. (2009) through analysing the orbital period distribution of newly identified CVs from SDSS (York et al. 2000). These systems were all identified spectroscopically (e.g. Szkody et al. 2002), and therefore not affected by the same biases/limitations as systems discovered through other means, e.g. DN outbursts




Figure 5. q vs M_2 plot for CVs. The grey regions are theoretically prohibited due to constraints put on M_1 . The dark grey regions cover unrealistically low white dwarf masses ($\lesssim 0.15 M_\odot$) and masses greater than the Chandrasekhar mass limit ($1.44 M_\odot$), while the light grey region is forbidden by the empirical consequential angular momentum loss (eCAML) model of Schreiber et al. (2016). The dashed grey line represents the mean value of M_1 from this work. The green and black points represent masses obtained from eclipse modelling of ULTRACAM/ULTRASPEC data, either from this work (green) or otherwise (black). The faint blue points represent measured CV masses from other methods: eclipse modelling of other data (circles), contact phase timing (squares), and radial velocity (triangles).

and X-ray emission (see Gänsicke et al. 2009 for more details). Spectroscopic identification, coupled with a survey depth of $g' \sim 19.5$, gives this particular sample the ability to provide the closest representation of the true orbital period distribution of CVs to date, a claim supported by the emergence of the long predicted-but-elusive period spike at the period minimum. Gänsicke et al. (2009) produced estimates for the location (82.4 ± 0.7 min) and width (FWHM = 5.7 min) of the period spike. Eight years on, the sample has increased and more P_{orb} measurements have become available, enabling the orbital period distribution – and in particular the properties of the period spike – to be reviewed.

The Gänsicke et al. (2009) sample consisted of 49 spectroscopically identified SDSS CVs below the period gap ($P_{\text{orb}} \lesssim 129$ min; Knigge 2006) with precise P_{orb} measurements (errors < 30 s). Precise P_{orb} measurements for an additional 23 systems (and updated measurements for a handful from the original sample) have since become available, increasing the sample to 72 systems. Of the new systems, six are eclipsing systems with observations using ULTRACAM/ULTRASPEC, 10 are from Thorstensen et al. (2015, 2017) and the remaining seven are from the Ritter & Kolb (2003) catalogue (v7.24; see references within). All systems were discovered by the SDSS (e.g. Szkody et al. 2011) except two, PHL 1445 and CSS110113, which were discovered by the 6dF Galaxy Survey (6dFGS; Jones et al. 2004).

Figure 6 shows the orbital period distribution of all 72 spectroscopically identified CVs in the form of both a histogram (red) and cumulative plot (blue). As with the Gänsicke et al. (2009) sample (dark red histogram), the new sample shows a clear accumulation of systems centred around ~ 82 min, which is clearly identifiable as the period spike. Estimating P_{spike} involved the fitting of a




Figure 6. Histogram (red) and cumulative plot (blue) for 72 spectroscopically identified (from SDSS and 6dFGS) CVs below the period gap with precise P_{orb} measurements (sub-30 s errors). For comparison, the sample of Gänsicke et al. (2009) is also shown (dark red histogram), in addition to the position and FWHM of the period spike estimated in the same study (black bar).

Gaussian distribution to the orbital period distribution between 77 and 87 min. An estimate of $P_{\text{spike}} = 82.7 \pm 0.4$ min ($\sigma = 2.35$ min, FWHM = 5.53 min) was obtained, which is largely unchanged from the Gänsicke et al. (2009) sample. This is not surprising, as the majority ($\sim 75\%$) of additional systems have $P_{\text{orb}} > 89$ min, and therefore do not belong to the period spike.

We note here that there is a hint of bi-modality in the period distribution of systems *below* the period gap, with a dearth of systems with orbital periods around 88 minutes. A Hartigan dip test (Hartigan & Hartigan 1985) reveals that this is not statistically significant.

5.3 Updating the Calibration of the superhump period excess-mass ratio Relation

During superoutburst the accretion disc is driven into an elliptical state by resonances between the donor star and material within the disc. Tidal interactions between the elliptical disc and the donor lead to periodic fluctuations in the elliptical, precessing, disc known as superhumps. The disc precesses at a slow rate, with a period (P_{prec}) significantly longer than P_{orb} . These two periods therefore both contribute to the formation of the superhump period (P_{sh}), which is simply the ‘beat period’ of P_{prec} and P_{orb} (Hellier 2001):

$$\frac{1}{P_{\text{sh}}} = \frac{1}{P_{\text{orb}}} - \frac{1}{P_{\text{prec}}}. \quad (1)$$

P_{sh} is therefore usually a few percent longer than P_{orb} , but does not stay constant throughout the superoutburst. In fact, a superoutburst can be split up into three distinct stages (A, B and C), with sharp transitions observed between each stage. Stage A represents the start of the superoutburst, with a long, stable P_{sh} . Stage B is the middle part of the superoutburst, with a shorter, unstable P_{sh} . The final stage (C) exhibits the shortest P_{sh} , which is stable once again (Olech et al. 2003; Kato et al. 2009). The general trend of decreasing P_{sh} across the superoutburst hints at an increas-

ing P_{prec} (from equation 1) and therefore a dwindling disc radius (Murray 2000).

The superhump excess (ϵ) is defined as $\epsilon = \frac{P_{\text{sh}} - P_{\text{orb}}}{P_{\text{orb}}}$, and is directly related to the mass ratio, q . A calibration of this relationship (e.g. Patterson et al. 2005; Knigge 2006) allows estimates of mass ratios for all superhumping systems. From this current work and the work of others (e.g. Savoury et al. 2011), new potential calibration systems have emerged, in addition to revised q values for existing calibration systems. Revised superhump periods have also been measured, courtesy of the SU UMa-type DNe survey of Kato et al. (2009, 2010, 2012, 2013, 2014a,b, 2015, 2016, 2017). With all of these new measurements becoming available since the work of Knigge (2006), it is appropriate to update the calibration of the $\epsilon(q)$ relation.

Table 3 contains all of the calibrating systems currently available⁴, along with their orbital and superhump periods (and references). The two superhump period columns, P_{sh}^{B} and P_{sh}^{C} , represent the superhump periods during stage B and stage C of superoutburst, respectively. All but the final four systems in Table 3 are SU UMa-type DNe that undergo superoutbursts. The other four systems are either Classical Novae or Novalikes that display permanent superhumps, and it is assumed these superhump periods resemble those of P_{sh}^{B} for SU UMa-type DNe.

For each system in Table 3, the superhump period excess was calculated for stage B (ϵ_{B}) and stage C (ϵ_{C}) depending on $P_{\text{sh}}^{\text{B}}/P_{\text{sh}}^{\text{C}}$ availability. Figure 7 shows ϵ_{B} plotted against q for the 24 calibration systems from Table 3 with available P_{sh}^{B} measurements. The dashed grey line shows the existing calibration from Knigge (2006), while the red line represents the following, updated linear calibration:

$$q(\epsilon_{\text{B}}) = (0.118 \pm 0.003) + (4.45 \pm 0.28) \times (\epsilon_{\text{B}} - 0.025). \quad (2)$$

This updated calibration was obtained through the same χ^2 minimisation technique employed by Knigge (2006) (see Appendix A of reference), and has an intrinsic dispersion (σ) of 0.012. While there is good coverage for systems with $0.1 < q < 0.2$, more calibration systems with q outside this range are required in order to further constrain the gradient. For example, due to its position in Figure 7, SDSS 1702 ($q \approx 0.25$) has a rather large influence on the gradient, so therefore more systems with precisely measured values of q greater than 0.2 are highly coveted. Unfortunately, this includes period gap systems, which are rare, and systems above the gap, for which precise measurements of q are hard to obtain. It is clear from Figure 7 that the new calibration has a steeper gradient than the existing one from Knigge (2006). A possible reason for this is the variation in measurement of P_{sh} between Patterson et al. (2005) and Kato et al. (2009); the sources of P_{sh} for both the existing and new calibration, respectively. Patterson et al. (2005) measures P_{sh} from ‘common’ superhumps, which typically cover stage B, but can also cover only a fraction of this stage or spread into stages A and C.

The same treatment was given to the 15 calibration systems in Table 3 with available P_{sh}^{C} measurements, producing

⁴ The calibration system KV UMa used by Knigge (2006) was not included on the basis of it being a low-mass X-ray binary, rather than a CV.




Figure 7. Measured ϵ_{B} and q values of superhumping and eclipsing CVs, with the same data point colour/shape scheme as Figure 5. The dashed grey line shows the existing linear calibration of the $\epsilon(q)$ relation for superhumping CVs from Knigge (2006), while the red line shows an updated calibration from this work. The red shaded region represents 1σ errors. The top plot shows the relationship for stage B superhumps, the bottom plot that for stage C superhumps.

the following linear relation (with $\sigma = 0.012$ again inferred):

$$q(\epsilon_{\text{C}}) = (0.135 \pm 0.004) + (5.0 \pm 0.7) \times (\epsilon_{\text{C}} - 0.025). \quad (3)$$

This relation is also shown in Figure 7.

5.4 Donor Masses and Radii of Superhumping CVs

Given our updating of the superhump-mass ratio relations above, we revisit the analysis of donor star properties in Knigge (2006) and Knigge et al. (2011). Firstly, P_{sh} values for all SU UMa-type DNe in the Patterson et al. (2005) sample (70 systems) were replaced by P_{sh}^{B} measurements from the SU UMa-type DNe survey of Kato et al. (2009, 2010, 2012, 2013, 2014a,b, 2015, 2016, 2017). For a number of systems, P_{orb} was also updated, either from measurements made by Kato et al., or additional studies (see references within Kato et al.). Values of ϵ_{B} were obtained from P_{sh}^{B} and P_{orb} , then subsequently converted into q via the newly calibrated $\epsilon_{\text{B}}(q)$ relation (equation 2). Equation 2 was also used to determine q for the eight systems displaying permanent superhumps. Assuming a constant white dwarf mass of

System	P_{orb} (d)	P_{sh}^{B} (d)	P_{sh}^{C} (d)	Ref.(s)
SDSS 1507	0.04625828(4)	0.046825(4)	–	1,2
SSS100615*	0.0587045(4) [§]	0.05972(9)	–	3
SDSS 1502	0.05890961(5)	0.060463(13)	0.060145(19)	1,4
SDSS 0903	0.059073543(9)	0.06036(5)	0.06007(5)	1,4
ASASSN-14ag	0.060310665(9) [§]	0.06206(6)	–	5
XZ Eri	0.061159491(5)	0.062807(18)	0.06265(12)	1,6
SDSS 1227	0.062959041(7)	0.064604(29)	0.06440(5)	1,7
OY Car*	0.06312092545(24) [§]	0.064653(28)	0.06444(5)	8
SSS130413*	0.0657692903(12) [§]	–	0.06751(24)	5
CSS110113*	0.0660508707(18) [§]	0.067583(26)	0.06731(4)	7
SDSS 1152*	0.0677497026(3) [§]	0.07036(4)	0.069914(19)	8
OU Vir	0.072706113(5)	0.074912(17)	–	1,6
IY UMa*	0.07390892818(21) [§]	0.076210(25)	0.075729(19)	4
Z Cha*	0.0744992631(3) [§]	0.07736(8)	0.076948(23)	5
SDSS 0901*	0.0778805321(5) [§]	0.08109(5)	0.08072(10)	9
DV UMa*	0.0858526308(7) [§]	0.08880(3)	0.08841(3)	6
SDSS 1702	0.10008209(9)	0.10507(8)	–	1,6
<hr/>				
WZ Sge	0.0566878460(3)	0.057204(5)	–	6,10
V2051 Oph	0.06242785751(8) [§]	0.06471(9)	0.06414(4)	5,11
HT Cas	0.0736471745(5) [§]	0.076333(5)	0.075886(5)	3,12
V4140 Sgr	0.0614296779(9)	0.06351(4)	0.06309(7)	6,11
<hr/>				
V348 Pup	0.101838931(14)	0.108567(2) [†]	–	13
V603 Aql	0.13820103(8)	0.14686(7) [†]	–	14,15
DW UMa	0.136606499(3)	0.14539(13) [†]	–	16,17
UU Aqr	0.1638049430	0.17510(18) [†]	–	18,19

Table 3. Orbital (P_{orb}) and superhump (P_{sh}) periods of the systems used to calibrate the $\epsilon(q)$ relation. The majority of systems are SU UMa-type DNe, however the bottom four are CNe/NLs. P_{sh}^{B} and P_{sh}^{C} are the periods for stage B and C superhumps, respectively. See Tables 2 and B1 for q values. References: (1) Savoury et al. (2011), (2) Patterson et al. (2017), (3) Kato et al. (2016), (4) Kato et al. (2010), (5) Kato et al. (2015), (6) Kato et al. (2009), (7) Kato et al. (2012), (8) Kato et al. (2017), (9) Kato et al. (2013), (10) Patterson (1998), (11) Baptista et al. (2003), (12) Horne et al. (1991), (13) Rolfe et al. (2000), (14) Peters & Thorstensen (2006), (15) Patterson et al. (1997), (16) Araujo-Betancor et al. (2003), (17) Patterson et al. (2002), (18) Baptista & Bortoletto (2008), (19) Patterson et al. (2005).

*Updated q value produced in this work (Table 2), [†]Superhump period from permanent superhumps, [§] P_{orb} from this work

(M_1) = 0.81 M_{\odot} , donor mass estimates were obtained for all systems in the superhumper sample.

As the donor fills its Roche lobe, the Eggleton (1983) approximation for the volume-averaged Roche lobe size, combined with Kepler’s 3rd law, can be used to obtain estimates for donor radii from q , M_2 and P_{orb} :

$$\frac{R_2}{R_{\odot}} = 0.2478 \left(\frac{M_2}{M_{\odot}} \right)^{1/3} P_{\text{orb}}^{2/3} \left[\frac{q^{1/3}(1+q)^{1/3}}{0.6q^{2/3} + \ln(1+q^{1/3})} \right], \quad (4)$$

where P_{orb} is in units of hrs. The Eggleton (1983) approximation for the volume-averaged Roche lobe size is the same one used to determine R_2 for systems that have been eclipse modelled, establishing consistency between the superhumping and eclipsing samples. It is important to note that Knigge et al. (2011) use a more complex, accurate approximation for the volume-averaged size of the Roche lobe based on the results of Sirotkin & Kim (2009), which represents the donor as a polytrope, rather than a point source. However, the advantage of using the Sirotkin & Kim (2009) approximation is small, with only a $\sim 1\%$ difference between the two approximations (Figure 3 of Knigge et al. 2011).

In addition to the 78 superhumper sample from Patterson et al. (2005), Kato et al. provide $P_{\text{sh}}^{\text{B,C}}$ and P_{orb} values for a further 147 systems. These systems were given

the same treatment as the Patterson et al. (2005) sample (outlined above). A handful of systems only have available P_{sh}^{C} values, in which case equation 3 was used. This brings the total number of superhumping systems with inferred donor properties to 225.

5.5 Updating the Semi-Empirical Mass-Radius Relation for CV Donor Stars

With donor masses and radii for 15 eclipsing systems in this work, a further 31 (mostly) eclipsing systems from the literature (see Table B1) and 225 superhumpers, it is possible to update the mass-radius relation for CV donor stars from Knigge (2006) and Knigge et al. (2011). The same fitting procedure used by Knigge (2006) was followed to update the mass-radius relation. Assumptions for some parameters in this model are required, since they are not well-constrained by the donor masses and radii. Assumptions for the donor mass within the period gap (M_{conv}), and the upper and lower ($P_{\text{gap,+}}$, $P_{\text{gap,-}}$) bounds of the period gap from Knigge et al. (2011) remained unchanged. We do adopt a smaller value for P_{bounce} (called P_{min} in Knigge et al. (2011)). P_{bounce} is the orbital period where the pre-bounce and post-bounce

power-law relationships intersect. Knigge et al. (2011) used the location of the period spike from Gänsicke et al. (2009) for P_{bounce} . However, real systems do not reach this orbital period, because the smooth track followed by real systems near period minimum is not well represented by two power laws. PHL 1445 (McAllister et al. 2015) is expected to be close to the absolute minimum period for main sequence CVs, and so its orbital period of 76.3 min is used for P_{bounce} here. The value of M_{bounce} shown above was determined from the optimal short-period fit.

$$\begin{aligned} M_{\text{bounce}} &= 0.063^{+0.005}_{-0.002} M_{\odot}, & P_{\text{bounce}} &= 76.3 \pm 1.0 \text{ min} \\ M_{\text{conv}} &= 0.20 \pm 0.02 M_{\odot}, & P_{\text{gap},-} &= 2.15 \pm 0.03 \text{ hrs}, \\ M_{\text{evol}} &\approx 0.6\text{--}0.8 M_{\odot}, & P_{\text{gap},+} &= 3.18 \pm 0.04 \text{ hrs}. \end{aligned}$$

The donor masses and radii for all but 12 systems were included in the fits. The majority of these systems were excluded due to being period gap systems (see grey box in bottom plot of Figure 8), while SDSS 1507 (outlying black data point in period bouncer regime) was excluded as it is known to be a Galactic halo object (Patterson et al. 2008; Uthas et al. 2011). The results from the three power law fits are shown in Figure 8, and take the following form:

$$\frac{R_2}{R_{\odot}} = \begin{cases} 0.109 \pm 0.003 \left(\frac{M_2}{M_{\text{bounce}}} \right)^{0.152 \pm 0.018} & M_2 < M_{\text{bounce}} \\ 0.225 \pm 0.008 \left(\frac{M_2}{M_{\text{conv}}} \right)^{0.636 \pm 0.012} & M_{\text{bounce}} < M_2 < M_{\text{conv}} \\ 0.293 \pm 0.010 \left(\frac{M_2}{M_{\text{evol}}} \right)^{0.69 \pm 0.05} & M_{\text{conv}} < M_2 < M_{\text{evol}}. \end{cases}$$

Comparing these results with Knigge et al. (2011), there is little change in the exponents of the mass-radius relation in both the long- and short-period regimes. One notable difference, however, is the amount of intrinsic scatter, σ_{int} , required for the short-period systems, reduced from approximately $0.02 R_{\odot}$ to $0.005 R_{\odot}$. The small scatter provides strong evidence for a very tight evolutionary path followed by non-evolved CV donors, implying little spread in AML loss rates for CVs with the same component masses. The scatter within the long-period regime, at $0.04 R_{\odot}$, is almost a factor of 10 larger than that at short periods. Figure 8 shows two outlying long-period systems with $R_2 \approx 0.40 R_{\odot}$, namely IP Peg (Copperwheat et al. 2010) and HS 0220+0603 (Rodríguez-Gil et al. 2015). The donors within these two systems are undersized for their masses, and may even be in thermal equilibrium, which is unexpected for a CV donor. It is possible that both IP Peg and HS 0220+0603 have donors in thermal equilibrium due to recently starting mass transfer.

The mass-radius relation for period-bouncers has changed significantly. The new power law exponent of 0.152 ± 0.018 is much smaller than that of Knigge et al. (2011), a consequence of using lower values for both M_{bounce} and P_{bounce} , in addition to the inclusion of many more period-bouncers in the new donor sample, which enables a better constraint of the power law in this regime. There has been a long-standing issue with the number of confirmed period-bounce CVs, which has always been much lower than the predicted 40–70% (Goliašch & Nelson 2015; Kolb 1993). Whilst the sample of donor masses collected here is far from homogeneous, and the presence of large numbers of superhumping systems introduces complicated selection effects,

we note here that 30% of our sample has a donor mass below $0.063 M_{\odot}$ and are therefore likely to be period-bouncers.

5.6 Comparison to theoretical CV evolution tracks

In addition to a broken-power-law mass-radius relation for CV donors, Knigge et al. (2011) present a theoretical evolutionary track, produced with the aim of quantifying the secular mass transfer rate in CVs. The track which best reproduces their donor sample requires reduced magnetic braking above the gap ($f_{\text{MB}} = 0.66 \pm 0.05$), but additional angular momentum loss below the gap ($f_{\text{GR}} = 2.47 \pm 0.22$). The donor sample presented in this work is shown in the M_2 - R_2 and P_{orb} - M_2 planes in Figures 9 and 10, respectively. Also shown is the ‘best fit’ track from Knigge et al. (2011), and the ‘standard’ track ($f_{\text{GR}} = f_{\text{MB}} = 1$). It is clear from these figures that the best-fit evolutionary track from Knigge et al. (2011) under predicts the donor mass at orbital periods below the period gap, and has a period minimum that is longer than that observed. This again implies that less additional angular momentum loss is needed below the period gap than suggested by Knigge et al. (2011). In contrast, we find that the ‘standard’ track provides a better fit to the donor sample immediately below the gap, where the donor mass is in the range $0.10\text{--}0.20 M_{\odot}$. This is most apparent in Figure 10. Although the standard track is a good fit to systems immediately below the gap, it diverges from the donor sequence at lower masses, and predicts a period minimum shorter than the observed value. Therefore, the donor properties in CVs appears to argue for an additional source of AML that is small compared to gravitational radiation just below the period gap, but becomes more significant at shorter orbital periods and/or donor masses.

The eCAML model of Schreiber et al. (2016) might provide something similar to the behaviour required. All models of CV evolution require a term ν , which expresses the AML which arises as a consequence of mass transfer. In the standard model, it is assumed that the mass lost from the white dwarf during nova eruptions carries with it the specific angular momentum of the white dwarf, leading to $\nu = M_2^2/(M_1 M)$, where M is the total mass of the system. In the eCAML model, an alternative form of $\nu \sim 0.35/M_1$ is proposed. We used equation 1 from Knigge et al. (2011) to roughly estimate the mass loss rates under the eCAML model at key points in the evolution of the donor. Just below the period gap, we take $M_1 = 0.82 M_{\odot}$, $M_2 = 0.15 M_{\odot}$ and we assume the donor is roughly in thermal equilibrium, so the mass-radius index is $\xi = 0.8$. This implies that in the eCAML model, mass loss rates just after the period gap are only around 35% higher than the ‘standard’, $f_{\text{GR}} = 1$, model. For systems near the period minimum, we take $M_1 = 0.82 M_{\odot}$, $M_2 = 0.065 M_{\odot}$ and $\xi = -1/3$, which suggests mass loss rates around 9 times higher than the $f_{\text{GR}} = 1$ case. Therefore, the eCAML model provides a mass loss law which is qualitatively similar to the one implied by CV donor properties. However, it is worth bearing in mind that the $\nu \sim 0.35/M_1$ prescription is not physically motivated. Schreiber et al. (2016) suggest that angular momentum loss during nova outbursts might produce a similar behaviour, but the frequency of nova outbursts will drop as the accretion rate falls. Therefore eCAML may be less important for CVs near the period minimum




Figure 8. Measured CV donor masses (M_2) and radii (R_2). The data point colour/shape scheme is the same as in Figure 5, but with additional superhumping systems (grey points), for which error bars have been omitted for clarity. The red line is the semi-empirical mass-radius relation from this work. The grey shaded region contains systems assumed to lie within the period gap, and are therefore not included in the updated broken-power-law fit.

than implied above. It will require a physically plausible model of CV evolution, including AML during nova outbursts, to determine if such a model can reproduce both the high white dwarf mass in CVs and the $P_{\text{orb}}-M_2$ locus of the donor stars.

Finally, we note that our results introduce a tension between the donor masses and radii, and the temperatures of white dwarfs in CVs. As described in Townsley & Gänsicke (2009), compressional heating of the white dwarfs due to accretion sets the equilibrium temperature of the white dwarf in a CV. The observed white dwarf temperature thus depends upon the accretion rate, averaged over the thermal timescale of the non-degenerate layer on the white dwarf surface (Townsley & Bildsten 2003). The best study of white dwarf temperatures in CVs to date is Pala et al. (2017), who show that the white dwarfs in CVs below the period gap imply AML rates approximately twice that implied by $f_{\text{GR}} = 1$. As discussed extensively in Section 4 of Knigge et al. (2011), one plausible explanation for the discrepancy is the presence of mass transfer rate fluctuations, coupled with the fact that the white dwarf temperature reflects the mass transfer rate averaged over much shorter timescales than the donor star radius. However, this would presumably lead to white dwarf temperatures scattered around the expected values; whereas they are systematically warmer than expected.

5.7 The Period Minimum

It is apparent from Figure 10 that the current donor sample contains a sufficiently large number of systems at the short-

est orbital periods to finally begin to reveal the locus of CVs evolving through the period minimum. The period minimum of the current donor sample covers an approximate period range of 76–82 min (1.27–1.37 hrs). Fitting a Gaussian distribution to the donor sample within this period range returned the following estimates for both the period minimum ($P_{\text{min}} = 79.6 \pm 0.2$ min) and its width (FWHM = 4.0 min). These estimates for P_{min} and its width are shown by the red vertical dashed line and shaded area within Figure 10.

It was briefly mentioned in Section 5.5 that the observed location of the period minimum appears to be slightly lower than the value $P_{\text{min}} = 81.8 \pm 0.9$ min predicted by the best-fit track of Knigge et al. (2011). The new measurement of P_{min} from the donor sample confirms this, with the two P_{min} estimates differing by approximately 2.4σ . A lower value of P_{min} than the existing estimate of Knigge et al. (2011) was previously hinted at by McAllister et al. (2015). Figure 10 shows that with the new estimate for P_{min} , PHL 1445 ($P_{\text{orb}} = 1.27$ hrs) and SDSS 1433 ($P_{\text{orb}} = 1.30$ hrs) are no longer troublesome outliers.

6 CONCLUSIONS

We present new measurements of the system parameters for 15 eclipsing CVs, six of which are published for the first time. We also compile a list of reliable system parameter determinations from the literature. We use these measurements to refine the calibration of the relationship between superhump period excess and mass ratio; allowing us to es-




Figure 9. Measured CV donor masses (M_2) and radii (R_2). The data point colour/shape scheme is the same as in Figure 8. The red and black lines represent the best-fit ($f_{\text{GR}} = 2.47 \pm 0.22$, $f_{\text{MB}} = 0.66 \pm 0.05$) and ‘standard’ ($f_{\text{GR}} = f_{\text{MB}} = 1$) evolutionary tracks from Knigge et al. (2011), respectively.

estimate the donor properties of 225 CVs showing superhump phenomena. This provides an extensive sample of CVs with known system parameters which we can use to test models of CV evolution.

We confirm the high average white dwarf mass in CVs, but we find no evidence for a trend in white dwarf mass with orbital period. Contrary to previous studies, we find that the donor properties of CVs immediately below the period gap are consistent with the standard model, in which AML due to magnetic braking is small compared to gravitational radiation. We do, however, find that CVs at shorter orbital periods and lower masses still require an additional source of AML. We argue that the eCAML model of Schreiber et al. (2016) predicts an AML law that is qualitatively similar to this behaviour. We find that, for systems below the period gap, donor radii at a given orbital period show a very small intrinsic scatter of only $0.005 R_{\odot}$, suggesting that most CVs below the gap follow a common evolutionary path. We estimate a value for the orbital period minimum of 79.6 ± 0.2 min, shorter than previously estimated by Knigge et al. (2011).

The CVs with donor properties estimated from superhumps show a sizeable fraction of systems which appear to have evolved past the period minimum. As a result, 30% of our sample appear to be post-period minimum systems. This hints that post-period minimum systems may be as common as models predict, but the superhump sample is strongly biased towards low mass ratios. The advent of Gaia means that detailed follow up of a relatively complete

volume-limited sample may resolve this question in the near future.

ACKNOWLEDGEMENTS

VSD, SPL, ULTRACAM and ULTRASPEC are supported by the STFC grant ST/R000964/1. This work has made use of data obtained at the Thai National Observatory on Doi Inthanon, operated by NARIT. VSD, TRM and SPL acknowledge the support of the Royal Society and the Leverhulme Trust for the operation of ULTRASPEC at the TNT. The WHT is operated on the island of La Palma by the Isaac Newton Group of Telescopes in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias. This work is based on observations collected at the European Organisation for Astronomical Research in the Southern Hemisphere.




Figure 10. Measured CV donor masses (M_2) as a function of orbital period (P_{orb}). The data point colour/shape scheme is the same as in Figure 8. The red and black lines represent the best-fit ($f_{\text{GR}} = 2.47 \pm 0.22$, $f_{\text{MB}} = 0.66 \pm 0.05$) and ‘standard’ ($f_{\text{GR}} = f_{\text{MB}} = 1$) evolutionary tracks from Knigge et al. (2011), respectively. The vertical dashed red line and shaded region is an estimate of the true P_{min} based on fitting a Gaussian distribution to the orbital periods in the range 76–82 mins.

REFERENCES

- Applegate J. H., 1992, *ApJ*, **385**, 621
- Araujo-Betancor S., et al., 2003, *ApJ*, **583**, 437
- Arenas J., Catalán M. S., Augusteijn T., Retter A., 2000, *MNRAS*, **311**, 135
- Arnold S., Berg R. A., Duthie J. G., 1976, *ApJ*, **206**, 790
- Bailer-Jones C. A. L., Rybizki J., Fouesneau M., Mantelet G., Andrae R., 2018, preprint, p. [arXiv:1804.10121](https://arxiv.org/abs/1804.10121) ([arXiv:1804.10121](https://arxiv.org/abs/1804.10121))
- Baptista R., Bortoletto A., 2008, *ApJ*, **676**, 1240
- Baptista R., Steiner J. E., Cieslinski D., 1994, *ApJ*, **433**, 332
- Baptista R., Borges B. W., Bond H. E., Jablonski F., Steiner J. E., Grauer A. D., 2003, *MNRAS*, **345**, 889
- Belloni D., Schreiber M. R., Zorotovic M., Ilkiewicz K., Hurley J. R., Giersz M., Lagos F., 2018, *MNRAS*, p. 1379
- Bergeron P., Wesemael F., Beauchamp A., 1995, *PASP*, **107**, 1047
- Borges B. W., Baptista R., 2005, *A&A*, **437**, 235
- Bours M. C. P., et al., 2016, *MNRAS*, **460**, 3873
- Copperwheat C. M., Marsh T. R., Dhillon V. S., Littlefair S. P., Hickman R., Gänsicke B. T., Southworth J., 2010, *MNRAS*, **402**, 1824
- Copperwheat C. M., et al., 2012, *MNRAS*, **421**, 149
- Dhillon V. S., et al., 2007, *MNRAS*, **378**, 825
- Dhillon V. S., et al., 2014, *MNRAS*, **444**, 4009
- Dillon M., et al., 2008, *MNRAS*, **386**, 1568
- Echevarría J., de la Fuente E., Costero R., 2007, *AJ*, **134**, 262
- Echevarría J., Ramírez-Torres A., Michel R., Hernández Santisteban J. V., 2016, *MNRAS*, **461**, 1576
- Eggleton P. P., 1983, *ApJ*, **268**, 368
- Gänsicke B. T., Long K. S., Barstow M. A., Hubeny I., 2006, *ApJ*, **639**, 1039
- Gänsicke B. T., et al., 2009, *MNRAS*, **397**, 2170
- Goliasch J., Nelson L., 2015, *ApJ*, **809**, 80
- Hardy L. K., et al., 2017, *MNRAS*, **465**, 4968
- Hartigan J. A., Hartigan P., 1985, *Ann. Statist.*, **13**, 70
- Hellier C., 2001, *Cataclysmic Variable Stars: How and Why they Vary*. Springer-Praxis, New York
- Hernández Santisteban J. V., Knigge C., Pretorius M. L., Sullivan M., Warner B., 2018, *MNRAS*, **473**, 3241
- Hernandez M. S., Zharikov S., Neustroev V., Tovmassian G., 2017, *MNRAS*, **470**, 1960
- Horne K., Wood J. H., Stiening R. F., 1991, *ApJ*, **378**, 271
- Horne K., Welsh W. F., Wade R. A., 1993, *ApJ*, **410**, 357
- Howell S. B., Warnock A., Mason K. O., Reichert G. A., Kreidl T. J., 1988, *MNRAS*, **233**, 79
- Jones D. H., et al., 2004, *MNRAS*, **355**, 747
- Kato T., et al., 2009, *PASJ*, **61**, S395
- Kato T., et al., 2010, *PASJ*, **62**, 1525
- Kato T., et al., 2012, *PASJ*, **64**, 21
- Kato T., et al., 2013, *PASJ*, **65**, 23
- Kato T., et al., 2014a, *PASJ*, **66**, 30
- Kato T., et al., 2014b, *PASJ*, **66**, 90
- Kato T., et al., 2015, *PASJ*, **67**, 105
- Kato T., et al., 2016, *PASJ*, **68**, 65
- Kato T., et al., 2017, *PASJ*, **69**, 75
- Knigge C., 2006, *MNRAS*, **373**, 484
- Knigge C., Baraffe I., Patterson J., 2011, *ApJS*, **194**, 28
- Kolb U., 1993, *A&A*, **271**, 149
- Lindgren L., et al., 2018, preprint, p. [arXiv:1804.09366](https://arxiv.org/abs/1804.09366) ([arXiv:1804.09366](https://arxiv.org/abs/1804.09366))
- Littlefair S. P., Dhillon V. S., Marsh T. R., Gänsicke B. T., Southworth J., Baraffe I., Watson C. A., Copperwheat C., 2008, *MNRAS*, **388**, 1582
- Littlefair S. P., Dhillon V. S., Gänsicke B. T., Bours M. C. P., Copperwheat C. M., Marsh T. R., 2014, *MNRAS*, **443**, 718
- Manser C. J., Gänsicke B. T., 2014, *MNRAS*, **442**, L23
- McAllister M. J., et al., 2015, *MNRAS*, **451**, 114
- McAllister M. J., et al., 2017a, *MNRAS*, **464**, 1353
- McAllister M. J., et al., 2017b, *MNRAS*, **467**, 1024
- Miszalski B., et al., 2016, *MNRAS*, **456**, 633
- Murray J. R., 2000, *MNRAS*, **314**, L1
- Nelemans G., Siess L., Repetto S., Toonen S., Phinney E. S., 2016, *ApJ*, **817**, 69
- Nogami D., Kato T., Baba H., Novák R., Lockley J. J., Somers M., 2001, *MNRAS*, **322**, 79
- Olech A., Schwarzenberg-Czerny A., Kedzierski P., Zloczewski K., Mularczyk K., Wisniewski M., 2003, *Acta Astron.*, **53**, 175
- Pala A. F., et al., 2017, *MNRAS*, **466**, 2855
- Patterson J., 1998, *PASP*, **110**, 1132
- Patterson J., Kemp J., Saad J., Skillman D. R., Harvey D., Fried R., Thorstensen J. R., Ashley R., 1997, *PASP*, **109**, 468
- Patterson J., et al., 2000, *PASP*, **112**, 1584
- Patterson J., et al., 2002, *PASP*, **114**, 1364
- Patterson J., et al., 2005, *PASP*, **117**, 1204
- Patterson J., Thorstensen J. R., Knigge C., 2008, *PASP*, **120**, 510
- Patterson J., et al., 2017, *Society for Astronomical Sciences Annual Symposium*, **37**, 1
- Peters C. S., Thorstensen J. R., 2006, *PASP*, **118**, 687

- Rappaport S., Joss P. C., Webbink R. F., 1982, *ApJ*, **254**, 616
- Rappaport S., Verbunt F., Joss P. C., 1983, *ApJ*, **275**, 713
- Ritter H., Kolb U., 2003, *A&A*, **404**, 301
- Robinson E. L., 1974, *ApJ*, **193**, 191
- Rodríguez-Gil P., Martínez-Pais I. G., Casares J., Villada M., van Zyl L., 2001, *MNRAS*, **328**, 903
- Rodríguez-Gil P., et al., 2015, *MNRAS*, **452**, 146
- Rolfe D. J., Haswell C. A., Patterson J., 2000, *MNRAS*, **317**, 759
- Savourey C. D. J., et al., 2011, *MNRAS*, **415**, 2025
- Savourey C. D. J., Littlefair S. P., Marsh T. R., Dhillon V. S., Parsons S. G., Copperwheat C. M., Steeghs D., 2012, *MNRAS*, **422**, 469
- Scaringì S., Groot P. J., Still M., 2013, *MNRAS*, **435**, L68
- Schreiber M. R., Zorotovic M., Wijnen T. P. G., 2016, *MNRAS*, **455**, L16
- Shafter A. W., Holland J. N., 2003, *PASP*, **115**, 1105
- Sing D. K., Green E. M., Howell S. B., Holberg J. B., Lopez-Morales M., Shaw J. S., Schmidt G. D., 2007, *A&A*, **474**, 951
- Sirotkin F. V., Kim W.-T., 2009, *ApJ*, **698**, 715
- Smith J. A., et al., 2002, *AJ*, **123**, 2121
- Smith A. J., Haswell C. A., Hynes R. I., 2006, *MNRAS*, **369**, 1537
- Southworth J., Marsh T. R., Gänsicke B. T., Aungwerojwit A., Hakala P., de Martino D., Lehto H., 2007, *MNRAS*, **382**, 1145
- Southworth J., Hickman R. D. G., Marsh T. R., Rebassa-Mansergas A., Gänsicke B. T., Copperwheat C. M., Rodríguez-Gil P., 2009, *A&A*, **507**, 929
- Southworth J., Copperwheat C. M., Gänsicke B. T., Pyrzas S., 2010, *A&A*, **510**, A100
- Steeeghs D., Marsh T., Knigge C., Maxted P. F. L., Kuulkers E., Skidmore W., 2001, *ApJ*, **562**, L145
- Steeeghs D., Perryman M. A. C., Reynolds A., de Bruijne J. H. J., Marsh T., Dhillon V. S., Peacock A., 2003, *MNRAS*, **339**, 810
- Steeeghs D., Howell S. B., Knigge C., Gänsicke B. T., Sion E. M., Welsh W. F., 2007, *ApJ*, **667**, 442
- Szkody P., Brownlee D. E., 1977, *ApJ*, **212**, L113
- Szkody P., et al., 2002, *AJ*, **123**, 430
- Szkody P., et al., 2011, *AJ*, **142**, 181
- Tappert C., Augusteijn T., Maza J., 2004, *MNRAS*, **354**, 321
- The Astropy Collaboration et al., 2018, preprint, p. [arXiv:1801.02634](https://arxiv.org/abs/1801.02634) ([arXiv:1801.02634](https://arxiv.org/abs/1801.02634))
- Thoroughgood T. D., Dhillon V. S., Watson C. A., Buckley D. A. H., Steeghs D., Stevenson M. J., 2004, *MNRAS*, **353**, 1135
- Thoroughgood T. D., et al., 2005, *MNRAS*, **357**, 881
- Thorstensen J. R., 2000, *PASP*, **112**, 1269
- Thorstensen J. R., Taylor C. J., Peters C. S., Skinner J. N., Southworth J., Gänsicke B. T., 2015, *AJ*, **149**, 128
- Thorstensen J. R., Alper E. H., Weil K. E., 2016, *AJ*, **152**, 226
- Thorstensen J. R., Ringwald F. A., Taylor C. J., Sheets H. A., Peters C. S., Skinner J. N., Alper E. H., Weil K. E., 2017, *Research Notes of the American Astronomical Society*, **1**, 29
- Tovmassian G., Stephania Hernandez M., González-Buitrago D., Zharikov S., García-Díaz M. T., 2014, *AJ*, **147**, 68
- Townsley D. M., Bildsten L., 2003, *ApJ*, **596**, L227
- Townsley D. M., Bildsten L., 2004, *ApJ*, **600**, 390
- Townsley D. M., Gänsicke B. T., 2009, *ApJ*, **693**, 1007
- Tulloch S. M., Rodríguez-Gil P., Dhillon V. S., 2009, *MNRAS*, **397**, L82
- Uthas H., Knigge C., Long K. S., Patterson J., Thorstensen J., 2011, *MNRAS*, **414**, L85
- Wade R. A., Horne K., 1988, *ApJ*, **324**, 411
- Welsh W. F., Froning C. S., Marsh T. R., Reimer T. W., Robinson E. L., Wood P. R., 2007, in Kang Y. W., Lee H.-W., Leung K.-C., Cheng K.-S., eds, *Astronomical Society of the Pacific Conference Series Vol. 362, The Seventh Pacific Rim Conference on Stellar Astrophysics*. p. 241
- Wood J. H., Horne K., 1990, *MNRAS*, **242**, 606
- Wood J., Horne K., Berriman G., Wade R., O'Donoghue D., Warner B., 1986, *MNRAS*, **219**, 629
- York D. G., et al., 2000, *AJ*, **120**, 1579
- Zorotovic M., Schreiber M. R., 2017, *MNRAS*, **466**, L63
- Zorotovic M., Schreiber M. R., Gänsicke B. T., 2011, *A&A*, **536**, A42
- van Amerongen S., Damen E., Groot M., Kraakman H., van Paradijs J., 1987, *MNRAS*, **225**, 93




Figure A1. Simultaneous eclipse model fit to six average CSS080623 eclipse light curves. See Section 4.1 for full details of what is plotted. Displayed in the top-right corner of each average eclipse plot is the date(s) and wavelength band each of the constituent eclipses were observed in.

APPENDIX A: FITS TO ECLIPSE LIGHTCURVES

The following figures show the full sets of eclipse model fits for the 15 additional systems from Section 4.1.

APPENDIX B: SYSTEM PARAMETERS FOR SUPPLEMENTARY SYSTEMS

The following table includes reliably determined system parameters for CVs from the literature.

APPENDIX C: JOURNAL OF OBSERVATIONS




Figure A2. Simultaneous eclipse model fit to five CTCV 1300 eclipse light curves. See Section 4.1 for full details of what is plotted. Displayed in the top-right corner of each eclipse plot is the cycle number of the eclipse and the wavelength band it was observed in.




Figure A3. Simultaneous eclipse model fit to three average CSS110113 eclipse light curves. See Section 4.1 for full details of what is plotted. Displayed in the top-right corner of each average eclipse plot is the date and wavelength band each of the constituent eclipses were observed in.

System	P_{orb} (d)	q	M_1 (M_{\odot})	M_2 (M_{\odot})	R_2 (R_{\odot})	Method	Ref.
SDSS J1433+1011	0.054240679(2)	0.0661(7)	0.865(5)	0.0571(7)	0.1074(4)	EM(U)	1
SDSS J1507+5230	0.04625828(4)	0.0647(18)	0.892(8)	0.0575(20)	0.0969(11)	EM(U)	1
SDSS J1035+0551	0.0570067(2)	0.0571(10)	0.835(9)	0.0475(12)	0.1047(8)	EM(U)	1
CTCV J2354–4700	0.065550270(1)	0.1097(8)	0.935(31)	0.101(3)	0.1463(16)	EM(U)	1
SDSS J1152+4049*	0.0677497026(3) [§]	0.155(6)	0.560(28)	0.087(6)	0.142(3)	EM(U)	1
SDSS J0903+3300	0.059073543(9)	0.113(4)	0.872(11)	0.099(4)	0.1358(20)	EM(U)	1
SDSS J1227+5139	0.062959041(7)	0.1115(16)	0.796(18)	0.0889(25)	0.1365(13)	EM(U)	1
XZ Eri	0.061159491(5)	0.118(3)	0.769(17)	0.091(4)	0.1350(18)	EM(U)	1
SDSS J1502+3334	0.05890961(5)	0.1099(7)	0.709(4)	0.0781(8)	0.1241(3)	EM(U)	1
SDSS J1501+5501*	0.05684126603(21) [§]	0.101(10)	0.767(27)	0.077(10)	0.122(5)	EM(U)	1
CTCV J1300–3052*	0.0889406998(17) [§]	0.240(21)	0.736(14)	0.177(21)	0.215(8)	EM(U)	1
OU Vir	0.072706113(5)	0.1641(13)	0.703(12)	0.1157(22)	0.1634(10)	EM(U)	1
DV UMa*	0.0858526308(7) [§]	0.1778(22)	1.098(24)	0.196(5)	0.2176(18)	EM(U)	1
SDSS J1702+3229	0.10008209(9)	0.248(5)	0.91(3)	0.223(10)	0.252(4)	EM(U)	1
PHL 1445	0.0529848884(13)	0.087(6)	0.73(3)	0.064(5)	0.109(4)	EM(U)	2
SDSS J1057+2759	0.0627919557(6)	0.0546(20)	0.800(15)	0.0436(20)	0.1086(17)	EM(U)	3
ASASSN-14ag	0.060310665(9)	0.149(16)	0.63(4)	0.093(13)	0.135(7)	EM(U)	4
KIS J1927+4447	0.165308(5)	0.570(11)	0.69(7)	0.39(4)	0.432(15)	EM(U)	5,6
IP Peg	0.1582061029(3)	0.48(1)	1.16(2)	0.55(2)	0.466(6)	EM(U)	7
EX Dra	0.20993718(2)	0.75(5)	0.71(4)	0.53(1)	0.565(4)	EM	8
SDSS J1006+2337*	0.185913107(13) [§]	0.51(8)	0.78(12)	0.40(10)	0.47(4)	EM	9
CSS111003 (Te 11)	0.120971471(9)	0.236(6)	1.18(11)	0.28(3)	0.314(11)	EM	10
HS 0220+0603	0.14920775(8)	0.54(3)	0.87(9)	0.47(5)	0.427(15)	EM	11
1RXS J0644+3344	0.26937431(22)	0.80(2)	0.73(7)	0.58(6)	0.690(24)	EM	12,13
SDSS J0756+0858	0.1369745(4)	0.47(9)	0.60(12)	0.28(5)	0.338(20)	EM	14

Table B1. System parameters for supplementary systems included in section 5 (Figures 5–10). The second-to-last column indicates the method used to obtain system parameters: EM – eclipse modelling (U – using ULTRACAM/ULTRASPEC data), CPT – contact phase timing, RV – radial velocity, GR – gravitational redshift, SM – spectrophotometric modelling. For consistency, all R_2 values were calculated using equation 4 (ensuring all systems follow the same period-density relation). References: (1) Savoury et al. (2011), (2) McAllister et al. (2015), (3) McAllister et al. (2017a), (4) McAllister et al. (2017b) (5) Scaringi et al. (2013), (6) Littlefair et al. (2014), (7) Copperwheat et al. (2010), (8) Shafter & Holland (2003), (9) Southworth et al. (2009), (10) Miszalski et al. (2016), (11) Rodríguez-Gil et al. (2015), (12) Sing et al. (2007), (13) Hernandez et al. (2017), (14) Tovmassian et al. (2014), (15) Steeghs et al. (2003), (16) Horne et al. (1991), (17) Wood & Horne (1990), (18) Littlefair et al. (2008), (19) Baptista et al. (2003), (20) Borges & Baptista (2005), (21) Araujo-Betancor et al. (2003), (22) Patterson et al. (2005), (23) Baptista & Bortoletto (2008), (24) Baptista et al. (1994), (25) Thorstensen (2000), (26) Wade & Horne (1988), (27) Echevarría et al. (2016), (28) Arnold et al. (1976), (29) Echevarría et al. (2007), (30) Horne et al. (1993), (31) Thoroughgood et al. (2005), (32) Rolfe et al. (2000), (33) Rodríguez-Gil et al. (2001), (34) Peters & Thorstensen (2006), (35) Arenas et al. (2000), (36) Robinson (1974), (37) Welsh et al. (2007), (38) Thoroughgood et al. (2004), (39) Patterson (1998), (40) Steeghs et al. (2001), (41) Steeghs et al. (2007), (42) van Amerongen et al. (1987), (43) Smith et al. (2006), (44) Szkody & Brownlee (1977), (45) Gänsicke et al. (2006).

*Updated system parameters produced in this work (Table 2), [§] P_{orb} from this work

System	P_{orb} (d)	q	M_1 (M_{\odot})	M_2 (M_{\odot})	R_2 (R_{\odot})	Method	Ref.
IY UMa*	0.07390892818(21) [§]	0.125(8)	0.79(4)	0.10(1)	0.158(5)	CPT	15
HT Cas	0.0736471745(5)	0.15(3)	0.61(4)	0.09(2)	0.152(11)	CPT	16
OY Car*	0.06312092545(24) [§]	0.102(3)	0.84(4)	0.086(5)	0.1354(26)	CPT	17,18
V2051 Oph	0.06242785751(8)	0.19(3)	0.78(6)	0.15(3)	0.161(11)	CPT	19
V4140 Sgr	0.0614296779(9)	0.125(15)	0.73(8)	0.092(16)	0.136(8)	CPT	19,20
DW UMa	0.136606499(3)	0.28(4)	0.73(3)	0.21(3)	0.304(14)	CPT	21,22
UU Aqr	0.1638049430	0.30(7)	0.67(14)	0.20(7)	0.34(4)	CPT	23,24
GY Cnc*	0.175442399(6) [§]	0.41(4)	0.82(14)	0.33(7)	0.42(3)	RV	25
Z Cha*	0.0744992631(3) [§]	0.149(4)	0.84(9)	0.125(14)	0.171(6)	RV	26
EX Hya	0.068233843(1)	0.13(2)	0.78(3)	0.10(2)	0.150(10)	RV	27
U Gem	0.17690617(6)	0.35(5)	1.20(5)	0.42(4)	0.456(14)	RV	28,29
DQ Her	0.193620897	0.66(4)	0.60(7)	0.40(5)	0.485(20)	RV	30
V347 Pup	0.231936060(6)	0.83(5)	0.63(4)	0.52(6)	0.603(23)	RV	31
V348 Pup	0.101838931(14)	0.31(6)	0.65(13)	0.20(4)	0.246(16)	RV	32,33
V603 Aql	0.13820103(8)	0.24(5)	1.2(2)	0.29(4)	0.341(16)	RV	34,35
EM Cyg	0.290909(4)	0.77(4)	1.00(12)	0.77(8)	0.797(28)	RV	36,37
AC Cnc	0.30047747(4)	1.02(4)	0.76(3)	0.77(5)	0.827(18)	RV	38
V363 Aur	0.32124187(8)	1.17(7)	0.90(6)	1.06(11)	0.97(4)	RV	38
WZ Sge	0.0566878460(3)	0.057(18)	0.85(4)	0.049(15)	0.105(11)	RV,GR	39,40,41
VW Hyi	0.074271038(14)	–	0.71(22)	–	–	GR	42,43
AM Her	0.128927(2)	–	0.78(15)	–	–	SM	44,45

Table B1. *Continued.*

Date	Object	Instrument Setup	Filter(s)	T_{mid} (MJD)	Cycle No.	Phase Coverage	T_{exp} (s)	$N_{u'}$	N_{exp}	Seeing (")	Flag
2010 May 12	CSS080623	NTT+UCAM	$u' g' r'$	55329.23459(3) ^h	0	-0.110-0.177	3.301	3	504	1.2-1.4	2
2010 May 17	CSS080623	NTT+UCAM	$u' g' r'$	55334.12012(3) ^h	82	-0.318-0.363	4.920	3	712	0.9-1.8	2
2010 May 17	CSS080623	NTT+UCAM	$u' g' r'$	55334.17971(3) ^h	83	-0.244-0.451	3.923	4	909	1.1-1.4	2
2010 May 17	CSS080623	NTT+UCAM	$u' g' r'$	55334.23925(3) ^h	84	-0.512-0.269	3.923	3	1021	1.2-1.6	2
2010 Jun 07	CSS080623	NTT+UCAM	$u' g' r'$	55355.03231(3) ^h	433	-0.254-0.226	3.818	2	644	1.0-1.1	2
2010 Jun 07	CSS080623	NTT+UCAM	$u' g' r'$	55355.15152(3) ^h	435	-0.318-0.250	3.818	2	764	1.0-1.8	2
2011 May 27	CSS080623	NTT+UCAM	$u' g' r'$	55709.05056(3) ^h	6375	-0.100-0.141	2.890	2	427	0.9-1.1	2
2011 May 30	CSS080623	NTT+UCAM	$u' g' r'$	55711.96995(3) ^h	6424	-0.192-0.146	3.941	3	440	1.1-1.3	2
2011 May 30	CSS080623	NTT+UCAM	$u' g' r'$	55712.02952(3) ^h	6425	-0.479-0.136	3.941	3	798	1.1-1.5	2
2011 May 30	CSS080623	NTT+UCAM	$u' g' r'$	55712.14867(3) ^h	6427	-0.505-0.279	3.941	3	1023	1.0-1.2	2
2011 Jan 18	CSS110113	NTT+UCAM	$u' g' i'$	55580.12190(10) ^h	-5479	-0.332-0.159	2.376	3	1168	0.9-1.0	8
2012 Jan 14	CSS110113	WHT+UCAM	$u' g' r'$	55940.95783(3) ^h	-16	-0.215-0.178	3.985	3	563	1.0-1.6	2
2012 Jan 15	CSS110113	WHT+UCAM	$u' g' r'$	55942.01465(3) ^h	0	-0.167-0.175	4.980	4	393	1.2-1.8	2
2012 Jan 16	CSS110113	WHT+UCAM	$u' g' r'$	55942.87330(3) ^h	13	-0.237-0.117	3.985	3	507	1.4-2.1	2
2012 Sep 09	CSS110113	WHT+UCAM	$u' g' r'$	56180.12801(3) ^h	3605	-0.197-0.114	3.987	3	448	1.4-1.8	5
2012 Oct 13	CSS110113	WHT+UCAM	$u' g' r'$	56214.07819(3) ^h	4119	-0.232-0.140	2.989	3	711	1.2-1.4	5
2014 Jan 02	CSS110113	WHT+UCAM	$u' g' r'$	56659.92157(3) ^h	10869	-0.194-0.091	3.980	3	408	0.9-1.4	5
2014 Jan 28	CSS110113	TNT+USPEC	$KG5$	56685.54929(3) ^h	11257	-0.030-0.174	9.352	-	126	1.3-2.0	7
2014 Jan 28	CSS110113	TNT+USPEC	$KG5$	56685.61535(3) ^h	11258	-0.186-0.185	9.352	-	227	1.4-1.7	7
2014 Jan 29	CSS110113	TNT+USPEC	g'	56686.54005(3) ^h	11272	-0.533-0.324	9.352	-	523	1.4-3.0	7
2014 Feb 01	CSS110113	TNT+USPEC	g'	56689.57840(3) ^h	11318	-0.106-0.358	8.958	-	296	1.9-2.1	7
2014 Mar 14	CSS110113	WHT+UCAM	$u' g' r'$	56730.86018(3) ^h	11943	-0.205-0.240	2.627	3	957	1.2-1.8	7
2007 Jun 10	CTCV 1300	VLT+UCAM	$u' g' r'$	54262.09916(3) ^h	0	-0.270-0.193	1.002	4	3463	1.2-2.3	1
2007 Jun 13	CTCV 1300	VLT+UCAM	$u' g' i'$	54265.12310(3) ^h	34	-0.261-0.144	1.952	3	1574	1.4-2.1	1
2010 Jun 07	CTCV 1300	NTT+UCAM	$u' g' r'$	55355.00260(5) ^h	12288	-0.142-0.120	3.938	3	511	0.9-1.1	6
2011 May 30	CTCV 1300	NTT+UCAM	$u' g' r'$	55712.18836(5) ^h	16304	-0.175-0.149	2.895	3	852	0.9-1.7	1
2003 May 23	GY Cnc	WHT+UCAM	$u' g' i'$	52782.93530(10) ^b	-17985	-0.085-0.128	1.628	1	1945	0.9-1.1	8
2012 Jan 11	GY Cnc	WHT+UCAM	$u' g' r'$	55938.26366(5) ^b	0	-0.082-0.140	3.974	3	842	1.0-1.2	6
2012 Jan 14	GY Cnc	WHT+UCAM	$u' g' r'$	55941.24626(5) ^b	17	-0.077-0.128	3.077	2	1005	1.2-1.8	6
2012 Jan 16	GY Cnc	WHT+UCAM	$u' g' r'$	55943.00068(5) ^b	27	-0.066-0.115	2.480	3	1096	0.9-1.6	6
2012 Jan 16	GY Cnc	WHT+UCAM	$u' g' r'$	55943.17605(5) ^b	28	-0.168-0.112	2.480	3	1692	1.1-1.5	1
2012 Jan 20	GY Cnc	WHT+UCAM	$u' g' r'$	55947.21130(5) ^b	51	-0.109-0.093	3.879	3	784	2.1-2.7	6
2013 Dec 30	GY Cnc	WHT+UCAM	$u' g' r'$	56657.22682(5) ^b	4098	-0.180-0.115	3.974	3	1120	0.9-1.2	6
2014 Jan 26	GY Cnc	TNT+USPEC	$KG5$	56683.71865(5) ^b	4249	-0.112-0.098	1.273	-	2462	1.0-1.6	1
2015 Feb 27	GY Cnc	TNT+USPEC	$KG5$	57080.56902(10) ^b	6511	-0.059-0.138	2.473	-	1200	1.6-2.2	6
2015 Dec 11	GY Cnc	TNT+USPEC	$KG5$	57367.76812(10) ^b	8148	-0.146-0.213	3.967	-	1370	1.3-1.6	6
2016 Mar 13	GY Cnc	TNT+USPEC	g'	57460.57718(10) ^b	8677	-0.111-0.123	3.926	-	898	1.1-1.6	6
2017 Feb 13	GY Cnc	TNT+USPEC	g'	57797.60241(15) ^b	10598	-0.185-0.161	1.766	-	2944	1.3-1.8	6
2003 May 20	DV UMa	WHT+UCAM	$u' g' i'$	52779.969152(20) ^h	-35	-0.092-0.177	5.921	1	339	1.2-2.7	2
2003 May 22	DV UMa	WHT+UCAM	$u' g' i'$	52781.943747(20) ^h	-12	-0.126-0.104	4.921	1	345	0.9-1.1	2
2003 May 23	DV UMa	WHT+UCAM	$u' g' i'$	52782.974025(20) ^h	0	-0.135-0.151	3.921	1	540	0.9-1.1	5
2007 Oct 19	DV UMa	WHT+UCAM	$u' g' r'$	54393.225867(10) ^h	18756	-0.172-0.180	2.754	2	940	1.2-2.4	2

Table C1. Journal of observations for systems modelled in this paper. The instrument setup column shows both the telescope and instrument (UCAM and USPEC refer to ULTRACAM and ULTRASPEC, respectively) used for each eclipse observation. T_{mid} represents the mid-eclipse time (taken to be that of the white dwarf, when visible), T_{exp} the exposure time and N_{exp} the number of exposures. $N_{u'}$ indicates the number of u' band frames which were co-added on-chip to reduce the impact of readout noise. The final column is a flag for eclipse status: (1) modelled individually, (2) modelled as constituent of average eclipse, (3) usable for individual modelling but not chosen, (4) obtained post-modelling but usable, (5) clear bright spot features but not included in average eclipse due to significant change in disc radius/contribution, (6) lack of bright spot ingress/merged ingresses, (7) poor SNR, (8) in outburst.

^hHeliocentric time in HMJD(UTC), ^bBarycentric time in BMJD(TDB).




Figure A4. Simultaneous eclipse model fit to three average DV UMa eclipse light curves. See Section 4.1 for full details of what is plotted. Displayed in the top-right corner of each average eclipse plot is the date and wavelength band each of the constituent eclipses were observed in.




Figure A5. Simultaneous eclipse model fit to five IY UMa eclipse light curves. See Section 4.1 for full details of what is plotted. Displayed in the top-right corner of each eclipse plot is the cycle number of the eclipse and the wavelength band it was observed in.




Figure A6. Simultaneous eclipse model fit to six OY Car eclipse light curves. See Section 4.1 for full details of what is plotted. Displayed in the top-right corner of each eclipse plot is the cycle number of the eclipse and the wavelength band it was observed in.




Figure A7. Simultaneous eclipse model fit to five average SDSS 0901 eclipse light curves. See Section 4.1 for full details of what is plotted. Displayed in the top-right corner of each average eclipse plot is the date and wavelength band each of the constituent eclipses were observed in.




Figure A8. Simultaneous eclipse model fit to four GY Cnc eclipse light curves. See Section 4.1 for full details of what is plotted. Displayed in the top-right corner of each eclipse plot is the cycle number of the eclipse and the wavelength band it was observed in.




Figure A9. Simultaneous eclipse model fit to four SDSS 1006 eclipse light curves. See Section 4.1 for full details of what is plotted. Displayed in the top-right corner of each eclipse plot is the cycle number of the eclipse and the wavelength band it was observed in.




Figure A10. Simultaneous eclipse model fit to four average SDSS 1152 eclipse light curves. See Section 4.1 for full details of what is plotted. Displayed in the top-right corner of each average eclipse plot is the date and wavelength band each of the constituent eclipses were observed in.




Figure A11. Simultaneous eclipse model fit to three average SSS100615 eclipse light curves. See Section 4.1 for full details of what is plotted. Displayed in the top-right corner of each average eclipse plot is the date and wavelength band each of the constituent eclipses were observed in.




Figure A12. Simultaneous eclipse model fit to six SDSS 1501 eclipse light curves. See Section 4.1 for full details of what is plotted. Displayed in the top-right corner of each eclipse plot is the cycle number of the eclipse and the wavelength band it was observed in.




Figure A13. Simultaneous eclipse model fit to six SSS130413 eclipse light curves. See Section 4.1 for full details of what is plotted. Displayed in the top-right corner of each eclipse plot is the cycle number of the eclipse and the wavelength band it was observed in.




Figure A15. Simultaneous eclipse model fit to six Z Cha eclipse light curves. See Section 4.1 for full details of what is plotted. Displayed in the top-right corner of each eclipse plot is the cycle number of the eclipse and the wavelength band it was observed in.




Figure A14. Simultaneous eclipse model fit to five V713 Cep eclipse light curves. See Section 4.1 for full details of what is plotted. Displayed in the top-right corner of each eclipse plot is the cycle number of the eclipse and the wavelength band it was observed in.

Date	Object	Instrument Setup	Filter(s)	T_{mid} (MJD)	Cycle No.	Phase Coverage	T_{exp} (s)	$N_{u'}$	N_{exp}	Seeing ('')	Flag
2014 Mar 30	IY UMa	TNT+USPEC	<i>KG5</i>	56746.639516(20) ^h	0	-0.102-0.325	2.190	-	1243	1.5-2.9	1
2014 Mar 30	IY UMa	TNT+USPEC	<i>KG5</i>	56746.713426(20) ^h	1	-0.060-0.201	2.190	-	763	1.6-2.0	1
2014 Mar 30	IY UMa	TNT+USPEC	<i>KG5</i>	56746.787335(20) ^h	2	-0.066-0.249	2.190	-	913	1.8-2.5	1
2014 Nov 30	IY UMa	TNT+USPEC	<i>KG5</i>	56991.94334(10) ^h	3319	-0.090-0.174	3.352	-	502	1.6-1.9	8
2015 Jan 03	IY UMa	TNT+USPEC	<i>g'</i>	57025.94133(3) ^h	3779	-0.203-0.176	3.352	-	713	1.1-1.4	1
2015 Jan 06	IY UMa	TNT+USPEC	<i>r'</i>	57028.89764(5) ^h	3819	-0.159-0.125	3.952	-	458	1.2-1.5	3
2015 Feb 23	IY UMa	TNT+USPEC	<i>r'</i>	57076.86456(3) ^h	4468	-0.130-0.249	3.852	-	632	1.9-2.3	1
2016 Mar 11	IY UMa	TNT+USPEC	<i>u'</i>	57458.67790(10) ^h	9634	-0.185-0.269	29.78	-	99	1.7-2.3	6
2016 Mar 13	IY UMa	TNT+USPEC	<i>u'</i>	57460.67369(10) ^h	9661	-0.195-0.287	25.35	-	122	1.5-2.0	3
2016 Mar 13	IY UMa	TNT+USPEC	<i>i'</i>	57460.74748(5) ^h	9662	-0.363-0.320	7.852	-	655	1.1-1.5	1
2010 Apr 27	OY Car	NTT+UCAM	<i>u' g' r'</i>	55314.104056(8) ^h	-632	-0.103-0.140	1.760	3	747	1.7-2.7	3
2010 Jun 06	OY Car	NTT+UCAM	<i>u' g' r'</i>	55353.996480(8) ^h	0	-0.119-0.170	1.424	3	3116	1.3-1.4	1
2010 Jun 07	OY Car	NTT+UCAM	<i>u' g' r'</i>	55355.069543(8) ^h	17	-0.293-0.249	1.369	3	2120	1.1-1.8	3
2010 Nov 18	OY Car	NTT+UCAM	<i>u' g' i'</i>	55519.310181(8) ^h	2619	-0.206-0.487	1.329	4	3894	1.3-2.7	1
2010 Dec 17	OY Car	NTT+UCAM	<i>u' g' i'</i>	55548.282678(8) ^h	3078	-0.189-0.126	2.814	2	606	0.8-1.0	1
2011 May 24	OY Car	NTT+UCAM	<i>u' g' r'</i>	55706.084989(8) ^h	5578	-0.368-0.180	1.329	1	2205	1.3-2.0	3
2016 Nov 10	OY Car	NTT+UCAM	<i>u' g' r'</i>	57703.294204(8) ^h	37219	-0.141-0.202	1.979	2	931	0.9-1.4	4
2006 Mar 09	SDSS 0901	WHT+UCAM	<i>u' g' r'</i>	53803.906350(20) ^h	-27455	-0.763-0.259	4.972	1	1374	1.2-2.0	2
2006 Mar 10	SDSS 0901	WHT+UCAM	<i>u' g' r'</i>	53804.996665(20) ^h	-27441	-0.135-0.092	4.972	1	307	1.1-1.4	2
2006 Mar 10	SDSS 0901	WHT+UCAM	<i>u' g' r'</i>	53805.152456(20) ^h	-27439	-0.258-0.178	4.972	1	590	1.2-1.7	2
2010 Jan 07	SDSS 0901	WHT+UCAM	<i>u' g' r'</i>	55203.96452(5) ^h	-9478	-0.289-0.137	1.677	4	1685	1.7-3.0	6
2012 Jan 15	SDSS 0901	WHT+UCAM	<i>u' g' r'</i>	55942.116352(20) ^h	0	-0.350-0.152	4.480	3	752	0.8-1.0	2
2012 Jan 15	SDSS 0901	WHT+UCAM	<i>u' g' r'</i>	55942.194233(20) ^h	1	-0.416-0.240	4.480	3	987	0.9-1.4	2
2012 Jan 16	SDSS 0901	WHT+UCAM	<i>u' g' r'</i>	55942.973064(20) ^h	11	-0.132-0.146	4.480	3	417	1.1-1.6	2
2012 Jan 16	SDSS 0901	WHT+UCAM	<i>u' g' r'</i>	55943.050921(20) ^h	12	-0.373-0.128	4.480	3	752	1.0-1.9	2
2012 Jan 16	SDSS 0901	WHT+UCAM	<i>u' g' r'</i>	55943.206687(20) ^h	14	-0.125-0.179	4.480	3	456	1.1-1.7	2
2012 Jan 16	SDSS 0901	WHT+UCAM	<i>u' g' r'</i>	55943.284578(20) ^h	15	-0.235-0.219	4.480	3	678	1.1-2.1	2
2012 Jan 15	SDSS 1006	WHT+UCAM	<i>u' g' r'</i>	55942.05221(10) ^h	-3984	-0.116-0.180	3.980	3	1187	0.8-1.7	6
2014 Jan 25	SDSS 1006	TNT+USPEC	<i>KG5</i>	56682.72940(20) ^h	0	-0.114-0.129	3.352	-	1165	1.6-2.3	1
2014 Jan 25	SDSS 1006	TNT+USPEC	<i>KG5</i>	56682.91552(20) ^h	1	-0.190-0.127	3.352	-	1514	1.6-2.0	6
2014 Jan 26	SDSS 1006	TNT+USPEC	<i>KG5</i>	56683.84494(20) ^h	6	-0.127-0.155	3.352	-	1342	1.2-1.4	3
2014 Jan 28	SDSS 1006	TNT+USPEC	<i>KG5</i>	56685.88995(20) ^h	17	-0.171-0.131	3.352	-	1446	1.2-1.4	1
2014 Feb 02	SDSS 1006	TNT+USPEC	<i>g'</i>	56690.72366(20) ^h	43	-0.267-0.176	5.892	-	1204	1.4-2.0	1
2014 Feb 02	SDSS 1006	TNT+USPEC	<i>r'</i>	56690.90953(20) ^h	44	-0.143-0.117	3.352	-	1244	1.3-2.1	1
2015 Dec 06	SDSS 1006	TNT+USPEC	<i>g'</i>	57362.79981(20) ^h	3658	-0.144-0.138	4.946	-	974	2.2-3.6	7
2016 Mar 14	SDSS 1006	TNT+USPEC	<i>g'</i>	57461.70591(20) ^h	4190	-0.266-0.202	9.640	-	783	1.0-3.1	7
2017 Feb 20	SDSS 1006	TNT+USPEC	<i>KG5</i>	57804.71547(20) ^h	6035	-0.239-0.208	4.970	-	1437	1.4-3.4	6
2017 Feb 21	SDSS 1006	TNT+USPEC	<i>g'</i>	57805.64520(20) ^h	6040	-0.198-0.152	5.470	-	1024	2.0-2.7	7
2010 Jan 07	SDSS 1152	WHT+UCAM	<i>u' g' r'</i>	55204.101280(10) ^h	0	-0.840-0.137	3.800	3	1492	2.4-4.0	2
2010 Jan 07	SDSS 1152	WHT+UCAM	<i>u' g' r'</i>	55204.169035(10) ^h	1	-0.274-0.116	3.800	3	600	1.2-2.6	2
2010 Jan 07	SDSS 1152	WHT+UCAM	<i>u' g' r'</i>	55204.236772(10) ^h	2	-0.151-0.119	3.800	3	415	1.5-3.0	2
2014 Mar 14	SDSS 1152	WHT+UCAM	<i>u' g' r'</i>	56730.908566(10) ^h	22536	-0.265-0.195	4.029	3	668	1.2-1.7	7
2014 Mar 30	SDSS 1152	TNT+USPEC	<i>KG5</i>	56746.694264(10) ^h	22769	-0.385-0.195	5.352	-	634	1.3-1.6	2
2014 Mar 30	SDSS 1152	TNT+USPEC	<i>KG5</i>	56746.762006(10) ^h	22770	-0.322-0.259	5.352	-	634	1.2-1.7	2
2014 Mar 30	SDSS 1152	TNT+USPEC	<i>KG5</i>	56746.829759(10) ^h	22771	-0.290-0.285	5.352	-	628	1.4-1.9	2

Table C1. *Continued.*

Date	Object	Instrument Setup	Filter(s)	T_{mid} (MJD)	Cycle No.	Phase Coverage	T_{exp} (s)	$N_{u'}$	N_{exp}	Seeing (")	Flag
2012 Apr 28	SDSS 1057	WHT+UCAM	$u' g' r'$	56046.002399(12) ^h	0	-0.581-0.149	4.021	3	981	1.2-2.7	2
2012 Apr 29	SDSS 1057	WHT+UCAM	$u' g' r'$	56046.944270(12) ^h	15	-0.239-0.228	4.021	3	628	1.1-2.0	2
2013 Dec 30	SDSS 1057	WHT+UCAM	$u' g' r'$	56657.28205(3) ^h	9735	-0.558-0.320	4.021	3	1178	1.0-1.6	7
2014 Jan 25	SDSS 1057	TNT+USPEC	$KG5$	56682.775595(12) ^h	10141	-0.316-0.064	4.877	-	422	1.4-2.7	7
2014 Nov 28	SDSS 1057	TNT+USPEC	$KG5$	56989.82829(3) ^h	15031	-0.225-0.138	3.945	-	498	1.3-2.5	7
2014 Nov 29	SDSS 1057	TNT+USPEC	$KG5$	56990.89570(3) ^h	15048	-0.158-0.143	4.945	-	331	0.9-1.4	7
2015 Feb 24	SDSS 1057	TNT+USPEC	$KG5$	57077.862577(12) ^h	16433	-0.219-0.281	11.852	-	230	1.4-2.1	2
2015 Feb 25	SDSS 1057	TNT+USPEC	$KG5$	57078.867265(12) ^h	16449	-0.207-0.169	11.946	-	172	2.0-2.4	2
2015 Mar 01	SDSS 1057	TNT+USPEC	$KG5$	57082.885950(12) ^h	16513	-0.101-0.138	11.852	-	111	1.4-1.8	2
2015 Jun 21	SDSS 1057	WHT+UCAM	$u' g' r'$	57194.906824(12) ^h	18297	-0.390-0.182	4.021	3	769	1.2-2.1	2
2015 Jun 22	SDSS 1057	WHT+UCAM	$u' g' r'$	57195.911476(12) ^h	18313	-0.170-0.171	4.021	3	460	1.2-2.3	2
2015 Jun 23	SDSS 1057	WHT+UCAM	$u' g' r'$	57196.916157(12) ^h	18329	-0.179-0.130	4.021	3	416	1.1-2.0	5
2004 May 17	SDSS 1501	WHT+UCAM	$u' g' r'$	53142.921635(10) ^h	-53411	-0.198-0.218	6.115	1	335	0.9-1.4	1
2006 Mar 04	SDSS 1501	WHT+UCAM	$u' g' r'$	53799.210838(10) ^h	-41865	-0.663-0.165	4.971	1	813	1.4-2.4	6
2006 Mar 05	SDSS 1501	WHT+UCAM	$u' g' r'$	53800.120302(10) ^h	-41849	-0.845-0.195	5.971	1	853	2.1-3.9	6
2006 Mar 07	SDSS 1501	WHT+UCAM	$u' g' r'$	53802.052900(10) ^h	-41815	-0.294-0.192	4.971	1	480	1.0-1.5	6
2006 Mar 07	SDSS 1501	WHT+UCAM	$u' g' r'$	53802.109742(10) ^h	-41814	-0.316-0.217	4.971	1	525	1.1-1.4	6
2006 Mar 07	SDSS 1501	WHT+UCAM	$u' g' r'$	53802.223433(10) ^h	-41812	-0.189-0.214	4.971	1	397	1.0-1.3	6
2006 Mar 08	SDSS 1501	WHT+UCAM	$u' g' r'$	53803.132876(10) ^h	-41796	-0.175-0.143	4.971	1	315	1.1-1.6	6
2006 Mar 08	SDSS 1501	WHT+UCAM	$u' g' r'$	53803.189718(10) ^h	-41795	-0.057-0.141	4.971	1	197	1.1-1.5	6
2006 Mar 08	SDSS 1501	WHT+UCAM	$u' g' r'$	53803.246575(10) ^h	-41794	-0.088-0.139	4.971	1	227	0.9-1.1	6
2010 Jan 07	SDSS 1501	WHT+UCAM	$u' g' r'$	55204.213149(15) ^h	-17147	-0.217-0.120	3.800	3	435	1.4-3.9	7
2010 Jan 07	SDSS 1501	WHT+UCAM	$u' g' r'$	55204.270013(15) ^h	-17146	-0.119-0.129	3.800	3	321	1.2-2.7	7
2012 Sep 08	SDSS 1501	WHT+UCAM	$u' g' r'$	56178.870508(10) ^h	0	-0.158-0.165	3.475	3	455	1.0-1.5	1
2014 Aug 03	SSS100615	WHT+UCAM	$u' g' r'$	56873.023626(5) ^h	0	-0.764-0.157	3.005	4	1538	1.0-1.2	2
2014 Aug 04	SSS100615	WHT+UCAM	$u' g' r'$	56874.021600(5) ^h	17	-0.084-0.162	3.005	3	416	1.0-1.1	2
2014 Aug 05	SSS100615	WHT+UCAM	$u' g' i'$	56874.960873(5) ^h	33	-0.114-0.103	5.056	3	219	1.7-2.3	2
2014 Jan 26	SSS130413	TNT+USPEC	$KG5$	56683.67392(5) ^h	0	-0.175-0.195	5.804	-	362	1.6-2.2	3
2014 Jan 26	SSS130413	TNT+USPEC	$KG5$	56683.73977(5) ^h	1	-0.085-0.141	5.804	-	222	1.5-2.0	1
2014 Jan 26	SSS130413	TNT+USPEC	$KG5$	56683.80551(5) ^h	2	-0.230-0.194	5.804	-	415	1.3-1.5	1
2014 Jan 27	SSS130413	TNT+USPEC	$KG5$	56684.72635(5) ^h	16	-0.602-0.349	5.804	-	928	1.8-2.9	3
2014 Jan 28	SSS130413	TNT+USPEC	$KG5$	56685.71280(5) ^h	31	-0.209-0.181	2.934	-	752	1.4-1.7	3
2014 Feb 01	SSS130413	TNT+USPEC	$KG5$	56689.65905(5) ^h	91	-0.291-0.191	2.934	-	931	1.9-2.2	3
2014 Feb 01	SSS130413	TNT+USPEC	$KG5$	56689.85631(5) ^h	94	-0.148-0.176	2.934	-	627	1.5-2.0	1
2014 Feb 02	SSS130413	TNT+USPEC	g'	56690.77702(5) ^h	108	-0.304-0.215	2.934	-	998	1.5-1.7	1
2014 Feb 02	SSS130413	TNT+USPEC	r'	56690.84278(5) ^h	109	-0.223-0.126	2.934	-	673	1.3-1.5	1
2014 Mar 23	SSS130413	TNT+USPEC	$KG5$	56739.64406(8) ^h	851	-0.177-0.149	2.934	-	628	1.3-1.6	7
2015 Jan 01	SSS130413	TNT+USPEC	i'	57023.76697(3) ^h	5171	-0.120-0.190	2.939	-	595	1.0-1.4	7
2016 Jan 29	SSS130413	TNT+USPEC	u'	57416.73848(5) ^h	11146	-0.388-0.161	9.252	-	338	2.7-3.6	7
2016 Jan 31	SSS130413	TNT+USPEC	u'	57418.71160(5) ^h	11176	-0.228-0.167	8.052	-	280	2.0-2.7	1
2017 Feb 12	SSS130413	TNT+USPEC	r'	57796.75351(10) ^h	16924	-0.656-0.244	8.052	-	670	1.8-2.7	7
2017 Feb 20	SSS130413	TNT+USPEC	g'	57804.77728(3) ^h	17046	-0.210-0.147	3.951	-	511	2.1-3.1	4
2017 Feb 21	SSS130413	TNT+USPEC	$KG5$	57805.69806(3) ^h	17060	-0.284-0.285	3.951	-	815	1.4-2.7	4
2017 Feb 24	SSS130413	TNT+USPEC	r'	57808.78922(3) ^h	17107	-0.305-0.186	3.952	-	705	1.2-1.6	4

Table C1. Continued.

Date	Object	Instrument Setup	Filter(s)	T_{mid} (MJD)	Cycle No.	Phase Coverage	T_{exp} (s)	$N_{u'}$	N_{exp}	Seeing ('')	Flag
2011 Aug 27	V713 Cep	WHT+UCAM	$u' g' i'$	55801.180379(20) ^h	-4399	-0.383-0.136	2.185	2	1737	1.2-1.6	6
2012 Sep 06	V713 Cep	WHT+UCAM	$u' g' r'$	56176.936374(20) ^h	0	-0.373-0.127	3.445	3	1065	1.0-1.2	3
2012 Sep 07	V713 Cep	WHT+UCAM	$u' g' r'$	56177.875986(20) ^h	11	-0.346-0.184	3.445	3	1126	1.0-1.2	1
2012 Sep 09	V713 Cep	WHT+UCAM	$u' g' r'$	56180.011450(20) ^h	36	-0.212-0.169	3.445	3	811	0.9-1.4	6
2013 Jul 14	V713 Cep	WHT+UCAM	$u' g' z'$	56488.201454(20) ^h	3644	-0.309-0.291	3.445	3	1277	1.0-1.2	3
2013 Jul 15	V713 Cep	WHT+UCAM	$u' g' i'$	56489.055641(20) ^h	3654	-0.124-0.162	3.445	3	609	1.1-1.3	3
2013 Jul 15	V713 Cep	WHT+UCAM	$u' g' i'$	56489.141063(20) ^h	3655	-0.105-0.111	3.445	3	460	0.9-4.5	1
2013 Jul 15	V713 Cep	WHT+UCAM	$u' g' i'$	56489.226484(20) ^h	3656	-0.429-0.110	3.445	3	1150	0.9-1.1	7
2013 Jul 25	V713 Cep	WHT+UCAM	$u' g' z'$	56499.220435(20) ^h	3773	-0.275-0.205	3.445	3	1022	1.1-1.3	6
2013 Aug 04	V713 Cep	WHT+UCAM	$u' g' r'$	56509.214395(20) ^h	3890	-0.323-0.308	3.445	3	1342	1.1-1.7	3
2013 Aug 05	V713 Cep	WHT+UCAM	$u' g' r'$	56510.154021(20) ^h	3901	-0.288-0.168	3.445	3	971	0.9-1.4	6
2014 Aug 02	V713 Cep	WHT+UCAM	$u' g' r'$	56872.07224(3) ^h	8138	-0.388-0.216	1.983	3	2224	1.4-3.0	6
2014 Aug 10	V713 Cep	WHT+UCAM	$u' g' i'$	56880.18702(3) ^h	8233	-0.182-0.155	3.445	3	719	1.1-2.4	7
2015 Jun 24	V713 Cep	WHT+UCAM	$u' g' r'$	57198.114652(20) ^h	11955	-0.213-0.070	3.445	3	603	0.8-1.0	6
2015 Sep 17	V713 Cep	WHT+UCAM	$u' g' r'$	57282.849778(20) ^h	12947	-0.047-0.118	4.985	3	245	1.1-1.5	6
2005 May 07	Z Cha	VLT+UCAM	$u' g' r'$	53498.011478(10) ^h	0	-0.147-0.142	0.476	1	4300	1.7-2.3	3
2005 May 10	Z Cha	VLT+UCAM	$u' g' r'$	53500.991449(10) ^h	40	-0.373-0.175	0.476	1	7632	1.3-4.8	6
2005 May 12	Z Cha	VLT+UCAM	$u' g' r'$	53503.002929(10) ^h	67	-0.071-0.113	0.476	1	10070	2.1-8.1	3
2005 May 15	Z Cha	VLT+UCAM	$u' g' r'$	53505.982900(10) ^h	107	-0.108-0.204	0.476	1	7007	1.7-2.7	3
2005 May 17	Z Cha	VLT+UCAM	$u' g' r'$	53507.994355(10) ^h	134	-0.586-0.176	0.476	1	9769	2.1-3.9	3
2005 May 21	Z Cha	VLT+UCAM	$u' g' r'$	53512.017323(10) ^h	188	-0.113-0.161	0.476	1	6674	1.8-3.6	3
2010 Apr 26	Z Cha	NTT+UCAM	$u' g' r'$	55313.03694(3) ^h	24363	-0.056-0.119	1.517	4	731	2.6-3.6	6
2010 Jun 06	Z Cha	NTT+UCAM	$u' g' r'$	55354.08603(10) ^h	24914	-0.337-0.090	1.331	3	2024	1.1-1.8	8
2010 Nov 26	Z Cha	NTT+UCAM	$u' g' i'$	55527.147891(20) ^h	27237	-0.063-0.097	1.381	3	1636	0.8-1.1	1
2010 Dec 02	Z Cha	NTT+UCAM	$u' g' r'$	55533.33134(10) ^h	27320	-0.153-0.265	1.324	3	1996	1.1-1.7	8
2010 Dec 11	Z Cha	NTT+UCAM	$u' g' i'$	55542.345728(20) ^h	27441	-0.169-0.171	1.331	3	1775	1.1-1.6	6
2010 Dec 17	Z Cha	NTT+UCAM	$u' g' i'$	55548.082204(20) ^h	27518	-0.136-0.128	2.874	2	589	1.1-2.0	1
2010 Dec 17	Z Cha	NTT+UCAM	$u' g' i'$	55548.305686(20) ^h	27521	-0.124-0.134	2.874	2	576	0.8-1.0	3
2011 May 19	Z Cha	NTT+UCAM	$u' g' r'$	55701.029233(20) ^h	29571	-0.295-0.202	2.855	2	1108	1.8-3.6	1

Table C1. *Continued.*