



UNIVERSITY OF LEEDS

This is a repository copy of *Study Protocol for the DETECTIVE Study: An International Collaborative Study To Develop Consensus Statements for Deferred Treatment with Curative Intent for Localised Prostate Cancer*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/145398/>

Version: Accepted Version

---

**Article:**

Lam, TBL, MacLennan, S, Plass, K et al. (37 more authors) (2019) Study Protocol for the DETECTIVE Study: An International Collaborative Study To Develop Consensus Statements for Deferred Treatment with Curative Intent for Localised Prostate Cancer. *European Urology*, 75 (4). pp. 699-702. ISSN 0302-2838

<https://doi.org/10.1016/j.eururo.2018.11.009>

---

© 2018 European Association of Urology. Published by Elsevier B.V. All rights reserved. Licensed under the Creative Commons Attribution-Non Commercial No Derivatives 4.0 International License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

## Brief Correspondence

### Study protocol for DETECTIVE Study: An international collaborative study to develop consensus statements for deferred treatment with curative intent for localised prostate cancer

Thomas B.L. Lam<sup>a,\*</sup>, Steven MacLennan<sup>a</sup>, Karin Plass<sup>b</sup>, Peter-Paul M. Willemse<sup>c</sup>, Malcolm D. Mason<sup>d</sup>, Philip Cornford<sup>e</sup>, James Donaldson<sup>a</sup>, Niall Davis<sup>f</sup>, Paolo Dell'Oglio<sup>g</sup>, Christian Fankhauser<sup>h</sup>, Nikos Grivas<sup>i</sup>, Alexandre Ingels<sup>j</sup>, Michael Lardas<sup>k</sup>, Matthew Liew<sup>l</sup>, Karl Pang<sup>m</sup>, Catherine Paterson<sup>n</sup>, M.I. Omar<sup>a</sup>, Fabio Zattoni<sup>o</sup>, Tim Buddingh<sup>p</sup>, Thomas Van den Broeck<sup>q</sup>, Marcus Cumberbatch<sup>m</sup>, Nicola Fossati<sup>g</sup>, Tobias Gross<sup>r</sup>, Lisa Moris<sup>q</sup>, Ivo G. Schoots<sup>s</sup>, Roderick C.N. van den Bergh<sup>t</sup>, Erik Briers<sup>u</sup>, Stefano Fanti<sup>v</sup>, Maria De Santis<sup>w</sup>, Silke Gillessen<sup>x</sup>, Jeremy P. Grummet<sup>y</sup>, Ann M. Henry<sup>z</sup>, Henk G. van der Poel<sup>t</sup>, Theo H. van der Kwast<sup>aa</sup>, Olivier Rouvière<sup>bb</sup>, Derya Tilki<sup>cc</sup>, Thomas Wiegel<sup>dd</sup>, *James N'Dow*<sup>a</sup>, Hendrik van Poppel<sup>q</sup>, Nicolas Mottet<sup>ee</sup>.

<sup>a</sup> Academic Urology Unit, University of Aberdeen, Aberdeen, UK; <sup>b</sup> EAU Guidelines Office, Arnhem, The Netherlands; <sup>c</sup> Department of Urology, University Utrecht, Utrecht, The Netherlands; <sup>d</sup> Division of Cancer & Genetics, School of Medicine Cardiff University, Velindre Cancer Centre, Cardiff, UK; <sup>e</sup> Royal Liverpool and Broadgreen Hospitals NHS Trust, Liverpool, UK; <sup>f</sup> Department of Urology, The Austin Hospital, Melbourne, Victoria, Australia; <sup>g</sup> Department of Urology, San Raffaele Hospital, Milan, Italy; <sup>h</sup> Department of Urology, University of Zurich, Zürich, Switzerland; <sup>i</sup> Department of Urology, G. Hatzikosta General Hospital, Ioánnina, Greece; <sup>j</sup> Department of Urology, Institut Montsouris, Paris, France; <sup>k</sup> Department of Urology, Leto Hospital, Athens, Greece; <sup>l</sup> Department of Urology, Wrightington, Wigan and Leigh NHS, Foundation Trust, Wigan, United Kingdom; <sup>m</sup> Department of Urology, The University of Sheffield, Sheffield, United Kingdom; <sup>n</sup> School of Nursing and Midwifery, Robert Gordon University, Aberdeen; <sup>o</sup> Department of Urology, University of Padova, Padova, Italy; <sup>p</sup> Department of Urology, Leiden University Medical Center, Leiden, The Netherlands; <sup>q</sup> Department of Urology, University Hospital K.U. Leuven, Leuven, Belgium; <sup>r</sup> Department of Urology, University of Bern, Inselspital, Bern, Switzerland; <sup>s</sup> Department of Radiology & Nuclear Medicine, Erasmus MC University Medical Center, Rotterdam, The Netherlands; <sup>t</sup> Department of Urology, Netherlands Cancer Institute, Amsterdam, The Netherlands; <sup>u</sup> Patient Advocate, Hasselt, Belgium; <sup>v</sup> Metropolitan Nuclear Medicine, St. Orsola-Malpighi Hospital, University of Bologna, Bologna, Italy; <sup>w</sup> Department of Urology, Medical University of Vienna, Vienna, Austria; <sup>x</sup> Department of Oncology/Hematology, Cantonal Hospital St. Gallen, St. Gallen, Switzerland; <sup>y</sup> Department of Surgery, Central Clinical School, Monash University, Melbourne, Australia; <sup>z</sup> Leeds Cancer Centre, St. James's University Hospital, Leeds, UK; <sup>aa</sup> Department of Pathology, Erasmus Medical Centre, Rotterdam, The Netherlands; <sup>bb</sup> Hospices Civils de Lyon, Radiology Department, Edouard Herriot Hospital, Lyon, France; <sup>cc</sup> Department of Urology, University Hospital Hamburg-Eppendorf, Hamburg, Germany; <sup>dd</sup> Department of Radiation Oncology, University Hospital Ulm, Ulm, Germany; <sup>ee</sup> Department of Urology, University Hospital, St. Etienne, France.

\* Corresponding author. Academic Urology Unit, University of Aberdeen, Foresterhill, Aberdeen AB25 2ZD, UK.

## Abstract:

Deferred active treatment (DAT) strategies, including active surveillance and active monitoring, are a recognised management option for men with localised low-risk prostate cancer. However, there is uncertainty due to heterogeneity of patient selection criteria, follow-up and monitoring characteristics, reclassification thresholds, and which outcome measures should be prioritised. This protocol describes a study led by the European Association of Urology (EAU) Prostate Cancer Guidelines Panel in conjunction with other guideline organisations and societies\* to develop consensus statements for all domains of deferred active treatment. The project is divided into 3 sequential phases: (1) Systematic review of studies reporting on DAT in order to summarise and define range of heterogeneity regarding all domains; (2) Two-round Delphi online survey involving a large, international panel of healthcare professionals (HCPs) and patients to initiate consensus; and (3) Consensus group meeting involving representatives from HCP and patient stakeholder groups to finalise the

consensus process. The consensus statements are expected to be adopted by clinical practice guidelines in order to standardise and guide practice for clinicians and researchers until better evidence emerges.

Patient summary: We describe a project aimed at standardising elements of practice in active surveillance/monitoring for early localised prostate cancer, because currently there is great variation and uncertainty regarding how best to conduct them. This will be achieved through a structured process of agreement (i.e. consensus) amongst a large, international panel of healthcare professionals and patients.

Deferred active treatment (DAT) strategies for men with localised prostate cancer have emerged as a viable alternative to radical intervention as we aim to avoid the consequences of over-treatment. Nevertheless, such strategies remain controversial, with significant uncertainty and heterogeneity in all domains, including criteria on patient selection, nature and timing of interventions during follow-up, criteria and thresholds for reclassification, and which outcome measures should be prioritised<sup>1-3</sup>. These are important barriers to the conduct and uptake of DAT by clinicians and patients, and makes it extremely difficult to compare the clinical effectiveness of different protocols against each other. In order to address these issues in a comprehensive, robust and systematic manner, the EAU Prostate Cancer Guidelines Panel, in partnership with other leading guideline authorities and societies\*, has commissioned a project to develop consensus statements in all domains relating to DAT in order to standardise clinical practice and research.

The specific objectives are to achieve consensus on the following domains: (1) Criteria on patient selection (including patient and disease characteristics, imaging criteria and type of biopsies); (2) Nature and timing of interventions during follow-up (such as repeat imaging and repeat biopsies); (3) Criteria and thresholds for reclassification; and (4) Type of outcome measures which should be prioritised.

To address these objectives, we will utilise transparent consensus methods involving a large, international cohort of stakeholders, broadly divided into two groups: (1) Healthcare professionals (HCPs) consisting of urologists, clinical or radiation oncologists, medical oncologists, radiologists, pathologists, primary care physicians, and nurse specialists; and (2) Patients. The research will be divided into 3 distinct but inter-related phases, and is expected to last 12 months.

**Phase 1** is a systematic review conducted according to PRISMA guidelines<sup>4</sup>. The aim to describe, explore and assess clinical heterogeneity in DAT studies which will inform the statements for the consensus processes. The review protocol has been published<sup>5</sup>. In brief, all prospective single-arm case series of DAT (including active surveillance and active monitoring but excluding watchful waiting), and all prospective comparative studies involving DAT will be included. The review will summarise eligibility and selection criteria, characteristics of monitoring and follow-up (including the type, frequency and timing of repeat imaging and repeat biopsies), reclassification definitions and thresholds, and primary outcomes measured in studies. English language articles published after 1990 will be included. Summary of findings tables including details of the pre-specified domains and sub-domains will be developed. From these tables, a list of statements organised according to the different domains and sub-domains relating to all aspects of DAT will be generated.

**Phase 2** will comprise of a two-round online Delphi survey involving a large, international cohort of key stakeholders (HCPs and patients). The consensus methods used have been described previously in consensus studies in prostate cancer<sup>6,7</sup>. HCPs involved with DAT identified through international specialist societies\* will be invited to participate. Patients throughout Europe with localised

prostate cancer and eligible for DAT will be recruited through patient charities\*. Up to 200 HCPs and 100 patients will be invited to participate. Patients will be asked to complete the patient-relevant parts of the survey only (i.e. identification of most important outcomes). Participants will be asked to vote based on their level of agreement, on a nine-point scale, ranging from strongly disagree (1) to strongly agree (9) (i.e. 1–3 disagree; 4–6 uncertain; 7–9 agree). There will also be an 'Unable to answer' option. An online questionnaire will be developed for the Delphi process using COMET Initiative DelphiManager<sup>8</sup>. Two iterative rounds will be conducted anonymously, with anonymised feedback provided to all participants at the end of each round showing the percent scoring at each response option. In Round 1, participants will have the opportunity to add further statements for incorporation into Round 2. With an anticipated response rate of 80% for both stakeholder groups, and expected completion rate of 60% for both rounds, the total number of participants involved is expected to be 144 (i.e. 96 HCPs and 48 patients). The results for each stakeholder group will be analysed and presented separately in each round. After the final round, statements scoring 'strongly agree' (i.e. 7–9) by  $\geq 70\%$  of participants AND with minimal disagreement scored by the rest (defined as  $< 15\%$  of participants scoring 'strongly disagree' i.e. 1-3) will be considered as reaching the threshold for 'consensus agree'. Conversely, statements scoring 'strongly disagree' (i.e. 1-3) by  $\geq 70\%$  of participants AND with minimal agreement scored by the rest (defined as  $< 15\%$  of participants scoring 'strongly agree' i.e. 7-9) will be considered as reaching the threshold for 'consensus disagree'. All other statements not falling in the above categories will be classified as 'equivocal'. Statements reaching consensus (either agree or disagree) will be collated for review in Phase 3, whilst equivocal statements will be brought forward for discussion and voting in Phase 3.

**Phase 3** is the final stage of the consensus process, involving a 1-day meeting attended by representatives of each stakeholder group and chaired by a non-voting methodologist and a clinician moderator. We will use structured discussion and live voting sessions. Representatives from each stakeholder groups and sub-groups (i.e. urologists, oncologists, radiologists, pathologists and patients) will be purposively sampled from those completing all rounds of the Delphi survey to ensure proportional representation. The voting panel will consist of 25 voting participants (i.e. 7 patients and 18 HCPs). Statements reaching consensus (either agree or disagree) from Phase 2 will be reviewed by the panel. Consensus decisions from the Delphi survey cannot be overturned by the panel without sound reasoning (e.g. misleading statements). Equivocal statements from Phase 2 will be discussed and voted on by the panel. Scoring thresholds will be the same as Phase 2 (i.e. level of agreement on a nine-point scale: 1–3 disagree; 4–6 uncertain; 7–9 agree; and 'Unable to answer'). Voting will be anonymous using Poll Everywhere<sup>9</sup> which participants can access during the meeting using personal computers and a shared IP address. Definitions of consensus will be the same as in Phase 2. Results for all statements will be conveyed in real-time, and final consensus statements will be prepared. A final list of consensus statements organised according to the domains and sub-domains of DAT will be issued.

The consensus statements are expected to be adopted by guideline developers and disseminated through clinical practice guidelines issued by the EAU Prostate Cancer Guidelines Panel and other organisations\*, and are intended to provide authoritative guidance to clinicians and researchers by standardising definitions, thresholds, terminology and characteristics of patient selection, monitoring, reclassification and change in management, and outcome measures which should be prioritised in programmes of deferred active treatment in clinical practice and research, at least until higher levels of evidence emerge such as from the GAP3 initiative<sup>10</sup>.

\* The list of official collaborators include the following organisations and patient-led societies:

European Association of Urology (EAU)  
European Association of Urology Nurses (EAUN)  
EAU PIONEER  
European Urology  
American Urological Association (AUA)  
Canadian Urological Association (CUA)  
European Society for Radiotherapy and Oncology (ESTRO)  
International Society of Urological Pathology (ISUP)  
Urological Society of Australia and New Zealand (USANZ)  
European Society of Urogenital Radiology (ESUR)  
EAU Section of Oncological Urology (ESOU)  
Urological Association of Asia (UAA)  
American Society for Radiation Oncology (ASTRO)  
American Society of Clinical Oncology (ASCO)  
European Forum for Primary Care (EFPC)  
EAU Research Foundation (EAU RF)  
UCAN UK  
Tackle Prostate Cancer UK  
Europa UOMO  
Movember Foundation

## References

1. Bruinsma, S.M., et al. Expert consensus document: Semantics in active surveillance for men with localized prostate cancer - results of a modified Delphi consensus procedure. *Nat Rev Urol*, 2017. 14: 312..
2. Mottet, N., Van den Bergh, RCN, Briers, E, Bourke, L, Cornford, P, De Santis, M, Gillissen, S, Govorov, A, Grummet, J, Henry, AM, Lam, TB, Mason, MD, van der Poel, HG, van der Kwast TH, Rouviere, O, Wiegel, T., EAU-ESTRO-ESUR-SIOG Guidelines on Prostate Cancer E.G. Office, Editor. 2018, European Association of Urology: Arnhem, The Netherlands
3. Donovan JL, Hamdy FC. Time for a "Radical" Change to Active Surveillance for Prostate Cancer? *Eur Urol*. 2018 Sep;74(3):281-282. doi: 10.1016/j.eururo.2018.05.009. Epub 2018 May 17. PMID: 297798604.
4. Moher, D., et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*, 2009. 6: e1000097.
5. Willemse, P., Lam, T, Mottet, N, Yuan, C, Donaldson J, Davis, N, Dell'Oglio, P, Fankhauser, C, Grivas, N, Ingels, A, Lardas, M, Liew, M, Pang, K, Paterson, C, Omar, I, Zattoni, F, Buddinh, T. Systematic review of deferred treatment with curative intent for localised prostate cancer to explore heterogeneity of definitions, thresholds and criteria and clinical effectiveness. 2018.
6. MacLennan, S., et al. A core outcome set for localised prostate cancer effectiveness trials. *BJU Int*, 2017. 120: E64.
7. van der Poel, H.G., et al. Sentinel node biopsy for prostate cancer: report from a consensus panel meeting. *BJU Int*, 2017. 120: 204.
8. COMET Initiative DelphiManager. 2018.
9. Everywhere, P. Poll Everywhere. <https://www.polleverywhere.com>
10. Bruinsma SM, Zhang L, Roobol MJ, Bangma CH, Steyerberg EW, Nieboer D, Van Hemelrijck M; Movember Foundation's Global Action Plan Prostate Cancer Active Surveillance (GAP3) consortium. The Movember Foundation's GAP3 cohort: a profile of the largest global prostate cancer active surveillance database to date. *BJU Int*. 2018 May;121(5):737-744. doi: 10.1111/bju.14106.