

This is a repository copy of *Evaluation of a nudge intervention providing simple feedback to clinicians of the consequence of radiation exposure on demand for computed tomography: a controlled study*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/145535/>

Version: Accepted Version

Article:

Lewis, S, Young, Ben orcid.org/0000-0001-6451-9360, Thurley, P et al. (7 more authors) (2019) Evaluation of a nudge intervention providing simple feedback to clinicians of the consequence of radiation exposure on demand for computed tomography: a controlled study. *Clinical Medicine*. pp. 290-293. ISSN 1473-4893

<https://doi.org/10.7861/clinmedicine.19-4-290>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Evaluation of a nudge intervention providing simple feedback to clinicians of the consequence of radiation exposure on demand for computerised tomography scans: a prospective, controlled study

Sarah Lewis* PhD
Ben Young* MSc
Pete Thurley FRCP
Dominic Shaw* PhD
Jo Cranwell** PhD
Rob Skelly FRCP
Tessa Langley* PhD
Mark Norwood MBA
Nigel Sturrock FRCP
Andrew W Fogarty* FRCP

Royal Derby Hospital, Uttoxeter Rd, Derby, DE22 3NE

*Division of Epidemiology and Public Health, University of Nottingham, Clinical Sciences Building, City Hospital, Nottingham NG5 1PB, UK

** Department for Health, Claverton Down, University of Bath, Bath, BA2 7AY.

Correspondence and requests for reprints to Andrew Fogarty

Email: andrew.fogarty@nottingham.ac.uk

Telephone: 0044 115 8231713

Funding: The Health Foundation

Key words: CT scan, nudge, age, demand, radiation, education

Word count: 1252 words

Author contributions: The study was designed by all of the authors except BY. The intervention was implemented by AF, PT, JC, NS and MN. The data were analysed by SL. All authors contributed to the data analysis and approved the final manuscript.

Conflicts of interest: None.

Abstract:

Computerised tomography (CT) is readily available in developed countries. As one of the side-effects includes an increased risk of cancer, interventions that may encourage more judicious use of CT scans are important. Behavioural economics theory includes the use of nudges that aim to help more informed decisions to be made, although these have been rarely used in hospitals to date. We aimed to evaluate the impact of a simple educational message appended to the CT scan report on subsequent numbers of CT scans completed using a controlled interrupted time series design based in two teaching hospitals in X. The intervention was the addition of a non-directional educational message on the risk of ionising radiation to all CT scan reports. There was a statistically significant reduction in the number of CT scans requested in the intervention hospital compared to the control hospital (-4.6%, 95% confidence intervals -7.4 to -1.7, $p = 0.002$) in the 12 months after the intervention was implemented. We conclude that a simple, non-directional nudge intervention has the capacity to modify clinician use of CT scans. This approach is cheap, and has potential in helping support doctors make informed decisions.

Introduction:

Computerised tomography (CT) scanning has undoubtedly been one of the major factors contributing to improvements in medical care over recent decades. As there is widespread variation in usage of diagnostic tests ¹, it is likely that some are unnecessary, which in the case of CT scans exposes the patient to avoidable ionising radiation. The estimates of the magnitude of this risk are uncertain; an estimate from 2007 suggested that use of CT scans in that year may account for 1.5 to 2% of new cancers in the USA ², although this would be expected to decrease as newer generation CT scanners deliver less ionising radiation exposure. Clinicians are not always aware of the long-term adverse health consequences of CT scanning ^{3,4}, suggesting that this is an area with potential for improvement. It was recently demonstrated that there is almost a three-fold increase in the risk of receiving a CT scan during the period of transition from paediatric to adult medical care ⁵, and one interpretation of these data is that paediatric clinicians may be better in using alternatives to CT scans than their adult counterparts.

Nudge theory is a category of behavioural economics that aims to use non-directional interventions to modify decision making ⁶. This generally aims to use opportunities to provide information or context that may allow better informed decision making, at a relatively low cost in terms of money or time. We hypothesised that providing referring clinicians with a simple statement about the consequences of radiation exposure may modify their subsequent decision making with regard to future use of CT scans. We tested this hypothesis in a busy teaching hospital in the UK using a prospective, controlled study design using the demand for CT scans as the primary outcome measure.

Methods:

Study design: We used a controlled interrupted time series design.

Study population: The setting was X, UK (X), with Y (Y) as a control hospital being the nearest large hospital serving a similar population. The X is a busy acute medical hospital that has 1100 beds ⁷. The study was an evaluation of a health service modification and no ethical approval was required.

Intervention: The intervention comprised the addition of the message below to the bottom of all inpatient and outpatient paper and electronic CT scan reports. It was designed to highlight the type of patient who is most at risk after exposure to ionising radiation.

“Message from the executive medical director: “did you know that a chest, abdo and pelvis CT scan in a 20 year old female population is associated with approximately a 1 in 300 risk of subsequent cancer? The equivalent risk is much lower in 90 year old men (less than 1 in 3000). Is there an equally effective alternative investigation that does not involve ionising radiation? If so, have you discussed all of the alternatives with your patient?” [Http://www.xrayrisk.com/index.php](http://www.xrayrisk.com/index.php) “

Outcome measures: Data on the number of weekly CT scans between February 2016 and February 2018 were collected, covering a period 12 months before the intervention was implemented to 12 months afterwards in X and Y. A sensitivity analysis omitted CT scans of the head as these involve a lower ionising radiation dose.

Statistical analysis: Segmented regression was used for the interrupted time series analysis, using a generalised linear model and assuming a Negative Binomial error distribution, that is, we fitted a model which was appropriate to count data with “overdispersion” where there is greater variability in the data than that normally expected in count data. We used the package `tscout` in R statistical software⁸. We initially fitted a full model for a controlled interrupted time series, with terms for the pre-intervention trend (to allow for any underlying baseline trend in CT scan rates), baseline difference between hospitals (to allow for a difference in mean CT scan rates between hospitals at baseline), difference in the pre-intervention trend between hospitals (to allow for any difference in the underlying trend between hospitals), change in level for Y (to allow for any change in CT scan rates in the control hospital post intervention due to confounding factors), change in trend for Y (to allow for any change in trend post intervention in the control hospital due to confounding factors), difference in the post-intervention change in level between X and Y (our measure of the effect of the intervention on the mean level of CT scan rates in the intervention hospital compared to any change in the control hospital), and difference in the post-intervention change in trend between X and Y (our measure of the effect of the intervention on the trend in CT scan rates in the intervention hospital compared to the control hospital). We therefore hypothesised an impact model for the intervention which might involve an immediate change in level or a gradual trend change in CT scan counts in hospital X, which were tested by the last two parameters described above. A dummy variable was fitted to allow for the annual drop in CT scan numbers over Christmas.

We also fitted parsimonious (simplest) models including only parameters that were statistically significant, identified by backward elimination, removing those terms consecutively that were least significant until all parameters were significant at the 5% level and no parameters were significant at the 5% level when added to the model.^{9,10} Models were checked for autocorrelation (that is, for

the fact that observations near in time may be correlated). An autoregressive process of order 4 was added to the final model to allow for each observation to be correlated with the previous 4 observations

Results:

The weekly numbers of CT scans in X (intervention institution) and Y (control institution) are shown in Figure 1. Rates were slightly higher in Y than X, but the underlying baseline trends were comparable.

The results of the resulting segmented regression analysis are shown in Table 1. In the full model, there was a significant difference in numbers of CT scans at baseline between hospitals, and a significant trend for increasing rates of CT scans at baseline, but no significant difference in the baseline trend between hospitals ($P = 0.8$), and the level ($P = 0.055$) and trend ($P = 0.3$) for Y did not change significantly post intervention. There was also no significant difference in the change in trend post intervention ($P = 0.3$), in other words there was no significant effect of the intervention on the trend in CT scan rates.

In the final parsimonious model, there were 29% fewer CT scans carried out in X than Y pre-intervention ($P < 0.001$), and a significant increasing trend in the number of CT scans at 0.1% increase per week in both hospitals. However, there was a statistically significant reduction in level of CT scans in X compared to Y post-intervention of 4.6% ($p = 0.002$). The results excluding head scans were very similar.

Discussion

Statement of principal findings

These data demonstrate that providing simple, non-directional feedback on the long-term consequences of exposure to ionising radiation is associated with a 4.6% decrease in the number of CT scans completed compared to the control hospital over one year.

Strengths and weaknesses of the study

This study has a number of strengths and limitations. The use of a prospective study design with a control group allowed the change in the number of CT scans delivered to be compared against the background of a trend of increasing demand for health care annually. The intervention was added to the radiology reporting system and the text of the message was placed underneath the CT scan

report, which all clinicians see when reviewing the scan result. The message on the CT scan report was designed collaboratively within a multidisciplinary team and piloted, but we are unable to conclude that it is optimal, and there is likely to be room for improvement in both the content and presentation of the message. As a consequence of designing and delivering 'nudge' informational messages in the context of health services research, there were a number of factors that required to be taken into consideration such as the opinions of important stakeholders such as the radiologists, and the limitations of the institutional information technology systems. The data that the educational ionising radiation message was based on is necessarily population based, as it is impossible to provide precise individual level data on radiation risk. The key assumption in the controlled interrupted time series is that the change in the level and/or trend in the outcome variable is presumed to be the same for both the control group and, counterfactually, for the treatment group had it not received the intervention. These hospitals were reasonably similar at baseline with similar increasing trends in CT scans at baseline, no significant change in level or trend in the control group at the point of the intervention, and we are not aware of any other reason to expect differences in the change in levels or trends over time between the hospitals other than the intervention. We are unable to exclude the possibility of a second intervention being implemented at the same time as our educational one, but as most of the study team work in X, this is unlikely. We have not conducted any sex or age stratified secondary subgroup analyses, as the primary hypothesis was that there was be a change in demand for CT scans at the institutional level, and we wished to avoid the risk of multiple hypothesis testing.

As there were 52 paired data points before and after the intervention, we have refrained from secondary data sub-analyses, to avoid the problem of multiple hypothesis testing within a limited number of datapoints, which also be relatively underpowered compared to our primary analysis. These findings represent a positive proof-of-concept study, and demonstrates that the ionising radiation educational message intervention is one that can be implemented, within the context of the UK health service. The definitive study would be a larger, randomised controlled trial of more healthcare institutions to provide proof that this intervention is generalisable to other similar hospitals. This would be a large undertaking, that would require substantial resources to deliver.

Meaning of the study: possible explanations and implications for clinicians and policymakers

Nudge theory has potential in helping doctors make more informed decisions when choosing the optimal imaging approach for their patients; weighing up the immediate benefits of diagnostic knowledge against the longer-term consequences. As the information provided is non-directional, and simply informative, the doctors' clinical autonomy is not infringed upon, as it is added to the context of many years' medical training and experience for many UK healthcare teams. Providing

the information after a scan has been completed and reported informed future practice, and had been demonstrated to be effective in modifying physicians' behaviour with regard to use of blood tests after cost feedback was provided with the results ¹¹. This approach was preferred to the alternative of delivering the information at the time of deciding to do the scan, when it could impact on the efficient delivery on clinical care. In the UK, the decision to request a CT scan for a patient is generally made in a face-to-face discussion with the patient, when the benefits and costs can be weighed up, and a collective decision made between the physician and the patient. The CT examination is then subsequently requested, when the decision to scan has already been made. As a consequence, feeding back an educational message on the longer-term health effects of exposure to ionising radiation was the only viable intervention in this context, although point-of care interventions can be implemented for other healthcare interventions, and have been effective in increasing prescribing of cheaper antibiotics when the price was presented at the point of antibiotics selection ¹².

Unanswered questions and future research


In summary, we have demonstrated that adding a simple educational ionising radiation awareness message to CT scan reports was associated with a 4.6% decrease in subsequent demand for CT scans compared to the control group or approximately 45 fewer CT scans delivered per week or 2340 less CT scans over a year. This nudge approach has potential to improve patient care when used prudently. However, it is unclear in which settings it will be efficacious and more controlled intervention studies are required across a broad range medical areas to develop the evidence base. We anticipate that informing physicians of the long-term risks of ionising radiation is unlikely to negatively impact on healthcare delivery, but future studies should consider both the benefits and the risks of such interventions.

References

1. Kennedy P, Leathley C, Hughes C. Clinical Practice Variation. *Med J Aust.* 2010;193:S97-S99.
2. Brenner D, Hall E. Computerised tomography - an increasing source of radiation exposure. *New Eng J Med.* 2007;357:2277-2284.
3. Puri S, Hu R, Quazi R, Voci S, Veasie P, Block R. Physician and midlevel providers' awareness of lifetime radiation attributable cancer risk from commonly performed CT scan and its relationship to their practice behavior. *Am J Roentgenology.* 2012;199:1328-1336.
4. Lee C, Haims A, Monico E, Brink J, Forman H. Diagnostic CT scans: Assessment of patient, physician, and radiologist awareness of radiation dose and possible risks. *Radiology.* 2004;231:393-398.
5. Thurley P, Crookdake J, Norwood M, Sturrock N, Fogarty A. Demand for CT scans increases during transition from paediatric to adult care: an observational study from 2009 to 2015. *British J Radiology.* 2017;90:20170467.
6. Thaler R, Sunstein C. *Nudge. Improving decisions about health, wealth and happiness.* London: Penguin; 2008.

7. Royal Derby Teaching Hospital Foundation Trust Report. Annual Report 2017-2018. <http://www.derbyhospitalsnhs.uk/EasysiteWeb/getresourceaxd?AssetID=342417&type=full&servicetype=Inline> (accessed 17/8/2018). 2018.
8. Liboschik T, Fokianos K, Fried R. An R package for analysis of count time series following generalized linear models. *J Statistical Software*. 2017;82:1-51. doi: 10.18637/jss.v18082.i18605.
9. Wagner AK, Soumerai SB, Zhang F, Ross-Degnan D. Segmented regression analysis of interrupted time series studies in medication use research. *Journal of Clinical Pharmacy and Therapeutics*. 2002;27(4):299-309.
10. Penfold RB, Zhang F. Use of Interrupted Time Series Analysis in Evaluating Health Care Quality Improvements. *Acad Pediatr*. 2013;13:38-44.
11. Fogarty A, Sturrock N, Premji K, Prinsloo P. Hospital clinicians' responsiveness to assay cost feedback: A prospective blinded controlled intervention study. *JAMA Internal Medicine*. 2013;173:1654-1655.
12. Langley T, Lacey J, Johnson A, et al. An evaluation of a price transparency intervention for two commonly prescribed medications on total institutional expenditure: a prospective study. *Future Healthcare J*. 2018;5:198-202.

Figure 1: Weekly total CT scans for X (interventional institution) and Y (control institution), before and after the intervention.



Intervention implementation in February 2017 shown as a dotted line.

Table 1: Change in the number of CT scans completed at X (intervention institution) compared to Y (control institution) after the informational feedback intervention was implemented

	Rate ratio (95% CI) for all CT scans (full model)	Rate ratio (95% CI) for all CT scans (parsimonious model)	Rate ratio (95% CI) for CT scans other than head (full model)	Rate ratio (95% CI) for CT scans other than head (parsimonious model)
Pre-intervention difference between X and Y	0.799 (0.753 to 0.847) P < 0.001	0.811 (0.773 to 0.852) P<0.001	0.779 (0.727 to 0.835) P < 0.001	0.794 (0.752 to 0.839) P<0.001
Pre-intervention trend (slope) for control Y (per week)	1.001 (1.000 to 1.002) P = 0.029	1.001 (1.001 to 1.002) P<0.001	1.001 (1.000 to 1.002) P = 0.14	1.002 (1.001 to 1.002) P<0.001
Pre-intervention difference in trend between X and Y	1.000 (0.999 to 1.002) P = 0.8	-	1.000 (0.999 to 1.002) P = 0.6	-
Change in level post-intervention for control Y	1.047 (0.999 to 1.097) P = 0.055	-	1.052 (0.993 to 1.114) P = 0.08	-
Change in trend post-intervention for control Y	0.999 (0.998 to 1.001) P = 0.3	-	1.000 (0.998 to 1.002) P = 0.7	-
Difference in the change in level between X and Y	0.904 (0.847 to 0.965) P = 0.002	0.954 (0.926 to 0.983) P=0.002	0.879 (0.811 to 0.953) P = 0.002	0.949 (0.915 to 0.985) P=0.005
Difference in the change in trend between X and Y	1.001 (0.999 to 1.003) P = 0.3	-	1.002 (0.999 to 1.004) P = 0.2	-

A dummy variable for the Christmas week was also included in the model.

Only statistically significant variables are included in the parsimonious models. Gaps in the table reflect that variables were not significant and hence were not included in the final model.

CI: Confidence intervals