



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/145621/>

Version: Published Version

---

**Article:**

Robertson, Lindsay and Nandhra, Sandip (2017) Laparoscopic surgery for elective abdominal aortic aneurysm repair. Cochrane Database of Systematic Reviews. CD012302. pp. 1-25. ISSN: 1469-493X

<https://doi.org/10.1002/14651858.CD012302.pub2>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



**Cochrane**  
**Library**

Cochrane Database of Systematic Reviews

## Laparoscopic surgery for elective abdominal aortic aneurysm repair (Review)

Robertson L, Nandhra S

Robertson L, Nandhra S.

Laparoscopic surgery for elective abdominal aortic aneurysm repair.

*Cochrane Database of Systematic Reviews* 2017, Issue 5. Art. No.: CD012302.

DOI: 10.1002/14651858.CD012302.pub2.

[www.cochranelibrary.com](http://www.cochranelibrary.com)

## TABLE OF CONTENTS

HEADER . . . . .	1
ABSTRACT . . . . .	1
PLAIN LANGUAGE SUMMARY . . . . .	2
SUMMARY OF FINDINGS FOR THE MAIN COMPARISON . . . . .	4
BACKGROUND . . . . .	6
OBJECTIVES . . . . .	7
METHODS . . . . .	7
RESULTS . . . . .	10
Figure 1. . . . .	10
Figure 2. . . . .	11
Figure 3. . . . .	12
DISCUSSION . . . . .	13
AUTHORS' CONCLUSIONS . . . . .	14
ACKNOWLEDGEMENTS . . . . .	14
REFERENCES . . . . .	14
CHARACTERISTICS OF STUDIES . . . . .	16
DATA AND ANALYSES . . . . .	19
Analysis 1.1. Comparison 1 Hand-assisted laparoscopic repair vs EVAR, Outcome 1 In-hospital mortality. . . . .	19
Analysis 1.2. Comparison 1 Hand-assisted laparoscopic repair vs EVAR, Outcome 2 Operative time (minutes). . . . .	20
Analysis 1.3. Comparison 1 Hand-assisted laparoscopic repair vs EVAR, Outcome 3 Major complications. . . . .	20
APPENDICES . . . . .	20
CONTRIBUTIONS OF AUTHORS . . . . .	22
DECLARATIONS OF INTEREST . . . . .	22
SOURCES OF SUPPORT . . . . .	22
DIFFERENCES BETWEEN PROTOCOL AND REVIEW . . . . .	23
INDEX TERMS . . . . .	23

[Intervention Review]

# Laparoscopic surgery for elective abdominal aortic aneurysm repair

Lindsay Robertson<sup>1</sup>, Sandip Nandhra<sup>2</sup>

<sup>1</sup>Department of Vascular Surgery, Freeman Hospital, Newcastle upon Tyne, UK. <sup>2</sup>Department of Vascular Surgery, Health Education North East, Durham, UK

Contact address: Lindsay Robertson, Department of Vascular Surgery, Freeman Hospital, Newcastle upon Tyne Hospitals NHS Foundation Trust, High Heaton, Newcastle upon Tyne, NE7 7DN, UK. [lindsay.robertson@nuth.nhs.uk](mailto:lindsay.robertson@nuth.nhs.uk), [lindsay.robertson@ed.ac.uk](mailto:lindsay.robertson@ed.ac.uk).

**Editorial group:** Cochrane Vascular Group.

**Publication status and date:** New, published in Issue 5, 2017.

**Citation:** Robertson L, Nandhra S. Laparoscopic surgery for elective abdominal aortic aneurysm repair. *Cochrane Database of Systematic Reviews* 2017, Issue 5. Art. No.: CD012302. DOI: 10.1002/14651858.CD012302.pub2.

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

## ABSTRACT

### Background

Abdominal aortic aneurysm (AAA) is an abnormal dilatation of the infradiaphragmatic aorta that is equal to or greater than 30 mm or a local dilatation of equal to or greater than 50% compared to the expected normal diameter of the artery. AAAs rarely occur in individuals under 50 years of age, but thereafter the prevalence dramatically increases with age, with men at a six-fold greater risk of developing an AAA than women. Prevalence of AAA has been reported to range from 1.3% in women aged 65 to 80 years to between 4% and 7.7% in men aged 65 to 80 years.

There is evidence that the risk of rupture increases as the aneurysm diameter increases from 50 mm to 60 mm. People with AAAs over 55 mm in diameter are therefore generally referred for consideration of repair, as the risk of rupture exceeds the risk of repair. The traditional treatment for AAA is open surgical repair (OSR) which involves a large abdominal incision and is associated with a significant risk of complications. Two less invasive procedures have recently become more widely used: endovascular aneurysm repair (EVAR) and laparoscopic repair. EVAR is carried out through sheaths inserted in the femoral artery in the groin; thereafter, a stent graft is placed within the aneurysm sac under radiological image guidance and anchored in place to form a new channel for blood flow. Laparoscopic repair involves the use of a laparoscope which is inserted through small cuts in the abdomen and the synthetic graft is sewn in place to replace the weakened area of the aorta. Laparoscopic AAA repair falls into two categories: hand-assisted laparoscopic surgery (HALS), where an incision is made to allow the surgeon's hand to assist in the repair; and total laparoscopic surgery (TLS). Both EVAR and laparoscopic repair are favourable over OSR as they are minimally invasive, less painful, associated with fewer complications and lower mortality rate and have a shorter duration of hospital stay.

Current evidence suggests that elective laparoscopic AAA repair has a favourable safety profile comparable with that of EVAR, with low conversion rates as well as similar mortality and morbidity rates. As a result, it has been suggested that elective laparoscopic AAA repair may have a role in treating those patients for whom EVAR is unsuitable.

### Objectives

To assess the effects of laparoscopic surgery for elective abdominal aortic aneurysm repair.

The primary objective of this review was to assess the perioperative mortality and operative time of laparoscopic (total and hand-assisted) surgical repair of abdominal aortic aneurysms (AAA) compared to traditional open surgical repair or EVAR. The secondary objective was to assess complication rates, all-cause mortality (> 30 days), hospital and intensive care unit (ICU) length of stay, conversion and re-intervention rates, and quality of life associated with laparoscopic (total and hand-assisted) surgical repair compared to traditional open surgical repair or EVAR.

## Search methods

The Cochrane Vascular Information Specialist (CIS) searched the Specialised Register (last searched August 2016) and CENTRAL (2016, Issue 7). In addition the CIS searched trials registries for details of ongoing or unpublished studies. We searched the reference lists of relevant articles retrieved by electronic searches for additional citations.

## Selection criteria

Randomised controlled trials and controlled clinical trials in which patients with an AAA underwent elective laparoscopic repair (total laparoscopic repair or hand-assisted laparoscopic repair) compared with either open surgical repair or EVAR.

## Data collection and analysis

Studies identified for potential inclusion were independently assessed for inclusion by at least two review authors.

## Main results

One randomised controlled trial with a total of 100 male participants was included in the review. The trial compared hand-assisted laparoscopic repair with EVAR and provided results for in-hospital mortality, operative time, length of hospital stay and lower limb ischaemia. The included study did not report on the other pre-planned outcomes of this review. No in-hospital deaths occurred in the study. Hand-associated laparoscopic repair was associated with a longer operative time (MD 53.00 minutes, 95% CI 36.49 to 69.51) than EVAR. The incidence of lower limb ischaemia was similar between the two treatment groups (risk ratio (RR) 0.50, 95% confidence interval (CI) 0.05 to 5.34). The mean length of hospital stay was 4.2 days and 3.4 days in the hand-assisted laparoscopic repair and EVAR groups respectively but standard deviations were not reported and therefore it was not possible to independently test the statistical significance of this result. The quality of evidence was downgraded for imprecision due to the inclusion of one small study; and wide confidence intervals and indirectness due to the study including male participants only. No study compared laparoscopic repair (total or hand-assisted) with open surgical repair or total laparoscopic surgical repair with EVAR.

## Authors' conclusions

There is insufficient evidence to draw any conclusions about effectiveness and safety of laparoscopic (total and hand-assisted) surgical repair of AAA versus open surgical repair or EVAR, because only one small randomised trial was eligible for inclusion in this review. High-quality randomised controlled trials are needed.

## PLAIN LANGUAGE SUMMARY

### Laparoscopic surgery for abdominal aortic aneurysm

#### Background

An abdominal aortic aneurysm (AAA) is an abnormal widening of the abdominal aorta, the main artery supplying blood to the organs in the abdomen and lower part of the body. Between 4% and 7% of men over 65 years of age have an AAA, but it is less common in women. Aneurysms over 55 mm in diameter carry a high risk of rupture which can lead to death; approximately 60% of people with a ruptured AAA die before reaching hospital. People with AAAs over 55 mm are generally referred for repair, as the risk of rupture exceeds the risk of repair. There are three methods of repairing an AAA: surgery, endovascular aneurysm repair (EVAR) and laparoscopic repair. Surgery involves making a large cut in the abdomen, after which the abdominal aorta is exposed and opened and a synthetic graft (tube) is sewn in place to replace the weakened area of the aorta. EVAR involves making a cut in the groin area, after which a stent graft is inserted in collapsed form and opened inside the aneurysm under x-ray guidance and held in place with a stent. Laparoscopic repair or 'keyhole' AAA surgery is carried out by making very small cuts in the patient's abdomen, after which a fine telescope (a laparoscope) is inserted through these cuts and the synthetic graft is sewn in place. The benefits of EVAR and laparoscopic repair are that they require smaller incisions, are less painful, have fewer complications, a lower mortality rate and shorter hospital stay than surgical repair. Current evidence suggests that EVAR is the preferred approach for AAA repair. However laparoscopic AAA repair has been suggested as a safe and effective alternative in treating those patients for whom EVAR is unsuitable. This review aimed to assess the effects of laparoscopic surgery for abdominal aortic aneurysms.

#### Study characteristics and key results

One randomised controlled trial (current until August 2016), studying 100 male participants and comparing hand-assisted laparoscopic repair with EVAR, was included in this review. No in-hospital deaths occurred during the study. The trial showed that hand-assisted

laparoscopic repair took longer to perform than EVAR but there was no difference in the number of patients with reduced blood flow to the leg following either treatment.

#### **Quality of evidence**

At present, there is a lack of randomised controlled trials examining the comparative effectiveness and safety of laparoscopic repair of AAA. The quality of the available evidence was imprecise due to the inclusion of one small study and wide confidence intervals; and indirect because the study includes male participants only.

#### **Conclusions**

Further research is required before conclusions can be made.

## SUMMARY OF FINDINGS FOR THE MAIN COMPARISON *[Explanation]*

Hand-assisted laparoscopic repair compared to EVAR for elective abdominal aortic aneurysm repair						
<b>Patient or population:</b> People undergoing elective abdominal aortic aneurysm repair						
<b>Setting:</b> Hospital						
<b>Intervention:</b> Hand-assisted laparoscopic AAA repair						
<b>Comparison:</b> EVAR						
Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No. of participants (studies)	Quality of the evidence (GRADE)	Comments
	Risk with EVAR	Risk with hand-assisted laparoscopic repair				
In-hospital mortality	see comment		Not estimable	100 (1 RCT)	⊕⊕○○ LOW <sup>12</sup>	0 participants died while in hospital in this study
Operative time (minutes)	The mean operative time was 125 minutes	The mean operative time was 53 minutes longer (36.49 longer to 69.51 longer)	-	100 (1 RCT)	⊕⊕○○ LOW <sup>12</sup>	
Major complications - lower limb ischaemia (up to 12 months)	Study population 40 per 1000	20 per 1000 (2 to 214)	RR 0.50 (0.05 to 5.34)	100 (1 RCT)	⊕⊕○○ LOW <sup>12</sup>	
Long-term complications (12 months or longer if reported)	see comment					outcome not reported
All-cause mortality/survival (> 30 days)	see comment					outcome not reported

Length of ICU stay (days)	see comment					outcome not reported
Overall length of hospital stay (days)	The mean length of hospital stay was 3.4 days	The mean length of hospital stay was 4.2 days	Not estimable	100 (1 RCT)	⊕⊕○○ LOW <sup>12</sup>	Standard deviations around the mean length of hospital stay were not reported

\***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

**CI:** confidence interval; **EVAR:** endovascular repair; **ICU:** intensive care unit; **RR:** risk ratio

**GRADE Working Group grades of evidence**

**High quality:** we are very confident that the true effect lies close to that of the estimate of the effect

**Moderate quality:** we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

**Low quality:** our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect

**Very low quality:** we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect

<sup>1</sup> Risk of bias was unclear for detection bias in the included studies but we did not consider it sufficient enough to downgrade the quality of the evidence

<sup>2</sup> Quality of evidence was downgraded for imprecision due to the inclusion of one small study and wide confidence intervals and indirectness due to the study including male participants only

## BACKGROUND

### Description of the condition

Abdominal aortic aneurysm (AAA) is an abnormal dilatation of the infradiaphragmatic aorta that is equal to or greater than 30 mm (Sakalihan 2005; Wanhainen 2008); or a local dilatation of equal to or greater than 50% compared to the expected normal diameter of the artery (Johnston 1991).

AAAs rarely occur in individuals under 50 years of age (Lederle 2000a), but thereafter the prevalence dramatically increases with age, with men at a six-fold greater risk of developing an AAA than women (Pleumeekers 1995; Scott 2002).

Prevalence of AAA has been reported to range from 1.3% in women aged 65 to 80 years to between 4% and 7.7% in men aged 65 to 80 years (Ashton 2002; Ashton 2007; Lindholt 2005; Nordon 2011; Norman 2004; Scott 2002). The annual incidence of AAA in Western populations has been estimated at between 0.4% and 0.67% (Forsdahl 2009; Lederle 2000b; Nordon 2011; Vardulaki 1999); however more recent evidence suggests that AAA incidence is decreasing, most likely because of a reduction in tobacco smoking (Anjum 2012).

The development of an AAA is often multifactorial, with a change in the composition of the collagen and elastin matrix in the arterial wall attributed to atherosclerosis and inflammation of the aortic wall (Shah 1997). Aside from age and sex, the main well-established modifiable risk factor is cigarette smoking, with smokers having a two- to three-fold increased risk of AAA compared to non-smokers (Lederle 1997; Lederle 2003). There is an apparent genetic correlation, with aneurysms tending to occur more frequently in close relatives of people who have suffered an AAA, but a mode of inheritance has not been demonstrated (Ballard 1999). The natural progression of AAA can vary considerably between individuals (Ernst 1993), with subsequent variation in presentation ranging from no symptoms to symptoms such as groin, back, or abdominal pain. Findings upon physical examination include pulsating abdominal masses, and co-existing aneurysmal popliteal or femoral arteries, and bruits. As the dilatation is often asymptomatic, the AAA can expand to such an extent that it ruptures, which is a surgical emergency. Approximately 60% of people with ruptured AAA die before reaching hospital (Ballard 1999), and even when it is possible to perform an emergency open surgical repair, the mortality rate remains high, at between 35% to 70% (Veith 2003). Newer techniques such as endovascular aneurysm repair have been shown to be more cost-effective in the emergency setting but do not confer a greater chance of survival (Sweeting 2015).

The decision to repair an AAA is made on an individual basis, balancing the risk of treatment against the risk of aneurysm rupture. There is evidence that the risk of rupture increases as the aneurysm diameter increases from 50 mm to 60 mm (Brewster 2003). People with an AAA over 55 mm in diameter are therefore

generally referred for consideration of repair, as the risk of rupture exceeds the risk of repair. Elective open surgical repair has a mortality of just over 5% (Bush 2006). Aneurysms below or equal to 55 mm are termed small AAAs and are at a low risk of rupture. Despite the improved mortality rates for elective versus emergency aneurysm repair, the management of patients with small AAAs is one of surveillance, whereby the AAA is routinely monitored for growth through ultrasound imaging (Filardo 2015). However, women with a small AAA have a higher rupture rate than men, so that a lower threshold (52 mm) is suggested (Brewster 2003).

### Description of the intervention

The traditional treatment for AAA is open surgical repair (OSR): the abdominal aorta which lies in the retroperitoneum is exposed and the aneurysm is clamped and opened; then a prosthetic graft is anastomosed proximally to the normal aorta and distally to the dilated aorta and the clamps are released with the return of normal blood flow. OSR involves a large abdominal incision and is associated with a significant risk of complications including myocardial infarction, arrhythmias, bleeding, injury to the bowel, limb ischaemia, embolus, infection, and kidney damage (Badger 2014). The 30-day mortality associated with open repair of AAA in the UK Small Aneurysm Trial was 5.8% (TUKSAT 1998), and reported rates range from 2% to 7% in otherwise fit individuals (Paravastu 2014).

A less invasive procedure known as endovascular aneurysm repair (EVAR) has recently become widely used. EVAR is carried out through sheaths inserted in the femoral artery in the groin. A modular covered stent graft is placed within the aneurysm sac, under radiological image guidance. The graft is manipulated into place using guidewires and anchored in place to form a new channel for blood flow (Parodi 1991). The aneurysm is excluded from inside to prevent further expansion and possible rupture. EVAR is favourable over OSR, as it is minimally invasive compared to major abdominal surgery of OSR. EVAR procedures are less painful and have a shorter recovery time and therefore a shorter duration of hospital stay (Paravastu 2014). Furthermore, in EVAR the blood flow to organs and the lower limbs is not disrupted to the same extent as in OSR, there are fewer respiratory side effects, and less blood is lost during the procedure (Paravastu 2014).

A third treatment for AAA is laparoscopic repair of AAA. There are two well-described methods of laparoscopic repair of AAA. The first is hand-assisted laparoscopic surgery (HALS), where an incision is made to allow the surgeon's hand to assist in the repair. The second, and more challenging technically, is total laparoscopic surgery (TLS) (NICE 2007). Both procedures require a laparoscope inserted through the abdominal wall via access ports and an induction of a pneumoperitoneum with further access ports. Laparoscopic instruments are used to dissect and clip the lumbar arteries and the inferior mesenteric artery (NICE 2007). The areas above and below the aneurysm are clamped, the sac of the

aneurysm is cut open, any thrombus is removed, and a prosthetic vascular graft is anastomosed to both sides of the aorta. The posterior parietal peritoneum or aneurysm sac is then closed over the graft (NICE 2007).

### Hand-assisted laparoscopic surgery

This is generally performed with the patient supine and the use of a mini-laparotomy incision at the midline for the placement of the laparoscopic port. A further incision is made (7 cm to 9 cm) in an appropriate position as determined by preoperative imaging, to allow the non-dominant hand of the surgeon to be introduced into the abdominal cavity. A seal and pneumoperitoneum is created. The table is then tilted to the right and the abdominal bowel loops are pushed away from the midline. Further laparoscopic ports are inserted to allow dissection of the aortic neck. The exposure of the aorta and the procedure follows much the same process as the open AAA repair. Once the proximal and distal anastomosis sites are exposed and dissected, the abdomen is deflated and the proximal clamps applied. Thereafter the aneurysm is opened and the distal back bleeding is stopped using occlusive balloons (such as a Fogarty catheter). The lumbar vessels are ligated using sutures in the standard fashion. The proximal anastomosis is performed using long but standard instruments with the mini-laparotomy incision being auto-retracted and moved cranially to allow visualisation. Following successful proximal anastomosis, the distal anastomoses (either single in a tube graft or bifurcated in a 'Y' graft) are performed. If the anastomosis is at the level of the external iliac then an oblique suprainguinal incision is required. The peritoneum is then irrigated, closed, and the mini-laparotomy incision closed (Ferrari 2006)

### Total laparoscopic surgery

This is performed through a transperitoneal left retro-renal approach. The patient is placed in a dorsal decubitus position with appropriate support and straps. A pneumoperitoneum is induced using a Veress needle. Multiple laparoscopic ports are inserted to allow the mobilisation of the left lateral colonic border, the kidney and spleen, so that they drop medially towards the patient's right side. The aorta is exposed and the left renal artery by fixation of the left kidney under traction sutures. The aorta is dissected and exposed from the iliac to the renal arteries taking care to ligate lumbar arteries and isolate the left ureter. The clamp is placed via the laparoscopic trocars on the proximal aorta, and the aortic wall is held under tension by a suture to allow opening of the aorta. The iliac arteries are then clamped using laparoscopic clamps. A longitudinal arteriotomy is then made in the aorta on the left side and thrombus evacuated. A tied tube graft is then sutured into the proximal neck. Next the distal anastomosis is performed after evacuating any new clot in the graft. Sutures and ties are performed intracorporally and the ports removed under vision with release of the pneumoperitoneum (Javerliat 2006)

### How the intervention might work

Current evidence suggests that EVAR is the preferred approach for AAA repair in suitable candidates, due to shorter hospital stay and lower perioperative morbidity and mortality rates, as opposed to an open surgical approach. Evidence also suggests that elective laparoscopic AAA repair has a favourable safety profile comparable with that of EVAR, with low conversion rates as well as similar mortality and morbidity rates (Ahmed 2014). As a result, it has been suggested that elective laparoscopic AAA repair may have a role in treating those patients for whom EVAR is unsuitable.

### Why it is important to do this review

There is currently no meta-analysis of the literature to help guide evidence-based recommendation for laparoscopic surgery for AAA. The aim of this review is to study the benefits and harms of laparoscopic surgery in people with AAA, by critical appraisal and meta-analysis of the existing literature.

## OBJECTIVES

To assess the effects of laparoscopic surgery for elective abdominal aortic aneurysm repair.

The primary objective of this review was to assess the perioperative mortality and operative time of laparoscopic (total and hand-assisted) surgical repair of abdominal aortic aneurysms (AAA) compared to traditional open surgical repair or EVAR. The secondary objective was to assess complication rates, all-cause mortality (> 30 days), hospital and intensive care unit (ICU) length of stay, conversion and re-intervention rates, and quality of life associated with laparoscopic (total and hand-assisted) surgical repair compared to traditional open surgical repair or EVAR.

## METHODS

### Criteria for considering studies for this review

#### Types of studies

We included randomised controlled trials (RCTs). If we had found no trials of this sort, we would also have considered controlled clinical trials (CCTs). We excluded case reports, case series, and retrospective studies. We applied no limitation on the language of publication or country.

## Types of participants

People of any age undergoing elective repair of an AAA.

## Types of interventions

Laparoscopic repair (total laparoscopic repair or hand-assisted laparoscopic repair) compared with either open surgical repair or EVAR.

We considered the following comparisons.

1. Laparoscopic repair (total or hand-assisted laparoscopic repair) versus open surgical repair.
2. Laparoscopic repair (total or hand-assisted laparoscopic repair) versus EVAR.

## Types of outcome measures

### Primary outcomes

1. 30-day or in-hospital mortality.
2. Operative time (minutes).

### Secondary outcomes

1. Major complications (e.g. myocardial infarction, stroke, renal failure, respiratory failure, bowel ischaemia, lower limb ischaemia, pneumonia, infection) up to 12 months.
2. Long-term (12 months or longer if reported) major complications.
3. All-cause mortality/survival (> 30 days).
4. Length of ICU stay (days).
5. Overall length of hospital stay (days).
6. Open conversion rates (conversion of the repair from a primarily laparoscopic approach to a total open surgical approach, inferring difficulty or complications).
7. Re-intervention rates and device-related complications.
8. Minor complications (e.g. haematoma, wound infection).
9. Completion of repair (successful completion of the anastomosis and repair by the intended method i.e. starting as a laparoscopic repair and completing this laparoscopically).
10. Quality of life.

## Search methods for identification of studies

### Electronic searches

The Cochrane Vascular Information Specialist (CIS) searched the following databases for relevant trials:

- The Cochrane Vascular Specialised Register (searched 10 August 2016).
- The Cochrane Central Register of Controlled Trials (CENTRAL; 2016, Issue 7) in the Cochrane Library (searched 10 August 2016).

See [Appendix 1](#) for details of the search strategy used to search CENTRAL.

The Cochrane Vascular Specialised Register is maintained by the CIS and is constructed from weekly electronic searches of MEDLINE Ovid, Embase Ovid, CINAHL, AMED, and through hand-searching relevant journals. The full list of the databases, journals and conference proceedings which have been searched, as well as the search strategies used, are described in the [Specialised Register](#) section of the Cochrane Vascular module in the Cochrane Library ([www.cochranelibrary.com](http://www.cochranelibrary.com)).

The CIS searched the following trial registries (10 August 2016) for details of ongoing and unpublished studies using the terms 'laparoscopic AND aneurysm';

- ClinicalTrials.gov ([www.clinicaltrials.gov](http://www.clinicaltrials.gov)).
- World Health Organization International Clinical Trials Registry Platform ([www.who.int/trialsearch](http://www.who.int/trialsearch)).
- ISRCTN Register ([www.isrctn.com/](http://www.isrctn.com/)).

See [Appendix 2](#) for details of the search strategies used.

### Searching other resources

We searched citations within identified studies. We planned, if needed, to contact authors of relevant papers by email to identify any unpublished randomised controlled trials.

## Data collection and analysis

### Selection of studies

Two review authors (LR, SN) independently reviewed the results of all searches to identify potentially eligible articles. The two review authors (LR, SN) discussed each study to confirm eligibility for inclusion in the systematic review, excluding those not fulfilling the criteria as described in [Criteria for considering studies for this review](#) and stating the reasons for exclusion in the review.

### Data extraction and management

Two review authors (LR, SN) independently extracted from each study information about the study characteristics, participants, interventions, duration of follow-up, and outcome parameters using standardised forms. We extracted data on the following items, where available.

1. Study design.
2. Number of study participants.
3. Details of participants, including age, sex, diameter of AAA, diagnosis (clinical or ultrasound), and presence of co-morbidities.
4. Stratification of low- and high-risk patients defined by co-morbidities, Vascular Physiological and Operative Severity Score for the enUmeration of Mortality and Morbidity (V(p)-Possum) or Glasgow Aneurysm Score (GAS) where appropriate.

5. Interventions, including type of repair, adverse events (major and minor), and length of hospital stay.
6. Outcome measures as stated above.
7. Length of follow-up.

### Assessment of risk of bias in included studies

Two review authors (LR, SN) independently assessed the design and execution of each study according to the following criteria: random sequence generation; allocation concealment of treatment; blinding of participants, personnel, and outcomes; incomplete outcome data; selective outcome reporting; and other sources of bias in accordance with Cochrane's tool for assessing risk of bias (Higgins 2011). We judged the studies as either low risk of bias, high risk of bias, or unclear (due to either lack of information or uncertainty over the potential for bias). We resolved any disagreements by consensus between the two review authors.

### Measures of treatment effect

We assessed dichotomous data using risk ratios (RRs) with 95% confidence intervals (CIs). We analysed continuous outcomes using mean differences (MDs) with 95% CIs where the scales were the same; and where scales were different but outcome measured was the same, we planned to use the standardised mean difference (SMD) with 95% CIs.

### Unit of analysis issues

We did not include cross-over trials. The individual participant was the unit of analysis.

### Dealing with missing data

Where information was missing, we contacted the authors of the relevant study. If unsuccessful, we planned to exclude the data from the meta-analysis but report it in the review. We intended to include outcome measures in this systematic review only if it was the intention of the study authors to perform the necessary assessments in all randomised participants. When fewer than 50% of the participants in a study had an acceptable follow-up for a particular outcome measure, we planned on not reporting the results of this outcome measure due to the associated high risk of attrition bias.

### Assessment of heterogeneity

It was intended that if the included studies were comparable with regard to age, sex, treatment, and used outcome definitions, we would perform a pooled analysis. We planned on assessing heterogeneity with the use of forest plots; and by a formal statistical test for heterogeneity, the  $I^2$  statistic. Substantial heterogeneity was defined as  $I^2$  greater than 50% (Higgins 2011). We planned on

exploring possible causes of heterogeneity and taking appropriate measures.

### Assessment of reporting biases

If more than 10 studies had been included in a meta-analysis, we would have constructed a funnel plot to graphically ascertain the existence of publication bias (Higgins 2011).

### Data synthesis

We entered the data into the Cochrane Review Manager 5 software (Review Manager 2014), and analysed them according to the guidelines of the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011). We planned to use a fixed-effect model where no substantial heterogeneity was found and a random-effects model if heterogeneity ( $I^2$  greater than 50%) was found.

### Subgroup analysis and investigation of heterogeneity

We planned on performing subgroup analysis according to the following.

- Age.
- Gender.
- Body mass index > 30 kg/m<sup>2</sup>.
- Diabetes.
- Type of laparoscopic repair (total or hand-assisted).
- Previous cardiopulmonary co-morbidities such as ischaemic heart disease, myocardial infarction, or chronic obstructive pulmonary disease (COPD) as determined by a preoperative cardiopulmonary exercise test (CPEX).
  - Participants at high risk of complications, identified where possible using well-validated and reliable scoring systems such as the GAS, V(p)-Possum, or other tests.

We intended to examine heterogeneity for these by visual inspection of forest plots and the  $I^2$  test.

### Sensitivity analysis

We planned on performing a sensitivity analysis by excluding studies at high risk of selection, performance, detection, attrition, and reporting bias.

### Summary of findings table

We presented the main findings of the review results concerning the quality of evidence, the magnitude of effect of the interventions examined, and the sum of available data for outcomes - 30-day or in-hospital mortality, operative time, major complications up to 12 months, long-term (12 months or longer if reported) major complications, all-cause mortality/survival (> 30 days), length of ICU stay, and overall length of hospital stay - in a 'Summary of findings' table, according to Higgins 2011 and Atkins 2004. Since

we planned to assess different comparisons of interventions, we intended developing a 'Summary of findings' table for each comparison. We used the GRADEpro (GRADEproGDT) software ([www.guidelinedevelopment.org/](http://www.guidelinedevelopment.org/)) to assist in the preparation of the 'Summary of findings' table.


## RESULTS

### Description of studies

#### Results of the search

See [Figure 1](#)

**Figure 1. Study flow diagram.**



### Included studies

One randomised controlled trial with 100 male participants was eligible for inclusion in the review (Veroux 2010); it compared hand-assisted laparoscopic repair with EVAR. Although the primary outcome of the study was perioperative sexual dysfunction, the study also measured in-hospital mortality, operative time, length of hospital stay and the major complication incidence of lower limb ischaemia (Veroux 2010).


### Excluded studies

One study comparing laparoscopic versus open AAA repair was withdrawn before enrolment and therefore excluded from this review (NCT00821145).


### Risk of bias in included studies

See Figure 2 and Figure 3.

**Figure 2. Risk of bias graph: review authors' judgements about each risk of bias item presented as percentages across all included studies.**



**Figure 3. Risk of bias summary: review authors' judgements about each risk of bias item for each included study.**



**Allocation**

The included study was judged to be at low risk of selection bias as the study authors reported that randomisation codes were computer-generated and sealed envelopes labelled with participant numbers were used and only opened when a participant was admitted to AAA repair.

authors did not report whether outcome assessors were blinded to treatment and we therefore judged this study to be at unclear risk of detection bias.

**Blinding**

The included study did not report whether participants were blinded to treatment. However, we judged that the study outcomes and outcome measurements reported by the included study were not likely to have been influenced by lack of blinding and therefore classified this study as low risk of performance bias. The study

**Incomplete outcome data**

The included study had no missing outcome data and was therefore judged to be at low risk of attrition bias.

**Selective reporting**

The included study reported all pre-specified outcomes and was therefore judged to be at low risk of reporting bias.

### Other potential sources of bias

The included study appeared to be free from other sources of bias.

### Effects of interventions

See: [Summary of findings for the main comparison](#) Hand-assisted laparoscopic repair compared to EVAR for elective abdominal aortic aneurysm repair

### Laparoscopic repair (total or hand-assisted) versus open surgical repair

We identified no studies which compared either total or hand-assisted laparoscopic repair with open surgical repair of AAA.

### Laparoscopic repair (total or hand-assisted) versus endovascular aneurysm repair

We identified one study which compared hand-assisted laparoscopic repair with EVAR. There were no cases of in-hospital mortality and the incidence of the major complication 'lower limb ischaemia' was similar between the two treatment groups (relative risk (RR) 0.50, 95% CI 0.05 to 5.34). Hand-assisted laparoscopic repair was associated with longer operative time than EVAR (mean difference (MD) 53.00 minutes, 95% CI 36.49 to 69.51). The study reported mean length of hospital stay as 4.2 days and 3.4 days in the hand-assisted and EVAR groups respectively, but standard deviations for the mean length of hospital stay were not reported and therefore it was not possible to independently test the statistical significance of this result. The study did not report on the other outcomes of this review (major complications other than lower limb ischaemia, long-term major complications, all-cause mortality/survival (> 30 days), length of ICU stay, open conversion rates, re-intervention rates, device-related complications, minor complications, completion of repair and quality of life). We identified no studies which compared total laparoscopic repair with EVAR.

## DISCUSSION

### Summary of main results

We identified one randomised controlled trial with a total of 100 male participants which fulfilled the eligibility criteria for inclusion in this review ([Veroux 2010](#)).

### Laparoscopic repair (total or hand-assisted) versus open surgical repair

We identified no studies which compared either total or hand-assisted laparoscopic repair with open surgical repair of AAA.

### Laparoscopic repair (total or hand-assisted) versus endovascular aneurysm repair

We identified one study which compared hand-assisted laparoscopic repair with EVAR. There were no cases of in-hospital mortality and the incidence of the major complication 'lower limb ischaemia' was similar between the two treatment groups. Hand-assisted laparoscopic repair was associated with a longer operative time than EVAR. The study reported mean length of hospital stay but standard deviations for the mean duration were not reported and therefore it was not possible to independently test the statistical significance of this result. The study did not report on the other outcomes of this review (major complications other than lower limb ischaemia, long-term major complications, all-cause mortality/survival (> 30 days), length of ICU stay, open conversion rates, re-intervention rates and device-related complications, minor complications, completion of repair and quality of life). We identified no studies which compared total laparoscopic repair with EVAR.

### Overall completeness and applicability of evidence

At present, there is no randomised controlled trial evidence regarding the efficacy and safety of laparoscopic repair compared with open surgical repair of AAA. Furthermore there is exceptionally limited randomised controlled trial evidence on laparoscopic repair compared with EVAR. Only one study on a total of 100 male participants met the inclusion criteria for this review ([Veroux 2010](#)). Furthermore, the study looked at mortality, operative time, length of hospital stay and lower leg ischaemia as its secondary outcomes. Other outcomes of interest for this review, such as long-term major complications, all-cause mortality (> 30 days), conversion and re-intervention rates, device-related complications and quality of life, were not studied and, therefore, remain unknown.

### Quality of the evidence

The only study included in this review was judged to be at low risk of bias for all domains except for detection bias where the risk was unclear due to insufficient information to permit a judgement of low or high risk.

For all outcomes, the quality of the evidence was downgraded to low due to the inclusion of only one study with a small sample size and wide confidence intervals. Furthermore, as the study only

included males, the quality of the evidence was downgraded further due to indirectness.

### Potential biases in the review process

None of the authors have any commercial or other conflict of interest. The Cochrane Vascular Information Specialist performed a comprehensive search of the literature; and review authors selected studies in accordance with recommendations provided in the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011). We resolved disagreements by discussion.

### Agreements and disagreements with other studies or reviews

To date, no other systematic review has assessed the efficacy and safety of laparoscopic repair of AAA.

Ahmed 2014 conducted a review to determine how elective laparoscopic abdominal aortic aneurysm (AAA) repair compares to endovascular aneurysm repair (EVAR) in terms of survival. Eight papers (five prospective studies, one retrospective study, one RCT and one systematic review) were deemed to be the best available evidence. The RCT included by Ahmed 2014 was also the only RCT included in our Cochrane Review (Veroux 2010). Ahmed

2014 concluded that laparoscopic AAA repair is just as safe as EVAR, with similar mortality and morbidity rates and length of hospital stay.

## AUTHORS' CONCLUSIONS

### Implications for practice

There is insufficient evidence to draw any conclusions about effectiveness and safety of laparoscopic repair of AAA versus open surgical repair or EVAR, because only one small randomised trial was eligible for inclusion in this review.

### Implications for research

This review highlights the gap in evidence for the use of laparoscopic repair of AAA. Future randomised controlled trials are required.

## ACKNOWLEDGEMENTS

We would like to thank the Cochrane Vascular Information Specialist for assistance with the search strategies.

## REFERENCES

### References to studies included in this review

#### Veroux 2010 *{published data only}*

Veroux P, D'Arrigo G, Veroux M, Giaquinta A, Lomeo A. Sexual dysfunction after elective endovascular or hand-assisted laparoscopic abdominal aneurysm repair. *European Journal of Vascular and Endovascular Surgery* 2010;**40**(1): 71–5.

### References to studies excluded from this review

#### NCT00821145 *{published data only}*

NCT00821145. Laparoscopic versus open abdominal aortic aneurysm (AAA) exclusion (LapAorta). Clinicaltrials.gov 2009. <https://clinicaltrials.gov/ct2/show/NCT00821145> (accessed 10 August 2016)

### Additional references

#### Ahmed 2014

Ahmed N, Gollop ND, Ellis J, Khan OA. How does elective laparoscopic abdominal aortic aneurysm repair compare to endovascular aneurysm repair?. *Interactive Cardiovascular and Thoracic Surgery* 2014;**18**(6):814–20.

#### Anjum 2012

Anjum A, von Allmen R, Greenhalgh R, Powell JT. Explaining the decrease in mortality from abdominal aortic aneurysm rupture. *British Journal of Surgery* 2012;**99**(5): 637–45. [DOI: 10.1002/bjs.8698]

#### Ashton 2002

Ashton HA, Buxton MJ, Day NE, Kim LG, Marteau TM, Scott RA, et al. The Multicentre Aneurysm Screening Study (MASS) into the effect of abdominal aortic aneurysm screening on mortality in men: a randomised controlled trial. *Lancet* 2002;**360**(9345):1531–9.

#### Ashton 2007

Ashton HA, Gao L, Kim LG, Druce PS, Thompson SG, Scott RA. Fifteen-year follow-up of a randomized clinical trial of ultrasonographic screening for abdominal aortic aneurysms. *British Journal of Surgery* 2007;**94**(6):696–701.

#### Atkins 2004

Atkins D, Best D, Briss PA, Eccles M, Falck-Ytter Y, Flottorp S, et al. Grading quality of evidence and strength of recommendations. *BMJ* 2004;**328**(7454):1490–4.

#### Badger 2014

Badger S, Bedenis R, Blair PH, Ellis P, Kee F, Harkin DW. Endovascular treatment for ruptured abdominal aortic

- aneurysm. *Cochrane Database of Systematic Reviews* 2014, Issue 7. [DOI: 10.1002/14651858.CD005261.pub3]
- Ballard 1999**  
Ballard DJ, Filardo G, Fowkes FG, Powell JT. Surgery for small asymptomatic abdominal aortic aneurysms. *Cochrane Database of Systematic Reviews* 1999, Issue 3. [DOI: 10.1002/14651858.CD001835]
- Brewster 2003**  
Brewster DC, Cronenwett JL, Hallett JW Jr, Krupski WC, Matsumura JS, Joint Council of the American Association for Vascular Surgery and Society for Vascular Surgery. Guidelines for the treatment of abdominal aortic aneurysms: Report of a subcommittee of the Joint Council of the American Association for Vascular Surgery and Society for Vascular Surgery. *Journal of Vascular Surgery* 2003;**37**(5): 1106–17.
- Bush 2006**  
Bush RL, Johnson ML, Collins TC, Henderson WG, Khuri SF, Yu HJ. Open versus endovascular abdominal aortic aneurysm repair in VA hospitals. *Journal of the American College of Surgeons* 2006;**202**(4):577–87.
- Ernst 1993**  
Ernst CB. Abdominal aortic aneurysm. *New England Journal of Medicine* 1993;**328**(16):1167–72.
- Ferrari 2006**  
Ferrari M, Adami D, Del Corso A, Berchiolli R, Pietrabissa A, Romagnani F, et al. Laparoscopy-assisted abdominal aortic aneurysm repair: early and middle-term results of a consecutive series of 122 cases. *Journal of Vascular Surgery* 2006;**43**(4):695–700.
- Filardo 2015**  
Filardo G, Powell JT, Martinez MAM, Ballard DJ. Surgery for small asymptomatic abdominal aortic aneurysms. *Cochrane Database of Systematic Reviews* 2015, Issue 2. [DOI: 10.1002/14651858.CD001835.pub4]
- Forsdahl 2009**  
Forsdahl SH, Singh K, Solberg S, Jacobsen BK. Risk factors for abdominal aortic aneurysms: a 7-year prospective study: the Tromsø study, 1994–2001. *Circulation* 2009;**119**(16): 2202–8.
- Higgins 2011**  
Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from [handbook.cochrane.org](http://handbook.cochrane.org).
- Javerliat 2006**  
Javerliat I, Di Centa I, Cerceau P, Alfonsi P, Goëau-Brissonnière O, Coggia M. Totally laparoscopic tube graft bypass for infrarenal aortic aneurysm: a well-established surgical technique. *Acta Chirurgica Belgica* 2006;**106**(2): 261–6.
- Johnston 1991**  
Johnston KW, Rutherford RB, Tilson MD, Shah DM, Hollier L, Stanley JC. Suggested standards for reporting on arterial aneurysms. Subcommittee on Reporting Standards for Arterial Aneurysms, Ad Hoc Committee on Reporting Standards, Society for Vascular Surgery and North American Chapter, International Society for Cardiovascular Surgery. *Journal of Vascular Surgery* 1991;**13**(3):452–8.
- Lederle 1997**  
Lederle FA, Johnson GR, Wilson SE, Chute EP, Littooy FN, Bandyk D, Aneurysm Detection and Management (ADAM) Veterans Affairs Cooperative Study Group. Prevalence and associations of abdominal aortic aneurysm detected through screening. *Annals of Internal Medicine* 1997;**126**(6):441–9.
- Lederle 2000a**  
Lederle FA, Johnson GR, Wilson SE, Chute EP, Hye RJ, Makaroun MS, Aneurysm Detection and Management Veterans Affairs Cooperative Study Investigators. The aneurysm detection and management study screening program: validation cohort and final results. *Archives of Internal Medicine* 2000;**160**(10):1425–30.
- Lederle 2000b**  
Lederle FA, Wilson SE, Johnson GR, Reinke DB, Littooy FN, Acher CW, et al. Immediate repair compared with surveillance of small abdominal aortic aneurysms. *New England Journal of Medicine* 2002;**346**(19):1437–44.
- Lederle 2003**  
Lederle FA, Nelson DB, Joseph AM. Smokers' relative risk for aortic aneurysm compared with other smoking-related diseases: a systematic review. *Journal of Vascular Surgery* 2003;**38**(2):329–34.
- Lindholt 2005**  
Lindholt JS, Juul S, Fasting H, Henneberg EW. Screening for abdominal aortic aneurysms: single centre randomised controlled trial. *BMJ* 2005;**330**(7494):750.
- NICE 2007**  
National Institute for Health and Clinical Excellence (NICE). Laparoscopic repair of abdominal aortic aneurysm. NICE interventional procedure guidance. [www.nice.org.uk/guidance/ipg229](http://www.nice.org.uk/guidance/ipg229) (date accessed 17 June 2015).
- Nordon 2011**  
Nordon IM, Hinchliffe RJ, Loftus IM, Thompson MM. Pathophysiology and epidemiology of abdominal aortic aneurysms. *Nature Reviews Cardiology* 2011;**8**(2):92–102.
- Norman 2004**  
Norman PE, Jamrozik K, Lawrence-Brown MM, Le MT, Spencer CA, Tuohy RJ, et al. Population based randomised controlled trial on impact of screening on mortality from abdominal aortic aneurysm. *BMJ* 2004;**329**(7477):1259.
- Paravastu 2014**  
Paravastu SCV, Jayarajasingam R, Cottam R, Palfreyman SJ, Michaels JA, Thomas SM. Endovascular repair of abdominal aortic aneurysm. *Cochrane Database of Systematic Reviews* 2014, Issue 1. [DOI: 10.1002/14651858.CD004178.pub2]
- Parodi 1991**  
Parodi JC, Palmaz JC, Barone HD. Transfemoral intraluminal graft implantation for abdominal aortic aneurysms. *Annals of Vascular Surgery* 1991;**5**(6):491–9.

**Pleumeekers 1995**

Pleumeekers HJ, Hoes AW, Van der Does E, Van Urk H, Hofman A, de Jong PT. Aneurysms of the abdominal aorta in older adults. The Rotterdam Study. *American Journal of Epidemiology* 1995;**142**(12):1291–9.

**Review Manager 2014 [Computer program]**

The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager 5 (RevMan 5). Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2013.

**Sakalihasan 2005**

Sakalihasan N, Limet R, Defawe OD. Abdominal aortic aneurysm. *Lancet* 2005;**365**(9470):1577–89.

**Scott 2002**

Scott RAP, Bridgewater S, Ashton HA. Randomised clinical trial of screening for abdominal aortic aneurysm in women. *British Journal of Surgery* 2002;**89**(3):283–5.

**Shah 1997**

Shah PK. Inflammation, metalloproteinases, and increased proteolysis: an emerging pathophysiological paradigm in aortic aneurysm. *Circulation* 1997;**96**(7):2115–7.

**Sweeting 2015**

Sweeting MJ, Ulug P, Powell JT, Desgranges P, Balm R, Ruptured Aneurysm Trialists. Ruptured aneurysm trials: the importance of longer-term outcomes and meta-analysis for 1-year mortality. *European Journal of Vascular and Endovascular Surgery* 2015;**50**(3):297–302. [DOI: 10.1016/j.ejvs.2015.04.015]

**TUKSAT 1998**

The UK Small Aneurysm Trial Participants. Mortality results for randomised controlled trial of early elective surgery or ultrasonographic surveillance for small abdominal aortic aneurysms. *Lancet* 1998;**352**(9141):1649–55.

**Vardulaki 1999**

Vardulaki KA, Prevost TC, Walker NM, Day NE, Wilink AB, Quick CR, et al. Incidence among men of asymptomatic aortic abdominal aneurysms: estimates from 500 screen detected cases. *Journal of Medical Screening* 1996;**6**(1):50–4.

**Veith 2003**

Veith FJ, Ohki T, Lipsitz EC, Suggs WD, Cynamon J. Treatment of ruptured abdominal aneurysms with stent grafts: a new gold standard?. *Seminars in Vascular Surgery* 2003;**16**(2):171–5.

**Wanhainen 2008**

Wanhainen A. How to define an abdominal aortic aneurysm - influence on epidemiology and clinical practice. *Scandinavian Journal of Surgery* 2008;**97**(2):105–9.

**References to other published versions of this review****Robertson 2016**

Robertson L, Nandhra S. Laparoscopic surgery for abdominal aortic aneurysm. *Cochrane Database of Systematic Reviews* 2016, Issue 7. [DOI: 10.1002/14651858.CD012302]

\* Indicates the major publication for the study

## CHARACTERISTICS OF STUDIES

### Characteristics of included studies [ordered by study ID]

#### Veroux 2010

Methods	Study design: single centre prospective randomised trial Length of follow-up: 2 years Dates of study: May 2006 to May 2008 Location: Italy Setting: hospital	
Participants	Number: 100: HALS 50, EVAR 50. Age, mean years: HALS 61.2 years, EVAR 69.6 years. Sex: male. Ethnic group: not stated. Inclusion criteria: men undergoing elective repair of AAA who did not have preoperative sexual dysfunction according to International Index of Erectile Function (IIEF) Exclusion criteria: emergency repair of ruptured aneurysm, juxtarenal aneurysm repair, previous prostate surgery and preoperative sexual dysfunction (class I and II) according to International Index of Erectile Function (IIEF) Diameter of AAA: HALS 5.9 ± 1.8 cm, EVAR 5.7 ± 2.1 cm. Diagnosis of AAA: not stated. Co-morbidities: chronic obstructive pulmonary disease: HALS 17, EVAR 24; coronary artery disease HALS 9, EVAR 8; acute myocardial infarction HALS 6, EVAR 7 Low/high risk patients: not stated.	
Interventions	Intervention 1: hand-assisted laparoscopic surgery (HALS). Intervention 2: endovascular aneurysm repair (EVAR).	
Outcomes	Primary: incidence of sexual dysfunction and retrograde ejaculation Secondary: perioperative mortality, leg ischaemia, duration of surgery (minutes) and length of hospital stay (days)	
Notes		
<b>Risk of bias</b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	Quote: "Randomisation codes were generated using a validated computer method, placed in sealed envelopes, each labelled with a patient number, and opened at patient admission for AAA repair"
Allocation concealment (selection bias)	Low risk	Quote: "Randomisation codes were placed in sealed envelopes, each labelled with a patient number, and opened at patient admission for AAA repair"

**Veroux 2010** (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Low risk	No blinding, but review authors judged that outcomes and outcome measurements are not likely to be influenced by lack of blinding
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to permit judgement of low or high risk of detection bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No missing outcome data.
Selective reporting (reporting bias)	Low risk	The study protocol is not available but it is clear that the published report includes all expected outcomes, including those that were pre-specified
Other bias	Low risk	Study appears to be free from other sources of bias.

AAA: abdominal aortic aneurysm

cm: centimetres

EVAR: endovascular assisted repair

HALS: hand-assisted laparoscopic repair

**Characteristics of excluded studies** [ordered by study ID]

Study	Reason for exclusion
<a href="#">NCT00821145</a>	Study was withdrawn prior to enrolment

## DATA AND ANALYSES

### Comparison 1. Hand-assisted laparoscopic repair vs EVAR

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 In-hospital mortality	1	100	Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2 Operative time (minutes)	1	100	Mean Difference (IV, Fixed, 95% CI)	53.0 [36.49, 69.51]
3 Major complications	1	100	Risk Ratio (M-H, Fixed, 95% CI)	0.5 [0.05, 5.34]
3.1 Lower limb ischaemia	1	100	Risk Ratio (M-H, Fixed, 95% CI)	0.5 [0.05, 5.34]

#### Analysis 1.1. Comparison 1 Hand-assisted laparoscopic repair vs EVAR, Outcome 1 In-hospital mortality.

Review: Laparoscopic surgery for elective abdominal aortic aneurysm repair

Comparison: 1 Hand-assisted laparoscopic repair vs EVAR

Outcome: 1 In-hospital mortality


Study or subgroup	HALS n/N	EVAR n/N	Risk Ratio M-H,Fixed,95% CI	Weight	Risk Ratio M-H,Fixed,95% CI
Veroux 2010	0/50	0/50			Not estimable
<b>Total (95% CI)</b>	<b>50</b>	<b>50</b>			<b>Not estimable</b>
Total events: 0 (HALS), 0 (EVAR)					
Heterogeneity: not applicable					
Test for overall effect: not applicable					
Test for subgroup differences: Not applicable					
			0.01 0.1	10 100	
			Favours HALS	Favours EVAR	

### Analysis 1.2. Comparison 1 Hand-assisted laparoscopic repair vs EVAR, Outcome 2 Operative time (minutes).

Review: Laparoscopic surgery for elective abdominal aortic aneurysm repair

Comparison: 1 Hand-assisted laparoscopic repair vs EVAR

Outcome: 2 Operative time (minutes)




### Analysis 1.3. Comparison 1 Hand-assisted laparoscopic repair vs EVAR, Outcome 3 Major complications.

Review: Laparoscopic surgery for elective abdominal aortic aneurysm repair

Comparison: 1 Hand-assisted laparoscopic repair vs EVAR

Outcome: 3 Major complications



## APPENDICES

### Appendix I. CENTRAL search strategy

#1	MESH DESCRIPTOR Aortic Aneurysm	100
#2	MESH DESCRIPTOR Aortic Aneurysm, Abdominal	417
#3	MESH DESCRIPTOR Aortic Aneurysm, Thoracic	52
#4	aort*:TI,AB,KY	6589
#5	(juxta renal):TI,AB,KY	0
#6	juxtarenal:TI,AB,KY	3
#7	(juxta renal or juxtarenal ):TI,AB,KY	3
#8	(pararenal or para renal ):TI,AB,KY	4
#9	(suprarenal or supra renal):TI,AB,KY	23
#10	(short neck* or shortneck*):TI,AB,KY	9
#11	(visceral aortic segment):TI,AB,KY	0
#12	thorac*:TI,AB,KY	8787
#13	abdominal:TI,AB,KY	19758
#14	#4 OR #5 OR #6 OR #7 OR #8 OR #9 OR #10 OR #11 OR #12 OR #13	32813
#15	aneur?sm*:TI,AB,KY	2525
#16	#14 AND #15	1038
#17	(JRAAA or JRAAAs or PAAA or PAAAs or TAAA or TAAAs or JPAA or JPAAAs or SRA or SRAs or SRAA or SRAAs):TI, AB,KY	69
#18	((aort* near3 (ballon* or dilat* or bulg* or expan*)):TI,AB, KY	77
#19	#1 OR #2 OR #3 OR #16 OR #17 OR #18	1142
#20	MESH DESCRIPTOR Laparoscopes EXPLODE ALL TREES	99

(Continued)

#21	MESH DESCRIPTOR Laparoscopy EXPLODE ALL TREES	4149
#22	laparosc*:TI,AB,KY	8960
#23	#20 OR #21 OR #22	8960
#24	#19 AND #23	10

## Appendix 2. Trial registries searches

ClinicalTrials.gov

4 studies found for: laparoscopic AND aneurysm

World Health Organization International Clinical Trials Registry Platform

3 studies found for: laparoscopic AND aneurysm

ISRCTN Register

2 studies found for: laparoscopic AND aneurysm

## CONTRIBUTIONS OF AUTHORS

LR: drafted the protocol, selected studies for inclusion and wrote the review.

SN: drafted the protocol, selected studies for inclusion and wrote the review.

## DECLARATIONS OF INTEREST

LR: none known.

SN: none known.

## SOURCES OF SUPPORT

### Internal sources

- No sources of support supplied

## External sources

- National Institute for Health Research (NIHR), UK.

This project was supported by the NIHR, via Cochrane Programme Grant funding (13/89/23) to the Cochrane Vascular Group. The views and opinions expressed therein are those of the authors and do not necessarily reflect those of the Systematic Reviews Programme, NIHR, NHS, or the Department of Health.

- Chief Scientist Office, Scottish Government Health Directorates, The Scottish Government, UK.

The Cochrane Vascular editorial base is supported by the Chief Scientist Office.

## DIFFERENCES BETWEEN PROTOCOL AND REVIEW

We have amended the title to reflect the objective of the review more accurately. For completeness and clarity we have also provided further details regarding the primary and secondary outcomes.

## INDEX TERMS

### Medical Subject Headings (MeSH)

Aortic Aneurysm, Abdominal [\*surgery]; Elective Surgical Procedures [adverse effects; \*methods]; Endovascular Procedures [adverse effects; \*methods]; Laparoscopy [adverse effects; \*methods]; Length of Stay; Operative Time; Randomized Controlled Trials as Topic

### MeSH check words

Humans; Male