

The relationship between regional pain with or without neuropathic symptoms and chronic widespread pain

John McBeth^{a,*}, Matthew R. Mulvey^b, Amir Rashid^a, James Anderson^a, Katie Druce^a

Abstract

This study was performed to test whether the risk of developing chronic widespread pain (CWP) in those with regional pain was augmented in those with symptoms of neuropathic pain (NP). Persons free of CWP completed the Douleur Neuropathique 4 (scores ≥ 3 indicating NP); demographics; Hospital Anxiety and Depression scale; Pittsburgh Sleep Quality Index; and pain medications. Participants were classified as having no pain, regional pain with no symptoms of NP ($\bar{N}P$), or regional pain with symptoms of NP (NP). At the 12-month follow-up, participants with CWP were identified. Logistic regression estimated the odds ratio, with 95% confidence intervals, of CWP in the NP and NP groups compared with no pain, and NP compared with $\bar{N}P$. Partial population attributable risks estimated the proportion of CWP attributable to baseline $\bar{N}P$ or NP exposure. One thousand one hundred sixty-two participants completed the baseline DN4 and provided pain data at follow-up: 523 (45.0%) had no baseline pain, 562 (48.4%) $\bar{N}P$, and 77 (6.6%) NP. One hundred fifty-three (13.2%) had CWP at 12 months: 19 (3.6%) no pain, 108 (19.2%) $\bar{N}P$, and 26 (33.8%) NP. $\bar{N}P$ (2.9 [1.9-4.3]) and NP (2.1 [1.1-4.0]) predicted CWP after adjusting for demographics, Hospital Anxiety and Depression scale, Pittsburgh Sleep Quality Index, and medications. The partial population attributable risk was 41.3% (25.2-54.0) for NP and 6.0% (0.1-11.6) for NP. The NP group were not more likely to develop CWP when compared directly with $\bar{N}P$ (1.5 [0.8-2.8]). Neuropathic pain was relatively rare and predicted a small number of new-onset CWP cases. Using these estimates, treatments targeting NP would at best prevent 6% of CWP cases.

Keywords: Chronic widespread pain, Fibromyalgia, Neuropathic pain, Cohort study, Prospective, Population risk

1. Introduction

Chronic widespread pain (CWP), the clinical hallmark of fibromyalgia,⁴⁷ is a significant health burden: it affects 11% of the general population,³¹ is associated with higher disability,⁴⁴ and increases mortality³⁰ most likely through reduced physical function.⁴⁰ Of the estimated 33% to 50%^{7,44} of the general population who have a painful regional musculoskeletal disorder (eg, back pain and knee pain), up to 20% develop CWP.^{20,33,34} Risk factors for CWP include sleep problems,¹ childhood chronic physical illness,³⁷ and trauma.²³ In a systematic review, the transition from regional musculoskeletal pain to CWP was predicted by pain independently of sex, increasing age, family history of pain, and depression.²⁸

Neuropathic pain (NP), pain distinguished from other pain conditions by characteristic symptoms such as “burning” or “freezing,”^{11,14} is present in up to 16% of persons with CWP,³⁵ double the population prevalence of 8%.¹⁵ The heritability of NP and CWP is correlated and the genetic predisposition to NP shared with CWP.³⁵ There are no prospective studies testing whether the risk of developing CWP in those with regional pain is augmented by the presence of NP.

This study tested the hypothesis that the risk of developing CWP would be augmented by the presence of pain with symptoms of NP.

2. Methods

In a prospective population-based cohort study (the Pain across the Adult Lifespan study⁸), participants were invited to complete a baseline questionnaire. Those free of CWP were identified and classified as having no pain, pain without neuropathic symptoms ($\bar{N}P$), or NP. All participants were followed up 12 months later, at which time they completed a second questionnaire and those with CWP identified. The relationship between pain status at baseline and CWP at follow-up was examined.

Ethical approval was granted by the North West 8 Research Ethics Committee—Greater Manchester East (reference number 10/H1013/29).

2.1. Baseline questionnaire

The baseline questionnaire collected data on the presence, location, and duration of pain. Participant’s reports of pain were used to identify those with prevalent CWP, classified using published criteria¹³ that have been used in previous

Sponsorships or competing interests that may be relevant to content are disclosed at the end of this article.

^a Versus Arthritis Centre for Epidemiology and the NIHR Manchester Musculoskeletal Biomedical Research Unit, Central Manchester University Hospitals NHS Foundation Trust; The University of Manchester, Manchester, United Kingdom, ^b Academic Unit of Palliative Care, Leeds Institute of Health Research, University of Leeds, Leeds, United Kingdom

*Corresponding author. Versus Arthritis Centre for Epidemiology, NIHR Manchester Musculoskeletal Biomedical Research Unit, Central Manchester University Hospitals NHS Foundation Trust, The University of Manchester, Manchester, M13 9PT, United Kingdom. Address: Tel.: +44 (0)161 275 5788. E-mail address: john.mc-beth@manchester.ac.uk (J. McBeth).

PAIN 160 (2019) 1817–1823

Copyright © 2019 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the International Association for the Study of Pain. This is an open access article distributed under the Creative Commons Attribution License 4.0 (CCBY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

<http://dx.doi.org/10.1097/j.pain.0000000000001568>

population-based surveys²⁴ (see below for details). The screening questionnaire also collected data on demographics (date of birth; sex; English Index of Multiple Deprivation 2007,³⁸ with higher scores indicating less deprivation; Hospital Anxiety and Depression (HAD)⁴⁸ depression and anxiety subscales (7 items each, scored 0–3, total score 0–21, higher scores represent higher probability of depression or anxiety); Pittsburgh Sleep Quality Index¹⁰ (19 items, scored 0–3, total score range 0–57); Rapid Assessment of Physical Activity⁴² (7 items, scored as sedentary, underactive, underactive regular, and active); self-reported pain medications (summed to give a total count); and signed consent.

2.2. Presence, distribution, and duration of musculoskeletal pain

The questionnaire included a detailed assessment of the presence of musculoskeletal pain. All participants were asked “During the past month, have you experienced any pain which has lasted at least 1 day or longer?” Respondents answering positively were asked to shade the location(s) of their pain on a four-view body manikin. Participants were then asked whether they had been aware of their pain for 3 months or longer (yes/no). Using their responses to these questions, participants were classified as having CWP using the definition in the American College of Rheumatology (ACR) 1990 classification criteria for fibromyalgia⁴⁷ that was operationalised by Croft et al.¹³ as requiring axial (cervical spine, thoracic spine, anterior chest, or low back) pain and pain in contralateral body quadrants, above and below the waist, and on the right and left hand sides of the body, which has been present for at least 3 months. Remaining participants were classified as having some pain (those participants reporting pain that did not satisfy the criteria for CWP) or no pain. The data collection tool (a blank body manikin) has been shown to collect valid and reliable pain data with high between-rater reliability scoring.²⁷ The original criteria for CWP defined by Croft et al.¹³ were based on the definition included in the 1990 classification criteria for fibromyalgia that required pain in the axial skeleton and in contralateral body quadrants. The criteria have been used in multiple population-based epidemiological studies and have face and construct validity (they identify persons with high care utilisation, poor sleep, depression, physical inactivity, and with higher tender point counts among many other findings)^{9,26} and have been shown to identify persons with a genetic susceptibility to developing CWP.^{29,39}

2.3. Neuropathic pain

Participants reporting the presence of pain were invited to report the sensory characteristics of their pain symptoms using the Douleur Neuropathique 4 (DN4) questionnaire.⁵ Participants were asked “Thinking about your single worst pain, does the pain have one or more of the following characteristics?”; burning, painful cold, electric shocks, tingling, pins and needles, numbness, and itching. For each characteristic responses were scored yes (= 1) or no (= 0). Participants scoring 3 or more were considered to have pain of predominantly neuropathic origin. The short version of the DN4 has been validated for use in self-reported questionnaires with 83% sensitivity and 90% specificity when compared to clinical diagnosis.⁴ The interrater reliability of the DN4 was demonstrated in the original French version with kappa values for individual items ranging from 0.66 to 0.96⁵ and has subsequently been demonstrated cross-culturally, although the level of supporting evidence is low.³²

2.4. Follow-up questionnaire

A follow-up questionnaire was mailed to those baseline participants who were free of CWP, 12 months after the date they returned a baseline questionnaire. The follow-up questionnaire included identical methods to assess pain as those used in the baseline survey. Based on their pain reports at follow-up, participants were classified as “CWP” for those who reported pain that satisfied the criteria for CWP or “Not CWP.”

2.5. Analysis

Logistic regression estimated the odds of CWP first in the \overline{NP} and NP groups compared with the no pain group (the referent group), second in the NP group compared with the \overline{NP} group (the referent group), and finally for the individual pain characteristics with those participants not reporting the characteristic being classified as the referent group. Results were expressed as odds ratios (ORs) with 95% confidence intervals (CIs). For those comparisons with significant ORs, population attributable risks (PARs) were calculated using formulae for cohort studies.^{21,41} These estimated the % of CWP that would be avoided if participants were not exposed to \overline{NP} or NP. In an evaluation of a preventive intervention in a multifactorial disease setting, the interest is in the percent of cases associated with the exposures to be modified, when other risk factors, possibly nonmodifiable, exist but do not change as a result of the intervention. The partial population attributable risk (PARp) was proposed to estimate this quantity. Under the assumption of no interaction of the index exposure effects with the background risk factors, the PARp is formulated as

$$\begin{aligned} \text{PARp} &= \frac{\sum_{s=1}^S \sum_{t=1}^T p_{st} R_{1s} RR_{2t} - \sum_{s=1}^S \sum_{t=1}^T p_{st} RR_{2t}}{\sum_{s=1}^S \sum_{t=1}^T p_{st} RR_{1s} RR_{2t}} \\ &= \frac{\sum_{t=1}^T p_{st} RR_{2t}}{\sum_{s=1}^S \sum_{t=1}^T p_{st} RR_{2t}} \end{aligned}$$

where t denotes a stratum of unique combinations of levels of all background risk factors which are not under study, $t = 1; \dots; T$ and RR_{2t} is the relative risk in combination t relative to the lowest risk level, where $RR_{2,1} = 1$. S indicates an index exposure group defined by each of the unique combinations of the levels of the index risk factors, that is, those risk factors to which the PARp applies, $s = 1; \dots; S$, and RR_{1s} is the relative risk corresponding to combinations relative to the lowest risk combination, $RR_{1,1} = 1$. The joint prevalence of exposure group s and stratum t is denoted by p_{st} , and $p_{\cdot t} = \sum_{s=1}^S p_{st}$.⁴¹

3. Results

3.1. Participation rates

Of 2654 persons who were mailed a baseline questionnaire, 1817 responded (68.5%, adjusted for 312 [11.8%] participants who had moved from their recorded address and 41 [1.5%] who had died) (**Fig. 1**). One thousand three hundred seventy-seven (75.8%) participants were free of CWP and were mailed a follow-up questionnaire, and 1174 (85.3%, adjusted for those who had moved or died) responded.

3.2. Rate of chronic widespread pain and associated factors

Of the 1174 participants at follow-up, 12 (1.0%) were missing pain data, leaving 1162 participants for analysis. Of the 1162




Figure 1. Flow diagram of participant recruitment and CWP rates. Primary care records were reviewed to identify those who were recorded as either having moved address or had died after the questionnaire mailing. CWP-free = CWP-free at baseline and follow-up; CWP = CWP-free at baseline and CWP at follow-up; †Proportions are of participants who sent baseline questionnaire. *Proportions are of participants who returned a baseline questionnaire. °Proportions are of participants free of CWP at baseline. °Proportions are of participants free of CWP at baseline and mailed a follow-up questionnaire. CWP, chronic widespread pain.

participants, 523 (45.0%) had no pain, 562 (48.4%) had $\overline{\text{NP}}$, and 77 (6.6%) had NP at baseline (**Table 1**). The proportion of participants classified as CWP at follow-up was significantly higher among those with $\overline{\text{NP}}$ (19.2%) and those with NP (33.8%), when compared with the proportion in those reporting no pain at baseline (3.6%). Baseline reports of NP characteristics assessed in the DN4 questionnaire were more common in those with CWP at follow-up ($P < 0.001$ for all comparisons). Chronic widespread pain was associated with having significantly more pain sites at baseline, higher pain medication use at baseline, female sex, and higher HAD anxiety, HAD depression, and sleep problem scores at baseline. There was no association with age, occupational status, or deprivation score (**Table 1**).

3.3. The relationship between neuropathic pain and chronic widespread pain

When compared to those with no pain at baseline, after adjusting for age and sex (**Table 2**), $\overline{\text{NP}}$ was associated with a 3-fold (OR = 2.9, 95% CI [2.0-4.2]) increased odds of CWP

at follow-up, and NP was associated with a 4-fold (3.9 [2.3-6.4]) increased odds. After adjusting for age, sex, psychosocial variables, demographics, and medication use, the association between $\overline{\text{NP}}$ and CWP persisted (2.9 [1.9, 4.3]), whereas the association between NP and CWP was attenuated (2.1 [1.1-4.0]). After adjusting for all covariates, the NP characteristics of burning, electric shocks, tingling, pins and needles, and numbness were all significantly associated with CWP.

After adjusting for age and sex, those with NP were twice (2.2 [1.3-3.6]) as likely to have CWP at follow-up when compared with the $\overline{\text{NP}}$ group (**Table 2**). However, the relationship between NP and CWP was attenuated and no longer significant (1.5 [0.8-2.8]) when adjusted for age, sex, psychosocial variables, demographics, and medication use.

3.4. Partial population attributable risks

The proportion of all cases of CWP that could be attributed to the presence of NP at baseline was 6.0% (95% CI [0.1-11.6]),

Table 1
Participant characteristics at baseline stratified by chronic widespread pain status at follow-up.

	Follow-up pain status		P†
	Not CWP; N = 1009 (86.8%)	CWP; N = 153 (13.2%)	
Neuropathic pain, no. of pain sites and pain medication use			
Presence of pain at baseline stratified by DN4 score*			
No pain	504 (49.9)	19 (12.4)	0.00
NP (regional pain, DN4 score 0-2)	454 (45.0)	108 (70.6)	
NP (regional pain, DN4 score 3-7)	51 (5.1)	26 (17.0)	
DN4 pain characteristics (n (%) reporting yes)*			
Burning	125 (12.4)	48 (31.4)	0.00
Painful cold	46 (4.6)	16 (10.5)	0.00
Electric shocks	75 (7.4)	33 (21.6)	0.00
Tingling	70 (6.9)	31 (20.3)	0.00
Pins and needles	49 (4.9)	23 (15.0)	0.00
Itching	23 (2.3)	8 (5.2)	0.04
Numbness	83 (8.2)	30 (19.6)	0.00
No. of pain sites	1 (0-3)	3 (2-5)	0.00
No. of current pain medications	1 (0-3)	2 (0-5)	0.00
Demographics			
Age	61 (53-67)	62 (54-67)	0.43
Sex*			
Women	564 (55.8)	99 (64.7)	0.04
Men	446 (44.2)	54 (35.3)	
Occupational status*			
Working full time	385 (38.2)	47 (31.1)	0.11
Working part time	156 (15.5)	26 (17.2)	
Unemployed due to ill health	18 (1.8)	7 (4.6)	
Other (including unemployed, n = 20)	449 (44.4)	71 (47.0)	
Deprivation quantiles*			
1	202 (20.0)	45 (29.4)	0.05
2	185 (18.3)	22 (14.4)	
3	320 (31.7)	41 (26.8)	
4	302 (29.9)	45 (29.4)	
Psychosocial variables			
HAD depression	2 (1-5)	4 (2-7)	0.00
HAD anxiety	4 (2-7)	6 (3-9)	0.00
Sleep problems	5 (2-9)	8 (4-13)	0.00

All values are median (interquartile range) unless indicated by * which are n (%) with column values totalling 100% (except where rounding equals <100%).

† P-values derived from χ^2 or Kruskal–Wallis tests as appropriate.

CWP, chronic widespread pain; DN4, Douleur Neuropathique 4; HAD, Hospital Anxiety and Depression scale; NP, neuropathic pain.

whereas 41.3% (25.2, 54.0) could be attributed to the presence of \overline{NP} (Table 2). The proportion of CWP attributable to the NP characteristics ranged from 1.9% to 5.0% (Table 2).

4. Discussion

4.1. Summary of findings

This study tested the hypothesis that the relationship between regional pain and CWP would be augmented by the presence of symptoms of NP. The data presented here do not support that hypothesis. When compared to those persons with no pain, having some pain was associated with an increased odds of developing CWP. The relationship with CWP initially seemed to be stronger among those persons with symptoms of NP, although the effect was attenuated when adjusted for age, sex, occupational status, deprivation, anxiety, depression, and sleep problems to the extent that \overline{NP} had a stronger relationship with CWP. The data suggest that, at a population level, the proportion of all new cases of CWP that could be attributed to

NP was modest, and was lower than the proportion attributable to \overline{NP} .

4.2. Setting these results in the context of current literature

The data presented here show that having regional pain increases the odds of CWP by 2-fold, irrespective of the presence of NP symptoms. Previous longitudinal studies have shown similar strengths of associations with arm pain (relative risk [RR] 2.4 [1.5-4.3]) and back pain (RR 2.5 [1.5-6.7])¹⁷ and pain at multiple sites (OR 2.6 [1.3-5.4])²² with CWP. In contrast to the findings presented here, some previous cross-sectional studies have supported the role of NP in CWP and associated disorders. Fishbain et al.¹⁶ reported that chronic pain patients including those with fibromyalgia scored above the cutoff on the Neuropathic Pain Scale. Fibromyalgia patients commonly report sensory symptoms characteristic of NP with 83% reporting deep-tissue hyperalgesia, 51% reporting burning, and 47% reporting prickling on the painDETECT questionnaire.^{2,18} There is suggestive evidence for an overlap between NP and CWP or

Table 2

Regional pain stratified by DN4 score and DN4 pain characteristics and CWP: results from multinomial logistic regression models.

	Adjustments				
	Age and sex	Age, sex, and psychosocial variables*	Age, sex, psychosocial variables, and demographics†	Age, sex, psychosocial variables, demographics, and medication use‡	
	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	PARp, % (95% CI)
Presence of pain at baseline stratified by DN4 score					
No pain	Referent	Referent	Referent	Referent	Referent
NP (DN4 score 0-2)	2.9 (2.0-4.2)	2.8 (1.9-4.1)	2.8 (1.9-4.2)	2.9 (1.9-4.3)	41.3 (25.2-54.0)
NP (DN4 score 3-7)	3.9 (2.3-6.4)	2.5 (1.4-4.4)	2.3 (1.2-4.2)	2.1 (1.1-4.0)	6.0 (0.1-11.6)
Presence of regional pain at baseline stratified by DN4 score					
NP (DN4 score 0-2)	Referent	Referent	Referent	Referent	Not estimated
NP (DN4 score 3-7)	2.2 (1.3-3.6)	1.7 (0.96-2.9)	1.6 (0.9-2.9)	1.5 (0.8-2.8)	
DN4 pain characteristics§					
Burning	3.2 (2.1-4.7)	2.7 (1.8-4.1)	2.7 (1.7-4.2)	2.6 (1.7-4.0)	5.0 (1.1-8.8)
Painful cold	2.5 (1.4-4.6)	1.9 (0.99-3.5)	1.9 (0.97-3.8)	1.8 (0.9-3.5)	1.1 (0.7-2.9)
Electric shocks	3.4 (2.2-5.4)	2.9 (1.8-4.6)	2.6 (1.5-4.4)	2.5 (1.4-4.2)	2.9 (0.0-5.7)
Tingling	3.4 (2.1-5.4)	2.3 (1.4-3.9)	2.3 (1.3-3.9)	2.2 (1.3-3.7)	2.8 (0.0-5.5)
Pins and needles	3.5 (2.1-5.9)	2.3 (1.3-4.1)	2.1 (1.1-3.9)	2.0 (1.1-3.7)	1.6 (0.1-3.8)
Itching	2.3 (1.0-5.3)	1.4 (0.6-3.4)	1.2 (0.5-3.2)	1.0 (0.4-2.8)	0.0 (0.0-1.2)
Numbness	2.8 (1.7-4.4)	2.0 (1.2-3.2)	1.8 (1.1-3.1)	1.8 (1.1-3.1)	1.9 (0.1-4.3)

* Adjustments were age, sex, and baseline measures of depression and sleep problems.

† Adjustments were as for previous model plus baseline measures of occupational status and deprivation.

‡ Adjustments were as for previous model plus baseline measures of number of current pain medications.

§ Referent group are participants not reporting the DN4 pain characteristic.

CI, confidence interval; CWP, chronic widespread pain; DN4, Douleur Neuropathique 4; NP, neuropathic pain; OR, odds ratio; PARp, partial population attributable risk estimated from fully adjusted model.

fibromyalgia; these disorders share common risk factors including age, sex, body mass index, smoking, and socioeconomic status.³⁵ However, these associations are not consistent: patients with painful diabetic neuropathy when compared to those with fibromyalgia were less likely to be female, had a higher BMI, were less likely to be depressed or anxious, and had better sleep patterns.²⁵

Fibromyalgia and CWP are clearly related, with fibromyalgia being a subset of persons with CWP.⁴⁶ However, fibromyalgia is not CWP, and one of the main differences may be the presence of NP. In a study of twins, 15% of those with CWP had features of NP, the heritability of NP and CWP were correlated, and the genetic predisposition to NP was shared with that of CWP.³⁵ These associations may be driven by the subset of twins who had fibromyalgia, and this is supported by the findings of similar results when the data set was restricted to the subgroup of twins with fibromyalgia.³⁵ The data presented here support the role of regional pain in the onset of CWP and clearly show that the relationship is independent of the presence of NP symptoms. It is possible that some participants with asymptomatic (ie, not painful) small fibre neuropathy could be more likely to develop CWP; however, we did not collect that data and are unable to address that question in the current study.

4.3. Methodological strengths and limitations

This study has several strengths. The cohort was a large, unselected group of persons free of CWP at baseline. The study used validated assessments of pain, NP, and all covariates. Previous studies that were cross-sectional were unable to determine the temporal association between the

presence of neuropathic features of acute localised pain and the transition to CWP. Here, participants who were free of CWP were identified and followed up over time to identify those who developed CWP, allowing the temporal relationship between having NP and developing CWP to be established. Previous studies have examined clinic populations where levels of psychological distress are higher than unselected general population samples, and high levels of psychological and emotional distress are known to be related to levels of pain reporting, somatosensory changes, and altered nociceptive processing.⁴⁵ Limitations were that participants were followed up once, 12 months after they completed their baseline questionnaire, and it was not clear what happened in the intervening months between baseline and follow-up. Although those reporting CWP at follow-up were identified and classified as the CWP group, it is likely that some participants may have developed CWP that then resolved before follow-up. We would have missed those incident cases, underestimated the rate of CWP in this cohort, and misclassified participants as Not CWP instead of CWP. How would that impact on the associations reported here? If the misclassification was random across baseline exposure status (ie, was independent of baseline pain status) it could act to attenuate the observed association and we may have underestimated the association between NP and the onset of CWP. Any underestimation of effect is unlikely to impact significantly on the pPAR estimates here because, although the PAR is based on the effect estimate, it is heavily influenced by how common a risk factor is in the population. Among those free of CWP, baseline NP was much more common, with 48.4% of participants having had NP and 6.6% having NP.

Several screening tools have been developed to assess NP characteristics. Previous studies using the DN4 and LANSS have estimated the prevalence of pain of predominately neuropathic origin to be between 7% and 8% of the general population.^{6,43} However, the use of NP screening tools in self-reported surveys of musculoskeletal pain is contentious because not all pains are the same.³⁶ NP screening tools have been primarily validated for use in subjects with single-site pain disorders (such as low back pain, diabetic neuropathy, or trigeminal neuralgia) and require the respondent to focus on a single painful region or area.^{3,5,18} Although the participants in this study did not have CWP at baseline, CWP by its nature is multisite and widespread, and the suitability of these screening tools for use in studies of multisite pain conditions is not clear.³⁶ A recent epidemiological survey of NP symptoms in the general population navigated this issue by asking participants to indicate the site(s) of any chronic pain on a body map and then to identify the most troublesome site of pain from a list of body regions, and subsequent questions on neuropathic symptoms were focused on this “most troublesome pain.”⁴³ That is, the method we used here, and although this method is not without its shortcomings, it does allow for the assessment of NP symptoms in population-based surveys of musculoskeletal pain which would otherwise not be possible.

4.4. Clinical implications

The effectiveness of pharmacological treatments for NP in patients with CWP disorders such as fibromyalgia are not clear with systematic reviews showing the supporting evidence for gabapentin is weak,¹² and there are no high-quality trials of oxycodone.¹⁹ In this study, NP was relatively rare and predicted a small number of new-onset CWP cases. Using these estimates, treatments targeting NP would at best prevent 6% of CWP cases. Chronic widespread pain is highly prevalent in the general population, and effective treatment of pain not of NP origin will have the biggest impact on population levels of CWP.

Conflict of interest statement

The authors have no conflict of interest to declare.

Acknowledgements

The authors express their gratitude to all study participants. This study was funded by Arthritis Research United Kingdom (Arthritis Research UK Centre for Epidemiology Centre Grant: 21755). J. McBeth had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Article history:

Received 24 October 2018

Received in revised form 11 March 2019

Accepted 15 March 2019

Available online 8 April 2019

References

- [1] Aili K, Andersson M, Bremander A, Haglund E, Larsson I, Bergman S. Sleep problems and fatigue as predictors for the onset of chronic widespread pain over a 5- and 18-year perspective. *BMC Musculoskelet Disord* 2018;19:390.
- [2] Amris K, Jespersen A, Bliddal H. Self-reported somatosensory symptoms of neuropathic pain in fibromyalgia and chronic widespread pain correlate with tender point count and pressure-pain thresholds. *PAIN* 2010;151:664–9.
- [3] Bennett M. The LANSS Pain Scale: the Leeds assessment of neuropathic symptoms and signs. *PAIN* 2001;92:147–57.
- [4] Bouhassira D, Attal N. All in one: is it possible to assess all dimensions of any pain with a simple questionnaire? *PAIN* 2009;144:7–8.
- [5] Bouhassira D, Attal N, Alchaar H, Boureau F, Brochet B, Bruxelle J, Cunin G, Fermanian J, Ginies P, Grun-Overdyking A, Jafari-Schluep H, Lanteri-Minet M, Laurent B, Mick G, Serrie A, Valade D, Vicaud E. Comparison of pain syndromes associated with nervous or somatic lesions and development of a new neuropathic pain diagnostic questionnaire (DN4). *PAIN* 2005;114:29–36.
- [6] Bouhassira D, Lanteri-Minet M, Attal N, Laurent B, Touboul C. Prevalence of chronic pain with neuropathic characteristics in the general population. *PAIN* 2008;136:380–7.
- [7] Brennan-Olsen SL, Cook S, Leech MT, Bowe SJ, Kowal P, Naidoo N, Ackerman IN, Page RS, Hosking SM, Pasco JA, Mohebbi M. Prevalence of arthritis according to age, sex and socioeconomic status in six low and middle income countries: analysis of data from the World Health Organization study on global AGEing and adult health (SAGE) Wave 1. *BMC Musculoskelet Disord* 2017;18:271.
- [8] Brown D, Mulvey M, Cordingley L, Rashid A, Horan M, Pendleton N, Duncan R, McBeth J. The relationship between psychological distress and multiple tender points across the adult lifespan. *Arch Gerontol Geriatr* 2016;63:102–7.
- [9] Burri A, Ogata S, Livshits G, Williams F. The association between chronic widespread musculoskeletal pain, depression and fatigue is genetically mediated. *PLoS One* 2015;10:e0140289.
- [10] Buysse DJ, Reynolds CF III, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res* 1989;28:193–213.
- [11] Caro XJ, Winter EF. The role and importance of small fiber neuropathy in fibromyalgia pain. *Curr Pain Headache Rep* 2015;19:55.
- [12] Cooper TE, Derry S, Wiffen PJ, Moore RA. Gabapentin for fibromyalgia pain in adults. *Cochrane Database Syst Rev* 2017;CD012188.
- [13] Croft P, Schollum J, Silman A. Population study of tender point counts and pain as evidence of fibromyalgia. *BMJ* 1994;309:696–9.
- [14] Dworkin RH, Fields HL. Fibromyalgia from the perspective of neuropathic pain. *J Rheumatol Suppl* 2005;75:1–5.
- [15] Fayaz A, Croft P, Langford RM, Donaldson LJ, Jones GT. Prevalence of chronic pain in the UK: a systematic review and meta-analysis of population studies. *BMJ Open* 2016;6:e010364.
- [16] Fishbain DA, Lewis JE, Cutler R, Cole B, Rosomoff HL, Rosomoff RS. Can the neuropathic pain scale discriminate between non-neuropathic and neuropathic pain? *Pain Med* 2008;9:149–60.
- [17] Forseth KO, Husby G, Gran JT, Forre O. Prognostic factors for the development of fibromyalgia in women with self-reported musculoskeletal pain. A prospective study. *J Rheumatol* 1999;26:2458–67.
- [18] Freynhagen R, Baron R, Gockel U, Tolle TR. painDETECT: a new screening questionnaire to identify neuropathic components in patients with back pain. *Curr Med Res Opin* 2006;22:1911–20.
- [19] Gaskell H, Moore RA, Derry S, Stannard C. Oxycodone for pain in fibromyalgia in adults. *Cochrane Database Syst Rev* 2016;CD012329.
- [20] Gupta A, Silman AJ, Ray D, Morriss R, Dickens C, MacFarlane GJ, Chiu YH, Nicholl B, McBeth J. The role of psychosocial factors in predicting the onset of chronic widespread pain: results from a prospective population-based study. *Rheumatology (Oxford)* 2007;46:666–71.
- [21] Heller RF, Dobson AJ, Attia J, Page J. Impact numbers: measures of risk factor impact on the whole population from case-control and cohort studies. *J Epidemiol Community Health* 2002;56:606–10.
- [22] Holm LW, Carroll LJ, Cassidy JD, Skilgate E, Ahlbom A. Widespread pain following whiplash-associated disorders: incidence, course, and risk factors. *J Rheumatol* 2007;34:193–200.
- [23] Hu J, Bortsov AV, Ballina L, Orrey DC, Swor RA, Peak D, Jones J, Rathlev N, Lee DC, Domeier R, Hendry P, Parry BA, McLean SA. Chronic widespread pain after motor vehicle collision typically occurs through immediate development and nonrecovery: results of an emergency department-based cohort study. *PAIN* 2016;157:438–44.
- [24] Hunt IM, Silman AJ, Benjamin S, McBeth J, Macfarlane GJ. The prevalence and associated features of chronic widespread pain in the community using the “Manchester” definition of chronic widespread pain. *Rheumatology (Oxford)* 1999;38:275–9.
- [25] Koroschetz J, Rehm SE, Gockel U, Brosz M, Freynhagen R, Tolle TR, Baron R. Fibromyalgia and neuropathic pain—differences and

- similarities. A comparison of 3057 patients with diabetic painful neuropathy and fibromyalgia. *BMC Neurol* 2011;11:55.
- [26] Lacey RJ, Belcher J, Rathod T, Wilkie R, Thomas E, McBeth J. Pain at multiple body sites and health-related quality of life in older adults: results from the North Staffordshire Osteoarthritis Project. *Rheumatology (Oxford)* 2014;53:2071–9.
- [27] Lacey RJ, Lewis M, Jordan K, Jinks C, Sim J. Interrater reliability of scoring of pain drawings in a self-report health survey. *Spine* 2005;30:E455–458.
- [28] Larsson B, Bjork J, Borsbo B, Gerdle B. A systematic review of risk factors associated with transitioning from regional musculoskeletal pain to chronic widespread pain. *Eur J Pain* 2012;16:1084–93.
- [29] Livshits G, Ni Lochlainn M, Malkin I, Bowyer R, Verdi S, Steves CJ, Williams FMK. Shared genetic influence on frailty and chronic widespread pain: a study from TwinsUK. *Age Ageing* 2018;47:119–25.
- [30] Macfarlane GJ, Barnish MS, Jones GT. Persons with chronic widespread pain experience excess mortality: longitudinal results from UK Biobank and meta-analysis. *Ann Rheum Dis* 2017;76:1815–22.
- [31] Mansfield KE, Sim J, Jordan JL, Jordan KP. A systematic review and meta-analysis of the prevalence of chronic widespread pain in the general population. *PAIN* 2016;157:55–64.
- [32] Mathieson S, Maher CG, Terwee CB, Folly de Campos T, Lin CW. Neuropathic pain screening questionnaires have limited measurement properties. A systematic review. *J Clin Epidemiol* 2015;68:957–66.
- [33] McBeth J, Harkness EF, Silman AJ, Macfarlane GJ. The role of workplace low-level mechanical trauma, posture and environment in the onset of chronic widespread pain. *Rheumatology (Oxford)* 2003;42:1486–94.
- [34] McBeth J, Macfarlane GJ, Benjamin S, Silman AJ. Features of somatization predict the onset of chronic widespread pain: results of a large population-based study. *Arthritis Rheum* 2001;44:940–6.
- [35] Momi SK, Fabiane SM, Lachance G, Livshits G, Williams FM. Neuropathic pain as part of chronic widespread pain: environmental and genetic influences. *PAIN* 2015;156:2100–6.
- [36] Mulvey MR, McBeth J. Comment on: “Self-reported somatosensory symptoms of neuropathic pain in fibromyalgia and chronic widespread pain correlated with tender point count and pressure-pain thresholds” by Amris et al. [*Pain*;151:664–669]. *PAIN* 2011;152:1684–5.
- [37] Muthuri SG, Kuh D, Bendayan R, Macfarlane GJ, Cooper R. Chronic physical illness in early life and risk of chronic widespread and regional pain at age 68: evidence from the 1946 British birth cohort. *PAIN* 2016;157:2382–9.
- [38] Noble M, McLennan D, Wilkinson K, Whitworth A, Barnes H, Dibben C. The English indices of deprivation 2007. London: Department for Communities and Local Government, 2008.
- [39] Peters MJ, Broer L, Willems HL, Eiriksdottir G, Hocking LJ, Holliday KL, Horan MA, Meulenbelt I, Neogi T, Popham M, Schmidt CO, Soni A, Valdes AM, Amin N, Dennison EM, Eijkelkamp N, Harris TB, Hart DJ, Hofman A, Huygen FJ, Jameson KA, Jones GT, Launer LJ, Kerkhof HJ, de Kruif M, McBeth J, Kloppenborg M, Ollier WE, Oostra B, Payton A, Rivadeneira F, Smith BH, Smith AV, Stolk L, Teumer A, Thomson W, Uitterlinden AG, Wang K, van Wingerden SH, Arden NK, Cooper C, Felson D, Gudnason V, Macfarlane GJ, Pendleton N, Slagboom PE, Spector TD, Volzke H, Kavelaars A, van Duijn CM, Williams FM, van Meurs JB. Genome-wide association study meta-analysis of chronic widespread pain: evidence for involvement of the 5p15.2 region. *Ann Rheum Dis* 2013;72:427–36.
- [40] Smith D, Wilkie R, Croft P, Parmar S, McBeth J. Pain and mortality: mechanisms for a relationship. *PAIN* 2018;159:1112–18.
- [41] Spiegelman D, Hertzmark E, Wand HC. Point and interval estimates of partial population attributable risks in cohort studies: examples and software. *Cancer Causes Control* 2007;18:571–9.
- [42] Topolski TD, LoGerfo J, Patrick DL, Williams B, Walwick J, Patrick MB. The Rapid assessment of physical activity (RAPA) among older adults. *Prev Chronic Dis* 2006;3:A118.
- [43] Torrance N, Smith BH, Bennett MI, Lee AJ. The epidemiology of chronic pain of predominantly neuropathic origin. Results from a general population survey. *J Pain* 2006;7:281–9.
- [44] Vos T, Flaxman AD, Naghavi M, Lozano R, Michaud C, Ezzati M, Shibuya K, Salomon JA, Abdalla S, Aboyans V, Abraham J, Ackerman I, Aggarwal R, Ahn SY, Ali MK, Alvarado M, Anderson HR, Anderson LM, Andrews KG, Atkinson C, Baddour LM, Bahalim AN, Barker-Collo S, Barrero LH, Bartels DH, Basáñez MG, Baxter A, Bell ML, Benjamin EJ, Bennett D, Bernabé E, Bhalla K, Bhandari B, Bikbov B, Bin Abdulhak A, Birbeck G, Black JA, Blencowe H, Blore JD, Blyth F, Bolliger I, Bonaventure A, Boufous S, Bourne R, Boussinesq M, Braithwaite T, Brayne C, Bridgett L, Brooker S, Brooks P, Brughu TS, Bryan-Hancock C, Bucello C, Buchbinder R, Buckle G, Budke CM, Burch M, Burney P, Burstein R, Calabria B, Campbell B, Canter CE, Carabin H, Carapetis J, Carmona L, Cella C, Charlson F, Chen H, Cheng AT, Chou D, Chugh SS, Coffeng LE, Colan SD, Colquhoun S, Colson KE, Condon J, Connor MD, Cooper LT, Corriere M, Cortinovis M, de Vaccaro KC, Couser W, Cowie BC, Criqui MH, Cross M, Dabhadkar KC, Dahiya M, Dahodwala N, Damsere-Derry J, Danaei G, Davis A, De Leo D, Degenhardt L, Dellavalle R, Delossantos A, Denenberg J, Derrett S, Des Jarlais DC, Dharmaratne SD, Dherani M, Diaz-Torne C, Dolk H, Dorsey ER, Driscoll T, Duber H, Ebel B, Edmond K, Elbaz A, Ali SE, Erskine H, Erwin PJ, Espindola P, Ewoigbokhan SE, Farzadfar F, Feigin V, Felson DT, Ferrari A, Ferri CP, Fèvre EM, Finucane MM, Flaxman S, Flood L, Foreman K, Forouzanfar MH, Fowkes FG, Franklin R, Fransen M, Freeman MK, Gabbe BJ, Gabriel SE, Gakidou E, Ganatra HA, Garcia B, Gaspari F, Gillum RF, Gmel G, Gosselin R, Grainger R, Groeger J, Guillemin F, Gunnell D, Gupta R, Haagsma J, Hagan H, Halasa YA, Hall W, Haring D, Haro JM, Harrison JE, Havmoller R, Hay RJ, Higashi H, Hill C, Hoen B, Hoffman H, Hotez PJ, Hoy D, Huang JJ, Ibeanusi SE, Jacobsen KB, James SL, Jarvis D, Jasrasaria R, Jayaraman S, Johns N, Jonas JB, Karthikeyan G, Kassebaum N, Kawakami N, Keren A, Khoo JP, King CH, Knowlton LM, Kobusingye O, Koranteng A, Krishnamurthi R, Lalloo R, Laslett LL, Lathlean T, Leasher JL, Lee YY, Leigh J, Lim SS, Limb E, Lin JK, Lipnick M, Lipshultz SE, Liu W, Loane M, Ohno SL, Lyons R, Ma J, Mabwejian J, MacIntyre MF, Malekzadeh R, Mallinger L, Manivannan S, Marcenos W, March L, Margolis DJ, Marks GB, Marks R, Matsumori A, Matzopoulos R, Mayosi BM, McAnulty JH, McDermott MM, McGill N, McGrath J, Medina-Mora ME, Meltzer M, Mensah GA, Merriman TR, Meyer AC, Miglioli V, Miller M, Miller TR, Mitchell PB, Mocumbi AO, Moffitt TE, Mokdad AA, Monasta L, Montico M, Moradi-Lakeh M, Moran A, Morawska L, Mori R, Murdoch ME, Mwaniki MK, Naidoo K, Nair MN, Naldi L, Narayan KM, Nelson PK, Nelson RG, Nevitt MC, Newton CR, Nolte S, Norman P, Norman R, O'Donnell M, O'Hanlon S, Olives C, Omer SB, Ortblad K, Osborne R, Ozgediz D, Page A, Pahari B, Pandian JD, Rivero AP, Patten SB, Pearce N, Padilla RP, Perez-Ruiz F, Perico N, Pesudovs K, Phillips D, Phillips MR, Pierce K, Pion S, Polanczyk GV, Polinder S, Pope CA III, Popova S, Porri E, Pourmalek F, Prince M, Pullan RL, Ramaiah KD, Ranganathan D, Razavi H, Regan M, Rehm JT, Rein DB, Remuzzi G, Richardson K, Rivara FP, Roberts T, Robinson C, De León FR, Ronfani L, Room R, Rosenfeld LC, Rushton L, Sacco RL, Saha S, Sampson U, Sanchez-Riera L, Sanman E, Schwebel DC, Scott JG, Segui-Gomez M, Shahraz S, Shepard DS, Shin H, Shivakoti R, Singh D, Singh GM, Singh JA, Singleton J, Sleet DA, Sliwa K, Smith E, Smith JL, Stapelberg NJ, Steer A, Steiner T, Stolk WA, Stovner LJ, Sudfeld C, Syed S, Tamburlini G, Tavakkoli M, Taylor HR, Taylor JA, Taylor WJ, Thomas B, Thomson WM, Thurston GD, Tleyjeh IM, Tonelli M, Towbin JA, Truelsen T, Tsilimbaris MK, Ubeda C, Undurraga EA, van der Werf MJ, van Os J, Vavilala MS, Venketasubramanian N, Wang M, Wang W, Wall K, Weatherall DJ, Weinstock MA, Weintraub R, Weisskopf MG, Weissman MM, White RA, Whiteford H, Wiersma ST, Wilkinson JD, Williams HC, Williams SR, Witt E, Wolfe F, Woolf AD, Wulf S, Yeh PH, Zaidi AK, Zheng ZJ, Zonies D, Lopez AD, Murray CJ, AlMazroa MA, Memish ZA. Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012;380:2163–96.
- [45] Wegener ST, Castillo RC, Haythornthwaite J, Mackenzie EJ, Bosse MJ; Group LS. Psychological distress mediates the effect of pain on function. *PAIN* 2011;152:1349–57.
- [46] Wolfe F, Clauw DJ, Fitzcharles MA, Goldenberg DL, Katz RS, Mease P, Russell AS, Russell IJ, Winfield JB, Yunus MB. The American College of Rheumatology preliminary diagnostic criteria for fibromyalgia and measurement of symptom severity. *Arthritis Care Res* 2010;62:600–10.
- [47] Wolfe F, Smythe HA, Yunus MB, Bennett RM, Bombardier C, Goldenberg DL, Tugwell P, Campbell SM, Abeles M, Clark P, Fam AG, Farber SJ, Fiechtner JJ, Franklin CM, Gatter RA, Hamaty D, Lessard J, Lichtbroun AS, Masi AT, Mccain GA, Reynolds WJ, Romano TJ, Jon Russell I, Sheon RP. The American College of Rheumatology 1990 criteria for the classification of fibromyalgia. Report of the multicenter criteria committee. *Arthritis Rheum* 1990;33:160–72.
- [48] Zigmund AS, Snaith RP. The hospital anxiety and depression scale. *Acta psychiatrica Scand* 1983;67:361–70.