



UNIVERSITY OF LEEDS

This is a repository copy of *Invited Review: IPCC, Agriculture and Food - A Case of Shifting Cultivation and History*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/146882/>

Version: Accepted Version

Article:

Porter, JR, Challinor, AJ orcid.org/0000-0002-8551-6617, Henriksen, CB et al. (3 more authors) (2019) *Invited Review: IPCC, Agriculture and Food - A Case of Shifting Cultivation and History*. *Global Change Biology*, 25 (8). pp. 2518-2529. ISSN 1354-1013

<https://doi.org/10.1111/gcb.14700>

© 2019 John Wiley & Sons Ltd. All rights reserved. This is the post-peer reviewed version of the following article: Porter, JR, Challinor, AJ, Henriksen, CB et al. (3 more authors) (2019) *Invited Review: IPCC, Agriculture and Food - A Case of Shifting Cultivation and History*. *Global Change Biology* 25 (8). pp. 2518-2529, which has been published in final form at <https://doi.org/10.1111/gcb.14700>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

1 **Running Title:** IPCC, Agriculture and Food

2 **Invited Review: IPCC, Agriculture and Food – A Case of Shifting Cultivation and History**

3 John R Porter^{1,2}, Andrew J. Challinor³, Christian Bugge Henriksen², S Mark Howden⁴, Pierre
4 Martre⁵, Pete Smith⁶

5 ¹CIHEAM-IAMM – SupAgro - MUSE Univ Montpellier, 34000 Montpellier, France.

6 ²Plant and Environmental Sciences, University of Copenhagen, Højbakkegård Alle 26, 2360
7 Taastrup, Denmark.

8 ³Institute for Climate and Atmospheric Science (ICAS), School of Earth and Environment,
9 University of Leeds, Leeds, LS2 9JT, UK.

10 ⁴Climate Change Institute, Australian National University, Canberra, 2600 ACT, Australia.

11 ⁵LEPSE, Université Montpellier, INRA, Montpellier SupAgro, 34000 Montpellier, France.

12 ⁶Institute of Biological and Environmental Sciences, University of Aberdeen, 23 St Machar
13 Drive, Aberdeen, AB24 3UU, UK.

14 **Contact Information** John R Porter; +45 29176961 or +336 3535 1498. Email: jrp@plen.ku.dk

15 **Keywords:** adaptation, climate change, food security, impact, IPCC, mitigation, policy.

16 **Paper type:** Invited Review.

17 **Abstract**

18 Since 1990 the Intergovernmental Panel on Climate Change (IPCC) has produced five
19 Assessment Reports (ARs), in which agriculture as the production of food for humans via crops
20 and livestock have featured in one form or another. A constructed data base of the *ca.* 2,100
21 cited experiments and simulations in the five ARs were analysed with respect to impacts on
22 yields via crop type, region and whether or not adaptation was included. Quantitative data on
23 impacts and adaptation in livestock farming have been extremely scarce in the ARs. The main
24 conclusions from impact and adaptation are that crop yields will decline but that responses have
25 large statistical variation. Mitigation assessments in the ARs have used both bottom-up and top-
26 down methods but need better to link emissions and their mitigation with food production and
27 security. Relevant policy options have become broader in later ARs and included more of the
28 social and non-production aspects of food security. Our overall conclusion is that agriculture
29 and food security, which are two of the most central, critical and imminent issues in climate
30 change, have been dealt with in an unfocussed and inconsistent manner between the IPCC five
31 ARs. This is partly a result of agriculture spanning two IPCC working groups but also the very
32 strong focus on projections from computer crop simulation modelling. For the future, we
33 suggest a need to examine interactions between themes such as crop resource use efficiencies
34 and to include all production and non-production aspects of food security in future roles for
35 integrated assessment models. (253 words).

36 **1 | Introduction**

37 Agriculture and the local, regional and global food system encompass what most people on
38 Earth do for a living. If one includes the downstream food system from the production to the
39 consumption of food by humans and other animals – the engagement of humans in food security
40 and food production systems dwarfs any other human activity; including computing,

41 pharmaceuticals, the media, energy industry, banking and academia - combined. Agriculture and
42 food production, distribution, marketing and consumption contribute about 30% of global gross
43 domestic product (Braun *et al.* 2017), and have easily higher returns on investment than any
44 economic corporation, sector or activity - but receive only about 5% of global research
45 investment (Pardey *et al.*, 2016). Agriculture and food systems however, are highly affected by
46 climate changes and also drive climate change through greenhouse gas emissions and land use
47 change.

48 The scientific bedrock of the agreement at the 21st Conference Of the Parties (COP21) of the
49 United Nations Framework Convention on Climate Change in Paris in December 2015 were the
50 5th Assessment Reports (AR5) of the Intergovernmental Panel on Climate Change (IPCC) from
51 2013 and 2014 (IPCC Assessment Reports are available at <https://www.ipcc.ch/reports/>). The
52 statement from COP21 reads '*Recognizing the fundamental priority of safeguarding food*
53 *security and the vulnerabilities of food production systems to the adverse impacts of*
54 *climate change*' acknowledging the central role of food security regionally and globally.
55 Important inter-disciplinary departures in the food security chapter of the IPCC (Porter *et al.*,
56 2014) were recognition of factors other than food production in food security: such factors
57 include food distribution and social and economic access to food, which all stand to be affected
58 by climate change and which have possibilities for adaptation. Food security and agriculture
59 have not always had such a clear or prominent position in IPCC ARs - with food security only
60 specified in AR5 and with agriculture often rolled in with forestry and forest products (AR1 and
61 AR4) or general ecosystem services (AR3). AR2 did examine impacts and adaptation of
62 agriculture. We regard the evolution of a food system perspective in IPCC AR5 as a very
63 positive development that we hope will be amplified in AR6 and future IPCC Special Reports.

64 This review aims to develop further, and in more detail, the recent paper by Porter *et al.* (2017)
65 on the link between the five IPCC Assessment Reports (AR1 to AR5) and agriculture. Space

66 constraints in that article prevented presentation of topics such as regional differences in
67 assessments of impacts, adaptation and mitigation linked to agriculture; the balance between
68 assessment of climate change and crops versus livestock; the methods used and how and why
69 assessments might develop in the future. Post AR5 (Porter *et al.*, 2014), the Royal Society of
70 London (Royal Society, 2017) published an update on climate change effects on food
71 production. Their conclusion was that post-AR5 studies have confirmed conclusions in AR5, but
72 new studies ‘point strongly to the importance of accounting for how land use and cropping
73 intensity might change’. Our review addresses the above gaps and addresses potentially policy-
74 relevant information that has become available since the AR5.

75 In addition, as this is an invited review, we have allowed ourselves the licence to include two
76 issues which we think are of importance for future assessment reports dealing with food,
77 agriculture and climate change. Section 6.1 presents ideas to improve the robustness of crop
78 models that have been the ‘work horses’ of many, perhaps too many, climate change
79 assessments. Models should check that they simulate accurately all crop and soil responses
80 underlying responses to climate and we suggest a way for doing this. Secondly, models should
81 be able to simulate accurately the interactions between resources such as radiation, water and
82 nutrients with and without changes in CO₂ level, an issue that rarely has been investigated (*cf.*
83 Teixeira *et al.*, 2014). Simulating such interactions correctly is particularly important when it
84 comes to examining adaptation options for crops. We show an example where this examination
85 has been done for a well-known and well-used data set. The second issue we raise is the
86 integrated assessment of adaptation and mitigation. Whilst crop models are generally responsive
87 to climate, the range of crops that can be simulated is not sufficiently broad for a full assessment
88 of food security – this is clear from the data presented in this paper. Further, disparities between
89 IAMs and crop models in spatial scale, treatment of uncertainty, data demand and representation
90 of agricultural management all limit the extent of crop model integration into IAMs that is

91 currently possible. We think that more than one approach is needed if we are to capture the
92 range of trade-offs and synergies that are important to food systems and relevant to policy
93 design and development. We need to recognise that emissions occur across the full range of
94 activities that deliver food security, not only agricultural production but with a focus on climate-
95 smart food systems.

96 **2 | Projected impacts**

97 To get an overview of the assessment of the projected impacts of climate change on crop yield
98 across the five IPCC Assessment Reports, across the different global regions and for the major
99 global crops, we compiled all data (2116 entries) on projected crop yield with and without
100 adaptation from AR1-AR5. We constructed a database with information about the AR volume,
101 crop type, global region and projected mean change and variation in yields with and without
102 adaptation. In this context adaptation refers to all adaptation measures investigated in the
103 scenarios throughout AR1 to AR5, including but not limited to altering sowing times, crop
104 cultivars and species, adjusting irrigation and fertilizer application, reducing tillage and
105 implementing technical measures to more effectively capture rainwater and reduce soil erosion.
106 Subsequently, the average mean change in yield with and without adaptation was calculated for
107 each IPCC Assessment Report, each global region and each major global crop (Tables 1-3). By
108 reviewing the constructed dataset it quickly becomes evident that the number of cases increases
109 almost exponentially (except from AR2 to AR3), thereby increasing the confidence level of the
110 results of each subsequent report. A striking omission across the five ARs is the almost
111 complete lack of quantitative data of the effects of climate change on livestock; no quantitative
112 data were presented from AR1 to AR3 and only 18 cases were reported in AR4 and AR5
113 combined (Rivera-Ferre *et al.* 2016).

114

Table 1. Mean percent change in average yield of all crops reported in AR1-AR5 with and without adaptation.

IPCC AR	Publication Year	With adaptation			Without adaptation		
		Number of cases	Mean change (%)	Standard deviation (%)	Number of cases	Mean change (%)	Standard deviation (%)
AR1	1990	6	9.0	11.5	28	3.4	33.0
AR2	1995	46	-0.2	23.1	53	-13.8	25.8
AR3	2001	57	-8.2	17.4	36	-5.2	23.4
AR4	2007	239	3.6	19.0	320	-4.0	17.7
AR5	2015	519	-3.9	17.2	812	-9.9	19.4

115 All IPCC Assessment Reports, except AR1 have projected a crop yield reduction without
116 adaptation (Table 1). The largest projected yield reduction was in AR2 with -13.8% followed by
117 -9.9% in AR5. When climate adaptations were included in the analysis, most assessment reports
118 also projected a yield reduction except for AR1 with a 9.0% yield increase and AR4 with a 3.6%
119 yield increase. However, the standard deviations in the projections are large, ranging from
120 11.5% to 33.0%.

Table 2. Mean percent change in average yield for different global regions summarized for AR1-AR5 with and without adaptation. When constructing the database, the results from AR1-AR5 were allocated to the IPCC AR5 global regions by following the following rules: data from Russia and former Soviet Union were allocated to the global region North Asia; data from Middle East and North Africa were allocated to the global region West Asia; data from Latin America and the Caribbean were allocated to Central and South America; data from south-east Mediterranean (Jordan, Egypt and Libya) were allocated to the global region Africa; data from Pacific Asia and Pacific OECD were allocated to the global region Australasia.

	With adaptation			Without adaptation		
	Number of cases	Mean change (%)	Standard deviation (%)	Number of cases	Mean change (%)	Standard deviation (%)
Africa	153	-4.2	19.8	274	-9.5	17.7
Australasia	38	6.9	17.7	38	-7.1	21.7
North America	109	1.2	17.3	167	-7.8	25.8
Central and South America	74	-12.6	17.7	91	-12.1	15.8
Europe	68	3.3	22.0	164	-4.3	21.3
North Asia	10	8.9	11.3	6	-14.0	17.7
East Asia	126	-1.5	14.6	175	-4.9	16.3
Central Asia	11	-3.9	18.4	9	-19.2	18.3
West Asia	8	-8.4	6.9	18	-5.0	11.7
South Asia	138	0.1	16.2	199	-11.7	18.9
South-east Asia	31	10.4	20.7	41	-0.6	14.0
Asia (unspecified)	18	-14.0	17.8	6	-2.3	11.2
Global	74	-6.4	17.5	37	-17.9	20.1

121 The standard deviation is also large for the mean change in yield for different global regions
122 (Table 2). Without adaptation Central Asia had yield change of -19.2%, followed by North Asia
123 with -14.0%, Central and South America with -12.1% and South Asia with -11.7% are the
124 regions with the largest projected yield decreases. With adaptation, South-east Asia, North Asia
125 and Australasia have the largest yield increase with +10.4%, +8.9% and +6.9%, respectively.

126

Table 3. Mean change in yield for different crops summarized for AR1-AR5 with and without adaptation.

							128
With adaptation			Without adaptation			129	
Crop	Mean		Standard	Mean		Standard	130
	Number	change	deviation	Number	change	deviation	
	of cases	(%) ^a	(%) ^a	of cases	(%)	(%)	131
Barley	1	-35.0	n/a	7	0.7	14.4	
Beans	1	45.0	n/a	12	-38.7	37.1	132
Cassava	0	n/a	n/a	21	-2.2	3.9	
Grass	4	11.8	24.	6	-8.5	45.7	133
Groundnut	3	34.0	17.	11	-6.6	12.5	134
Maize	303	-5.6	16.	281	-10.8	18.2	
Millet	2	-27.0	13.	111	-9.3	20.4	135
Potato	0	n/a	n/a	19	-2.0	17.4	136
Rice	140	3.4	15.	231	-5.3	14.7	
Sorghum	2	-23.5	37.	21	-9.1	7.8	137
Soybean	73	-12.8	17.	83	-16.9	27.0	138
Sugarcane	0	n/a	n/a	18	-2.5	9.8	
Sunflower	0	n/a	n/a	10	-3.1	6.1	139
Sweet potato	0	n/a	n/a	5	-2.2	7.2	
Wheat	225	1.9	21.	343	-7.0	20.6	140
							141

142 For major global crops (Table 3), it is evident that the crops most severely affected by climate
143 change without adaptation are soybean and maize with yield reductions of -16.7% and -10.8%,
144 respectively. The yield reduction for beans is even larger but only based on a single observation.
145 For protein crops, this yield reduction is particularly alarming given their potential to replace

146 meat-based protein with both health and greenhouse gas emissions benefits (Tilman and Clark
147 2014). Also, besides maize, some of the other major staple crops for the Southern Hemisphere
148 are projected to have significant yield reductions without adaptation, e.g. -10.8% for maize, -
149 9.3% for millet, -9.1% for sorghum and -16.9% for soybean. Even with adaptation, large yield
150 reductions are projected for maize (-5.6%), millet (-27.0%), sorghum (-23.5%) and soybean
151 (12.8%). Considering that these three crops cover 60% of the area cultivated with cereal crops in
152 Africa and provide 67% of the cereal yield on the continent (Macauley, 2015), a yield reduction
153 of this magnitude would have severe consequences. Overall, adaptation is not projected to have
154 a very large effect on reducing or even reversing yield reductions for the major global crops.
155 Large yield increases can be seen for beans, groundnut and grass, but these results are only
156 based on few observations. Based on this analysis, rice and wheat, with yield increases of +3.4%
157 and +1.9%, seems to be the only major global crops to benefit from adaptation efforts. It is
158 worth noting that only 15 crops are included in the IPCC Assessment Reports. It is evident that a
159 more accurate assessment of food security would require that a much larger number of crops are
160 investigated.

161 **3 | Adaptation**

162 From the first IPCC Assessment onwards, a systems approach has been applied to the analysis
163 of climate impacts and adaptation relating to agriculture, food production and, more recently,
164 food systems. However, both the supporting literature and the emphasis and framing of this have
165 changed significantly over the five IPCC ARs, with a relative increase in the number of studies
166 including adaptations to impacts. In AR1, there was relatively little quantitative literature on
167 climate change impacts and so a conceptual systems approach was used to identify the likely
168 impacts and their interlinkages. These included suggestions that changing crop yields could lead
169 to potential changes in geographical distribution of cropping. The coverage was of average
170 agricultural production, paleo-analogues and basic physiological responses such as laboratory

171 responses of plants to CO₂ to support scenarios of future impacts. The main focus was on cereal
172 crops rather than livestock or other food-producing systems such as horticulture. Studies were
173 almost exclusively drawn from the temperate zones and from developed nations. Subsequent
174 IPCC assessments of climate impacts on production of the major crops (wheat, rice, maize and
175 soybean) have significantly increased in complexity, drawing from the expanding literature
176 base. The increase in the number and coverage of studies has successively allowed tabulation of
177 crop responses (AR3), and then meta-analyses initially developing simple relationships (AR4)
178 and subsequently statistical relationships between variables (AR5; Challinor *et al.* 2014). In
179 particular the crop modelling studies have evolved from simple, often site-based scenarios
180 driven by fixed temperature and rainfall changes (e.g. +3°C and -20% rainfall) towards
181 integration of downscaled GCM data in grid-based or multi-site, regional assessments.
182 Nevertheless, the focus of the IPCC remained on mean yield change and it was only in AR3 was
183 there inclusion of a focus on changes in yield variability and, in AR5, the nutritional quality of
184 crops. Whilst there are regional and global crop production studies there are have been few
185 impact studies which have used a value chain or a food systems perspective. Developing country
186 studies remain relatively under-represented in terms of population (Table 2), even though
187 developing countries were identified as early as AR2 that they were likely to be the most
188 negatively affected. Similarly, even though AR2 concluded that elevated atmospheric CO₂
189 concentrations would have beneficial impacts on crop production, there remained active debate
190 in AR5 about the degree to which this may affect crop yields and quality.

191 As noted above, there has been relatively little quantitative treatment of livestock (Rivera-Ferre
192 *et al.* 2016), other field-crops, horticulture and viticulture across IPCC reports with coverage
193 being largely restricted to either generic, system-level responses or site-specific cases, largely
194 because of the relative lack of studies using somewhat-comparable modelling or other analysis
195 methods in contrast to the mechanistic and other crop models, which have enabled meta-

196 analysis, cross-model comparison and assessment of uncertainties (Rosenzweig *et al.* 2014). The
197 treatment of weeds, pests and disease impacts are also inconsistently dealt with across the
198 reports for the same reasons.

199 The aggregation of climate change impacts on food production systems to broad-scale economic
200 and food price impact was also initiated in the AR2 with results reported from two economic
201 models (Rosenzweig and Parry 1994, Reilly *et al.*, 1994. Successive IPCC Assessment Reports
202 have synthesised the rapidly developing literature to not only address global and regional
203 impacts of climate change on prices, production and trade but also the uncertainties in model
204 results and the reasons behind these (e.g. Nelson *et al.* 2014).

205 Adaptation to the sorts of climate change impacts noted above is a fundamental part of risk
206 management. Agriculture and food producers as well as value chain managers, consumers and
207 policy makers have shown considerable ability to adapt to climate changes both currently and
208 going back into history; for instance, the establishment of grapevines in England in Roman
209 times or the settlement of Greenland in medieval times. The expectation that adaptation of food
210 systems is likely to be both feasible and attractive has resulted in coverage from AR1 onwards.
211 However, the framing, scope, likely effectiveness and analytical methods used in IPCC reports
212 has changed significantly since then (Table 4). There remain many gaps in terms of adaptation
213 of food systems including, but not limited to, the need to include assessment of more
214 transformative adaptations, adaptation of value chains and of regional food systems. Other
215 important issues for the future include how to address the multitudinous barriers to adaptation,
216 developing the pathways to not only build adaptive capacity but to also move this into
217 adaptation actions, developing policies and programs to establish effective monitoring,
218 evaluation and attribution of adaptation and assessments and to more effectively address net
219 greenhouse gas emission reduction within adaptation strategies. This latter point is starting to be

220 addressed in the IPCC AR6 cycle, being covered by two Special Reports (www.ipcc.ch/reports)
221 as well as within the main Assessment Reports.

222

223

Table 4. The framing, scope and analysis methods used to address climate adaptation in agricultural and food systems in successive IPCC Assessment Reports (ARs).

IPCC	Framing, scope and analysis methods used
Assessment	
AR1 1990	<i>Framing:</i> Three adaptation domains - physiological adaptation, farm level management ‘adjustment’ and responses arising from policy at regional, national and international levels. These were expressed in terms of enabling farming systems to reach a new equilibrium in response to altered climates. <i>Scope:</i> Farm-level, production focus not food systems. <i>Analysis:</i> Generally, adaptations were described qualitatively using historical analogues or first principles approaches rather than quantified responses.
AR2 1995	<i>Framing:</i> Spontaneous or planned adaptation, in response to or anticipation of climate change. <i>Scoping:</i> Farm-level production system focus not food systems with brief reference to global economic analyses of producer surplus, which included with and without adaptation. <i>Analysis:</i> Few quantitative adaptation studies although most adaptation options were raised based on a systems view. However, these were mostly incremental such as agronomic adjustments although there were some systemic adaptations (<i>sensu</i> Rickards and Howden, 2012) such as the introduction of new species.

There was a recognition that successful adaptation depends upon technological advances, institutional arrangements, availability of financing and information exchange as well as adaptive capacity and alignment of the options with farmer needs so as to enhance adoption paths. Additionally, there was recognition of the possibility of policy maladaptation.

AR3 *Framing:* No specific framing, focused on farm-level, agronomic changes.

2001 *Scope:* Farm-level production focus not food systems, with examples of integrated regional economic analyses of impacts and adaptation.

Analysis: As well as qualitative discussion of options such as crop breeding to adjust to elevated CO₂ and temperatures, there were more quantitative analyses of cropping system adaptations allowing both tabular and figure summaries of the modelled effectiveness of adaptation. However, there was a critique that methodologically, there had been little progress since the previous IPCC Assessment with the adaptation strategies being modelled limited to a small subset of the possible options and unrealistic assumptions regarding the degree and effectiveness of farmer adoption. There was recognition of adaptation costs including transition costs, dislocation costs and capital and operational costs.

There was however, limited coverage of livestock adaptation with discussion of a range of management adaptations to reduce the effects of heat waves but few quantified or modelled analyses to draw from.

- AR4** *Framing:* Autonomous and planned adaptation modes.
- 2007** *Scope:* Recognition of the importance of a food systems approach but the focus remained on agricultural production.
- Analysis:* Discussion of a broader range of possible adaptation options for both cropping and livestock using a more structured approach particularly drawing off the burgeoning literature on cropping system impacts and adaptations. This allowed more geographically explicit analyses as well as a meta-analysis of impacts and adaptation as a function of temperature increase. However, most adaptation options addressed were still incremental in nature, reflecting in part limitations of the modelling approaches being used. There was a critique of the failure to provide generalised knowledge of adaptive capacity, of adoption pathways and barriers to these and of a more comprehensive range of adaptation strategies especially beyond simple, single agronomic changes. There was still limited evaluation of the costs of adaptation or of consequences of adaptation in relation to the environment and the natural resource base.
- AR5** *Framing:* incremental to transformational adaptation.
- 2014** *Scope:* Food systems approach although much of the literature able to be synthesised was on food production only.
- Analysis:* Discussion of a broad range of possible adaptation options and their adoption paths for both cropping and livestock using a consistent framing. The further increase in the literature on cropping system impacts and adaptations allowed 1) an improved meta-analyses of impacts and adaptation as a function of temperature providing finer-grained information across the major crops, by broad region and disaggregating results to allow assessment of the effectiveness of different agronomic adaptation options; and 2) a meta-analysis

of the possible increase in crop yield variability over time. Livestock adaptations were not able to be dealt with as comprehensively as cropping systems due to limitations in the literature. There was increased recognition of the importance of institutional limits and adoption barriers but some other issues identified as shortcomings in prior IPCC Assessments remain largely unaddressed (e.g. adaptation costs, lack of methodological innovation and diversity in adaptation analysis).

224

225 **4 Mitigation**

226 For mitigation potential in the agriculture sector, methods have changed markedly over the
227 course of the IPCC Assessment Reports. Bottom-up methods, assessing mitigation potential
228 practice-by-practice using data on land areas and livestock numbers available, were used in AR1
229 and AR2. AR3 largely replaced this approach with a top-down assessment from integrated
230 assessment models (IAMs). For both AR4 and AR5, both bottom-up and top-down estimates
231 were included in conjunction. IAMs have the advantage that they can consider mitigation
232 options across sectors and select least-cost options and pathways for mitigation, which bottom-
233 up approaches cannot. Their disadvantage, however, is the limited number of agricultural
234 options that they include, which are mostly confined to non-CO₂ greenhouse gases. Bottom-up
235 methods, on the other hand, capture the rich detail of the agricultural practices available
236 (Bennetzen *et al.*, 2016) but are unable to consider mitigation across sectors, so estimates of
237 economic potential are more uncertain. The combination of top-down and bottom-up approaches
238 will likely prove useful again in AR6.

239 Chapters dealing with climate change mitigation in the IPCC Assessment Reports have been
240 weak in linking emissions with the primary purpose of agriculture, i.e. producing food. For

241 example, demand-side measures to limit greenhouse gas emissions through changes in human
242 diet or through waste reduction were not considered in detail until AR5 following the IAASTD
243 report (McIntyre *et al*, 2009) and other publications. Systematic changes in the food system
244 have been under-represented compared to technical interventions, such as changes in
245 fertilisation, livestock feed-additives and changes in tillage practice, on farm. This is perhaps
246 driven by the sectoral approach taken in most assessments. For example, greenhouse gas
247 emission reductions through fossil fuel offsets by production of bioenergy are not accounted for
248 in the agriculture sector, so are not reported in the agriculture or land chapters. Reduced energy
249 consumption in agriculture is not reported in the agricultural and land sector, nor any emission
250 reductions associated with improved packaging, transport, distribution and storage. Taking an
251 approach based on the sectors from which emissions are reported is logical, but does not
252 encourage food systems approaches to addressing emission reduction goals. Future assessments
253 will need to take a more holistic view of the food system, and go beyond the accounting /
254 reporting sectors considered to date.

255 Another persistent issue across IPCC Assessment Reports arises from the structure in which
256 assessments are conducted, with Working Group 1 focussing on the physical science basis of
257 climate change, Working Group 2 focussing on impacts of climate change and adaptation, and
258 Working Group 3 focussing on mitigation. The chapters dealing with agriculture and land in
259 each Assessment Report are written by different authors and appear in different volumes,
260 corresponding to each Working Group. While efforts are made to encourage cross-working
261 group / cross-volume collaboration and consistency, results have been uneven, with a number of
262 disconnects in emphasis across the volumes.

263 The IPCC Special Report on Climate Change and Land, under production as part of the AR6
264 cycle and due in 2019, offers an opportunity to address some of the issues raised above. Firstly,
265 it is a joint action across the three Working Groups, thereby including experts from more

266 disciplines than usually found within Working Groups. Secondly, it considers a wide range of
267 land and climate change related issues, including mitigation, adaptation, desertification, land
268 degradation, sustainable land management and food security. With an emphasis on integrated
269 response options to address all of these challenges, considering synergies and trade-offs, it
270 necessarily takes a broader view of land, agriculture, food systems and the interventions
271 available to address the considerable challenges facing humanity now and in the future. While
272 examining all of these factors together is extremely challenging, due to the complexity of the
273 sectors involved, the importance of food and agriculture and climate change for the future of
274 humanity means it is a challenge that must be met. Future IPCC Assessment Reports could learn
275 from the experience of producing this Special Report – to take a broader view of the issues
276 facing land and agriculture, and to facilitate cross Working Group integration.

277 **5 | Policy**

278 The policy elements of climate adaptation and mitigation in relation to agriculture and food
279 systems have been addressed unevenly and incompletely over the various Assessment Reports.
280 AR1 acknowledged the importance of a range of policies (listing food price, land-use, forest
281 resources, extension and water transfers) but required more information in relation to potential
282 responses. AR2 expanded the list to include research, land-use planning, water pricing and
283 allocation, disaster vulnerability assessment, transport and trade policy and policies countries
284 use to encourage or control production, limit food prices and manage resource inputs to
285 agriculture. There was a brief critical analysis of how policies may discourage adaptation
286 strategies and acknowledgement of the political, economic and cultural factors at play but
287 overall very little concrete guidance in relation to policy design and development. In contrast,
288 the AR3 and AR4 provided few linkages to policy and it was not until AR5 that more policy-
289 relevant suggestions were developed. These included *inter alia* capacity building across the food
290 system via support of monitoring and communication, systems analysis, extension capacity and


291 industry and regional networks that develop social capital and share information, supporting
292 community partnerships in developing food and forage banks, enhancing investment in
293 irrigation infrastructure and efficient water use technologies, revising land tenure arrangements
294 (including attention to well-defined property rights), establishment of accessible, efficiently
295 functioning markets for inputs and outputs (seed, fertiliser, labour, water, products, greenhouse
296 gases emissions, etc.) and for financial services, including insurance. There was also
297 introduction of ideas relating to modes of operation such as policy ‘mainstreaming’ and policy
298 analysis methodologies such as the need for multi-level assessment. Importantly, these policy
299 inclusions in AR5 were consistent with moving away from the previous ‘agricultural
300 production’ focus to a more ‘food systems’ focus but nevertheless did not substantially progress
301 the integrated treatment of climate adaptation and mitigation.

302 **6 | Future improvements in examining impacts and adaptation**

303 **6.1. | Assessing crop growth models skills to predict interactions between resource use** 304 **efficiencies**

305 The main types of models used in IPCC impact assessments on crop production fall into the
306 category of crop simulation models, that attempt to predict yields based on bio-climatic inputs
307 and are mostly site-based; statistical relationships have also been used (Porter *et al.*, 2014). Such
308 models are only just being used to examine CO₂ and other effects on yield and its protein
309 concentration (Asseng *et al.*, 2019) even though this topic has been a persistent theme in the
310 ARs. Thus as suggestions, we wish to highlight the need to analyse the interactions between
311 resource use efficiencies to change the consistency of crop models and better understand
312 cropping systems response to climate change as a topic, focused on modelling. We think this is
313 an important topic for future assessment of climate impacts, adaptation and mitigation within the
314 land-sector and agriculture and their position and role in climate change.

315 Bennetzen *et al.* (2016) showed via a historical deconstruction analysis, using a modified Kaya
316 identity analysis (Kaya and Yokoburi, 1997), that greenhouse gas emissions from agriculture
317 have decoupled from food production since 1970, and give grounds for optimism that
318 agriculture can make a substantial contribution to reducing global emissions as well as helping
319 to store carbon in land. A reduction of emissions per unit product means that the utilization
320 efficiency of the principle inputs into food production, namely water and fertilizer, has
321 increased. At the same time crop simulation models have been used extensively to project the
322 impacts of changes in CO₂, temperature, rainfall and other factors for global and regional
323 productivity of crops (e.g. Ruane *et al.*, 2017). Resource utilisation efficiencies do not operate in
324 isolation; that is to say that there are interactions between, for example, a crop's utilisation
325 efficiency of water, nitrogen and photosynthetically active short-wave radiation. How far these
326 interactions of resource utilisation efficiencies are incorporated into crop models is unclear and
327 needs testing, together with a critical need to design and make experiments to test the models.
328 Models should not get the 'right' answers for the 'wrong' reasons such as via cancellation of
329 errors (Challinor *et al.*, 2014; Martre *et al.*, 2015).




330

331 **Figure 1.** Decomposition of water use efficiency. In (a) identity showing the relationship
 332 between water (WUE) and radiation (RUE) use efficiencies and water, nitrogen and light
 333 utilization trade-offs. In (b) four quadrants visual representation of the identity shown in (a). In
 334 quadrants 1 and 2, the thick lines are the upper limits of RUE and WUE, respectively. In
 335 quadrant 1, the plateau is the potential grain yield defined as the grain yield that can be attained
 336 by current cultivars grown in an environment to which it is adapted with water, nutrients and
 337 other abiotic and biotic factors controlled effectively (Evans and Fisher, 1999). In quadrants 3
 338 and 4, the thick lines are critical N uptake, defined as the minimum N uptake for achieving
 339 maximum above ground biomass at the upper limits WUE and RUE, respectively.

340 To this end, we propose a methodology based on mathematical identities (Porter *et al.*, 2013)
 341 that decomposes water and nitrogen utilisation efficiencies and portrays their interactions or
 342 trade-offs with water utilisation efficiency. The ideas stem originally from the work of CT de
 343 Wit and his colleagues at Wageningen, NL and have been developed by others (Teixera *et al.*,
 344 2014; Sadras, 2016) but has seemingly not as yet penetrated crop modelling as an issue for
 345 climate change impacts (Ruane *et al.*, 2017) . The identity for water utilisation efficiency

346 (WUE) and its graphical portrayal (Figure 1) show a possible relationship between WUE and
 347 radiation utilisation efficiency (RUE). Questions for that need responses from crop models
 348 including ‘what are the modelled upper limits for RUE and WUE in ambient and changed
 349 climate pathways and how do they compare with observations?’ and ‘In comparison with a
 350 control treatment, how do the utilisation efficiencies change and interact?’ Crop models should
 351 be able to populate such analyses and we give an example (Figure 2) using the *SiriusQuality*
 352 wheat model (Martre *et al.*, 2006; Martre and Dambreville, 2018;
 353 <http://www1.clermont.inra.fr/siriusquality/>). The simulations are of a four-year CO₂ enrichment
 354 experiment on spring wheat at Maricopa, USA (Kimball *et al.*, 2017) in which the crops were
 355 grown in ambient and elevated CO₂ for combinations of either high or low levels of nitrogen
 356 and of either full or reduced irrigation (see Figure 2 caption for details).



357

358 **Figure 2.** Effect of nitrogen supply, water supply, and atmospheric CO₂ concentration on
359 resource use efficiency and trade-offs illustrating the identity in Figure 1a. A Free air CO₂
360 enrichment experiment conducted over a four years period with a spring wheat cultivar at
361 Maricopa, AZ, USA (Kimball *et al.*, 2017) was simulated with the wheat simulation model
362 *SiriusQuality* (Martre *et al.*, 2006; Martre and Dambreville, 2018). In the first two years wheat
363 crops were grown with high (38.9 g N m⁻²) and low (7.6 g N m⁻²) nitrogen supply under ambient
364 (370 ppm; aCO₂) and elevated (550 ppm ; eCO₂) atmospheric CO₂ concentration. In the
365 following two years a fully irrigated (665 mm) and a water deficit (330 mm) treatments were
366 factorized with the same two CO₂ treatments. In (a) and (b), black dashed lines are upper limits
367 of grain yield calculated with potential radiation use efficiency (2.93 g above ground DM MJ⁻¹
368 PAR; Sinclair and Muchow, 1999), harvest index (0.6; Foulkes *et al.*, 2011), and water use
369 efficiency (2.2 g grain DM m⁻² mm⁻¹; Sadras and Angus, 2006) for wheat, and orange dashed
370 lines are RUE and WUE isopleths calculated with measured data for the control treatment,
371 respectively. In (c) and (d), dashed lines are critical crop N uptake defined as the minimum N
372 uptake for achieving maximum above ground biomass calculated using the RUE and WUE
373 shown in (a) and (b) and the N dilution curve for wheat (Justes *et al.*, 1994). The solid lines
374 between eCO₂ and aCO₂ are drawn to improve the reading of the figure.

375

376 The upper part of Figure 2 shows measured and simulated resource utilisation for radiation
377 (Figure 2a) and water (Figure 2b) when quantified as intercepted PAR or evapotranspiration
378 against crop grain yield. The black dotted lines shows the theoretical potential RUE and WUE
379 and the orange dashed line shows these utilisation efficiencies for the control treatment in
380 ambient CO₂ and with ample water and nitrogen supplies. Points above the orange lines mean
381 that utilisation efficiency is increased relative to control and *vice versa*. Points above the black
382 lines would be above the theoretical resource efficiencies and would therefore be suspicious.
383 Under ambient CO₂, simulations agreed reasonably well with the field measurements but the
384 model underestimated RUE and WUE under water deficit. A higher CO₂ concentration
385 increased both utilisation efficiencies. The model simulated well the effect of elevated CO₂ on
386 RUE but it overestimated the effect of elevated CO₂ on WUE (+23% vs. +14%). Terms 3 and 4
387 in Figure 1a, which measure the trade-offs between N, radiation, and evapotranspiration, are
388 shown in the lower part of Figure 2. The dashed lines show critical N uptake (that is, the
389 minimum crop N uptake for achieving maximum above ground biomass) considering the

390 theoretical potential utilisation efficiencies (black lines) and those for the control treatments
391 (orange lines). For the control and the water deficit treatment, crop N was close to the critical N
392 uptake, especially under elevated CO₂. The increase of crop N uptake under elevated CO₂ is
393 consistent with the reported higher crop N demand under elevated CO₂ (Rogers *et al.*, 2006).
394 Points for the low N treatment were significantly above the critical N uptake curve, showing that
395 N uptake relative to radiation and water use was significantly reduced in real and simulated crop
396 growth.

397 Our conclusions from this very preliminary analysis using a single crop model are that models
398 should be examined for their ability to represent resource use efficiencies under ambient and
399 elevated CO₂ concentrations and, more importantly, how models portray the trade-offs between
400 resources. The upper part of Figure 2 can also be used to estimate resource co-limitation if the
401 upper-limit of resource utilization efficiency can be defined (Cossani *et al.*, 2010). Theory
402 developed in ecology predicts that plant growth is maximized when all resources are equally
403 non-limiting (Sperfeld, 2016) and several experimental and modelling studies have shown that
404 crop yield is often co-limited by water and N (Cossani and Sadras, 2018), and theory from
405 ecology have been introduced in agricultural science and can provide a theoretical framework to
406 test model consistency and help understanding uncertainties when crop models are used in IAMs
407 studies such as those used recently in the IPCC. The identity used here as an illustration of the
408 proposed approach can be easily modified to account for N utilization efficiency and other
409 identities can be worked out (including abiotic factors) to fit the aim of a study. Such work
410 cannot be solely model-based but requires the analysis of existing experiments and where
411 necessary the making of new experiments to test our models. Such experiments are rare, partly
412 because experiments are often designed in the absence of clear theoretical deductive analysis.
413 For example, even in the very comprehensive Maricopa FACE experiment used here, an
414 emphasis on the interactions between water and N resource utilisation efficiencies would have

415 resulted in parallel measurement of N as well as water uptake, while only water uptake was
416 measured.

417 **6.2. | Impacts, adaptation and mitigation in integrated assessment studies**

418 Our second point to improve future impacts and assessments analyses concerns how impacts,
419 adaptation and mitigation have historically been assessed by different communities using
420 different methods. This history is reflected in the structure of the IPCC reports, with each of
421 these, especially mitigation, being treated separately. This separation has also been reflected in
422 many policy domains. Recent progress and trends have helped to break down these silos. One
423 example of this is climate-smart agriculture. This idea was borne from the need to integrate
424 climate adaptation and mitigation. Early progress in climate-smart agriculture came through
425 intellectual and political leadership (Lipper *et al.*, 2014), with the evidence base supporting the
426 identification of specific climate-smart agriculture practices coming later (e.g. Rosenzweig *et*
427 *al.*, 2016). Similarly, introduction of carbon taxes, carbon prices or greenhouse gas footprint
428 labelling and similar programs necessitates re-evaluation of risk and returns in all components of
429 food systems which could include addressing the implications of increasingly frequent
430 disruptions from climate extremes (Lim Camacho *et al.* 2017).

431 IAMs are one way of assessing the integration of adaptation and mitigation. Efforts to include
432 agriculture in IAMs is relatively new and a number of challenges need to be addressed (Ewert *et*
433 *al.*, 2015). Whilst crop models are generally responsive to climate, the range of crops that can be
434 simulated is not sufficiently broad for a full assessment of food security. Further, disparities
435 between IAMs and crop models in spatial scale, treatment of uncertainty, data demand and
436 representation of agricultural management all limit the extent of crop model integration into
437 IAMs that is currently possible. Whilst significant progress is being made with these challenges
438 (Ruane *et al.*, 2017), it is likely that more than one approach is needed if we are to capture the


439 range of trade-offs and synergies that are important to food systems (Vermeulen *et al.*, 2013)
440 and relevant to policy design and development in the huge variety of contexts that exist globally.
441 One particularly important challenge, for any holistic approach to food systems and climate
442 change, is to develop a framing for research that recognises that emissions occur across the full
443 range of activities that deliver food security, not only agricultural production (Whitfield *et al.*,
444 2018). Thus the idea of climate-smart food systems has emerged as way to take a more
445 comprehensive look at how climate, food and human activities are interrelated.

446 Progress in climate-smart food systems can be expected to come from a number of promising
447 avenues. IAMs have the potential to be an important tool for allowing a broader and more
448 complete view of agricultural impacts, adaptation and mitigation but as argued earlier, can be
449 limited in their ability to include locally-important factors. Risk assessment methods provide
450 another set of approaches (Challinor *et al.*, 2018a). Working with stakeholders and using
451 multiple methods to identify the timing of key risks is one approach that has been shown to
452 work within constrained systems (Challinor *et al.*, 2016) but is not without its costs and risks
453 (Cvitanovic *et al.* 2019). The review of Challinor *et al.* (2018b) found increasing transparency
454 and inter-comparability in risk assessments to be an important aspect to future work. While
455 studies often address uncertainty, the nature of the treatment and the assumptions underlying
456 that analysis are often unclear. Paraphrasing ESM3 from Wesselink *et al.* (2015), we can list
457 some sources of this lack of clarity: the question of whether and how observations been used,
458 and if so whether measurement uncertainty been accounted for; which uncertainties in model
459 inputs (e.g. initial conditions, boundary conditions, physical constants, driving variables) and
460 model structure (e.g. inaccuracy in model equations, spatial and temporal discretization) have
461 been assessed?; have intrinsic and non-measurable stochastic variability (e.g. fundamental limits
462 to predictability resulting from chaotic processes) and uncertainty resulting from explicit
463 variation of model parameters (i.e. potential over- or under- estimation of uncertainty when

464 producing a perturbed-parameter ensemble) been assessed? Uncertainty also arises from
465 insufficient ensemble size (i.e. potential under-estimation of uncertainty due to not capturing the
466 full range of possible model responses) and the use (or not) of expert judgement.

467 Whitfield *et al.* (2018) set out an agenda for climate-smart food systems research, arguing that a
468 number of fundamental questions need to be answered, including: what is climate smartness and
469 how do we measure it?; what trade-offs emerge from climate-smart practices?; how do theory-
470 based climate-smart actions differ across spatial scales?; which climate-smart actions are
471 feasible and attractive?; in which systems and at which scales is climate smartness evident?; and
472 finally, how can diet choices contribute to the climate smartness of the food system in the long
473 term?

474 Issues of spatial scale play a key role in agriculture and climate change, as highlighted by
475 Whitfield *et al.* (2018) for climate-smart food systems, and by many authors for the narrow and
476 older field of crop-climate modelling (Hansen and Jones, 2000; van Bussel *et al.*, 2011,
477 Challinor *et al.*, 2015). Food systems cross international boundaries and recent work has
478 highlighted how climate risks cross both sectors and international boundaries. Challinor *et al.*,
479 (2018b) and The Royal Society (2017) concluded that complex risk transmission mechanisms of
480 this sort cannot be assessed using existing impacts, adaptation and mitigation research alone.
481 Rather, a range of approaches are needed, including expert judgement, interactive scenario
482 building, global systems science, innovative use of climate and integrated assessment models,
483 and methods to understand societal responses to climate risk (Figure 3). These are the types of
484 issues and approaches addressed by policy design and development groups in government and in
485 industry and there is likely much to learn from them in relation to developing effective climate-
486 smart food systems: integrating policy, practice and research.



487

488 **Figure 3.** The range of approaches that are needed, including expert judgement, interactive
 489 scenario building, global systems science, innovative use of climate and integrated assessment
 490 models, and methods to understand, project societal responses to climate risks.

491 **7 | Conclusion**

492 The IPCC ARs have evolved over 34 years since AR1. During this time, several themes have
 493 become apparent, which we have tried to identify in this review. There has been a plethora of
 494 modelling studies on the impacts, with and without adaptation, on a wide range of crops and in
 495 many regions. Results from these more than 2100 studies show consistently both the adverse
 496 effect of climate change on a basic element of food security, namely food production and the
 497 significant potential value of adaptation in reducing these impacts. Over the IPCC cycles, an
 498 increasing array of mitigation approaches has been treated by both top-down and bottom-up
 499 approaches and the range of adaptation options considered has both become more nuanced and
 500 broader. These are positive evolutions in the synthesis and evaluation of research that is the role
 501 of the IPCC authors and reviewers. However, there are large remaining gaps – particularly with

502 respect to impacts, adaptation and mitigation in the livestock sector. The lack of quantitative
503 data on livestock in the five ARs was a shock for us as ‘historical’ reviewers which needs
504 addressing as does an increased attention to non-production aspects of food systems. We also
505 suggest a couple of ‘closer to now’ issues on the interactions between resource use efficiencies
506 and the future role of IAMs that may become important in the context of climate change
507 assessment in the near term. In the longer term, future directions for research in agriculture and
508 food will be to ask as much about efficiency and food demand issues as the past has been
509 concerned with adequacy of food supply and environmental outcomes. Thus, issues such as
510 human nutrition and health, diet and obesity, food waste, circular and local food systems could
511 become dominant themes for food systems research and thereby the foci for future IPCC
512 Assessment Reports.

513 **Acknowledgements**

514 J.R. Porter was supported by the Agropolis Fondation under the reference ID 1502-602 through
515 the « Investissements d’avenir » programme (Labex Agro:ANR-10-LABX-0001-01), under the
516 frame of I-SITE MUSE (ANR-16-IDEX-0006»).

517 P. Martre was supported by the EU project H2020 SolACE (grant agreement no. 727247).

518 **References**

519 Asseng, S., Martre, P., Maiorano, A., *et al.* (2019). Climate change impact and adaptation for
520 wheat protein. *Global Change Biology*, 25, 155-173.

521 Braun, J. von, Gulati, A., Kharas, H. (2017). Key policy actions for sustainable land and water
522 use to serve people. *Economics*, 11, 2017-2032.

523 Bennetzen, E.H., Smith, P., Porter, J.R. (2016). Decoupling of greenhouse gas emissions from
524 global agricultural production: 1970–2050. *Global Change Biology*, 22, 763–781.

525 Challinor, A.J., Adger, W.N., Benton, T.G., *et al.* (2018b). Transmission of climate risks across
526 sectors and borders. *Philosophical Transactions of the Royal Society A: Mathematical, Physical*
527 *and Engineering Sciences*, 376, Article: 20170301.

528 Challinor, A.J., Koehler, A.K., Ramirez-Villegas, J., *et al.* (2016). Current warming will reduce
529 yields unless maize breeding and seed systems adapt immediately. *Nature Climate Change*, 6,
530 954-958.

531 Challinor, A., Martre, P., Asseng, S., *et al.* (2014). Making the most of climate impacts
532 ensembles. *Nature Climate Change*, 4, 77-80.

533 Challinor, A.J., Müller, C., Asseng, S., *et al.* (2018a). Improving the use of crop models for risk
534 assessment and climate change adaptation, *Agricultural Systems*, 159, 296-306.

535 Challinor, A.J., Parkes, B., Ramirez-Villegas, J. (2015). Crop yield response to climate change
536 varies with cropping intensity. *Global Change Biology*, 21, 1679-88.

537 Cossani, C.M., Slafer, G.A., Savin, R. (2010). Co-limitation of nitrogen and water, and yield
538 and resource-use efficiencies of wheat and barley. *Crop and Pasture Science*, 61, 844-851.

539 Cossani, C.M., Sadras, V.O. (2018). Water–nitrogen colimitation in grain crops. *Advances in*
540 *Agronomy*, 150, 231-274.

541 Cvitanovic, C., Howden, M., Colvin, R.M., *et al.* (2019). Maximising the benefits of
542 participatory climate adaptation research by understanding and managing the associated
543 challenges and risks. *Environmental Science and Policy*, 94, 20-31.

544 Evans, L.T., Fischer, R.A. (1999). Yield potential: Its definition, measurement, and significance.
545 *Crop Science*, 39, 1544-1551.

546 Ewert, F., Rötter, R.P., Bindi, M., *et al.* (2015). Crop modelling for integrated assessment of risk
547 to food production from climate change. *Environmental Modelling and Software*, 72, 287-303.

548 Foulkes, M.J., Slafer, G.A., Davies, W.J. *et al.* (2011). Raising yield potential of wheat. III.
549 Optimizing partitioning to grain while maintaining lodging resistance. *Journal of Experimental*
550 *Botany*, 62, 469-486.

551 Hansen, J.W., Jones, J.W. (2000). Scaling-up crop models for climate variability applications.
552 *Agricultural Systems*, 65, 43-72.

553 Justes, E., Mary, B., Meynard, J.M., *et al.*, (1994) Determination of a critical nitrogen dilution
554 curve for winter wheat crops. *Annals of Botany*, 74, 397-407.

555 Kaya, Y., Yokoburi, K. (1997). *Environment, energy, and economy: strategies for*
556 *sustainability*. Tokyo: United Nations University Press.

557 Kimball, B.A., Pinter Jr, P.J., Lamorte, R.L., *et al.* (2017). Data from the Arizona FACE (Free-
558 Air CO₂ Enrichment) experiments on wheat at ample and limiting levels of water and nitrogen.
559 *Open Data Journal for Agricultural Research*, 3, 29-38.

560 Lim-Camacho, L., Plagányi, E.E., Crimp, S.J., *et al.* (2017). Complex resource supply chains
561 display higher resilience to simulated climate shocks. *Global Environmental Change*, 46, 126-
562 138.

563 Lipper, L., Thornton, P., Campbell, B.M., *et al.* (2014). Climate-smart agriculture for food
564 security. *Nature Climate Change*, 4, 1068-1072.

565 Macauley, H. (2015). Cereal crops: Rice, maize, millet, sorghum, wheat. *Feeding Africa*. Abdou
566 Diouf International Conference Center, Dakar, Senegal

567 Martre, P., Jamieson, P.D., Semenov, M.A., *et al.* (2006). Modelling protein content and
568 composition in relation to crop nitrogen dynamics for wheat. *European Journal of Agronomy*,
569 25, 138-154.

570 Martre, P., Dambreville, A. (2018). A model of leaf coordination to scale-up leaf expansion
571 from the organ to the canopy. *Plant Physiology*, 176, 704-716.

572 Martre, P., Wallach, D., Asseng, S. *et al.* (2015). Multimodel ensembles of wheat growth: many
573 models are better than one. *Global Change Biology*, 21, 911-925.

574 McIntyre, Beverly D., Herren, Hans R., Wakhungu, Judi, and Watson, Robert T., (eds.)
575 *International Assessment of Agricultural Knowledge, Science and Technology for Development*
576 *Synthesis Report*. International Assessment of Agricultural Knowledge, Science and Technology
577 for Development, Washinton, DC, USA, pp. 1-11.

578 Nelson, G.C., Valin, H., Sands, R.D., *et al.* (2014). Economic response in agriculture to climate
579 change. *Proceedings of the National Academy of Sciences*, 111, 3274-3279.

580 Pardey, P.G., Chan-Kang, C., Beddow, J.M., *et al.* (2016). Agricultural R&D is on the move.
581 *Nature*, 537, 301-3.

582 Porter, J.R., Christensen, S. (2013). Deconstructing crop processes and models via identities.
583 *Plant Cell and Environment*, 36, 1919-1925.

584 Porter, J.R., Xie, L., Challinor, A.J., *et al.* (2014). Food security and food production systems.
585 In: C.B. Field, V.R. Barros, D.J. Dokken, K.J. Mach, M.D. Mastrandrea, T.E. Bilir, M.
586 Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma, E.S. Kissel, A.N. Levy, S.
587 MacCracken, P.R. Mastrandrea, & L.L. White (Eds), *Climate change 2014: Impacts*,
588 *adaptation, and vulnerability. Part A: Global and sectoral aspects. Contribution of Working*

589 *Group II to the fifth Assessment Report of the Intergovernmental Panel on Climate Change* (pp.
590 485-533). Cambridge: Cambridge University Press.

591 Porter, J.R., Howden, S.M., Smith P. (2017). Considering agriculture in IPCC assessments.
592 *Nature Climate Change*, 7, 680-683.

593 Reilly, J., Hohmann, N., Kane, S. (1994). Climate change and agricultural trade: who benefits,
594 who loses? *Global Environmental Change*, 4, 24-36.

595 Rosenzweig, C., Parry, M.L. (1994) Potential impact of climate change on world food supply.
596 *Nature*, 367, 133-138.

597 .

598 Rickards, L., Howden, S.M. (2012) Transformational adaptation: agriculture and climate
599 change. *Crop and Pasture Science*, 63, 240-250.

600 Rogers, A., Gibon, Y., Stitt, M., *et al.* (2006). Increased C availability at elevated carbon
601 dioxide concentration improves N assimilation in a legume. *Plant, Cell and Environment*, 29,
602 1651-1658.

603 Rivera-Ferre, M.G., López-i-Gelats, F., Howden S.M., *et al.* (2016). Re-framing the climate
604 change debate in the livestock sector: mitigation and adaptation options. *WIREs Climate*
605 *Change*, 7, 869-892.

606 Rosenzweig, C., Elliott, J., Deryng, D., *et al.* (2014). Assessing agricultural risks of climate
607 change in the 21st century in a global gridded crop model intercomparison. *Proceedings of the*
608 *National Academy of Sciences*, 111, 3268-3273.

609 Rosenstock, T.S., Lamanna, C., Chesterman, S., *et al.* (2016). *The scientific basis of climate-*
610 *smart agriculture: A systematic review protocol*. CCAFS Working Paper no. 138. Copenhagen,

611 Denmark: CGIAR Research Program on Climate Change, Agriculture and Food Security
612 (CCAFS).

613 Royal Society of London (2017). *Climate updates – What have we learnt since the IPCC 5th*
614 *Assessment Report?* The Royal Society of London, UK.

615 Ruane, A.C., Rosenzweig, C., Asseng, S., *et al.* (2017). An AgMIP framework for improved
616 agricultural representation in integrated assessment models. *Environmental Research Letters*,
617 12, 125003-125016.

618 Sinclair, T.R., Muchow, R.C. (1999). Radiation use efficiency. *Advances in Agronomy*, 65, 216-
619 265.

620 Sperfeld, E., Raubenheimer, D., Wacker, A., (2016). Bridging factorial and gradient concepts of
621 resource co-limitation: towards a general framework applied to consumers. *Ecological Letters*,
622 19, 201–215.

623 Teixeira, A.I., George, M., Herreman, T. (2014). The impact of water and nitrogen limitation on
624 maize biomass and resource-use efficiencies for radiation, water and nitrogen. *Field Crops*
625 *Research*, 168, 109–118.

626 Tilman, D., Clark, M. (2014). Global diets link environmental sustainability and human health.
627 *Nature*, 515, 518–522.

628 Van Bussel, L.G.I., Ewert, F., Leffelaar, P.A. (2011). Effects of data aggregation on simulations
629 of crop phenology. *Agriculture, Ecosystems and Environment*, 142, 75-84.

630 Vermeulen, S.J., Challinor, A.J., Thornton, P., *et al.* (2013). Addressing uncertainty in
631 adaptation planning for agriculture. *Proceedings of the National Academy of Sciences*, 110,
632 8357-8362.

- 633 Wesselink, A., Challinor, A.J., Watson, J., *et al.* (2015). Equipped to deal with uncertainty in
634 climate and impacts predictions: lessons from internal peer review. *Climatic Change*, 132, 1-14
- 635 Whitfield, S., Challinor, A.J., Rees, R.M. (2018). Frontiers in climate smart food systems:
636 Outlining the research space. *Frontiers in Sustainable Food Systems*, 2, 2.