

This is a repository copy of *An analogue of the Perelomov-Popov formula for the Lie superalgebra $Q(N)$* .

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/146939/>

Version: Accepted Version

Article:

Nazarov, Maxim and Grigoryev, Timofey (2019) An analogue of the Perelomov-Popov formula for the Lie superalgebra $Q(N)$. *Functional Analysis and Its Applications*. pp. 304-308. ISSN 0016-2663

<https://doi.org/10.1134/S0016266319040075>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

AN ANALOGUE OF THE PERELOMOV-POPOV FORMULA FOR THE LIE SUPERALGEBRA $\mathfrak{q}(N)$

T. A. GRIGORYEV AND M. L. NAZAROV

ABSTRACT. We study the center of the universal enveloping algebra of the strange Lie superalgebra $\mathfrak{q}(N)$. We obtain an analogue of the well known Perelomov-Popov formula [6] for central elements of this algebra – an expression of the central characters through the highest weight parameters.

1. INTRODUCTION

1.1. **Lie superalgebra $\mathfrak{q}(N)$.** The strange Lie superalgebra $\mathfrak{q}(N)$ can be realized as a subalgebra in the general linear Lie superalgebra $\mathfrak{gl}(N|N)$ over the complex field, see for instance [2]. If the indices i and j range over $-N, \dots, -1, 1, \dots, N$ then the $4N^2$ elements E_{ij} span the algebra $\mathfrak{gl}(N|N)$ as a vector space. The Lie superbracket on $\mathfrak{gl}(N|N)$ is defined by

$$[E_{ij}, E_{kl}] = \delta_{jk} E_{il} - (-1)^{(\bar{i}+\bar{j})(\bar{k}+\bar{l})} \delta_{il} E_{kj}$$

where

$$\bar{k} = \begin{cases} 0 & \text{if } k > 0, \\ 1 & \text{if } k < 0. \end{cases}$$

Then $\mathfrak{q}(N) \subset \mathfrak{gl}(N|N)$ is the subalgebra of fixed points of the involution

$$\eta : E_{ij} \mapsto E_{-i-j}.$$

Hence as a vector space $\mathfrak{q}(N)$ is spanned by the $2N^2$ elements

$$F_{ij} = E_{ij} + E_{-i-j}$$

with $i > 0$. The Lie superbracket on $\mathfrak{q}(N)$ is then described by

$$(1) \quad [F_{ij}, F_{kl}] = \delta_{kj} F_{il} - (-1)^{(\bar{i}+\bar{j})(\bar{k}+\bar{l})} \delta_{il} F_{kj} + \delta_{k-j} F_{-il} - (-1)^{(\bar{i}+\bar{j})(\bar{k}+\bar{l})} \delta_{-il} F_{k-j}.$$

1.2. **Casimir elements in $U(\mathfrak{q}(N))$.** Unless otherwise stated, we will be assuming that the indices in the expressions below range over $-N, \dots, -1, 1, \dots, N$. For any $n = 1, 2, \dots$ consider the elements of the universal enveloping algebra $U(\mathfrak{q}(N))$ first proposed in [7]:

$$(2) \quad C_{ij}^{(n)} = \sum_{k_1, \dots, k_{n-1}} F_{ik_1} (-1)^{\bar{k}_1} F_{k_1 k_2} (-1)^{\bar{k}_2} \cdots F_{k_{n-2} k_{n-1}} (-1)^{\bar{k}_{n-1}} F_{k_{n-1} j}.$$

Taking into account the recurrence relations

$$C_{ij}^{(n+1)} = \sum_k F_{ik} (-1)^{\bar{k}} C_{kj}^{(n)}$$

one can find the supercommutator

$$[F_{ij}, C_{kl}^{(n)}] = \delta_{kj} C_{il}^{(n)} - (-1)^{(\bar{i}+\bar{j})(\bar{k}+\bar{l})} \delta_{il} C_{kj}^{(n)} + \delta_{k-j} C_{-il}^{(n)} - (-1)^{(\bar{i}+\bar{j})(\bar{k}+\bar{l})} \delta_{-il} C_{k-j}^{(n)}$$

which is similar to the superbracket (1) between the generators of $\mathfrak{q}(N)$. It is then easy to see that the elements

$$c_n = \sum_i C_{ii}^{(n)}$$

are central in $U(\mathfrak{q}(N))$. Moreover, by using the relation

$$C_{-i-j}^{(n)} = (-1)^{n-1} C_{ij}^{(n)}$$

following from (2) we see that $c_n = 0$ if n is even. This is why we will be only interested in the c_n with n odd. These are the *Casimir elements* for the Lie superalgebra $\mathfrak{q}(N)$ as introduced in [3]. It was shown in [4] that these elements generate the center of $U(\mathfrak{q}(N))$.

1.3. Harish-Chandra homomorphism. Let v be a singular vector of any irreducible finite-dimensional representation of the Lie superalgebra $\mathfrak{q}(N)$ relative to the natural triangular decomposition $\mathfrak{q}(N) = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$ where

$$\begin{aligned} \mathfrak{n}_- &= \text{span} \{ F_{ij} \mid |i| > |j| \}, \\ \mathfrak{h} &= \text{span} \{ F_{ij} \mid |i| = |j| \}, \\ \mathfrak{n}_+ &= \text{span} \{ F_{ij} \mid |i| < |j| \}. \end{aligned}$$

Then the following equalities hold:

$$(3) \quad F_{ij} \cdot v = 0 \quad \text{for} \quad |i| < |j|,$$

$$(4) \quad F_{ii} \cdot v = \lambda_i v \quad \text{for} \quad i > 0.$$

Here the λ_i are the eigenvalues of the elements $F_{ii} = F_{-i-i}$ of the even part of the Cartan subalgebra $\mathfrak{h}_0 = \text{span} \{ F_{ii} \mid i > 0 \}$. They depend on the particular representation of $\mathfrak{q}(N)$. Let $\lambda \in \mathfrak{h}_0^*$ be the highest weight of the representation so that $\lambda(F_{ii}) = \lambda_i$ for $i > 0$.

The generators of $\mathfrak{q}(N)$ in the summands of (2) can always be rearranged in such a way that in each monomial left-to-right first go the lowering operators, then the operators from Cartan subalgebra and last the raising operators. Here the lowering operators are elements of \mathfrak{n}_- and the raising operators are elements of \mathfrak{n}_+ . The operators from Cartan subalgebra can also be rearranged so that elements from its even part \mathfrak{h}_0 go after those from its odd part $\mathfrak{h}_1 = \text{span} \{ F_{-ii} \mid i > 0 \}$. It suffices to use the supercommutation relations (1) to achieve this. The part of the resulting sum which belongs to $U(\mathfrak{h})$ is well-defined. For the sum (2) with $i = j$ this is its image under the *Harish-Chandra homomorphism* χ , see [2].

The subspace formed by the vectors of an irreducible representation of $\mathfrak{q}(N)$ satisfying (3),(4) is called the singular subspace. The peculiarity of $\mathfrak{q}(N)$ shows in the fact that the singular subspace of an irreducible representation is not one-dimensional, but is irreducible over the Cartan subalgebra $\mathfrak{h} = \mathfrak{h}_0 \oplus \mathfrak{h}_1$ for which we have $U(\mathfrak{h}) = S(\mathfrak{h}_0) \otimes \wedge(\mathfrak{h}_1)$, see [5].

Due to (2) the Casimir elements are even and therefore commute with the whole algebra $U(\mathfrak{q}(N))$ in the usual non \mathbb{Z}_2 -graded sense. This implies that they act as scalar operators in the irreducible representations. This allows us to consider their eigenvalues when acting on some fixed singular vector instead and the computations in this case are rather simple.

Let n be odd and v be a singular vector of an irreducible representation of $\mathfrak{q}(N)$. Then

$$(5) \quad c_n \cdot v = \chi(c_n) \cdot v = \chi(c_n) \Big|_{F_{ii}=\lambda_i} v \quad \text{for} \quad i > 0.$$

For $i > 0$ we will automatically substitute $F_{ii} \mapsto \lambda_i$ after applying the homomorphism χ as we did in (5). Hence we will be describing the action on the singular vector explicitly.

2. COMPUTATIONS

2.1. Recurrence relations. Here we will derive a recurrence relation for the images of the elements $C_{ii}^{(n)}$ with $n = 1, 3, \dots$ under the Harish-Chandra homomorphism. For brevity we will denote by G_i the element $F_{-ii} = F_{i-i}$ of the odd part of the Cartan subalgebra.

Proposition 1. *For $i > 0$ we have $\chi(G_i^2) = \lambda_i$.*

Proof. We have $G_i^2 \cdot v = F_{-ii}^2 \cdot v = \frac{1}{2} [F_{-ii}, F_{-ii}] \cdot v = F_{ii} \cdot v = \lambda_i v$ for $i > 0$.

Proposition 2. *We have $C_{ij}^{(n)} \cdot v = 0$ whenever $|i| < |j|$.*

Proof. Suppose that $|i| < |j|$. Then $C_{ij}^{(1)} \cdot v = F_{ij} \cdot v = 0$. Let us use the induction on n :

$$C_{ij}^{(n+1)} \cdot v = \sum_k F_{ik} (-1)^{\bar{k}} C_{kj}^{(n)} \cdot v = \sum_{|k| \geq |j|} (-1)^{\bar{k}} [F_{ik}, C_{kj}^{(n)}] \cdot v = \sum_{|k| \geq |j|} (-1)^{\bar{k}} C_{ij}^{(n)} \cdot v = 0.$$

Proposition 3. *For $i > 0$ we have*

$$\chi(C_{ii}^{(n+1)}) = \lambda_i \chi(C_{ii}^{(n)}) - G_i \chi(C_{-ii}^{(n)}) - \sum_{|k| > i} \chi(C_{kk}^{(n)}).$$

Proof. For $i > 0$ the vector $C_{ii}^{(n+1)} \cdot v$ equals

$$\begin{aligned} \sum_{|k| \geq i} F_{ik} (-1)^{\bar{k}} C_{ki}^{(n)} \cdot v &= (F_{ii} C_{ii}^{(n)} - F_{i-i} C_{-ii}^{(n)}) \cdot v + \sum_{|k| > i} (-1)^{\bar{k}} [F_{ik}, C_{ki}^{(n)}] \cdot v = \\ &= (\lambda_i C_{ii}^{(n)} - G_i C_{-ii}^{(n)}) \cdot v + \sum_{|k| > i} (-1)^{\bar{k}} (C_{ii}^{(n)} - (-1)^{\bar{k}} C_{kk}^{(n)}) \cdot v = \\ &= (\lambda_i C_{ii}^{(n)} - G_i C_{-ii}^{(n)} - \sum_{|k| > i} C_{kk}^{(n)}) \cdot v. \end{aligned}$$

Corollary 1. *For $i > 0$ and $m = 1, 2, \dots$ we get $\chi(C_{ii}^{(2m+1)}) = \lambda_i \chi(C_{ii}^{(2m)}) - G_i \chi(C_{-ii}^{(2m)})$.*

Proposition 4. *For $i > 0$ we have*

$$\chi(C_{-ii}^{(n+1)}) = G_i \chi(C_{ii}^{(n)}) - \lambda_i \chi(C_{-ii}^{(n)}) - \sum_{|k| > i} (-1)^{\bar{k}} \chi(C_{k-k}^{(n)}).$$

Proof. For $i > 0$ the vector $C_{-ii}^{(n+1)} \cdot v$ equals

$$\begin{aligned} \sum_{|k| \geq i} F_{-ik} (-1)^{\bar{k}} C_{ki}^{(n)} \cdot v &= (F_{-ii} C_{ii}^{(n)} - F_{ii} C_{-ii}^{(n)}) \cdot v + \sum_{|k| > i} (-1)^{\bar{k}} [F_{-ik}, C_{ki}^{(n)}] \cdot v = \\ &= (G_i C_{ii}^{(n)} - \lambda_i C_{-ii}^{(n)}) \cdot v + \sum_{|k| > i} (-1)^{\bar{k}} (C_{-ii}^{(n)} - (-1)^{\bar{k}(\bar{k}+1)} C_{k-k}^{(n)}) \cdot v = \\ &= (G_i C_{ii}^{(n)} - \lambda_i C_{-ii}^{(n)}) \cdot v - \sum_{|k| > i} (-1)^{\bar{k}} C_{k-k}^{(n)} \cdot v. \end{aligned}$$

Corollary 2. *For $i > 0$ and $m = 1, 2, \dots$ we get $\chi(C_{-ii}^{(2m)}) = G_i \chi(C_{ii}^{(2m-1)}) - \lambda_i \chi(C_{-ii}^{(2m-1)})$.*

Proposition 5. *For $i > 0$ and $m = 1, 2, \dots$ we have the relation*

$$\chi(C_{ii}^{(2m+1)}) = \chi(C_{-i-i}^{(2m+1)}) = \lambda_i (\lambda_i - 1) \chi(C_{ii}^{(2m-1)}) - 2\lambda_i \sum_{j > i} \chi(C_{jj}^{(2m-1)}).$$

Proof. If $i > 0$ then the vector $C_{ii}^{(2m+1)} \cdot v$ equals

$$\begin{aligned} & (\lambda_i C_{ii}^{(2m)} - G_i C_{-ii}^{(2m)}) \cdot v = \\ & \lambda_i (\lambda_i C_{ii}^{(2m-1)} - G_i C_{-ii}^{(2m-1)} - \sum_{|j|>i} C_{jj}^{(2m-1)}) \cdot v - G_i (G_i C_{ii}^{(2m-1)} - \lambda_i C_{-ii}^{(2m-1)}) \cdot v = \\ & (\lambda_i^2 - G_i^2) C_{ii}^{(2m-1)} \cdot v - \lambda_i \sum_{|j|>i} C_{jj}^{(2m-1)} \cdot v = \lambda_i (\lambda_i - 1) C_{ii}^{(2m-1)} \cdot v - 2\lambda_i \sum_{j>i} C_{jj}^{(2m-1)} \cdot v. \end{aligned}$$

Corollary 3. For $i > 0$ and $m = 0, 1, 2, \dots$ we have $\chi(C_{ii}^{(2m+1)}) = \sum_{j=1}^N (A^m)_{ij} \lambda_j$ where

$$A = \begin{pmatrix} \lambda_1(\lambda_1 - 1) & -2\lambda_1 & \cdots & -2\lambda_1 \\ 0 & \lambda_2(\lambda_2 - 1) & \cdots & -2\lambda_2 \\ \vdots & & \ddots & \vdots \\ 0 & \cdots & 0 & \lambda_N(\lambda_N - 1) \end{pmatrix}.$$

Proof. This follows from Proposition 5 by taking into account that $\chi(C_{ii}^{(1)}) = \chi(F_{ii}) = \lambda_i$.

2.2. Generating functions. In order to compute $\chi(c_{2m+1})$ more explicitly, for each $i > 0$ consider the generating function

$$(6) \quad \mu_i(u) = \sum_{m=0}^{\infty} \chi(C_{ii}^{(2m+1)}) u^{-2m-1} = u \sum_{j=1}^N ((u^2 - A)^{-1})_{ij} \lambda_j$$

and write

$$A = \Lambda(\Lambda - 1) - 2\Lambda\Delta(1 - \Delta)^{-1}$$

where $\Delta_{ij} = \delta_{i,j-1}$ and $\Lambda = \text{diag}(\lambda_i)$. Denote

$$\Pi = (u^2 - \Lambda(\Lambda + 1))(u^2 - \Lambda(\Lambda - 1))^{-1}.$$

Then

$$(u^2 - A)^{-1} = \frac{1}{2} (1 - \Pi) (1 - \Delta\Pi)^{-1} (1 - \Delta)\Lambda^{-1}.$$

The last two factors here cause all summands but the last in the sum (6) cancel, leaving

$$\begin{aligned} \mu_i(u) &= \frac{u}{2} (1 - \Pi_{ii}) ((1 - \Delta\Pi)^{-1})_{iN} = \\ & \frac{u}{2} (1 - \Pi_{ii}) (\Delta\Pi)_{ii+1} (\Delta\Pi)_{ii+2} \cdots (\Delta\Pi)_{N-1N} = \frac{u}{2} (1 - \Pi_{ii}) \prod_{j>i} \Pi_{jj}. \end{aligned}$$

This can also be written in more straightforward way:

$$\mu_i(u) = \frac{u\lambda_i}{u^2 - \lambda_i(\lambda_i - 1)} \prod_{j>i} \frac{u^2 - \lambda_j(\lambda_j + 1)}{u^2 - \lambda_j(\lambda_j - 1)}.$$

In order to find a generating function $\mu(u)$ of the images of the central elements under χ , we should now sum all the above obtained expressions for $\mu_i(u)$ over the positive values of the index i and recall that $C_{-i-i}^{(2m+1)} = C_{ii}^{(2m+1)}$. By making further cancellations we get

$$\mu(u) = \sum_{m=0}^{\infty} \chi(c_{2m+1}) u^{-2m-1} = 2 \sum_{i=1}^N \mu_i(u) = u \left(1 - \prod_{j=1}^N \Pi_{jj} \right).$$

Finally, we obtain

$$\mu(u) = u \left(1 - \prod_{i=1}^N \frac{u^2 - \lambda_i(\lambda_i + 1)}{u^2 - \lambda_i(\lambda_i - 1)} \right).$$

2.3. Images of the central elements. Introduce a new variable $z = u^{-2}$ and define

$$\tilde{\mu}(z) = u \mu(u) = \sum_{m=0}^{\infty} \chi(c_{2m+1}) z^m = \frac{1}{z} \left(1 - \prod_{i=1}^N \frac{1 - z\lambda_i(\lambda_i + 1)}{1 - z\lambda_i(\lambda_i - 1)} \right).$$

From the above definition it immediately follows that $\chi(c_{2m+1}) = \text{Res}_0 \tilde{\mu}(z) z^{-m-1} dz$. Combining this with our explicit expression for $\tilde{\mu}(z)$ we obtain that $\chi(c_{2m+1})$ equals

$$- \sum_{i=1}^N \text{Res}_{(\lambda_i(\lambda_i-1))^{-1}} \tilde{\mu}(z) z^{-m-1} dz = \sum_{i=1}^N \text{Res}_{(\lambda_i(\lambda_i-1))^{-1}} \prod_{j=1}^N \frac{1 - z\lambda_j(\lambda_j + 1)}{1 - z\lambda_j(\lambda_j - 1)} z^{-m-2} dz$$

where the regularity of the form at the infinity is taken into account. Finally for $m \geq 0$

$$\chi(c_{2m+1}) = 2 \sum_{i=1}^N \lambda_i^{m+1} (\lambda_i - 1)^m \prod_{j \neq i} \frac{\lambda_i(\lambda_i - 1) - \lambda_j(\lambda_j + 1)}{\lambda_i(\lambda_i - 1) - \lambda_j(\lambda_j - 1)}.$$

Note that despite the fact that the last expression is formally a rational function of λ , its denominator always cancels which allows us to regard $\chi(c_{2m+1})$ as a polynomial of λ .

REMARKS

The analogue of the Perelomov-Popov formula for the Lie superalgebra $\mathfrak{q}(N)$ presented here was obtained by the second named author about 30 years ago but left unpublished. Independently but by the same method, this analogue was obtained in [1] and recently by the first named author. Publishing this note gives us an opportunity to review the history of this analogue. We thank Jonathan Brundan, Maria Gorelik, Alexandre Kirillov, Grigori Olshanski, Ivan Penkov, Vera Serganova and Alexander Sergeev for helpful conversations.

REFERENCES

- [1] J. Brundan and A. Kleshchev, *Modular representations of the supergroup $Q(n)$, I*, J. Algebra **260** (2003), 64–98.
- [2] S. Cheng and W. Wang, *Dualities and representations of Lie superalgebras*, AMS, Providence, 2013.
- [3] D. Leites and A. Sergeev, *Casimir operators for Lie superalgebras*, in: E. Ivanov *et al* (eds.), “Supersymmetries and Quantum Symmetries”, JINR, Dubna, 2000, pp. 409–411.
- [4] M. Nazarov and A. Sergeev, *Centralizer construction of the Yangian of the queer Lie superalgebra*, in: J. Bernstein *et al* (eds.), “Studies in Lie Theory”, Boston, Birkhäuser, 2006, pp. 417–441.
- [5] I. B. Penkov, *Characters of typical irreducible finite-dimensional $\mathfrak{q}(n)$ -modules*, Funct. Anal. Appl. **20** (1986), 30–37.
- [6] A. M. Perelomov and V. S. Popov, *Casimir operators for semisimple Lie groups*, Math. USSR Izv. **2** (1968), 1313–1335.
- [7] A. Sergeev, *The centre of enveloping algebra for Lie superalgebra $Q(n, \mathbb{C})$* , Lett. Math. Phys. **7** (1983), 177–179.

T. A. GRIGORYEV: SKOLKOVO INSTITUTE OF SCIENCE AND TECHNOLOGY, SKOLKOVO INNOVATION CENTER, MOSCOW 143026, RUSSIA AND NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF ECONOMICS, DEPARTMENT OF MATHEMATICS, MOSCOW 119048, RUSSIA

M. L. NAZAROV: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, YORK YO10 5DD, UNITED KINGDOM